POLYNOMIAL SOLUTIONS TO DIFFERENCE EQUATIONS CONNECTED TO PAINLEVE' II-VI

Gert Almkvist

1 Introduction.

Around 1989 the following problem circulated: Given

$$a_0 = a_1 = a_2 = a_3 = 1$$

and

$$a_n a_{n-4} = a_{n-1} a_{n-3} + a_{n-2}^2$$

for $n \ge 4$. Show that all a_n are integers (the *Somos sequence*). Soon various solutions appeared (see Gale [1]).

The more general difference equation

$$a_n a_{n-k} = \sum_{j=1}^{[k/2]} a_{n-j} a_{n-k+j}$$

where

$$a_0 = a_1 = \dots = a_{k-1} = 1$$

has integer solutions for k = 4, 5, 6, 7 but not for k > 7. Why 7? There does not seem to be an explanation for this.

Here we shall instead consider similar nonlinear difference equations involving polynomials connected to rational solutions of Painleve' equations II-VI. We have the following table of equations

 P_{II} : Yablonskii-Vorob'ev polynomials [2], [4].

$$P_0 = 1, P_1 = x.$$

$$P_{n+1}P_{n-1} = -4(P_n P_n'' - P_n'^2) + xP_n^2$$

 P_{III} : Umemura polynomials [3]

$$P_0 = P_1 = 1$$

$$P_{n+1}P_{n-1} = -x^4(P_n P_n'' - P_n'^2) - x^3 P_n P_n' + (cx+1)P_n^2$$

 $\mathbf{P}_{IV}:[2].$

$$P_0 = P_1 = 1$$

$$P_{n+1}P_{n-1} = P_n P_n'' - P_n'^2 + (x^2 + 2n - 1)P_n^2$$

and

$$P_0 = 1, P_1 = x$$

$$P_{n+1}P_{n-1} = P_n P_n'' - P_n'^2 + (x^2 + 2n)P_n^2$$

$$\mathbf{P}_V:[5].$$

$$P_0 = P_1 = 1$$

$$P_{n+1}P_{n-1} = x(P_n P_n'' - P_n'^2) + P_n P_n' + (\frac{x}{8} - v + \frac{3n}{8})P_n^2$$

 $\mathbf{P}_{VI}:[6].$

$$P_0 = P_1 = 1$$

$$P_{n+1}P_{n-1} = \frac{(x^2 - 4)^2}{4} (P_n P_n'' - P_n'^2) + \frac{x(x^2 - 4)}{4} P_n P_n' + (cx + (n - \frac{1}{2})^2) P_n^2$$

E.g.

$$y = \frac{P_n(1/x, c)P_{n+1}(1/x, c-1)}{P_n(1/x, c-1)P_{n+1}(1/x, c)}$$

is a rational solution of \mathbf{P}_{III}

$$y'' = \frac{y'^{2}}{y} - \frac{y'}{x} + \frac{ay^{2} + b}{x} + y^{3} - \frac{1}{y}$$

where (see [3])

$$a = 2n - 1 + 2c$$
$$b = 2n + 1 - 2c$$

In this paper we consider the more general equation

$$P_{n+1}P_{n-1} = f(P_n P_n'' - P_n'^2) + gP_n P_n' + hP_n^2$$

with

$$P_0 = P_1 = 1$$

We prove the following

Theorem1 Assume that f, g, h are polynomials without common factor and independent of n. Assume also that consecutive P_n :s are without common factor. Then all P_n are polynomials if and only if

$$ff'' - f'^2 + 3f'g - 2fg' - 2g^2 = 0 \qquad (*)$$

Theorem2 The only solutions of (*) are the following

(i)

 \boldsymbol{f} an arbitrary polynomial

$$g = \frac{f'}{2}$$

(ii)

$$f = (ax+b)^k$$
$$g = a(ax+b)^{k-1}$$

It turns out that these results also hold when

$$h = n + p(x)$$

which occurs in the \mathbf{P}_{IV} and \mathbf{P}_{V} cases. For the \mathbf{P}_{VI} case we have

$$h = n(n-1) + p(x)$$

then we get the condition

$$ff'' - f'^{2} + 3f'g - 2fg' - 2g^{2} + 2f = 0$$

which seems to have only solutions with $\deg(f) \leq 4$.

2. The proofs.

To shorten the proof we use the notation

$$P = P_n$$
$$R = P_{n-1}$$
$$S = P_{n-2}$$

We have to show that

$$f(PP^{\prime\prime}-P^{\prime~2})+gPP^{\prime}+hP^{2}$$

is divisible by ${\cal R}$. We have

$$SP = f(RR'' - R'^2) + gRR' + hR^2 \equiv -fR'^2 \mod R$$

It follows

$$S'P + SP' = f'(RR'' - R'^{2}) + f(RR''' - R'R'') + g'RR' + gR'^{2} + gRR'' + h'r^{2} + 2hRR'$$

and

$$S^2 P' \equiv R'^2 (S'f - Sf' + Sg) - R'R''Sf \qquad mod \ R$$

Differentiating once more we get

$$S''P + 2S'P' + SP'' \equiv R'^{2}(-f'' + 2g' + 2h) + R'R''(-2f' + 3g) - R''^{2}f$$

and

$$S^{3}P'' \equiv R'^{2} \left\{ S^{2}(-f'' + 2g' + 2h) - S'^{2}2f + SS'(2f' - 2g) + SS''f \right\}$$
$$+ R'^{3}R'' \left\{ S^{2}(-2f' + 3g) + SS'2f \right\} + R''^{2} \left\{ -S^{2}f \right\}$$

Hence

$$S^{4}(PP'' - P'^{2}) \equiv R'^{4} \left\{ S^{2}(ff'' - 2fg' + 2f'g - f'^{2} - 2fh - g^{2}) + S'^{2}f^{2} - SS''f^{2} \right\}$$
$$-R'^{3}R''S^{2}fg$$

and

$$S^{4}PP' \equiv R'^{4} \left\{ S^{2}(ff' - fg) - SS'f^{2} \right\} + R'^{3}R''S^{2}f^{2}$$

$$S^4P^2 \equiv R^{\prime \ 4}S^2f^2$$

Adding up we obtain

$$S^{4} \left\{ f(PP'' - P'^{2}) + gPP' + hP^{2} \right\} \equiv$$
$$R'^{4} f \left\{ S^{2}(ff'' - f'^{2} + 3f'g - 2fg' - 2g^{2}) - f(f(SS'' - S'^{2}) + gSS' + hS^{2}) \right\} \equiv 0 \qquad mod \ R$$

if and only if

$$ff'' - f'^2 + 3f'g - 2fg' - 2g^2 = 0$$

since

$$f(SS'' - S'^2) + gSS' + hS^2$$

is divisible by R.

We also use that (R, S) = 1 and (R, R') = 1. If $(R, R') \neq 1$ then R would have a double root causing $(P, R) \neq 1$ considered in C[x].

Remark: In the cases P_{II} and P_{III} one can assume that P_n and P_{n+1} have no common roots so the assumptions in Theorem 1 are fulfilled. But experiments show that Theorem 1 is true without any assum

ptions on the P_n :s.

Proof of Theorem 2: One checks that

$$g = \frac{f'}{2}$$

solves

$$ff'' - f'^{2} + 3f'g - 2fg' - 2g^{2} = 0$$

To find all solutions we make the substitution

$$g = u + \frac{f'}{2}$$

Then we get

$$fu' - \frac{f'}{2}u + u^2 = 0$$

Put

$$w = \frac{1}{u}$$

Then

$$\sqrt{f}w' + \frac{w}{2\sqrt{f}} = (\sqrt{f}w)' = \frac{1}{\sqrt{f}}$$

with the solution

$$w = \frac{1}{\sqrt{f}} \left\{ \int \frac{dx}{\sqrt{f}} + C \right\}$$

and

$$u = \frac{\sqrt{f}}{\int \frac{dx}{\sqrt{f}} + C}$$

The only way u can become a polynomial is when C = 0 and

$$f = (ax+b)^k$$

where k > 0 is an integer. Then

$$\int \frac{dx}{\sqrt{f}} = \int (ax+b)^{-k/2} dx = \frac{(ax+b)^{1-k/2}}{a(1-k/2)}$$

 $\quad \text{and} \quad$

$$u = a(1 - \frac{k}{2})(ax + b)^{k-1}$$

Finally

$$g = u + \frac{f'}{2} = a(ax+b)^{k-1}$$

If

$$h = h_n = n + p(x)$$

we will end up with the following coefficient of S^2

$$-2h_{n-1} + h_n = -2(n-1+p(x)) + n + p(x) = -(n-2+p(x)) = -h_{n-2}$$

so nothing is changed in the proof of Theorem 1.

If

$$h = h_n = n(n-1) + p(x)$$

 then

$$-2h_{n-1} + h_n = -2((n-1)(n-2) + p(x)) + n(n-1) + p(x) =$$
$$-((n-2)(n-3) + p(x)) + 2 = -h_{n-2} + 2$$

which means that the condition (*) is changed to

$$ff'' - f'^{2} + 3f'g - 2fg' - 2g^{2} + 2 = 0$$

If you try to find polynomial solutions with $\deg(f) > 4$ you will run into contradictions. E.g. there are the following solutions

(i)

$$f = x^2 + ax + b$$

$$g = 2x + a$$

(ii)

$$f = (x^2 - 1)^2$$

$$g = x(x^2 - 1)$$

(iii)

$$f = (x^2 + 2x)^2$$

 $g = (x + 1)(x^2 + 2x)$
(iv)
 $f = \frac{(x^2 - 4)^2}{4}$
 $g = \frac{x(x^2 - 4)}{4}$

References:

1.D.Gale, The strange and surprising saga of the Somos sequences, Math. Intelligencer 13 (1991), no 3, 40-42 and no 4, 49-50.

2.S.Fukutani, K.Okamoto, H.Umemura, Special polynomials and the Hirota bilinear relations of the second and the fourth Painleve' equations, Nagoya Math. J. 159 (2000), 179-200.

3.K.Kajiwara, T. Masuda, On the Umemura polynomials for the Painleve' III equation, Physics Letters A 260 (1999), 462-467

 $4.M.Kaneko,\,H.Ochiau,\,On$ the coefficients of Yablonskii-Vorob'ev polynomials, QA 0205178

5.M.Noumi, Y.Yamada, Umemura polynomials for the Painleve' V equation, Physics Letters A 247 (1998), 65-69

6.M.Taneda, Polynomials associated with an algebraic solution of the sixth Painleve' equation, Japan. J. Math. (N.S.) 27 (2001), 257-274.