RELATIVE HYPERBOLICITY AND ARTIN GROUPS

ILYA KAPOVICH AND PAUL SCHUPP

ABSTRACT. Let $G = \langle a_1, \ldots, a_n \, | \, a_i a_j a_i \cdots = a_j a_i a_j \ldots, i < j \rangle$ be an Artin group and let $m_{ij} = m_{ji}$ be the length of each of the sides of the defining relation involving a_i and a_j . We show if all $m_{ij} \geq 7$ then G is relatively hyperbolic in the sense of Farb with respect to the collection of its two-generator subgroups $\langle a_i, a_j \rangle$ for which $m_{ij} < \infty$.

1. Introduction

The notion of a word-hyperbolic group introduced by Gromov [18] has played a pivotal role in the development of geometric group theory for the last fifteen years. In [18, 19] Gromov also suggested a way of generalizing this notion to that of a group relatively hyperbolic with respect to a collection of subgroups called parabolic subgroups. Several researchers proposed ways of making these ideas precise. First, Farb [15] defined relative hyperbolicity by requiring that the Cayley graph of a group with cosets of the parabolic subgroups "coned-off" is a hyperbolic metric space. Later Bowditch [5] gave a rigorous interpretation of Gromov's original approach which mimics the case of a geometrically finite Kleinian group. Recently Yaman [31] showed that the Gromov-Bowditch version of relative hyperbolicity can be characterized in terms of convergence group actions. Bowditch had earlier obtained such a characterization for word-hyperbolic groups [6]. Szczepański [29] investigated the relationship between different versions of relative hyperbolicity. He proved that relative hyperbolicity in the sense of Gromov-Bowditch-Yaman implies that of Farb but not vice versa. Bowditch [5] observed that relative hyperbolicity in the sense of Farb together with what Farb termed the "bounded coset penetration property" implies relative hyperbolicity in the sense of Gromov-Bowditch-Yaman. A number of interesting results regarding relatively hyperbolic groups are obtained in [7, 8, 9, 17, 12, 13, 16, 21, 23, 30].

In particular, Szczepański [30] recently provided a way of constructing groups relatively hyperbolic in the sense of Farb by mimicking the hyperbolization of polyhedra construction.

In this paper we study the class of Artin groups, that is, groups given by a presentation of the form:

(†)
$$G = \langle a_1, \dots, a_n | u_{ij} = u_{ji}, \text{ for } 1 \le i < j \le n \rangle$$

where for $i \neq j$

$$u_{ij} := \underbrace{a_i a_j a_i \dots}_{m_{ij} \text{ terms}}$$

and where $m_{ij} = m_{ji}$ for each i < j. We allow $m_{ij} = \infty$ in which case the relation $u_{ij} = u_{ji}$ is omitted from presentation (†).

In the theory of Artin groups a subgroup of an Artin group G generated by a subset of $\{a_1, \ldots, a_n\}$ is called a *parabolic subgroup* (see, for example, [27]). It follows from the results of Appel-Schupp [2] that if in (\dagger) all $m_{ij} \geq 4$ then for any i < j the two-generator parabolic subgroup $G_{ij} := \langle a_i, a_j \rangle \leq G$ is itself an Artin group with the presentation

$$G_{ij} = \langle a_i, a_j \mid u_{ij} = u_{ji} \rangle.$$

Although we will not use the fact, it is easy to see that for an even $m_{ij} = 2k$ the group G_{ij} is isomorphic to the Baumslag-Solitar group $B(k,k) = \langle x,y|y^{-1}x^ky = x^k\rangle$. Similarly, if $m_{ij} = 2k+1$ is odd, then G_{ij} is isomorphic to the torus-knot group $\langle x,y|x^{2k+1}=y^2\rangle$.

An Artin group is said to be of extra large type if all $m_{ij} \geq 4$ in presentation (†). Even before the general theory of hyperbolic groups Appel and Schupp [2] proved theorems about Artin groups of extra large type by showing that they were "relatively small cancellation" with respect to the collection of subgroups G_{ij} . Our main result here is:

Theorem A. Let G be an Artin group given by presentation (†) above. Assume that for all i < j we have $m_{ij} \geq 7$. Then G is relatively hyperbolic in the sense of Farb with respect to the collection of subgroups $\{G_{ij}|m_{ij} < \infty\}$, where $G_{ij} = \langle a_i, a_j \rangle \leq G$.

Thus Theorem A asserts that G is relatively hyperbolic with respect to the family of non-free two-generator parabolic subgroups. This provides additional justification for using the term "parabolic subgroup" in the context of Artin groups.

Since we do not actually use it, we refer to Farb's paper [15] for the precise definition of the "bounded coset penetration property." But

it is easy to see that if two subgroups H_1 , H_2 of a group G have infinite intersection then G does not satisfy the bounded coset penetration property with respect to the collection (H_1, H_2) (or with respect to any larger collection of subgroups). Thus for an Artin group G as in Theorem A if there are distinct i, j, k such that m_{ij} and m_{ik} are finite then G does not have the bounded coset penetration property with respect to the collection of subgroups G_{ij} . This raises the interesting question of finding different conditions which ensure good group-theoretic properties in the presence of Farb's definition of relative hyperbolicty. It seems plausible that most Artin groups satisfy some sort of a "nested" version of the bounded coset penetration property, especially since many of these groups are known to be CAT(0) and biautomatic (see for example, [3, 4, 10, 25, 28]).

2. ARTIN GROUPS AND SMALL CANCELLATION THEORY

In this section we explain how one can apply small cancellation theory to study Artin groups, even though the given finite presentation (†) does not have good small cancellation properties. In order to do this we assume that the reader is familiar with the basics of standard small cancellation theory. For definitions and terminology see Lyndon and Schupp [22].

Let R_{ij} be the symmetrized set of words in $F(a_i, a_j)$ generated by the defining relator, $u_{ij}u_{ji}^{-1}$, and its inverse. The cancellation between two noninverse elements of R_{ij} can be almost half of each of the words but *not* an entire half. Consider, for example, a product such as

$$(a_1a_2a_1a_2a_1^{-1}a_2^{-1}a_1^{-1}a_2^{-1})(a_2a_1a_2a_1^{-1}a_2^{-1}a_1^{-1}a_2^{-1}a_1)$$

It is not difficult to verify that the set R_{ij} satisfies the "flat" small cancellation condition C(4) - T(4). To exploit this fact we must use the right normal form for elements of $F(a_1, \ldots, a_n)$.

Convention 2.1. For the remainder of this paper we denote $F := F(a_1, ..., a_n)$. In the free group F every nontrivial reduced word w has a unique normal form with exponents

$$w = a_{h_1}^{n_1} ... a_{h_s}^{n_s}$$

where each $h_t \neq h_{t+1}$ and each $n_t \neq 0$. The integer s is the syllable length of w and is written ||w||. The subwords $a_{h_i}^{n_i}$ are called the syllables of w.

Notation 2.2. For each i < j such that $m_{ij} < \infty$ let \mathcal{R}_{ij} be the set of all nontrivial cyclically reduced words in $F(a_i, a_j)$ which are equal to the identity in the group G_{ij} .

By introducing "strips" to study the fine geometry of C(4)-T(4) diagrams, Appel and Schupp [2] proved:

Proposition 2.3. Suppose $2 \leq m_{ij} < \infty$ and let w be a word from \mathcal{R}_{ij} . Then $||w|| \geq 2m_{ij}$.

Notation 2.4. To deal with the Artin group G we now switch to the following *infinite* presentation of G:

$$(\ddagger) \qquad \qquad G = \langle a_1, \dots, a_n \, | \, \mathcal{R} \rangle.$$

where

$$\mathcal{R} = \bigcup_{m_{ij} < \infty} \mathcal{R}_{ij}.$$

Remark 2.5. The point of shifting to the infinite presentation (‡) is that it allows a strong use of minimality. In considering van Kampen diagrams for a word w = 1 in G, it suffices to consider minimal diagrams, that is, diagrams with as few as regions as possible. In a minimal diagram Δ over (‡) distinct regions labelled by relators from the same \mathcal{R}_{ij} cannot have even a vertex in common since they could then be combined into a single region, contradicting minimality. Also, any common edge shared by regions labelled from some \mathcal{R}_{ij} and \mathcal{R}_{il} where $j \neq l$ must be labelled by a power of the generator a_i , that is, the edge label has syllable length one. If G is of extra large type, Proposition 2.3 will show that minimal diagrams for the presentation (‡) have "hyperbolic" C(8)- geometry.

Notation 2.6. As in [22], if Δ is a map and D is a region of Δ , then i(D) will denote the interior degree of D, that is, the number of Δ -interior edges of D, where each edge occurring twice in a boundary cycle of D is counted twice in i(D). Recall also that a region D of Δ is said to be a boundary region if $\partial D \cap \partial \Delta \neq \emptyset$. Note that a boundary region need not contain a boundary edge of Δ as the intersection may contain only vertices of $\partial \Delta$. We will say that a boundary region D is a simple boundary region of Δ if the intersection $\partial D \cap \partial \Delta \neq \emptyset$ is connected and if the edges of $\partial D \cap \partial \Delta$ are consecutively traversed in some boundary cycle of Δ .

Lemma 4.1 in Ch. V of [22] asserts that in a (6,3)-map the boundary of every region is a simple closed curve. In particular this means that if

D is a simple boundary region of a (6,3)-map Δ then the intersection $\partial D \cap \partial \Delta$ is consecutively traversed in some boundary cycle of D.

We shall need the following "basic fact" of small cancellation theory [22]:

Proposition 2.7. Let Δ be a (6,3)-map without vertices of degree one and with no simple boundary regions of interior degree zero. Then at least one of the following occurs:

- (a) The map Δ contains at least two simple boundary regions of interior degree one.
- (b) The map Δ has at least three simple boundary regions of interior degree at most three.

Proposition 2.3 and Proposition 2.7 together imply [2]:

Proposition 2.8. Suppose G is given by (\dagger) where all $m_{ij} \geq 4$. Then:

- (1) Every minimal diagram over the infinite presentation (\ddagger) satisfies C(8) and every interior edge in such a diagram is labeled by a power of some generator a_i .
- (2) If w is a nontrivial freely reduced word representing 1 in G then w contains a subword v such that v is also a subword of some $r \in \mathcal{R}_{ij}$ with r = vu, $||u|| \leq 3$ and $||v|| \geq 2m_{ij} 3$.
- (3) If $m_{ij}, m_{ik} < \infty$ for some $k \neq j$ then $G_{ij} \cap G_{ik} = \langle a_i \rangle$. Moreover, if $\{i, j\} \cap \{t, k\} = \emptyset$ then $G_{ij} \cap G_{tk} = \{1\}$ in G.

Convention 2.9. For the remainder of this section we assume that G is given by presentation (†) where all $m_{ij} \geq 4$.

Definition 2.10. Let w be a freely reduced word in $F(a_1, \ldots, a_n)$. We say that w is Artin-reduced if w does not contain a subword v such that v is also a subword of some $r \in \mathcal{R}_{ij}$ with r = vu, $||u|| \leq 3$.

Similarly, a word w is strongly Artin-reduced if w does not contain a subword v such that v is also a subword of some $r \in \mathcal{R}_{ij}$ with r = vu, $||u|| \leq 4$.

An equality diagram for a pair of Artin-reduced words has the standard structure, consisting of "islands" in which each region "cuts through" the diagram from "top to bottom" as described precisely in the following lemma.

Lemma 2.11. Suppose Δ is a minimal diagram over (\ddagger) such that the boundary cycle of Δ corresponds to the relation $u =_G v$ where both u and v are Artin-reduced. Then Δ is a union of several (possibly none) linearly arranged disk components connected by (possibly degenerate) arcs, as shown in Figure 1. Moreover, each of these disk components

corresponds to an equality $u' =_G v'$, where u' is a subword of u and v' is a subword of v, and the component has the form shown in Figure 2 satisfying the following properties:

- (1) Each region Q_s corresponds to a relator from (\ddagger) .
- (2) Each interior edge p_s is labeled by a single syllable $a_{is}^{n_s}$ and has one endpoint on the "top" part of $\partial \Delta$ corresponding to u and its other endpoint on the "bottom" part of $\partial \Delta$ corresponding to v.
- (3) Each region Q_s has at least one boundary edge contained in the top part of $\partial \Delta$ and at least one boundary edge contained in the bottom part of $\partial \Delta$.




FIGURE 1. Equality diagram

Proof. By removing interior vertices of degree two and combining edge labels, we follow the usual convention that Δ has no interior vertices of degree two. Since u and v are Artin-reduced, it is clear that Δ is indeed a union of linearly arranged disk components, each corresponding to an equality diagram for $u' =_G v'$ for some subwords u' of u and v' of v, as shown in Figure 1. Also, each of the disk components is a minimal diagram over (\ddagger) .

Recall that by Proposition 2.8 in a minimal diagram over (‡) no two regions corresponding to relators from the same \mathcal{R}_{ij} have a common edge. Moreover, if $r \in \mathcal{R}_{ij}, r' \in \mathcal{R}_{il}$, where $l \neq j$, have a nontrivial common subword, this subword must be a power of a_i . Thus every interior edge of a minimal diagram over (‡) is labeled by some a_i^s .

Consider an individual disk component Δ_0 corresponding to an equality $u' =_G v'$, where u' is a subword of u and v' is a subword of v.

Let x_1 be the common initial vertex of the paths labeled u' and v' in the boundary of Δ_0 . Similarly, let x_2 be their common terminal




Figure 2. A disk component in an equality diagram




FIGURE 3. Equality diagram Δ_0 for $u' =_G v'$ in Lemma 2.11

vertex. If Δ_0 has only one region, Lemma 2.11 certainly holds. Thus we now suppose that Δ_0 has at least two regions. Since Δ_0 is minimal over presentation (‡), by Proposition 2.8 Δ_0 is a C(8)-diagram. Hence we can apply part Proposition 2.7 to Δ_0 . Since u' and v' are Artinreduced, case (b) of Proposition 2.7 cannot occur. Thus case (a) takes place and there are two distinct boundary regions Q and Q' in Δ_0 each with interior degree one and with a single syllable written on its interior edge, as shown in Figure 3. Denote those edges by q and q' accordingly.

It is clear that x_1 belongs to one of these regions and x_2 belongs to the other. Suppose x_1 is in Q and x_2 is in Q'. If Δ_0 consists of two or three regions, the statement of the lemma obviously holds. Otherwise let Δ' be obtained from Δ_0 by first removing the regions Q, Q', and, if their removal creates any vertices of degree one, then successively removing vertices of degree one and their incident edges until no such vertices remain. If Δ' has fewer than two regions then the statement of the lemma holds.

Otherwise the diagram Δ' contains at least two regions and we can apply Proposition 2.7 to it. Again, clearly case (b) of Proposition 2.7 cannot occur and case (a) takes place. Let S, S' be two distinct boundary regions in Δ' , each of interior degree one and thus with a single syllable written on the interior edge. Since u and v are weakly Artinreduced, one of these regions, say S, contains the edge q and the other region S' contains q'.

We claim that each of S, S' has an edge in common with both the top and the bottom portions of the boundary of Δ_0 . If this is not the case then either u' or v' contains a subword of r_{ij}^* missing at most two letters, contradicting the assumption that u' and v' are Artin-reduced. Hence we can remove S and S' from Δ' and repeatedly apply the same argument to the remaining diagram.

This yields the statement of Lemma 2.11.

3. Coned-off Cayley graphs and relative hyperbolicity

Recall that a geodesic path in a metric space (X,d) is an isometric embedding $\gamma:[a,b]\to X$, where $[a,b]\subseteq\mathbb{R}$. In this situation we call the set $\gamma([a,b])$ a geodesic segment from $x=\gamma(a)$ to $y=\gamma(b)$ in X and denote it by [x,y]. A metric space (X,d) is said to be geodesic if for any $x,y\in X$ there exists a geodesic segment [x,y] in X. The general notion of a "hyperbolic metric space" is, of course, due to Gromov [18]:

Definition 3.1 (Hyperbolic Metric Space). A geodesic metric space (X,d) is said to be hyperbolic if there is a number $\delta \geq 0$ such that for any three points $x,y,z\in X$, for any geodesic segments [x,y],[x,z] and [y,z] and for any point $p\in [x,y]$ there is a point $q\in [x,z]\cup [y,z]$ such that

$$d(p,q) \leq \delta.$$

We refer the reader to [1, 11, 20, 18] for the background material on hyperbolic spaces and hyperbolic groups.

We need to recall Farb's definitions of coned-off Cayley graphs and relative hyperbolicity [15]:

Definition 3.2 (Coned-off Cayley Graph). [15] Let G be a group with a finite generating set A and let H_1, \ldots, H_t be a family of subgroups of G. Let $\Gamma = \Gamma_A(G)$ be the Cayley graph of G with respect to A.

The coned-off Cayley graph $X = \widehat{\Gamma}_A(G; H_1, \dots, H_t)$ is obtained from Γ as follows: for each coset gH_i (where $g \in G$, $1 \le i \le t$) we add a new vertex $v(gH_i)$ to Γ and for every element $g' \in gH_i$ add an edge of length 1/2 from $v(gH_i)$ to g'.

The vertices $v(gH_i)$ are referred to as cone vertices and the edges from $v(gH_i)$ to elements of gH_i are called cone edges.

Definition 3.3 (Relative Hyperbolicity). [15] Let G be a finitely generated group and let H_1, \ldots, H_t be a family of subgroups of G. We say that G is relatively hyperbolic in the sense of Farb with respect to the t-tuple of subgroups (H_1, \ldots, H_t) if for some finite generating set A of G the coned-off Cayley graph $X = \widehat{\Gamma}_A(G; H_1, \ldots, H_t)$ is a hyperbolic metric space.

Recall that a finitely generated group G is hyperbolic if for some (and hence for any) finite generating set A of G the Cayley graph $\Gamma(G, A)$ is a hyperbolic metric space. Thus G is hyperbolic if and only if it is relatively hyperbolic with respect to the trivial subgroup $H = \{1\}$.

It is shown in [15] that G being relatively hyperbolic with respect to the given collection of subgroups does not depend on the choice of the generating set A.

4. Geodesics in the coned-off Cayley graph of an Artin group

Let G be given by presentation (†) where $m_{ij} \geq 7$ for all $i \neq j$. Let $\Gamma = \Gamma_A(G)$ where $A = \{a_1, \ldots, a_n\}$. Let $X = \widehat{\Gamma}_A(G; \{G_{ij} | m_{ij} < \infty\})$ be the coned-off Cayley graph of G. If e is an edge of a graph, we will denote by o(e) the initial vertex of e and by t(e) the terminal vertex of e. These notations and conventions will be fixed for the remainder of the paper.

Convention 4.1. Let $g \in G$ be an arbitrary element and let α be a geodesic in X from $1 \in G$ to g. The goal of this section is to show that α is 4-close in X to a path γ from 1 to g in Γ whose label is strongly Artin-reduced. As an intermediate step in constructing this path γ we will first define an auxiliary path β . Unless specified otherwise, we shall fix g and α for the remainder of this section.

4.1. Construction of the path β . The path α can be subdivided as a concatenation of paths

$$\alpha = \alpha_1 \dots \alpha_t$$

where $t \geq 1$ and where each α_k is either a path in Γ from a vertex of Γ to a vertex of Γ , in which case α_k is referred to as a Γ -block, or $\alpha_k = e_k e_k'$, where e_k and e_k' are cone-edges separated by a cone-vertex, in which case α_k is referred to as a cone-block. Thus $t(e_k) = o(e_k')$ is a cone-vertex of X and $t(e_k') = o(e_k)g_k$ for some $g_k \in G_{ij} - \{1\}$. Moreover, we can assume that the Γ -blocks are chosen maximally, so that no two such blocks are consecutive in α . For each cone-block α_k choose a freely reduced word v_k in G_{ij} of the smallest possible syllable length representing g_k . Note that if g_k belongs to two distinct G_{ij} -subgroups and thus is a power of the generator then the pair $\{i, j\}$ may not be uniquely defined by g_k , but in that case the word v_k of syllable length one will be uniquely determined by g_k .

Remark 4.2. The choice of the word v_k for the cone-block α_k as a word in the corresponding G_{ij} which is "geodesic" with respect to syllable length is a crucial feature of our argument.

If α_k is a Γ -block then the label v_k of α_k is an A-geodesic word.

Notation 4.3. The word

$$v = v_1 \dots v_t$$

certainly represents the element g. We denote the path in Γ from 1 to g with label v by β . Thus

$$\beta = \beta_1 \dots \beta_t$$

where β_k is the path in Γ with label v_k from the initial point of α_k to the terminal point of α_k .

The following lemma is a straightforward consequence of the definitions:

Lemma 4.4. Let α and β be as above. Then:

- (1) The label v_k of each Γ -block α_k is a Γ -geodesic word.
- (2) If v' is a subword of the label v_k of some Γ -block such that v' is a word in some G_{ij} then $|v'| \leq 1$.
- (3) Suppose that α_{k+1} is a Γ -block and that α_k is a cone-block with $v_k \in G_{ij}$. Then the first syllable of v_{k+1} is not a power of either a_i or a_j .

- (4) Suppose that α_k is a Γ -block and that α_{k+1} is a cone-block with $v_{k+1} \in G_{ij}$. Then the last syllable of v_k is not a power of either a_i or a_j .
- (5) If α_k, α_{k+1} are cone-blocks and $v_k \in G_{ij}, v_{k+1} \in G_{st}$ then $\{i, j\} \neq \{s, t\}.$
- (6) The paths α and β are 2-Hausdorff close in X.

Lemma 4.5. Suppose that $\alpha_{k-1}, \alpha_k, \alpha_{k+1}$ are cone-blocks with $v_{k-1} \in G_{ij}$, $v_k \in G_{js}$, $v_{k+1} \in G_{sq}$. If the first syllable of v_k is a power of a_j and the last syllable of v_k is a power of a_s then $||v_k|| \geq 3$.

Proof. If $||v_k|| = 2$ then we can replace the path $\alpha_{k-1}\alpha_k\alpha_{k+1}$ of length 3 in X by a path of length 2 in X consisting of two pairs of cone-edges: the first corresponding to $G_{i,j}$ and the second corresponding to G_{sq} . This contradicts the assumption that α is a geodesic in X.

4.2. Construction of the path γ . Note that the word v is not necessarily reduced. Indeed, it is possible that α_k, α_{k+1} are two consecutive cone blocks such that the last syllable of v_k and the first syllable of v_{k+1} are powers of the same a_i . (The above lemmas imply that this is the only way in which v may fail to be freely reduced). We will modify v and β to rectify this problem by "condensing syllables" from left to right as follows.

We define a sequence of words u_1, \ldots, u_t inductively.

Put $u_1 := v_1$. Suppose $1 \le k < t$ and u_1, \ldots, u_k are already defined. We have two cases to consider:

(a) Suppose α_k is a cone-block with $v_k \in G_{ij}$.

Let $u_k = z_k s_k$ where s_k is the last syllable of u_k .

If α_{k+1} is also a cone-block, let $v_{k+1} = s_{k+1}y_{k+1}$ where s_{k+1} is the first syllable of v_{k+1} . If the last syllable of v_k and the first syllable of v_{k+1} are not powers of the same generator, then let $u_{k+1} = v_{k+1}$. If the last syllable of v_k and the first syllable of v_{k+1} are powers of the same generator, then let u'_k be the reduced form of $u_k s_{k+1}$ and redefine u_k to be u'_k . Set $u_{k+1} := y_{k+1}$.

If α_{k+1} is a Γ -block, put $u_{k+1} := v_{k+1}$.

(b) If α_k is a Γ -block, put $u_{k+1} := v_{k+1}$.

Notation 4.6. We set

$$(\spadesuit) \qquad \qquad u := u_1 \dots u_t,$$

and denote by γ the path in Γ from 1 to g with label u. Also, denote by γ_k the subpath of γ corresponding to u_k , so that

$$\gamma = \gamma_1 \dots \gamma_t$$
.

We summarize the relevant properties of u and γ :

Lemma 4.7. The following hold:

- (1) The word u as in (\spadesuit) is freely reduced.
- (2) Suppose α_k is a cone-block with $u_k \in G_{ij}$. Then for any word w in G_{ij} representing the same element as u_k we have $||u_k|| \le ||w|| + 1$.
- (3) For any k < t we have $||u_k u_{k+1}|| = ||u_k|| + ||u_{k+1}||$.
- (4) For any Γ -block α_k we have $\gamma_k = \beta_k = \alpha_k$ and $u_k = v_k$.
- (5) The paths γ and β are 2-Hausdorff close in X.

Lemma 4.8. Let w be a reduced word in some G_{ij} , $m_{ij} < \infty$ such that $||w|| \ge 4$. Suppose w is a subword of u and let w' be obtained from w by removing the first and the last syllables of w. Then w' is a subword of some u_k such that α_k is a cone-block of α .

Proof. Suppose first that the occurrence of w in u overlaps some $u_k = v_k$ corresponding to a Γ-block α_k of α . Then some syllable s of w is a subword of this u_k . Recall that no two Γ-blocks are adjacent in α . Part 2 of Lemma 4.4 implies that neither of the syllables preceding or following s of w can overlap u_k . Let $u_k = asb$. Among the words a, b choose the one of largest syllable length. By symmetry we may assume that $||a|| \ge ||b||$. Hence $||a|| \ge 2$ since $||w|| \ge 4$. Since ||a|| > 0 then α_{k-1} is a cone-block. Also, since $||a|| \ge 2$ and w is a word in G_{ij} , we conclude that both s (which is the first syllable of u_k) and u_{k-1} are words in G_{ij} (this is true even if $||u_{k-1}|| = 1$). This is impossible by Lemma 4.4.

Thus w does not overlap any u_k corresponding to Γ -blocks α_k .

Assume now that there is some u_k contained in w such that w also overlaps u_{k-1} and u_{k+1} . Each of α_{k-1} , α_k and α_{k+1} is a cone-block. If $||u_k|| = 1$ then either the part of w following u_k or the part of w preceding u_k has syllable length at least two. Since w is a word in G_{ij} this yield a contradiction with part 5 of Lemma 4.4 and the definition of u. Hence $||u_k|| \geq 2$. If either the either the part of w following u_k or the part of w preceding u_k has syllable length at least two, then either both v_{k-1}, v_k or both v_k, v_{k+1} belong to G_{ij} . Again, this is impossible by part 5 of Lemma 4.4. The statement of Lemma 4.8 now follows.

Suppose now that there is no u_k contained in w such that w also overlaps u_{k-1} and u_{k+1} . Thus w overlaps at most two of the words u_k . If w is contained in a single u_k , the statement of Lemma 4.8 obviously holds. Assume now that w is a subword of $u_k u_{k+1}$ and that w overlaps both of these words. If both overlaps have syllable length at least two, then $g_k, g_{k+1} \in G_{ij}$, contrary to part 5 of Lemma 4.4. If one of

the overlaps has syllable length one, then all but the first or the last syllable of w is contained in either u_k or u_{k+1} and the statement of Lemma 4.8 holds.

Proposition 4.9. The word u is strongly Artin reduced and the paths α and γ are 4-Hausdorff close in X.

Proof. The paths α and γ are 4-close since each is 2-close to the path β by part 6 of Lemma 4.4 and by part 5 of Lemma 4.7.

We now show that u is strongly Artin reduced. If not, there is a nontrivial relator $r \in \mathcal{R}_{ij}$ for some $m_{ij} < \infty$ and a subword w of u such that r = wy and $||y|| \le 4$. Thus $w = y^{-1}$ in G.

It is now that we use the assumption that all $m_{ij} \geq 7$. By Proposition 2.3 this condition implies that $||r|| \geq 14$ and hence $||w|| \geq 10$. Let $w = s_1 w' s_2$ where s_1, s_2 are respectively the first and the last syllables of w. Then $||w'|| \geq 8$ and by Lemma 4.8 w' is a subword of some u_k such that α_k is a cone-block of α .

Moreover, if we write u_k as $u_k = zw'z'$ then $||u_k|| = ||z|| + ||w'|| + ||z'||$. Using the relator r, we have

$$w' =_G s_1^{-1} y^{-1} s_2^{-1}$$
 and hence $u_k =_G z s_1^{-1} y^{-1} s_2^{-1} z'$

But

$$||zs_1^{-1}y^{-1}s_2^{-1}z'|| \leq ||z|| + 1 + 4 + 1 + ||z'|| < ||z|| + ||w'|| - 1 + ||z'|| = ||u_k|| - 1,$$

where the last inequality holds since $||w'|| \ge 8$. Hence there exists a word representing the same element as u_k but with syllable length less than $||u_k|| - 1$.

This contradicts part 2 of Lemma 4.7.

5. Proof of the main result

We recall the following useful fact due to Papasoglu [26]:

Proposition 5.1 (Thin bigons criterion). Let C be a connected graph with the simplicial metric d. Then C is hyperbolic if and only if there is some $\delta > 0$ such that for any points $x, y \in C$ (possibly inside edges) any two geodesic paths from x to y in C are δ -Hausdorff close.

Remark 5.2. The above result was stated in [26] only for Cayley graphs of finitely generated groups. However, it is easy to see that Papasoglu's proof [26] does not use the Cayley graph assumption and works for any connected graph with the simplicial metric. This was noted, for example, in [24].

We are now ready to prove Theorem A.

Theorem A. Let G be an Artin group given by presentation (\dagger) . Suppose that for all i < j we have $m_{ij} \ge 7$.

Then G is relatively hyperbolic in the sense of Farb with respect to the collection of subgroups $\{G_{ij}|m_{ij}<\infty\}$.

Proof. Put $A = \{a_1, \ldots, a_n\}$. Let $\Gamma = \Gamma_A(G)$ and let

$$X = \widehat{\Gamma}_A(G; \{G_{ij} | m_{ij} < \infty\})$$

be the coned-off Cayley graph of G. We need to show that X is hyperbolic.

Note that if we barycentrically subdivide each of the Γ -edges of X to obtain the graph X', then twice the metric of X coincides with the simplicial metric for X'. Hence to establish Theorem A by using Papasoglu's criterion it suffices to show that geodesic bigons in X (with endpoints possibly inside edges) are uniformly thin. Moreover, the definition of X implies that it is enough to prove the following:

Claim. There exists a constant K > 0 with the following property. Let α_1, α_2 be geodesics in X from $h_1 \in G$ to $g_1 \in G$ and from $h_2 \in G$ to $g_2 \in G$ respectively such that:

- (1) Either $h_1^{-1}h_2 \in G_{ij}$ for some $m_{ij} < \infty$ or $d_{\Gamma}(h_1, h_2) \le 1$. (2) Either $g_2^{-1}g_1 \in G_{sl}$ for some $m_{sl} < \infty$ or $d_{\Gamma}(g_1, g_2) \le 1$.

Then α_1 and α_2 are K-Hausdorff close in w.

To verify the Claim, let $a, b \in G$ be such that $h_1 = h_2 a$ and $g_1 = g_2 b$. By Proposition 4.9 for l = 1, 2 there is a path γ_l in Γ from h_l to g_l such that the label U_l of γ_l is a strongly Artin-reduced word and such that the paths γ_l and α_l are 4-close in X, as shown in Figure 4.

If $d_{\Gamma}(h_1, h_2) \leq 1$ then a is either trivial or a single letter of $A \cup A^{-1}$. Let W_1 be the freely reduced form of the word aU_1 . Since U_1 is strongly Artin-reduced, the word W_1 is Artin-reduced.

Assume now that $d_{\Gamma}(h_1, h_2) \geq 2$, so that $a \in G_{ij}$ for some $m_{ij} < \infty$. If a belongs to the cyclic subgroup generated by one of the letters of A (in which case G_{ij} may not be uniquely determined by a), say $a = a_i^q$, we let W_1 be the freely reduced from of the word $a_i^q U_1$. Again, the word W_1 is Artin-reduced because U_1 is strongly Artin-reduced. Assume now that a is not a power of the generator, so that $a \in G_{ij}$ for a unique pair $\{i,j\}$. Let $U_1 = Y_1Z_1$ where Y_1 is the maximal initial segment of U_1 which is a word in G_{ij} . Let Y'_1 be the reduced word of smallest possible syllable length in G_{ij} representing the element aY_1 . Then Y'_1 and Z_1 are strongly Artin-reduced and the word $W_1 = Y_1'Z_1$ is Artin-reduced.

In either of the above cases, using the cone-vertex $v(h_1G_{ij})$ it is easy to see that the path γ_1 is 2-close in X to the path λ_1 in Γ from h_2 to g_1 labeled by W_1 .




FIGURE 4. Thinness of bigons

Similarly, by considering the product U_2b we can find an Artinreduced word W_2 and a path λ_2 from h_2 to g_1 in Γ with label W_2 such that γ_2 and λ_2 are 2-close in X.

Now consider an equality diagram for the equality $W_1 =_G W_2$ of smallest area over the infinite presentation (‡) of G. Applying Lemma 2.11 to this diagram and using cone vertices we see that λ_1 and λ_2 are 2-close in X. Adding the distances between the paths considered in Figure 4 we see that α_1 and α_2 are 14-close in X and the Claim is verified. \square

Remark 5.3. The value K=14 obtained in the above proof of Theorem A using Papasolglu's criterion does not depend on the particular group G. Thus all the coned-off Cayley graphs X of the groups G covered by Theorem A are all δ -hyperbolic for some $\delta > 0$ which is independent of X and hence of the choice of G. So the relative hyperbolicity of G with respect to G_{ij} 's is in a sense uniform over the class of Artin groups covered by the Theorem.

References

 J. Alonso, T. Brady, D. Cooper, V. Ferlini, M. Lustig, M. Mihalik, M. Shapiro and H. Short, *Notes on hyperbolic groups*, In: "Group theory from a geometrical viewpoint", Proceedings of the workshop held in Trieste, World Scientific Publishing Co., 1991

- [2] K. Appel and P. Schupp, Artin groups and infinite Coxeter groups, Invent. Math. 72 (1983), no. 2, 201–220
- [3] J. Altobelli, and R. Charney, A geometric rational form for Artin groups of FC type. Geom. Dedicata 79 (2000), no. 3, 277–289
- [4] N. Brady, and J. Crisp, Two-dimensional Artin groups with CAT(0) dimension three. Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part I (Haifa, 2000). Geom. Dedicata 94 (2002), 185–214
- [5] B. H. Bowditch, *Relatively hyperbolic groups*, preprint, University of Southampton, 1997
- [6] B. H. Bowditch, A topological characterisation of hyperbolic groups. J. Amer. Math. Soc. 11 (1998), no. 3, 643–667
- [7] B. H. Bowditch, Treelike structures arising from continua and convergence groups. Mem. Amer. Math. Soc. 139 (1999), no. 662
- [8] B. H. Bowditch, *Peripheral splittings of groups*, Trans. Amer. Math. Soc. **353** (2001), no. 10, 4057–4082
- [9] J. Brock and B. Farb, Curvature and rank of Teichmüller space, preprint, 2001
- [10] R. Charney, and M. Davis, The $K(\pi,1)$ -problem for hyperplane complements associated to infinite reflection groups. J. Amer. Math. Soc. 8 (1995), no. 3, 597–627
- [11] M. Coornaert, T. Delzant, and A. Papadopoulos, Géométrie et théorie des groupes. Les groupes hyperboliques de Gromov. Lecture Notes in Mathematics, 1441; Springer-Verlag, Berlin, 1990
- [12] F. Dahmani, Combination of Convergence Groups, preprint, 2002
- [13] F. Dahmani, Classifying Space and Boundary for Relatively Hyperbolic Groups, Proc. London Math. Soc. 86 (2003), no. 3, 666–684
- [14] D. Epstein, J. Cannon, D. Holt, S. Levy, M. Paterson, W. Thurston, Word Processing in Groups, Jones and Bartlett, Boston, 1992
- [15] B. Farb, Relatively hyperbolic groups, Geom. Funct. Anal. 8 (1998), 810–840
- [16] F. Gautero and M. Lustig, Relative hyperbolicity of (one-ended hyperbolic)-by-cyclic groups. preprint, 2003
- [17] B. Goldfarb, Novikov conjectures and relative hyperbolicity. Math. Scand. **85** (1999), no. 2, 169–183
- [18] M. Gromov, *Hyperbolic Groups*, in "Essays in Group Theory (G.M.Gersten, editor)", MSRI publ. **8**, 1987, 75–263
- [19] M. Gromov, Asymptotic invariants of infinite groups. Geometric group theory, Vol. 2 (Sussex, 1991), 1–295, London Math. Soc. Lecture Note Ser., 182, Cambridge Univ. Press, Cambridge, 1993
- [20] E. Ghys and P. de la Harpe (editors), Sur les groupes hyperboliques d'aprés Mikhael Gromov, Birkhäuser, Progress in Mathematics series, vol. 83, 1990.
- [21] C. Hruska, Nonpositively curved 2-complexes with isolated flats, preprint, 2003
- [22] R. Lyndon and P. Schupp, Combinatorial Group Theory, Springer-Verlag, 1977
- [23] H. Masur and Y. Minsky, Geometry of the complex of curves. I. Hyperbolicity, Invent. Math. 138 (1999), no. 1, 103–149

- [24] W. Neumann and M. Shapiro, A Short Course in Geometric Group Theory, Notes for the ANU Workshop January/February 1996, www.math.columbia.edu/~neumann/preprints/canberra.ps
- [25] G. Niblo, and L. Reeves, The geometry of cube complexes and the complexity of their fundamental groups. Topology 37 (1998), no. 3, 621–633
- [26] P. Papasoglu, Strongly geodesically automatic groups are hyperbolic, Invent. Math. 121 (1995), no. 2, 323–334
- [27] L. Paris, Parabolic subgroups of Artin groups. J. Algebra 196 (1997), no. 2, 369–399
- [28] D. Peifer, Artin groups of extra large type are biautomatic. J. Pure Appl. Algebra 110 (1996), 15-56
- [29] A. Szczepański, Relatively hyperbolic groups. Michigan Math. J. 45 (1998), no. 3, 611–618
- [30] A. Szczepański, Examples of Relatively Hyperbolic Groups, preprint, 2002 Geometrica Dedicata 93 (2002), no. 1, 139–142
- [31] A. Yaman, A topological characterisation of relatively hyperbolic groups, preprint, University of Southampton 2001

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, 1409 WEST GREEN STREET, URBANA, IL 61801, USA

E-mail address: kapovich@math.uiuc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, 1409 WEST GREEN STREET, URBANA, IL 61801, USA

E-mail address: schupp@math.uiuc.edu