Periods of Abelian Varieties

J.S. Milne*

February 27, 2003. Submitted version.

Abstract

We prove various characterizations of the period torsor of abelian varieties.

Contents	
----------	--

Introduction		1
1	Preliminaries	4
2	Periods of abelian varieties with complex multiplication	13
3	Periods of abelian varieties	22
References		33

Introduction

For an abelian variety A over \mathbb{Q} , $H^1(A(\mathbb{C}), \mathbb{C})$ has two \mathbb{Q} -structures, that provided by singular cohomology $H^1(A(\mathbb{C}), \mathbb{Q})$ and that provided by de Rham cohomology $\mathbb{H}^1(A_{\text{Zar}}, \Omega_{A/k}^{\bullet})$. The *periods* of A are the coefficients of the transition matrix from a basis for one \mathbb{Q} -space to a basis for the other. It is known (Deligne 1982) that Hodge classes on A impose algebraic relations on the periods, and it is conjectured that these are the only such relations. Thus, there appears to be no hope of obtaining an explicit description of the periods, but one may still hope to characterize some of the objects attached to them. The singular and de Rham cohomologies define fibre functors on the category of motives based on the abelian varieties over \mathbb{Q} , and the difference of these functors is measured by the period torsor P^{AV} . In this paper, we obtain various characterizations of P^{AV} . Beyond its intrinsic interest, the period torsor controls the rationality of automorphic vector bundles, and therefore of holomorphic automorphic forms (see Milne 1988; Milne 1990, III 4).

^{*}Partially supported by the National Science Foundation.

The first problem one runs into is that P^{AV} is a torsor for an affine group scheme G over \mathbb{Q} for the flat (more specifically, the f.p.q.c) topology. Such torsors are classified by the flat cohomology group $H^1(\mathbb{Q}, G)$ rather than a more familiar Galois cohomology group. For an algebraic quotient G_n of G, the two cohomology groups coincide. In §1 we prove that the canonical map

$$H^1(\mathbb{Q},G) \to \lim H^1(\mathbb{Q},G_n)$$

is surjective, and that the fibre of the map containing the class of a G-torsor P is

$$\underbrace{\lim}^{1}(P \wedge^{G} G_{n})(\mathbb{Q})$$

(nonabelian higher inverse limit). The limits are over the set of algebraic quotients G_n of G. We show, moreover, that the full group $H^1(\mathbb{Q}, G)$ can be interpreted as a Galois cohomology group, but it is Galois cohomology defined using cochains that are continuous relative to the inverse limit topology on $\lim_{i \to G} G_n(\mathbb{Q}^{al})$ (discrete topology on each $G_n(\mathbb{Q}^{al})$). It is important to note that these results depend crucially on the fact that the algebraic quotients of G form a countable set — I do not even know how to define a nonabelian higher inverse limit except for countable coefficient sets. The remainder of §1 reviews results, more-or-less known, concerning nonabelian higher inverse limits and the classification of morphisms of torsors.

In §2, we take up the problem of characterizing P^{CM} , the period torsor for the category of motives based on abelian varieties of (potential) CM-type over \mathbb{Q} . Since, as Deligne has pointed out, the period torsor P^{Art} attached to the subcategory of Artin motives can be explicitly described, it is natural rather to consider the "relative" problem of characterizing the morphism $P^{CM} \rightarrow P^{Art}$ of torsors. Objects of this type are classified by the flat cohomology group $H^1(\mathbb{Q}, fS)$ where fS is a certain twist of the Serre group. As the Serre group is commutative, the main result of §1 simplifies to an exact sequence

$$0 \to \varprojlim{}^{1}{}_{f}S_{n}(\mathbb{Q}) \to H^{1}(\mathbb{Q}, {}_{f}S) \to \varprojlim{} H^{1}(\mathbb{Q}, {}_{f}S_{n}) \to 0.$$

Blasius (unpublished) showed that $\lim_{t \to T} H^1(\mathbb{Q}, {}_fS_n)$ satisfies a Hasse principle, and Wintenberger (1990) showed that $\lim_{t \to T} H^1(\mathbb{Q}, {}_fS_n) = 0$. If $\lim_{t \to T} {}^1{}_fS_n(\mathbb{Q})$ were also zero, then $P^{\mathsf{CM}} \to P^{\mathsf{Art}}$ would be characterized up to isomorphism (inducing the identity on P^{Art}) by its cohomology class. Alas, it is not zero — in fact, we show that $\lim_{t \to T} {}^1{}_fS_n(\mathbb{Q})$ is uncountable. Our proof of this uses an old theorem of Scholz and Reichardt on the embedding problem for Galois groups of number fields. It would be interesting to have more information on $\lim_{t \to T} {}^1{}_fS_n(\mathbb{Q})$.

In §3, we take up the problem of chacterizing P^{AV} . Again, it is more natural to consider the relative problem of characterizing $P^{AV} \rightarrow P^{CM}$. Among other results, we prove that the isomorphism class of $P^{AV} \rightarrow P^{CM}$ is uniquely determined by its classes over \mathbb{Q}_l $(l = 2, 3, ..., \infty)$, about which a great deal is known.

Blasius and Borovoi (1999) study the problem of characterizing $P^{\mathcal{H}}$ where $\mathcal{H} \subsetneq AV$ is the category of motives based on a certain class of abelian varieties over \mathbb{Q} whose Mumford-Tate groups have simply connected derived group. Unfortunately, they make the (false!) assumption that the flat cohomology groups coincide with the inverse-limit Galois cohomology groups, i.e., effectively they set all \lim^{1} s equal to zero.¹ If this were their

¹This alone accounts for most of the relative brevity of their paper.

only error, their main Theorem 1.5 would hold for suitably large finite sets of abelian varieties satisfying their condition, but they also misidentify the cohomology class that must be proved trivial². Nevertheless, by combining (1.14) of this paper with their arguments, one obtains the following theorem (3.25): the class of $P^{\mathcal{H}} \to P^{\mathsf{CM}}$ is uniquely determined by its class over \mathbb{R} . This observation began my work on this paper.

Notations and conventions

"Variety" means geometrically reduced scheme of finite type over a field. Semisimple algebraic groups are connected and "simple" for an algebraic group means "noncommutative and having no proper closed connected normal subgroup $\neq 1$ ". The identity component of a group scheme G over a field is denoted by G° . For a connected (pro)reductive group G over a field, ZG is the centre of G, G^{ad} is the adjoint group G/ZG of G, G^{der} is the derived group of G, and G^{ab} is the largest commutative quotient G/G^{der} of G. The universal covering of a semisimple group G is denoted $\tilde{G} \to G$.

The algebraic closure of \mathbb{Q} in \mathbb{C} is denoted by \mathbb{Q}^{al} , and (except in §1) $\Gamma = \operatorname{Gal}(\mathbb{Q}^{al}/\mathbb{Q})$. We set $\operatorname{Gal}(\mathbb{C}/\mathbb{R}) = \{1, \iota\}$. A CM-field is any field E algebraic over \mathbb{Q} admitting a non-trivial involution ι_E such that $\iota \circ \rho = \rho \circ \iota_E$ for all $\rho \colon E \to \mathbb{C}$.

All categories of motives will be defined using absolute Hodge classes as the correspondences (Deligne 1982a; Deligne and Milne 1982, §6).

We sometimes use [x] to denote an equivalence or isomorphism class containing x. The notation $X \approx Y$ means that X and Y are isomorphic, and $X \cong Y$ means that X and Y are canonically isomorphic (or that a particular isomorphism is given).

²With their notations, in order to prove their theorem, they would have to show in **5.1** that the class of $\mathcal{P} \to \mathcal{P}^{CM}$ in $H^1(\mathbb{Q}, (\mathcal{G}_{DR}^\circ)^{der})$ is trivial (see (1.10) below). In fact, they prove only the weaker statement that the image of the class in $H^1(\mathbb{Q}, \mathcal{G}_{DR}^\circ)$ is trivial, which, in fact, is all their hypotheses imply, even when one ignores the \lim^1 terms.

1 Preliminaries

Inverse limits

For an inverse system of groups indexed by (\mathbb{N}, \leq) ,

$$(A_n, u_n)_{n \in \mathbb{N}} = (A_0 \leftarrow \dots \leftarrow A_{n-1} \xleftarrow{u_n} A_n \leftarrow \dots),$$

define $\lim_{n \to \infty} A_n$ to be the set of orbits for the left action of the group $\prod_n A_n$ on the set $\prod_n A_n$.

$$\prod_n A_n \times \prod_n A_n \to \prod_n A_n$$

(..., a_n, ...) \mapsto (..., a_n \cdot x_n \cdot u_{n+1}(a_{n+1})^{-1}, ...).

This is a set, pointed by the orbit of $\underline{1} = (1, 1, ...)$. Note that

$$\varprojlim A_n = \{a \in \prod A_n \mid a \cdot \underline{1} = \underline{1}\}$$

Let $(A_n)_{n\in\mathbb{N}}\to (B_n)_{n\in\mathbb{N}}$ be an inverse systems of injective homomorphisms. From

$$0 \to (A_n)_{n \in \mathbb{N}} \to (B_n)_{n \in \mathbb{N}} \to (B_n/A_n)_{n \in \mathbb{N}} \to 0$$

we obtain an exact sequence

$$1 \to \varprojlim A_n \to \varprojlim B_n \to \varprojlim (B_n/A_n) \to \varprojlim^1 A_n \to \varprojlim^1 B_n \tag{1}$$

of groups and pointed sets. Exactness at $\varprojlim(B_n/A_n)$ means that the fibres of $\varprojlim(B_n/A_n) \rightarrow \varprojlim^1 A_n$ are the orbits for the natural action of $\varprojlim B_n$ on $\varprojlim(B_n/A_n)$. When each A_n is normal in B_n , so that $C_n =_{df} B_n/A_n$ is a group, (1) can be extended to an exact sequence

$$1 \to \varprojlim A_n \to \varprojlim B_n \to \varprojlim C_n \to \varprojlim^1 A_n \to \varprojlim^1 B_n \to \varprojlim^1 C_n \to 1.$$
 (2)

Exactness at $\varprojlim^1 A_n$ means that the fibres of $\varprojlim^1 A_n \to \varprojlim^1 B_n$ are the orbits for the natural action of $\varprojlim C_n$ on $\varprojlim^1 A_n$.

Recall that an inverse system $(X_n)_{n \in \mathbb{N}}$ of sets (or groups) is said to satisfy the condition (ML) if, for each m, the decreasing chain in X_m of the images of the X_n for $n \ge m$ is eventually constant.

Proposition 1.1. Let $(A_n, u_n)_{n \in \mathbb{N}}$ be an inverse system of groups.

- (a) If (A_n, u_n) satisfies (ML), then $\lim^1 A_n = 0$.
- (b) If the A_n are countable and $(A_n, u_n)_{n \in \mathbb{N}}$ fails (ML), then $\underline{\lim}^1 A_n$ is uncountable.

Proof. (a) The action³

$$(\ldots, a_n, \ldots) \cdot (\ldots, x_n, \ldots) = (\ldots, a_n \cdot x_n \cdot (ua_{n+1})^{-1}, \ldots)$$

³We usually omit the subscript on transition maps.

of the group $G_{N+1} =_{df} \prod_{0 \le n \le N+1} A_n$ on the set $S_N =_{df} \prod_{0 \le n \le N} A_n$ is transitive, and the projection $(a_n)_n \mapsto a_{N+1}$ gives an isomorphism from the stabilizer of any $x \in S_N$ onto A_{N+1} . Let $x, y \in \prod_{n \in \mathbb{N}} A_n$, and let

$$P_N = \left\{ g \in G_{N+1} \mid gx^N = y^N \right\}.$$

where x^N and y^N are the images of x and y in S_N . We have to show that $\varprojlim P_N$ is nonempty. The observations in the first sentence show that P_N is nonempty and that A_{N+1} acts simply transitively on it. It follows that the inverse system (P_N) satisfies (ML). Let $Q_N = \bigcap_i \operatorname{Im}(P_{N+i} \to P_N)$. Then each Q_N is nonempty, and $(Q_N)_{N \in \mathbb{N}}$ is an inverse system with surjective transition maps. Hence, $\varprojlim Q_N$ is (obviously) nonempty, and any element of it is an element of $\lim P_N$.

(b) If (A_n) fails (ML), then there exists an m such that infinitely many of the groups

$$B_i \stackrel{\text{df}}{=} \operatorname{Im}(A_{m+i} \to A_m)$$

are distinct. As

$$\varprojlim_i {}^1A_{m+i} \to \varprojlim_i {}^1B_i$$

is surjective (see (2)), it suffices to show that $\lim_{i \to \infty} B_i$ is uncountable. This is accomplished by the next lemma (applied with $A = A_m$).

Lemma 1.2. Let $\cdots \supset A_n \supset A_{n+1} \supset \cdots$ be a sequence of subgroups of a countable group *A*. If infinitely many of the A_n are distinct, then $\lim^{1} A_n$ is uncountable.

Proof. From (1) applied to the inverse system $(A_n \hookrightarrow A)_{n \in \mathbb{N}}$, we obtain a bijection

$$A \setminus (\varprojlim A/A_n) \to \varprojlim^1 A_n.$$

As a map of sets, $A/A_{n+1} \rightarrow A/A_n$ is isomorphic to the projection map

$$A/A_n \times A_n/A_{n+1} \to A/A_n,$$

and so $\lim A/A_n \approx \prod A_n/A_{n+1}$ (as sets), which is uncountable.

Remark 1.3. The above statements apply to inverse systems indexed by any directed set *I* containing an infinite countable cofinal set, because such an *I* will also contain a cofinal set isomorphic to (\mathbb{N}, \leq) .

Notes. The definition of $\lim_{x \to 1}^{1}$ for nonabelian groups and the sequence (2) can be found in Bousfield and Kan 1972, IX §2. In the commutative case, statement (a) of Proposition 1.1 is proved in Atiyah 1961 and statement (b) in Gray 1966.

Torsors

Let E be a category with finite fibred products (in particular, a final object S) endowed with a topology in the sense of Grothendieck (see Bucur and Deleanu 1968, Chapter 2). Thus, E is a site. By "torsor" we mean "right torsor".

1 PRELIMINARIES

1.4. For a sheaf of groups A on E, a right A-sheaf X, and a left A-sheaf Y, $X \wedge^A Y$ denotes the contracted product of X and Y, i.e., the quotient sheaf of $X \times Y$ by the diagonal action of A, $(x, y)a = (xa, a^{-1}y)$. When $A \to B$ is a homomorphism of sheaves of groups, $X \wedge^A B$ is the B-sheaf obtained from X by extension of the structure group. In this last case, if X is an A-torsor, then $X \wedge^A B$ is a B-torsor.

1.5. For an *A*-torsor *P* and a left *A*-sheaf *X*, define

$${}^{P}X = P \wedge^{A} X.$$

When X is a sheaf of groups and A acts by group homomorphisms, ${}^{P}X$ is a sheaf of groups. For example, when we let A act on itself by inner automorphisms, ${}^{P}A$ is the inner form of A defined by P. There is a natural left action of ${}^{P}A$ on P, which makes P into a left ${}^{P}A$ -torsor and induces an isomorphism

$${}^{P}A \to \mathcal{A}ut_{A}(P).$$
 (3)

Let P(A) denote the category of A-torsors and $H^1(S, A)$ the set of isomorphism classes of objects in P(A) (pointed by the class of the trivial torsor A_A).

1.6. Let $v: B \to C$ be a homomorphism of sheaves of groups on E, and let Q be a C-torsor. Define $P(B \to C; Q)$ to be the category whose objects are the v-morphisms of torsors $P \to Q$ and whose morphisms $Hom(P \to Q, P' \to Q)$ are the B-morphisms $P \to P'$ giving a commutative triangle. Let $H^1(S, B \to C; Q)$ denote the set of isomorphism classes in $P(B \to C; Q)$. When $Q = C_C$, we drop it from the notation; then $H^1(S, B \to C)$ is pointed by the class of $B_B \to C_C$. The category $P(B \to 0)$ is canonically equivalent with the category of B-torsors, and so

$$H^1(S, B \to 0) \cong H^1(S, B).$$

Let $A = \text{Ker}(B \to C)$. Then A is stable under the action of B on itself by inner automorphisms, and for any object $P \to Q$ of $P(B \to C; Q)$,

$$\mathcal{A}ut(P \to Q) \cong P \wedge^B A.$$

1.7. Let $v: B \to C$ be a surjective homomorphism with kernel A. To give a B-torsor P with vP trivialised by $e \in (vP)(S)$ amounts to giving the A-torsor $f^{-1}(e)$: the natural functor

$$\mathsf{P}(A \to 0) \to \mathsf{P}(B \to C)$$

is an equivalence.

1.8. Let $v: B \to C$ be a homomorphism of sheaves of groups on E. A *B*-torsor *P* allows us to twist *v*:

$${}^{P}v \colon {}^{P}B \to {}^{P}C, \quad {}^{P}C \stackrel{\mathrm{df}}{=} P \wedge {}^{B}C.$$

Here a local section b of B acts on C by $c \mapsto (vb)c(vb)^{-1}$. Let Q = vP. Then ${}^{P}C \cong {}^{Q}C$.

1.9. Let $v: B \to C$ be a homomorphism of sheaves of groups on E. A *B*-torsor *P* can be regarded as a $({}^{P}B, B)$ -bitorsor (see (3)). There is a functor

$$\mathsf{P}(^{P}B \to {}^{Q}C) \to \mathsf{P}(B \to C;Q) \tag{4}$$

sending $P' \to Q'$ to $P' \wedge^{P_B} P \to Q' \wedge^{Q_B} Q$. In particular, the neutral object of $P(^PB \to ^QC)$ is sent to the object $P \to Q$ of $P(B \to C; Q)$. Let P^{opp} denote the $(B, ^PB)$ -bitorsor with the same underlying sheaf as P but with local sections b and b' of B and PB acting as $(b, b') \cdot p = b'^{-1} \cdot p \cdot b^{-1}$. The functor

$$\mathsf{P}(B \to C; Q) \to \mathsf{P}(^{P}B \to {}^{Q}C) \tag{5}$$

sending $P' \to Q$ to $(P' \to Q) \wedge^B P^{\text{opp}}$ is a quasi-inverse to the functor in (4). Therefore, both functors are equivalences of categories.

Proposition 1.10. Let

$$1 \to A \to B \xrightarrow{v} C \to 0 \tag{6}$$

be an exact sequence of sheaves of groups on E, and let $P \rightarrow Q$ be a v-morphism of torsors. There is a natural bijection

$$H^1(S, {}^{P}A) \to H^1(S, B \to C; Q)$$

sending the neutral element of $H^1(S, {}^{P}A)$ to the element $[P \to Q]$ of $H^1(S, B \to C; Q)$.

Proof. We can use P to twist the sequence (6):

$$1 \to {}^{P}A \to {}^{P}B \to {}^{Q}C \to 1, \quad {}^{P}A = P \wedge^{B}A.$$

Now combine

$$H^1(S, {}^{P}A) \xrightarrow{1.7} H^1(S, {}^{P}B \to {}^{Q}C) \xrightarrow{(4)} H^1(S, B \to C; Q).$$

Remark 1.11. If in the proposition A is commutative, then the action of B on A factors through an action of C on A, and so

$${}^{P}A \stackrel{\mathrm{df}}{=} P \wedge^{B} A \cong P \wedge^{B} C \wedge^{C} A \cong Q \wedge^{C} A \stackrel{\mathrm{df}}{=} {}^{Q}A.$$

Notes. The basic definitions 1.4–1.5 are from Giraud 1971. The remaining statements can be found, or are hinted at, in Deligne 1979a, 2.4.3–2.4.4. See also Breen 1990. (The main ideas go back to Dedecker and Grothendieck in the 1950s.)

Cohomology and inverse limits

We now fix an affine scheme S and let E be the category of affine schemes over S endowed with the fpqc topology (that for which the covering families are the finite surjective families of flat morphisms).

Throughout this subsection, $(G_n, u_n)_n$ is an inverse system, indexed by (\mathbb{N}, \leq) , of flat affine group schemes of finite type over S with faithfully flat transition maps, and $G = \lim G_n$. Thus, G is a flat affine group scheme over S.

Proposition 1.12. The map $[P] \mapsto ([P \wedge^G G_n])_{n \ge 0}$

$$H^{1}(S,G) \to \varprojlim_{n} H^{1}(S,G_{n})$$
(7)

is surjective. For a G-torsor P, the fibre of the map containing [P] is $\varprojlim^1 G'_n(S)$ where G'_n is the inner form $P \wedge^G G_n$ of G_n .

Proof. A class c in $\lim H^1(S, G_n)$ is represented by an inverse system

$$P_0 \leftarrow P_1 \leftarrow \cdots \leftarrow P_n \leftarrow \cdots$$

with P_n a G_n -torsor. The inverse limit of this system is a G-torsor mapping to c.

Let P' and P be G-torsors such that $P'_n \approx P_n$ for all n, and choose isomorphisms $a_n \colon P'_n \to P_n$. Consider

$$\begin{array}{ccc} P'_{n+1} & \xrightarrow{a_{n+1}} & P_{n+1} \\ \downarrow^{v'} & & \downarrow^{v} \\ P'_n & \xrightarrow{a_n} & P_n. \end{array}$$

There is a unique isomorphism $b_n \colon P'_n \to P_n$ for which the diagram commutes, i.e., such that

$$b_n \circ v' = v \circ a_{n+1}.$$

Let e_n be the element of $Aut(P_n)$ such that $e_n \circ b_n = a_n$; then

$$e_n \circ v \circ a_{n+1} = a_n \circ v'.$$

Replacing $(a_n)_{n\geq 0}$ with $(c_n \circ a_n)_{n\geq 0}$ replaces $(e_n)_{n\geq 0}$ with $(c_n \cdot e_n \cdot u'c_{n+1}^{-1})_{n\geq 0}$ where u' is the transition map $\operatorname{Aut}(P_{n+1}) \to \operatorname{Aut}(P_n)$. Thus, the class of $(e_n)_{n\geq 0}$ in $\lim_{n \to \infty} \operatorname{Aut}(P_n)$ is independent of the choice of the a_n . Similarly, it depends only on the isomorphism class of P'. Therefore, we have a well-defined map from the fibre containing [P] to $\lim_{n\to\infty} \operatorname{Aut}(P_n)$, and it is straightforward to check that it is a bijection. Finally, (3) allows us to replace $\operatorname{Aut}(P_n)$ with $G'_n(S)$.

Corollary 1.13. When the G_n are commutative, there is an exact sequence

$$0 \to \varprojlim^1 G_n(S) \to H^1(S, G) \to \varprojlim H^1(S, G_n) \to 0.$$

Proof. In this case, $G'_n = G_n$.

Corollary 1.14. For any countable family $(G_i)_{i \in I}$ of flat affine group schemes,

$$H^1(S, \prod G_i) = \prod H^1(S, G_i).$$

Proof. This is certainly true for finite families. Thus, we may assume that I is infinite, and equals \mathbb{N} . Let $A_n = \prod_{0 \le i \le n} G_i$. For any $\prod_{i \ge 0} G_i$ -torsor P, the projection maps ${}^{P}A_n(S) \to {}^{P}A_{n-1}(S)$ admit sections, and so are surjective. Therefore $\varprojlim^{1}{}^{P}A_n(S) = 0$ (1.1a), and it follows that

$$H^{1}(S,G) \stackrel{1.12}{\cong} \varprojlim_{n} H^{1}(S,A_{n}) \cong \varprojlim_{n} \prod_{0 \le i \le n} H^{1}(S,G_{i}) \cong \prod_{i \ge 0} H^{1}(S,G_{i}).$$

Remark 1.15. Let $S = \text{Spec}(\mathbb{Q})$. Although the maps $u_n \colon G_n \to G_{n-1}$ are surjective, typically the maps $G_n(\mathbb{Q}) \to G_{n-1}(\mathbb{Q})$ will not be. In fact, typically, the inverse system $(G_n(\mathbb{Q}))_n$ will not satisfy (ML) and so $\varprojlim^1 G_n(\mathbb{Q})$ will be uncountable (1.1b). For example, consider a tower of distinct subfields of \mathbb{Q}^{al} ,

$$\mathbb{Q} \subset F_1 \subset \cdots \subset F_{n-1} \subset F_n \subset \cdots, \quad [F_n : \mathbb{Q}] < \infty.$$

There is an inverse system (G_n, u_n) with surjective transition maps for which G_n is the \mathbb{Q} -torus obtained from \mathbb{G}_{m/F_n} by restriction of scalars and u_n is the norm map. Then

$$(G_n(\mathbb{Q}), u_n(\mathbb{Q}))_{n \in \mathbb{N}} = (F_n^{\times}, \operatorname{Nm}_{F_n/F_{n-1}})_{n \in \mathbb{N}},$$

which fails (ML),⁴ and so $\underline{\lim}^1 G_n(\mathbb{Q})$ is uncountable.

Comparison with Galois cohomology

We now let S be the spectrum of a field k, and we let $H^1(k, -)$ denote $H^1(S, -)$. Choose a separable closure k^{sep} of k, and let $\Gamma = \text{Gal}(k^{\text{sep}}/k)$.

Proposition 1.16. For any smooth algebraic group N over k, there is a canonical isomorphism

$$H^1(k, N) \to H^1(\Gamma, N(k^{sep})).$$

Proof. An *N*-torsor *P* is represented by an algebraic variety over *k*, and hence acquires a point *p* over some subfield of k^{sep} of finite degree over *k*. The formula

$$\tau p = p \cdot a_{\tau} \tag{8}$$

defines a continuous crossed homomorphism $a_{\tau} \colon \Gamma \to N(k^{\text{sep}})$ whose cohomology class is independent of the choice of p and depends only on the isomorphism class of P. Thus, we have a well-defined map $H^1(k, N) \to H^1(\Gamma, N(k^{\text{sep}}))$, and it follows from descent theory that this is an isomorphism.

⁴To see this, use that, for a finite extension E/F of number fields and a finite prime v of F, $\operatorname{ord}_v(\operatorname{Nm} E^{\times})$ is the ideal in \mathbb{Z} generated by the residue class degrees of the primes of E lying over v.

1.17. Let N be a smooth algebraic group over k, and let $f: \Gamma \to Aut(N)(k^{\text{sep}})$ be a continuous crossed homomorphism (discrete topology on $Aut(N)(k^{\text{sep}})$). The "twist" of N by f is a smooth algebraic group $_fN$ over k such that $_fN(k^{\text{sep}}) = N(k^{\text{sep}})$ but with $\tau \in \Gamma$ acting according to the rule

$$\tau * x = f(\tau) \cdot \tau x.$$

When we let N act on itself by inner automorphisms, a crossed homomorphism $f: \Gamma \to N(k^{\text{sep}})$ defines a twist $_fN$ of N with $\tau \in \Gamma$ acting on $_fN(k^{\text{sep}})$ by

$$\tau * x = f(\tau) \cdot \tau x \cdot f(\tau)^{-1}.$$

1.18. Let $G = \varprojlim(G_n, u_n)$ be as in the preceding subsection but with G_n now a smooth algebraic group over k, and define $H^1_{\text{cts}}(\Gamma, G)$ be the cohomology set computed using crossed homomorphisms $\Gamma \to G(k^{\text{sep}})$ that are continuous for the profinite topology on Γ and the inverse limit topology on $G(k^{\text{sep}}) = \varprojlim G_n(k^{\text{sep}})$ (discrete topology on $G_n(k^{\text{sep}})$). Thus, giving a continuous crossed homomorphism $f \colon \Gamma \to G(k^{\text{sep}})$ amounts to giving a compatible family of continuous crossed homomorphisms $f_n \colon \Gamma \to G_n(k^{\text{sep}})$.

Proposition 1.19. The map

$$H^1_{cts}(\Gamma, G) \to \underline{\lim} H^1(\Gamma, G_n)$$

sending [f] to $([f_n])_{n\geq 0}$ is surjective. The fibre of the map containing [f] equals $\varprojlim^1 G'_n(k)$ where $G'_n = {}_f G_n$.

Proof. Each class c in $\varprojlim H^1(S, G_n)$ is represented by a family $(f_n)_{n\geq 0}$ of crossed homomorphisms, which can be chosen so that $f_{n-1} = u_n \circ f_n$. The f_n define a continuous crossed homomorphism $f: \Gamma \to G(k^{\text{sep}})$ mapping to c.

Let f' and f be continuous crossed homomorphisms such that $f'_n \sim f_n$ for all n, and choose $a_n \in G_n(k^{\text{sep}})$ in such a way that

$$f'(\tau)_n = a_n^{-1} \cdot f(\tau)_n \cdot \tau a_n.$$
(9)

Define $e_n \in G_n(k^{sep})$ by the equation

$$e_n \cdot ua_{n+1} = a_n.$$

On applying u to the equation $(9)_{n+1}$, we obtain the equation

$$f'(\tau)_n = (ua_{n+1})^{-1} \cdot f(\tau)_n \cdot \tau(ua_{n+1}),$$

or,

$$f'(\tau)_n = a_n^{-1} \cdot e_n \cdot f(\tau)_n \cdot \tau e_n^{-1} \cdot \tau a_n.$$

On comparing this with (9), we find that

$$e_n = f_n(\tau) \cdot \tau e_n \cdot f_n(\tau)^{-1},$$

1 PRELIMINARIES

ie., that

$$e_n \in ({}_f G_n)(k^{\operatorname{sep}})^{\Gamma} = ({}_f G_n)(k).$$

The element a_n can be replaced by $c_n \cdot a_n$, where c_n is any element of $({}_fG_n)(k)$. When this is done for each n, $(e_n)_{n\geq 0}$ is replaced by $(c_n \cdot e_n \cdot (uc_{n+1})^{-1})_{n\geq 0}$. Thus, the class of (e_n) in $\lim_{k \to 0} {}^1({}_fG_n)(k)$ is independent of the choice of the a_n . Similarly, it depends only on the cohomology class of f'. Therefore, we have a well-defined map from the fibre containing [f] to $\lim_{k \to 0} {}^1({}_fG_n)(k)$, and it is straightforward to check that this is a bijection.

Proposition 1.20. There is a canonical isomorphism of pointed sets

$$H^1(k,G) \to H^1_{cts}(\Gamma,G).$$

Proof. Let P be a G-torsor, and let $P_n = u_n P$. Then $P(k^{\text{sep}}) = \varprojlim P_n(k^{\text{sep}})$, which, because the maps $P_{n+1}(k^{\text{sep}}) \to P_n(k^{\text{sep}})$ are surjective, is nonempty. Choose a $p \in P(k^{\text{sep}})$. Then the formula

$$\tau p = p \cdot f(\tau)$$

defines a continuous crossed homomorphism $f: \Gamma \to G(k^{\text{sep}})$ whose cohomology class is independent of the choice of p and of the choice of P in its isomorphism class. Therefore, we have a well-defined map $H^1(k, G) \to H^1_{\text{cts}}(\Gamma, G)$. Since this map is compatible with the maps in Propositions 1.12 and 1.19, they, together with (1.16) show that it is a bijection. \Box

Remark 1.21. Let S be the spectrum of a field. The following conditions on an affine group scheme G over S are equivalent:

- (a) the set of closed normal subgroup schemes $H \subset G$ such that G/H is of finite type over S is infinite and countable;
- (b) $G = \lim_{n \to \infty} (G_n, u_n)$ where $(G_n, u_n)_n$ is an inverse system, indexed by (\mathbb{N}, \leq) , of affine algebraic groups over S with surjective transition maps.

An affine group scheme G satisfying these conditions will be said to be *separable*. When G is separable, any inverse system satisfying (b) is cofinal in the inverse system of all algebraic quotients of G. Therefore, the former can be replaced by the latter, which makes Proposition 1.19 more canonical.

Notes. Proposition 1.16 is a standard result. In the commutative case, Proposition 1.19 is proved in Tate 1976.

Application to periods

Let $Mot(\mathbb{Q})$ denote the category of motives based on all smooth projective varieties over \mathbb{Q} . Let $CM(\mathbb{Q})$ be the Tannakian subcategory of $Mot(\mathbb{Q})$ generated by the zero-dimensional varieties over \mathbb{Q} and the abelian varieties of CM-type (see §2). Let $G^{Mot} = Aut^{\otimes}(\omega_B)$ and $P^{Mot} = Isom^{\otimes}(\omega_B, \omega_{dR})$, and define G^{CM} and P^{CM} similarly. From the inclusion $CM(\mathbb{Q}) \subset Mot(\mathbb{Q})$, we obtain a faithfully flat homomorphism $G^{Mot} \to G^{CM}$. **Theorem 1.22.** If the kernel of $G^{Mot} \to G^{CM}$ is an inverse limit of simply connected semisimple groups, then the isomorphism class of $P^{Mot} \to P^{CM}$ in $P(G^{Mot} \to G^{CM}; P^{CM})$ is uniquely determined by its class over \mathbb{R} .

Proof. Let $G = \text{Ker}(G^{\text{Mot}} \to G^{\text{CM}})$. The condition on G implies that it is, in fact, a product of semisimple groups each of which is simply connected. Moreover, the product is countable because $\text{Mot}(\mathbb{Q})$ is generated as a Tannakian category by a countable set of varieties. According to Proposition 1.10, the isomorphism classes in $P(G^{\text{Mot}} \to G^{\text{CM}}; P^{\text{CM}})$ are classified by $H^1(\mathbb{Q}, G')$ where $G' = P^{\text{Mot}} \wedge^{G^{\text{Mot}}} G$. As G' is a form of G, it also is a countable product of simply connected semisimple groups, and so the proposition follows from Corollary 1.14 and the theorem of Kneser, Harder, and Chernousov (see 3.6 below). □

Remark 1.23. (a) The group G^{Mot} is proreductive because $Mot(\mathbb{Q})$ is semisimple. If Deligne's hope (1979b, 0.10) that every Hodge class is an absolute Hodge class is true, then $(G^{Mot})^{\circ}$ is the group attached to the category of motives over \mathbb{Q}^{al} (see Deligne and Milne 1982, 6.22, 6.23, where the hypothesis was inadvertently omitted); moreover, the group G in the above proof is the kernel of the canonical homomorphism from $(G^{Mot})^{\circ}$ to the Serre group (ibid. p220), and it is the derived group of $(G^{Mot})^{\circ}$; it is therefore an inverse limit of semisimple groups.

(b) It is generally hoped that the derived group of $(G^{Mot})^{\circ}$ is simply connected — see the question in Serre 1994, 8.1.

Remark 1.24. Let $G = \text{Ker}(G^{\text{Mot}} \to G^{\text{CM}})$, and let $H \subset G$ be the intersection of the kernels of the homomorphisms from G onto simply connected semisimple algebraic groups over \mathbb{Q} . Let $\text{Mot}^H(\mathbb{Q})$ be the subcategory of $\text{Mot}(\mathbb{Q})$ of objects on which H acts trivially. Then, with the obvious notations, the isomorphism class of $P^{\text{Mot}^H} \to P^{\text{CM}}$ in $P(G^{\text{Mot}^H} \to G^{\text{CM}}; P^{\text{CM}})$ is uniquely determined by its class over \mathbb{R} . The preceding remark indicates that it is reasonable to hope that H = 0.

2 Periods of abelian varieties with complex multiplication

Periods of zero-dimensional varieties

The category of motives based on the zero-dimensional varieties over \mathbb{Q} is denoted $\operatorname{Art}(\mathbb{Q})$ and is called the category of Artin motives over \mathbb{Q} . The Betti fibre functor ω_B defines an equivalence of $\operatorname{Art}(\mathbb{Q})$ with the category of continuous representations of Γ on finitedimensional \mathbb{Q} -vector spaces (Deligne and Milne 1982, 6.17) from which it follows that $\operatorname{Aut}^{\otimes}(\omega_B)$ is the constant profinite group scheme Γ with $\Gamma(\mathbb{Q}) = \Gamma(\mathbb{Q}^{\operatorname{al}}) = \Gamma$. The de Rham fibre functor is $hX \mapsto \Gamma(X, \mathcal{O}_X)$, from which the next statement follows easily.

Theorem 2.1. Let P^{Art} be the period torsor for $\operatorname{Art}(\mathbb{Q})$. Then $P^{\text{Art}} \cong \operatorname{Spec} \mathbb{Q}^{al}$ with its natural action of Γ , and p^{Art} is the obvious \mathbb{Q}^{al} -point of $\operatorname{Spec} \mathbb{Q}^{al}$. Thus, the period point p^{Art} has coordinates in \mathbb{Q}^{al} , and the cocycle corresponding to the pair $(P^{\text{Art}}, p^{\text{Art}})$ is the crossed homomorphism $\Gamma \to \Gamma(\mathbb{Q}^{al}), \tau \mapsto \tau$.

Note that the theorem determines the pair (P^{Art}, p^{Art}) uniquely up to a unique isomorphism.

Throughout this section, we use f to denote the (crossed) homomorphism in the theorem.

Notations for tori

For a finite étale \mathbb{Q} -algebra A, let

$$\Sigma_{A/\mathbb{Q}} = \operatorname{Hom}_{\mathbb{Q}\operatorname{-alg}}(A, \mathbb{Q}^{\operatorname{al}}),$$

$$\Gamma_{A/\mathbb{Q}} = \operatorname{Aut}_{\mathbb{Q}\operatorname{-alg}}(A),$$

and let $(\mathbb{G}_m)_{A/\mathbb{Q}}$ be the torus over \mathbb{Q} obtained from $\mathbb{G}_{m/A}$ by restriction of scalars. Thus

$$(\mathbb{G}_m)_{A/\mathbb{Q}}(R) = (A \otimes R)^{\times}$$

for all \mathbb{Q} -algebras R. For an infinite field extension K/k,

$$(\mathbb{G}_m)_{K/k} = \varprojlim (\mathbb{G}_m)_{K'/\mathbb{Q}}, \quad K' \subset K, \quad [K':\mathbb{Q}] < \infty.$$

2.2. There is an equivalence $T \mapsto X^*(T) =_{df} \operatorname{Hom}(T_{/\mathbb{Q}^{al}}, \mathbb{G}_m)$ from the category of tori over \mathbb{Q} to the category of finitely generated free \mathbb{Z} -modules endowed with a continuous left action of Γ .

2.3. There is an equivalence $A \mapsto \Sigma_{A/\mathbb{Q}}$ from the category of finite étale \mathbb{Q} -algebras to the category of finite sets endowed with a continuous left action of Γ . A quasi-inverse is provided by

$$\Sigma \mapsto A(\Sigma) \stackrel{\text{df}}{=} \operatorname{Hom}_{\Gamma}(\Sigma, \mathbb{Q}^{al}).$$

If $\Sigma \leftrightarrow A$, then the decomposition $\Sigma = \amalg \Sigma_i$ of Σ into orbits corresponds to the decomposition $A = \prod A(\Sigma_i)$ of A into a product of fields. If Γ acts transitively on Σ , then the choice of an $e \in \Sigma$ determines isomorphisms $\Gamma/\Gamma_e \to \Sigma$ and $A(\Sigma) \to (\mathbb{Q}^{\mathrm{al}})^{\Gamma_e}$. Here $\Gamma_e = \{\tau \in \Gamma \mid \tau e = e\}$.

2.4. On combining these equivalences, we see that there is a fully faithful functor $\Sigma \mapsto T^{\Sigma}$ from the category of finite sets endowed with a continuous left Γ -action to the category of tori, for which $T^{\Sigma} = (\mathbb{G}_m)_{A(\Sigma)/\mathbb{Q}}, X^*(T^{\Sigma}) = \mathbb{Z}[\Sigma]$ (free \mathbb{Z} -module on Σ with the natural left action of Γ), and $T^{\Sigma_1 \sqcup \Sigma_2} = T^{\Sigma_1} \times T^{\Sigma_2}$.

Abelian varieties with complex multiplication

For an abelian variety A defined over a subfield k of \mathbb{C} , the Mumford-Tate group of A is

$$\mathrm{MT}(A) = \mathcal{A}ut^{\otimes}(\omega_{\mathbf{B}}|\langle A_{/\mathbb{C}}\rangle^{\otimes}).$$

where $\langle A_{/\mathbb{C}} \rangle^{\otimes}$ be the category of motives based on $A_{/\mathbb{C}}$ and the projective spaces. There is a canonical cocharacter μ^A of MT(A) that splits the Hodge filtration on $H^1_B(A)$. When MT(A) is commutative (hence a torus), we say that A is of *CM-type*⁵, and we define the *reflex field* of A to be the field of definition of μ^A . It is a CM-subfield of \mathbb{C} .

For subfields k, K of \mathbb{C} with K a CM-field, we define $CM^{K}(k)$ to be the category of motives based on

- the abelian varieties of CM-type over k with reflex field contained in K,
- the projective spaces, and
- the zero-dimensional varieties.

When $K = \mathbb{Q}^{cm}$, the composite of all CM-subfields of \mathbb{Q}^{al} , we omit it from the notation.

The category $CM(\mathbb{C})$

2.5. Let K be a CM-subfield of \mathbb{C} . With the notation of (2.4), $(\mathbb{G}_m)_{K/\mathbb{Q}} = T^{\Sigma_{K/\mathbb{Q}}}$ and $X^*((\mathbb{G}_m)_{K/\mathbb{Q}}) = \mathbb{Z}[\Sigma_{K/\mathbb{Q}}]$. The group $S^K = Aut^{\otimes}(\omega_B | \mathsf{CM}^K(\mathbb{C}))$ is called the *Serre* group for K. When K has finite degree over \mathbb{Q} , S^K is the quotient of $(\mathbb{G}_m)_{K/\mathbb{Q}}$ such that

$$X^*(S^K) = \{ n \in \mathbb{Z}[\Sigma_{K/\mathbb{Q}}] \mid n + \iota n = \text{constant} \}.$$

Thus, there is an exact sequence

$$0 \to (\mathbb{G}_m)_{F/\mathbb{Q}} \to (\mathbb{G}_m)_{K/\mathbb{Q}} \times \mathbb{G}_m \to S^K \to 0$$
(10)

where F is the largest totally real subfield of K. We have

$$S = \lim_{K \in \mathcal{K}} S^K$$

where \mathcal{K} is the set of all CM-subfields of \mathbb{C} finite and Galois over \mathbb{Q} .

2.6. For an abelian variety A over \mathbb{C} , there is a canonical surjection $S \to MT(A)$, which factors through S^K if and only if A has reflex field contained in K.

Notes. For more on Mumford-Tate groups, see Deligne 1982a, §3.

⁵Other authors, and this author at other times, say that A is potentially of CM-type.

The category $CM(\mathbb{Q}^{al})$

2.7. The functor the category of abelian varieties of CM-type up to isogeny over \mathbb{Q}^{al} to the similar category over \mathbb{C} is an equivalence. For any CM-subfield K of \mathbb{C} , the functor "extension of scalars" $CM^{K}(\mathbb{Q}^{al}) \to CM^{K}(\mathbb{C})$ is an equivalence of tensor categories.

The period torsor

2.8. Let *K* be a CM-subfield of \mathbb{C} , Galois over \mathbb{Q} . Then Γ acts on the terms of the sequence (10) in such a way that the action on the sequence of \mathbb{Q} -points

$$1 \to F^{\times} \to K^{\times} \times \mathbb{Q}^{\times} \to S^{K}(\mathbb{Q}) \to 1$$

is the obvious one. Let

$$G_{\mathbf{B}}^{\mathsf{CM},K} = \mathcal{A}ut^{\otimes}(\omega_{\mathbf{B}}|\mathsf{CM}^{K}(\mathbb{Q})).$$

The tensor functors

$$\mathsf{CM}^{K}(\mathbb{Q}^{\mathrm{al}}) \leftarrow \mathsf{CM}^{K}(\mathbb{Q}) \leftarrow \mathsf{Art}(\mathbb{Q})$$
 (11)

define homomorphisms

$$1 \to S^K \to G_{\mathbf{B}}^{\mathsf{CM},K} \xrightarrow{v} \Gamma \to 1.$$
(12)

This sequence is exact and the action of Γ on S^K it defines is that described above (Deligne 1982b, Lemme 1, Lemme 2).

2.9. Let $P^{CM,K}$ be the period torsor for $CM^{K}(\mathbb{Q})$. The second functor in (11) defines a v-morphism $P^{CM,K} \to P^{Art}$, and since P^{Art} is known (Theorem 2.1), in order to determine $P^{CM,K}$ it suffices to determine $P^{CM,K} \to P^{Art}$ as an object of $P(G_B^{CM,K} \to \Gamma; P^{Art})$. The first step is to correctly identify the cohomology group classifying the isomorphism classes of objects in this category.

Proposition 2.10. Let $f: \Gamma \to \Gamma(\mathbb{Q})$ be as in Theorem 2.1, and let ${}_{f}S^{K}$ be the twist of S^{K} by f. There is a natural one-to-one correspondence between the set of isomorphism classes in $\mathsf{P}(G_{B}^{\mathsf{CM},K} \to \Gamma; P^{\mathsf{Art}})$ and $H^{1}(\mathbb{Q}, {}_{f}S^{K})$ (fpqc cohomology).

Proof. Apply Proposition 1.10, Remark 1.11, and Theorem 2.1.

In particular, when $K = \mathbb{Q}^{cm}$, the isomorphism classes of objects $P \to P^{Art}$ are classified by the group $H^1(\mathbb{Q}, {}_fS)$. According to Corollary 1.13, there is an exact sequence

$$0 \to \varprojlim_{K \in \mathcal{K}}^{1} {}_{f}S^{K}(\mathbb{Q}) \to H^{1}(\mathbb{Q}, {}_{f}S) \to \varprojlim_{K \in \mathcal{K}} H^{1}(\mathbb{Q}, {}_{f}S^{K}) \to 0.$$

We shall show:

Proposition 2.11. The group $\varprojlim_{K \in \mathcal{K}} H^1(\mathbb{Q}, {}_fS^K) = 0$, but $\varprojlim_{K \in \mathcal{K}} {}^1f^K(\mathbb{Q})$ is uncountable. Therefore,

$$H^1(\mathbb{Q}, {}_fS) \cong \varprojlim_{K \in \mathcal{K}}^1 {}_fS^K(\mathbb{Q})$$

and is uncountable.

Twisting $(\mathbb{G}_m)_{L/\mathbb{Q}}$

2.12. Let *L* be a finite extension of \mathbb{Q} , and let $T = (\mathbb{G}_m)_{L/\mathbb{Q}}$.

- (a) The (left) action of $\tau \in \Gamma$ on $\Sigma_{L/\mathbb{Q}}$, $\tau\sigma = \tau \circ \sigma$, corresponds to the natural (left) action of Γ on $X^*(T) = \mathbb{Z}[\Sigma_{L/\mathbb{Q}}]$ (see 2.4).
- (b) Assume $L \subset \mathbb{Q}^{al}$ and is Galois over \mathbb{Q} . Identify $\Sigma_{L/\mathbb{Q}}$ with $\Gamma_{L/\mathbb{Q}}$. There is then a natural (left) action of Γ on T (as a torus over \mathbb{Q}), which defines a (left) action of Γ on $X^*(T) = \mathbb{Z}[\Gamma_{L/\mathbb{Q}}]$. The former gives the obvious action of Γ on $T(\mathbb{Q}) = L^{\times}$, and the latter corresponds to the (left) action of $\tau \in \Gamma$ on $\Gamma_{L/\mathbb{Q}}$, $\tau\sigma = \sigma \circ \tau^{-1}$.

Lemma 2.13. For a subfield L of \mathbb{Q}^{al} , finite and Galois over \mathbb{Q} , let ${}_{f}\Gamma_{L/\mathbb{Q}}$ denote the twist of $\Gamma_{L/\mathbb{Q}}$ by f (so that $\tau \in \Gamma$ acts by $\tau \sigma = \tau \circ \sigma \circ \tau^{-1}$ (Serre 1964, 5.3)), and let

$$B(L) = A({}_{f}\Gamma_{L/\mathbb{O}}).$$

Then

$$f(\mathbb{G}_m)_{L/\mathbb{Q}}\cong (\mathbb{G}_m)_{B(L)/\mathbb{Q}}$$

Proof. Clearly, $_f(\mathbb{G}_m)_{L/\mathbb{Q}} \cong T^{_f\Gamma_{L/\mathbb{Q}}}$, which equals $(\mathbb{G}_m)_{A(_f\Gamma_{L/\mathbb{Q}})/\mathbb{Q}}$.

Note that the orbits of Γ acting on ${}_{f}\Gamma_{L/\mathbb{Q}}$ are the conjugacy classes C in $\Gamma_{L/\mathbb{Q}}$, and so (2.3, 2.4)

$$B(L) = \prod_C A(C), \quad (\mathbb{G}_m)_{B(L)/\mathbb{Q}} = \prod_C T^C.$$

For any $\sigma \in C$, $A(C) \cong L^{Z(\sigma)}$ where $Z(\sigma)$ is the centralizer of σ in $\Gamma_{L/\mathbb{Q}}$.

The tori ${}_{f}S^{K}$ and ${}_{f}\bar{S}^{K}$

Let $K \in \mathcal{K}$. Let $w \colon \mathbb{G}_m \to S^K$ be the weight homomorphism, and let $\bar{S}^K = S^K / w(\mathbb{G}_m)$. Then

$$X^*(\bar{S}^K) = \{ n \in \mathbb{Z}[\varSigma_K] \mid n + \iota n = 1 \},\$$

and there is an exact sequence

$$1 \to (\mathbb{G}_m)_{F/\mathbb{Q}} \to (\mathbb{G}_m)_{K/\mathbb{Q}} \to \bar{S}^K \to 1.$$
(13)

П

The group Γ acts on the exact sequence (13), and so we can twist the sequence by f to obtain an exact sequence

$$1 \to {}_f(\mathbb{G}_m)_{F/\mathbb{Q}} \to {}_f(\mathbb{G}_m)_{K/\mathbb{Q}} \to {}_f\bar{S}^K \to 1.$$
(14)

Lemma 2.14. If K contains a quadratic imaginary field k, then

$$_f \bar{S}^K \approx (\mathbb{G}_m)_{B(F)/\mathbb{Q}}$$

where F is the largest totally real subfield of K.

Proof. As

$$\Gamma_{K/\mathbb{Q}} = \Gamma_{F/\mathbb{Q}} \times \Gamma_{k/\mathbb{Q}}, \quad \Gamma_{k/\mathbb{Q}} = \{1, \iota\},\$$

for each conjugacy class C in $\Gamma_{F/\mathbb{Q}}$, there are exactly two conjugacy classes in $\Gamma_{K/\mathbb{Q}}$ mapping to it, namely,

$$C_1 = \{(\tau, 1) \mid \tau \in C\}, \quad C_\iota = \{(\tau, \iota) \mid \tau \in C\}.$$

Therefore, (14) can be written

$$1 \to \prod_C T^C \to \prod_C (T^{C_1} \times T^{C_{\iota}}) \to {}_f \bar{S}^K \to 1$$

(product over the conjugacy classes in $\Gamma_{F/\mathbb{Q}}$). Since

$$C \cong C_1 \cong C_\iota$$
 (as Γ -sets),

each of the maps

$$\prod_C T^C \to \prod_C T^{C_1}, \quad \prod_C T^{C_1} \to {}_f \bar{S}^K$$

is an isomorphism.

The group $\varprojlim_{K \in \mathcal{K}} H^1(\mathbb{Q}, {}_fS^K)$.

Proposition 2.15 (Wintenberger 1990, 1.3). If K contains a quadratic imaginary number field, then $H^1(\mathbb{Q}, {}_fS^K) = 0$.

Proof. From the exact sequence

$$1 \to \mathbb{G}_m \to {}_f S^K \to {}_f \bar{S}^K \to 1 \tag{15}$$

we see that it suffices to show that $H^1(\mathbb{Q}, {}_f\bar{S}^K) = 0$, but this follows from Lemma 2.14 and Hilbert's Theorem 90.

Corollary 2.16. The group $\varprojlim_{K \in \mathcal{K}} H^1(\mathbb{Q}, {}_f\bar{S}^K) = 0.$

Proof. The CM-fields satisfying the hypothesis of the proposition are cofinal in \mathcal{K} .

Remark 2.17. Wintenberger (1990, p3) shows by example that the proposition is false without the hypothesis on K.

The group $\lim_{K \in \mathcal{K}} {}^{1}_{f} S^{K}(\mathbb{Q})$.

Proposition 2.18. The group $\varprojlim_{K \in \mathcal{K}}^{1} fS^{K}(\mathbb{Q})$ is uncountable.

From (2) applied to the cohomology sequence of (15), we see that

$$\varprojlim_{K\in\mathcal{K}}^{1}{}_{f}S^{K}(\mathbb{Q}) \xrightarrow{\cong} \varprojlim_{K\in\mathcal{K}}^{1}{}_{f}\bar{S}^{K}(\mathbb{Q}),$$

and so we compute the second group.

Lemma 2.19. Let \mathcal{F} be the set of totally real subfields of \mathbb{C} that are finite and Galois over \mathbb{Q} . Then the inverse system $({}_f \bar{S}^K)_{K \in \mathcal{K}}$ is equivalent with the inverse system $((\mathbb{G}_m)_{B(F)/\mathbb{Q}})_{F \in \mathcal{F}}$. Thus,

$$\varprojlim_{K\in\mathcal{K}}^{1}{}_{f}\bar{S}^{K}(\mathbb{Q})\approx\varprojlim_{F\in\mathcal{F}}^{1}B(F)^{\times}$$

(on the right, the transition maps are the norm maps).

Proof. Fix a quadratic imaginary field k. The fields $K \in \mathcal{K}$ containing k form a cofinal set. Once k has been fixed, the isomorphism in Lemma 2.14 becomes canonical. In particular, it is natural for the norm maps.

Note that the isomorphism in the lemma depends only on the choice of k.

Example 2.20. If $\Gamma_{F/\mathbb{Q}} = \operatorname{Gal}(F/\mathbb{Q})$ is commutative, B(F) is a product of copies of \mathbb{Q} indexed by the elements of $\Gamma_{F/\mathbb{Q}}$. Thus, $(\mathbb{G}_m)_{B(F)/\mathbb{Q}} \approx \mathbb{G}_m^{[F:\mathbb{Q}]}$.

Example 2.21. Let $F = F_1 \cdot F_2$ with $F_1, F_2 \in \mathcal{F}$ and $F_1 \cap F_2 = \mathbb{Q}$. Then

 $\operatorname{Gal}(F/\mathbb{Q}) \cong \operatorname{Gal}(F_1/\mathbb{Q}) \times \operatorname{Gal}(F_2/\mathbb{Q})$

and a conjugacy class C in $Gal(F_1/\mathbb{Q})$ is the image of the conjugacy class $C \times \{1\}$ in $Gal(F/\mathbb{Q})$. Therefore, $B(F_1)$ is a direct factor of B(F), and so the norm map

$$B(F)^{\times} \to B(F_1)^{\times}$$

is surjective.

Example 2.22. Fix an odd prime l, and write C_l for any cyclic group of order l. Define G to be the semi-direct product $N \rtimes_{\theta} Q$ of $N = C_l \times C_l$ (generators a, b) with $Q = C_l$ (generator c) relative to the homomorphism $\theta \colon C_l \to \operatorname{Aut}(C_l \times C_l)$ for which

$$\theta(c^i) = \begin{pmatrix} 1 & 0\\ i & 1 \end{pmatrix}$$
, i.e., $\theta(c^i)(a) = ab^i$, $\theta(c^i)(b) = bai$

Then G has generators a, b, c, and relations

$$a^{l} = b^{l} = c^{l} = 1, \quad ab = cac^{-1}, \quad [b, a] = 1 = [b, c].$$

All elements $\neq 1$ in G have order l, and the centre of G is $\langle b \rangle$.

The inverse image under $G \to Q$ of the conjugacy class $\{c\}$ breaks up into l conjugacy classes, namely,

$$\{a^{j}c, ba^{j}c, \dots, b^{l-1}a^{j}c\}, \quad 0 \le j \le l-1,$$

(because $a^{-1}ca = bc$). The centralizer of $a^{j}c$ in G is $\langle b, a^{j}c \rangle$, which has order l^{2} .

Let E be an extension of \mathbb{Q} with Galois group G, and let $F_0 = E^N$. Then (2.20)

$$B(F_0) \cong F_0^{\{1\}} \times F_0^{\{c\}} \times \dots \times F_0^{\{c^{l-1}\}}, \quad F_0^{\{c^i\}} = \mathbb{Q},$$

and the inverse image of $F_0^{\{c\}}$ in B(E) under the norm map is

$$E^{\langle b,c\rangle} \times E^{\langle b,ac\rangle} \times \dots \times E^{\langle b,a^{l-1}c\rangle}$$

Lemma 2.23. The group $\varprojlim_{F \in \mathcal{F}}^1 B(F)^{\times}$ is uncountable.

Proof. According to Proposition 1.1b, it suffices to show that the inverse system $(B(F)^{\times})_{F \in \mathcal{F}}$ fails (ML). Examples 2.20 and 2.21 show that we shall need to consider nonabelian Galois groups and nonsplit extensions.

Fix an odd prime number l, and choose a prime number p_0 that splits completely in $\mathbb{Q}[\sqrt[l]{1}]$. Then $l|p_0 - 1$, and so there is a surjective homomorphism

$$(\mathbb{Z}/p_0\mathbb{Z})^{\times} \to C_l.$$

Let F_0 be the subfield of $\mathbb{Q}[\sqrt[p_0]{1}]$ fixed by the kernel of one such homomorphism.

To prove that $(B(F)^{\times})_{F \in \mathcal{F}}$ fails (ML) it suffices to show that, for each $L \in \mathcal{F}$ containing F_0 , there exists an $E \in \mathcal{F}$ containing F_0 and such that

$$\operatorname{Nm}_{B(E)/B(F_0)}(B(E)^{\times}) \not\supseteq \operatorname{Nm}_{B(L)/B(F_0)}(B(L)^{\times});$$

for then $E \cdot L \in \mathcal{F}$, but

$$\operatorname{Nm}_{B(E \cdot L)/B(F_0)}(B(E \cdot L)^{\times}) \neq \operatorname{Nm}_{B(L)/B(F_0)}(B(L)^{\times}).$$

Choose a prime p_1 that splits completely in $L[\sqrt[l]{1}, \sqrt[l]{p_0}]$, and construct a cyclic extension F_1 of \mathbb{Q} of degree l by choosing a surjective homomorphism $(\mathbb{Z}/p_1\mathbb{Z})^{\times} \to C_l$, as before. Then $F =_{df} F_0 \cdot F_1$ has Galois group

$$\operatorname{Gal}(F/\mathbb{Q}) \cong C_l \times C_l.$$

Let G be as in Example 2.22, and consider the extension

$$1 \to \langle b \rangle \to G \to \langle a, c \rangle \to 1.$$

Let

$$\alpha \colon G/\langle b \rangle \to \operatorname{Gal}(F/\mathbb{Q})$$

be the isomorphism sending c to a generator of $\operatorname{Gal}(F_0/\mathbb{Q})$ and a to a generator of $\operatorname{Gal}(F_1/\mathbb{Q})$. The only primes ramifying in F are p_0 and p_1 , and for $p = p_0$ or p_1

- *l* divides p-1 (because both primes split in $\mathbb{Q}[\sqrt[l]{1}]$);

- for all primes v of F dividing p, F_v is totally ramified over \mathbb{Q}_p (p_i is totally ramified in F_i ; p_1 splits completely in F_0 by construction; p_0 splits completely in F_1 because it becomes an l^{th} power in \mathbb{F}_{p_1}).

Now an argument of Scholz and Reichardt (see Serre 1992, Theorem 2.1.3) shows that there exists a Galois extension E of \mathbb{Q} containing F for which there is a commutative diagram



Note that E, being Galois of odd degree over \mathbb{Q} , must be totally real.



I claim that the image of $B(E)^{\times}$ in $F_0^{\{c\}\times}$ does not contain the image of $B(L)^{\times}$. In order to show this, it suffices to show that the image of $(B(E) \otimes \mathbb{Q}_{p_1})^{\times}$ in $(F_0^{\{c\}} \otimes \mathbb{Q}_{p_1})^{\times} = \mathbb{Q}_{p_1}^{\times}$ does not contain the image of $(B(L) \otimes \mathbb{Q}_{p_1})^{\times}$. But, because p_1 splits completely in L, the second group is $\mathbb{Q}_{p_1}^{\times}$. On the other hand, p_1 is totally ramified in each field $E^{\langle b, a^j c \rangle}$ (because $E^{\langle b, a^j c \rangle} \cdot F_0 = F$, $(p_1) = \mathfrak{p}_1^l \cdots \mathfrak{p}_l^l$ in F, and p_1 splits completely in F_0), and so, for each $v|p, E_v^{\langle b, a^j c \rangle}$ is the (unique) tamely ramified cyclic extension of \mathbb{Q}_{p_1} of degree l. Thus, the image of $\prod_v E_v^{\langle b, a^j c \rangle \times}$ in $\mathbb{Q}_{p_1}^{\times}$ does not contain $\mathbb{Z}_{p_1}^{\times}$.

Characterizing the period torsor

Theorem 2.24. Let K be a CM-subfield of \mathbb{C} , Galois over \mathbb{Q} .

- (a) If $[K:\mathbb{Q}] < \infty$, then $\mathsf{P}(G_B^{\mathsf{CM},K} \to \Gamma; P^{\mathsf{Art}})$ contains exactly one isomorphism class, which is represented by $P^{\mathsf{CM},K} \to P^{\mathsf{Art}}$.
- (b) If $K = \mathbb{Q}^{cm}$, then $\mathsf{P}(G_B^{\mathsf{CM},K} \to \Gamma; P^{\mathsf{Art}})$ contains uncountably many isomorphism classes.

Proof. According to Proposition 2.10, the isomorphism classes in $P(G_B^{CM,K} \to \Gamma; P^{Art})$ are classified by $H^1(\mathbb{Q}, {}_fS^K)$. Therefore, (a) and (b) follow respectively from Proposition 2.15 and Proposition 2.11.

Remark 2.25. In fact, we have shown that the uncountable group $\lim_{K \to K} f^{K}(\mathbb{Q})$ acts simply transitively on the set of isomorphism classes of objects in $\mathsf{P}(G_{\mathsf{B}}^{\mathsf{CM}} \to \Gamma; P^{\mathsf{Art}})$. To make this explicit, let K_0 be a quadratic imaginary field, and let

$$K_0 \subset \cdots \subset K_n \subset \cdots \subset \mathbb{Q}^{\mathrm{cm}}$$

be a sequence of CM-fields with union \mathbb{Q}^{cm} . Let $S_n = S^{K_n}$ and $P_n^{CM} = P^{CM,K_n}$. From an element $s = (s_n)_{n \in \mathbb{N}}$ of $\prod_f S_n(\mathbb{Q})$ we obtain a *v*-morphism of torsors $P^{CM}(s) \to P^{Art}$ by modifying the transition maps in P^{CM} : define

$$P^{\mathsf{CM}}(s) = \lim_{n \in \mathbb{N}} (P_n^{\mathsf{CM}}, v_n \circ s_n).$$

Then, the isomorphism class of $P^{\mathsf{CM}}(s) \to P^{\mathsf{Art}}$ depends only on the class of (s_n) in $\lim_{t \to n} {}^1_f S_n(\mathbb{Q})$, distinct classes in $\lim_{t \to n} {}^1_f S_n(\mathbb{Q})$ give nonisomorphic objects in $\mathsf{P}(G_B^{\mathsf{CM}} \to \overline{\Gamma}; P^{\mathsf{Art}})$, and every object in $\mathsf{P}(G_B^{\mathsf{CM}} \to \Gamma; P^{\mathsf{Art}})$ is isomorphic to $P^{\mathsf{CM}}(s) \to P^{\mathsf{Art}}$ for some $s \in \prod_f S_n(\mathbb{Q})$.

Remark 2.26. There remains the problem of characterizing the isomorphism class of $P^{CM} \rightarrow P^{Art}$. One may hope that it is uniquely determined by its isomorphism classes in $P((G_B^{CM})_{/\mathbb{Q}_l} \rightarrow \Gamma_{/\mathbb{Q}_l}, P_{/\mathbb{Q}_l}^{Art})$ for $l = 2, 3, ..., \infty$, i.e., that the kernel of

$$H^1(\mathbb{Q}, {}_fS) \to \prod_l H^1(\mathbb{Q}_l, {}_fS)$$

is zero, but the calculations I have made in this direction do not look promising. Note that ${}_{f}S_{/\mathbb{R}}$ is a countable product of copies of \mathbb{G}_{m} , and so $H^{1}(\mathbb{R}, {}_{f}S) = 0$.

3 Periods of abelian varieties

For a class of abelian varieties \mathcal{A} over \mathbb{Q} , $\langle \mathcal{A} \rangle^{\otimes}$ denotes the category of motives based on the abelian varieties in \mathcal{A} , the projective spaces, and the zero-dimensional varieties. Let

$$G_{\mathbf{B}}^{\mathcal{A}} = \mathcal{A}ut^{\otimes}(\omega_{\mathbf{B}}|\langle \mathcal{A} \rangle^{\otimes}),$$

$$G_{\mathbf{dR}}^{\mathcal{A}} = \mathcal{A}ut^{\otimes}(\omega_{\mathbf{dR}}|\langle \mathcal{A} \rangle^{\otimes}),$$

$$P^{\mathcal{A}} = \mathcal{I}som^{\otimes}(\omega_{\mathbf{B}}, \omega_{\mathbf{dR}}).$$

When $\mathcal{A} = \{A\}$, we write A for \mathcal{A} .

3.1. The inclusion of the Artin motives into $\langle \mathcal{A} \rangle^{\otimes}$ defines a homomorphism

$$G_{\mathbf{B}}^{\mathcal{A}} \to \Gamma.$$

This homomorphism is surjective, and its kernel is the identity component $(G_B^A)^\circ$ of G_B^A (Deligne and Milne 1982, 6.23)⁶. In particular, for a single abelian variety A, there is an exact sequence

$$1 \to \mathrm{MT}(A) \to G^A_{\mathrm{B}} \to \Gamma \to 1$$

3.2. For a single abelian variety A, there is a unique homomorphism

$$S \to \mathrm{MT}(A)^{\mathrm{ab}}_{/\mathbb{R}}$$

sending h_{can} onto $(h^A)^{\text{ab}}$ where, as usual, $h^A \colon \mathbb{S} \to \mathrm{MT}(A)_{/\mathbb{R}}$ is the homomorphism defining the Hodge structure on $H^*(A, \mathbb{Q})$. This homomorphism is surjective, and it factors through S^K if and only if K contains the reflex field of $(\mathrm{MT}(A), h^A)^{\text{ab}}$.

3.3. On combining the last two statements, we see that if A is an abelian variety over \mathbb{Q} such that $S^K \to MT(A)^{ab}$ is an isomorphism for some K, then

$$1 \to \mathrm{MT}(A)^{\mathrm{der}} \to G^A_{\mathrm{B}} \to G^{\mathrm{CM},K}_{\mathrm{B}} \to 1$$

is exact.

Some Hasse principles

We first prove an elementary structure theorem.

Lemma 3.4. Every semisimple group H over \mathbb{Q} such that $H_{/\mathbb{Q}^{dl}}$ is a product of simple groups is isomorphic to a product of groups H_i of the form $H_i = \operatorname{Res}_{F_i/\mathbb{Q}} N_i$ with F_i a number field and N_i an absolutely simple group over F_i .

⁶Which applies because of Deligne's theorem (Deligne 1982a) that all Hodge classes on abelian varieties are absolutely Hodge.

3 PERIODS OF ABELIAN VARIETIES

Proof. To give a semisimple group over \mathbb{Q} is the same as to give a semisimple group H over \mathbb{Q}^{al} together with a descent datum $(\alpha_{\sigma})_{\sigma\in\Gamma}$. Here α_{σ} is an isomorphism $\sigma H \to H$, $\alpha_{\sigma} \circ \sigma \alpha_{\tau} = \alpha_{\sigma\tau}$ for all σ and τ , and there is a continuity condition.

First consider a pair $(H, (\alpha_{\sigma})_{\sigma \in \Gamma})$ with H an adjoint group. Write $H = \prod_{i \in I} H_i$ as a product of simple groups. For each $\sigma \in \Gamma$, there is a permutation (also denoted σ) of I such that α_{σ} is a product of isomorphisms

$$\alpha_{\sigma}(i) \colon \sigma H_i \to H_{\sigma i}.$$

Let J be an orbit of Γ in I, let $j \in J$, and let $\Gamma_j = \{\sigma \in \Gamma \mid \sigma j = j\}$. Then $(\alpha_{\sigma}(j))_{\sigma \in \Gamma_j}$ is a descent datum on H_j , and $(\prod_{i \in J} \alpha_{\sigma}(i))_{\sigma \in \Gamma}$ is a descent datum on $\prod_{i \in J} H_i$. The first defines a model N_j of H_j over $F_j =_{df} \mathbb{Q}^{\Gamma_j}$, which is absolutely simple, and the second defines a model M_J of $\prod_{i \in J} H_i$ over \mathbb{Q} , which is isomorphic to $\operatorname{Res}_{F_j/\mathbb{Q}} N_j$. Now $\prod_{J \in \Gamma \setminus I} M_J$ is a semisimple group over \mathbb{Q} giving rise to $(H, (\alpha_{\sigma})_{\sigma \in \Gamma})$ over \mathbb{Q}^{al} .

Next consider a pair $(H, (\alpha_{\sigma})_{\sigma \in \Gamma})$ with H a product $H = \prod_{i \in I} H_i$ of simple groups. Then $(\alpha_{\sigma})_{\sigma}$ defines a descent datum $(\alpha_{\sigma}^{ad})_{\sigma}$ on H^{ad} , and, as above, α_{σ}^{ad} is a product of isomorphisms $\alpha_{\sigma}^{ad}(i) : \sigma H_i^{ad} \to H_{\sigma i}^{ad}$. Consider

Here $\alpha_{\sigma}(i)$ is the composite

$$\sigma H_i \hookrightarrow \sigma H \xrightarrow{\alpha_{\sigma}} H \xrightarrow{\text{project}} H_i.$$

Because the diagram commutes, α_{σ} and $\prod_{i \in I} \alpha_{\sigma}(i)$ differ by a map from σH into the centre of H, which must be trivial because σH is connected. Thus, $\alpha_{\sigma} = \prod_{i \in I} \alpha_{\sigma}(i)$, and the same argument as in the preceding paragraph completes the proof of the lemma.

We define the *index* (of connectivity) of a semisimple algebraic group H to be the degree of the universal covering $\tilde{H} \to H$. Thus, for an isogeny $a: H' \to H$ of semisimple groups,

$$\operatorname{index}(H') \leq \operatorname{index}(H)$$

with equality if and only if a is an isomorphism.

Consider the following condition on a semisimple algebraic group H over \mathbb{Q} :

(*) $H_{\mathbb{O}^{al}}$ is a product of simple groups of index 1 or 2.

Proposition 3.5. For any semisimple group H over \mathbb{Q} satisfying (*), the map

$$H^1(\mathbb{Q}, H) \to \prod_{l=2,\dots,\infty} H^1(\mathbb{Q}_l, H)$$

is injective.

Proof. According to (3.4), $H \approx \prod \operatorname{Res}_{F_i/\mathbb{Q}} N_i$ with each N_i absolutely simple of index 1 or 2, and

$$H^1(\mathbb{Q}, H) \approx \prod_i H^1(\mathbb{Q}, H_i) \approx \prod_i H^1(F_i, N_i).$$

Therefore, the proposition follows from the next two lemmas.

Lemma 3.6. For any simply connected semisimple group H over a number field F, the map

$$H^1(F,H) \to \prod_{v \text{ real}} H^1(F_v,H)$$

is bijective.

Proof. This is the theorem of Kneser, Harder, and Chernousov — see Platonov and Rapinchuk 1994, Theorem 6.6, p286. \Box

Lemma 3.7. For any semisimple group H of index 2 over a number field F,

$$H^1(F,H) \to \prod_v H^1(F_v,H)$$
 (v runs over all primes of F)

is injective.

Proof. Platonov and Rapinchuk (1994, Remark p337) note that the map has trivial kernel. The lemma can now be proved by a twisting argument, because any form of a semisimple group of index 2 again has index 2. \Box

Proposition 3.8. Let H be a semisimple group over \mathbb{Q} satisfying (*). If two cohomology classes in $H^1(\mathbb{Q}, \tilde{H})$ have the same image in $H^1(\mathbb{R}, H)$, then they have the same image in $H^1(\mathbb{Q}, H)$. In other words, $H^1(\mathbb{Q}, H) \to H^1(\mathbb{R}, H)$ is injective on the image of $H^1(\mathbb{Q}, \tilde{H}) \to H^1(\mathbb{Q}, H)$.

Proof. In the diagram

the rows are exact in the sense that the fibres of the second map are the orbits of the natural action of the first group on the middle set. The first vertical arrow is surjective, and the second is injective. A diagram chase completes the proof. \Box

Remark 3.9. The proof of Proposition 3.8 shows that for any central extension

$$1 \to Z \to G' \to G \to 1$$

of algebraic groups, the map $H^1(\mathbb{Q}, G) \to H^1(\mathbb{R}, G)$ is injective on the image of $H^1(\mathbb{Q}, G')$ in $H^1(\mathbb{Q}, G)$ provided

- (a) the map $H^1(\mathbb{Q}, Z) \to H^1(\mathbb{R}, Z)$ is surjective;
- (b) the map $H^1(\mathbb{Q}, G') \to H^1(\mathbb{R}, G')$ is injective.

Proposition 3.10. Let G be connected reductive group over \mathbb{Q} such that G^{der} is simply connected.

- (a) If $H^1(\mathbb{Q}, G^{ab}) \to \prod_l H^1(\mathbb{Q}_l, G^{ab})$ is injective, then so also is $H^1(\mathbb{Q}, G) \to \prod_l H^1(\mathbb{Q}_l, G)$ (product over $l = 2, 3, 5, ..., \infty$).
- (b) If $H^1(\mathbb{Q}, G^{ab}) \to H^1(\mathbb{R}, G^{ab})$ is injective, then so also is $H^1(\mathbb{Q}, G) \to H^1(\mathbb{R}, G)$.

Proof. (a) See Deligne 1971, 5.12. (Recall that, because G^{der} is simply connected, $H^1(\mathbb{Q}_l, G^{der}) = 0$ for $l \neq \infty$. Using this, we obtain a commutative diagram with exact rows

The image of $G(\mathbb{R})$ in $G^{ab}(\mathbb{R})$ contains its identity component, and the real approximation theorem shows that $G^{ab}(\mathbb{R})$ maps onto $\pi_0(G^{ab}(\mathbb{R}))$. Now a diagram chase shows that any element of $H^1(\mathbb{Q}, G)$ that is locally zero is zero. To show that two elements c and c' of $H^1(\mathbb{Q}, G)$ are equal when they are locally, choose a torsor representing c, and twist the groups by it.)

(b) Similar, but easier.

Characterizing $P^{\text{AV}} \rightarrow P^{\text{CM}}$

Let $AV(\mathbb{Q}) = \langle \mathcal{A} \rangle^{\otimes}$ with \mathcal{A} the class of all abelian varieties over \mathbb{Q} .

Theorem 3.11. Let A be an abelian variety over \mathbb{Q} . If $MT(A)^{der}$ satisfies (*) and $^7 MT(A)^{ab} \cong S^K$ for some K, then the isomorphism class of $P^A \to P^{\mathsf{CM},K}$ is uniquely determined by its classes over \mathbb{Q}_l $(l = 2, 3, ..., \infty)$.

Proof. According to Proposition 1.10, the isomorphism classes in $P(G_B^A \to G_B^{CM,K}; P^{CM,K})$ are classified by $H^1(\mathbb{Q}, H)$ where H is the twist of $Ker(G_B^A \to G_B^{CM,K})$ by the period torsor. But according to (3.3), this kernel is $MT(A)^{der}$, and so the theorem follows from Proposition 3.5.

The next result shows that the abelian varieties over \mathbb{Q} satisfying the conditions of Theorem 3.11 are cofinal among all abelian varieties over \mathbb{Q} for the relation $A \prec B$ if hA is isomorphic to an object of $\langle B \rangle^{\otimes}$.

Proposition 3.12. Let A be an abelian variety over \mathbb{Q} . For any sufficiently large CMsubfield K of \mathbb{C} , there exists an abelian variety B over \mathbb{Q} such that $MT(A \times B)^{der}$ satisfies (*) and $MT(A \times B)^{ab} \cong S^K$.

This will be proved in the next subsubsection.

⁷We don't really need to assume $MT(A)^{ab} \cong S^K$ — it only makes the statement a little more pleasant.

Theorem 3.13. The isomorphism class of $P^{AV} \rightarrow P^{CM}$ is uniquely determined by its classes over \mathbb{Q}_l $(l = 2, 3, ..., \infty)$.

Proof. According to Proposition 1.10, the isomorphism classes in $P(G_B^{AV} \to G_B^{CM}; P^{CM})$ are classified by $H^1(\mathbb{Q}, H)$ where H is the twist of $H' = \text{Ker}(G_B^{AV} \to G_B^{CM})$ by the period torsor. Proposition 3.12 implies that

$$H' \cong \lim \operatorname{MT}(A)^{\operatorname{der}}$$

where A runs over the abelian varieties satisfying the hypothesis of Theorem 3.11. Therefore H is a countable product of simple algebraic groups satisfying (*), and so we can apply Corollary 1.14 and Proposition 3.5. \Box

Proof of Proposition 3.12

We shall say that an algebraic group H over a field k of characteristic zero is a *special* orthogonal group if there exists vector space V over k of dimension n > 8 and a nondegenerate quadratic form q on V such that $H \approx SO(V, q)$. Such an H is a connected and simple, and there are isogenies of degree 2

$$\tilde{H} \to H \to H^{\mathrm{ad}}$$

with \tilde{H} the simply connected covering group of H (a spinor group). Let $Z = Z\tilde{H}$. When n is even $Z(k^{\text{al}}) \approx C_2 \times C_2$.

3.14. Recall from Deligne 1979a the following statements.

- (a) Let A be an abelian variety over C. Each Q-simple factor H of MT(A)^{ad} is of type A, B, C, D^ℝ, or D^ℍ (notations as in ibid. 2.3.8).
- (b) Let A be an abelian variety over \mathbb{C} , and let G = MT(A). The homomorphism $h: \mathbb{S} \to G_{\mathbb{R}}$ defined by the Hodge structure on $H^*(A, \mathbb{Q})$ satisfies the following conditions:
 - i) the Hodge structure on the Lie algebra g of G defined by Ad ∘h: S → GL(g) is of type {(1,-1), (0,0), (-1,1)};
 - ii) adh(i) is a Cartan involution on G^{ad} ;
 - iii) h generates G (i.e., there is no proper closed subgroup $G' \subset G$ such that $h(\mathbb{S}) \subset G'_{\mathbb{R}}$).
- (c) Let H be a simple adjoint group over \mathbb{Q} , and let $h: \mathbb{S}/\mathbb{G}_m \to H_{\mathbb{R}}$ be a homomorphism satisfying the conditions (b) i), ii), iii).
 - i) If H is of type A, B, C, or D^ℝ, there exists an abelian variety A over C such that MT(A)^{der} is simply connected and (MT(A), h^A)^{ad} ≈ (H, h) (apply ibid. 2.3.10).

ii) Suppose H is of type D^H. Then H = Res_{F/Q} N for some absolutely simple group N over a totally real field F, and we let H' = Res_{F/Q} N' where N' → N is the double covering of N that is an inner form of a special orthogonal group (ibid. 2.3.8). Any homomorphism H̃ → MT(A) sending h to (h^A)^{ad} factors through H' (ibid. 1.3.10). There exists an abelian variety A over C with MT(A)^{der} ≈ H' and (MT(A), h^A)^{ad} ≈ (H, h) (apply ibid. 2.3.10).

We shall need the following condition on a semisimple group H over \mathbb{Q} :

(**) *H* is a product of simple groups; a simple factor of $H_{\mathbb{R}}$ has index 2 if it is of type $D^{\mathbb{H}}$ and index 1 otherwise.

Definition 3.15. An abelian variety defined over a subfield of \mathbb{C} is *maximal* if $MT(A)^{der}$ satisfies (**).

Lemma 3.16. A finite product of maximal abelian varieties is maximal.

Proof. Let $A = \prod_j A_j$ where each A_j is maximal. The canonical map $MT(A) \to \prod MT(A_j)$ is injective and its composite with any projection $\prod MT(A_j) \to MT(A_j)$ is surjective (Deligne 1982a, §3). For a judicious choice of simple factors H_i of $\prod MT(A_j)^{der}$, the homomorphism $MT(A)^{ad} \to \prod H_i^{ad}$ will be an isomorphism. Consider

$$\prod \tilde{H}_i \to \operatorname{MT}(A)^{\operatorname{der}} \to \prod H_i.$$

Then $\prod \tilde{H}_i \to MT(A)^{\text{der}}$ factors through $\prod H'_i$ where H'_i is as in (3.14cii) when H_i is of type $D^{\mathbb{H}}$ and equals \tilde{H}_i otherwise. Consider

$$\prod H'_i \twoheadrightarrow \mathrm{MT}(A)^{\mathrm{der}} \twoheadrightarrow \prod H_i.$$

As $\prod H'_i$ and $\prod H_i$ have the same index, the composite is an isomorphism, and so $MT(A)^{der} \rightarrow \prod H_i$ is an isomorphism.

Remark 3.17. Let A and B be abelian varieties such that $MT(A \times B)^{ad} \cong MT(B)^{ad}$. If B is maximal, then so also is $A \times B$. This can be proved the same way as Lemma 3.16 — one only has to observe that, because of the condition on the adjoint groups, the H_i in the proof can be chosen to be factors of $MT(B)^{der}$.

Lemma 3.18. For any abelian variety A over \mathbb{C} , there exists an abelian variety B over \mathbb{C} such that $(MT(B), h^B)^{ad} \approx (MT(A), h^A)^{ad}$ and B is maximal.

Proof. According to (3.14), for each simple factor (H, h) of $(MT(A), h^A)^{ad}$, there exists an abelian variety B(H) such that $(MT(B(H)), h^{B(H)})^{ad} \approx (H, h)$ and $MT(B(H))^{der}$ is the covering H' of H in (3.14cii) if H is of type $D^{\mathbb{H}}$ and is simply connected otherwise. Take $B = \prod B(H)$. Because each B(H) is maximal, so is B (3.16).

Recall that, when $A \prec B$, there is a canonical surjection $c_{A,B} \colon MT(B) \to MT(A)$ carrying h^B to h^A .

Lemma 3.19. An abelian variety A over \mathbb{C} is maximal if and only if, for all B such that $A \prec B$ and $c_{A,B}^{ad}$: $MT(B)^{ad} \to MT(A)^{ad}$ is an isomorphism, $c_{A,B}^{der}$: $MT(B)^{der} \to MT(A)^{der}$ is an isomorphism.

Proof. Let B' be as in (3.18), and let $B = A \times B'$. Then $A \prec B$ and $MT(B)^{ad} \to MT(A)^{ad}$ is an isomorphism, but $MT(B)^{der} \to MT(A)^{der}$ is an isomorphism only if A is maximal. This proves the "if".

Let A be maximal. Then it is clear from (3.14c) that it satisfies the condition.

Lemma 3.20. Let A be an abelian variety over \mathbb{C} , and let τ be an automorphism of \mathbb{C} . Then A is maximal if and only if τA is maximal.

Proof. Recall that $MT(A) = Aut^{\otimes}(\omega_B)$ and $MT(\tau A) = Aut^{\otimes}(\omega_{\tau})$ where ω_{τ} is the fibre functor on $AV(\mathbb{C})$ sending hX to $H^*(\tau X, \mathbb{Q})$. Therefore, from the theory of Tannakian categories, $MT(\tau A) \cong {}^{P}MT(A)$ where $P = \mathcal{I}som(\omega_B, \omega_{\tau})$.

If A is not maximal, then there exists an abelian variety B such that $A \prec B$ and $MT(A)^{ad} \cong MT(B)^{ad}$ but $MT(A)^{der} \ncong MT(B)^{der}$. Clearly τB has the same properties relative to τA , which proves that τA is not maximal.

Lemma 3.21. Let A_1 and A_2 be abelian varieties over \mathbb{C} such that $(MT(A_1), h^{A_1})^{ad} \approx (MT(A_2), h^{A_2})^{ad}$. If A_1 can be defined over \mathbb{Q}^{al} , then so also can A_2 .

Proof. See Blasius and Borovoi 1999, 3.3.

Lemma 3.22. For any CM-subfield K of \mathbb{C} of finite degree over \mathbb{Q} , there exists an abelian variety A over \mathbb{C} such that the canonical map $S^K \to MT(A)$ is an isomorphism. If K is Galois over \mathbb{Q} , then A may be chosen to be defined over \mathbb{Q} .

Proof. ⁸First recall that the CM-types on K generate $X^*(S^K)$: in fact, if $\psi = \tau_1 + \cdots + \tau_g$ is one CM-type on K, then the CM-types $\psi_i = \tau_i + \sum_{j \neq i} \iota \tau_j$ $(i = 1, \ldots, g)$ and $\bar{\phi} = \sum \iota \tau_j$ form a basis for the \mathbb{Z} -module $X^*(S^K)$.

Let *B* be a simple abelian variety over \mathbb{C} of CM-type (E, ϕ) . For each $\rho: E \to \mathbb{Q}^{al}$ and $\tau \in \operatorname{Gal}(\mathbb{Q}^{\operatorname{cm}}/\mathbb{Q})$, define $\psi_{\rho}(\tau) = \phi(\tau^{-1} \circ \rho)$. Then, as ρ runs over the embeddings of *E* into \mathbb{Q}^{al} , ψ_{ρ} runs over a Γ -orbit of CM-types on $\mathbb{Q}^{\operatorname{cm}}$. The map $B \mapsto \{\psi_{\rho}\}$ defines a bijection from the set of isogeny classes of simple abelian varieties of CM-type over \mathbb{Q}^{al} to the set of Γ -orbits of CM-types on $\mathbb{Q}^{\operatorname{cm}}$. A $\tau \in \Gamma$ fixes the reflex field *K* of *B* if and only if $\tau \phi = \phi$. Each ψ_{ρ} is the extension to $\mathbb{Q}^{\operatorname{cm}}$ of a primitive CM-type on *K*, and the kernel of $S^K \to \operatorname{MT}(B)$ is the intersection of the kernels of the ψ_{ρ} (see Milne 1999, §2). For any automorphism σ of \mathbb{C} , σB has reflex field σK .

Choose a finite set of B's with reflex field contained in K such that the corresponding ψ 's generate S^K , and let A be their product. The canonical map $S^K \to MT(A)$ is then an isomorphism. According to (2.7), we may take A to be defined over a subfield k of \mathbb{C} of finite degree over \mathbb{Q} . Then, if K is Galois over \mathbb{Q} , $A_* =_{df} \operatorname{Res}_{k/\mathbb{Q}} A$ has reflex field contained in K, and the canonical map $S^K \to MT(A_*)$ is an isomorphism. \Box

⁸For slightly weaker results, see Milne 1990, I 4.6; Wei 1994, 1.5.1; Borovoi and Blasius 1999, 3.5.

PROOF OF PROPOSITION 3.12. Let A be an abelian variety over \mathbb{Q} . Choose B as in (3.18). After (3.21), we may assume B is defined over a number field F. Then $B_* = \prod_{\sigma \in \Sigma_{F/\mathbb{Q}}} \sigma B$ is defined over \mathbb{Q} . Let K be any CM-subfield of \mathbb{Q}^{al} containing the reflex field of $(MT(A \times B_*), h^{A \times B_*})^{ab}$ and finite and Galois over \mathbb{Q} , and choose C as in (3.22). Consider the abelian variety $A \times B_* \times C$. It is defined over \mathbb{Q} , and (3.20, 3.16, 3.17) show that it is maximal. By assumption, the reflex field of $(MT(A \times B_* \times C), h^{A \times B_* \times C})^{ab}$ is contained K, and so the canonical homomorphism $S \to MT(A \times B_* \times C)^{ab}$ factors through S^K . But $MT(A \times B_* \times C)$ surjects onto $MT(C) \cong S^K$, and so $S^K \cong MT(A \times B_* \times C)^{ab}$. This completes the proof of Proposition 3.12.

Characterizing $P^{\mathcal{H}} \rightarrow P^{\mathsf{CM}}$

Theorem 3.23. Let A be an abelian variety over \mathbb{Q} . If $MT(A)^{der}$ is simply connected and $MT(A)^{ab} \cong S^K$ for some K, then the isomorphism class of $P^A \to P^{\mathsf{CM},K}$ is uniquely determined by its class over \mathbb{R} .

Proof. According to Proposition 1.10 and (3.3), the isomorphism classes in $P(G_B^A \to G_B^{CM,K}; P^{CM,K})$ are classified by $H^1(\mathbb{Q}, H)$, where H is the twist of $MT(A)^{der}$ by the period torsor. Therefore H is simply connected, and so this follows from the theorem of Kneser, Harder, and Chernousov (3.6).

Let \mathcal{H} denote the class of abelian varieties A over \mathbb{Q} such that $MT(A)^{ad}$ has no factor of type $D^{\mathbb{H}}$. Note that, because of (3.14cii), any abelian variety for which $MT(A)^{der}$ is simply connected lies in \mathcal{H} . The next result shows that the abelian varieties satisfying the conditions of (3.23) are cofinal among all abelian varieties in \mathcal{H} .

Proposition 3.24 (Blasius and Borovoi 1999, 3.2). Let A be an abelian variety in \mathcal{H} . For any sufficiently large CM-field K, there exists an abelian variety B over \mathbb{Q} such that $MT(A \times B)^{ab} \cong S^K$ and $MT(A \times B)^{der}$ is simply connected.

Proof. When A is in \mathcal{H} , the proof of Proposition 3.12 can be modified to show that B can be chosen in such a way that $MT(A \times B)^{der}$ is simply connected — in fact, this significantly simplifies the proof.

Theorem 3.25. The isomorphism class of $P^{\mathcal{H}} \to P^{\mathsf{CM}}$ is uniquely determined by its class over \mathbb{R} .

Proof. Similar to that of Theorem 3.13.

Notes. The main theorem of Blasius and Borovoi 1999 (Theorem 1.5) states that the isomorphism class of $P^{\mathcal{H}} \to P^{\mathsf{CM}}$ is determined by the cohomology class of $P^{\mathcal{H}}$ in $\lim_{A \in \mathcal{H}} H^1(\mathbb{R}, G^A_{\mathrm{dR}})$. As noted in the introduction, this statement is false.⁹ Similarly, their

$$H^{1}(\mathbb{Q}, (G_{\mathrm{dR}}^{\mathcal{H}})^{\circ \mathrm{der}}) \to H^{1}(\mathbb{Q}, (G_{\mathrm{dR}}^{\mathcal{H}})^{\circ}) \to \varprojlim_{A \in \mathcal{H}} H^{1}(\mathbb{Q}, (G_{\mathrm{dR}}^{A})^{\circ}),$$

but they are not injective.

⁹They prove (ibid. 5.2) that their conditions determine the cohomology class of $P^{\mathcal{H}}$ in $\lim_{\substack{K \in \mathcal{H} \\ P^{\mathcal{H}} \to P^{\mathsf{CM}}}} H^1(\mathbb{Q}, (G^A_{\mathrm{dR}})^\circ)$, whereas what is needed is that the conditions determine the cohomology class of $P^{\mathcal{H}}$ in $H^1(\mathbb{Q}, (G^A_{\mathrm{dR}})^\circ)$. There are homomorphisms

statement (ibid 1.6) concerning the existence of an analogue of their Theorem 1.5 for the category of all motives is false. The correct statement is Theorem 1.22 of this paper.

Characterizing $P^{AV} \rightarrow P^{CM}$ in terms of a lifting property

Let A be an variety over \mathbb{Q} such that $MT(A) \cong S^K$ for some CM-field K. Recall that $MT(A)^{der} = Ker(G^A \to G_B^{\mathsf{CM},K})$. We say that $P^A \to P^{\mathsf{CM},K}$ has the *lifting property* if there exists a surjective homomorphism $\tilde{G} \to G_B^A$ such that the kernel of $\tilde{G} \to G_B^{\mathsf{CM},K}$ is the universal covering group of $MT(A)^{der}$ and P^A lifts to \tilde{G} .

Theorem 3.26. Let A be an abelian variety over \mathbb{Q} such that $MT(A)^{der}$ satisfies (*) and $S^K \cong MT(A)^{ab}$ for some CM-subfield K of \mathbb{C} . Up to isomorphism, there exists at most one object $P \to P^{\mathsf{CM},K}$ in $\mathsf{P}(G^A \to G_B^{\mathsf{CM},K}; P^{\mathsf{CM},K})$ such that

(a) $P \rightarrow P^{CM,K}$ has the lifting property, and

(b)
$$(P \to P^{\mathsf{CM},K})_{\mathbb{R}} \approx (P^A \to P^{\mathsf{CM},K})_{\mathbb{R}}$$

Proof. Apply Proposition 3.8 to $H = MT(A)^{der}$.

The notion of a lifting property extends in an obvious fashion to infinite sets of abelian varieties.

Theorem 3.27. Up to isomorphism, there exists at most one object $P \to P^{CM}$ in $P(G_B^{AV} \to G_B^{CM}; P^{CM})$ such that

(a) $P \to P^{\mathsf{CM}}$ has the lifting property, and

(b)
$$(P \to P^{\mathsf{CM}})_{\mathbb{R}} \approx (P^{\mathsf{AV}} \to P^{\mathsf{CM}})_{\mathbb{R}}$$

Proof. Similar to the proof of Theorem 3.13.

Remark 3.28. Deligne's hope that all Shimura varieties with rational weight are moduli varieties for motives (Deligne 1979a, p248) implies that $P^{AV} \rightarrow P^{CM}$ has the lifting property over \mathbb{C} . It would be interesting to prove this unconditionally.

Characterizing $P^{\mathsf{AV}} \to P^{\mathsf{Art}}$

Theorem 3.29. Let A be an abelian variety over \mathbb{Q} such that $MT(A)^{ab} \cong S^K$ for some CM-subfield K of \mathbb{C} .

- (a) If $MT(A)^{der}$ is simply connected, then the isomorphism class of $P^A \to P^{Art}$ is uniquely determined by its class over \mathbb{R} .
- (b) If $MT(A)^{der}$ satisfies (*), then, up to isomorphism, there exists at most one object $P \to P^{Art}$ in $P(G_B^A \to \Gamma; P^{Art})$ having the lifting property and isomorphic to $P^A \to P^{Art}$ over \mathbb{R} .

Proof. (a) According to Proposition 1.10, the isomorphism classes in $P(G_B^A \to \Gamma; P^{Art})$ are classified by $H^1(\mathbb{Q}, G)$, where G is the twist of

$$\operatorname{MT}(A) \stackrel{3.1}{=} \operatorname{Ker}(G_{\mathrm{B}}^{A} \to \Gamma)$$

by the period torsor. Then $G^{ab} = {}_{f}S^{K}$, and so G satisfies the hypotheses of (3.10b).

(b) In this case, G satisfies the hypotheses of (3.9).

Remark 3.30. For any abelian variety A over \mathbb{Q} , the map $G^A_{d\mathbb{R}}(\mathbb{R}) \to {}_f\Gamma(\mathbb{R})$ is surjective (Blasius and Borovoi 1999, 4.4), and so

$$H^1(\mathbb{R}, (G^A_{\mathrm{dR}})^\circ) \to H^1(\mathbb{R}, G^A_{\mathrm{dR}})$$

is injective. Therefore, for $P \to P^{\text{Art}}$ in $\mathsf{P}(G^A_{\mathsf{B}} \to \Gamma; P^{\text{Art}})$, the isomorphism class of $P \to P^{\text{Art}}$ over \mathbb{R} is determined by the isomorphism class of P over \mathbb{R} .

Theorem 3.31. (a) There are uncountably many isomorphism classes in $P(G_B^{\mathcal{H}} \to \Gamma; P^{Art})$ that become isomorphic to $P^{\mathcal{H}} \to P^{Art}$ over \mathbb{R} .

(b) If there exists one object $P \to P^{\text{Art}}$ in $\mathsf{P}(G_B^{\text{AV}} \to \Gamma; P^{\text{Art}})$ having the lifting property and isomorphic to $P^{\text{AV}} \to P^{\text{Art}}$ over \mathbb{R} , then there are uncountably many.

Proof. (a) According to Proposition 1.10, the isomorphism classes in $P(G_B^{\mathcal{H}} \to \Gamma; P^{Art})$ are classified by $H^1(\mathbb{Q}, (G_{dR}^{\mathcal{H}})^\circ)$, where $(G_{dR}^{\mathcal{H}})^\circ$ is the twist of

$$(G_{\mathrm{B}}^{\mathcal{H}})^{\circ} \stackrel{3.1}{=} \operatorname{Ker}(G_{\mathrm{B}}^{A} \to \Gamma)$$

by the period torsor. For an $A \in \mathcal{H}$ with $MT(A)^{ab} \cong S^K$, consider

$$0 \to (G^A_{\mathrm{dR}})^{\circ \mathrm{der}}(\mathbb{Q}) \to (G^A_{\mathrm{dR}})^{\circ}(\mathbb{Q}) \to {}_fS^K(\mathbb{Q}) \to H^1(\mathbb{Q}, (G^A_{\mathrm{dR}})^{\circ \mathrm{der}}).$$

Because $(G_{dR}^{\mathcal{H}})^{\circ der}$ is a countable product of algebraic groups,

$$\varprojlim^{1}(G_{\mathrm{dR}}^{A})^{\mathrm{oder}}(\mathbb{Q}) = 0 = \varprojlim^{1} H^{1}(\mathbb{Q}, (G_{\mathrm{dR}}^{A})^{\mathrm{oder}})$$

(by 1.14). Therefore,

$$\varprojlim^1(G^A_{\mathrm{dR}})^{\circ}(\mathbb{Q}) \cong \varprojlim^1{}_f S^K(\mathbb{Q})$$

which is uncountable (2.18). Similarly,

$$\varprojlim^1(G^A_{\mathrm{dR}})^{\circ}(\mathbb{R}) \cong \varprojlim^1{}_f S^K(\mathbb{R}),$$

which is zero because ${}_{f}S_{\mathbb{R}}^{K}$ is a product of copies of \mathbb{G}_{m} . In view of Proposition 1.12, this completes the proof.

(b) Similar.

Remark 3.32. I expect that there are uncountably many distinct isomorphism classes in $P(G_B^{AV} \to \Gamma; P^{Art})$ that become equal to the class of $P^{AV} \to P^{Art}$ over every field \mathbb{Q}_l .

Characterizing *P*^{AV}

Theorem 3.33. Let A be an abelian variety over \mathbb{Q} such that $MT(A)^{ab} \cong S^K$ for some CM-field K.

- (a) If $MT(A)^{der}$ is simply connected, then, up to isomorphism, P^A is the only G^A_B -torsor P such that
 - i) $P \wedge^{G_B^A} \Gamma \approx P^{\text{Art.}}$ and
 - ii) $P_{\mathbb{R}} \approx P^A_{\mathbb{R}}$.
- (b) If $MT(A)^{der}$ satisfies (*), then, up to isomorphism, there exists at most one G_R^A -torsor P such that
 - i) $P \wedge^{G_B^A} \Gamma \approx P^{\operatorname{Art}}$
 - ii) there exists a surjective homomorphism $u: G' \to G^A_B$ such that $u^{der}: G'^{der} \to (G^A_B)^{der}$ is the universal covering group of $(G^A_B)^{der}$ and P = uP' for some G'torsor, and

iii)
$$P_{\mathbb{R}} \approx P_{\mathbb{R}}^A$$
.

Proof. (a) Let P satisfy the conditions (i) and (ii), and choose a morphism $P \to P^{Art}$. Then Theorem 3.29 and Remark 3.30 show that $(P \rightarrow P^{Art}) \approx (P^A \rightarrow P^{Art})$.

(b) Similar to (a).

Other fields

Hasse principles are known to hold for some fields other than number fields. For example, Scheiderer (1996) proves a Hasse principle for the Galois cohomology groups of connected linear algebraic groups over perfect fields with virtual cohomological dimension < 1. However, if the field k is not countable, the affine group scheme G attached to the category of abelian motives over k will not be a countable inverse limit of algebraic groups. In particular, the relation of the flat cohomology group of G to the Galois cohomology groups of its algebraic quotients is unknown, and so (pace Flicker 2001) such results do not imply Hasse principles for period torsors.

References

- ATIYAH, M. F. 1961. Characters and cohomology of finite groups. Inst. Hautes Études Sci. Publ. Math. No. 9, 23–64.
- BLASIUS, D., AND M. BOROVOI. 1999. On period torsors. In Automorphic forms, automorphic representations, and arithmetic (Fort Worth, TX, 1996), 1–8, Proc. Sympos. Pure Math., 66, Part 1, Amer. Math. Soc., Providence, RI.
- BOUSFIELD, A. K., AND D. M. KAN. 1972. Homotopy limits, completions and localizations. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York.
- BREEN, L. 1990. Bitorseurs et cohomologie non abélienne. In The Grothendieck Festschrift, Vol. I, 401–476, Progr. Math., 86, Birkhäuser Boston, Boston, MA.
- BUCUR, I., AND A. DELEANU. 1968. Introduction to the theory of categories and functors. With the collaboration of Peter J. Hilton and Nicolae Popescu. Pure and Applied Mathematics, Vol. XIX Interscience Publication John Wiley & Sons, Ltd., London-New York-Sydney.
- DELIGNE, P. 1971. Travaux de Shimura. Séminaire Bourbaki, 23ème année (1970/71), Exp. No. 389 (Preliminary version).
- DELIGNE, P. 1979a. Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques. In Automorphic forms, representations and *L*-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, pp. 247–289, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I..
- DELIGNE, P. 1979b. Valeurs de fonctions L et périodes d'intégrales. With an appendix by N. Koblitz and A. Ogus. Proc. Sympos. Pure Math., XXXIII, Automorphic forms, representations and Lfunctions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, pp. 313–346, Amer. Math. Soc., Providence, R.I., 1979.
- DELIGNE, P. 1982a. (Notes by J. Milne), Hodge cycles on abelian varieties. In Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics, 900. Springer-Verlag, Berlin-New York, pp9–100.
- DELIGNE, P., 1982b. Motifs et groupe de Taniyama. In Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics, 900. Springer-Verlag, Berlin-New York, pp261–279.
- DELIGNE, P., AND J.S. MILNE. 1982. Tannakian categories. In Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics, 900. Springer-Verlag, Berlin-New York, pp101–228.
- FLICKER, Y. Z. 2001. Motivic torsors. Israel J. Math. 122, 61-77.
- GIRAUD, J. 1971. Cohomologie non abélienne. Die Grundlehren der mathematischen Wissenschaften, Band 179. Springer-Verlag, Berlin-New York.
- GRAY, B. I. 1966. Spaces of the same *n*-type, for all *n*. Topology 5, 241–243.
- GROTHENDIECK, A. 1966. On the de Rham cohomology of algebraic varieties. Inst. Hautes Études Sci. Publ. Math. No. 29, 95–103.
- MILNE, J. S. 1988. Automorphic vector bundles on connected Shimura varieties. Invent. Math. 92, no. 1, 91–128.
- MILNE, J. S. 1990. Canonical models of (mixed) Shimura varieties and automorphic vector bundles. Automorphic forms, Shimura varieties, and *L*-functions, Vol. I (Ann Arbor, MI, 1988), 283–414, Perspect. Math., 10, Academic Press, Boston, MA.
- MILNE, J. S. 1999. Lefschetz motives and the Tate conjecture. Compositio Math. 117, no. 1, 45-76.

- PLATONOV, V. AND A. RAPINCHUK. 1994. Algebraic groups and number theory. Translated from the 1991 Russian original by Rachel Rowen. Pure and Applied Mathematics, 139. Academic Press, Inc., Boston, MA.
- SCHEIDERER, C. 1996. Hasse principles and approximation theorems for homogeneous spaces over fields of virtual cohomological dimension one. Invent. Math. 125, no. 2, 307–365.
- SERRE, J.-P. 1964. Cohomologie galoisienne. Cours au Collège de France, 1962-1963. Seconde édition. Lecture Notes in Mathematics 5 Springer-Verlag, Berlin-Heidelberg-New York.
- SERRE, J.-P. 1992. Topics in Galois theory. Lecture notes prepared by Henri Damon [Henri Darmon]. With a foreword by Darmon and the author. Research Notes in Mathematics, 1. Jones and Bartlett Publishers, Boston, MA.
- SERRE, J.-P. 1994. Propriétés conjecturales des groupes de Galois motiviques et des représentations *l*-adiques. In Motives (Seattle, WA, 1991), 377–400, Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI.
- TATE, J. 1976. Relations between K_2 and Galois cohomology. Invent. Math. 36, 257–274.
- WEI, W. 1994. Moduli fields of CM-motives applied to Hilbert's 12-th problem, Preprint http://www.mathematik.uni-bielefeld.de/sfb343/preprints/pr94070.ps.gz.
- WINTENBERGER, J.-P. 1990. Torseurs pour les motifs et pour les représentations *p*-adiques potentiellement de type CM, Math. Ann. 288, no. 1, 1–8.

2679 Bedford Rd, Ann Arbor, MI 48104, USA. E-mail: math@jmilne.org; Home page: www.jmilne.org/math/