On a Theorem of Lenstra and Schoof

B.V. Petrenko¹

Department of Mathematics, University of Illinois, 1409 West Green Street, Urbana, Illinois 61801, U.S.A.

Abstract

We give a detailed proof of Theorem 1.15 from a well-known paper "Primitive normal bases for finite fields" by H.W. Lenstra Jr. and R.J. Schoof. We are not aware of any other proofs. Let L/K be a finite-dimensional Galois field extension and B the set of all normal bases of this extension. Theorem 1.15 describes the group of all γ in the multiplicative group of L such that $\gamma B = B$.

Key words: Normal basis, Wedderburn's Theorems, group ring, primitive normal basis.

PACS: 12F10, 16K20, 20C05, 11T30, 12E20

1 Introduction

The celebrated paper of Lenstra and Schoof [6] solved a difficult problem. It proved that every extension of finite fields has a primitive normal basis. This means that for any extension of finite fields $\mathbb{F}_{q^m}/\mathbb{F}_q$, there exists $\alpha \in \mathbb{F}_{q^m}$ such that α generates the multiplicative group $\mathbb{F}_{q^m}^*$ of \mathbb{F}_{q^m} , and $\{\alpha^{q^j} \mid j \in \{1, ..., m\}\}$ is an \mathbb{F}_q -basis of \mathbb{F}_{q^m} . Several particular cases have been done earlier by Carlitz in [1] and [2], and Davenport in [5]. Statement 1.15 is a step in the proof of the main result of [6]. We recall this statement below for the convenience of the reader. We call it Theorem 1 for reference purposes.

Email address: petrenko@uiuc.edu (B.V. Petrenko).

¹ I thank Hendrik Lenstra and Everett Dade for their ideas and help.

Theorem 1 Let $K \subseteq L$ be a finite extension of fields with Galois group G. Let

$$B = \{ \alpha \in L \mid (\tau(\alpha))_{\tau \in G} \text{ is a basis of } L \text{ over } K \},$$

and denote by w the number |G|th roots of unity in K^* . Then for $\gamma \in L^*$ the following assertions are equivalent:

- (i) $\gamma B \subseteq B$.
- (ii) $\gamma B = B$.
- (iii) $\frac{\tau(\gamma)}{\gamma} \in K^*$ for all $\tau \in G$.
- (iv) $\gamma^w \in K^*$.

The set of all $\gamma \in L^*$ satisfying these conditions is a subgroup of L^* containing K^* and C/K^* is isomorphic to the group of all homomorphisms $G \to K^*$.

This theorem is stated in [6] without a proof and used there in the case when L is a finite field. Because we are not aware of any proof of this result, we prove it below using the Wedderburn's Theorems (see, for example, [3]). This is the main result of the present paper. It is established in Section 4. Our proof is based on a private communication of Lenstra [6].

2 Notation

If W is a set, we denote its cardinality by |W|.

F is a field

 F^* is the multiplicative group of F

A is a finite-dimensional associative algebra over F with a two-sided identity 1_A . We assume that $0_A \neq 1_A$.

- J(A) is the Jacobson radical of A.
- U(A) is the unit group of A, i.e. the set of $u \in A$ having two-sided multiplicative inverses u^{-1} .
- nil(A) the set of all nilpotent elements of A.

H is a hyperplane in A, i.e. an F-subspace of A of codimension one.

 \mathbb{F}_2 is a field with two elements.

 $M_n(R)$ is the ring of $n \times n$ matrices with entries in a ring R.

 $E_{i,j}$ is the matrix in $M_n(R)$ all of whose entries are 0_R except the (i,j)-entry which is 1_R . It will always be clear what R and n are in any particular situation.

 ωFG is the augmentation ideal of the group algebra FG, where F is a field and G is a group.

Tr is the map $\operatorname{Tr}: L \to K, t \mapsto \sum_{\tau \in G} \tau(t)$.

N is the kernel of the map Tr, defined above.

 \mathbb{N} is the set of positive integers.

3 Preliminary Results

We begin by stating three main theorems of this section. Their proofs are based on a private communication of Dade [4].

Theorem 2 If H is a hyperplane in an F-algebra A, then $H \cap U(A) \neq \emptyset$ except possibly when either

- (i) H is a two-sided ideal of A, or
- (ii) |F| = 2 and A has a factor algebra isomorphic to $\mathbb{F}_2 \oplus \mathbb{F}_2 \oplus \mathbb{F}_2$.

Theorem 3 If H is a hyperplane in an F-algebra A, then $H \setminus U(A) \neq \emptyset$ except possibly when

(iii) |F| = 2 and A has a factor algebra isomorphic to $\mathbb{F}_2 \oplus \mathbb{F}_2$.

Theorem 4 If H is a hyperplane in the group algebra FG of a finite group G over a field F, then $H \cap U(A) \neq \emptyset$ and $H \setminus U(A) \neq \emptyset$ except when H is a two-sided ideal of A.

We see that Theorems 2 and 3 imply Theorem 4 except possibly when |F| = 2. Lemma 5 below shows that the group algebra \mathbb{F}_2G of any group G has only one two-sided ideal of codimension one, namely $\omega \mathbb{F}_2G$. Therefore, Theorems 2 and 3 imply Theorem 4 when |F| = 2 also. **Lemma 5** The group algebra \mathbb{F}_2G of any group G has only one two-sided ideal of codimension one, namely $\omega \mathbb{F}_2G$.

PROOF. Let I be a two-sided ideal of \mathbb{F}_2G with $\dim_{\mathbb{F}_2}\mathbb{F}_2G/I=1$. Then

$$\mathbb{F}_2 G \xrightarrow{\pi, \text{onto}} \mathbb{F}_2 G/I \xrightarrow{\epsilon, \cong} \mathbb{F}_2$$

where $\pi(t) = t + I$, $t \in \mathbb{F}_2G$, and ε is the unique field isomorphism. We see that $G \subseteq U(\mathbb{F}_2G)$ and $\varepsilon \circ \pi(U(\mathbb{F}_2G)) = \{1\}$. Therefore, $\pi(g) = \pi(1)$ for any $g \in G$, i.e. $g - 1 \in I$. Since $\omega \mathbb{F}_2G = \sum_{g \in G} \mathbb{F}_2(g - 1)$, we have that $\omega \mathbb{F}_2G \subseteq I$. We know that $\omega \mathbb{F}_2G$ is a maximal two-sided ideal of \mathbb{F}_2G . Therefore, $I = \omega \mathbb{F}_2G$.

A key step in proving Theorems 2 and 3 is the following

Lemma 6 Let H be a hyperplane in an F-algebra A. We do not assume that $dim_F A$ is finite in this lemma. Suppose that either $U(A) \subseteq H$ or $U(A) \cap H = \emptyset$, then $nil(A) \subseteq H$.

PROOF.

- (1) Suppose that $U(A) \subseteq H$. Then $1_A \in H$. If $t \in \text{nil}(A)$, then $1_A + t \in U(A) \subseteq H$. Hence $t = (1_A + t) 1_A \in H$.
- (2) Suppose that $U(A) \cap H = \emptyset$. Then $1_A \notin H$. Suppose that there exists $t \in \operatorname{nil}(A) \setminus H$. Since $\dim_F A/H = 1$, we see that $1_A + H$ and t + H are F-linearly dependent. Therefore, $1_A + \alpha t \in H$ for some $\alpha \in F$. At the same time $\alpha t \in \operatorname{nil}(A)$, so that $1_A + \alpha t \in U(A)$. We conclude that $U(A) \cap H \neq \emptyset$, a contradiction.

Next we prove Theorem 2 for simple algebras.

Lemma 7 Let H be a hyperplane in a simple F-algebra A such that $H \cap U(A) = \emptyset$. Then $A \cong F$ and $H = \{0\}$.

PROOF. By Wedderburn's Theorems, $A \cong M_n(D)$ for some division F-algebra D.

If n = 1, then $A \cong D$, so that $U(A) = A \setminus \{0\}$. Since $H \cap U(A) = \emptyset$, we see that $H = \{0\}$. Then $\dim_F A/H = 1$ implies that $A \cong F$.

Next we show that $n \geq 2$ cannot occur. Suppose not. Let $\varphi : A \to M_n(D)$ be an isomorphism of F-algebras. By Lemma 6, $\operatorname{nil}(M_n(D)) \subseteq \varphi(H)$. Hence $E_{i,j} \in \varphi(H)$ for all $i \neq j$. Therefore,

$$M = E_{n,1} + E_{1,2} + \dots + E_{n-1,n} = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \dots & \ddots & \vdots \\ 0 & \dots & \dots & 0 & 1 \\ 1 & 0 & \dots & \dots & 0 \end{pmatrix} \in \varphi(H).$$

However, M is a permutation matrix and therefore is invertible in $M_n(F) \subseteq M_n(D)$. Since $\varphi^{-1}(M) \subseteq H$, we conclude that $H \cap U(A) \neq \emptyset$, contradicting initial assumption.

Theorem 3 for simple algebras is

Lemma 8 No hyperplane in any simple F-algebra A can contain U(A).

PROOF. Suppose not. Let A be a simple F-algebra and H a hyperplane in A such that $U(A) \subseteq H$. Let $\varphi : A \to M_n(D)$ be an isomorphism of F-algebras, for some $n \in \mathbb{N}$ and a division F-algebra D.

If n = 1, then $A \cong D$ and $A \setminus \{0\} = U(A) \subseteq H$. This is impossible.

Let us consider the case $n \geq 2$. By Lemma 6, $\operatorname{nil}(M_n(D)) \subseteq \varphi(H)$. Therefore, $E_{i,j} \in \varphi(H)$ for all $i \neq j$. We also note that

$$E_{i,i} + E_{i,j} - E_{j,i} - E_{j,j} \in \varphi(H)$$

because the square of this matrix is zero. Thus

$$E_{i,i} - E_{j,j} = (E_{i,i} + E_{i,j} - E_{j,i} - E_{j,j}) - E_{i,j} + E_{j,i} \in \varphi(H).$$

Define $tr: M_n(D) \to D, (d_{i,j}) \mapsto \sum_{l=1}^n d_{l,l}$. Then the above discussion shows that $Ker(tr) \subseteq \varphi(H)$. Hence, $\varphi(H) = tr^{-1}(H_1)$ for some F-hyperplane H_1 of D. Let $d \in D \setminus H_1$. Then the matrix

$$Q = (dE_{1,1} + E_{n,1}) + E_{1,2} + E_{2,3} + \dots + E_{n-1,n} = \begin{pmatrix} d & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & 0 & 1 \\ 1 & 0 & \dots & \dots & 0 \end{pmatrix}$$

has the two-sided inverse

$$Q^{-1} = (E_{1,2} - dE_{2,n}) + E_{2,1} + E_{3,2} + \dots + E_{n,n-1} = \begin{pmatrix} 0 & 0 & 0 & \dots & 1 \\ 1 & 0 & 0 & \dots & -d \\ 0 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & \ddots & 0 & 0 \\ 0 & 0 & \dots & \dots & 1 & 0 \end{pmatrix}$$

and $tr(Q) = d \notin H_1$, so that $Q \in \varphi(U(A) \setminus H)$. Hence $\varphi^{-1}(Q) \in U(A) \setminus H$, a contradiction.

Now we are ready to prove Theorems 2 and 3. They hold when $\dim_F A = 1$. Below we prove these theorems simultaneously by induction on $\dim_F A$.

PROOF OF THEOREMS 2 AND 3. We can assume that $\dim_F A \geq 2$ and that Theorems 2 and 3 hold for all strictly smaller values of $\dim_F A$.

Assume that Theorem 2 is false for some hyperplane H in A. Then $H \cap U(A) = \emptyset$, H is not a two-sided ideal of A, and A does not satisfy (ii).

We claim that $J(A)=\{0\}$. Suppose not. By Lemma 6, $\mathrm{nil}(A)\subseteq H$. So $J(A)\subseteq H$. Define

$$\pi: A \to A' = A/J(A), a \mapsto a + J(A).$$

Put $H' = \pi(H)$. Since $J(A) \subseteq H$, we see that that $H = \pi^{-1}(H')$ and H' is a hyperplane in A'. Theorem 2 holds for A' and H'. Let $u' \in H' \cap U(A')$. Then u' = u + J(A) for some $u \in H$. At the same time, $u \in U(A)$ because J(A) is nilpotent. Therefore $H \cap U(A) \neq \emptyset$, a contradiction.

By Wedderburn's Theorems, $A = A_1 \dotplus ... \dotplus A_n$, where A_i are two-sided ideals each of which is a simple F-subalgebra of $A, n \in \mathbb{N}$.

If n=1, then $A=A_1$ is simple. Since $H\cap U(A)=\emptyset$, Lemma 7 implies that $H=\{0\}$ and therefore H is a two-sided ideal of A. This contradicts our assumptions. Hence $n\geq 2$.

We claim that $A_i \nsubseteq H$ for all $i \in \{1, ..., n\}$. Suppose not. Then $A_1 \subseteq H$ without loss of generality. Define $A' = \sum_{j=2}^n A_j$, $H' = H \cap A'$ and

$$\pi: A \to A', a_1 + ... + a_n, a_i \in A_i, i \in \{1, ..., n\}.$$

We see that $H' = \pi(H)$ is a hyperplane of A' and $Ker(\pi) = A_1$. As before, H' is not a two-sided ideal of A', and A' does not satisfy (ii). Since $\dim_F A' < \dim_F A$, by inductive hypothesis, there exists $u' \in H' \cap U(A') \subseteq H$. Choose any $u_1 \in U(A_1)$. Then $u = u' + u_1 \in U(A) \cap H = \emptyset$, a contradiction.

Because $A_n \nsubseteq H$, we see that $H_n = H \cap A_n$ is a hyperplane in A_n . Because $A'' = A_1 \dotplus ... \dotplus A_{n-1} \nsubseteq H$, we see that $H'' = H \cap A''$ is a hyperplane in A''. If there exists $u_n \in H_n \cap U(A_n)$ and $u'' \in H'' \cap U(A'')$, then $u = u_n + u'' \in H \cap U(A) = \emptyset$, a contradiction. So either $H_n \cap U(A_n) = \emptyset$ or $H'' \cap U(A'') = \emptyset$.

Suppose that $H_n \cap U(A_n) = \emptyset$. Then $A_n \cong F$ by Lemma 7. Therefore $A = A'' \dotplus A_n \cong A'' \oplus F$. Since A does not satisfy (ii), it follows that A'' does not satisfy (iii). Theorem 3 holds for A'' and A'' because $\dim_F A'' < \dim_F A$. Therefore, there exists $u'' \in U(A'') \setminus H''$. Since $\dim_F A/H = 1$ and $u'' \notin H$, we have that A/H = (u'' + H)F. Let $u_n \in A_n \setminus H_n$. Then $u_n \notin H$, so that $A/H = (u_n + H)F$. Since u'' + H and $u_n + H$ are F-linearly dependent, we have that $u'' + \alpha u_n \in H$ for some $0 \neq \alpha \in F$. However, $A_n \cong F$ implies that $\alpha u_n \in U(A_n)$. Then $u'' + \alpha u_n \in U(A'') + U(A_n) = U(A)$. It follows that $U(A) \cap H \neq \emptyset$, a contradiction. We conclude that $H'' \cap U(A'') = \emptyset$.

Lemma 8 gives us some $u_n \in U(A_n) \setminus H_n$. Let $u'' \in U(A'')$. Because $u'', u_n \in A \setminus H$, there is $0 \neq \alpha \in F$ such that $u = u'' + \alpha u_n \in H$. Then $u \in U(A) \cap H = \emptyset$, a contradiction. We conclude that Theorem 2 holds for A and H.

It remains to prove Theorem 3 for A and H. Assume it is false. Then $U(A) \subseteq H$ and A does not satisfy (iii). By Lemma 6, $\operatorname{nil}(A) \subseteq H$. As in the above proof, this implies that $J(A) = \{0\}$, so that A is the internal direct sum $A_1 \dotplus \dots \dotplus A_n$ of $n \in \mathbb{N}$ two-sided ideals A_i each of which is a simple F-subalgebra of A. As in the above proof, $A_i \nsubseteq H$ for all $i \in \{1, ..., n\}$. Moreover, $n \geq 2$ by Lemma 8.

The intersections $H_n = H \cap A_n$ and $H'' = H \cap A''$ are now hyperplanes in A_n and $A'' = A_1 \dotplus ... \dotplus A_{n-1}$, respectively. Lemma 8 gives us some $u_n \in U(A_n) \backslash H_n$, and hence $u_n \notin H$. If there is some $u'' \in H'' \cap U(A'')$, then $u = u_n + u'' \in U(A) \backslash H$. This contradicts our assumption that $U(A) \subseteq H$. Hence $H'' \cap U(A'') = \emptyset$. Because A does not satisfy (iii), its epimorphic image A'' does not satisfy (ii). So Theorem 2, for the algebra A'' with $\dim_F A'' < \dim_F A$, tells us that H'' is a two-sided ideal of A''. Since $\dim_F A''/H'' = 1$, we conclude that one of the direct summands $A_1, ..., A_{n-1}$ of A'', say A_{n-1} , is isomorphic to F, and that $H'' = A_1 \dotplus ... \dotplus A_{n-2}$. But no A_i is contained in H and $H'' \subseteq H$. Therefore $H'' = \{0\}$, n = 2, and $A'' = A_{n-1} = A_1 \cong F$.

We claim that $A_n \cong F$ and $H_n = \{0\}$. If there exists $u_n \in H_n \cap U(A_n)$, then $u'' + u_n \in U(A) \setminus H$ for any $0 \neq u'' \in A''$. This contradicts our assumption that $U(A) \subseteq H$. Hence $H_n \cap U(A_n) = \emptyset$. Then Lemma 7 implies that $A_n \cong F$ and $H_n = \{0\}$.

Now $A = A'' \dotplus A_n \cong F \oplus F$. If $|F| \geq 3$, then no one-dimensional subspace of $F \oplus F$ can contain $U(F \oplus F)$. Hence |F| = 2, so that $A \cong \mathbb{F}_2 \oplus \mathbb{F}_2$. Thus A satisfies (iii), a contradiction. This contradiction shows that A must satisfy Theorems 2 and 3. So the simultaneous proof of these theorems is complete.

4 Main Result

PROOF OF THEOREM 1. We note that once we prove that (i) \iff (ii) \iff (iii) \iff (iv), then we can conclude that the groups C/K^* and $\operatorname{Hom}_{\mathbb{Z}}(G, K^*)$ are isomorphic via the map $(\gamma K^*)(\tau) = \frac{\tau(\gamma)}{\gamma}, \gamma \in C$. This map is surjective because the group $\operatorname{H}^1(G, L^*)$ is trivial (see, for example, [8]).

We see that (ii) \Longrightarrow (i).

Next we prove that (i) \Longrightarrow (ii). We begin by making the following observations.

- (1) If $\gamma_1, \gamma_2 \in L$ are such that $\gamma_1 B, \gamma_2 B \subseteq B$, then $B \supseteq \gamma_1 B \supseteq \gamma_1 \gamma_2 B$.
- (2) If $\alpha \in K^*$, then $\alpha B = B$.
- (3) If $\tau \in G$, then $\tau(B) = B$.

Let $\gamma_1 = \gamma$ and $\gamma_2 = \prod_{1 \neq \tau \in G} \tau(\gamma)$. Then $\gamma_1 B, \gamma_2 B \subseteq B$, and at the same time, $\gamma_1 \gamma_2 B = B$ since $\gamma_1 \gamma_2 \in K^*$. Therefore, $B \supseteq \gamma_1 B \supseteq \gamma_1 \gamma_2 B = B$.

We have proved that (i) \iff (ii).

We now show that (iii) \Longrightarrow (iv). Let $\gamma_{\tau} = \frac{\tau(\gamma)}{\gamma}$, then $\gamma_{\tau} \in K^*$. Let $S_{\gamma} = \{\gamma_{\tau} \mid \tau \in G\}$. Then $\gamma_{\tau_1}\gamma_{\tau_2} = \gamma_{\tau_1\circ\tau_2}$ for all $\tau_1, \tau_2 \in G$. Therefore, S_{γ} is a group under multiplication. It is a subgroup of the group

$$S = \{ \mu \in K^* \mid \mu^{|G|} = 1 \}$$

of order w. Hence, by Lagrange's Theorem, it would be sufficient to show that $\gamma^{|S_{\gamma}|} \in K^*$. This is indeed the case: $1 = \gamma_{\tau}^{|S_{\gamma}|} = \frac{\tau(\gamma^{|S_{\gamma}|})}{\gamma^{|S_{\gamma}|}}$ for any $\tau \in G$.

Next we show that (iv) \Longrightarrow (iii). From (iv) we conclude that $\left(\frac{\tau(\gamma)}{\gamma}\right)^w = 1$ for any $\tau \in G$. We claim that $\frac{\tau(\gamma)}{\gamma} \in S$ (since $S \subseteq K^*$, we would be done). Indeed, each element of S is a solution of the equation $x^w - 1 = 0$. This equation has at most w solutions in any field extension of K. On the other hand, |S| = w, so that S is the set of all solutions of this equation in any field extension of K. Because $\left(\frac{\tau(\gamma)}{\gamma}\right)^w = 1$, we conclude that $\frac{\tau(\gamma)}{\gamma} \in S$.

We have proved that (iii) \iff (iv).

We note that (iii) \Longrightarrow (i) because any element of K is fixed by any element of G and the map

$$\hat{\gamma}: L \to L, t \mapsto t\gamma$$

is a bijection.

Finally, we show that (ii) \Longrightarrow (iii). This would prove that

$$(i) \Longleftrightarrow (ii) \Longleftrightarrow (iii) \Longleftrightarrow (iv).$$

We know that $B \neq \emptyset$ by the Normal Basis Theorem (see, for example, [8]). Let $a \in B$. Define the map

$$\widetilde{a}: KG \to L, \sum_{\tau \in G} \alpha_{\tau} \tau \mapsto \sum_{\tau \in G} \alpha_{\tau} \tau(a).$$

Then

- (1) \tilde{a} is an isomorphism of KG-modules.
- (2) $\tilde{a}(\omega KG) = N$.
- (3) There exists a unique K-linear map Γ which makes the following diagram commutative:

$$\begin{array}{ccc}
L & \xrightarrow{\widehat{\gamma}} & L \\
\widetilde{a} & & \widetilde{a} \\
KG & \xrightarrow{\Gamma} & KG.
\end{array}$$

We claim that γN is a KG-submodule of L. Because $\tilde{a}^{-1}(\gamma N) = \Gamma(\omega KG)$, Theorem 4 would imply our claim if $\Gamma(\omega KG) \cap U(KG) = \emptyset$. Suppose that $\Gamma(\omega KG) \cap U(KG) \neq \emptyset$. Let $r \in \Gamma(\omega KG) \cap U(KG)$. Then

- (1) $\{\tau r \mid \tau \in G\}$ is a K-basis of KG.
- (2) $\tilde{a}(r) = \gamma \tilde{a}(r')$, where $r' \in \omega KG$ is such that $r = \Gamma(r')$.

Therefore, $\{\tau(\tilde{a}(r)) \mid \tau \in G\}$ is a K-basis of L. On the other hand, since $\tilde{a}(r') \in N$, a proper KG-submodule of L, we see that $\{\tau(\tilde{a}(r')) \mid \tau \in G\}$ is not a K-basis of L. We conclude that $\gamma^{-1}B \nsubseteq B$, contradicting (ii). Hence, we have proved that γN is a KG-submodule of L.

Because $N, \gamma N$ are KG-submodules of L, we conclude that, for any $\tau \in G$,

$$\gamma N = \tau(\gamma N) = \tau(\gamma)\tau(N) = \tau(\gamma)N.$$

Therefore, N is a vector space over the field $K' = K(\frac{\tau(\gamma)}{\gamma})$. Consequently, $\dim_K K'$ divides $\dim_K N = \dim_K L - 1$. Because L is a K'-vector space, we conclude that $\dim_K K'$ divides $\dim_K L$. Hence $\dim_K K'$ divides $\gcd(\dim_K L - 1, \dim_K L) = 1$, i.e. $K(\frac{\tau(\gamma)}{\gamma}) = K$ for any $\tau \in G$. The proof is complete.

References

- [1] Carlitz L., Primitive roots in a finite field, Trans. Amer. Math. Soc. 73 (1952), pp. 373-382.
- [2] Carlitz L., Some problems involving primitive roots in a finite field, Proc. Nat. Acad. Sci. U.S.A., 38 (1952) pp. 314-318, 618.
- [3] Cohn P.M., Algebra, Vol. 2, John Wiley & Sons, 1989.
- [4] Dade E.C., Private communication (2002).
- [5] Davenport H., Bases for finite fields, J. London Math. Soc., 43 (1968), pp. 21-39, 44 (1969), p.378.
- [6] Lenstra H.W. Jr. and Schoof R.J., Primitive normal bases for finite fields, Mathematics of Computation, 48 (1987), Number 177, pp. 217-231.
- [7] Lenstra H.W. Jr., Private communication (2002).
- [8] Milne J.S., Fields and Galois Theory (http://www.jmilne.org/math/).