AN EXTENSION THEOREM FOR SEPARATELY MEROMORPHIC FUNCTIONS WITH PLURIPOLAR SINGULARITIES

MAREK JARNICKI AND PETER PFLUG

ABSTRACT. Let $D_j \subset \mathbb{C}^{n_j}$ be a pseudoconvex domain and let $A_j \subset D_j$ be a locally pluriregular set, $j=1,\ldots,N$. Put

$$X := \bigcup_{j=1}^{N} A_1 \times \ldots \times A_{j-1} \times D_j \times A_{j+1} \times \ldots \times A_N.$$

Let $M \subset X$ be relatively closed. For any $j \in \{1, \ldots, N\}$ let Σ_j be the set of all $(z', z'') \in (A_1 \times \ldots \times A_{j-1}) \times (A_{j+1} \times \ldots \times A_N)$ such that the fiber $M_{(z', \cdot, z'')} := \{z_j \in \mathbb{C}^{n_j} : (z', z_j, z'') \in M\}$ is not pluripolar. Assume that $\Sigma_1, \ldots, \Sigma_N$ are pluripolar. Put

$$X' := \bigcup_{j=1}^{N} \{ (z', z_j, z'') \in (A_1 \times \ldots \times A_{j-1}) \times D_j \times (A_{j+1} \times \ldots \times A_N) :$$
$$(z', z'') \notin \Sigma_j \}.$$

Then (Theorem 1.3) there exists a relatively closed pluripolar subset $\widehat{M} \subset \widehat{X}$ of the 'envelope of holomorphy' \widehat{X} of X such that:

- $\bullet \ \widehat{M} \cap X' \subset M,$
- every function f separately meromorphic on $X \setminus M$ (Definition 1.2) extends to a (uniquely determined) function \widehat{f} meromorphic on $\widehat{X} \setminus \widehat{M}$,
- if f is separately holomorphic on $X \setminus M$, then \widehat{f} is holomorphic on $\widehat{X} \setminus \widehat{M}$, and
- \widehat{M} is singular with respect to the family of all functions \widehat{f} . The case of separately holomorphic functions was solved in [Jar-Pfl 2002b]. In the case where $N=2, M=\varnothing$, the above result will be strengthened in Theorem 1.4.

1. Introduction. Main results.

We keep the main notation from [Jar-Pfl 2002b]:

¹⁹⁹¹ Mathematics Subject Classification. 32D15, 32D10.

The first author was supported in part by the KBN grant No. 5 P03A 033 21. The paper was finished during the RiP stay of both authors at Mathematisches Forschungsinstitut in Oberwolfach. We like to thank these institutions.

• Let $N \in \mathbb{N}$, $N \geq 2$, and let $\emptyset \neq A_j \subset D_j \subset \mathbb{C}^{n_j}$, where D_j is a domain, $j = 1, \ldots, N$. We define an N-fold cross

$$X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$$

$$:= \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times D_j \times A_{j+1} \times \dots \times A_N \subset \mathbb{C}^{n_1 + \dots + n_N} = \mathbb{C}^n.$$

• For an open set $\Omega \subset \mathbb{C}^n$ and $A \subset \Omega$ let

$$h_{A,\Omega} := \sup\{u : u \in \mathcal{PSH}(\Omega), u \leq 1 \text{ on } \Omega, u \leq 0 \text{ on } A\},\$$

where $\mathcal{PSH}(\Omega)$ is the set of all functions plurisubharmonic on Ω . Put

$$\omega_{A,\Omega} := \lim_{k \to +\infty} h_{A \cap \Omega_k,\Omega_k}^*,$$

where $(\Omega_k)_{k=1}^{\infty}$ is a sequence of relatively compact open sets $\Omega_k \subset \Omega_{k+1} \in \Omega$ with $\bigcup_{k=1}^{\infty} \Omega_k = \Omega$ (h^* denotes the upper semicontinuous regularization of h).

• For an N-fold cross $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$ put

$$\widehat{X} := \{(z_1, \dots, z_N) \in D_1 \times \dots \times D_N : \sum_{j=1}^N \omega_{A_j, D_j}(z_j) < 1\}.$$

- We say that a subset $\emptyset \neq A \subset \mathbb{C}^n$ is locally pluriregular if $h_{A \cap \Omega, \Omega}^*(a) = 0$ for any $a \in A$ and for any open neighborhood Ω of a.
- Suppose that $S_j \subset (A_1 \times ... \times A_{j-1}) \times (A_{j+1} \times ... \times A_N), j = 1, ..., N$. Define the generalized N-fold cross

$$T = \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; S_1, \dots, S_N) := \bigcup_{j=1}^N \{ (z', z_j, z'') \in (A_1 \times \dots \times A_{j-1}) \times D_j \times (A_{j+1} \times \dots \times A_N) : (z', z'') \notin S_j \}.$$
 (1)

• Let $M \subset T$ be a relatively closed set. We say that a function $f: T \setminus M \longrightarrow \mathbb{C}$ is separately holomorphic $(f \in \mathcal{O}_s(T \setminus M))$ if for any $j \in \{1, \ldots, N\}$ and $(a', a'') \in (A_1 \times \ldots \times A_{j-1}) \times (A_{j+1} \times \ldots \times A_N) \setminus S_j$ the function $f(a', \cdot, a'')$ is holomorphic in the open set $D_j \setminus M_{(a', \cdot, a'')}$, where $M_{(a', \cdot, a'')} := \{z_j \in \mathbb{C}^{n_j} : (a', z_j, a'') \in M\}$ (2). Notice that the definition applies to the case where T = X is an N-fold cross $(S_1 = \cdots = S_N = \varnothing)$.

The following general extension theorem for separately holomorphic functions with singularities was proved in [Jar-Pfl 2002a] and [Jar-Pfl 2002b].

Theorem 1.1. Let $D_j \subset \mathbb{C}^{n_j}$ be a pseudoconvex domain, let $A_j \subset D_j$ be a locally pluriregular set, j = 1, ..., N, and let $M \subset X$ be a relatively closed subset of the N-fold cross $X := \mathbb{X}(A_1, ..., A_N; D_1, ..., D_N)$. Assume that for each $j \in \{1, ..., N\}$

⁽¹⁾ Observe that $\mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N) = \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; \varnothing, \dots, \varnothing)$. Moreover, if N = 2, then $\mathbb{T}(A_1, A_2; D_1, D_2; S_1, S_2) = \mathbb{X}(A_1 \setminus S_2, A_2 \setminus S_1; D_1, D_2)$.

⁽²⁾ Observe that the above condition is empty if $M_{(a',\cdot,a'')} = D_j$.

the set $\Sigma_j = \Sigma_j(M)$ of all points $(z', z'') \in (A_1 \times \ldots \times A_{j-1}) \times (A_{j+1} \times \ldots \times A_N)$ such that the fiber $M_{(z', \ldots z'')}$ is not pluripolar is pluripolar. Put

$$X' = X'(M) := \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; \Sigma_1, \dots, \Sigma_N).$$

Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:

- $\widehat{M} \cap X' \subset M$,
- for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X' \setminus M$,
 - \widehat{M} is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$.

In particular, $\hat{X} \setminus \hat{M}$ is the envelope of holomorphy of $X \setminus M$ with respect to the space of separately holomorphic functions.

Moreover:

- (a) if M is pluripolar, then $\Sigma_1, \ldots, \Sigma_N$ are pluripolar (3),
- (b) if $M = X \cap \widetilde{M}$, where \widetilde{M} is an analytic subset of an open connected neighborhood of X, then \widehat{M} is analytic,
- (c) if $M = X \cap \widetilde{M}$, where \widetilde{M} is an analytic subset of \widehat{X} , then \widehat{M} is the union of all pure (n-1)-dimensional irreducible components of \widetilde{M} (4).

Some special cases of the above theorem were studied by many authors — see the references in [Jar-Pfl 2002b].

It is known that the envelope of holomorphy (of any Riemann domain over \mathbb{C}^n) coincides with the envelope of meromorphy (cf. [Jar-Pfl 2000], Th. 3.6.6). Thus it is natural to conjecture that in the above situation the domain $\widehat{X}\setminus \widehat{M}$ is also the envelope of meromorphy of $X\setminus M$ with respect to separate meromorphic functions. The case $M=\varnothing$ was studied in [Sak 1957], [Kaz 1976], [Kaz 1978], [Kaz 1984], [Shi 1986], and [Shi 1989].

Definition 1.2. Let $T = \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; S_1, \dots, S_N)$ be a generalized N-fold cross. Let $M \subset T$, $S \subset T \setminus M$ be relatively closed. We say that a function $f: (T \setminus M) \setminus S \longrightarrow \mathbb{C}$ is separately meromorphic on $T \setminus M$ $(f \in \mathcal{M}_s(T \setminus M))$ if for any $j \in \{1, \dots, N\}$ and $(a', a'') \in (A_1 \times \dots \times A_{j-1}) \times (A_{j+1} \times \dots \times A_N) \setminus S_j$ with $(M \cup S)_{(a', \cdot, a'')} \neq D_j$, there exists a function $f(a', \cdot, a'') \in \mathcal{M}(D_j \setminus M_{(a', \cdot, a'')})$ such that $f(a', \cdot, a'') = f(a', \cdot, a'')$ on $D_j \setminus (M \cup S)_{(a', \cdot, a'')}$.

Observe that $f \in \mathcal{O}_s(T \setminus (M \cup S))$ (5).

The main results of the paper are the following two theorems.

Theorem 1.3. Let $(A_j, D_j)_{j=1}^N$, X, M, and \widehat{M} be as in Theorem 1.1. Let $S \subset X \setminus M$ be relatively closed and let $f: (X \setminus M) \setminus S \longrightarrow \mathbb{C}$ be a separate meromorphic function on $X \setminus M$ such that

(*) the sets $\Sigma_1(S), \ldots, \Sigma_N(S)$ are pluripolar.

⁽³⁾ And, consequently, the assumption of the theorem is always satisfied for pluripolar sets.

 $[\]widehat{M} = \emptyset$ In particular, $M = \emptyset \Longrightarrow \widehat{M} = \emptyset$.

⁽⁵⁾ Note that $M \cup S$ is relatively closed in T.

Put $Q_f := M \cup S$. Then there exists exactly one $\widehat{f} \in \mathcal{M}(\widehat{X} \setminus \widehat{M})$ such that:

- $\hat{f} \in \mathcal{O}(\hat{X} \setminus \hat{Q}_f)$, where the set \hat{Q}_f is constructed via Theorem 1.1 (in the same way as \widehat{M} for M) (6),
 - $\widehat{f} = f$ on $X'_f \setminus Q_f$, where

$$X_f' := \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; \Sigma_1(Q_f), \dots, \Sigma_N(Q_f)).$$

Consequently, the envelope of $X \setminus M$ with respect to separately meromorphic functions satisfying (*) coincides with its envelope of separate holomorphy.

In the case where N=2, $M=\varnothing$, the above result may be strengthened as follows.

Theorem 1.4. Let $D \subset \mathbb{C}^p$, $G \subset \mathbb{C}^q$ be pseudoconvex domains, let $\emptyset \neq A \subset D$, $\emptyset \neq B \subset G$ be locally pluriregular sets, and let

$$X := \mathbb{X}(A, B; D, G) = (A \times G) \cup (D \times B).$$

Let $S \subset X$ be a relatively closed set. Assume that:

(1.4.1) for every $(a,b) \in A \times B$ we have $\operatorname{int}_{\mathbb{C}^q} S_{(a,\cdot)} = \emptyset$, $\operatorname{int}_{\mathbb{C}^p} S_{(\cdot,b)} = \emptyset$,

 $(1.4.2) A \times B \subset \overline{(A \times B) \setminus S} (7)$

there exist exhaustions $(D_j)_{j=1}^{\infty}$ and $(G_j)_{j=1}^{\infty}$ of D and G, respectively, such that: (1.4.3) D_j , G_j are relatively closed pseudoconvex subdomains of D and G, respectively,

(1.4.4) $A_j := A \cap D_j \neq \emptyset$, $B_j := B \cap G_j \neq \emptyset$, (1.4.5) for every $(a,b) \in A_j \times B_j$ we have $B_j \setminus S_{(a,\cdot)} \neq \emptyset$, $A_j \setminus S_{(\cdot,b)} \neq \emptyset$,

Then for every function $f: X \setminus S \longrightarrow \mathbb{C}$ which is separately meromorphic on X there exists a function $\hat{f} \in \mathcal{M}(\hat{X})$ such that $\hat{f} = f$ on $X \setminus S$.

2. Auxiliary results.

Remark 2.1. (a) ([Kli 1991], Corollary 4.8.4) If $A, B \subset \mathbb{C}^n$ are plurithin at a point $a \in \mathbb{C}^n$ (8), then $A \cup B$ is plurithin at a.

- (b) ([Arm-Gar 2001], Th. 7.2.2) Every polar set $P \subset \mathbb{C}$ is thin at any point $a \in \mathbb{C}$.
- (c) If $A \subset \mathbb{C}$ is not thin at a point $a \in \overline{A}$, then for any polar set $P \subset \mathbb{C}$, the set $A \setminus P$ is not thin at a ((c) follows directly from (a) and (b)).
- (d) If $A \subset \mathbb{C}^n$ is locally pluriregular at a point $a \in \overline{A}$, then A is not plurithin at a. If $A \subset \mathbb{C}$ is not thin at a point $a \in \overline{A}$, then A is locally regular at a.

⁽⁶⁾ Note that $\widehat{M} \subset \widehat{Q}_f$.

⁽⁷⁾ In particular, for every $(a,b) \in A \times B$ and for every neighborhood $U \subset \mathbb{C}^p \times \mathbb{C}^q$ of (a,b)the set $(A \times B) \cap U \setminus S$ is not pluripolar.

⁽⁸⁾ We say that a set $A \subset \mathbb{C}^n$ is plurithin at a point $a \in \mathbb{C}^n$ if either $a \notin \overline{A}$ or $a \in \overline{A}$ and $\limsup_{A\setminus\{a\}\ni z\to a} u(z) < u(a)$ for a function u plurisubharmonic in a neighborhood of a.

Indeed, suppose that $A \subset \mathbb{C}^n$ is locally pluriregular at a and

$$\lim_{A \setminus \{a\} \ni z \to a} u(z) < c < u(a)$$

for some $u \in \mathcal{PSH}(V)$, where V is an open neighborhood of a. We may assume that $u \leq 0$ on V. Take an open neighborhood $U \subset V$ of a such that u < c on $(A \setminus \{a\}) \cap U$. Put $v := \frac{u}{-c} + 1$. Then $v \leq 1$ on U and $v \leq 0$ on $(A \setminus \{a\}) \cap U$. Hence $v \leq h^*_{(A \setminus \{a\}) \cap U, U} = h^*_{A \cap U, U}$ on U. In particular, $0 = v(a) = \frac{u(a)}{-c} + 1 < 0$; contradiction.

Now, suppose that $A \subset \mathbb{C}$ is not thin at a and $h^*_{A\cap U,U}(a) > 0$ for some neighborhood U of a. Let $P \subset U$ be a polar set such that $h^*_{A\cap U,U} = h_{A\cap U,U}$ on $U \setminus P$ (cf. [Jar-Pfl 2000] Th. 2.1.41). In particular, $h^*_{A\cap U,U} = 0$ on $A \setminus P$. By (c), the set $A \setminus P$ is not thin at a. Hence $0 < h^*_{A\cap U,U}(a) = \limsup_{A \setminus P \ni z \to a} h^*_{A\cap U,U}(z) = 0$; contradiction.

- (e) ([Arm-Gar 2001], Th. 7.3.9) If $A \subset \mathbb{C}$ is thin at a point $a \in \overline{A}$, then there is a sequence $r_k \searrow 0$ such that $\{z \in A : |z a| = r_k\} = \emptyset, k = 1, 2, \dots$
- (f) ([Bed-Tay 1982], Corollary 10.5) For a non-pluripolar set $A \subset \mathbb{C}^n$ let A^* denote the set of all $a \in \overline{A}$ such that A is locally pluriregular at a. Then $A \setminus A^*$ is pluripolar.

3. Corollaries from Theorem 1.4.

Let E denote the unit disc. For $a \in \mathbb{C}^k$, r > 0, let $\Delta_a(r) = \Delta_a^k(r)$ be the polydisc with center at a and the radius r.

Corollary 3.1 (Cf. [Sak 1957]). Let $S \subset E \times E$ be a relatively closed set such that:

- int $S = \emptyset$,
- for every domain $U \subset E \times E$ the set $U \setminus S$ is connected $\binom{9}{}$. Let A (resp. B) denote the set of all $a \in E$ (resp. $b \in E$) such that $\operatorname{int}_{\mathbb{C}} S_{(a,\cdot)} = \emptyset$ (resp. $\operatorname{int}_{\mathbb{C}} S_{(\cdot,b)} = \emptyset$). Put $X := \mathbb{X}(A,B;E,E) = (A \times E) \cup (E \times B)$.

Then for every function $f: X \setminus S \longrightarrow \mathbb{C}$ which is separately meromorphic on X, there exists an $\widehat{f} \in \mathcal{M}(E \times E)$ such that $\widehat{f} = f$ on $X \setminus S$.

Remark 3.2. Notice that the original proof of the above result is not correct: the proof of Theorem 1 in [Sak 1957] contains an essential gap. Namely, on p. 78 the author claims that for any domain $U \subset E \times E$ the set $(A \times B) \cap U \setminus S$ contains an open polydisc. The following example shows that this is in general impossible.

Let $(\mathbb{Q} + i\mathbb{Q}) \cap E = \{q_1, q_2, \dots\}$, $S := \bigcup_{k=1}^{\infty} \{q_k\} \times \overline{\Delta}_{1-1/k}(1/k^2) \subset E \times E$. Then S satisfies all the assumptions of Corollary 3.1 but in this case the interior of $A = E \setminus (\mathbb{Q} + i\mathbb{Q})$ is empty.

Proof. First we check that the sets A and B are not thin at any point of E (in particular, they are dense in E).

⁽⁹⁾ We shortly say that S does not separate domains.

Indeed, suppose that A is thin at a point $a \in E$. By Remark 2.1(e), there exist a circle $C \subset E$ such that $C \cap A = \emptyset$. Using a Baire category argument, we conclude that there exist a non-empty open arc $\Gamma \subset C$ and an open disc $\Delta \subset E$ such that the 3-dimensional real surface $\Gamma \times \Delta$ is contained in S. Hence, since S is nowhere dense and does not separate domains, we get a contradiction.

Consequently, by Remark 2.1(d), the sets A and B are locally regular and $h_{A,E}^* = h_{B,E}^* = 0$. In particular, $\widehat{X} = E \times E$.

Now, using the fact that A and B are dense in E, one can easily check that all the assumptions of Theorem 1.4 (D = G = E) are satisfied with arbitrary exhaustions $D_j := \Delta_0(r_j), G_j := \Delta_0(r_j), 0 < r_j \nearrow 1$, which satisfy condition

Remark 3.3. (a) E. Sakai claims in [Sak 1957] that also the following n-dimensional version of Corollary 3.1 is true. We do not know how to prove it.

Let $S \subset E^n$ be relatively closed such that int $S = \emptyset$ and S does not separate domains. Let $f: E^n \setminus S \longrightarrow \mathbb{C}$ be such that for any $j \in \{1, \ldots, n\}$ and for any $(a', a'') \in E^{j-1} \times E^{n-j}$ for which $\operatorname{int}_{\mathbb{C}} S_{(a', \cdot, a'')} = \emptyset$, the function $f_{(a', \cdot, a'')}$ extends meromorphically to $E^{(10)}$. Then f extends meromorphically to E^n .

In particular, we would like to ask whether for any set $A \subset E^k$ which is plurithin at $0 \in E^k$ there exists a non-empty relatively open subset Γ of a real hypersurface such that $\Gamma \subset E^k \setminus A$ (cf. the proof of Corollary 3.1).

(b) We also do not know whether the following generalization of Corollary 3.1 is true.

Let $D \subset \mathbb{C}^p$, $G \subset \mathbb{C}^q$ be pseudoconvex domains and let $S \subset D \times G$ be a relatively closed set such that int $S = \emptyset$ and S does not separate domains. Let A (resp. B) denote the set of all $a \in D$ (resp. $b \in G$) such that $\operatorname{int}_{\mathbb{C}^q} S_{(a,\cdot)} = \emptyset$ (resp. $\operatorname{int}_{\mathbb{C}^p} S_{(\cdot,b)} = \varnothing$). Put $X := \mathbb{X}(A,B;D,G) = (A \times G) \cup (D \times B)$. Then for every function $f: X \setminus S \longrightarrow \mathbb{C}$ which is separately meromorphic on X, there exists an $\widehat{f} \in \mathcal{M}(D \times G)$ such that $\widehat{f} = f$ on $X \setminus S$.

Corollary 3.4 (Cf. [Shi 1989], Th. 2). Let D, G, A, B, X be as in Theorem 1.4. Assume that $S \subset X$ is a relatively closed set such that

- the set $D \setminus A$ is of zero Lebesgue measure,
- for every $a \in A$ the fiber $S_{(a,\cdot)}$ is pluripolar,

• for every $b \in B$ the fiber $S_{(\cdot,b)}$ is of zero Lebesgue measure. Then for every function $f: X \setminus S \longrightarrow \mathbb{C}$ which is separately meromorphic on X, there exists an $\widehat{f} \in \mathcal{M}(D \times G)$ such that $\widehat{f} = f$ on $X \setminus S$.

Proof. One can easily check that all the assumptions of Theorem 1.4 are satisfied (with arbitrary exhaustions satisfying (1.4.3-4)). It remains to observe that $h_{A,D}^* \equiv 0$ (because $h_{A,D}^* = 0$ on A and the set $D \setminus A$ is of zero measure). Hence $\widehat{X} = D \times G.$

⁽¹⁰⁾ That is, f is separately meromorphic on the n-fold generalized cross T := $\mathbb{T}(E,\ldots,E;E,\ldots,E;S_1,\ldots,S_n)$, where S_j denote the set of all $(a',a'')\in E^{j-1}\times E^{n-j}$ for which int_C $S_{(a',\cdot,a'')} \neq \emptyset$, j = 1, ..., n; cf. Definition 1.2.

4. Rothstein Theorem.

Theorem 4.1 (Cf. [Rot 1950]). Let $f \in \mathcal{M}(E^p \times E^q)$. Assume that $A \subset E^p$ be a locally pluriregular set such that for any $a \in E^p$ we have $(P_f)_{(a,\cdot)} \neq E^q$, where P_f denote the polar set of f, i.e. P_f is the union of the set of all poles of f and the set of all indeterminancy points of f (11). Let $G \subset \mathbb{C}^p$ be a domain such that $E^q \subset G$. Assume that for every $a \in A$ the function $f(a,\cdot)$ extends meromorphically to G. Then there exists an open neighborhood Ω of $(E^p \times E^q) \cup (A \times G)$ and a function $\widehat{f} \in \mathcal{M}(\Omega)$ such that $\widehat{f} = f$ on $E^p \times E^q$.

We present a sketch of the proof.

(1) The case where $A = E^p$ (12), q = 1, $G = \Delta_0(R)$ (R > 1), and $f \in \mathcal{O}(E^p \times E)$:

The proof may be found for instance in [Siu 1974].

(2) The case where $A = E^p$, q = 1, and $G = \Delta_0^q(R)$:

Recall that $(P_f)_{(a,\cdot)} \neq E^q$ for any $a \in E^p$, and therefore, for any $a \in E^p$ there exists a $b \in E^q$ such that f is holomorphic in a neighborhood of (a,b). By applying locally (1), we get the required result.

(3) The case where $A = E^p$ and $G = \Delta_0^q(R)$:

Let R_0 denote the radius of the maximal polydisc $\Delta_0^q(R_0)$ such that f extends meromorphically to $E^p \times \Delta_0^q(R_0)$. We only need to show that $R_0 \geq R$. Obviously $R_0 \geq 1$. Suppose that $R_0 < R$.

Let S_q be the set of all $(z,w') \in E^p \times \Delta_0^{q-1}(R_0)$ such that $(P_f)_{(z,w',\cdot)} = E$. It is well known that S_q is an analytic subset of $E^p \times \Delta_0^{q-1}(R_0)$. Moreover, our assumptions imply that $S_q \neq E^p \times \Delta_0^{q-1}(R_0)$. Applying locally the Rothstein theorem to $(E^p \times \Delta_0^{q-1}(R_0) \setminus S_q) \times \Delta_0(R) \subset \mathbb{C}^{p+q-1} \times \mathbb{C}$, we conclude that f extends meromorphically to $((E^p \times \Delta_0^{q-1}(R_0) \setminus S_q) \times \Delta_0(R)) \cup (E^p \times \Delta_0^q(R_0))$. Observe that, by the Levi extension theorem ([Jar-Pfl 2000], Prop. 3.4.5), the envelope of holomorphy of $((E^p \times \Delta_0^{q-1}(R_0) \times \Delta_0(R)) \setminus (S_q \times \Delta_0(R))) \cup (E^p \times \Delta_0^q(R_0))$ equals $E^p \times \Delta_0^{q-1}(R_0) \times \Delta_0(R)$. Consequently, the function f extends meromorphically to $E^p \times \Delta_0^{q-1}(R_0) \times \Delta_0(R)$. Repeating the same argument with respect to other variables in \mathbb{C}^q , we conclude that f extends meromorphically to the domain $E^p \times H$, where

$$H = \bigcup_{j=1}^{q} \Delta_0^{j-1}(R_0) \times \Delta_0(R) \times \Delta_0^{q-j}(R_0).$$

The envelope of holomorphy of $E^p \times H$ has the form $E^p \times \widehat{H}$, where \widehat{H} contains a polydisc $\Delta_0^q(R_0')$ with $R_0' > R_0$. Thus f extends meromorphically to $E^p \times \Delta_0^q(R_0')$; contradiction — cf. the proof of Lemma 12 in [Jar-Pfl 2002b].

⁽¹¹⁾ Note that P_f is analytic and $f \in \mathcal{O}(E^p \times E^q \setminus P_f)$.

 $[\]binom{12}{2}$ Observe that if $A = E^p$, then we have to prove that f extends meromorphically to $E^p \times G$.

(4) The case where $A \subset E^p$ is locally pluriregular and $G = \Delta_0^q(R)$:

For every $z \in E^p$, let $\rho_f(z)$ denote the radius of the maximal polydisc $\Delta_0^q(r)$ such that $f(z,\cdot)$ extends meromorphically to $\Delta_0^q(r)$. Obviously, $\rho_f \geq 1$ on E^p and $\rho_f \geq R$ on A.

Using (3), one can easily conclude that f extends meromorphically to the Hartogs domain

$$D := \{ (z, w) \in E^p \times \mathbb{C}^q : |w| < (\rho_f)_*(z) \}.$$

Let $\widetilde{f} \in \mathcal{M}(D)$ be the meromorphic extension of f.

Moreover, $-\log(\rho_f)_* \in \mathcal{PSH}(E^p)$.

Indeed, let \widehat{D} denote the envelope of holomorphy of D. It is known that $\widehat{D} \subset E^p \times \mathbb{C}^q$ is a Hartogs domain with complete q-circled fibers ([Jar-Pfl 2000], Remark 3.1.2(h)). Moreover, \widetilde{f} extends meromorphically to \widehat{D} ([Jar-Pfl 2000], Th. 3.6.6). In particular,

$$(\rho_f)_*(z) = \inf\{\delta_{\widehat{D},(0,\xi)}(z,0) : \xi \in \mathbb{C}^q, |\xi| = 1\}, z \in E^p,$$

where

$$\delta_{\widehat{D},(0,\xi)}(z,0) = \sup\{r > 0 : (z,0) + \Delta_0(r)(0,\xi) \subset \widehat{D}\}.$$

Consequently, $-\log(\rho_f)_* \in \mathcal{PSH}(E^p)$ ([Jar-Pfl 2000], Th. 2.2.9(iv)).

Thus $-\log(\rho_f)_* \in \mathcal{PSH}(E^p)$. Recall that $\rho_f \geq R$ on A. Hence, using the local pluriregularity of A, we conclude that $(\rho_f)_* \geq R$ on A (13). Thus $A \times \Delta_0^q(R) \subset D$, and therefore D is the required neighborhood.

(5) The general case where $A \subset E^p$ is locally pluriregular and G is arbitrary: Fix an $a \in A$. Let G_0 denote the set of all $b \in G$ such that there exist $r_b > 0$ and $f_b \in \mathcal{M}(\Delta_{(a,b)}(r_b))$, $\Delta_{(a,b)}(r_b) \subset E^p \times G$, such that:

$$\forall_{\alpha \in A \cap \Delta_a(r_b)} : f_b(\alpha, \cdot) = \widetilde{f(\alpha, \cdot)} \text{ on } \Delta_b(r_b) \ (^{14}).$$

Obviously G_0 is open, $G_0 \neq \emptyset$ ($E^q \subset G_0$). Using the Rothstein theorem with $G = \Delta_0^q(R)$, one can prove that G_0 is closed in G. Thus $G_0 = G$.

Moreover, one can also prove that if $\Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''}) \neq \emptyset$, then $f_{b'} = f_{b''}$ on $\Delta_{(a,b')}(r_{b'}) \cap \Delta_{(a,b'')}(r_{b''})$. This gives a meromorphic extension of f to an open neighborhood of $\{a\} \times G$. Since a was arbitrary, we get the required neighborhood Ω .

The proof of the Rothstein theorem is completed.

⁽¹³⁾ Suppose that $h_{A,E^p}^* = h_{A,E^p}$ on $E^p \setminus P$, where P is pluripolar. Put $u := \frac{-\log(\rho_f)_*}{\log R} + 1$. Then $u \le 1$ and $u \le 0$ on $A \setminus P$. Consequently, $u \le h_{A \setminus P,E^p}^* = h_{A,E^p}^*$. In particular, $u \le 0$ on A, i.e. $(\rho_f)_* \ge R$ on A.

⁽¹⁴⁾ As before, $\widetilde{f(\alpha,\cdot)}$ denotes the meromorphic extension of $f(\alpha,\cdot)$.

5. Proof of Theorem 1.3.

Fix a function $f \in \mathcal{M}_s(X \setminus M) \cap \mathcal{O}_s(X \setminus Q_f)$. By Theorem 1.1 there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{Q}_f)$ with $\widehat{f} = f$ on $X'_f \setminus Q_f$. It remains to prove that $\widehat{f} \in \mathcal{M}(\widehat{X} \setminus \widehat{M})$.

It is sufficient to prove that $\widehat{f} \in \mathcal{M}(\Omega \setminus \widehat{M})$, where $\Omega \subset \widehat{X}$ is an open neighborhood of X'_f .

Indeed, by virtue of Lemma 9 from [Jar-Pfl 2002b] and the Chirka theorem (cf. [Chi 1993], see also [Jar-Pfl 2002b], Th. 6), the envelope of holomorphy of $\Omega \setminus \widehat{M}$ coincides with $\widehat{X} \setminus \widehat{M}$. Consequently, the function \widehat{f} extends meromorphically to $\widehat{X} \setminus \widehat{M}$ (cf. [Jar-Pfl 2000], Th. 3.6.6).

Fix a $j \in \{1, ..., N\}$ and a point

$$(a', a'') \in (A_1 \times \ldots \times A_{j-1}) \times (A_{j+1} \times \ldots \times A_N) \setminus \Sigma_j(Q_f).$$

Take an $a_j \in D_j \setminus (Q_f)_{(a',a'')}$ and let r > 0 be such that $\Delta_a(r) \subset \widehat{X} \setminus \widehat{Q}_f$, where $a = (a', a_j, a'')$. Take a $D_j' \in D_j \setminus \widehat{M}_{(a',a'')}$ with $a_j \in D_j'$. We may assume that $\Delta_{(a',a'')}(r) \times D_j' \subset \widehat{X} \setminus \widehat{M}$ and $\Delta_{a_j}(r) \subset D_j'$. By the Rothstein theorem 4.1 with $p := n_1 + \dots + n_{j-1} + n_{j+1} + \dots + n_N$, $q := n_j$,

$$A := ((A_1 \times \ldots \times A_{i-1}) \times (A_{i+1} \times \ldots \times A_N)) \cap \Delta_{(a',a'')}(r),$$

we get an open set $\Omega_a \supset A \times D'_j$ such that \hat{f} extends meromorphically to Ω_a . The proof of Theorem 1.3 is completed.

6. Proof of Theorem 1.4.

It suffices to prove that for each j there exists an open neighborhood Ω_j of the cross $X_j := \mathbb{X}(A_j, B_j; D_j, G_j) = (A_j \times G_j) \cup (D_j \times B_j)$ such that there exists an $\widetilde{f}_j \in \mathcal{M}(\Omega_j)$ with $\widetilde{f}_j = f$ on $X_j \setminus S$.

Indeed, we may assume that $\Omega_j \subset \widehat{X}_j$. Observe that $\widehat{X}_j \nearrow \widehat{X}$. By Lemma 9 from [Jar-Pfl 2002b] the envelope of holomorphy of Ω_j equals \widehat{X}_j . Hence, by Theorem 3.6.6 from [Jar-Pfl 2000], the function \widehat{f}_j extends to a function $\widehat{f}_j \in \mathcal{M}(\widehat{X}_j)$. Since $X_j \setminus S$ is not pluripolar (by (1.4.2)), we conclude that $\widehat{f}_j = \widehat{f}_{j+1}$ on \widehat{X}_j . Finally, we glue up the functions $(\widehat{f}_j)_{j=1}^{\infty}$ and we get the required extension.

Fix $(a,b) \in A_j \times B_j \setminus S$ and let r > 0 be such that $\Delta_{(a,b)}(r) \subset D_j \times G_j \setminus S$. Define $Y := \mathbb{X}(A \cap \Delta_a(r), B \cap \Delta_b(r); \Delta_a(r), \Delta_b(r))$. Then $f \in \mathcal{O}_s(Y)$ and hence, by Theorem 1.1, $f|_Y$ extends holomorphically on \widehat{Y} . In particular, f extends holomorphically to an open neighborhood of (a,b).

By the Rothstein theorem 4.1, we get an open set

$$\Omega_{i,a,b} = (\Delta_a(r_{a,b}) \times G_i) \cup (D_i \times \Delta_b(r_{a,b})) \subset D_i \times G_i$$

for which there exists a function $\widehat{f}_{j,a,b} \in \mathcal{M}(\Omega_{j,a,b})$ such that $\widehat{f}_{j,a,b} = f$ on $X \cap \Omega_{j,a,b} \setminus S$.

Now we show that if $\Omega_{j,a,b} \cap \Omega_{j,a',b'} \neq \emptyset$, then $\widehat{f}_{j,a,b} = \widehat{f}_{j,a',b'}$ on $\Omega_{j,a,b} \cap \Omega_{j,a',b'}$. Observe that

$$\begin{split} \Omega_{j,a,b} \cap \Omega_{j,a',b'} &= \left(\Delta_a(r_{a,b}) \cap \Delta_{a'}(r_{a',b'}) \right) \times G_j \\ &\quad \cup \Delta_a(r_{a,b}) \times \Delta_{b'}(r_{a',b'}) \\ &\quad \cup \Delta_{a'}(r_{a',b'}) \times \Delta_b(r_{a,b}) \\ &\quad \cup D_j \times \left(\Delta_b(r_{a,b}) \cap \Delta_{b'}(r_{a',b'}) \right). \end{split}$$

First observe that $\widehat{f}_{j,a,b} = f = \widehat{f}_{j,a',b'}$ on $(A_j \times B_j) \cap (\Delta_a(r_{a,b}) \times \Delta_{b'}(r_{a',b'})) \setminus S$. Hence, by (1.4.2), $\widehat{f}_{j,a,b} = \widehat{f}_{j,a',b'}$ on $\Delta_a(r_{a,b}) \times \Delta_{b'}(r_{a',b'})$. The same argument works on $\Delta_{a'}(r_{a',b'}) \times \Delta_b(r_{a,b})$.

If $\Delta_a(r_{a,b}) \cap \Delta_{a'}(r_{a',b'}) \neq \emptyset$, then for any $\beta \in B_j$ we have $\widehat{f}_{j,a,b}(\cdot,\beta) = f(\cdot,\beta)$ on $A_j \cap \Delta_a(r_{a,b}) \setminus S_{(\cdot,\beta)}$. Hence $\widehat{f}_{j,a,b}(\cdot,\beta) = \widehat{f}(\cdot,\beta)$ on $\Delta_a(r_{a,b})$, and, consequently, $\widehat{f}_{j,a,b}(\cdot,\beta) = \widehat{f}_{(\cdot,\beta)}(\cdot,\beta) = \widehat{f}_{j,a',b'}(\cdot,\beta)$ on $\Delta_a(r_{a,b}) \cap \Delta_{a'}(r_{a',b'})$ for any $\beta \in B_j$. The identity principle implies that $\widehat{f}_{j,a,b} = \widehat{f}_{j,a',b'}$ on $(\Delta_a(r_{a,b}) \cap \Delta_{a'}(r_{a',b'})) \times G_j$. The same argument works on $D_j \times (\Delta_b(r_{a,b}) \cap \Delta_{b'}(r_{a',b'}))$.

It remains to observe that, by (1.4.5), $\Omega_j := \bigcup_{(a,b) \in A_j \times B_j \setminus S} \Omega_{j,a,b}$ is an open neighborhood of X_j .

The proof of Theorem 1.4 is completed.

References

[Arm-Gar 2001] D. H. Armitage, S. J. Gardiner, Classical Potential Theory, Springer Verlag

[Bed-Tay 1982] E. Bedford, B. A. Taylor, A new capacity for plurisubharmonic functions, Acta Math. 149 (1982), 1–40.

[Chi 1993] E. M. Chirka, The extension of pluripolar singularity sets, Proc. Steklov Inst. Math. 200 (1993), 369–373.

[Jar-Pfl 2000] M. Jarnicki, P. Pflug, Extension of Holomorphic Functions, de Gruyter Expositions in Mathematics 34, Walter de Gruyter 2000.

[Jar-Pfl 2002a] M. Jarnicki, P. Pflug, An extension theorem for separately holomorphic functions with analytic singularities, Ann. Pol. Math. (2002), to appear.

[Jar-Pfl 2002b] M. Jarnicki, P. Pflug, An extension theorem for separately holomorphic functions with pluripolar singularities, Trans. Amer. Math. Soc., to appear.

[Kaz 1976] M. V. Kazaryan, On separately meromorphic functions of several complex variables, Mat. Sb. 9 (141) (1976), 538-547.

[Kaz 1978] M. V. Kazaryan, On separately meromorphic functions, Dokl. Akad. Nauk Arm. SSR 17 (1978), 69–73.

[Kaz 1984] M. V. Kazaryan, Meromorphic continuation with respect to groups of variables, Mat. Sb. 125 (167) (1984), 384–397.

[Kli 1991] M. Klimek, Pluripotential Theory, Oxford University Press, 1991.

[Rot 1950] W. Rothstein, Ein neuer Beweis des Hartogsschen Hauptsatzes und seine Ausdehnung auf meromorphe Funktionen, Math. Z. 53 (1950), 84–95.

[Sak 1957] E. Sakai, A note on meromorphic functions in several complex variables, Memoirs of the Faculty of Science, Kyusyu Univ. 11 (1957), 75–80.

[Shi 1986] B. Shiffman, Complete characterization of holomorphic chains of codimension one, Math. Ann. 274 (1986), 233–256.

 $[Shi\ 1989] \hspace{1cm} \hbox{B. Shiffman, } \textit{Separate analyticity and Hartogs theorems}, \ \hbox{Indiana Univ. Math}.$

J. 38 (1989), 943-957.

[Siu 1974] Y. T. Siu, Techniques of Extension of Analytic Objects, Lecture Notes in Pure

and Appl. Math. 8, Marcel Dekker, 1974.

JAGIELLONIAN UNIVERSITY, INSTITUTE OF MATHEMATICS

REYMONTA 4, 30-059 KRAKÓW, POLAND $E\text{-}mail\ address:}$ jarnicki@im.uj.edu.pl

CARL VON OSSIETZKY UNIVERSITÄT OLDENBURG, FACHBEREICH MATHEMATIK

 $\begin{array}{l} {\it Postfach~2503,~D-26111~Oldenburg,~Germany} \\ {\it E-mail~address:} \ \ {\it pflug@mathematik.uni-oldenburg.de} \end{array}$