Adrian Vasiu

Surjectivity Criteria for p-adic Representations, Part I

Received: 10.09.2002 Revised version: 27.01.2003 Manuscripta Mathematica, MM-No:577 Published online: 15.10.2003

ABSTRACT. We prove general surjectivity criteria for p-adic representations. In particular, we classify all adjoint and simply connected group schemes G over the Witt ring W(k) of a finite field k such that the reduction epimorphism $G(W_2(k)) \rightarrow G(k)$ has a

section.

§1. Introduction

Let p be a rational prime. Let $d, r \in \mathbb{N}$. Let $q := p^r$. Let A be an abelian variety of dimension d over a number field E. The action of $Gal(\bar{E}/E)$ on the \bar{E} -valued points of A of p-power order defines a p-adic representation

$$\rho_{A,p}: \operatorname{Gal}(\bar{E}/E) \to GL(T_p(A))(\mathbb{Z}_p) = GL_{2d}(\mathbb{Z}_p),$$

where $T_p(A)$ is the Tate module of A. If A is a semistable elliptic curve over \mathbb{Q} (so d=1 and $E=\mathbb{Q}$), then Serre proved that for $p \geq 3$ the homomorphism $\rho_{A,p}$ is surjective iff its reduction mod p is irreducible (cf. [23, Prop. 21], [24, p. IV-23-24] and [26, p. 519]).

Similar results are expected to hold for many homomorphisms of the form

$$\rho: \operatorname{Gal}(E) \to G(W(k)),$$

with G a reductive group scheme over the Witt ring W(k) of $k := \mathbb{F}_q$ (like when the Zariski closure $G_{A,p}$ of $\mathrm{Im}(\rho_{A,p})$ in $GL(T_p(A))$ is a reductive group scheme and ρ is the factorization of $\rho_{A,p}$ through $G_{A,p}(\mathbb{Z}_p)$. Cases involving q=p and groups G which are extensions of \mathbb{G}_m by products of Weil restrictions of SL_2 groups were treated for $p \geq 5$ in [22]. If $f: G \hookrightarrow GL_n$ is a monomorphism of interest, then in general the assumption that the reduction mod p of $f(W(k)) \circ \rho$ is an irreducible representation over k, is too weak to produce significant results. So the goal of this paper is to get criteria under which ρ is surjective if its reduction mod p (or occasionally mod 4 for p=2) is surjective.

Adrian Vasiu, Mathematics Department, University of Arizona, 617 N. Santa Rita, P.O. Box 210089, Tucson, AZ-85721, USA. e-mail: adrian@math.arizona.edu

 $Mathematics\ Subject\ Classification\ (2000):$ Primary 11S23, 14L17, 17B45, 20G05 and 20G40

A reductive group scheme F over a connected affine scheme $\operatorname{Spec}(R)$ is assumed to have connected fibres. Let F^{der} , Z(F), F^{ab} and F^{ad} be the derived group, the center, the maximal commutative quotient and respectively the adjoint group of F. So $F^{\operatorname{ad}} = F/Z(F)$ and $F/F^{\operatorname{der}} = F^{\operatorname{ab}}$. Let $Z^0(F)$ be the maximal torus of Z(F). Let F^{sc} be the simply connected semisimple group cover of F^{der} . Let $c(F^{\operatorname{der}})$ be the degree of the central isogeny $F^{\operatorname{sc}} \to F^{\operatorname{der}}$. If S is a closed subgroup of F, then $\operatorname{Lie}(S)$ is the R-Lie algebra of S. We now review the contents of this paper.

- 1.1. Adjoint representations. If the reduction mod p of ρ is surjective, then the study of the surjectivity of ρ is intimately interrelated with the study of the adjoint representation AD_{G_k} of G(k) on $\text{Lie}(G_k)$. Though not stated explicitly, this principle is present in disguise in [24, p. IV-23-24] and [21, 2.1]. In order to apply it in §4, in §3 we assume G is semisimple, we work just with G_k (without mentioning G), and we deal with the classification of subrepresentations of AD_{G_k} . If G_k is split, simply connected and G_k^{ad} is absolutely simple, then such a classification was obtained in [13]. The main goal of §3 is to extend loc. cit. to the more general context of Weil restrictions of scalars (see [2, 7.6]) of semisimple groups having absolutely simple adjoints (see 3.4, 3.10 and 3.11). The methods we use are similar to the ones of [13] except that we rely more on the work of Curtis and Steinberg on representations over \bar{k} of finite groups of Lie type and on the work of Humphreys and Hogeweij on ideals of $\text{Lie}(G_{\bar{k}}^{\text{der}})$ (see [17] and [14]; see also [18, 0.13] and [20, §1]). Whenever possible we rely also on [13]. Basic properties of Lie algebras and Weil restrictions are recalled in 2.2 and 2.3.
- **1.2. The problem.** For $s \in \mathbb{N}$ let $W_s(k) := W(k)/p^sW(k)$. Let $\operatorname{Lie}_{\mathbb{F}_p}(G_k)$ be $\operatorname{Lie}(G_k)$ but viewed just as an abelian group identified with $\operatorname{Ker}(G(W_2(k)) \to G(k))$. So if H is a normal subgroup of G_k , then we also view $\operatorname{Lie}_{\mathbb{F}_p}(H)$ as a subgroup of $G(W_2(k))$. Let K be a closed subgroup of G(W(k)) surjecting onto $(G/Z^0(G))(k)$ (we think of it as the image of some ρ). Let $K_2 := \operatorname{Im}(K \to G(W_2(k)))$. The problem we deal with is to find conditions which imply that K surjects onto $(G/Z^0(G))(W(k))$. It splits into two cases: G is or is not semisimple. In this Part I we deal with the first case and in Part II we will deal with the second case and with applications of it to abelian varieties. The group G is semisimple iff the torus $Z^0(G)$ is trivial. If G is semisimple and p|c(G), then there are proper, closed subgroups of G(W(k)) surjecting onto G(k) (cf. 4.1.1). On the other hand, we also have the following general result:
- **1.3. Main Theorem.** We assume that G is semisimple, that g.c.d.(p, c(G)) = 1 and that K surjects onto G(k). We also assume that one of the following five conditions holds:
 - **a)** $q \ge 5$;
- **b)** q = 3 and for each normal subgroup H of G_k which is a PGL_2 or an SL_2 group we have $Lie_{\mathbb{F}_3}(H) \cap K_2 \neq \{0\}$;
 - c) q = 4 and G_k^{ad} has no simple factor which is a PGL_2 group;
 - d) q = 2 and the following two additional conditions hold:
- no simple factor of G_k^{ad} is a PGL_2 group, a Weil restriction from \mathbb{F}_4 to k of a PGL_2 group, or an SU_4^{ad} group;

- for each normal subgroup H of G_k whose adjoint is a PGL_3 group, a SU_3^{ad} group or split of G_2 Lie type, we have $Lie_{\mathbb{F}_2}(H) \cap K_2 \neq \{0\}$;

e)
$$p = 2$$
 and $K_2 = G(W_2(k))$.

Then we have K = G(W(k)).

The case $G = SL_2$ for $p \ge 3$ is due to Lenstra (unpublished computations with 3×3 matrices). Serre proved the case $p \ge 5$ for SL_n and Sp_{2n} groups over \mathbb{Z}_p and the mod 8 variant of e) for SL_n groups over \mathbb{Z}_2 (see [24, IV] and [26, p. 52]). Most of the extra assumptions of b) to d) were known to be needed before (for instance, cf. [12, Sect. 4] in connection to d) for the G_2 Lie type). Though the case q >> 0 of 1.3 is considered well known, we do not know any other concrete literature pertaining to 1.3.

1.4. On the proof of 1.3. The proof of 1.3 is presented in 4.7. Its main ingredients are 4.3 to 4.5 and most of §3. In 4.5 we list all isomorphism classes of adjoint groups $G = G^{\rm ad}$ for which the short exact sequence $0 \to \operatorname{Lie}_{\mathbb{F}_p}(G_k) \to G(W_2(k)) \to G(k) \to 0$ of abstract groups has a section, i.e. the epimorphism $G(W_2(k)) \to G(k)$ has a right inverse. They are the eight ones showing up concretely in 1.3 b) to d). Though most of them are well known, we were not able to trace a reference pertaining to the complete classification of 4.5. The main idea of 4.5 is the following approach of inductive nature (see 4.4). Let $\gamma_G \in H^2(G(k), \operatorname{Lie}_{\mathbb{F}_p}(G_k))$ be the class defining the mentioned short exact sequence, with $\operatorname{Lie}_{\mathbb{F}_p}(G_k)$ viewed as a left G(k)-module via AD_{G_k} . If $\gamma_G = 0$, then all images of restrictions of γ_G are 0 classes and so $\gamma_{G_0^{\rm ad}} = 0$ for any semisimple subgroup G_0 of G normalized by a maximal torus of G (see 4.3.4). But if there is a simple factor of G whose isomorphism class is not in the list, then we can choose G_0 such that the direct computations of 4.4 show that $\gamma_{G_0^{\rm ad}} \neq 0$; so $\gamma_G \neq 0$.

We now detail how 4.5 and §3 get combined to prove 1.3 for $p \ge 5$. Serre's method of [24, IV] can be adapted to get that for $p \ge 5$ it is enough to show that $K_2 = G(W_2(k))$ (see 4.1.2). Based on 4.5 we know that $\text{Lie}_{\mathbb{F}_p}(G_k) \cap K_2$ is not included in $\text{Lie}_{\mathbb{F}_p}(Z(G_k))$. So $\text{Lie}_{\mathbb{F}_p}(G_k) \cap K_2 = \text{Lie}_{\mathbb{F}_p}(G_k)$, cf. 3.7.1 and 3.10 2). So $K_2 = G(W_2(k))$.

Though §3 and §4 handle also the cases p = 2 and p = 3, §4 does not bring anything new to Serre's result recalled before 1.1; however, one can adapt our results to elliptic curves or to [22] in order to get meaningful results in mixed characteristics (0, 2) and (0, 3) (for instance, cf. 1.3 c) to e)). This Part I originated from seminar talks in Berkeley of Ribet and Lenstra; the first draft of §3 was a letter to Lenstra.

§2. Preliminaries

In 2.1 we list our notations and conventions. In 2.2 and 2.3 we recall simple properties of Lie algebras and respectively of Weil restrictions of scalars.

2.1. Notations and conventions. Always $n \in \mathbb{N}$. We denote by k_1 a finite field extension of $k = \mathbb{F}_q$. We abbreviate absolutely simple as a.s. and simply connected as s.c. Let R, F and S be as in §1. We say F^{ad} is simple (resp. is a.s.) if (resp. if each geometric fibre of) it has no proper, normal subgroup of positive relative dimension. If

M is a free R-module of finite rank, then GL(M) (SL(M), etc.) are viewed as reductive group schemes over R. So GL(M)(R) is the group of R-linear automorphisms of M.

Let F be semisimple and $\operatorname{Spec}(R)$ connected. Let o(F) be the order of Z(F) as a finite, flat, group scheme. So $o(SL_2)=2$. We have $c(F)=o(F^{\operatorname{sc}})/o(F)$. See item (VIII) of [5, planches I to IX] for o(F)'s of s.c. semisimple groups having a.s. adjoints. If X_* (resp. X_{R_1} or X) is a scheme over $\operatorname{Spec}(R_1)$, then X_{*R} (resp. X_R) is its pull back to $\operatorname{Spec}(R)$. All modules are left modules. A representation of a Lie algebra or a finite group LG is also referred as an LG-module. See [25, p. 132] and [4, 16.3 to 16.6] for Lang theorem on connected, affine groups over finite fields. See [29] and [11] for the Lie types of semisimple groups over k having a.s. adjoints. For Curtis and Steinberg theory of representations of finite groups of Lie type we refer to [9], [10] and [28, Theorem 1.3]; see also [11, 2.8] for an overview and [3, 6.4, 7.2 and 7.3] for the case of Chevalley groups.

See [5, §4] and [16, §11] for the classification of connected Dynkin diagrams. We say F is of (or has) isotypic $DT \in \{A_n, B_n, C_n | n \in \mathbb{N}\} \cup \{D_n | n \geq 3\} \cup \{E_6, E_7, E_8, F_4, G_2\}$ Dynkin type if the connected Dynkin diagram of any simple factor of every geometric fibre of F^{ad} is DT; if F^{ad} is a.s. we drop the word isotypic. We use the standard notations for classical reductive group schemes over k, W(k) or \mathbb{C} (see [1]). So $PGL_n = GL_n^{\mathrm{ad}} = SL_n^{\mathrm{ad}}$, $PGSp_{2n} = Sp_{2n}^{\mathrm{ad}} = GSp_{2n}^{\mathrm{ad}}$, etc. Also GU_n is the non-split form of GL_n over k or W(k), $SU_n = GU_n^{\mathrm{der}}$ and $PGU_n := SU_n^{\mathrm{ad}} = GU_n^{\mathrm{ad}}$.

2.2. Lie algebras. Let x be an independent variable. As an R-module, we identify Lie(S) with the tangent space of S at the identity section, i.e. with $\text{Ker}(S(R[x]/x^2) \to S(R))$, where the R-epimorphism $R[x]/(x^2) \to R$ takes x into 0. If $y, z \in \text{Lie}(S)$, then the Lie bracket [y, z] is $yzy^{-1}z^{-1}$, the product being taken inside $\text{Ker}(S(R[x]/x^2) \to S(R))$. We now assume that S is a smooth group scheme over R. So Lie(S) is a free R-module of rank equal to the relative dimension of S. The representation of S on GL(Lie(S)) defined by inner conjugation is called the adjoint representation. Let

$$AD_S: S(R) \to GL(\operatorname{Lie}_R(S))(R)$$

be the adjoint representation evaluated at R.

2.3. Weil restrictions of scalars. Let $i_1: R_1 \hookrightarrow R$ be a finite, flat \mathbb{Z} -monomorphism. Let $\operatorname{Lie}_{R_1}(S)$ be $\operatorname{Lie}(S)$ but viewed as an R_1 -Lie algebra. If $R_1 = \mathbb{F}_p$, then we often view $\operatorname{Lie}_{\mathbb{F}_p}(S)$ just as an abelian group; if also $R = \mathbb{F}_p$, then we often drop the lower right index \mathbb{F}_p . Let $\operatorname{Res}_{R/R_1}S$ be the affine group scheme over R_1 obtained from S through the Weil restriction of scalars (see [2, 7.6] and [7, 1.5]). So $\operatorname{Res}_{R/R_1}S$ is defined by the functorial group identification

(1)
$$\operatorname{Hom}_{\operatorname{Spec}(R_1)}(Y, \operatorname{Res}_{R/R_1} S) = \operatorname{Hom}_{\operatorname{Spec}(R)}(Y \times_{\operatorname{Spec}(R_1)} \operatorname{Spec}(R), S),$$

where Y is an arbitrary $\operatorname{Spec}(R_1)$ -scheme. Based on 2.2 and (1) we get a canonical and functorial identification

(2)
$$\operatorname{Lie}(\operatorname{Res}_{R/R_1} S) = \operatorname{Lie}_{R_1}(S).$$

If $i_2: R_2 \hookrightarrow R_1$ is a second finite, flat \mathbb{Z} -monomorphism, then we have a canonical and functorial identification

$$\operatorname{Res}_{R_1/R_2} \operatorname{Res}_{R/R_1} S = \operatorname{Res}_{R/R_2} S.$$

2.3.1. Proposition. We assume there is a finite subgroup C of $Aut_{R_1}(R)$ such that we have an R-isomorphism $R \otimes_{R_1} R = \prod_{c \in C} R \otimes_{R_c} R$. Then we have

$$(Res_{R/R_1}S)_R = \prod_{c \in C} S \times_{Spec(R)} {}_cSpec(R).$$

If S_1 is a closed (resp. smooth, flat, semisimple or reductive) subgroup of S, then $Res_{R/R_1}S_1$ is a closed (resp. smooth, flat, semisimple or reductive) subgroup of $Res_{R/R_1}S$. If S is a reductive group scheme, then any maximal torus (resp. Borel subgroup) J of $Res_{R/R_1}S$ is of the form $Res_{R/R_1}T$ (resp. $Res_{R/R_1}B$), where T (resp. B) is a maximal torus (resp. Borel subgroup) of S.

Proof: Formula (3) follows from (1) and our hypothesis. See [2, Prop. 5 of 7.6] for the closed subgroup, smooth or flat part. Let now S be a reductive group scheme. We know that $\operatorname{Res}_{R/R_1}S$ is affine, flat and smooth. So in order to show that it is a reductive group scheme, it is enough to show that its geometric fibres are so. So it is enough to show that the fibres of $(\operatorname{Res}_{R/R_1}S)_R$ are reductive group schemes. But this is so as $(\operatorname{Res}_{R/R_1}S)_R$ is a reductive group scheme, cf. (3). Similarly we argue that $\operatorname{Res}_{R/R_1}S$ is semisimple if S is so.

To check the last part, we first remark that J_R is a maximal torus (resp. Borel subgroup) of $(\operatorname{Res}_{R/R_1}S)_R$. So based on (3), it is uniquely determined by the projection T (resp. B) of J_R on the factor $S = S \times_{\operatorname{Spec}(R) 1_R} \operatorname{Spec}(R)$ of $(\operatorname{Res}_{R/R_1}S)_R$. So $J = \operatorname{Res}_{R/R_1}T$ (resp. $J = \operatorname{Res}_{R/R_1}B$). This ends the proof.

The next structure Theorem (see [29, 3.1.2]) will play important roles in §3 and §4.

2.3.2. Theorem. We assume R is a field. Then any adjoint (resp. s.c.) group F over R is isomorphic to a product of adjoint (resp. s.c.) groups of the form $Res_{\tilde{R}/R}\tilde{F}$, where \tilde{R} is a separable finite field extension of R and \tilde{F} is an adjoint (resp. s.c.) group over \tilde{R} having an a.s. adjoint. So if F^{ad} is simple, then F is of isotypic Dynkin type.

§3. Adjoint representations over finite fields

Let H be a semisimple group over $k = \mathbb{F}_q$. In §3 we study H, Lie(H) and AD_H . In 3.3 we recall all cases when Lie(H) is simple, i.e. it has no proper ideal defined over k. These cases are precisely the ones when AD_H is irreducible, cf. 3.4. In 3.5 to 3.11 we work in the context of Weil restrictions. In 3.1, 3.2 and 3.5 to 3.8 we include preliminary material on H and Lie(H). See 3.7.1 for a refinement of 3.4 obtained by working with

$$\mathfrak{L}_H := \operatorname{Im}(\operatorname{Lie}(H^{\operatorname{sc}}) \to \operatorname{Lie}(H))$$

instead of Lie(H). In 3.9 we list semisimple subgroups of H to be used in 3.10, 3.11 and §4. In 3.10 and 3.11 we study subrepresentations of AD_H ; these sections extend [13].

- **3.1.** On Z(H). The finite group scheme Z(H) is of multiplicative type, cf. [27, Vol. III, 4.1.7 of p. 173]. So the finite group Z(H)(k) is of order prime to p. If p|o(H), then Lie(Z(H)) is a proper ideal of Lie(H) normalized by H and so AD_H is reducible. If g.c.d.(p,o(H))=1, then the central isogeny $H\to H^{\text{ad}}$ is étale and so by identifying the tangent spaces in the origin we get $\text{Lie}(H^{\text{ad}})=\text{Lie}(H)$. Similarly, if g.c.d.(p,c(H))=1 we have $\text{Lie}(H^{\text{sc}})=\text{Lie}(H)$.
- **3.2. Lemma.** We assume that H^{ad} is a.s. and $Lie(H^{\mathrm{ad}})$ is not a.s. Then $Lie(H^{\mathrm{ad}})$ has a proper, characteristic ideal and so AD_H is reducible.

Proof: If $p \ge 5$, or if p = 3 and $H^{\rm ad}$ is not of G_2 Lie type, or if p = 2 and $H^{\rm ad}$ is not of F_4 Lie type, then $[{\rm Lie}(H^{\rm ad}), {\rm Lie}(H^{\rm ad})]$ is a characteristic ideal of ${\rm Lie}(H^{\rm ad})$ of codimension 1 or 2 (cf. [18, 0.13] applied to $H^{\rm ad}_{\bar k}$). In the excluded two cases the group $H^{\rm ad}$ is split and (cf. loc. cit.) ${\rm Lie}(H^{\rm ad})$ has a unique proper ideal: it corresponds to root vectors of short roots. So ${\rm Lie}(H^{\rm ad})$ has a proper, characteristic ideal and so AD_H is reducible.

- **3.3. Proposition.** The Lie algebra Lie(H) is not simple iff one of the following four conditions holds:
 - i) H^{ad} is not simple;
 - ii) H^{ad} is simple of isotypic A_{pn-1} Dynkin type;
- iii) p = 2 and H^{ad} is simple of an isotypic Dynkin type belonging to the union $\{B_n, C_n | n \in \mathbb{N}\} \cup \{D_n | n \geq 3\} \cup \{E_7, F_4\};$
 - iv) p = 3 and H^{ad} is simple of isotypic E_6 or G_2 Dynkin type.

Proof: If i) to iv) do not hold, then g.c.d.(p, o(H)) = 1 and so $\text{Lie}(H) = \text{Lie}(H^{\text{ad}})$. If Lie(H) is simple, then g.c.d.(p, o(H)) = 1 and so $\text{Lie}(H) = \text{Lie}(H^{\text{ad}})$ (cf. 3.1) and H is simple (cf. 2.3.2). So it suffices to prove the Proposition under the extra assumption that H is adjoint and simple. Let k_1 be such that $H = \text{Res}_{k_1/k}H_1$, where H_1 is an a.s. adjoint group over k_1 (cf. 2.3.2). We have $\text{Lie}(H) = \text{Lie}_k(H_1)$, cf. (2). We show that $\text{Lie}(H_1)$ is simple iff Lie(H) is simple. The "if" part is obvious. To check the "only if" part let I be a non-zero ideal of Lie(H). So $I \otimes_k k_1$ is a non-zero ideal of

(4)
$$\operatorname{Lie}(H) \otimes_k k_1 = \operatorname{Lie}_k(H_1) \otimes_k k_1 \tilde{\to} \bigoplus_{\tau \in \operatorname{Gal}(k_1/k)} \operatorname{Lie}(H_1) \otimes_{k_1 \tau} k_1$$

and so a direct sum of some of these direct summands indexed by elements of $Gal(k_1/k)$. But the only non-zero such direct sum defined over k is $Lie_k(H_1) \otimes_k k_1$. So I = Lie(H).

Also, $Lie(H_1)$ is a.s. iff it is simple (cf. 3.2 for the if part). So Lie(H) is simple iff $Lie(H_1)$ is a.s. So the Proposition follows from [18, 0.13].

3.4. Proposition. Let H be a semisimple group over k. Then AD_H is irreducible iff Lie(H) is simple (i.e. iff none of the four conditions of 3.3 holds).

Proof: We assume AD_H is irreducible. So g.c.d.(p, o(H)) = 1 and $Lie(H) = Lie(H^{ad})$, cf. 3.1. Moreover H^{ad} is simple, cf. 2.3.2. So it suffices to show that Lie(H) is simple under the extra assumption that H^{ad} is a.s., cf. the proof of 3.3. So Lie(H) is a.s., cf. 3.2.

We now assume Lie(H) is simple. So H^{ad} is simple and $g.c.d.(p, o(H^{\text{sc}})) = 1$, cf. 3.3. So to show that AD_H is irreducible we can assume that $H = \text{Res}_{k_1/k}H_1$, with H_1 a s.c.

semisimple group over k_1 having an a.s. adjoint. So $\text{Lie}(H_1) = \text{Lie}(H_1^{\text{ad}})$ are a.s., cf. 3.3. Let DT be the Dynkin type of H_1 . The maximal weight ϖ of the adjoint representation of the complex, simple Lie algebra of DT Dynkin type is the maximal root. So if DT is A_n (resp. is B_n with $n \geq 3$, C_n with $n \geq 2$, D_n with $n \geq 4$, E_6 , E_7 , E_8 , F_4 or G_2), then with the notations of [5, planche I to IX] ϖ is $\varpi_1 + \varpi_n$ (resp. is ϖ_2 , $2\varpi_1$, ϖ_2 , ϖ_2 , ϖ_1 , ϖ_8 , ϖ_1 or ϖ_2). From Curtis and Steinberg theory and the last two sentences we get that AD_{H_1} is absolutely irreducible. Let I be a non-trivial, irreducible subrepresentation of AD_H . The extension of AD_H to k_1 is a direct sum of $[k_1 : k]$ copies of AD_{H_1} , cf. (4). So due to the absolute irreducibility of AD_{H_1} , $I \otimes_k k_1$ is a $\text{Lie}(H_1)$ -submodule of $\text{Lie}(H) \otimes_k k_1$. So as $(I \otimes_k k_1) \cap \text{Lie}(H) = I$, I is an ideal of Lie(H). So I = Lie(H). This ends the proof.

3.5. The basic setting. Until §4 we assume that

$$H = \operatorname{Res}_{k_1/k} H_1,$$

where H_1 is a semisimple group over k_1 such that H_1^{ad} is a.s. So $\text{Lie}(H) = \text{Lie}_k(H_1)$. Let

DT

be the Dynkin type of H_1 . Let k_2 and k_3 be the quadratic and respectively the cubic field extension of k_1 . The group H_1 has a Borel subgroup B_1 (this is part of Lang theorem). Let T_1 be a maximal torus of B_1 , cf. [4, (i) of Theorem 18.2]. The group H_1 is split iff T_1 is. So there is a smallest field extension l of k_1 such that H_{1l} is split. Let

(5)
$$\operatorname{Lie}(H_{1l}) = \operatorname{Lie}(T_{1l}) \bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha}$$

be the Weyl decomposition of $\text{Lie}(H_{1l})$ with respect to T_{1l} . So each \mathfrak{g}_{α} is a 1 dimensional l-vector space normalized by T_{1l} and Φ is an irreducible root system of characters of T_{1l} . Let $\Phi^+ := \{\alpha \in \Phi | \mathfrak{g}_{\alpha} \subset \text{Lie}(B_{1l})\}$. Let Δ be the basis of Φ included in Φ^+ . We use [5, planches I to IX] to denote the elements of Δ , Φ^+ and Φ ; so $\Delta = \{\alpha_1, ..., \alpha_{|\Delta|}\}$, where $|\Delta|$ is the number of elements of DT. Warning: if $l = k_1$ (resp. l = k), then in connection to (5) we often drop the lower right index l (resp. 1l and so also T_1 becomes T).

We denote also by Δ the Dynkin diagram of DT defined by the elements of Δ . Let ϕ be the Frobenius automorphism of \bar{l} fixing k_1 . It acts on Φ via the rule: if $\alpha \in \Phi$, then we have $\mathfrak{g}_{\phi(\alpha)} = (1_{\mathrm{Lie}(H_1)} \otimes \phi)(\mathfrak{g}_{\alpha})$. Let $e \in \mathbb{N}$ be the order of the permutation of Δ defined by ϕ . For $m \in \mathbb{N}$, ϕ^m fixes $\mathrm{Lie}(T_{1l})$ iff e|m. So $[l:k_1]=e$. But e is the order of an element of $\mathrm{Aut}(\Delta)$ and so $e \in \{1,2,3\}$. More precisely, we have $l=k_1$ if $e \in \{B_n, C_n|n \geq 1\} \cup \{E_7, E_8, F_4, G_2\}$, $l \in \{k_1, k_2\}$ if $DT \in \{A_n|n \geq 2\} \cup \{D_n|n \geq 5\} \cup \{E_6\}$ and $l \in \{k_1, k_2, k_3\}$ if $LT = D_4$. If e = 1 let LT := DT. If $e \in \{2,3\}$ let $LT := {}^eDT$. We recall that the pair (Δ, ϕ) or just

LT

is called the Lie type of H_1 and that H_1^{ad} and H_1^{sc} are uniquely determined by it (see [29]). For $\alpha \in \Phi$ let $H(|\alpha|)$ be the semisimple subgroup of H_{1l} generated by the two \mathbb{G}_a subgroups $\mathbb{G}_{a,\alpha}$ and $\mathbb{G}_{a,-\alpha}$ of H_{1l} normalized by T_{1l} and having \mathfrak{g}_{α} and respectively $\mathfrak{g}_{-\alpha}$ as their Lie algebras. So $H(|\alpha|)^{\text{ad}}$ is a PGL_2 group and H_{1l} is generated by all these $H(|\alpha|)$'s. All these can be deduced from [4, §13 of Ch. IV] via descent from \bar{l} to l.

- **3.6. Proposition.** Let $\beta \in \Phi \setminus \{-\alpha, \alpha\}$. Let $\Psi_1(\alpha, \beta) := \{i\alpha + \beta | i \in \mathbb{N}, i\alpha + \beta \in \Phi\}$. If $\alpha + \beta \notin \Phi$ let $\mathfrak{g}_{\alpha+\beta} := \{0\}$. Let $g \in \mathbb{G}_{a,\alpha}(l)$ be a non-identity element. Let $x \in \mathfrak{g}_{\beta} \setminus \{0\}$.
 - a) We have $AD_{H_{1l}}(g)(x) x \in \bigoplus_{\delta \in \Psi_1(\alpha,\beta)} \mathfrak{g}_{\delta}$.
- **b)** We always have an inclusion $[\mathfrak{g}_{\alpha},\mathfrak{g}_{\beta}] \subset \mathfrak{g}_{\alpha+\beta}$. It is not an equality iff one of the following disjoint three conditions holds:
 - i) p = 3, $DT = G_2$, and α and β are short roots forming an angle of 60° ;
 - ii) p = 2, $DT = G_2$, and α and β are short roots forming an angle of 120° ;
 - iii) p = 2, $DT \in \{B_n, C_n | n \ge 2\} \cup \{F_4\}$, α and β are short, $\alpha \perp \beta$ and $\alpha + \beta \in \Phi$.
- c) If none of the three conditions of b) holds and if $\alpha + \beta \in \Phi$, then the component of $AD_{H_{1l}}(g)(x) x$ in $\mathfrak{g}_{\alpha+\beta}$ is non-zero.

Proof: Let $\kappa(\alpha, \beta) := \Phi \cap \{i\alpha + j\beta | (i,j) \in \mathbb{N}^2\}$. It is known that there are isomorphisms $x_{\delta} : \mathbb{G}_a \stackrel{\sim}{\to} \mathbb{G}_{a,\delta} \ (\delta \in \kappa(\alpha,\beta))$, an ordering on \mathbb{N}^2 , and integers $m_{\alpha,\beta;i,j} \ ((i,j) \in \mathbb{N}^2)$ such that for all $s, t \in \mathbb{G}_a(\bar{l}) = \bar{l}$ we have (cf. [27, Vol. III, p. 321] or [8, 4.2])

(6)
$$x_{\alpha}(s)x_{\beta}(t)x_{\alpha}(s^{-1})x_{\beta}(t^{-1}) = \prod_{(i,j)\in\mathbb{N}^2, i\alpha+j\beta\in\kappa(\alpha,\beta)} x_{i\alpha+j\beta}(m_{\alpha,\beta;i,j}s^it^j).$$

Taking the derivative in (6) with respect to t at the identity element, we get that the difference $AD_{H_1\bar{l}}(x_{\alpha}(s))(x) - x$ belongs to $(\bigoplus_{i \in \mathbb{N}, i\alpha + \beta \in \Psi_1(\alpha,\beta)} \mathfrak{g}_{i\alpha+\beta}) \otimes_l \otimes \bar{l}$ and its component in $\mathfrak{g}_{i\alpha+\beta} \otimes_l \otimes \bar{l}$ is a fixed generator of $\mathfrak{g}_{i\alpha+\beta} \otimes_l \otimes \bar{l}$ times $m_{\alpha,\beta;i,1}s^i$ (cf. [4, 3.16 of Ch. 1]). Similarly, taking the derivative in (6) with respect to s and t at the identity element, we get that $[\mathfrak{g}_{\alpha},\mathfrak{g}_{\beta}] = m_{\alpha,\beta;1,1}\mathfrak{g}_{\alpha+\beta} \subset \mathfrak{g}_{\alpha+\beta}$. But $m_{\alpha,\beta;1,1} = \pm (m+1)$, where $m \in \mathbb{Z}$ is the greatest integer such that $\beta - m\alpha \in \Phi$ (see [27, Vol. III, p. 321] or [8, 4.2 and 4.3]). So the second part of b) and c) follow from the fact that we have p|m+1 iff one of the three conditions of b) holds (see loc. cit.). This ends the proof.

Let B_1^{opp} be the Borel subgroup of H_1 which is the opposite of B_1 with respect to T_1 . Let U_1 and U_1^{opp} be the unipotent radicals of B_1 and respectively of B_1^{opp} . Let L_H be the Lie subalgebra of $\text{Lie}(H) = \text{Lie}_k(H_1)$ generated by $\text{Lie}_k(U_1)$ and $\text{Lie}_k(U_1^{\text{opp}})$.

- **3.7. Proposition. 1)** The Lie algebra L_H is an ideal of Lie(H) containing [Lie(H), Lie(H)]. If p > 2 and $l = k_1$, then $L_H = \sum_{\alpha \in \Phi^+} Lie_k(H(|\alpha|))$.
- **2)** Always $L_H = \mathfrak{L}_H$. Also, if p > 2 or if p = 2 and $DT \notin \{C_n | n \in \mathbb{N}\}$, then $L_H = [Lie(H), Lie(H)]$.
 - 3) We have $\mathfrak{L}_{H^{\mathrm{ad}}} = L_{H^{\mathrm{ad}}} = [Lie(H^{\mathrm{ad}}), Lie(H^{\mathrm{ad}})].$

Proof: Tensoring with l we can assume $k = k_1 = l$; so $H = H_1$ is split. The first part of 1) is obvious. If $p \ge 3$, then $\text{Lie}(H(|\alpha|)) = \mathfrak{g}_{\alpha} \oplus \mathfrak{g}_{-\alpha} \oplus [\mathfrak{g}_{\alpha}, \mathfrak{g}_{-\alpha}]$. So the last part of 1) follows from the inclusions of 3.6 b). We now prove 2). Both U_1 and U_1^{opp} are naturally identified with unipotent subgroups of H_1^{sc} and so $L_{\tilde{H}}$ makes sense if $\tilde{H} \to H$ is an arbitrary central isogeny. Always $L_{\tilde{H}}$ surjects onto L_H and so to prove 2) we can assume that H is s.c.

If p > 2 or if p = 2 and $DT \notin \{C_n | n \in \mathbb{N}\}$, then Lie(H) = [Lie(H), Lie(H)] (cf. [18, 0.13]) and so 2) follows from 1). So we are left to show that $L_H = \text{Lie}(H)$ if p = 2 and $DT = C_n$. But then H has a subgroup SL(H) which is a product of SL_2 groups and has T as a maximal torus; so $\text{Lie}(T) \subset L_H$ and so $L_H = \text{Lie}(H)$. This proves 2).

Based on 2) and its proof, its suffices to prove 3) under the extra assumptions that p = 2 and $DT = C_n$. As $L_{H^{\mathrm{ad}}} = \mathfrak{L}_{H^{\mathrm{ad}}}$, the ideal $L_{H^{\mathrm{ad}}}$ of $\mathrm{Lie}(H^{\mathrm{ad}})$ is of codimension 1 and contains $[\mathrm{Lie}(H^{\mathrm{ad}}), \mathrm{Lie}(H^{\mathrm{ad}})]$. By reasons of dimensions we get $L_{H^{\mathrm{ad}}} = [\mathrm{Lie}(H^{\mathrm{ad}}), \mathrm{Lie}(H^{\mathrm{ad}})]$ (see [18, 0.13]). This ends the proof.

3.7.1. Corollary. We assume g.c.d.(p, o(H)) = 1. If p = 3 we also assume that $DT \neq G_2$ and if p = 2 we also assume that $DT \notin \{B_n, C_n | n \geq 2\} \cup \{F_4\}$. Then the natural representation of $H^{\text{sc}}(k) = H_1^{\text{sc}}(k_1)$ on $\mathfrak{L}_{H^{\text{ad}}} = \mathfrak{L}_H = [Lie(H), Lie(H)]$ is irreducible.

Proof: The H_1 -module \mathfrak{L}_{H_1} is a.s. Argument: the case $DT = A_1$ is trivial and the case $DT \neq A_1$ is a consequence of the fact that \mathfrak{L}_{H_1} is an a.s. $\text{Lie}(H_1)$ -module (see [18, 0.13]). The maximal weight of the representation of H_{1l} on $\mathfrak{L}_{H_{1l}}$ is ϖ of the proof of 3.4. The rest of the proof is entirely the same as the proof of 3.4.

We come back to (5). Let $\alpha \in \Phi$. Let $T(|\alpha|)$ be the maximal subtorus of T_{1l} centralizing \mathfrak{g}_{α} . So $[\mathfrak{g}_{\alpha}, \operatorname{Lie}(T(|\alpha|))] = \{0\}$. We have a short exact sequence

(7)
$$0 \to T(|\alpha|) \to TH(|\alpha|) \to \tilde{H}(|\alpha|) \to 0,$$

where $TH(|\alpha|)$ is the reductive subgroup of H_{1l} generated by $T(|\alpha|)$ and $H(|\alpha|)$ and where $\tilde{H}(|\alpha|)$ is either $H(|\alpha|)$ or its adjoint.

3.8. Lemma. The group $H(|\alpha|)$ is a PGL_2 group iff $DT = B_n$, H_1 is adjoint and α is short. Also, if H_1 is adjoint, then $\tilde{H}(|\alpha|) = H(|\alpha|)^{ad}$.

Proof: For $\beta \in \Phi$ let $ST(\alpha, \beta) := \{i\alpha + \beta | i \in \mathbb{Z}\} \cap \Phi$ be the α -string through β . It is of the form $\{i\alpha + \beta | i \in \{-s, -s + 1, ..., t\}\}$, for some $s, t \in \mathbb{N} \cup \{0\}$ with $s + t \leq 3$ (see [15, p. 45]). The set $\Phi \setminus \{\alpha, -\alpha\}$ is a disjoint union of α -strings. So let $\Phi(\alpha)$ be a subset of $\Phi \setminus \{\alpha, -\alpha\}$ such that we have a disjoint union $\Phi \setminus \{\alpha, -\alpha\} = \bigcup_{\beta \in \Phi(\alpha)} ST(\alpha, \beta)$. For $\beta \in \Phi(\alpha)$ let $V_{\alpha,\beta} := \bigoplus_{\delta \in ST(\alpha,\beta)} \mathfrak{g}_{\delta}$. It is an $H(|\alpha|)$ -module, cf. 3.6 a). We get a direct sum decomposition of $H(|\alpha|)$ -modules $\operatorname{Lie}(H_{1l}) = \operatorname{Lie}(TH(|\alpha|)) \bigoplus \bigoplus_{\beta \in \Phi(\alpha)} V_{\alpha,\beta}$.

If $H(|\alpha|)$ is a PGL_2 group, then $\dim_l(V_{\alpha,\beta}) \in \{1,3\}$, $\forall \beta \in \Phi(\alpha)$, and the converse holds if H_1 is adjoint. Using [5, planches I to IX] and [15, Lemma C of p. 53] we easily get that $\dim_l(V_{\alpha,\beta}) \in \{1,3\}$, $\forall \beta \in \Phi(\alpha)$, iff $DT = B_n$ and α is short. But if H_1 is s.c., then $H(|\alpha|)$ is an SL_2 group as we can check by reduction to the B_1 and $B_2 = C_2$ Dynkin types. This proves the first part.

If H_1 is adjoint, then the subgroup of T_{1l} fixing \mathfrak{g}_{α} is a subtorus of T_{1l} of codimension 1 and so it is $T(|\alpha|)$. So $Z(H(|\alpha|))$ is a subgroup of $T(|\alpha|)$ and so $\tilde{H}(|\alpha|) = H(|\alpha|)^{\mathrm{ad}}$. This ends the proof.

3.8.1. Remark. If $(\Psi_1(\alpha,\beta) \cup \Psi_1(-\alpha,\beta)) \neq \emptyset$, then $\Phi \cap (\mathbb{Z}\alpha \oplus \mathbb{Z}\beta)$ is the root system of a Dynkin type $DT_2 := DT(\alpha,\beta) \in \{A_2,B_2,G_2\}$. If $DT_2 = G_2$, then $DT = G_2$. If $DT_2 = B_2$, then $DT \in \{B_n,C_n|n\geq 2\} \cup \{F_4\}$. Let $x \in \mathfrak{g}_{\alpha}$. If $DT_2 = A_2$, then $\Psi_1(\alpha,\beta)$ has at most 1 element and so $\operatorname{ad}(x)^2$ annihilates $V_{\alpha,\beta}$ (cf. 3.6 a)). Similarly if $DT_2 = B_2$

(resp. $DT_2 = G_2$), then $\Psi_1(\alpha, \beta)$ has at most 2 (resp. 3) elements and so $ad(x)^3$ (resp. $ad(x)^4$) annihilates $V_{\alpha,\beta}$.

3.9. Subgroups. We list semisimple subgroups of H_1 normalized by T_1 . By taking the $\operatorname{Res}_{k_1/k}$ of them we get semisimple subgroups of H. So except for the first two Cases below (which are samples) we will just mention semisimple subgroups of H_1 and not of H.

Let Φ_0 be a subset of Φ having the following three properties:

- (1) it is invariant under ϕ of 3.5;
- (2) it is stable under additions, i.e. we have $\Phi \cap \{\alpha + \beta | \alpha, \beta \in \Phi_0\} \subset \Phi_0$;
- (3) it is symmetric, i.e. we have $\Phi_0 = -\Phi_0$.

Let H_{0l} be the subgroup of H_{1l} generated by $H(|\alpha|)$, where $\alpha \in \Phi_0$. It is invariant under ϕ and T_{1l} and so it the extension to l of a subgroup H_0 of H_1 normalized by T_1 . The subgroup TH_0 of H_1 generated by T_1 and H_0 is reductive, cf. [27, Vol. III, 5.4.7 and 5.10.1 of Exp. XXII] applied to TH_{0l} . But H_0 is a subgroup of $(TH_0)^{\text{der}}$ and contains the unipotent radical of any Borel subgroup of TH_0 normalized by T_1 . So $H_0 = (TH_0)^{\text{der}}$ and so H_0 is semisimple. A subset Φ_0^+ of $\Phi_0 \cap \Phi^+$ is said to generate Φ_0 , if each element of $\Phi_0 \cap \Phi^+$ is a linear combination with non-negative integer coefficients of elements of Φ_0^+ . We refer to H_0 as the semisimple subgroup of H_1 associated to Φ_0 or generated by Φ_0^+ .

In this paragraph we assume $l = k_1$. If $DT \notin \{A_1, B_2, G_2\}$, then we can choose Φ_0 such that H_0 is of A_2 Lie type. If $DT \notin \{A_1, A_2, B_2, B_3, C_3, D_4, F_4, G_2\}$ (resp. $DT \in \{B_n, C_n | n \ge 2\} \cup \{F_4\}$), then we can choose Φ_0 such that H_0 is of A_3 (resp. C_2) Lie type. We assume now that $DT \in \{B_n, C_n | n \ge 2\} \cup \{F_4, G_2\}$. Let H_1^{long} be the semisimple subgroup of H_1 generated by $\{\alpha \in \Phi^+ | \alpha \text{ is long}\}$. Warning: the group H_1^{long} as well as most of the below semisimple subgroups of H_1 are not generated by subsets of Δ . If $n \ge 3$ and $DT = B_n$, then H_1^{long} is of D_n Lie type. If $n \ge 2$, H_1 is s.c. and $DT = C_n$, then H_1^{long} is an SL_2^n group. If $DT = F_4$ (resp. $DT = G_2$), then H_1^{long} is of A_3 (resp. A_2) Lie type. These Lie types can be read out from [5, planches II, III, VIII and IX].

Until 3.10 we use (5) with $l \in \{k_2, k_3\}$. If ϖ is as in the proof of 3.4, then the subgroup $H(|\varpi|)$ of H_{1l} is the pull back of an SL_2 subgroup of H_1 (cf. also 3.8 applied to H_{1l}). If $LT = {}^2A_{n+4}$ and $\Phi_0^+ = \{\alpha_1, \alpha_2, \alpha_{n+3}, \alpha_{n+4}\}$, then H_0 is isogenous to the $\operatorname{Res}_{k_2/k_1}$ of an SL_3 group. If $LT = {}^2A_4$ and $\Phi_0^+ = \{\alpha_1, \alpha_4\}$, then H_0 is the $\operatorname{Res}_{k_2/k_1}$ of an SL_2 group. We assume now that $LT = {}^2D_{n+3}$. If n = 1 we assume ϕ permutes α_3 and α_4 . If n > 1 (resp. n = 1) and $\Phi_0^+ = \{\alpha_1, \alpha_2, \alpha_3\}$ (resp. $\Phi_0^+ = \{\alpha_1, \alpha_2, \alpha_2 + \alpha_3 + \alpha_4\}$), then H_0^{ad} is a PGL_4 group. To introduce other semisimple subgroups of H_1 we consider five Cases.

Case 1: $LT = {}^2A_{2n+1}$. Let $H_1(n+1)$ be the SL_2 subgroup of H_1 such that $H_1(n+1)_l = H(|\alpha_{n+1}|)$. So $H(n+1) := \operatorname{Res}_{k_1/k} H_1(n+1)$ is a subgroup of H. Let $i \in \{1,...,n\}$. Let $\alpha_i^1 := \sum_{j=i}^{n+1} \alpha_j$ and $\alpha_i^2 := \sum_{j=n+1}^{2n+2-i} \alpha_j$. The subgroups $H(|\alpha_i^1|)$ and $H(|\alpha_i^2|)$ of H_{1k_2} are SL_2 groups and commute. We first assume that either n > 1 or $H_1 = H_1^{\rm sc}$. The product $H(|\alpha_i^1|) \times_{k_2} H(|\alpha_i^2|)$ is naturally a subgroup of H_{1k_2} , invariant under the action of $\operatorname{Gal}(k_2/k_1)$ on H_{1k_2} . So it is the extension to k_2 of a subgroup $H_1(i)$ of H_1 . As $\operatorname{Gal}(k_2/k_1)$ takes $\operatorname{Lie}(H(|\alpha_i^1|))$ into $\operatorname{Lie}(H(|\alpha_i^2|))$, the group $H_1(i)$ is the $\operatorname{Res}_{k_2/k_1}$ of an SL_2 group. So $H(i) := \operatorname{Res}_{k_1/k} H_1(i)$ is a subgroup of H which is the $\operatorname{Res}_{k_2/k}$ of an SL_2 group. If n = 1 and $H_1 \neq H_1^{\rm sc}$, then let $H_1(1)$ be the semisimple subgroup of H_1 such that we have

a natural central isogeny $H(|\alpha_1^1|) \times_{k_2} H(|\alpha_1^2|) \to H_1(1)_{k_2}$ of degree 2; the adjoint of the subgroup $H(1) := \operatorname{Res}_{k_1/k} H_1(1)$ of H is the $\operatorname{Res}_{k_2/k}$ of a PGL_2 group.

Case 2: $LT = {}^2A_{2n}$. For $i \in \{1, ..., n-1\}$ let $\alpha_i^1 := \sum_{j=i}^{n+1} \alpha_j$ and $\alpha_i^2 := \sum_{j=n}^{2n+1-i} \alpha_j$. Let $H_1(i)$ and $H_1(n+1)$ be subgroups of H_1 such that $H_1(i)_{k_2} = H(|\alpha_i^1|) \times_{k_2} H(|\alpha_i^2|)$ and $H_1(n+1)_{k_2} = H(|\alpha_n + \alpha_{n+1}|)$. The adjoint of the semisimple subgroup $H_1(n)$ of H_1 generated by $\{\alpha_n, \alpha_{n+1}\}$ is a PGU_3 group. So for $i \in \{1, ..., n+1\}$, the group $H(i) := \operatorname{Res}_{k_1/k} H_1(i)$ is a subgroup of H. If $i \in \{1, ..., n-1\}$ (resp. if i = n+1), then H(i) is the $\operatorname{Res}_{k_2/k}$ (resp. $\operatorname{Res}_{k_1/k}$) of an SL_2 group.

Case 3: $LT = {}^{2}E_{6}$. Let $H_{1}(1)'$ (resp. $H_{1}(4)$) be the semisimple subgroup of H_{1} generated by $\{\alpha_{1}, \alpha_{3}, \alpha_{4}, \alpha_{5}, \alpha_{6}\}$ (resp. $\{\alpha_{2}\}$). So $H_{1}(1)'$ is of ${}^{2}A_{5}$ Lie type and $H_{1}(4)$ is an SL_{2} group, cf. 3.8. We apply Case 1 to $H_{1}(1)'$: we get semisimple subgroups $H_{1}(1)$, $H_{1}(2)$ and $H_{1}(3)$ of $H_{1}(1)'$ and so also of H_{1} .

Case 4: $LT = {}^2D_{n+3}$. If n = 1 we assume ϕ permutes α_3 and α_4 . For $i \in \{1, ..., n+2\}$ let $H_1(i)$ be the semisimple subgroup of H_1 which over k_2 is generated by $H(|\alpha|)$, with $\alpha \in \{\sum_{j=i}^{n+2} \alpha_j, \alpha_{n+3} + \sum_{j=i}^{n+1} \alpha_j\}$. So $H_1(i)^{\text{sc}}$ is the Res_{k_2/k_1} of an SL_2 group.

Case 5: $LT = {}^3D_4$. Let $H_1(1)$ (resp. $H_1(2)$) be the semisimple subgroup of H_1 which over k_3 is generated by $H(|\alpha|)$, with $\alpha \in \{\alpha_1 + \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_4, \alpha_2 + \alpha_3 + \alpha_4\}$ (resp. with $\alpha = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$). So $H_1(2)$ is an SL_2 group (cf. 3.8) and $H_1(1)^{\text{sc}}$ is the Res_{k_3/k_1} of an SL_2 group.

3.9.1. Lemma. We refer to Case 4 (resp. Case 5). If $H = H^{sc}$, then each $H_1(i)$ (resp. then $H_1(1)$) are (resp. is) s.c.

Proof: To check this we can work with H_l . Even better, based on the existence of Chevalley group schemes over \mathbb{Z} we can work over \mathbb{C} . Based on [6, §13 of Ch. VIII] we need to show that any standard (resp. standard composite) monomorphism $SO_4 \hookrightarrow SO_{2n+6}$ (resp. $SL_2 \times SO_4 \hookrightarrow SO_4 \times SO_4 \hookrightarrow SO_8$) over \mathbb{C} , lifts to a monomorphism $\operatorname{Spin}_4 \hookrightarrow \operatorname{Spin}_{2n+6}$ (resp. $SL_2^3 = SL_2 \times \operatorname{Spin}_4 \hookrightarrow \operatorname{Spin}_8$). To check this last statement we can work just in the context of SO_{2n+6} . The restriction of the spin representation of $\operatorname{Lie}(SO_{2m})$ to $\operatorname{Lie}(SO_{2m-2})$ via a standard monomorphism $SO_{2m-2} \hookrightarrow SO_{2m}$ is a disjoint union of two spin representations of $\operatorname{Lie}(SO_{2m-2})$, $\forall m \in \mathbb{N}$, $m \geq 4$ (see the description of spin representations in loc. cit.). From this the statement on lifts follows. This ends the proof.

3.9.2. Lemma. The Lie subalgebra LS_H of $Lie_k(H_1) = Lie(H)$ generated by all intersections $Lie_k(H_1(i) \cap U_1)$'s and $Lie_k(H_1(i) \cap U_1^{\text{opp}})$'s is $\mathfrak{L}_H = L_H$.

Proof: Tensoring with l over k, this follows easily from 3.6 b).

3.9.3. Simple properties. Let $TH_1(i)$ be the reductive subgroup of H_1 generated by T_1 and $H_1(i)$. Let $T_1(i)$ be the maximal torus of $Z(TH_1(i))$. We have a short exact sequence

(8)
$$0 \to \operatorname{Lie}(T_1(i)) \to \operatorname{Lie}(TH_1(i)) \to \operatorname{Lie}(\tilde{H}_1(i)) \to 0,$$

where $\tilde{H}_1(i)$ is the quotient of $H_1(i)$ by $Z(H_1(i)) \cap T_1(i)$. If H is adjoint, then as in the proof of 3.8 we argue that in fact $\tilde{H}_1(i) = H_1(i)^{\text{ad}}$. Let $S_1(i) := \text{Lie}(TH_1(i))$. Let

$$S_2(i) := \operatorname{Lie}(H_1) \cap (\bigoplus_{\alpha \in \Phi, \, \mathfrak{g}_{\alpha} \not\subset \operatorname{Lie}(H_1(i)_l)} \mathfrak{g}_{\alpha}).$$

We have a direct sum decomposition $\text{Lie}(H_1) = S_1(i) \oplus S_2(i)$ of $H_1(i)$ -modules.

We assume now that $H_1(i)^{\operatorname{ad}}$ is the $\operatorname{Res}_{l/k_1}$ of a PGL_2 group. Let $x \in \operatorname{Lie}(U_1) \cap \operatorname{Lie}(H_1(i))$. In the first four Cases (resp. in Case 5) we can write $x = x_1 + x_2$ (resp. $x = x_1 + x_2 + x_3$), where $x_i \in \mathfrak{g}_{\alpha_i}$ for some $\alpha_i \in \Phi^+$. This writing is unique up to ordering. The restriction of $\operatorname{ad}(x_i)^2$ to $S_2(i)$ is 0, cf. the $DT_2 = A_2$ part of 3.8.1. As $H_1(i)^{\operatorname{ad}}$ is the $\operatorname{Res}_{l/k_1}$ of a PGL_2 group we have $\operatorname{ad}(x_1)\operatorname{ad}(x_2) = \operatorname{ad}(x_2)\operatorname{ad}(x_1)$ (resp. $\operatorname{ad}(x_i)\operatorname{ad}(x_j) = \operatorname{ad}(x_j)\operatorname{ad}(x_i)$, $\forall i, j \in \{1, 2, 3\}$). Also in Case 5 we have $\operatorname{ad}(x_1)\operatorname{ad}(x_2)\operatorname{ad}(x_3) = 0$. So

$$h_x := \sum_{s=0}^{\infty} \frac{\operatorname{ad}(x)^s}{s!} = \sum_{s=0}^{2} \frac{\operatorname{ad}(x)^s}{s!}$$

is a well defined automorphism of $S_2(i)$. As in characteristic 0 we can identify h_x with the automorphism of $S_2(i)$ defined by an element of $H_1(i)(k_1)$.

We assume now that $LT = {}^2A_{2n}$. So $H_1(n)$ is of 2A_2 Lie type. Let $x \in \text{Lie}(H_1(n)) \cap \mathfrak{g}_{\alpha_n} \oplus \mathfrak{g}_{\alpha_{n+1}}$. We write $x = x_1 + x_2$, with $x_1 \in \mathfrak{g}_{\alpha_n}$ and $x_2 \in \mathfrak{g}_{\alpha_{n+1}}$. The restrictions of $\text{ad}(x_1)$ and $\text{ad}(x_2)$ to $S_2(n)$ do not commute but $\text{ad}(x_{i_1})\text{ad}(x_{i_2})\text{ad}(x_{i_3})$ restricted to $S_2(n)$ is $0, \forall i_1, i_2, i_3 \in \{1, 2\}$. So for $p \geq 3$ we can define h_x as above.

- **3.10. Theorem. 1)** If g.c.d.(p, o(H)) = 1, then no 1 dimensional k-vector subspace of Lie(H) is normalized by $AD_H(Im(H^{sc}(k) \to H(k)))$.
- 2) We assume that p|o(H) and g.c.d.(p, c(H)) = 1. If q = 2, we also assume that H is not an SL_2 group. Then there is no proper k-vector subspace V of Lie(H) normalized by $AD_H(Im(H^{sc}(k) \to H(k)))$ and such that Lie(H) = V + Lie(Z(H)).
- 3) We assume that $DT = D_{2n+2}$, that $o(H_1) = 2$ and that p = 2. Then there is no proper k-vector subspace V_0 of [Lie(H), Lie(H)] normalized by $AD_H(Im(H^{sc}(k) \to H(k)))$ and such that $[Lie(H), Lie(H)] = V_0 + Lie(Z(H))$.
- **4)** We assume that p divides both c(H) and o(H). If p = 2 we also assume that $DT \neq D_{2n+2}$. Then the adjoint representation of $H^{sc}(k)$ on Lie(H) is the direct sum of an irreducible, non-trivial representation and of a trivial representation.
- **5)** If p = 2 and $DT = C_n$ we assume that H_1 is adjoint. Then the $H_1(k_1)$ -module $Lie(H_1)/[Lie(H_1), Lie(H_1)]$ is trivial and of dimension over k_1 at most 2.

Proof: If $l \neq k_1$ we use the notations of 3.9. To prove 1) we can assume H is adjoint and $k_1 = k$. Let $u \in \text{Lie}(H)$ be such that ku is normalized by $AD_H(\text{Im}(H^{\text{sc}}(k) \to H(k)))$. The case $DT = A_1$ is well known (we can assume that q = p and cf. 3.4 that p = 2). So we can assume $DT \neq A_1$. We first assume $l \neq k$. Let $x \in \text{Lie}(H_1(i) \cap U_1)$, with $H_1(i)$ a semisimple subgroup of $H = H_1$ listed in the five Cases of 3.9. We write $u = u_1 + u_2$, where $u_s \in S_s(i)$ ($s \in \{1, 2\}$). The group $H_1(i)(k_1)$ normalizes ku_1 and ku_2 , cf. 3.9.3.

We assume now that $DT \neq A_{2n}$. So $H_1(i)$ is of isotypic A_1 Dynkin type. We show that $[x, u_2] = 0$. Let $a \in l \setminus \{0, 1\}$. If $H_1(i)^{\text{ad}}$ is a PGL_2 group, then the restriction of $\text{ad}(x)^2$ to $S_2(i)$ is 0 (cf. 3.8.1 applied to H_{1l}) and so the automorphism of $S_2(i)$ induced by $\sum_{s=0}^{1} \text{ad}(x)^s$ is defined by an element of $H_1(i)(k)$. So $u_2 + [x, u_2] \in ku_2$ and so as ad(x) is nilpotent we get $[x, u_2] = 0$. If $H_1(i)^{\text{ad}}$ is the $\text{Res}_{l/k}$ of a PGL_2 group, then as h_x and h_{ax} (of 3.9.3) normalize ku_2 we similarly get that $[x, u_2] = 0$. We now show that $[x, u_1] = 0$.

We use (8). If p > 2, then from 3.4 we get that the image of u_1 in $\text{Lie}(\tilde{H}_1(i))$ is 0; so $[x, u_1] = 0$. If p = 2, then $\tilde{H}_1(i)(k_1) = H_1(i)^{\text{ad}}(k_1)$ and as $\text{Lie}(H_1(i)^{\text{ad}})$ has no non-zero elements normalized by $H_1(i)^{\text{ad}}(k_1)$ (cf. the case $DT = A_1$) we get $[x, u_1] = 0$.

So [x, u] = 0. Similarly we get [y, u] = 0, where $y \in \text{Lie}(H_1(i) \cap U_1^{\text{opp}})$. So u is annihilated by \mathfrak{L}_H , cf. 3.9.2. So u as an element of $\text{Lie}(H_l)$ annihilates \mathfrak{g}_{α} , $\forall \alpha \in \Phi$. As $l \neq k$ none of the three conditions of 3.6 b) holds and so from 3.6 b) we get that the component of u in \mathfrak{g}_{α} with respect to (5) is $0, \forall \alpha \in \Phi$. So $u \in \text{Lie}(T_l)$. But as H is adjoint, no element of $\text{Lie}(T_l)$ annihilates \mathfrak{g}_{α} , $\forall \alpha \in \Phi$. So u = 0.

We assume now that $DT = A_{2n}$. If g.c.d.(p, 2n + 1) = 1, then the equality u = 0 is implied by 3.4. So we can assume $p \ge 3$. If $i \ne n$, then as above we get [x, u] = 0. If i = n we assume that x is as in the end of 3.9.3. As h_x and h_{2x} normalize ku_2 , as above we get $[x, u_2] = 0$. So as in the previous paragraph to show that u = 0 it is enough to show that $[x, u_1] = 0$. To show this we can assume n = 1 (so $\text{Lie}(H) = S_1(1)$ and $u = u_1$). As u is annihilated by $\mathfrak{g}_{\alpha_1+\alpha_2}$ (cf. the case i = n+1) from 3.6 b) we get that the component of u in \mathfrak{g}_{α} with respect to (5) is 0, $\forall \alpha \in \Phi$. So $u \in \text{Lie}(T)$. Let $g \in U_1(k_1) \setminus \mathbb{G}_{a,\alpha_1+\alpha_2}(k_1)$. As $AD_H(g)(ku) \subset ku$, we get that u is annihilated by \mathfrak{g}_{α_1} and \mathfrak{g}_{α_2} . So u = 0.

We are left to prove 1) in the case when l=k. If all roots of Φ have equal lengths, then the above part involving u_1 and u_2 for $DT \neq A_{2n}$ applies entirely. If DT is F_4 or G_2 , then $H^{\mathrm{ad}} = H^{\mathrm{sc}}$ and so [13, Hauptsatz] applies. So to prove 1) for l=k, we can assume $DT \in \{B_n, C_n | n \geq 2\}$. We can also assume p=2, cf. 3.4. As in the mentioned part involving u_1 and u_2 we get that $[u, \mathfrak{g}_{\beta}] = \{0\}$, provided $\beta \in \Phi$ is long. For any short root $\alpha \in \Phi$ there is a long root $\beta \in \Phi$ such that $\alpha + \beta \in \Phi$; so the component of u in \mathfrak{g}_{α} with respect to (5) is 0, cf. 3.6 b). For any long root $\beta \in \Phi$ there is a short root $\alpha \in \Phi$ such that $\alpha + \beta \in \Phi$ is also short. So the component of u in \mathfrak{g}_{β} is also 0, as otherwise the component of $AD_H(g)(u) - u$ in $\mathfrak{g}_{\alpha+\beta}$ is non-zero (cf. 3.6 c)); here $g \in \mathbb{G}_{a,\alpha}(k_1)$ is an arbitrary non-identity element. So $u \in \mathrm{Lie}(T)$. As H is adjoint, all $\tilde{H}(|\alpha|)$ groups are PGL_2 groups (cf. 3.8). So from (7) and the $DT = A_1$ case we get that $u \in \mathrm{Lie}(T(|\alpha|))$, $\forall \alpha \in \Phi$. As H is adjoint we have $\cap_{\alpha \in \Phi} \mathrm{Lie}(T(|\alpha|) = \{0\}$. So u = 0. This ends the proof of 1).

To prove 2) we can assume H is s.c. We assume such a V does exist and we show that this leads to a contradiction. We first show that we can assume $k_1 = k$. We write $H_{k_1} = \tilde{H}_1 \times_{k_1} \tilde{H}_2$, where \tilde{H}_1 and \tilde{H}_2 are normal, s.c. subgroups of H_{k_1} , the adjoint of \tilde{H}_1 being simple. Let $\tilde{V}_1 := \operatorname{Ker}(V \otimes_k k_1 \to \operatorname{Lie}(\tilde{H}_2^{\operatorname{ad}}))$. Any trivial $\tilde{H}_1(k_1)$ -submodule of \tilde{V}_1 is contained in $\operatorname{Lie}(Z(H_{k_1}))$, cf. 1). We get $\tilde{V}_1 = \tilde{V}_1 \cap \operatorname{Lie}(\tilde{H}_1) + \tilde{V}_1 \cap \operatorname{Lie}(Z(H_{k_1}))$. From this and the inclusion $\operatorname{Lie}(\tilde{H}_1) \subset \tilde{V}_1 + \operatorname{Lie}(Z(H_{k_1}))$ we get $\operatorname{Lie}(\tilde{H}_1) = \tilde{V}_1 \cap \operatorname{Lie}(\tilde{H}_1) + \operatorname{Lie}(Z(\tilde{H}_1))$. So if we have $\operatorname{Lie}(\tilde{H}_1) = \tilde{V}_1 \cap \operatorname{Lie}(\tilde{H}_1)$ for any such \tilde{H}_1 , then $V \otimes_k k_1 = \operatorname{Lie}(H_{k_1})$ and this contradicts the fact that V is a proper k-vector subspace of $\operatorname{Lie}(H)$. So we can assume $k_1 = k$; so $H = H_1$. If DT is not (resp. is) D_{2n+2} , then $\dim_k(\operatorname{Lie}(Z(H)))$ is 1 (resp. is 2). So the number $\dim_k(\operatorname{Lie}(H)/V)$ is 1 (resp. is 1 or 2). We can assume $\dim_k(\operatorname{Lie}(H)/V) = 1$.

We consider first the case $p \ge 3$. So either $DT = A_{pn-1}$ or p = 3 and $DT = E_6$. If l = k, then $\dim_k(\operatorname{Lie}(H(|\alpha|)) \cap V) \ge 2$ and so $\operatorname{Lie}_k(H(|\alpha|)) \cap V = \operatorname{Lie}_k(H(|\alpha|))$ (cf. 3.4), $\forall \alpha \in \Phi$. So $\operatorname{Lie}(H) = \mathfrak{L}_H \subset V$, cf. 3.7 1) and 2). Contradiction. If $l \ne k$, then $l = k_2$. Using similar intersections $V \cap \operatorname{Lie}_k(H_1(i))$ we get (cf. Cases 1, 2 and 3 of 3.9) that to prove $V = \operatorname{Lie}(H)$ we can assume that $DT = A_2$, p = 3 and we just need to show

that $\operatorname{Lie}(H_1(2)) \subset V$. The $H_1(2)(k)$ -module $\operatorname{Lie}(H)$ is semisimple: it is the direct sum of $\operatorname{Lie}(H_1(2))$ with $\operatorname{Lie}(Z(H))$ and with two k-vector spaces of dimension 2 included in $\mathfrak{g}_{-\alpha_1} \oplus \mathfrak{g}_{-\alpha_2} \oplus \mathfrak{g}_{\alpha_1} \oplus \mathfrak{g}_{\alpha_2}$. So as V is normalized by $H_1(2)(k)$, by reasons of dimensions we get that V contains these two k-vector spaces and $\operatorname{Lie}(H_1(2))$. Let $g \in (U_1 \setminus H_1(2))(k)$. Let $x \in V \cap (\mathfrak{g}_{-\alpha_1} \oplus \mathfrak{g}_{-\alpha_2} \oplus \mathfrak{g}_{\alpha_1} \oplus \mathfrak{g}_{\alpha_2}) \setminus \{0\}$. It is easy to see that we can choose g and x such that the component of $AD_H(g)(x) - x$ in $\operatorname{Lie}(Z(H))$ is non-zero. So as $AD_H(g)(x) - x \in V$ we get $\operatorname{Lie}(Z(H)) \subset V$. So $\operatorname{Lie}(H_1(2)) \subset V$. So $V = \operatorname{Lie}(H)$. Contradiction.

We are left with the case p=2. The case $DT \in \{A_1, E_7\}$ follows from [13, Haupt-satz]. So we can assume $DT \in \{A_{2n-1}, B_n, C_n | n \geq 2\} \cup \{D_{n+3} | n \in \mathbb{N}\}$. As in the previous paragraph we just need to show that $\operatorname{Lie}_k(H_1(i)) \cap V = \operatorname{Lie}_k(H_1(i))$, for any $H_1(i)$ as in 3.9. The case $l=k \neq \mathbb{F}_2$ is as above. If $l=k=\mathbb{F}_2$, then 2) follows from loc. cit. So we can assume $l \neq k$. So DT is A_{2n-1} or D_{n+3} and we are in one of the Cases 1, 4 or 5 of 3.9. In Case 4 all $H_1(i)$'s are $\operatorname{Res}_{k_2/k}$ of SL_2 groups, cf. 3.9.1; so as $l=k_2 \neq \mathbb{F}_2$ we get $\operatorname{Lie}_k(H_1(i)) \cap V = \operatorname{Lie}_k(H_1(i))$. The same applies to Cases 1 and 5, except for the situation when $l=k_3$ and $k_1=k=\mathbb{F}_2$; so $DT=D_4$. We now refer to this situation and we use the notations of Case 5 of 3.9. It is enough to show that $V \cap \operatorname{Lie}(H_1(2)) = \operatorname{Lie}(H_1(2))$. As $H_1(2)$ is an SL_2 group (cf. 3.9.1), we have $\{0, x+h, y+h, x+y\} \subset V \cap \operatorname{Lie}(H_1(2))$, where x, y and h are the non-zero elements of $\operatorname{Lie}(H_1(2))$ such that $x \in \mathfrak{g}_{\alpha_1+\alpha_2+\alpha_3+\alpha_4}$, $y \in \mathfrak{g}_{-\alpha_1-\alpha_2-\alpha_3-\alpha_4}$ and $h \in \operatorname{Lie}(T)$. If $g \in \mathbb{G}_{a,\alpha_2}(l) \cap H(k)$ is a non-identity element, then $u := AD_H(g)(x+y) - (x+y)$ is the non-zero element of $\mathfrak{g}_{\alpha_1+2\alpha_2+\alpha_3+\alpha_4} \cap \operatorname{Lie}(H)$ (cf. 3.6 c)). Similarly, if $w \in \mathbb{G}_{a,-\alpha_2}(l) \cap H(k)$ is the non-identity element, then $x = AD_H(w)(u) - u$. So $V \cap \operatorname{Lie}(H_1(2))$ contains x. So $V \cap \operatorname{Lie}(H_1(2)) = \operatorname{Lie}(H_1(2))$. This ends the proof of 2).

We prove 3). As $Z(H_1^{\text{sc}}) = \mu_2^2$ and $Z(H_1) = \mu_2$ we have $\dim_{k_1}(\text{Lie}(H_1^{\text{ad}})/\mathfrak{L}_{H_1^{\text{ad}}}) = 2$ and $\dim_{k_1}(\text{Lie}(H_1)/\mathfrak{L}_{H_1}) = 1$. So $\text{Lie}(Z(H_1)) \subset \mathfrak{L}_{H_1}$. So 3) follows from 2) applied to the inverse image V of V_0 in $\text{Lie}(H^{\text{sc}})$. We prove 4). Either $DT = A_{p^2n-1}$ or p = 2 and $DT = D_{2n+3}$. We have a direct sum decomposition of H-modules $\text{Lie}(H) = \text{Lie}(Z(H)) \oplus [\text{Lie}(H), \text{Lie}(H)] = \text{Lie}(Z(H)) \oplus [\text{Lie}(H^{\text{ad}}), \text{Lie}(H^{\text{ad}})]$, cf. [18, 0.13]. So 4) follows from 3.7.1. To prove 5) we can assume l = k. We have $\dim_k(\text{Lie}(H)/\mathfrak{L}_H) \le 2$ and the equality can take place only for p = 2, cf. loc. cit. As $\dim_k(H) \ge 3$ and as for p = 2 the group H is not an SL_2 group, the representation of H on $\text{Lie}(H)/\mathfrak{L}_H$ is trivial. So H(k) acts trivially on $\text{Lie}(H)/\mathfrak{L}_H$. This ends the proof.

3.11. On exceptional ideals. Until 3.12 we assume that H is adjoint and that either p=2 and $DT \in \{B_n, C_n | n \in \mathbb{N}\} \cup \{F_4\}$ or p=3 and $DT=G_2$. So $l=k_1$. It is known that H_1 is the extension to k_1 of a split, adjoint group H_0 over k, cf. [27, Vol. III, p. 410]. Let I (resp. I^{sc}) be the ideal of $\text{Lie}(H_1)$ (resp. $\text{Lie}(H_1^{\text{sc}})$) generated by \mathfrak{g}_{α} , with $\alpha \in \Phi$ short. We have $I=I_0 \otimes_k k_1$ (resp. $I^{\text{sc}} = I_0^{\text{sc}} \otimes_k k_1$), with I_0 (resp. I_0^{sc}) as the similarly defined ideal of $\text{Lie}(H_0)$ (resp. $\text{Lie}(H_0^{\text{sc}})$). As I^{sc} and I are normalized by H_1 we have $I=\text{Im}(I^{\text{sc}} \to \text{Lie}(H_1))$. So I is an a.s. $H_1(k_1)$ -module, cf. [13, Hauptsatz]. Similarly, I_0 is an a.s. $H_0(k_0)$ -module. Let I_k (resp. I_k^{sc}) be I (resp. I^{sc}) but viewed as a k-ideal or as a k-vector subspace of Lie(H).

3.11.1. Proposition. The only simple H(k)-submodule of Lie(H) is I_k .

Proof: Let $I_{H_0} \otimes_k k_1 : H_0(k) \to GL(I_0 \otimes_k k_1)(k)$ be the representation defined by the restriction of AD_H to $H_0(k) \triangleleft H_0(k_1) = H_1(k_1) = H(k)$. It is semisimple, being a direct

sum of $[k_1:k]$ copies of the irreducible representation $I_{H_0}:H_0(k)\to GL(I_0)(k)$ defined by AD_{H_0} . As I_0 is an a.s. $H_0(k_0)$ -module, any non-trivial subrepresentation of $I_{H_0}\otimes_k k_1$ is of the form $IRR=I_0\otimes_k W$, with W a non-trivial k-vector subspace of k_1 . We choose B_1 and T_1 such that T_1 is the extension to k_1 of a maximal torus T_0 of H_0 . So $T_1(k_1)=T_0(k_1)$ is included in $H_0(k_1)$. So W is stable under multiplications with elements of k_1^* which belong to the image of the homomorphism $T_1(k_1)\to \mathbb{G}_m(k_1)$ associated to a character $T_1\to \mathbb{G}_m$ defining a short root of Φ . So W is k_1^* -stable and so $W=k_1$. So $IRR=I_0\otimes_k k_1$. So I_k is a simple H(k)-module. From loc. cit. and (4) we get that any simple H(k)-submodule of \mathfrak{L}_H \mathfrak{L}_H is contained in $I_k\otimes_k k_1$. So I_k is the only simple H(k)-submodule of \mathfrak{L}_H . Any other simple H(k)-submodule of Lie(H) would be trivial (cf. 3.10 5)) and so does not exist (cf. 3.10 1)). This ends the proof.

- **3.11.2. Theorem. 1)** If DT is B_n or C_n we assume n is odd and at least 3. Then the H(k)-module \mathfrak{L}_H/I_k is simple and non-trivial. Moreover, the short exact sequence of H(k)-modules $0 \to I_k \to \mathfrak{L}_H \to \mathfrak{L}_H/I_k \to 0$ does not split.
- **2)** If $DT = C_n$ with n odd and at least 3, then there are no 1 dimensional k-vector subspaces of $Lie(H)/I_k$ normalized by H(k).
- 3) Let $H_1^{\rm sclong}$ be the analogue of $H_1^{\rm long}$ of 3.9 but for $H_1^{\rm sc}$ instead of H_1 . We assume $DT = C_n$ (resp. $DT = B_n$). If n = 1 we also assume $k \neq \mathbb{F}_2$. Then $[Lie(H^{\rm sc}), Lie(H^{\rm sc})]$ (resp. $I_k + Lie_k(Z(H_1^{\rm sclong}))$) is the only maximal H(k)-submodule of $Lie(H^{\rm sc})$.

Proof: The first part of 1) is argued as in 3.11.1 for I_k . The second part of 1) follows from 3.11.1. To prove 2) we can assume $k = k_1$. So $H = H_1$ and $I = I_k$. The group $H^{\text{long}} := H_1^{\text{long}}$ is the quotient of the group $SL(H) = SL_2^n$ of the proof of 3.7 by $Z(H^{\text{sc}})$. Let T_{sc} be the inverse image of T in H^{sc} . We have $\dim_k(\text{Lie}(H^{\text{sc}})/I^{\text{sc}}) = 2n$, cf. [13, Hauptsatz]. This implies $\text{Lie}(T_{\text{sc}}) \subset I^{\text{sc}}$ and so Lie(H)/I (resp. \mathfrak{L}_H/I) is 2n+1 (resp. 2n) dimensional. So Lie(H)/I is identified naturally with a Lie subalgebra of $\text{Lie}((H^{\text{long}})^{\text{ad}})$ in such a way that \mathfrak{L}_H/I corresponds to $\mathfrak{L}_{(H^{\text{long}})^{\text{ad}}}$. But $\text{Lie}((H^{\text{long}})^{\text{ad}})$ has no 1 dimensional k-vector subspace normalized by the subgroup $H^{\text{long}}(k)$ of H(k), cf. 3.10 1). So 2) holds.

We now prove 3). We deal just with the $DT = C_n$ case as the $DT = B_n$ case is entirely the same. The H(k)-module $\text{Lie}(H^{\text{sc}})/[\text{Lie}(H^{\text{sc}}), \text{Lie}(H^{\text{sc}})]$ is simple. Argument: the case n=1 follows from 3.7.1 and the case n>1 is argued in the same way 3.11.1 was, the only difference being that $T_1 \to \mathbb{G}_m$ defines this time a long root of Φ . Let M be an H(k)-submodule of $\text{Lie}(H^{\text{sc}})$ not contained in $[\text{Lie}(H^{\text{sc}}), \text{Lie}(H^{\text{sc}})]$. It is enough to show that $M = \text{Lie}(H^{\text{sc}})$. We show that the assumption $M \neq \text{Lie}(H^{\text{sc}})$ leads to a contradiction. Let \tilde{M} be a maximal $H_1(k_1)$ -submodule of $\text{Lie}(H^{\text{sc}}) = \text{Lie}(H^{\text{sc}}) \otimes_k k_1$ containing $M \otimes_k k_1$. From (4) and loc. cit. we get that $[\text{Lie}(H^{\text{sc}}), \text{Lie}(H^{\text{sc}})] \otimes_k k_1 \subset \tilde{M}$. As M surjects onto $\text{Lie}(H^{\text{sc}})/[\text{Lie}(H^{\text{sc}}), \text{Lie}(H^{\text{sc}})]$, \tilde{M} surjects onto $(\text{Lie}(H^{\text{sc}})/[\text{Lie}(H^{\text{sc}}), \text{Lie}(H^{\text{sc}})]) \otimes_k k_1$. So $\tilde{M} = \text{Lie}(H^{\text{sc}})$. Contradiction. So $M = \text{Lie}(H^{\text{sc}})$. This ends the proof.

From 3.7.1 and 3.11.2 3) we get:

3.12. Corollary. In general, the $H^{\text{sc}}(k)$ -module $\mathfrak{L}_{H^{\text{ad}}}$ has a unique maximal submodule $\mathfrak{I}_{H^{\text{ad}}}$. Moreover, the $H^{\text{sc}}(k)$ -module $\mathfrak{L}_{H^{\text{ad}}}/\mathfrak{I}_{H^{\text{ad}}}$ is non-trivial.

$\S 4$. The *p*-adic context

Let $k = \mathbb{F}_q$ and $s \in \mathbb{N}$. Let G be a reductive group scheme over W(k). Let \mathbb{A}^1 be the affine line over $B(k) := W(k)[\frac{1}{p}]$. The group G(B(k)) is endowed with the coarsest topology making all maps $G(B(k)) \to \mathbb{A}^1(B(k)) = B(k)$ induced by morphisms $G_{B(k)} \to \mathbb{A}^1$ to be continuous. Let K be a closed subgroup of G(W(k)).

4.1. Problem. Find practical conditions on G, K and p which imply K = G(W(k)).

This Problem was first considered in the context of SL_n groups over \mathbb{Z}_p in [24, Ch. IV]. After mentioning two general properties (see 4.1.1 and 4.1.2), in 4.1.3 and 4.1.4 we identify a key subpart (question) of this Problem. In 4.2 we list some simple properties of G. In 4.3 we include an approach of inductive nature which allows to get information on $\text{Im}(K \to G(W_2(k)))$ from information on semisimple subgroups of G. In 4.4 to 4.7 we assume G is semisimple. Section 4.5 solves question 4.1.4 for adjoint and s.c. groups G. The computations needed for this are gathered in 4.4. In 4.6 we include a supplement to 4.5 related to the exceptional ideals. In 4.7 we prove 1.3.

We now point out that in general we do need some conditions on G, K and p.

4.1.1. Example. We assume there is an isogeny $f: G^1 \to G$ over W(k) of order p^n . So $\operatorname{Ker}(f)$ is a finite, flat subgroup of $Z(G^1)$ of multiplicative type (see [27, Vol. III, 4.1.7 of p. 173]). Being of order p^n it is connected. So the Lie homomorphism $\operatorname{Lie}(G_k^1) \to \operatorname{Lie}(G_k)$ has non-trivial kernel and so it is not an epimorphism. Moreover, the group $\operatorname{Ker}(f)(\bar{k})$ is trivial and so the homomorphism $f(k): G^1(k) \to G(k)$ is an isomorphism. So the image of the homomorphism $f(W(k)): G^1(W(k)) \to G(W(k))$ is a proper subgroup of G(W(k)) surjecting onto G(k).

We identify $\operatorname{Ker}(G(W_{s+1}(k)) \to G(W_s(k)))$ and $\operatorname{Lie}_{\mathbb{F}_p}(G_k)$ as G(k)-modules. Let $L_{s+1} := \operatorname{Im}(K \to G(W_{s+1}(k))) \cap \operatorname{Ker}(G(W_{s+1}(k)) \to G(W_s(k))).$

4.1.2. Lemma. If p = 2 (resp. p > 2) let m := 3 (resp. m := 2). If K surjects onto $G(W_m(k))$, then K = G(W(k)).

Proof: This is just the generalization of [24, Exc. 1 p. IV-27]. We show that in the abstract context of G we can appeal to matrix computations as in loc. cit. Let $\rho_{B(k)}: G_{B(k)} \hookrightarrow GL(W)$ be a faithful representation, with W a finite dimensional B(k)-vector space. So $\rho_{B(k)}$ extends to a representation $\rho: G \to GL(L)$, with L a W(k)-lattice of W (cf. [19, 10.4 of Part I]). The morphism ρ is a closed embedding, cf. [30, c) of Proposition 3.1.2.1]. So each element $g \in \text{Ker}(G(W(k)) \to G(W_s(k)))$ mod p^{s+1} is congruent to $1_L + p^s x$, where $x \in \text{Lie}(G) \subset \text{End}(L)$. So if $s \geq 3$ (resp. $s \geq 2$), then the image of g in $G(W_{s+1}(k))$ is the p-th power of any element of $G(W_{s+1}(k))$ whose image in $G(W_s(k))$ is the same as the one of $1_L + p^{s-1}x \mod p^s$. So by induction on $s \geq 3$ (resp. on $s \geq 2$) we get that $L_{s+1} = \text{Ker}(G(W_{s+1}(k)) \to G(W_s(k)))$ and so that K surjects onto $G(W_{s+1}(k))$. As s is arbitrary and K is compact we get K = G(W(k)). This ends the proof.

If p=2, then 4.1.2 does not hold in general if we replace m=3 by m=2. Example: $G=\mathbb{G}_m$ and q=2. See 4.7.1 below for more examples.

4.1.3. The class γ_G . We view $\text{Lie}_{\mathbb{F}_p}(G_k)$ as a G(k)-module via AD_{G_k} . Let

$$\gamma_G \in H^2(G(k), \mathrm{Lie}_{\mathbb{F}_n}(G_k))$$

be the class defining the standard short exact sequence

(9)
$$0 \to \operatorname{Lie}_{\mathbb{F}_p}(G_k) \to G(W_2(k)) \to G(k) \to 0.$$

Lemma 4.1.2 points out that for the study of 4.1 we need to understand what possibilities for L_2 (resp. for L_2 and L_3) we have if p > 2 (resp. if p = 2). The study of L_2 and so of 4.1 is intimately interrelated to the following question.

- **4.1.4. Question.** When does the short exact sequence (9) have a section (i.e. when is $\gamma_G = 0$)?
- **4.2.** Proposition. 1) Any maximal torus T_k of a Borel subgroup B_k of G_k lifts to a maximal torus T of G. Moreover, there is a unique Borel subgroup B of G containing T and having B_k as its special fibre.
- 2) The map $G \to G_k$ establishes a bijection between isomorphism classes of reductive (resp. semisimple) group schemes over W(k) and isomorphism classes of reductive (resp. semisimple) group schemes over k.
- **3)** If G is an adjoint (resp. a s.c. semisimple) group, then it is a product of Weil restrictions of a.s. adjoint groups (resp. of s.c. semisimple groups having a.s. adjoints) over Witt rings of finite field extensions of k. So if G^{ad} is simple, then G is of isotypic Dynkin type.

Proof: As W(k) is complete, the first part of 1) follows from [27, Vol. II, p. 47–48]: by induction on $s \in \mathbb{N}$ we lift T_k to a maximal torus $T_{W_s(k)}$ of $G_{W_s(k)}$. To prove the second part of 1) we can assume that T is split and so the existence and the uniqueness of B follow from [27, Vol. III, 5.5.1 and 5.5.5 of Exp. XXII]. See [27, Vol. III, Prop. 1.21 of p. 336–337] for 2). Part 3) follows from 2.3.2 and 2).

- **4.3.** An inductive approach. Until end let T and B be as in 4.2 1). Let G_{0k} be a semisimple subgroup of G_k normalized by T_k . Let T_{0k} (resp. T_{00k}) be the maximal subtorus of T_k which is a torus of G_{0k} (resp. which centralizes G_{0k}). It lifts uniquely to a subtorus T_0 (resp. T_{00}) of T, cf. [27, Vol. II, p. 47–48].
- **4.3.1. Lemma.** The group G_{0k} lifts to at most one closed, semisimple subgroup G_0 of G normalized by T.

Proof: Let l be the smallest field extension of k such that T_l is split. So $T_{W(l)}$ is also split. We consider the Weyl decomposition

(10)
$$\operatorname{Lie}(G_{W(l)}) = \operatorname{Lie}(T_{W(l)}) \bigoplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha}$$

of $\operatorname{Lie}(G_{W(l)})$ with respect to $T_{W(l)}$. Warning: whenever T is split (i.e. k=l) we drop the left index W(l). Let $\Phi_0 := \{\alpha \in \Phi | \mathfrak{g}_{\alpha} \otimes_{W(l)} l \subset \operatorname{Lie}(G_{0l})\}$. It suffices to prove the Lemma under the extra assumptions that T is split and that T_{00} is trivial (otherwise we replace G by the derived subgroup of the centralizer of T_{00} in G). So G is semisimple and the kernel of the natural homomorphism from T_k into the identity component $\operatorname{Aut}^0(G_{0k})$ of the group scheme of automorphisms of G_{0k} is finite. We have $\operatorname{Aut}^0(G_{0k}) = G_{0k}^{\operatorname{ad}}$, cf. [27, Vol. III, Prop. 2.15 of p. 343]. So T_k and G_{0k} have equal ranks. More precisely, T_k is the maximal torus of G_{0k} whose image in $Aut^0(G_{0k})$ is the same as of T_k . So $T_0 = T$. As G_{0k} is a semisimple group having T_k as a maximal torus, Φ_0 is symmetric and invariant under the natural action of the group $Aut_{W(k)}(W(l))$ on Φ . If G_0 exists, then it is generated by T and by the \mathbb{G}_a subgroups of G normalized by T and whose Lie algebras are \mathfrak{g}_{α} , with $\alpha \in \Phi_0$. So as Φ_0 is determined uniquely by G_{0k} , the Lemma follows.

4.3.2. Proposition. There is a closed, semisimple subgroup G_0 of G lifting G_{0k} iff Φ_0 is closed under addition, i.e. we have $\Phi \cap \{\alpha + \beta | \alpha, \beta \in \Phi_0\} \subset \Phi_0$. So G_0 exists if for any simple factor H of G_k^{ad} none of the three conditions of 3.6 b) holds.

Proof: We can assume that T is split and T_{00} is trivial. So T is a maximal torus of G_0 . If G_0 exists, then the direct sum $\operatorname{Lie}(T) \bigoplus \bigoplus_{\alpha \in \Phi_0} \mathfrak{g}_{\alpha}$ is a Lie subalgebra of $\operatorname{Lie}(G)$. From Chevalley rule in characteristic 0 (see [27, Vol. III, 6.5 of p. 322]) we get that Φ_0 is closed under addition. So G_0 exists and is a reductive group scheme, cf. [27, Vol. III, 5.3.4, 5.4.7 and 5.10.1 of Exp. XXII]. So G_0 is semisimple as G_{0k} is so. We assume now that for any simple factor H of G_k^{ad} none of the three conditions of 3.6 b) holds. So as $\operatorname{Lie}(G_{0k})$ is a Lie subalgebra of $\operatorname{Lie}(G_k)$, from the equality part of 3.6 b) we get that Φ_0 is closed under addition. So G_0 exists. This ends the proof.

We assume G_0 exists. So $\Phi_0 = -\Phi_0$ is closed under addition. Let TG_0 be the closed subgroup of G generated by T and G_0 . It is a reductive group scheme, cf. loc. cit.

4.3.3. Lemma. We have a unique direct sum decomposition $Lie(G) = Lie(TG_0) \oplus V_0$ of TG_0 -modules. The resulting direct sum decomposition $Lie_{\mathbb{F}_p}(G_k) = Lie_{\mathbb{F}_p}(TG_{0k}) \oplus V_0/pV_0$ is $G_0(k)$ -invariant.

Proof: To check the first part we can assume that T is split and G is semisimple. As Φ_0 is closed under addition, $V_0 := \bigoplus_{\alpha \in \Phi \setminus \Phi_0} \mathfrak{g}_{\alpha}$ is a $\operatorname{Lie}(TG_0)$ -module and so a TG_0 -module. But V_0 is the unique supplement of $\operatorname{Lie}(TG_0)$ in $\operatorname{Lie}(G)$ normalized by T, cf. (10). So the first part holds. The second part follows from the first part. This ends the proof.

4.3.4. Key Lemma. If $\gamma_{G_0^{\text{ad}}} \neq 0$, then $\gamma_G \neq 0$.

Proof: It is enough to show that if $\gamma_G = 0$, then $\gamma_{G_0^{\operatorname{ad}}} = 0$. Let $\gamma_{G_0,G}^0 \in H^2(G_0(k), \operatorname{Lie}_{\mathbb{F}_p}(G_k))$ be the restriction of γ_G via the monomorphism $G_0(k) \hookrightarrow G(k)$. The component of $\gamma_{G_0,G}^0$ in $H^2(G_0(k), \operatorname{Lie}_{\mathbb{F}_p}(TG_{0k}))$ with respect to the decomposition of the second part of 4.3.3, is the 0 class. So $TG_0(W_2(k))$ has a subgroup S_0 mapping isomorphically into $G_0(k)$. The order o of $Z(G_0)(k)$ is prime to p. So the image of S_0 in $(TG_0)^{\operatorname{ad}}(W_2(k)) = G_0^{\operatorname{ad}}(W_2(k))$ is a subgroup mapping isomorphically into $\operatorname{Im}(G_0(k) \to G_0^{\operatorname{ad}}(k))$. The index of this last group in $G_0^{\operatorname{ad}}(k)$ is o. So as $\operatorname{Lie}_{\mathbb{F}_p}(G_{0k}^{\operatorname{ad}})$ is a p-group, we get $\gamma_{G_0^{\operatorname{ad}}} = 0$. This ends the proof.

4.3.5. Lemma. We assume that q=2 and $\gamma_G \neq 0$. Then $\gamma_{G_W(\mathbb{F}_4)} \neq 0$.

Proof: Let $a \in \mathbb{F}_4 \setminus k$. We have a direct sum decomposition $\operatorname{Lie}_{\mathbb{F}_2}(G_{\mathbb{F}_4}) = \operatorname{Lie}(G_k) \oplus a\operatorname{Lie}(G_k)$ of G(k)-modules. So the restriction of $\gamma_{G_{W(\mathbb{F}_4)}}$ via the monomorphism $G(k) \hookrightarrow G(\mathbb{F}_4)$ is the direct sum of γ_G and of a class in $H^2(G(k), a\operatorname{Lie}(G_k))$. So $\gamma_{G_{W(\mathbb{F}_4)}} \neq 0$.

4.4. Computations. Until end we assume G is semisimple. For $y \in W_2(k)$ let $\bar{y} \in k$ be its reduction mod p. For $\alpha \in k^*$ let $t_{\alpha} \in W_2(k)$ be the reduction mod p^2 of the Teichmüller

lift of α . We start answering 4.1.4 for adjoint and s.c. groups. In 4.4.1 we include a general Proposition which among other things allows us to assume that $q \leq 4$. In 4.4.2 (resp. 4.4.3 to 4.4.9) we deal with four (resp. seven) special cases corresponding to $q \leq 4$ and s.c. (resp. a.s. adjoint) groups G of small rank. All of them are minimal cases in the sense that 4.4.1 does not apply to them and moreover for any closed, semisimple subgroup G_0 of G normalized by T and different from G, we have $\gamma_{G_0^{\rm ad}} = 0$. Sections 4.3 and 4.4 allow us to list in 4.5 all cases when $\gamma_G = 0$ and G is adjoint or s.c. In other words, all other minimal such cases are "handled" by 4.4.5, 4.3.5 and the available literature (see proof of 4.5).

The identity (resp. zero) $n \times n$ matrix is denoted as I_n (resp. as 0_n). For $i, j \in \{1, ..., n\}$ let E_{ij} be the $n \times n$ matrix whose all entries are 0 except the ij entry which is 1. We list matrices by rows in increasing order (the first row, then the second row, etc.).

4.4.1. Proposition. We have $\gamma_G \neq 0$ if any one of the following three conditions hold:

- $q \ge 5$;
- -q=3 and G_k^{ad} has a non-split simple factor which is not a PGU_3 group;
- $-q \in \{2,4\}$ and G_k^{ad} has simple factors which do not split over \mathbb{F}_4 and are not the $Res_{W(\mathbb{F}_4)/W(k)}$ of a PGU_3 group.

Proof: We can assume G^{ad} is adjoint and simple. Our hypotheses imply that G has a closed subgroup G_0 normalized by T and isogenous to the Weil restriction of an SL_2 group over the Witt ring of a finite field with at least 5 elements. Argument: we can assume G is a.s. (cf. 4.2 3)) and we can deal just with special fibres (cf. 4.2 2) and 4.3.2); so the statement follows from 3.9 and 4.3.2. So based on 4.3.4, if $p \ge 3$ (resp. p = 2) it suffices to show that $\gamma_G \ne 0$ for $q \ge 5$ and G an SL_2 (resp. a PGL_2) group. If $p \ge 5$ and $g \in M_2(W_2(k))$, then $(I_2 + E_{12} + py)^p = I_2 + pE_{12} \ne I_2$. So $G(W_2(k))$ has no element of order p specializing to a non-identity k-valued point of a \mathbb{G}_a subgroup of G. So $\gamma_G \ne 0$ if $p \ge 5$.

Let now p=3; so $r\geq 2$. For $\alpha\in k^*$ let $X_\alpha:=I_2+t_\alpha E_{12}+3x_\alpha E_{21}+3Y_\alpha\in M_2(W(k))$, with $x_\alpha\in W_2(k)$ and with $Y_\alpha\in M_2(W_2(k))$ such that its 21 entry is 0. Let $\beta\in k^*$. The matrix equations $X_\alpha^3=X_\beta^3=I_2$ and $X_\alpha X_\beta=X_\beta X_\alpha$ inside $M_2(W_2(k))$, get translated into equations with coefficients in k involving the reductions mod p of x_α , x_β and of the entries of Y_α and Y_β . We need just few of these equations. Identifying the 12 (resp. 11) entries of the equations $X_\alpha^3=X_\beta^3=I_2$ (resp. $X_\alpha X_\beta=X_\beta X_\alpha$) we get $1+\alpha \bar{x}_\alpha=1+\beta \bar{x}_\beta=0$ (resp. $\alpha \bar{x}_\beta=\beta \bar{x}_\alpha$). So $\alpha^2=\beta^2$. But as $r\geq 2$ there are elements α , $\beta\in k^*$ such that $\alpha^2\neq\beta^2$. So $\gamma_G\neq 0$ if G is an SL_2 group, p=3 and $r\geq 2$.

Let now p = 2; so $r \ge 3$. As the homomorphism $GL_2(W_2(k)) \to PGL_2(W_2(k))$ is an epimorphism, we can use again matrix computations. For $\delta \in k$ let

$$X_{\delta} := I_2 + t_{\delta} E_{12} + 2Y_{\delta},$$

with $Y_{\delta} \in M_2(W_2(k))$. Let (a_{δ}, b_{δ}) and (c_{δ}, d_{δ}) be the rows of Y_{δ} . As $r \geq 3$ we can choose $\alpha, \beta \in k \setminus \{0, 1\}$ such that $1 + \alpha + \beta \neq 0$ and $\alpha \neq \beta$. The conditions that the images of $X_1, X_{\alpha}, X_{\beta}$ in $PGL_2(W_2(k))$ are of order 2 and commute with each other imply that we have the power 2 equations $X_1^2 = (1 + 2v_1)I_2, X_{\alpha}^2 = (1 + 2v_{\alpha})I_2$ and $X_{\beta}^2 = (1 + 2v_{\beta})I_2$ and the commuting equations $X_1X_{\alpha} = (1 + 2v_{1\alpha})X_{\alpha}X_1, X_1X_{\beta} = (1 + 2v_{1\beta})X_{\beta}X_1$ and

 $X_{\alpha}X_{\beta} = (1 + 2v_{\alpha\beta})X_{\beta}X_{\alpha}$, where v_1 , v_{α} , v_{β} , $v_{1\alpha}$, $v_{1\beta}$ and $v_{\alpha\beta}$ belong to $W_2(k)$. Among the equations we get are the following ones:

(11)
$$\bar{a}_1 + \bar{d}_1 = \bar{a}_{\alpha} + \bar{d}_{\alpha} = \bar{a}_{\beta} + \bar{d}_{\beta} = 1;$$

$$\bar{v}_{1\alpha} = \bar{v}_{1\beta} = \bar{v}_{\alpha\beta} = 1;$$

(13)
$$1 + \bar{c}_{\alpha} + \alpha \bar{c}_{1} = 1 + \bar{c}_{\beta} + \beta \bar{c}_{1} = 1 + \alpha \bar{c}_{\beta} + \beta \bar{c}_{\alpha} = 0.$$

The equations (11) are obtained by identifying the 12 entries of the power 2 equations (we have $X_{\delta}^2 = I_2 + 2t_{\delta}E_{12} + 2t_{\delta}[Y_{\delta}, E_{12}]$, etc.). The equations (12) are obtained by identifying the 12 entries of the commuting equations and by inserting (11) into the resulting equations. The equations (13) are obtained by identifying the 11 entries of the commuting equations and by inserting (12) into the resulting equations. As $1 + \alpha + \beta \neq 0$, the system (13) of equations in the variables \bar{c}_1 , \bar{c}_{α} and \bar{c}_{β} has no solution in k. So $\gamma_G \neq 0$ if p = 2, $r \geq 3$ and G is a PGL_2 group. This ends the proof.

4.4.2. Applications. Let $W_0 := W(\mathbb{F}_4)^2$. We refer to the proof of 4.4.1 with q = 4, $\alpha \in k \setminus \mathbb{F}_2$ and without mentioning β . The equations $X_1^2 = X_{\alpha}^2 = I_2$ and $X_1 X_{\alpha} = X_{\alpha} X_1$ have no solution, cf. the value of $v_{1\alpha}$ in (12). So the class $\gamma_4 \in H^2(SL(W_0)(k), \text{Lie}_{\mathbb{F}_2}(GL(W_0/pW_0)))$ defined naturally by the inverse image of $SL(W_0)(k)$ in $GL(W_0)(W_2(k))$ is non-zero. So also $\gamma_G \neq 0$ for G an SL_2 group over W(k). For most applications 4.3.4 suffices. We now use γ_4 to get a variant of 4.3.4 in two situations to be used later on.

First we assume that q=2 and that G is an SU_4 (resp. SU_5) group. Let G_0 be such that $G_{0W(\mathbb{F}_4)}$ is generated by the \mathbb{G}_a subgroups of $G_{W(\mathbb{F}_4)}$ normalized by $T_{W(\mathbb{F}_4)}$ and corresponding as in Case 2 (resp. Case 1) of 3.9 to the roots $\pm \alpha_1$ and $\pm \alpha_3$ (resp. $\pm \alpha_1$ and $\pm \alpha_4$) of Φ . It is the $\operatorname{Res}_{W(\mathbb{F}_4)/W(k)}$ of an SL_2 group G_{02} . The composite monomorphism $G_0(W(k)) = G_{02}(W(\mathbb{F}_4)) \hookrightarrow G(W(k)) \hookrightarrow G(W(\mathbb{F}_4))$ corresponds to the direct sum W of two copies of the standard rank 2 representation W_0 of G_{02} with the trivial rank 0 (resp. 1) representation of G_{02} over $W(\mathbb{F}_4)$. We show that the assumption $\gamma_G = 0$ leads to a contradiction. The image $\gamma^1_{G_0,G}$ of $\gamma^0_{G_0,G}$ in $H^2(G_{02}(\mathbb{F}_4), \operatorname{Lie}_{\mathbb{F}_2}(GL(W/2W)))$ is also 0. This image is defined by the natural monomorphisms $\operatorname{Lie}(G_k) \hookrightarrow \operatorname{Lie}_{\mathbb{F}_2}(GL(W/2W))$ of $G_0(k) = G_{02}(\mathbb{F}_4)$ -modules. But the component of $\gamma^1_{G_0,G}$ in $H^2(G_{02}(\mathbb{F}_4), (\operatorname{Lie}_{\mathbb{F}_2}(GL(W_0/2W_0)))^4)$ can be identified with four copies of γ_4 and so it is non-zero. Contradiction. So $\gamma_G \neq 0$.

Second we assume that q=4 and G is an SU_3 group. Let G_0 be the unique SL_2 subgroup of G normalized by T. The composite monomorphism $G_0(W(k)) \hookrightarrow G(W(\mathbb{F}_{16})) = SL_3(W(\mathbb{F}_{16}))$ corresponds to the direct sum of W_0^2 with the trivial rank 2 representation of G_0 over W(k). So as in the previous paragraph we get that $\gamma_G \neq 0$.

4.4.3. On PGL_3 over \mathbb{Z}_3 . We assume that q=3 and G is a PGL_3 group. We show that $\gamma_G \neq 0$. We work inside $M_3(W_2(k))$. Let $X_1 := I_3 + E_{12} + 3Y_1$ and $X_2 := I_3 + E_{23} + 3Y_2$, with $Y_1, Y_2 \in M_3(W_2(k))$. Let $X_3 := X_1X_2X_1^2X_2^2$. We have $X_3 = I_3 + E_{13} + 3Y_3$, with $Y_3 \in M_3(W_2(k))$. It is enough to show that we can not choose Y_1 and Y_2 such that X_3^3 is of the form βI_3 , with $\beta \in \text{Ker}(\mathbb{G}_m(W_2(k)) \to \mathbb{G}_m(k))$. As $X_3^3 = I_3 + 3E_{13} + 3E_{13}Y_3E_{13}$, it is enough to show that the 31 entry of X_3 (and so also of Y_3 mod 3) is 0.

Let $=_{31}$ be the equivalence relation on $M_3(W_2(k))$ such that two matrices are in relation $=_{31}$ iff their 31 entries are equal. For $Z \in M_3(W_2(k))$ we have

(14)
$$E_{12}Z =_{31} E_{13}Z =_{31} E_{23}Z =_{31} ZE_{12} =_{31} ZE_{23} =_{31} 0_3.$$

So we compute

$$X_{3} =_{31} (I_{3} + E_{12} + E_{23} + 3E_{12}Y_{2} + E_{12}E_{23} + 3Y_{1} + 3Y_{2} + 3Y_{1}E_{23})(I_{3} + 2E_{12} + 3E_{12}Y_{1} + 6Y_{1} + 3Y_{1}E_{12})$$

$$(I_{3} + 2E_{23} + 3E_{23}Y_{2} + 6Y_{2} + 3Y_{2}E_{23}) =_{31} (I_{3} + 3Y_{1} + 3Y_{2} + 3Y_{1}E_{23})(I_{3} + 2E_{12} + 3E_{12}Y_{1} + 6Y_{1} + 3Y_{1}E_{12})$$

$$(I_{3} + 3E_{23}Y_{2} + 6Y_{2}) =_{31} (I_{3} + 3Y_{1} + 3Y_{2} + 3Y_{1}E_{23})(I_{3} + 3E_{23}Y_{2} + 6Y_{2} + 2E_{12} + 6E_{12}E_{23}Y_{2} + 3E_{12}Y_{2} +$$

$$+3E_{12}Y_{1} + 6Y_{1} + 3Y_{1}E_{12}) =_{31} (I_{3} + 3Y_{1} + 3Y_{2} + 3Y_{1}E_{23})(I_{3} + 3E_{23}Y_{2} + 6Y_{2} + 6E_{13}Y_{2} + 3E_{12}Y_{2} +$$

$$+3E_{12}Y_{1} + 6Y_{1}) =_{31} I_{3} + 3E_{23}Y_{2} + 6Y_{2} + 6E_{13}Y_{2} + 3E_{12}Y_{2} + 3E_{12}Y_{1} + 6Y_{1} + 3Y_{1} + 3Y_{2} + 3Y_{1}E_{23}.$$
So $X_{3} =_{31} I_{3} + 9Y_{1} + 9Y_{2} =_{31} I_{3}$ and so $\gamma_{G} \neq 0$.

4.4.4. On $PGSp_4$ over \mathbb{Z}_3 . We assume that q=3 and G is a $PGSp_4$ group. To show that $\gamma_G \neq 0$ we follow the pattern of 4.4.3. We choose the alternating form ψ on $W(k)^4$ such that for the standard W(k)-basis $\{e_1, ..., e_4\}$ of $W(k)^4$ and for $i, j \in \{1, ..., 4\}$, j > i, we have $\psi(e_i, e_j) = 1$ if j - i = 2 and $\psi(e_i, e_j) = 0$ if $j - i \neq 2$. We take $X_1 := I_4 + E_{12} - E_{43} + 3Y_1$, $X_2 := I_3 - E_{14} - E_{23} + 3Y_2$, with Y_1 and Y_2 as in 4.4.3. Defining X_3 as in 4.4.3, we have $X_3 = I_3 + E_{13} + 3Y_3$. As in 4.4.3 it is enough to show that the 31 entry of X_3 is 0. But the computations of 4.4.3 apply, once we remark that similar to (14) for $Z \in M_4(W_2(k))$ and for the equivalence relation $=_{31}$ on $M_4(W_2(k))$ defined as in 4.4.3, we have $(E_{12} - E_{43})Z =_{31} (-E_{14} - E_{23})Z =_{31} Z(E_{12} - E_{43}) =_{31} Z(-E_{14} - E_{23}) =_{31} 0_4$.

So $\gamma_G \neq 0$. Warning: here X_1 , X_2 and X_3 are "related" to the roots α_1 , $\alpha_1 + \alpha_2$ and respectively $2\alpha_1 + \alpha_2$ of the C_2 Dynkin type; the similar computations for the roots α_1 , α_2 and $\alpha_1 + \alpha_2$ do not imply that $\gamma_G \neq 0$.

4.4.5. On $PGSp_4$ over \mathbb{Z}_2 . We assume that q=2 and G is a $PGSp_4$ group. Let ψ be as in 4.4.4 (but with q=2). Let $GSp_4:=GSp(W(k)^4,\psi)$. We show that $\gamma_G\neq 0$. The group $GSp_4(k)$ is the symmetric group S_6 (see [1]). It can be checked that G(W(k)) has a subgroup isomorphic to A_6 and so in this case the following computations are more involved. Let $I:=[\mathrm{Lie}(G_k^{\mathrm{sc}}),\mathrm{Lie}(G_k^{\mathrm{sc}})]$. We view it as a $GSp_4(k)$ -module and so also as a normal subgroup of $GSp_4(W_2(k))$. As $GSp_4(k)=G(k)$ and as $\mathrm{Lie}(Z(GSp_4)_k)\subset I$, it is enough to show that the short exact sequence

(15)
$$0 \to \operatorname{Lie}(GSp_{4k})/I \to GSp_4(W_2(k))/I \to GSp_4(k) \to 0$$

does not have a section. We show that the assumption that (15) has a section leads to a contradiction. Let S be a subgroup of $GSp_4(W_2(k))/I$ mapping isomorphically into $GSp_4(k)$.

If $U \in GSp_4(W_2(k))$ let \bar{U} be the reduction mod p of U and let \tilde{U} be the image of U in $GSp_4(W_2(k))/I$. Below $x_i, y_i, z_i, w_i \in W_2(k)$, where $i \in \{1, ..., 5\}$. Let $X \in GSp_4(W_2(k))$ whose rows are $(1 + 2x_5 \ 0 \ 2x_1 \ 0)$, $(0 \ 1 + 2x_5 \ 0 \ 1 + 2x_2 + 2x_5)$, $(2x_3 \ 0 \ 1 \ 0)$ and $(0 \ 2x_4 \ 0 \ 1 + 2x_5 \ 0 \ 1)$

 $2x_4$). Let $Y \in GSp_4(W_2(k))$ whose rows are $(1 + 2y_5 \ 0 \ 1 + 2y_1 + 2y_5 \ 0)$, $(0 \ 1 + 2y_5 \ 0 \ 2y_2)$, $(2y_3 \ 0 \ 1 + 2y_3 \ 0)$ and $(0 \ 2y_4 \ 0 \ 1)$. Let $Z \in GSp_4(W_2(k))$ whose rows are $(1 + 2z_5 \ 0 \ 2z_1 \ 2z_1)$, $(-1 + 2z_5 \ 1 + 2z_5 \ 0 \ 2z_2)$, $(2z_3 \ 0 \ 1 \ 1)$ and $(2z_4 \ 2z_4 \ 0 \ 1)$. Let $W \in GSp_4(W_2(k))$ whose rows are $(1 + 2w_5 \ 0 \ 2w_1 \ 1 + 2w_5)$, $(0 \ 1 + 2w_5 \ - 1 + 2w_5 \ 2w_2)$, $(2w_3 \ 0 \ 1 \ 2w_3)$ and $(0 \ 2w_4 \ 2w_4 \ 1)$. The subgroup of $GSp_4(k)$ generated by $\bar{X}, \bar{Y}, \bar{Z}$ and \bar{W} is a 2-Sylow subgroup.

The subgroup of $\text{Lie}(GSp_{4k})$ generated by E_{13} , E_{24} , E_{31} , E_{42} and $E_{11}+E_{22}$ is a direct supplement of I. So for $U \in \{X,Y,Z,W\}$, \tilde{U} is the general element of $GSp_4(W_2(k))/I$ lifting \bar{U} . We now choose \tilde{X} , \tilde{Y} , \tilde{Z} and \tilde{W} to generate a 2-Sylow subgroup of S. So \tilde{X}^2 , \tilde{Y}^2 , \tilde{Z}^2 , \tilde{W}^2 , $(\tilde{Z}\tilde{W})^2$, $(\tilde{W}\tilde{X})^2$, $(\tilde{X}\tilde{Y})^2$ and $(\tilde{Z}\tilde{X})^2$ are all identity elements.

Both \bar{Z} and \bar{W} are associated to short roots of the C_2 Lie type. So from (6) we get that they belong to the commutator subgroup of G(k). So also \tilde{Z} and \tilde{W} belong to the commutator subgroup of S and so to $G^{\rm sc}(W_2(k))/I$. So $2z_5=2w_5=0$. Looking at the 31 entry of Z^2 (and so of \tilde{Z}^2) we get $2z_4=0$. Looking at the 24 entry of W^2 we get $2w_3=0$. Looking at the 13 entry of $(ZW)^2$ we get $2w_4=0$. Looking at the 13 and 24 entries of $(WX)^2$ and using that $2w_3=2w_4=2w_5=0$, we get that $2x_4=0$ and respectively that $2x_3+2x_4+2x_5=2$. Looking at the 24 entry of X^2 we get $2x_4+2x_5=2$. So $2x_3=0$ and $2x_5=2$. Looking at the 13 entry of Y^2 we get $2y_3+2y_5=2$. Looking at the 13 entry of $(XY)^2$ we get $2x_3+2x_5+2y_3+2y_5=2$. So $2y_3+2y_5=0$. As we also have $2y_3+2y_5=2$, we reached a contradiction. So (15) does not have a section and so also $\gamma_G \neq 0$.

4.4.6. On PGL_4 over \mathbb{Z}_2 . We assume that q=2 and G is a PGL_4 group. We identify $G^{\operatorname{sc}}=SL(W(k)^4)$. We show that the assumption $\gamma_G=0$ leads to a contradiction. Let S be a subgroup of $G(W_2(k))$ mapping isomorphically into G(k). So S is a simple group, cf. [11, 2.2.7]. The group $G(W_2(k))/\mathfrak{L}_{G_k}$ is isomorphic to $S \times \mathbb{Z}/2\mathbb{Z}$, cf. 3.10 5). As there is no epimorphism from S onto $\mathbb{Z}/2\mathbb{Z}$, the images of S and $G^{\operatorname{sc}}(W_2(k))$ in $G(W_2(k))/\mathfrak{L}_{G_k}$ coincide. So the inverse image S_1 of S in $G^{\operatorname{sc}}(W_2(k))$ surjects onto S. Let $SP := \{(i,j)|i,j\in\{1,2,3,4\} \text{ and } i\neq j\}$. For $(i,j)\in SP$ let $u_{ij}\in S_1$ be such that its image in S is the reduction mod S of S in S or S and S is the reduction mod S of S in S and S is the reduction mod S of S in S in S in S is the reduction mod S of S in S in S in S in S is the reduction mod S of S in S in S in S in S is the reduction mod S of S in S in S in S in S in S in S is the reduction mod S in S

Let $X_1:=u_{23}u_{34}u_{23}u_{34}=I_4+E_{24}+2Y_1$ and $X_2:=u_{13}=I_4+E_{13}+2Y_2$, where $Y_1,\,Y_2\in M_2(W_2(k))$. We know that $X_1,\,X_2\in SL_4(W(k)^4),\,X_1^2=X_2^2=I_4$ and $X_1X_2=X_2X_1$. We take $x_y\in W_2(k)$, where $x\in\{a,b,...,p\}$ and $y\in\{1,2\}$. As $X_1^2=I_4+2E_{24}+2[Y_1,E_{24}]=I_4$, the rows of X_1 are $(1+2a_1\ 0\ 2c_1\ 2d_1),\,(2e_1\ 1+2f_1\ 2g_1\ 1+2h_1),\,(2i_1\ 0\ 1+2k_1\ 2l_1)$ and $(0\ 0\ 0\ 3+2f_1)$. Similarly, as $X_2^2=I_4$ we get that the rows of X_2 are $(1+2a_2\ 2b_2\ 1+2c_2\ 2d_2),\,(0\ 1+2f_2\ 2g_2\ 2h_2),\,(0\ 0\ 3+2a_2\ 0)$ and $(0\ 2n_2\ 2o_2\ 1+2p_2)$. Identifying the 24 entries of X_2X_1 and X_1X_2 we get $(1+2h_1)(1+2f_2)+2h_2(3+2f_1)=(1+2f_1)2h_2+(1+2h_1)(1+2p_2)$. So $2f_2=2p_2$. As $2f_2=2p_2$ we have $\det(X_2)=(1+2a_2)(3+2a_2)(1+2f_2)(1+2p_2)=3$. So $X_2\notin G^{\rm sc}(W_2(k))$. Contradiction. So $\gamma_G\neq 0$.

4.4.7. On PGL_3 over $W(\mathbb{F}_4)$. We assume that q=4 and G^{ad} is a PGL_3 group. We show that $\gamma_G \neq 0$. As $c(G^{\operatorname{sc}})$ is odd we can assume G is an SL_3 group. Let $\alpha, \beta \in k^*$, with $\alpha \neq \beta$. Let $X_1 := I_3 + t_{\alpha}E_{12} + 2Y_1$ and $X_2 := I_3 + t_{\beta}E_{12} + 2Y_2$, where $Y_1, Y_2 \in M_3(W_2(k))$. If $X_1^2 = X_2^2 = I_3$, then Y_j has the rows $(2a_j \ 2b_j \ 2c_j) \ (0 \ 2 + 2a_j \ 0)$ and $(0 \ 2h_j \ 2i_j)$. Here $j \in \{1, 2\}$ and all $a_j, ..., i_j \in W_2(k)$. So as $\alpha \neq \beta$ we get that the 12 entries of X_1X_2 and X_2X_1 are distinct. So $X_1X_2 \neq X_2X_1$ and so $\gamma_G \neq 0$.

4.4.8. On PGU_3 over \mathbb{Z}_2 . We show that $\gamma_G = 0$ if q = 2 and G^{ad} is a PGU_3 group. We can assume G is s.c. Let U be the unipotent radical of B. As $G_{W(\mathbb{F}_4)}$ splits, it is the SLgroup of $M := W(\mathbb{F}_4)^3$ and $T_{W(\mathbb{F}_4)}$ splits. So we can choose a $W(\mathbb{F}_4)$ -basis $\mathcal{B} = \{e_1, e_2, e_3\}$ of M such that the $W(\mathbb{F}_4)$ -spans $\langle e_1 \rangle$ and $\langle e_1, e_2 \rangle$ are normalized by $B_{W(\mathbb{F}_4)}$ and the $W(\mathbb{F}_4)$ -spans $\langle e_1 \rangle$, $\langle e_2 \rangle$ and $\langle e_3 \rangle$ are normalized by $T_{W(\mathbb{F}_4)}$. In what follows the matrices of elements of $G(W(\mathbb{F}_4))$ are computed with respect to \mathcal{B} . We can assume that \mathcal{B} is such that the automorphism of $U_{W(\mathbb{F}_4)}$ defined by the non-identity element τ of $\operatorname{Gal}(\mathbb{F}_4/k) = \operatorname{Gal}(B(\mathbb{F}_4)/B(k))$ takes $A \in U(W(\mathbb{F}_4))$ whose rows are $(1 \, x \, y)$, $(0 \, 1 \, z)$ and (001) into the element $\tau(A) \in U(W(\mathbb{F}_4))$ whose rows are $(1\tau(z)\tau(xz-y)), (01\tau(x))$ and $(0\ 0\ 1)$. We have $U(W(k)) := \{A \in U(W(\mathbb{F}_4)) | \tau(A) = A\}$. As the G(k)-submodule $\operatorname{Lie}(G_k)$ of $\operatorname{Lie}_{\mathbb{F}_2}(G_{\mathbb{F}_4})$ has a direct supplement and as U(k) is a Sylow 2-subgroup of G(k), it is enough to show that the pull back of the standard short exact sequence $0 \to \mathrm{Lie}_{\mathbb{F}_2}(G_{\mathbb{F}_4}) \to$ $G(W_2(\mathbb{F}_4)) \to G(\mathbb{F}_4) \to 0$ via the composite monomorphism $U(k) \hookrightarrow G(k) \hookrightarrow G(\mathbb{F}_4)$ has a section. Even better, as the G(k)-submodule $\mathrm{Lie}_{\mathbb{F}_2}(G_{\mathbb{F}_4})$ of $\mathrm{Lie}_{\mathbb{F}_2}(GL(M/2M))$ has $\operatorname{Lie}_{\mathbb{F}_2}(Z(GL(M/2M)))$ as a direct supplement, it is enough to deal with the image of the resulting short exact sequence via the monomorphism $\operatorname{Lie}_{\mathbb{F}_2}(G_{\mathbb{F}_4}) \hookrightarrow \operatorname{Lie}_{\mathbb{F}_2}(GL(M/2M))$.

Let $a \in \mathbb{F}_4$ be such that $\mathbb{F}_4 = \{0, 1, a, a+1\}$. Let \tilde{t}_a and \tilde{t}_{a+1} be elements of $W_2(\mathbb{F}_4)$ lifting a and respectively a+1 and such that we have the following two identities

(16)
$$\tilde{t}_{a+1} + \tilde{t}_a = \tilde{t}_a \tilde{t}_{a+1} = 1.$$

Let X_1, X_2 and $X_3 \in GL(M)(W_2(\mathbb{F}_4))$ be defined as follows. The rows of X_1 (resp. of X_2) are $(1 \, 1 \, \tilde{t}_a)$, $(2 \, 3 + 2 \, \tilde{t}_a \, 3)$ and $(0 \, 0 \, 1)$ (resp. are $(1 \, \tilde{t}_a \, \tilde{t}_a)$, $(2 + 2 \, \tilde{t}_a \, 1 \, \tilde{t}_{a+1})$ and $(0 \, 0 \, 3 + 2 \, \tilde{t}_a)$). The rows of X_3 are $(3 \, 2 \, \tilde{t}_a \, 3 + 2 \, \tilde{t}_a)$, $(0 \, 3 \, 0)$ and $(0 \, 0 \, 1)$. It is easy to see that $X_3^2 = I_3$ and $X_1^2 = X_2^2 = (X_1 X_2)^2 = X_3$. We include here just the only two computations which appeal to (16). The 13 entry of X_2^2 is $\tilde{t}_a + \tilde{t}_a \, \tilde{t}_{a+1} + 3 \, \tilde{t}_a + 2 (\tilde{t}_a)^2 = 1 + (2 \, \tilde{t}_a)^2 = 3 + 2 \, \tilde{t}_a$. The rows of $X_1 X_2$ are $(3 + 2 \, \tilde{t}_a \, 1 + \tilde{t}_a \, 2 (\tilde{t}_a)^2 + \tilde{t}_{a+1})$, $(2 \, \tilde{t}_a \, 3 \, 3 + 3 \, \tilde{t}_{a+1})$ and $(0 \, 0 \, 3 + 2 \, \tilde{t}_a)$. So the 13 entry of $(X_1 X_2)^2$ is $2(3 + 2 \, \tilde{t}_a)[2(\tilde{t}_a)^2 + \tilde{t}_{a+1}] + (1 + \tilde{t}_a)(3 + 3 \, \tilde{t}_{a+1}) = \tilde{t}_{a+1} + 3 + 3 \, \tilde{t}_a + 3 \, \tilde{t}_a \, \tilde{t}_{a+1} = 3 + 2 \, \tilde{t}_a$. So the subgroup of $GL(W)(W_2(\mathbb{F}_4))$ generated by X_1, X_2, X_3 is a quaternion group

So the subgroup of $GL(W)(W_2(\mathbb{F}_4))$ generated by X_1, X_2, X_3 is a quaternion group of order 8. Its reduction mod 2 is U(k). So $\gamma_G = 0$ for the present case.

- **4.4.9.** On PGU_3 over \mathbb{Z}_3 . We assume that q=3 and G^{ad} is a PGU_3 group. We show that $\gamma_G \neq 0$. Let M, \mathcal{B} , U, τ be as in 4.4.8 but with q=3 and with \mathbb{F}_4 replaced by \mathbb{F}_9 . Let $a \in \mathbb{F}_9$ be such that $a^2+1=0$. Let $X_1, X_2, X_3, Y_1, Y_2, Y_3 \in M_3(W(\mathbb{F}_9))$ be such that $X_1=I_3+t_aE_{12}+t_{2a}E_{23}+2E_{13}+3Y_1, X_2=I_3+t_{a+1}E_{12}+t_{2a+1}E_{23}+E_{13}+3Y_2$ and $X_3=X_1X_2X_1^2X_2^2=I_3+t_{2a}E_{13}+3Y_3$. The reductions mod 3 of X_1, X_2 and X_3 define elements of U(k). As in 4.4.3 and 4.4.4 we get that the 31 entry of Y_3 mod 3 is 0. So X_3^3 is not a scalar multiple of I_3 . So $\gamma_G \neq 0$.
- **4.5. Theorem.** We assume G is adjoint. Then $\gamma_G = 0$ (resp. $\gamma_{G^{\text{sc}}} = 0$) iff $q \le 4$ (resp. $q \le 3$) and G (resp. G^{sc}) is a product of adjoint (resp. s.c.) groups of the following type:

F2 PGL_2 , PGL_3 , $Res_{W(\mathbb{F}_4)/\mathbb{Z}_2}PGL_2$, PGU_3 , PGU_4 and split of G_2 Dynkin type (resp. SL_3 , SU_3 and split of G_2 Dynkin type) if q=2;

F3 PGL_2 (resp. SL_2) if q=3;

F4 PGL_2 if q = 4.

Proof: If $\gamma_{G^{\text{sc}}} = 0$, then $\gamma_G = 0$. It suffices to prove the Theorem for a simple, adjoint group G, cf. 4.2 3) and 4.3.4. We first assume $\gamma_G = 0$. Let k_1 be such that $G = \text{Res}_{W(k_1)/W(k)}G_1$, where G_1 is an a.s. adjoint group (cf. 4.2 3)). Let T_1 (resp. B_1) be the maximal torus (resp. Borel subgroup) of G_1 such that $\text{Res}_{W(k_1)/W(k)}T_1 = T$ (resp. $\text{Res}_{W(k_1)/W(k)}B_1 = B$), cf. 2.3.1. Let G_{01k_1} be a closed, semisimple subgroup of G_{1k_1} normalized by T_{1k_1} and obtained as in 3.9 (i.e. working with respect to B_{1k_1}). It lifts to a semisimple subgroup G_{01} of G_1 normalized by T_1 , cf. 4.3.2 and 3.9 (2). So $G_0 := \text{Res}_{W(k_1)/W(k)}G_{01}$ is a closed, semisimple subgroup of G (cf. 2.3.1) normalized by T.

As $\gamma_G = 0$ we have $p \le 3$, cf. 4.4.1. If p = 3, then q = 3 and G is split (cf. 4.4.1 and 4.4.9). We show that G is a PGL_2 group. If it is not, then we can take G_0 to be split of A_2 or C_2 Dynkin type (even if G is of G_2 Dynkin type). So $\gamma_{G_0^{\text{ad}}} \ne 0$ (cf. 4.4.3 and 4.4.4) and so $\gamma_G \ne 0$ (cf. 4.3.4). We reached a contradiction. So G is a PGL_2 groups. But $\gamma_G = 0$ and $\gamma_{G^{\text{sc}}} = 0$ for such a G, cf. [24, p. IV-27-28].

Until end of the proof we assume p=2. As $\gamma_G=0$, k is a subfield of \mathbb{F}_4 and either G splits over $W(\mathbb{F}_4)$ or is the $\mathrm{Res}_{W(\mathbb{F}_4)/W(k)}$ of a PGU_3 group (cf. 4.4.1).

- i) The group G_0^{ad} can not be a PGL_4 group over \mathbb{Z}_2 , cf. 4.4.6 and 4.3.4. So also G_0^{ad} can not be the $\mathrm{Res}_{W(\mathbb{F}_4)/W(k)}$ of a PGL_4 group, cf. 4.3.4 and 4.3.5. So G is not the Weil restriction of a split group of E_6 Dynkin type, cf. 3.9 and 4.3.2. Also G is not a PGU_{n+7} group, a PGL_{n+3} group, or of isotypic D_{n+3} , E_7 or E_8 Dynkin type (cf. 3.9 and 4.3.2).
- ii) The group G_0^{ad} can not be a $PGSp_4$ group over \mathbb{Z}_2 , cf. 4.4.5 and 4.3.4. So also G_0^{ad} can not be the $\text{Res}_{W(\mathbb{F}_4)/W(k)}$ of a $PGSp_4$ group, cf. 4.3.4 and 4.3.5. So G is not of isotypic B_n with $n \geq 2$, C_n with $n \geq 2$ or F_4 Dynkin type (cf. 3.9 and 4.3.2).
- iii) The group G_0^{ad} can not be the $\operatorname{Res}_{W(\mathbb{F}_4)/\mathbb{Z}_2}$ of a PGL_3 group, cf. 4.4.7 and 4.3.4. So G is not a PGU_6 or a PGU_7 group, cf. 3.9 and 4.3.2.
- iv) The group G_0 can not be a PGU_6 group over \mathbb{Z}_2 , cf. iii) and 4.3.4. So G is not non-split of E_6 Dynkin type, cf. Case 3 of 3.9 and 4.3.2.
- v) The groups G_0 and G are not the $\operatorname{Res}_{W(\mathbb{F}_4)/W(k)}$ of a PGL_3 group, cf. 4.4.7 and 4.3.4. So G is also not $\operatorname{Res}_{W(\mathbb{F}_4)/W(k)}$ of a split group of G_2 Dynkin type, cf. 3.9 and 4.3.2.
- vi) The group G is not a PGU_5 group or the $\operatorname{Res}_{W(\mathbb{F}_4)/W(k)}$ of a PGU_3 group, cf. 4.4.2 and the fact that c(G) is odd.

So G is among the adjoint groups listed in F2 and F4. If q=2 and G is split of G_2 Dynkin type, then the epimorphism G(W(k)) G(k) has a right inverse (cf. [12, Sect. 4]). It is well known that this also holds if q=2 (resp. q=4) and G is a PGL_n group with $n \in \{2,3\}$ (resp. n=2). Let SO_6^- be the semisimple group over \mathbb{Z}_2 which is an isogeny cover of degree 2 of a PGU_4 group. The epimorphism $SO_6^-(\mathbb{Z}_2) SO_6^-(\mathbb{F}_2)$ has a right inverse (see [1, p. 26]; $SO_6^-(\mathbb{F}_2)$ is the Weyl group W_{E_6}). Based on all these and 4.4.8, we get that F2 and F4 list all cases when p=2, G is a simple adjoint group and $\gamma_G=0$.

The passage from adjoint groups to s.c. groups for p=2 is easy. It is well known that $\gamma_{G^{\text{sc}}} \neq 0$ if $q \in \{2,4\}$ and G is a PGL_2 group (see 4.4.2 for q=4). So $\gamma_{G^{\text{sc}}} = 0$ iff q=2 and G^{sc} is a product of s.c. semisimple groups having a.s. adjoints of A_2 or G_2 Dynkin type (cf. 4.4.2 for the exclusion of SU_4 groups for q=2). This ends the proof.

4.5.1. Remark. We have $\gamma_G = 0$ if q = 2 and G is the quotient of $\operatorname{Res}_{W(\mathbb{F}_4)/\mathbb{Z}_2} SL_2$ by μ_2 (this can be deduced either by just adapting (11) to (13) or from [1, p. 26] via the

standard embedding of G into SO_6^-).

- **4.6.** A supplement to **4.5.** Until 4.7 we assume G is split, s.c. and G^{ad} is a.s. We also assume that either p=3 and G is of G_2 Dynkin type or p=2 and G is of B_n , C_n or F_4 Dynkin type. If G is an SL_2 group, then we also assume that $q \geq 8$. Let I be the maximal G(k)-submodule of $\operatorname{Lie}_{\mathbb{F}_p}(G_k)$, cf. 3.11.2 1) and 3). We refer to (10). Let G_0 be the closed, semisimple subgroup of G generated by the \mathbb{G}_a subgroups of G having \mathfrak{g}_{α} 's as their Lie algebras, with $\alpha \in \Phi$ a long root (see 3.9 and 4.3.2).
- **4.6.1.** Lemma. We have identifications of Lie algebras

(17)
$$\operatorname{Lie}(G_k)/I = \operatorname{Lie}(G_{0k})/\operatorname{Lie}(Z(G_{0k})) = \mathfrak{L}_{G_{0k}^{\mathrm{ad}}}.$$

Proof: The intersection $I \cap \text{Lie}(G_{0k})$ is fixed by any maximal torus of G_{0k} and so also by G_{0k} . So $I \cap \text{Lie}(G_{0k}) \subset \text{Lie}(Z(G_{0k}))$, cf. 3.10 1) applied to G_{0k}^{ad} . A simple computation involving $\dim_k(I)$ (see [13, p. 409]) and $\dim_k(G_{0k})$ (see 3.9 for the Lie type of G_{0k}^{ad}) shows that $\dim_k(I \cap \text{Lie}(G_{0k})) \geq \dim_k(Z(G_{0k}^{\text{sc}}))$. Example: if G is of C_n Dynkin type, then $\dim_k(I \cap \text{Lie}(G_{0k})) \geq \dim_k(Z(G_{0k}^{\text{sc}})) \geq \dim_k(Z(G_{0k}^{\text{sc}})) \geq \dim_k(Z(G_{0k}))$, we get $I \cap \text{Lie}(G_{0k}) = \text{Lie}(Z(G_{0k}))$ and $\dim_k(Z(G_{0k})) = \dim_k(Z(G_{0k}^{\text{sc}}))$. So $\text{Lie}(G_{0k})/\text{Lie}(Z(G_{0k})) = \text{Lie}(G_{0k}^{\text{sc}})/\text{Lie}(Z(G_{0k}^{\text{sc}})) = \mathfrak{L}_{G_{0k}^{\text{ad}}}$ (cf. also proof of 3.10 4) for when p = 2 and G_{0k} is of A_3 or D_{2n+3} Lie type). By reasons of dimensions the monomorphism $\text{Lie}(G_{0k})/\text{Lie}(Z(G_{0k})) = \text{Lie}(G_{0k})/I \cap \text{Lie}(G_{0k}) \hookrightarrow \text{Lie}(G_k)/I$ is onto. So (17) holds. This ends the proof.

4.6.2. Proposition. We view I as a normal subgroup of $G(W_2(k))$. Then the short exact sequence

(18)
$$0 \to Lie_{\mathbb{F}_p}(G_k)/I \to G(W_2(k))/I \to G(k) \to 0$$

does not have a section.

Proof: Its pull back via the monomorphism $G_0(k) \hookrightarrow G(k)$ is the short exact sequence

$$(19) \quad 0 \to \mathrm{Lie}_{\mathbb{F}_p}(G_{0k})/\mathrm{Lie}_{\mathbb{F}_p}(Z(G_{0k})) \to \mathrm{Im}(G_0(W_2(k))) \to G_0^{\mathrm{ad}}(W_2(k))) \to G_0(k) \to 0,$$

- cf. (17). We show that the assumption that (18) has a section leads to a contradiction. So (19) has a section too. So $\gamma_{G_0^{\rm ad}} = 0$. So $q \le 4$ and G is of isotypic C_n Dynkin type with $n \ge 2$ (cf. 4.5; see 3.9 for the structure of $G_{0k}^{\rm ad}$). To reach a contradiction we can assume q = 2 (to be compared with 4.3.5 and the proof of 3.11.1). Using a standard monomorphism $\mathbb{G}_m^{n-2} \times Sp_4 \hookrightarrow Sp_{2n}$ over W(k), similar to 4.3.3 and 4.3.4 we argue that we can assume n = 2. But as (15) does not have a section (see 4.4.5), (18) does not have a section for q = 2 and G an Sp_4 group. Contradiction. This ends the proof.
- **4.7.** Proof of 1.3. We now prove 1.3. Let G and K be as in 1.3. Let L_2 and L_3 be as before 4.1.2. As K surjects onto G(k), both L_2 and L_3 are G(k)-submodules of $\operatorname{Lie}_{\mathbb{F}_p}(G_k)$. Let K^{sc} be the inverse image of K in $G^{\operatorname{sc}}(W(k))$. We have $\operatorname{Im}(K^{\operatorname{sc}} \to G(k)) = \operatorname{Im}(G^{\operatorname{sc}}(k) \to G(k))$. So $\operatorname{Im}(K^{\operatorname{sc}} \to G^{\operatorname{sc}}(k))$ is a normal subgroup of $G^{\operatorname{sc}}(k)$ of index dividing the order of $Z(G^{\operatorname{sc}})(k)$ and so prime to p. So as $G^{\operatorname{sc}}(k)$ is generated by elements of order p (cf. [11,

2.2.6 (f)]), K^{sc} surjects onto $G^{\text{sc}}(k)$. As g.c.d.(p, c(G)) = 1, the isogeny $G^{\text{sc}} \to G$ is étale. So $\text{Ker}(G^{\text{sc}}(W(k)) \to G^{\text{sc}}(k)) = \text{Ker}(G(W(k)) \to G(k))$. So if $K^{\text{sc}} = G^{\text{sc}}(W(k))$, then K = G(W(k)). So to show that K = G(W(k)) we can assume $G = G^{\text{sc}}$. Let G^{0} be a direct factor of G such that G^{0ad} is simple.

We first assume that 1.3 e) holds; so p=2 and $L_2=\operatorname{Lie}_{\mathbb{F}_p}(G_k)$. To prove that K=G(W(k)) it suffices to show that L_3 is $\operatorname{Lie}_{\mathbb{F}_p}(G_k)=\operatorname{Ker}(G(W_3(k))\to G(W_2(k)))$, cf. 4.1.2. We consider a faithful representation $\rho:G\hookrightarrow GL(L)$ as in the proof of 4.1.2. Let $x,\ y\in\operatorname{Lie}(G)$. Let h_x and $h_y\in K$ be such that we have $h_x=1_L+2x+4\tilde{x}$ and $h_y=1_L+2y+4\tilde{y}$, where $\tilde{x},\ \tilde{y}\in\operatorname{End}(L)$. We compute

$$(20) h_x h_y h_x^{-1} h_y^{-1} = (1_L + 2x + 4\tilde{x} + 2y + 4xy + 4\tilde{y})(1_L + 2x + 4\tilde{x})^{-1}(1_L + 2y + 4\tilde{y})^{-1} =$$

$$= (1_L + 2y + 4xy - 4yx + 4\tilde{y})(1_L + 2y + 4\tilde{y})^{-1} = 1_L + 4(xy - yx).$$

So L_3 contains $[\operatorname{Lie}_{\mathbb{F}_2}(G_k), \operatorname{Lie}_{\mathbb{F}_2}(G_k)]$. So L_3 contains also $\operatorname{Lie}_{\mathbb{F}_2}(G_k^0)$ except when G^0 is of isotypic C_n Dynkin type, cf. 3.7 2). Let now G^0 be of isotypic C_n Dynkin type. Let k_1 be such that $G^0 = \operatorname{Res}_{W(k_1)/W(k)}G_1^0$, with G_1^0 a s.c. semisimple group over $W(k_1)$ whose adjoint is a.s. (cf. 4.2 3)). The group G_1^0 is split.

Let G_{10}^0 be a closed SL_2 subgroup of G_1^0 normalized by a maximal split torus T_1^0 of G_1^0 . We identify $G_{10}^0(W_3(k_1))$ with a subgroup of $GL_2(W_3(k_1))$. We denote also by t_{α} the reduction mod $8=p^3$ of the Teichmüler lift of $\alpha \in k_1 \setminus \{0\}$. Let j_2 (resp. j_4) be the image in $GL(L)(W_3(k))$ through ρ of the element $I_2 + 2t_{\alpha}E_{12}$ (resp. $I_2 + 4t_{\alpha}E_{12}$) of $G_{10}^0(W_3(k_1)) \triangleleft G(W(k))$. For $y \in \operatorname{End}(L/8L)$ we have $(j_2 + 4y)^2 = j_4 \neq I_2$. So L_3 contains the Lie algebra of any \mathbb{G}_a subgroup of $G_{10k_1}^0$ normalized by $T_{1k_1}^0$. So as the intersection $L_3 \cap \operatorname{Lie}_{\mathbb{F}_2}(G_{10k_1}^0)$ is $G_{10}^0(k_1)$ -invariant, it is $\operatorname{Lie}_{\mathbb{F}_2}(G_{10k_1}^0)$. So L_3 contains $\operatorname{Lie}_{\mathbb{F}_2}(G_{1k_1}^0) = \operatorname{Lie}_{\mathbb{F}_2}(G_k^0)$, cf. 3.7 1) and 2). So $L_3 = \operatorname{Lie}_{\mathbb{F}_p}(G_k)$ and so K = G(W(k)).

We now assume that one of the conditions a) to d) of 1.3 holds. To prove that K = G(W(k)) it is enough to show that $L_2 = \operatorname{Lie}_{\mathbb{F}_p}(G_k)$, cf. 4.1.2 and the above part referring to 1.3 e). We first assume G^{ad} is simple. Either $\gamma_{G^{\operatorname{ad}}} \neq 0$ or $L_2 \not\subset \operatorname{Lie}_{\mathbb{F}_p}(Z(G_k))$, cf. 4.5 and our hypotheses. So the group $\tilde{L}_2 := \operatorname{Im}(L_2 \to \operatorname{Lie}_{\mathbb{F}_p}(G_k^{\operatorname{ad}}))$ is non-trivial. If G is (resp. is not) the Weil restriction of a s.c. group having an a.s. simple adjoint and among the s.c. groups of 4.6, then from 4.6.2 (resp. 3.7.1) we get that \tilde{L}_2 is not included in the unique maximal G(k)-submodule of $\operatorname{Im}(\operatorname{Lie}_{\mathbb{F}_p}(G_k) \to \operatorname{Lie}_{\mathbb{F}_p}(G_k^{\operatorname{ad}}))$ (see 3.12). So $\tilde{L}_2 = \operatorname{Im}(\operatorname{Lie}_{\mathbb{F}_p}(G_k) \to \operatorname{Lie}_{\mathbb{F}_p}(G_k^{\operatorname{ad}}))$. So $L_2 = \operatorname{Lie}_{\mathbb{F}_p}(G_k)$, cf. 3.10 2). So K = G(W(k)).

We now treat the general case; so G^{ad} is not simple. We know that K projects onto $G^0(W(k))$ and that L_2 projects onto $\mathrm{Lie}_{\mathbb{F}_p}(G_k^0)$. If $x \in L_2$ and $g \in G^0(k)$, then $AD_{G_k}(g)(x) - x \in L_2 \cap \mathrm{Lie}_{\mathbb{F}_p}(G_k^0)$. From this and 3.12 we get that there is a G(k)-submodule of $L_2 \cap \mathrm{Lie}_{\mathbb{F}_p}(G_k^0)$ surjecting onto $\mathrm{Im}(\mathrm{Lie}_{\mathbb{F}_p}(G_k^0) \to \mathrm{Lie}_{\mathbb{F}_p}(G_k^{\mathrm{oad}}))$. So from 3.10 2) applied to $\mathrm{Res}_{k/\mathbb{F}_p}G_k^0$ we get that the direct factor $\mathrm{Lie}_{\mathbb{F}_p}(G_k^0)$ of $\mathrm{Lie}_{\mathbb{F}_p}(G_k)$ is contained in L_2 . As G^0 is an arbitrary direct factor of G having a simple adjoint, we get that L_2 contains $\mathrm{Lie}_{\mathbb{F}_p}(G_k)$ and so it is $\mathrm{Lie}_{\mathbb{F}_p}(G_k)$. This ends the proof of 1.3.

4.7.1. Remark. We assume that p=2, $G=G^{\mathrm{ad}}$ and 2|c(G). Then there are proper, closed subgroups of G(W(k)) surjecting onto $G(W_2(k))$. One checks this using a short exact sequence $0 \to \mathrm{Lie}_{\mathbb{F}_2}(G_k)/[\mathrm{Lie}_{\mathbb{F}_2}(G_k), \mathrm{Lie}_{\mathbb{F}_2}(G_k)] \to G(W_3(k))/I \to G(W_2(k)) \to 0$

and the fact that for any torus \tilde{T} over W(k) the square homomorphism $\operatorname{Ker}(\tilde{T}(W(k)) \to \tilde{T}(k)) \to \operatorname{Ker}(\tilde{T}(W_3(k)) \to \tilde{T}(W_2(k)))$ is not surjective.

Acknowledgements. We would like to thank Serre for two e-mail replies; the first one was the starting point of 4.4 and 4.5 which prove all its expectations and the second one led to a better presentation and to precise references in the proof of 4.5. We would also like to thank D. Ulmer and the referee for many useful comments and U of Arizona for good conditions for the writing of the paper.

References

- [1] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of finite groups*, Oxford Univ. Press, Eynsham, 1985
- [2] S. Bosch, W. Lütkebohmert, M. Raynaud, Néron models, Springer-Verlag, 1990
- [3] A. Borel, Properties and linear representations of Chevalley groups, LNM 131, Springer-Verlag, 1970, pp. 1–55
- [4] A. Borel, Linear algebraic groups, Grad. Text Math. 126, Springer-Verlag, 1991
- [5] N. Bourbaki, Groupes et algèbres de Lie, Chapitre 4–6, Hermann, 1968
- [6] N. Bourbaki, Groupes et algèbres de Lie, Chapitre 7–8, Hermann, 1975
- [7] F. Bruhat and J. Tits, Groupes réductifs sur un corps local, Publ. Math. I.H.E.S. **60**, 5–184 (1984)
- [8] A. Borel and J. Tits, Homomorphismes "abstraits" de groupes algèbriques simples, Ann. of Math. (3) **97**, 499–571 (1973)
- [9] C. W. Curtis, Representations of Lie algebras of classical Lie type with applications to linear groups, J. Math. Mech. 9, 307–326 (1960)
- [10] C. W. Curtis, On projective representations of certain finite groups, Proc. Am. Math. Soc. 11, 852–860, A. M. S. 1960
- [11] D. Gorenstein, R. Lyons and R. Soloman, The classification of the finite simple groups, Number 3, Math. Surv. and Monog., Vol. 40, A. M. S. 1994
- [12] B. Gross, Groups over Z, Inv. Math. **124**, 263–279 (1996)
- [13] G. Hiss, Die adjungierten Darstellungen der Chevalley-Gruppen, Arch. Math. **42**, 408–416 (1982)
- [14] G. M. D. Hogeweij, Almost-classical Lie algebras, Indag. Math. 44, I. 441–452, II. 453–460 (1982)
- [15] J. E. Humphreys, Introduction to Lie algebras and representation theory, Grad. Texts Math. 9, Springer-Verlag, 1975
- [16] J. E. Humphreys, *Linear algebraic groups*, Grad. Texts Math. **21**, Springer–Verlag, 1975
- [17] J. E. Humphreys, Algebraic groups and modular Lie algebras, Mem. Am. Math. Soc., No. **71**, A. M. S., 1976
- [18] J. E. Humphreys, Conjugacy classes in semisimple algebraic groups, Math. Surv. and Monog., Vol. 43, A. M. S., 1995
- [19] J. C. Jantzen, Representations of algebraic groups, Academic Press 1987
- [20] R. Pink, Compact subgroups of linear algebraic groups, J. of Algebra **206**, 438–504 (1998)

- [21] K. Ribet, On l-adic representations attached to modular forms, Inv. Math., Vol. 28, 245–275 (1975)
- [22] K. Ribet, Images of semistable Galois representations, Pac. J. of Math., Vol. 3 (3), 277–297 (1997)
- [23] J. -P. Serre, Propriétés galoisiennes des points d'ordre fini des courbes elliptiques, Inv. Math. 15, 259–331 (1972)
- [24] J. -P. Serre, Abelian l-adic representations and elliptic curves, Addison-Wesley Publ. Co. 1989
- [25] J.-P. Serre, Galois Cohomology, Springer-Verlag 1997
- [26] J.-P. Serre, Collected papers, Vol. IV, Springer-Verlag 2000
- [27] M. Demazure, A. Grothendieck, et al. Schemes en groupes, Vol. I to III, LNM 151–153, Springer–Verlag 1970
- [28] R. Steinberg, Representations of algebraic groups, Nagoya Math. J. 22, 33–56 (1963)
- [29] J. Tits, Classification of algebraic semisimple groups, Proc. Sympos. Pure Math., Vol. 9, 33–62, A. M. S. 1966
- [30] A. Vasiu, Integral canonical models of Shimura varieties of preabelian type, Asian J. Math., Vol. 3 (2), 401–518 (1999)