# ADEM-CARTAN OPERADS

DAVID CHATAUR\* UNIVERSITÉ D'ANGERS MURIEL LIVERNET UNIVERSITÉ PARIS 13

ABSTRACT. In this paper, we introduce Adem-Cartan operads and prove that the cohomology of any algebra over such an operad is an unstable level algebra over the extended Steenrod algebra. Moreover we prove that this cohomology is endowed with secondary cohomological operations.

**MSC (2000):** 55-xx; 55S10; 18D50. **Keywords:** cohomology operations, Adem-Cartan relations, operads,  $E_{\infty}$ -structures, level algebras.

## INTRODUCTION

The Steenrod Algebra  $\mathcal{A}_p$  is one of the main computational tool of homotopy theory. Steenrod's operations were first introduced by N.E. Steenrod in 1947 [22] for p = 2 and for an odd prime in 1952 [23]. The relations between these cohomological operations were determined by J. Adem [2] and H. Cartan [6]. Cartan's proof relies on the computation of the singular cohomology of the Eilenberg-Mac Lane spaces at the prime p. Adem's proof is based on the computation of the homology of the symmetric group  $\Sigma_{p^2}$  acting on  $p^2$  elements at the prime p. In the sixties, Adams introduced secondary cohomological operations [1], which are an efficient tool to deal with realisability problems of unstable modules over  $\mathcal{A}_2$ . In this paper we extend these results to a more algebraically framework, at the prime 2.

The purpose of this paper is to describe a family of operads, named Adem-Cartan operads, such that the cohomology of an algebra over such an operad is an unstable level algebra over the extended Steenrod algebra  $\mathcal{B}_2$  (see corollary 3.3.5). By a *level algebra*, we mean a commutative algebra (not necessarily associative) satisfying the following

 $<sup>^*</sup>$  Partially supported by Marie Curie grant #HPMF-CT-2001-01179 and european TMR "Modern Homotopy Theory" #HPRN-CT-1999-00119 and Bourse postdoctorale Région pays de loire

Date: October 31, 2018.

4-terms relation:

$$(a * b) * (c * d) = (a * c) * (b * d).$$

There is an operad, denoted by  $\mathcal{L}ev$ , such that level algebras are algebras over the operad  $\mathcal{L}ev$ . An unstable algebra over  $\mathcal{B}_2$  is a  $\mathcal{B}_2$ module satisfying the usual conditions, namely the Cartan formula and Adem relations. In order to describe the family of Adem-Cartan operads, the cofibrant operad  $\mathcal{L}ev^{AC}$  is introduced, obtained by the process of attaching cells from the standard bar resolution of  $\mathbb{F}_2$ , such that  $\mathcal{L}ev^{AC} \to \mathcal{L}ev$  is a fibration. More precisely, the operad  $\mathcal{L}ev^{AC}$ is generated by some 2-cells,  $e_i \in \mathcal{L}ev^{AC}(2)^{-i}$ , corresponding to the usual  $\cup_i$ -products and some 4-cells  $G_n^m \in \mathcal{L}ev^{AC}(4)^{-n}$ . An Adem-Cartan operad is an operad  $\mathcal{O}$  together with a morphism of operads  $\mathcal{L}ev^{AC} \to \mathcal{O}$  satisfying some conditions (see definition 2.3.1). We prove that the cohomology of a  $\mathcal{O}$ -algebra is an unstable algebra over  $\mathcal{B}_2$ . Namely, the cells  $e_i$  are responsible for the existence of Steenrod squares whereas the cells  $G_n^m$  are responsible for the relations between them (Adem and Cartan relations). Moreover the cohomology of a  $\mathcal{L}ev^{AC}$ algebra A is a  $\mathcal{L}ev^{AC}$ -algebra itself (this structure is however not natural). Hence the 4-cells  $G_{n+1}^m$  yield secondary cohomological operations  $\theta^{m,n}: H^q(A) \to H^{4q-n-1}(A)$ . We prove that given an Adem relation  $\sum_{i} \operatorname{Sq}^{m_{i}} \operatorname{Sq}^{n_{i}}(a) = 0$ , there exist maps  $\psi^{m,n}$  from  $\cap_{i} \operatorname{Sq}^{n_{i}} \subset H^{*}(A)$  to  $\overline{H^*}(A) / \sum_i \operatorname{Im} \operatorname{Sq}^{m_i}$  defined at the cochain level. We prove that these two maps coincide, that is  $\psi^{m,n}(a) = [\theta^{m,n}(a)]$ , for  $a \in \bigcap_i \operatorname{Sq}^{n_i}$  (see theorem 4.1.2)

Note that we recover some classical results on topological spaces. Since  $E_{\infty}$ -operads are Adem-Cartan operads (see 2.3.2), any algebra over an  $E_{\infty}$ -operad is an unstable algebra over the extended Steenrod algebra (see [18], [15]). Furthermore since the cochain complex of a topological space  $C^*(X; \mathbb{F}_2)$  is an algebra over a  $E_{\infty}$ -operad (see [12]), then it has secondary cohomological operations. Thanks to the work of Kristensen [14], we prove that there operations coincide with Adams operations. Moreover, we can extend these operations in a non natural way to secondary cohomogical operations on the whole cohomology (see theorems 4.2.2 and 4.2.3).

The paper is presented as follows. Section 1 contains the background needed. In Section 2, we set the construction of the operad  $\mathcal{L}ev^{AC}$ , define Adem-Cartan operads and prove that  $E_{\infty}$ -operads lie in this family. Section 3 is devoted to the fondamental theorem 3.3.4 and its corollary 3.3.5. Section 4 is concerned with secondary cohomological

 $\mathbf{2}$ 

operations and section 5 is devoted to proofs of technical lemmas stated in the different sections.

## 1. Recollections

The ground field is  $\mathbb{F}_2$ . In this article, a *vector space* means a differential  $\mathbb{Z}$ -graded vector space over  $\mathbb{F}_2$ , where the differential is of degree 1.

The symbol  $\Sigma_n$  denotes the symmetric group acting on n elements. Any  $\sigma \in \Sigma_n$  is written  $(\sigma(1) \dots \sigma(n))$ .

1.1. **Operads.** ([9], [10], [15], [16]) A (right)  $\Sigma_n$ -module is a (right)  $\mathbb{F}_2[\Sigma_n]$ -differential graded module. A  $\Sigma$ -module  $\mathcal{M} = {\mathcal{M}(n)}_{n>0}$  is a collection of  $\Sigma_n$ -modules. Any  $\Sigma_n$ -module M gives rise to a  $\Sigma$ -module  $\mathcal{M}$  by setting  $\mathcal{M}(q) = 0$  if  $q \neq n$  and  $\mathcal{M}(n) = M$ .

An operad is a right  $\Sigma$ -module  $\{\mathcal{O}(n)\}_{n>0}$  such that  $\mathcal{O}(1) = \mathbb{F}_2$ , together with composition products:

$$\mathcal{O}(n) \otimes \mathcal{O}(i_1) \otimes \ldots \otimes \mathcal{O}(i_n) \longrightarrow \mathcal{O}(i_1 + \ldots + i_n)$$
$$o \otimes o_1 \otimes \ldots \otimes o_n \quad \mapsto \quad o(o_1, \ldots, o_n).$$

These compositions are subject to associativity conditions, unitary conditions and equivariance conditions with respect to the action of the symmetric group. The equivariance conditions write:

$$o(a_1 \cdot \tau_1, \dots, a_n \cdot \tau_n) = o(a_1, \dots, a_n) \cdot (\tau_1 \times \dots \times \tau_n)$$
  
$$(o \cdot \sigma)(a_1, \dots, a_n) = o(a_{\sigma^{-1}(1)}, \dots, a_{\sigma^{-1}(n)}) \cdot \sigma(i_1, \dots, i_n),$$

where  $\tau_1 \times \ldots \times \tau_n$  is the permutation of  $\Sigma_{i_1+\ldots+i_n}$  such that  $\tau_1$  operates on the first  $i_1$  terms,  $\tau_2$  on the next  $i_2$  terms and so on; the permutation  $\sigma(i_1,\ldots,i_n)$  in  $\Sigma_{i_1+\ldots+i_n}$  operates as  $\sigma$  on the *n*-blocks of size  $i_k$ .

For instance, for any  $\sigma, \mu, \nu \in \mathcal{O}(2)$ , one has

$$(\sigma \cdot (21))(\mu, \nu) = \sigma(\nu, \mu) \cdot (3412)$$
(1.1)

There is another definition of operads via  $\circ_i$  operations

$$\circ_i: \mathcal{O}(n) \otimes \mathcal{O}(m) \longrightarrow \mathcal{O}(n+m-1),$$

where  $p \circ_i q$  is p composed with n-1 copies of the unit  $1 \in \mathcal{O}(1)$  and with q at the *i*-th position.

The forgetful functor from the category of operads to the category of  $\Sigma$ -modules has a left adjoint : the *free operad functor*, denoted by  $\mathcal{F}$ ree.

An algebra over an operad  $\mathcal{O}$  or a  $\mathcal{O}$ -algebra A is a vector space together with evaluation maps

$$\mathcal{O}(n) \otimes A^{\otimes n} \longrightarrow A$$
$$o \otimes a_1 \otimes \ldots \otimes a_n \mapsto o(a_1, \ldots, a_n)$$

These evaluation maps are subject to associativity conditions and equivariance conditions. These equivariance conditions write:

$$(o \cdot \sigma)(a_1, \ldots, a_n) = o(a_{\sigma^{-1}(1)}, \ldots, a_{\sigma^{-1}(n)}).$$

A graded algebra over an operad  $\mathcal{O}$  is a  $\mathcal{O}$ -algebra whose differential is null.

1.2. **Operadic cellular attachment.** ([4], [11]) The category of operads is a closed model category. Weak equivalences are quasi-isomorphisms (i.e. isomorphisms in cohomology of the underlying vector spaces) and fibrations are epimorphisms. Cofibrations can be defined by the left lifting property with respect to the acyclic fibrations. For background material on closed model categories we refer to [8], [13] and [21]. Any morphism of operads

$$\mathcal{P} \longrightarrow \mathcal{Q}$$

can be factorized by a cofibration (>--->) followed by an acyclic fibration ( $\xrightarrow{\sim}$ ). This factorization can be realized using the inductive process of attaching cells (the category of operads is cofibrantly generated [11]). An operad is *cofibrant* if the morphism from the initial object  $\mathcal{F}$ ree(0) to the operad is a cofibration. In order to produce a *cofibrant replacement* to an operad  $\mathcal{O}$ , one applies the inductive process of attaching cells to the canonical morphism  $\mathcal{F}$ ree(0)  $\longrightarrow \mathcal{O}$ .

Let  $S_p^n$  be the free  $\Sigma_p$ -module generated by  $\delta t$  in degree n considered as a  $\Sigma$ -module. Let  $D_p^{n-1}$  be the free  $\Sigma_p$ -module generated by t in degree n-1 and dt in degree n, the differential sending t to dt. We have a canonical inclusion  $i_n: S_p^n \longrightarrow D_p^{n-1}$  of  $\Sigma$ -modules (sending  $\delta t$ to dt). Let  $f: S_p^n \longrightarrow \mathcal{O}$  be a morphism of  $\Sigma$ -modules. The cell  $D_p^{n-1}$ is attached to  $\mathcal{O}$  along the morphism f via the following push-out:

$$\begin{array}{c|c} \mathcal{F}\mathrm{ree}(S_p^n) & \xrightarrow{i_n} \mathcal{F}\mathrm{ree}(D_p^{n-1}) \\ \mathcal{F}\mathrm{ree}(f) & \downarrow \\ \mathcal{O} & \xrightarrow{i} \mathcal{O} \coprod_{\tau} \mathcal{F}\mathrm{ree}(S_p^{n-1}) \end{array}$$

The main point of this process is that  $f(\delta t)$ , which was a cycle in  $\mathcal{O}$ , becomes a boundary in  $\mathcal{O} \coprod_{\tau} \mathcal{F} \operatorname{ree}(S_p^{n-1})$ .

By iterating this process of cellular attachment, one gets a quasi-free extension:  $\mathcal{O} \xrightarrow{i} \mathcal{O} \coprod_{\tau} \mathcal{F} \operatorname{ree}(V)$ , that is if we forget the differential on V then  $\mathcal{O} \coprod_{\tau} \mathcal{F} \operatorname{ree}(V)$  is the coproduct of  $\mathcal{O}$  by a free operad over a free graded  $\Sigma$ -module V. Note that any cofibration is a retract of a quasi-free extension. A quasi-free operad is an operad which is free over a free  $\Sigma$ -module if we forget the differential.

The following proposition will be fundamental for our applications.

1.2.1. **Proposition.** Let V be a free graded  $\Sigma_p$ -module together with  $d_V: V \longrightarrow \mathcal{O}(p) \oplus V$ 

such that  $d_V + d_{\mathcal{O}}$  is of square zero. Then if V is bounded above, the morphism  $\mathcal{O} \to \mathcal{O} \coprod_{\tau} \mathcal{F} \operatorname{ree}(V)$  is a cofibration.

**Proof.** The boundness assumption is needed in order to build a map

$$\mathcal{O} \longrightarrow \mathcal{O} \coprod_{\tau} \mathcal{F} \operatorname{ree}(V)$$

by induction on the degree of V, using the cellular attachment process. In order to add cells the assumption  $(d_V + d_O)^2 = 0$  is needed.  $\Box$ 

1.3. Homotopy invariance principle. Let  $\mathcal{O}$  be a cofibrant operad. The category of  $\mathcal{O}$ -algebras is also a closed model category where weak equivalences are quasi-isomorphisms and fibrations are epimorphisms [4] and [11].

Recall that the category of vector spaces is a closed model category, where weak equivalences are quasi-isomorphisms and fibrations are epimorphisms. In this category all objects are fibrant and cofibrant.

The following theorem is stated in [7]; its proof relies on a general result of Berger and Moerdijk about transfer of algebraic structure in closed model categories [4]. Such a result was proved in caracteristic zero by Markl [19] and for topological spaces by Boardman and Vogt [5].

# 1.3.1. Theorem. (Homotopy invariance principle.) [7]

Let  $\mathcal{O}$  be a cofibrant operad and assume that the morphism of vector spaces

$$f: X \longrightarrow Y$$

is a weak equivalence between vector spaces. Assume that X is a  $\mathcal{O}$ -algebra. Then Y is also provided with a  $\mathcal{O}$ -algebra structure. For

any cofibrant replacement  $\tilde{X}$  of X, there exists a sequence of quasiisomorphisms of  $\mathcal{O}$ -algebras:

$$X \longleftarrow \tilde{X} \longrightarrow Y.$$

such that the diagram commute in cohomology:



1.3.2. Corollary. Let  $\mathcal{O}$  be a cofibrant operad. Let C be a  $\mathcal{O}$ -algebra and let H be its cohomology. Then H is a  $\mathcal{O}$ -algebra and there is a sequence of quasi-isomorphisms of  $\mathcal{O}$ -algebras



where  $\tilde{C}$  is a cofibrant replacement of C and such that  $H^*(\Phi) = H^*(\psi)$ .

**Proof.** Set  $X = C, Y = H^*(C)$ . Choose a decomposition of vector spaces of  $X = Z \oplus Q$ , where Z is the kernel of the differential and let  $f: X \to Y$  be the composite of the projections onto Z and onto H. Then f is a weak equivalence of vector spaces, since  $H^*(f) = \text{Id}$ , and we can apply the previous theorem.

## 2. Adem-Cartan operads

In this section, the operad  $\mathcal{L}ev^{AC}$  is built, as a first step towards a resolution of the operad  $\mathcal{L}ev$ . The latter operad governs level algebras which are commutative algebras satisfying the 4-terms relation (2.1). The idea of introducing level algebras instead of commutative algebras in order to deal with Cartan and Adem relations is inspired by the fact that these relations are not conditioned by the associativity of the product. In this section we define also the family of Adem-Cartan operads and prove that  $E_{\infty}$ -operads belong to this family.

2.1. The operad  $\mathcal{L}ev.$  A *level algebra* A is a vector space together with a commutative product \* (non necessarily associative) satisfying the relation

$$(a * b) * (c * d) = (a * c) * (b * d), \ \forall a, b, c, d \in A.$$
(2.1)

2.1.1. **Definition.** Let  $\mathbb{F}_2$  be the trivial representation of  $\Sigma_2$  (generated by the operation  $\mu$ ) and  $R_{\mathcal{L}ev}$  be the sub- $\Sigma_4$ -module of  $\mathcal{F}ree(\mathbb{F}_2)(4)$ generated by the elements  $\mu(\mu, \mu) \cdot (\mathrm{Id} + \sigma)$  for all  $\sigma \in \Sigma_4$ . Then the operad  $\mathcal{L}ev$  is the operad

$$\mathcal{L}$$
ev =  $\mathcal{F}$ ree $(\mathbb{F}_2)/ < R_{\mathcal{L}$ ev > 1

Algebras over this operad are level algebras.

2.1.2. **Remark.** Since a commutative and associative algebra is trivially a level algebra, there is a morphism of operads

$$\mathcal{L}\mathrm{ev}\longrightarrow\mathcal{C}\mathrm{om},$$

where  $\mathcal{C}$ om denotes the operad defining commutative and associative algebras.

2.1.3. **Definition.** For any  $\Sigma_2$ -module M, the vector space

 $\mathcal{F}$ ree(M)(4) is a direct sum of two  $\Sigma_4$ -modules: the one indexed by trees of *shape 1* denoted by  $\mathbb{T}_1$ , that is the  $\Sigma_4$ -module generated by all the compositions  $\mu \circ_1 \gamma$  for  $\mu \in M$  and  $\gamma \in \mathcal{F}$ ree(M)(3) and the one indexed by trees of *shape 2* denoted by  $\mathbb{T}_2$ , that is the  $\Sigma_4$ -module generated by all the compositions  $\mu(\nu, \eta)$  for  $\mu, \nu, \eta \in M$ .



As an example, since there is only one generator  $\mu \in \mathcal{F}ree(\mathbb{F}_2)(2)$ , the dimension of  $\mathbb{T}_1(\mathcal{F}ree(\mathbb{F}_2))$  is 12 and the dimension of  $\mathbb{T}_2(\mathcal{F}ree(\mathbb{F}_2))$  is 3, whereas the dimension of  $\mathbb{T}_1(\mathcal{L}ev)$  is 12 and the dimension of  $\mathbb{T}_2(\mathcal{L}ev)$  is 1.

2.2. Construction of  $\mathcal{L}ev^{AC}$ . The construction of this operad is done by the process of attaching cells (see 1.2) to the standard bar resolution  $\mathcal{E}$  of  $\mathbb{F}_2[\Sigma_2]$ . 2.2.1. The standard bar resolution. Let (21) be the non trivial permutation of  $\Sigma_2$ . The standard bar resolution of  $\Sigma_2$  over  $\mathbb{F}_2$  is given by

$$\mathcal{E}^{-i} = \begin{cases} < e_i, e_i \cdot (21) > & \text{if } i \ge 0, \\ 0, & \text{if } i < 0, \end{cases}$$

and 
$$d(e_i) = e_{i-1} + e_{i-1} \cdot (21)$$
, with  $e_{-1} = 0$ .

Denote again by  $\mathcal{E}$  the free operad generated by the  $\Sigma$ -module  $\mathcal{E}$ .

Since  $\mathcal{L}ev = \mathcal{F}ree(\mathbb{F}_2) / \langle R_{\mathcal{L}ev} \rangle$ , there is a fibration of operads:

$$p: \mathcal{E} \xrightarrow{\sim} \mathcal{F}ree(\mathbb{F}_2) \longrightarrow \mathcal{L}ev.$$

By definition of  $\mathcal{L}ev$ , p(n) is clearly a quasi-isomorphism for n < 4. Furthermore p(4) is a quasi-isomorphism on pieces of shape 1 (see 2.1.3).

Note that for any operad  $\mathcal{O}$  such that  $\mathcal{O}(2)$  is a  $\Sigma_2$ -projective resolution of  $\mathbb{F}_2$ , there exists a morphism of operads  $m : \mathcal{E} \to \mathcal{O}$  such that m(2) is a homotopy equivalence. Then the image  $m(e_i) \in \mathcal{O}(2)$  is non zero. In the sequel, this image will be denoted also by  $e_i$ .

2.2.2. Notation. –We use May's convention: for any integers i and j the symbol (i, j) denotes  $\frac{(i+j)!}{i!j!} \in \mathbb{F}_2$ , if  $i \ge 0$  and  $j \ge 0$  and (i, j) = 0 otherwise. If the 2-adic expansion of i and j is  $i = \sum a_k 2^k$  and  $j = \sum b_k 2^k$  then  $(i, j) = 0 \in \mathbb{F}_2$  if and only if there exists k such that  $a_k = b_k = 1$ .

For any  $\sigma, \tau, \nu \in \mathcal{E}(2)$  we denote  $\sigma \cdot (21)(\tau, \nu)$  by  $\sigma(\tau, \nu) \cdot F$  which is equal the to  $\sigma(\nu, \tau) \cdot (3412)$  (see (1.1)).

## 2.2.3. Definition.

a) Let define some elements  $[u_n^m]_x \in \mathcal{E}(4)^{-n-x}$ , for m > 0 which are a combination of elements of shape 2. For any m such that  $2^k \leq m \leq 2^{k+1} - 1$ ,

$$[u_0^m]_x = e_x \cdot (21)^{m-1}(e_0, e_0), \text{ and for } n > 0,$$
  

$$[u_n^m]_x = \sum_{i=0}^{2^{k+1}-1} \sum_{0 \le 2^{k+1}\delta - i \le n} (n-m+i, m-1)[$$
  

$$(i,m)e_x \cdot (21)^{m-1}(e_{2^{k+1}\delta - i}, e_{n+i-2^{k+1}\delta}) +$$
  

$$(i-1,m)e_x \cdot (21)^{m-1}(e_{2^{k+1}\delta - i}, e_{n+i-2^{k+1}\delta} \cdot (21))]$$
(2.2)

### ADEM-CARTAN OPERADS

b) Let define some elements  $\alpha_{n,p} \in \mathcal{E}(4)^{-n}$ , for  $p \in \mathbb{Z}$ 

$$\alpha_{n,p} = \sum_{s=0}^{n-p} [u_{s+p}^{s+1}]_{n-s-p} + \sum_{s=0}^{n-p-1} [u_{s+p+1}^{s+1}]_{n-s-p-1} \cdot (3412)$$
(2.3)

2.2.4. **Proposition.** The  $[u_n^m]_x$ 's satisfy the following properties:

$$[u_n^m]_x = 0, \text{ for } 0 < n < m \tag{2.4}$$

$$[u_m^m]_x = e_x \cdot (21)^{m-1}(e_0, e_m) + e_x \cdot (21)^{m-1}(e_m, e_0 \cdot (21))$$
(2.5)

$$d[u_{n+1}^{m+1}]_x = [u_n^{m+1}]_x (\mathrm{Id} + (2143)) + [u_n^m]_x (F + (4321)) + [u_{n+1}^{m+1}]_{x-1} (\mathrm{Id} + F)$$
(2.6)

The  $\alpha_{n,p}$ 's satisfy the following property:

$$\alpha_{n,p} = 0, \text{ if } p < 0,$$

$$d\alpha_{n,p} = \alpha_{n-1,p-1} \cdot (\mathrm{Id} + (2143)) + \alpha_{n-1,p} \cdot (\mathrm{Id} + (4321))$$
(2.7)

The proof of this proposition is postponed to the last section.

2.2.5. **Examples.** By definition we have for all  $n \ge 0$ 

$$[u_n^1]_x = \sum_k e_x(e_{2k}, e_{n-2k}) + \sum_l e_x(e_{2l+1}, e_{n-2l-1} \cdot (21))$$
$$= e_x(\psi(e_n))$$

where  $\psi$  is the coproduct in the standard bar resolution  $\mathcal{E}$  of  $\mathbb{F}_2$  (see [18], [3]). And for  $n \geq 0$ , one has

$$[u_n^2]_x = \begin{cases} \sum_{0 \le 2\delta \le n} e_x \cdot (21)(e_{2\delta}, e_{n-2\delta} \cdot (21)^{\delta}) & \text{if n even} \\ \sum_{0 \le 4\delta - 1 \le n} e_x \cdot (21)(e_{4\delta - 1}, e_{n+1 - 4\delta}((12) + (21))) & \text{if n odd} \end{cases}$$

2.2.6. **Theorem.** There exists a cofibrant operad  $\mathcal{L}ev^{AC}$  satisfying the following properties:

- a)  $\mathcal{L}ev^{AC}(2) = \mathcal{E};$
- b) there is a fibration  $f : \mathcal{L}ev^{AC} \to \mathcal{L}ev$  such that f(n) is a quasiisomorphism for n < 4 and induces an isomorphism  $H^0(\mathcal{L}ev^{AC}(n)) \simeq \mathcal{L}ev(n);$

c) there are elements 
$$G_n^m \in \mathcal{L}ev^{AC}(4)$$
 of degree  $-n$ , satisfying  
 $G_n^m = 0$  for  $m \le 0$  or  $n < m$   
 $\mathrm{d}G_m^m = G_{m-1}^{m-1} \cdot (3214)(\mathrm{Id} + (2143)) + G_{m-1}^{m-2} \cdot (\mathrm{Id} + (4321))$   
 $+ \alpha_{m-1,m-1-p} + \alpha_{m-1,p} \cdot (3214)$   
and for  $n > m$   
 $\mathrm{d}G_n^m = G_{n-1}^m \cdot (\mathrm{Id} + (2143)) + G_{n-1}^{m-2} \cdot (\mathrm{Id} + (4321))$   
 $+ \alpha_{n-1,n-1-p} + \alpha_{n-1,p} \cdot (3214)$   
where  $p$  is the integer part of  $\frac{m-1}{2}$ .

**Proof.** The proof consists of building a sequence of cofibrant operads  $\mathcal{L}ev_m^{AC}$ ,  $m \geq 0$ , satisfying a) and b) where  $\mathcal{L}ev_m^{AC} \to \mathcal{L}ev_{m+1}^{AC}$  is a cofibration obtained by operadic cellular attachment. More precisely  $\mathcal{L}ev_0^{AC} = \mathcal{E}$ ; assume  $\mathcal{L}ev_m^{AC}$  is built and let  $V^{m+1}$  be the free graded  $\Sigma_4$ -module generated by elements  $G_n^{m+1}$  of degree -n for  $n \geq m+1$  with  $d: V^{m+1} \to \mathcal{L}ev_m^{AC}(4) \oplus V^{m+1}$  defined by the relation (2.8). If  $d^2 = 0$ , according to proposition 1.2.1, then

$$\mathcal{L}ev_m^{AC} \to \mathcal{L}ev_m^{AC} \coprod_{\tau} \mathcal{F}ree(V^{m+1}) =: \mathcal{L}ev_{m+1}^{AC}$$

is a cofibration. The main ingredient to prove that  $d^2 = 0$  is the relation (2.7).

$$\begin{aligned} First \ case: \ n > m + 1; \ then \ n - 1 > m \ and \ n - 1 > m - 2 \\ (d)^2(G_n^m) &= d(G_{n-1}^m \cdot (\mathrm{Id} + (2143)) + G_{n-1}^{m-2} \cdot (\mathrm{Id} + (4321))) \\ &+ d(\alpha_{n-1,n-1-p} + \alpha_{n-1,p} \cdot (3214)) \\ &= G_{n-2}^m \cdot (\mathrm{Id} + (2143))^2 + G_{n-2}^{m-2} \cdot (\mathrm{Id} + (4321))(\mathrm{Id} + (2143)) \\ &+ G_{n-2}^{m-2} \cdot (\mathrm{Id} + (2143))(\mathrm{Id} + (4321)) + G_{n-2}^{m-4} \cdot (\mathrm{Id} + (4321))^2 \\ &+ \alpha_{n-2,n-2-p} \cdot (\mathrm{Id} + (2143)) + \alpha_{n-2,p} \cdot (3214)(\mathrm{Id} + (2143)) \\ &+ \alpha_{n-2,n-2-(p-1)} \cdot (\mathrm{Id} + (4321)) \\ &+ \alpha_{n-2,p-1} \cdot (3214)(\mathrm{Id} + (4321)) \\ &+ d(\alpha_{n-1,n-1-p} + \alpha_{n-1,p} \cdot (3214)) \end{aligned}$$

Since (2143) and (4321) commute and are permutations of order 2 the 2 first lines vanish. Furthermore

$$d\alpha_{n-1,n-1-p} = \alpha_{n-2,n-2-p} \cdot (\mathrm{Id} + (2143)) + \alpha_{n-2,n-1-p} \cdot (\mathrm{Id} + (4321)),$$

and 
$$(3214)(\mathrm{Id} + (4321)) = (\mathrm{Id} + (2143))(3214)$$
 implies

$$d\alpha_{n-1,p} \cdot (3214) = \alpha_{n-2,p-1} \cdot (3214)(\mathrm{Id} + (4321)) + \alpha_{n-2,p} \cdot (3214)(\mathrm{Id} + (2143))$$

which yields  $(d)^2(G_n^m) = 0$  for n > m + 1. Second case: n = m + 1, then n - 1 = m and n - 1 > m - 2; thus terms in  $\alpha_{u,v}$  will vanish as before and  $2 + \alpha m - 2$ 

$$(d)^{2}(G_{m+1}^{m}) = G_{m-1}^{m-1} \cdot (3214)(\mathrm{Id} + (2143))^{2} + G_{m-1}^{m-2} \cdot (\mathrm{Id} + (4321))(\mathrm{Id} + (2143)) + G_{m-1}^{m-2} \cdot (\mathrm{Id} + (2143))(\mathrm{Id} + (4321)) + G_{m-1}^{m-4} \cdot (\mathrm{Id} + (4321))^{2} = 0$$

$$\begin{split} Last \ case: \ n &= m, \ \text{then} \\ (d)^2(G_m^m) = &d(G_{m-1}^{m-1} \cdot (3214)(\text{Id} + (2143)) + G_{m-1}^{m-2} \cdot (\text{Id} + (4321))) \\ &+ d(\alpha_{m-1,m-1-p} + \alpha_{m-1,p} \cdot (3214)) \\ = &G_{m-2}^{m-2} \cdot [(3214)(\text{Id} + (2143))]^2 + G_{m-2}^{m-3} \cdot (\text{Id} + (4321))(3214)(\text{Id} + (2143)) \\ &+ G_{m-2}^{m-2} \cdot (\text{Id} + (2143))(\text{Id} + (4321)) + G_{m-2}^{m-4} \cdot (\text{Id} + (4321))^2 \\ &+ \alpha_{m-2,m-2-p'} \cdot (3214)(\text{Id} + (2143)) + \alpha_{m-2,p'} \cdot (3214)^2(\text{Id} + (2143)) \\ &+ \alpha_{m-2,m-2-(p-1)} \cdot (\text{Id} + (4321)) + \alpha_{m-2,p-1} \cdot (3214)(\text{Id} + (4321)) \\ &+ d(\alpha_{m-1,m-1-p} + \alpha_{m-1,p} \cdot (3214)) \end{split}$$

where p' is the integer part of  $\frac{m-2}{2}$  and p the integer part of  $\frac{m-1}{2}$ . It's easy to check that the 2 first lines vanish. Furthermore, if m = 2k + 1 is odd then p = k, p' = k - 1 and the last lines write

$$+ \alpha_{2k-1,k} (\mathrm{Id} + (4321)) \cdot (3214) + \alpha_{2k-1,k-1} (\mathrm{Id} + (2143)) + \alpha_{2k-1,k} \cdot (\mathrm{Id} + (4321)) + \alpha_{2k-1,k-1} \cdot (\mathrm{Id} + (2143)) (3214) + \mathrm{d}(\alpha_{2k,k} + \alpha_{2k,k} \cdot (3214)) = 0,$$

and if 
$$m = 2k + 2$$
 is even, then  $p = k$ ,  $p' = k$  and the last lines write  
+  $\alpha_{2k,k}(\mathrm{Id} + (4321)) \cdot (3214) + \alpha_{2k,k}(\mathrm{Id} + (2143))$   
+  $\alpha_{2k,k+1} \cdot (\mathrm{Id} + (4321)) + \alpha_{2k,k-1} \cdot (\mathrm{Id} + (2143))(3214)$   
+  $\mathrm{d}(\alpha_{2k+1,k+1} + \alpha_{2k+1,k} \cdot (3214)) = 0$ 

Let us prove that the operads  $\mathcal{L}ev_m^{AC}$  satisfy b). According to 2.2.1  $\mathcal{E}(n) \to \mathcal{L}ev(n)$  is a quasi-isomorphism for n < 4, so is  $\mathcal{L}ev_m^{AC}(n) \to \mathcal{L}ev_m^{AC}(n)$  $\mathcal{L}ev(n)$  for all m.

To prove that  $f_m : \mathcal{L}ev_m^{AC} \to \mathcal{L}ev$  induces an isomorphism  $H^0(f_m)$ for all m > 0 it suffices to prove that  $H^0(f_1(4))$  is an isomorphism; if so, since the relations defining  $\mathcal{L}ev$  are generated by a  $\Sigma_4$ -module, then  $H^0(\operatorname{Lev}_1^{AC}(n)) \to \operatorname{Lev}(n)$  is an isomorphism, for all n; if m > 1, we do not introduce cells in degree -1, hence  $H^0(f_m(n))$  is an isomorphism for all n.

$$H^{0}(\mathcal{L}ev_{1}^{AC}(4)) = \mathbb{T}_{1}(\mathcal{E}^{0}(4))/d(\mathbb{T}_{1}(\mathcal{E}^{1}(4))) \bigoplus_{\mathbb{T}_{2}(\mathcal{E}^{0}(4))/d[\mathbb{T}_{2}(\mathcal{E}^{1}(4)) \oplus G_{1}^{1} \cdot \mathbb{F}_{2}(\Sigma_{4})]}$$

the first summand being isomorphic to  $\mathbb{T}_1(\mathcal{L}ev(4))$  (see 2.2.1). To prove that the second summand is isomorphic to  $\mathbb{T}_2(\mathcal{L}ev(4))$  it suffices to prove that it is 1-dimensional (see 2.1.3). Let  $X = e_0(e_0, e_0) \in \mathcal{E}^0(4)$ and  $\overline{X}$  its class in the second summand.

- i) For all  $\sigma$  in the dihedral group  $D_1$  ( $\sigma$  satisfies { $\sigma(1), \sigma(2)$ }  $\subset$  $\{1,2\} \cup \{3,4\}$ , there exists  $u = e_i(e_j, e_k), i + j + k = 1$ , such that  $du = X + X \cdot \sigma$ . Hence for all  $\sigma \in D_1$  we have  $\overline{X \cdot \sigma} = \overline{X}$ .
- ii) Since  $dG_1^1 = X + X \cdot (3214)$ , for all  $\sigma \in D_1$  we have  $\overline{X \cdot (3214)\sigma} =$  $\overline{X}$ , i.e. for all  $\tau \in D_2 = \{\tau | \{\tau(1), \tau(2)\} \subset \{1, 4\} \cup \{2, 3\}\}$  we have  $\overline{X \cdot \tau} = \overline{X}$ .
- iii) Finally equality (3124) = (2134)(3214) implies  $X \cdot (3124)\sigma =$  $\overline{X}$  for all  $\sigma \in D_1$ , or for all  $\tau \in D_3 = \{\tau | \{\tau(1), \tau(2)\} \subset \{1, 3\} \cup$  $\{2,4\}\}, \overline{X\cdot\tau} = \overline{X}.$

Since  $\Sigma_4 = D_1 \cup D_2 \cup D_3$  we get the result. Finally we define  $\mathcal{L}ev^{AC}$  to be the limit over m of  $\mathcal{L}ev^{AC}_m$ . Then  $\mathcal{L}ev^{AC}$  satisfies a), b) and c). 

2.2.7. **Definition.** An Adem-Cartan algebra is an algebra over  $\mathcal{L}ev^{AC}$ .

### 2.3. Adem-Cartan operads.

2.3.1. **Definition.** An Adem-Cartan operad  $\mathcal{O}$  is an operad such that  $\mathcal{O}(2)$  is a  $\Sigma_2$ -projective resolution of  $\mathbb{F}_2$  together with a morphism of operads  $m : \mathcal{L}ev^{AC} \to \mathcal{O}$  such that m(2) is a quasi-isomorphism. More precisely, there exists non trivial  $e_i \in \mathcal{O}(2)^{-i}$  and  $G_n^m \in \mathcal{O}(4)^{-n}$ satisfying the relations (2.8).

2.3.2.  $E_{\infty}$ -operads. According to remark 2.1.2, there is a morphism  $\mathcal{L}ev \longrightarrow \mathcal{C}om$ . An  $E_{\infty}$ -operad E is a  $\Sigma_*$ -projective resolution of the operad  $\mathcal{C}om$ : for every r, E(r) is  $\Sigma_r$ -projective and there exists an acyclic fibration  $E \to \mathcal{C}om$ . The structure of closed model category on operads implies that for any cofibrant operad  $\mathcal{O} \to \mathcal{C}om$  there exists a morphism  $\mathcal{O} \to E$  such that the following diagram commutes:



For instance there exists a morphism  $\mathcal{L}ev^{AC} \to E$  such that the previous diagram commutes. Hence any  $E_{\infty}$ -operad is an Adem-Cartan operad. In particular the algebraic Barratt-Eccles operad  $\mathcal{BE}$  (studied in [3]) is an Adem-Cartan operad.

# 3. Adem-Cartan algebras and the extended Steenrod Algebra

The aim of this section is to prove that the cohomology of an algebra over an Adem-Cartan operad carries an action of the extended Steenrod algebra.

## 3.1. The extended Steenrod algebra $\mathcal{B}_2$ .

3.1.1. Generalized Steenrod powers. ([18], [17]) The extended Steenrod algebra, denoted by  $\mathcal{B}_2$ , is a graded associative algebra over  $\mathbb{F}_2$  generated by the generalized Steenrod squares  $\operatorname{Sq}^i$  of degree  $i \in \mathbb{Z}$ . These generators satisfy the Adem relation, if t < 2s:

$$\operatorname{Sq}^{t}\operatorname{Sq}^{s} = \sum_{i} {s-i-1 \choose t-2i} \operatorname{Sq}^{s+t-i}\operatorname{Sq}^{i}.$$

Note that negative Steenrod squares are allowed and that  $Sq^0 = Id$  is not assumed. If  $\mathcal{A}_2$  denotes the classical Steenrod algebra, then

$$\mathcal{A}_2 \cong rac{\mathcal{B}_2}{<\operatorname{Sq}^0 + \operatorname{Id} >}$$

3.1.2. **Definition.** As in the classical case, an *unstable module over*  $\mathcal{B}_2$ , is a graded  $\mathcal{B}_2$ -module together with the instability condition.

$$\operatorname{Sq}^n(x) = 0$$
 if  $|x| < n$ 

An unstable level algebra over  $\mathcal{B}_2$  is a graded level algebra (A, \*) which is an unstable module over  $\mathcal{B}_2$ , such that

$$\begin{cases} \operatorname{Sq}^{|x|}(x) = x * x, \\ \operatorname{Sq}^{s}(x * y) = \sum \operatorname{Sq}^{t}(x) * \operatorname{Sq}^{s-t}(y) \quad \text{(Cartan relation)} \end{cases}$$

3.1.3. **Remark.** The category of unstable algebras over  $\mathcal{A}_2$  is a full subcategory of the category of unstable level algebras over  $\mathcal{B}_2$ .

3.2. Cup-i products. Let A be an algebra over an operad  $\mathcal{O}$  such that  $\mathcal{O}(2)$  is a  $\Sigma_2$ -projective resolution of  $\mathbb{F}_2$ . The evaluation map

$$\mathcal{O}(2) \otimes A^{\otimes 2} \longrightarrow A,$$

defines cup-i products  $a \cup_i b = e_i(a, b)$  ( $e_i$ 's were defined in 2.2.1). Steenrod squares are defined by  $\operatorname{Sq}^{r}(a) = a \cup_{|a|-r} a = e_{|a|-r}(a, a)$ . Following P. May [18], define  $D_n(a) = e_n(a, a) = \operatorname{Sq}^{|a|-n}(a)$ . In that terminology, Adem relations read

$$\sum_{k} (k, v - 2k) D_{w-v+2k} D_{v-k}(a) = \sum_{l} (l, w - 2l) D_{v-w+2l} D_{w-l}(a),$$
(3.1)

and Cartan relation read

$$D_n(x*y) = \sum_{k=0}^n D_k(x) * D_{n-k}(y).$$
(3.2)

Note that at this stage, we do not know that an  $\mathcal{O}$ -algebra satisfies these two relations.

### 3.3. Main theorem.

3.3.1. Lemma. Let A be a graded algebra  $(d_A = 0)$  over an operad  $\mathcal{O}$ . Assume that  $o \in \mathcal{O}(n)$  is a boundary. Then  $o(a_1, \ldots, a_n) = 0$ , for all  $a_1, \ldots, a_n$  in A.

**Proof.** There exists  $\omega \in \mathcal{O}(n)$  such that  $d\omega = o$ . The Leibniz rule implies that

$$d_A(\omega(a_1,\ldots,a_n)) = o(a_1,\ldots,a_n) + \sum_i \omega(a_1,\ldots,d_A a_i,\ldots,a_n)$$
  
the result follows because  $d_A = 0$ .

and the result follows because  $d_A = 0$ .

3.3.2. Lemma. Let  $\mathcal{O}$  be an operad such that  $\mathcal{O}(2)$  is a  $\Sigma_2$ -projetive resolution of  $\mathbb{F}_2$ . For every graded  $\mathcal{O}$ -algebra A and for every  $a \in A$ ,

$$[u_n^m]_x(a, a, a, a) = (n - 2m, 2m - 1)D_x D_{\frac{n}{2}}(a)$$

where  $[u_n^m]_x$  denotes the image of  $[u_n^m]_x \in \mathcal{E}$  by  $\mathcal{E} \to \mathcal{O}$ .

The proof is postponed to the last section.

3.3.3. **Proposition.** Let A be a graded Adem-Cartan algebra, then it is a level algebra.

**Proof.** For any operad  $\mathcal{O}$  and any  $\mathcal{O}$ -algebra A,  $H^*(A)$  is a  $H^*(\mathcal{O})$ -algebra hence a  $H^0(\mathcal{O})$ -algebra. Since  $H^0(\mathcal{L}ev^{AC}) = \mathcal{L}ev$  and  $H^*(A) = A$ , one gets the result.

3.3.4. **Theorem.** Let A be a graded Adem-Cartan algebra then A is an unstable level algebra over the extended Steenrod algebra.

**Proof.** We have already proved in proposition 3.3.3 that a graded Adem-Cartan algebra is a level algebra. Assume first that it is a module over  $\mathcal{B}_2$  (Adem relation (3.1)). Hence the instability condition reads

$$\operatorname{Sq}^{n}(x) = x \cup_{|x|=n} x = 0, \text{ if } |x| - n < 0.$$

The next equality is also immediate

$$\operatorname{Sq}^{|x|}(x) = x \cup_0 x = e_0(x, x) = x * x.$$

The Cartan relation is given by  $dG_n^1$ ; according to lemma 3.3.1,  $dG_{n+1}^1(x, x, y, y) = 0$ . Using relations (2.8), (2.3) and (2.5), one has  $0 = G_n^1 \cdot (\mathrm{Id} + (2143))(x, x, y, y)$   $+ \alpha_{n,n}(x, x, y, y) + \alpha_{n,0} \cdot (3214)(x, x, y, y)$   $= 2G_n^1(x, x, y, y) + [u_n^1]_0(x, x, y, y)$   $+ \sum_s [u_s^{s+1}]_{n-s}(y, x, x, y) + [u_{s+1}^{s+1}]_{n-s-1} \cdot (3412)(y, x, x, y)$   $= \sum_{l=0}^n e_0(e_l, e_{n-l} \cdot (21)^l)(x, x, y, y) + [u_0^1]_n(y, x, x, y)$  $+ \sum_{s=0}^{n-1} e_{n-s-1} \cdot (21)^s(x * y, y \cup_{s+1} x) + e_{n-s-1} \cdot (21)^s(x \cup_{s+1} y, x * y)$ 

Using the commutativity of \* and  $\cup_i$  for all i, one gets

$$dG_{n+1}^{1}(x, x, y, y) = \sum_{l=0}^{n} D_{l}(x) * D_{n-l}(y) + D_{n}(x * y)$$

which gives the Cartan relation (3.2).

The proof of Adem relation (3.1) relies on lemma 3.3.2, and on the relation  $dG_{n+1}^m(a, a, a, a) = 0$ . Combined with the relation (2.8) one

gets  $\alpha_{n,p}(a, a, a, a) = \alpha_{n,n-p}(a, a, a, a)$  that is

$$\begin{split} \sum_{s} [u_{p+s}^{s}(a, a, a, a) + u_{p+s}^{s+1}(a, a, a, a)]_{n-p-s} = \\ \sum_{t} [u_{n-p+t}^{t}(a, a, a, a) + u_{n-p+t}^{t+1}(a, a, a, a)]_{p-t} \\ \Rightarrow \sum_{t} [(p-s, 2s-1) + (p-s-2, 2s+1)]D_{n-p-s}D_{\frac{p+s}{2}}(a) = \\ \sum_{t} [(n-p-t, 2t-1) + (n-p-t-2, 2t+1)]D_{p-t}D_{\frac{n-p+t}{2}}(a). \end{split}$$
But  $(x, y-2) + (x-2, y) = (x, y)$ , hence  
$$\sum_{s} (p-s, 2s+1)D_{n-p-s}D_{\frac{p+s}{2}}(a) = \end{split}$$

$$\sum_{t} (n - p - t, 2t + 1) D_{p-t} D_{\frac{n-p+t}{2}}(a).$$

Since the first term is zero as soon as p-s is odd, we can set s = p-2l, and also t = n - p - 2k. As a consequence,

$$\sum_{l} (2l, 2p - 4l + 1) D_{n-2p+2l} D_{p-l}(a) = \sum_{k} (2k, 2n - 2p - 4k + 1) D_{2p-n+2k} D_{n-p-k}(a)$$

Using the 2-adic expansion, one gets

$$2l, 2p - 4l + 1) = (l, p - 2l);$$
(3.3)

by setting w := p and v := n - p, one obtains

$$\sum_{l} (l, w - 2l) D_{v-w+2l} D_{w-l}(a) = \sum_{k} (k, v - 2k) D_{w-v+2k} D_{v-k}(a)$$
  
ich is the Adem relation (3.1).

which is the Adem relation (3.1).

3.3.5. Corollary. Let  $\mathcal{O}$  be an Adem-Cartan operad. The cohomology of any  $\mathcal{O}$ -algebra is an unstable level algebra over  $\mathcal{B}_2$ .

**Proof.** By definition 2.3.1 there is a morphism of operads  $\mathcal{L}ev^{AC} \to \mathcal{O}$ , then it suffices to prove the theorem for  $\mathcal{O} = \mathcal{L}ev^{AC}$ . Let A be a  $\mathcal{L}ev^{AC}$ -algebra, then  $H^*(A)$  is a level algebra. In order to prove the Adem-Cartan relations we compute the boundaries of  $G_n^1(a, a, b, b)$  and  $G_n^m(a, a, a, a)$  for cocycles a and b that represents classes [a] and [b] in the cohomology. These boundaries give the Adem-Cartan relations between  $e_o(a, b)$ ,  $e_i(a, a)$ ,  $e_j(b, b)$  which represent [a] \* [b],  $D_i([a])$  and  $D_j([b])$  respectively.

In particular for algebras over a  $E_{\infty}$ -operad, we recover the results of May ([18] and [15]).

3.3.6. **Example.** Given an  $\mathcal{E}$ -algebra, a natural question is to know whether it is possible to extend this structure into a structure of Adem-Cartan algebra. The following example shows that it can not be done by imposing the triviality of all  $G_n^m$ 's. Let us consider the torus  $\mathbb{T}^2 = S^1 \times S^1$ . Here is the algebraic model of

Let us consider the torus  $\mathbb{T}^2 = S^1 \times S^1$ . Here is the algebraic model of the normalized singular cochains of  $\mathbb{T}^2$  that we use:

$$A_{\mathbb{T}^2} = C^*(S^1) \otimes C^*(S^1).$$

The vector space  $A_{\mathbb{T}_2}$  is generated by:  $1 := 1 \otimes 1$  in degree zero,  $\alpha := a \otimes 1$  and  $\beta := 1 \otimes b$  in degree 1 and  $\alpha\beta = a \otimes b$  in degree 2. The differential is trivial on  $A_{\mathbb{T}^2}$ .

The  $\mathcal{E}$ -structure on  $C^*(S^1)$  is given by  $e_0(1,1) = 1$ ,  $e_1(a,a) = a$ where a is the generator in degree 1 and all the others are zero (see [3]). Hence the  $\mathcal{E}$ -structure on  $A_{\mathbb{T}^2}$  is given by the coproduct  $\psi$  on  $\mathcal{E}$ . Since the differential is zero one has  $d(G_3^1(\alpha, \beta, \alpha, \beta)) = 0$ . But  $dG_3^1 = G_2^1(\mathrm{Id} + (2143)) + \alpha_{2,2} + \alpha_{2,0} \cdot (3214)$ . Then, using the definition of the  $\alpha_{n,p}$ 's and the commutativity of the  $e_i$ 's, one has

$$d(G_3^1(\alpha, \beta, \alpha, \beta)) = G_2^1(\alpha, \beta, \alpha, \beta) + G_2^1(\beta, \alpha, \beta, \alpha) + e_0(e_1, e_1 \cdot (21))(\alpha, \beta, \alpha, \beta) + e_2(e_0, e_0)(\alpha, \beta, \alpha, \beta) = G_2^1(\alpha, \beta, \alpha, \beta) + G_2^1(\beta, \alpha, \beta, \alpha) + e_1(\alpha, \beta)^2 + e_2(\alpha\beta, \alpha\beta)$$

But 
$$e_1(\alpha, \beta) = \psi(e_1)(a \otimes 1, 1 \otimes b)$$
  
 $= (e_0 \otimes e_1 + e_1 \otimes e_0 \cdot (21))((a, 1) \otimes (1, b))$   
 $= e_0(a, 1) \otimes e_1(1, b) + e_1(a, 1) \otimes e_0(b, 1) = 0$   
and  $e_2(\alpha\beta, \alpha\beta) = (e_0 \otimes e_2 + e_1 \otimes e_1 \cdot (21) + e_2 \otimes e_0)((a, a) \otimes (b, b))$   
 $= a \otimes b = \alpha\beta$ 

thus,  $G_2^1(\alpha, \beta, \alpha, \beta) + G_2^1(\beta, \alpha, \beta, \alpha) = \alpha\beta$ which proves that the action of  $G_2^1$  is non zero.

### 4. Operadic secondary cohomological operations.

In this section we prove that there exist secondary cohomological operations on the cohomology of an Adem-Cartan algebra A, and that these operations coincide with the Adams operations in case  $A = C^*(X, \mathbb{F}_2)$  is the singular cochain complex of a space X.

4.1. Secondary cohomological operations on Adem-Cartan algebras. Let  $\mathcal{O}$  be an Adem-Cartan operad and A be an  $\mathcal{O}$ -algebra. Then A is a  $\mathcal{L}ev^{AC}$ -algebra, hence  $H^*(A)$  is endowed with a (non natural) structure of  $\mathcal{L}ev^{AC}$ -algebra (see corollary 1.3.2). Hence for any (m, p) there are morphisms

$$\theta^{m,p}: H^n(A) \to H^{4n-p-1}(A)$$

given by  $\theta^{m,p}(x) = G_{p+1}^m(x, x, x, x).$ 

Besides from corollary 3.3.5, the cohomology  $H^*(A)$  is an unstable level algebra over  $\mathcal{B}_2$ . Let  $x \in H^n(A)$  and

$$R_{Ad}(x) = \sum_{i} Sq^{m_i} Sq^{n_i}(x) = 0$$

be an Adem relation with  $x \in \bigcap_{i} \operatorname{Ker}(\operatorname{Sq}^{n_{i}})$ . Let  $c \in A^{n}$ , dc = 0 representing x. Then there exists (m, p) such that  $(dG_{p+1}^{m})(c, c, c, c) = R_{Ad}(c)$ :  $p = 3n - m_{i} - n_{i}$  and m depends on n and  $m_{i} + n_{i}$ . Since  $\operatorname{Sq}^{n_{i}}(x) = 0$ , there exists  $b_{i} \in A^{n+n_{i}-1}$  such that  $db_{i} = e_{n-n_{i}}(c, c)$ . The element

$$b = \sum_{i} e_{n-m_i+n_i}(1, e_{n-n_i})(b_i, c, c) + e_{n-m_i+n_i-1}(b_i, b_i)$$

satisfies  $d(G_{p+1}^m(c,c,c,c)+b) = 0.$ 

4.1.1. **Proposition.** The class of  $G_{p+1}^m(c, c, c, c) + b$  in  $H^{m_i+n_i+n-1}(A) / \sum_i Im(\operatorname{Sq}^{m_i})$  does not depend on the choices of the  $b'_i$ s and c.

**Proof.** First it does not depend on the choices of the  $b_i$ 's. Let  $b'_i \in A^{n+n_i-1}$  such that  $db'_i = db_i = e_{n-n_i}(c, c)$ , and

$$b' = \sum_{i} e_{n-m_i+n_i}(1, e_{n-n_i})(b'_i, c, c) + e_{n-m_i+n_i-1}(b'_i, b'_i)$$

Then d(b + b') = 0 and the following relation implies the result:

$$b + b' = \sum_{i} \operatorname{Sq}^{m_i}(b_i + b'_i) + \operatorname{d}e_{n-m_i+n_i}(b_i + b'_i, b_i).$$

Secondly it does not depend on the choice of a representant c of x. Using the homotopy invariance principle as in 1.3.2, there is a zig-zag of acyclic fibrations of Adem-Cartan algebras



with  $H^*(f) = H^*(g)$ . Let c and c' be two cocycles that represent x, and let  $u, u' \in \tilde{A}$  such that du = 0, f(u) = c, g(u) = x and the same for u', c'. Since  $\operatorname{Sq}^{n_i}(x) = 0$ , there exists  $v_i, v'_i$  such that  $dv_i = \operatorname{Sq}^{n_i}(u)$ and  $dv'_i = \operatorname{Sq}^{n_i}(u')$ . Let v, v' defined as b. Let us prove that

$$[G_{p+1}^m(c,c,c,c) + f(v)] = [G_{p+1}^m(c',c',c',c') + f(v')]$$

in  $H^{n+m_i+n_i-1}(A)/\sum_i Im(\operatorname{Sq}^{m_i})$ . This is equivalent to prove that g(v) + g(v') is in the sum of all  $Im(\operatorname{Sq}^{m_i})$ . But

$$v = \sum_{i} e_{n-m_i+n_i}(v_i, \mathrm{d}v_i) + \mathrm{Sq}^{m_i}(v_i)$$

and  $g(dv_i) = 0$  imply that  $g(v) = \sum_i \operatorname{Sq}^{m_i}(g(v_i)) \in \sum Im(\operatorname{Sq}^{m_i})$ , which proves the result.

As a consequence, we have defined a map

$$\psi^{m,p} : \bigcap_{i} \operatorname{Ker}(\operatorname{Sq}^{n_{i}}) \subset H^{n}(A) \longrightarrow H^{n+m_{i}+n_{i}-1}(A) / \sum_{i} Im(\operatorname{Sq}^{m_{i}})$$
$$x \qquad \mapsto \qquad [G_{p+1}^{m}(c,c,c,c)+b]$$

4.1.2. **Proposition.** Let  $\iota : \bigcap_{i} \operatorname{Ker}(\operatorname{Sq}^{n_{i}}) \to H^{n}(A)$  be the canonical inclusion and  $\pi : H^{n+m_{i}+n_{i}-1}(A) \to H^{n+m_{i}+n_{i}-1}(A) / \sum_{i} \operatorname{Im}(\operatorname{Sq}^{m_{i}})$  be the canonical projection. Then

$$\pi\theta^{m,p}\iota = \psi^{m,p}.$$

**Proof.** Using the proof of the previous proposition, it only remains to show that  $[G_{p+1}^m(c, c, c, c) + b] = G_{p+1}^m(x, x, x, x)$  in  $H^{m_i+n_i+n-1}(A) / \sum_i Im(\operatorname{Sq}^{m_i})$  for  $x \in \bigcap_i \operatorname{Ker}(\operatorname{Sq}^{n_i})$ . This is equivalent to prove that g(v) is in  $\sum_i Im(\operatorname{Sq}^{m_i})$  for v with f(v) = b, which has been already proved.  $\Box$ 

4.2. Adams operations. Let  $C^*(X; \mathbb{F}_2)$  denotes the singular cochains complex of a topological space X and  $H^*(X, \mathbb{F}_2)$  its cohomology. We recall that  $C^*(X; \mathbb{F}_2)$  is an algebra over a  $E_{\infty}$ -operad [12], hence an Adem-Cartan algebra. Adams defined in an axiomatic way stable secondary cohomological operations [1]. His approach is topological, and uses the theory of socalled "universal examples". These operations correspond to Adem relations

$$R_{Ad} = \sum_{i} \operatorname{Sq}^{m_i} \operatorname{Sq}^{n_i},$$

and are denoted by  $\Phi$ . Let recall Adams axioms:

**Axiom 1.** For any  $u \in H^n(X; \mathbb{F}_2)$ ,  $\Phi(u)$  is defined if and only if  $\operatorname{Sq}^{n_i}(u) = 0$  for all  $n_i$ .

Axiom 2. If  $\Phi(u)$  is defined then

$$\Phi(u) \in H^{m_i + n_i + n - 1}(X; \mathbb{F}_2) / \sum_i Im(\mathrm{Sq}^{m_i}).$$

Axiom 3. The operation  $\Phi$  is natural.

**Axiom 4.** Let (X, A) be a pair of topological spaces, we have the long exact sequence

$$\dots H^{n-1}(A; \mathbb{F}_2) \xrightarrow{\delta^*} H^n(X, A; \mathbb{F}_2) \xrightarrow{j^*} H^n(X; \mathbb{F}_2) \xrightarrow{i^*} H^n(A; \mathbb{F}_2) \xrightarrow{\delta^*} \dots$$

let  $v \in H^n(X, A; \mathbb{F}_2)$  be a class such that  $\phi$  is defined on  $j^*(v) \in H^n(X; \mathbb{F}_2)$ . Let  $w_i \in H^*(A; \mathbb{F}_2)$  such that  $\delta^*(w_i) = \operatorname{Sq}^{n_i}(v)$ . Then, we have

$$i^*\Phi(j^*(v)) = \sum_i \operatorname{Sq}^{m_i}(w_i) \in H^*(A; \mathbb{F}_2)/i^*(\sum_i \operatorname{Im}(\operatorname{Sq}^{m_i})).$$

Axiom 5. The operation  $\Phi$  commutes with suspension.

Later on, Kristensen proved that these operations can be defined at the cochain level, using the existence of a coboundary which creates the stable secondary cohomological operation defined by Adams ([14], chapter 6). More precisely, for an Adem relation  $R_{Ad}$  and a class  $x \in \bigcap_i \operatorname{Sq}^{n_i}$ , Kristensen defines cochain operations  $\theta$  such that the differential of  $\theta(c)$  (c is a representant of x) gives a cocycle representing an Adem relation  $R_{Ad}(x)$ . If one chooses  $b_i$  such that  $db_i = \operatorname{Sq}^{n_i}(c)$ , then one gets a cocycle, and a cohomology class

$$Qu^{r}(c) = [\theta(c) + \sum_{i} (e_{n-m_{i}+n_{i}}(1, e_{n-n_{i}})(b_{i}, c, c) + e_{n-m_{i}+n_{i}-1}(b_{i}, b_{i}))].$$

Then,

4.2.1. **Theorem.** (Kristensen, theorem 6.1 of [14]) Any operation  $x \mapsto Qu^r(c)$  satisfies axiom 1-5 of Adams.

4.2.2. Corollary. The maps  $\psi^{m,p}$  coincide with the stable secondary cohomological operations of Adams.

**Proof.** The proof relies on theorem 4.2.1 with  $\theta(c) = G_{n+1}^m(c, c, c, c)$ .

4.2.3. **Theorem.** The stable secondary cohomological operations  $\phi$ of Adams extend to maps  $\theta^{m,p}$  :  $H^n(X, \mathbb{F}_2) \to H^{n+m_i+n_i-1}(X, \mathbb{F}_2)$ . More precisely, if we denote by  $\iota$  :  $\bigcap_i \operatorname{Ker}(\operatorname{Sq}^{n_i}) \to H^n(X, \mathbb{F}_2)$  and  $\pi : H^{n+m_i+n_i-1}(X, \mathbb{F}_2) \to H^{m_i+n_i+n-1}(X, \mathbb{F}_2) / \sum_i \operatorname{Im}(\operatorname{Sq}^{m_i})$  then  $\phi = \pi \theta^{m,p} \iota.$ 

**Proof.** It is the translation of theorem 4.1.2 for  $A = C^*(X, \mathbb{F}_2)$ .

### 5. Proof of technical lemmas

In this section proposition 2.2.4 and lemma 3.3.2 are proved.

## 5.1. Lemma. For any $i, j \leq 2^p - 1$ one has

$$(i, j) = 0$$
, if  $i + j \ge 2^p$ , and  
 $(i, j) = (2^p - i - j - 1, j)$ .

**Proof.** Let  $\sum_{l=0}^{p-1} a_l 2^l$  and  $\sum_{l=0}^{p-1} b_l 2^l$  be the 2-adic expansion of i and j respectively. Recall that (i, j) = 1 if and only if the 2-adic expansion of i + j is  $\sum (a_l + b_l) 2^l$ . If  $i + j \ge 2^p$  this is not the case, thus (i, j) = 0. If (i, j) = 1 then the 2-adic expansion of  $2^p - 1 - i - j$  is  $\sum_{l=0}^{p-1} (1 - a_l - b_l) 2^l$ , thus the 2-adic expansion of  $(2^p - 1 - i - j) + j$  has for coefficients  $(1 - a_l - b_l) + (b_l)$ . Consequently  $(2^p - 1 - i - j, j) = 1$ . The converse is true by symmetry.

Note that the first assertion is a consequence of the second one, because if  $(i + j) \ge 2^p$  then  $(2^p - 1 - i - j) < 0$ , and  $(\alpha, \beta) = 0$  if  $\alpha < 0$  or  $\beta < 0$ .

5.2. **Proof of lemma 3.3.2.** Using the commutativity of  $\cup_x$  one has

$$[u_n^m]_x(a, a, a, a) = \sum_{i=0}^{2^{k+1}-1} \sum_{0 \le 2^{k+1}\delta - i \le n} (n-m+i, m-1)(i, m) D_{2^{k+1}\delta - i}(a) \cup_x D_{n+i-2^{k+1}\delta}(a) + (n-m+i, m-1)(i-1, m) D_{2^{k+1}\delta - i}(a) \cup_x D_{n+i-2^{k+1}\delta}(a) = \sum_{i,\delta} (n-m+i, m-1)(i, m-1) D_{2^{k+1}\delta - i}(a) \cup_x D_{n+i-2^{k+1}\delta}(a).$$

Let  $0 \leq j \leq 2^{k+1} - 1$  such that  $n + i \equiv -j$   $[2^{k+1}]$ , then there exists  $\delta'$  such that  $n + i - 2^{k+1}\delta = 2^{k+1}\delta' - j$ , and lemma 5.1 implies

$$(n - m + i, m - 1) = (2^{k+1}(\delta + \delta') - m - j, m - 1)$$
  
= (j, m - 1) (5.1)  
$$(n - m + j, m - 1) = (i, m - 1)$$

As a consequence, if  $i \neq j$  or i = j and  $\delta \neq \delta'$ , the 2 following terms in  $[u_n^m]_x(a, a, a, a)$ 

$$(n-m+i,m-1)(i,m-1)D_{2^{k+1}\delta-i}(a) \cup_x D_{2^{k+1}\delta'-j}(a) + (n-m+j,m-1)(j,m-1)D_{2^{k+1}\delta'-j}(a) \cup_x D_{2^{k+1}\delta-i}(a)$$

vanish. Hence if there exists  $(i,\delta)$  such that  $2^{k+1}\delta-i=n-(2^{k+1}\delta-i),$  then

$$[u_n^m]_x(a, a, a, a) = (n - m + i, m - 1)^2 D_x D_{\frac{n}{2}}(a).$$

Relation (3.3) implies (n - m + i, m - 1) = (2n - 2m + 2i, 2m - 1) = (n - 2m, 2m - 1). Furthermore, if (n - 2m, 2m - 1) = 1, then n is even and we can pick  $0 \le r \le 2^{k+1} - 1$  such that  $r \equiv -\frac{n}{2} [2^{k+1}]$ .

5.3. **Proof of proposition 2.2.4.** We have to prove the relations (2.4), (2.5), (2.6) and (2.7). The relation (2.7) is straightforward using relation (2.6) and definition (2.3).

Proof of relations (2.4) and (2.5)– Assume  $n \leq m$ . The condition  $0 \leq 2^{k+1}\delta - i \leq n \leq m \leq 2^{k+1} - 1$  implies  $\delta$  equals 0 or 1. If  $\delta = 0$ , then i = 0. If  $\delta = 1$ , then  $i + m \geq 2^{k+1}$  and (i, m) = 0 by lemma 5.1. So  $[u_n^m]_x$  writes

$$[u_n^m]_x = (n - m, m - 1)e_x \cdot (21)^{m-1}(e_0, e_n) + \sum_{i=0}^{2^{k+1}-1} (n - m + i, m - 1)(i - 1, m)e_x \cdot (21)^{m-1}(e_{2^{k+1}-i}, e_{n+i-2^{k+1}} \cdot (21))$$

If n < m, then (n - m, m - 1) = 0 and  $i - 1 + m \ge 2^{k+1}$  implies (i - 1, m) = 0, which proves relation (2.4).

If n = m, (i - 1, m) = 0 for all  $i \neq 2^{k+1} - m$  and  $(2^{k+1} - m, m - 1)(2^{k+1} - 1 - m, m) = (0, m - 1)(0, m)$  by virtue of lemma 5.1. This proves relation (2.5).

Proof of relation (2.6) – For the convenience of the reader, let

$$B_{x,[m],n}^{\delta,i} = e_x \cdot (21)^{m-1} (e_{2^{k+1}\delta-i}, e_{n+i-2^{k+1}\delta})$$

where [m] means  $m \mod 2$ , then

$$[u_n^m]_x = \sum_{i,\delta} (n-m+i,m-1)[(i,m)B_{x,[m],n}^{\delta,i} + (i-1,m)B_{x,[m],n}^{\delta,i} \cdot (1243)].$$

Remarks:

- a) Let  $P = \{ \mathrm{Id}, (2134), (2143), (1243) \} \subset \Sigma_4$ . Then the set  $\mathcal{F} = \{ B_{x,[m],n}^{\delta,i} \cdot \sigma, B_{x',[m+1],n'}^{\delta',j} \cdot \tau, \forall x, n, x', n', i, i', \delta, \delta', \forall \sigma, \tau \in P \}$  form a free system in  $\mathcal{E}(4)$ .
- b) For any  $\sigma \in P$ ,  $B_{x,[m],n}^{\delta,i} \cdot \sigma F = B_{x,[m],n}^{\delta,i} \cdot F\sigma = B_{x,[m+1],n}^{\delta,i} \cdot \sigma$ .

There are two cases to consider: if  $2^k \leq m \leq 2^{k+1} - 2$  (then  $m+1 \leq 2^{k+1} - 1$ ) or if  $m = 2^{k+1} - 1$ . Since computation are long but not difficult, we'll present only the first case.

$$d[u_{n+1}^{m+1}]_{x+1} = \sum_{i,\delta} \underbrace{(n-m+i,m)(i,m+1)}_{a_i} [B_{x+1,[m+1],n}^{\delta,i+1} \cdot (\mathrm{Id} + (2134)) + B_{x+1,[m+1],n}^{\delta,i} \cdot (\mathrm{Id} + (1243)) + B_{x,[m+1],n+1}^{\delta,i} \cdot (\mathrm{Id} + F)] + \underbrace{(n-m+i,m)(i-1,m+1)}_{b_i} [B_{x+1,[m+1],n}^{\delta,i+1} \cdot ((1243) + (2143)) + B_{x+1,[m+1],n}^{\delta,i} \cdot (\mathrm{Id} + F)(1243)] + B_{x+1,[m+1],n}^{\delta,i} \cdot (\mathrm{Id} + (1243)) + B_{x,[m+1],n+1}^{\delta,i} \cdot (\mathrm{Id} + F)(1243)]$$

$$[u_n^{m+1}]_{x+1}(\mathrm{Id} + ((2143))) = \sum_{i,\delta} \underbrace{(n-m+i-1,m)(i,m+1)}_{c_i} B_{x+1,[m+1],n}^{\delta,i}(\mathrm{Id} + (2143)) + \underbrace{(n-m+i-1,m)(i-1,m+1)}_{d_i} B_{x+1,[m+1],n}^{\delta,i}((1243) + (2134))$$

$$\begin{split} [u_n^m]_{x+1}(F+(4321)) &= \\ & \sum_{i,\delta} \underbrace{(n-m+i,m-1)(i,m)}_{e_i} B_{x+1,[m],n}^{\delta,i}(F+(4321)) \\ & + \underbrace{(n-m+i,m-1)(i-1,m)}_{f_i} B_{x+1,[m],n}^{\delta,i}(1243)(F+(4321)) \end{split}$$

Note that

 $B_{x+1,[m],n+1}^{\delta,i} \cdot (4321) = e_{x+1} \cdot (21)^m (e_{n+i-2^{k+1}\delta}, e_{2^{k+1}\delta-i}) \cdot (3412)(4321)$ Hence by using relation (5.1) we get

$$\sum_{i,\delta} (n-m+i,m-1)(i,m) B_{x+1,[m],n}^{\delta,i} \cdot (4321) = \sum_{j,\delta'} \underbrace{(n-m+j-1,m)(j,m-1)}_{m_j} B_{x+1,[m+1],n}^{\delta',j} (2143)$$

and

$$\sum_{i,\delta} (n-m+i,m-1)(i-1,m)B_{x+1,[m],n}^{\delta,i}(1243)(4321) = \sum_{j,\delta'} \underbrace{(n-m+j,m)(j,m-1)}_{l_j} B_{x+1,[m+1],n}^{\delta',j}(1243)$$

$$[u_{n+1}^{m+1}]_{x}(1+F) = \sum_{i,\delta} \underbrace{(n-m+i,m)(i,m+1)}_{g_{i}} B_{x,[m+1],n+1}^{\delta,i}(\mathrm{Id}+F) + \underbrace{(n-m+i,m)(i-1,m+1)}_{h_{i}} B_{x,[m+1],n+1}^{\delta,i}(1243)(\mathrm{Id}+F)$$

Thus to prove relation (2.6), it suffices to prove that the sum of all the coefficients of elements of  $\mathcal{F}$  vanishes. For instance, the coefficient of  $B_{x+1,[m+1],n}^{\delta,i} \cdot \text{Id is } a_{i-1} + a_i + b_i + c_i + e_i \text{ that is}$ (n-m+i-1,m)(i-1,m+1) + (n-m+i,m)(i,m+1) + (n-m+i,m)(i,m+1) + (n-m+i,m)(i-1,m+1) + (n-m+i-1,m)(i,m+1) + (n-m+i,m-1)(i,m)

$$=(n-m+i,m-1)(i,m)$$
  
=(n-m+i,m-1)(i-1,m+1) + (n-m+i,m-1)(i,m+1) + (n-m+i,m-1)(i,m) = 0.

The coefficient of  $B_{x+1,[m+1],n}^{\delta,i} \cdot (2143)$  is  $b_{i-1} + c_i + m_i$  that is (n-m+i-1,m)(i-2,m+1) + (n-m+i-1,m)(i,m+1)+ (n-m+i-1,m)(i,m-1) = 0.

All the others computation go the same.

#### ADEM-CARTAN OPERADS

Acknowledgements. The authors would like to thank Benoit Fresse for fruitful discussions. They are especially indebted to Lionel Schwartz for pointing them the importance of level algebras in algebraic topology. The first author would like to thank Marc Aubry, Yves Felix and Jean-Claude Thomas for their support, the paper was written during his stay at the University of Louvain-La-Neuve and at the CRM (Barcelona). The second author would like to thank the University of Louvain-La-Neuve and the CRM for their hosting.

### References

- J. F. ADAMS: On the non-existence of elements of Hopf invariant one, Ann. of Math. (2)72, (1960), 20–104.
- J. ADEM: The iteration of the Steenrod squares in algebraic topology, Proc. Nat. Acad. Sci. U. S. A. 38, (1952), 720–726.
- [3] C. BERGER, B. FRESSE: Combinatorial operad actions on cochains, Preprint (2001), math.AT/0204326.
- [4] C. BERGER, I. MOERDIJK: Axiomatic homotopy theory for operads, Prépublication de l'université de Nice, 651 (2002).
- [5] J. M. BOARDMAN, R. VOGT: Homotopy invariant algebraic structures on topological spaces, Lecture Notes in Mathematics, 347, Springer-Verlag, 1973.
- [6] H. CARTAN: Sur l'itération des opérations de Steenrod, Comment. Math. Helv. 29, (1955), 40–58.
- [7] D. CHATAUR: Deformations of  $E_{\infty}$ -algebras, Preprint (2001).
- [8] W.G. DWYER, J. SPALINSKI: Homotopy theory and model categories, in "Handbook of algebraic topology", North-Holland (1995), 73–126.
- [9] E. GETZLER, J.D.S. JONES: Operads, homotopy algebras and iterated integrals for double loop spaces, preprint (1994), hep-th/9403055.
- [10] V. GINZBURG, M. KAPRANOV: Koszul duality for operads, Duke Math. J. (1)76, (1994), 203–272.
- [11] V. HINICH: Homological algebra of homotopy algebras, Comm. Algebra (10)25, (1997), 3291–3323.
- [12] V. HINICH-V. SCHECHTMANN: On homotopy limit of homotopy algebras, in "K-theory, arithmetic and geometry (Moscow, 1984–1986)", Lecture Notes in Math., **1289** (1987), 240-264.
- [13] M. HOVEY; Model categories, Mathematical Surveys and Monographs, American Mathematical Society 63, (1999).
- [14] L. KRISTENSEN; On secondary cohomology operations, Math. Scand. 12, (1963), 57–82.
- [15] I. KRIZ-J.P. MAY: Operads, algebras, modules and motives, Astérisque, 233 (1995).
- [16] J. L. LODAY:La renaissance des opérades, in "Séminaire Bourbaki 1994-1995", Astérisque, 236, (1996), 47–74.
- [17] M.MANDELL: $E_{\infty}$ -algebras and p-adic homotopy theory, Topology (1) 40, (2001), 43–94.

- [18] J.P. MAY: A general algebraic approach to Steenrod operations, in "the Steenrod algebra and its applications" Lecture Notes in Math. 168 (1970), 153–231.
- [19] M. MARKL: homotopy algebras are homotopy algebras, preprint (1999), math.AT/9907138.
- [20] J.E. MCCLURE, J.H. SMITH: a solution of Deligne's Hochschild cohomology conjecture, in "Recent progress in homotopy theory (Baltimore, MD, 2000)" Contemp. Math. 293 (2002), 153–193,
- [21] D. QUILLEN: Homotopical Algebra, Lecture Notes in Math. 43 (1967).
- [22] N. E. STEENROD: Products of cocycles and extensions of mappings, Ann. of Math. (2) 48, (1947), 290–320.
- [23] N. E. STEENROD: Reduced powers of cohomology classes, Ann. of Math. (2)56, (1952), 47–67

E-mail address: dchataur@tonton.univ-angers.fr, livernet@math.univ-paris13.fr







