ON NORMAL STRATIFIED PSEUDOMANIFOLDS

G. PADILLA

Devoted to the victims of the natural tragedy in Vargas, Dec.15/1999, who died under the rage of Waraira Repano.

ABSTRACT. Any pl-stratified pseudomanifod can be normalized preserving its intersection homology. In this paper we extend this result for any topological stratified pseudomanifold and for a family of perversities which is larger than usual. Our construction is functorial. We also give a detailed description of the normalizer's stratification in terms of the initial stratified pseudomanifold.

FOREWORD

For an entire version of this article the reader should go better to Extracta Math. Vol. 15 (3), 383-412.

A stratified pseudomanifold is normal if its links are connected. A normalization of a stratified pseudomanifold X is a normal stratified pseudomanifold X^N together with a finite-to-one projection $\mathfrak{n}: X^N \to X$ preserving the intersection homology. Recall that intersection homology is the suitable algebraic tool for the stratified point of view: it was first introduced by Goresky and MacPherson in the pl-category and later extended for any topological stratified pseudomanifold [3], [4]. Following Borel the map \mathfrak{n} is usually required to satisfy the following property: For each $x \in X$ there is a distinguished neighborhood U such that the points of $\mathfrak{n}^{-1}(x)$ are in correspondence with the connected components of the regular part of U. A normalization satisfying the above condition always exists for any pl-stratified pseudomanifold [1], [6]. In this article we study the main properties of the map \mathfrak{n} . More precisely, we prove that \mathfrak{n} can be required to satisfy a stronger condition: it is a locally trivial stratified morphism preserving the conical structure transverse to the strata. We make an explicit construction of such a normalization for any topological stratified pseudomanifold. Our construction is functorial, thus unique. We exhibit the relationship between the stratifications of X and X^N . Finally we prove that the normalization preserves the intersection homology with the family of perversities given in [7], see also [5], [9]. This family of perversities is larger than the usual one. With little adjust our procedure holds also in the C^{∞} category.

Date: May 9/2002, revised on June 15/2002.

¹⁹⁹¹ Mathematics Subject Classification. Primary 55N33; Secondary 35S35.

Key words and phrases. Intersection Homology, Stratified Pseudomanifolds.

This paper was supported by the the ECOSNord project, Euskal Herria Unibertsitatea's Math Department, the University of Artois' LaboGA and the Universidad Central de Venezuela's C.D.C.H. The author would like to thank M. Saralegi for encouraging us to look at this problem, and also R. Popper for some helpful conversations.

2 G. PADILLA

References

- [1] BOREL, A. et al. Intersection Cohomology. PM#50(1984) Birkhäuser.
- [2] BREDON, G. Topology and Geometry. GTM#139 (1993) Springer-Verlag.
- [3] GORESKY, M. & R. MacPherson. Intersection Homology Theory. Topology #19 (1980), 135-162.
- [4] GORESKY, M. & R. MacPherson. Intersection Homology II. Invent. Math. #71 (1983), 77-129.
- [5] KING, H. Topology invariance of intersection homology without sheaves. Topoloy Appl. 20 (1984), 1091-1106.
- [6] MACCRORY, C. Poincaré duality in spaces with singularities. Ph.D. Thesis. Brandeis University. (1972).
- [7] MACPHERSON, R. Intersection homology and perverse sheaves. Colloquium Lectures, Annual Meeting of the AMS, San Francisco, June 1991.
- [8] PFLAUM, M. Analytic and geometric study of Stratified Spaces. (2000) Springer-Verlag.
- [9] SARALEGI, M. Homological Properties of Stratified Spaces. Illinois J. Math. #38 (1994), 47-70.

Universidad Central de Venezuela, Caracas $E\text{-}mail\ address$: gabrielp@euler.ciens.ucv.ve