

The serial harness interacting with a wall

P.A. Ferrari¹ L.R.G. Fontes¹ B. Niederhauser^{*,1}
M. Vachkovskaia²

¹Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, CEP 05508–090, São Paulo SP, Brazil.

E-mails: pablo@ime.usp.br, lrenato@ime.usp.br, beat@ime.usp.br
<http://www.ime.usp.br/~pablo>

²Instituto de Matemática, Estatística e Computação Científica, Universidade Estadual de Campinas. Caixa Postal 6065, CEP 13083–970, Campinas SP, Brazil.

E-mail: marinav@ime.unicamp.br

Abstract The serial harnesses introduced by Hammersley describe the motion of a hypersurface of dimension d embedded in a space of dimension $d + 1$. The height assigned to each site i of \mathbb{Z}^d is updated by taking a weighted average of the heights of some of the neighbors of i plus a “noise” (a centered random variable). The surface interacts by exclusion with a “wall” located at level zero: the updated heights are not allowed to go below zero. We show that for any distribution of the noise variables and in all dimensions, the surface delocalizes. This phenomenon is related to the so called “entropic repulsion”. For some classes of noise distributions, characterized by their tail, we give explicit bounds on the speed of the repulsion.

Key words and phrases harness, surface dynamics, entropic repulsion

AMS 1991 subject classifications. 60K35 82B 82C

1 Introduction and results

The harnesses are stochastic processes introduced by Hammersley (1965) to model the long range correlations in the cristaline structure of metals. The serial harness models the time evolution of a hypersurface of dimension d embedded in a $d + 1$ dimensional space. At each site $i \in \mathbb{Z}^d$ at time $n \geq 0$ a quantity $Y_n(i) \in \mathbb{R}$ stays for the height of the surface. The initial configuration is the flat surface $Y_0(i) = 0$ for all i . The evolution is described

*Supported by FAPESP (00/05134–5)

as follows. At each (discrete) moment $n \geq 0$ the height at each site is substituted by a weighted average of the heights at the previous moment plus a symmetric random variable.

Let $\mathcal{P} = \{p(i, j)\}_{i, j \in \mathbb{Z}^d}$ be a stochastic matrix, i.e. $p(i, j) \geq 0$ and $\sum_j p(i, j) = 1$, which satisfies $p(i, j) = p(0, j - i) =: p(j - i)$ (homogeneity), $\sum_j j p(j) = 0$, and $p(j) = 0$ for all $|j| > v$ for some v (finiteness). Assume also that \mathcal{P} is truly d -dimensional: $\{j \in \mathbb{Z}^d : p(j) \neq 0\}$ generates \mathbb{Z}^d .

Let $\mathcal{E} = (\varepsilon, (\varepsilon_n(i), i \in \mathbb{Z}^d), n \in \mathbb{Z})$ be a family of i.i.d. integrable symmetric random variables. Let \mathbb{P} and \mathbb{E} denote the probability and expectation in the probability space generated by \mathcal{E} . (We use preliminary $n \in \mathbb{N}$ in the definitions but later it will be useful to have $n \in \mathbb{Z}$.)

The *serial harness* $(Y_n, n \geq 0)$ is the discrete-time Markov process in $\mathbb{R}^{\mathbb{Z}^d}$ defined by

$$Y_n(i) = \begin{cases} 0, & \text{if } n = 0, \\ \sum_{j \in \mathbb{Z}^d} p(i, j) Y_{n-1}(j) + \varepsilon_n(i), & \text{if } n \geq 1. \end{cases} \quad (1.1)$$

Here $Y_n(i)$ denotes the height of the serial harness at site i at time n . In other words, the evolution is given by

$$Y_n = \mathcal{P}Y_{n-1} + \varepsilon_n. \quad (1.2)$$

where $\varepsilon_n = (\varepsilon_n(i), i \in \mathbb{Z}^d)$. Since the “noise variable” ε is symmetric and thus has zero mean, we have that $\mathbb{E}Y_n(i) = 0$ for all i, n . We can interpret $p(i, j)$ as transition probabilities of a random walk on \mathbb{Z}^d ; let $p_m(i, j)$ be its m -step transition probabilities. By homogeneity, $p_m(i, j) = p_m(0, j - i) =: p_m(j - i)$. Iterating (1.1),

$$Y_n(i) = \sum_{r=1}^n \sum_{j \in \mathbb{Z}^d} p_{n-r}(i, j) \varepsilon_r(j) \stackrel{d}{=} \sum_{r=0}^{n-1} \sum_{j \in \mathbb{Z}^d} p_r(j) \varepsilon_r(j), \quad (1.3)$$

for all $n \geq 1, i \in \mathbb{Z}^d$, where $\stackrel{d}{=}$ means equidistributed. Hammersley (1967) obtained that

$$\mathbb{E}(Y_n(i))^2 = \sigma^2 s(n) \quad (1.4)$$

where σ^2 is the variance of ε and

$$s(n) := \sum_{r=0}^{n-1} \sum_{j \in \mathbb{Z}^d} p_r(j)^2. \quad (1.5)$$

is the expected number of encounters up to time n of two independent copies of a random walk starting at 0 with transition probabilities \mathcal{P} . For completeness we derive (1.4) in (6.11) later. Since $s(n) \sim \sqrt{n}$ for $d = 1$, $s(n) \sim \log n$ for $d = 2$ and $s(n)$ is uniformly bounded in n for $d \geq 3$ (see, for example, Spitzer (1976)), the surface delocalizes in dimensions $d \leq 2$ and stays localized in dimensions $d \geq 3$. Toom (1997) studies localization of the surface and surface-differences in function of the decay of the distribution of ε .

We consider the serial harness interacting by exclusion with a wall located at the origin. The *wall process* $(W_n, n \geq 0)$ is the Markov process in $(\mathbb{R}^+)^{\mathbb{Z}^d}$ defined by

$$W_n(i) = \begin{cases} 0, & \text{if } n = 0, \\ \left(\sum_{j \in \mathbb{Z}^d} p(i, j) W_{n-1}(j) + \varepsilon_n(i) \right)^+, & \text{if } n \geq 1, \end{cases} \quad (1.6)$$

for $i \in \mathbb{Z}^d$, where for $a \in \mathbb{R}$, $a^+ = a \vee 0 = \max(a, 0)$; this can be reexpressed as

$$W_n = (\mathcal{P}W_{n-1} + \varepsilon_n)^+. \quad (1.7)$$

We say that the law of a random surface Z is an *invariant measure* for the wall process if $Z \stackrel{d}{=} (\varepsilon_0 + \mathcal{P}Z)^+$, with ε_0 and Z independent. We show in Section 2 that

$$W_n \leq W_{n+1} \text{ stochastically.} \quad (1.8)$$

This implies that W_n is stochastically non-decreasing and thus their laws converge to a limit (that could give positive weight to infinity). If the limit is nondegenerate, then it is an invariant measure for the wall process. Monotonicity (1.8) implies in particular

$$\mu_n := \mathbb{E}W_n(0)$$

is nondecreasing and thus converges either to a finite limit or to ∞ . Our first result is general and rules out the former possibility, showing however that μ_n goes to infinity slower than n .

Theorem 1.1 (a) *There is no nondegenerate invariant measure for the wall process (W_n) ; (b) $W_n \rightarrow \infty$ in probability; (c) $\mu_n \rightarrow \infty$ as $n \rightarrow \infty$; (d) $\mu_n/n \rightarrow 0$ as $n \rightarrow \infty$.*

This theorem is proven in Section 2.

Let F be the law of ε , $\bar{F}(x) = \mathbb{P}(\varepsilon > x)$ and define

$$\mathcal{L}_\alpha^- := \{F : \bar{F}(x) \leq ce^{-c'x^\alpha}, x > 0, \text{ for some positive } c, c'\} \quad (1.9)$$

$$\mathcal{L}_\alpha^+ := \{F : \bar{F}(x) \geq ce^{-c'x^\alpha}, x > 0, \text{ for some positive } c, c'\} \quad (1.10)$$

and

$$\mathcal{L}_\alpha := \mathcal{L}_\alpha^- \cap \mathcal{L}_\alpha^+ \quad (1.11)$$

We next state our main result. It consists of upper and lower bounds for μ_n for different noise distributions.

Theorem 1.2 *There exist constants c and C that may depend on the dimension such that*

(i) *for $d = 1$ if $F \in \mathcal{L}_1^-$,*

$$cn^{1/4} \leq \mu_n \leq Cn^{1/4} \sqrt{\log n}; \quad (1.12)$$

(ii) for $d = 2$, if $F \in \mathcal{L}_\alpha$, for some $\alpha \geq 1$,

$$c(\log n)^{\frac{1}{\alpha}\vee\frac{1}{2}} \leq \mu_n \leq C \log n; \quad (1.13)$$

(iii) for $d \geq 3$, if $F \in \mathcal{L}_\alpha$, for some $1 \leq \alpha \neq 1 + d/2$,

$$c(\log n)^{\frac{1}{\alpha}} \leq \mu_n \leq C(\log n)^{\frac{1}{\alpha}\vee\frac{2}{2+d}}; \quad (1.14)$$

(iv) for $d \geq 3$ if $F \in \mathcal{L}_{1+d/2}$,

$$c(\log n)^{\frac{2}{2+d}} \leq \mu_n \leq C(\log n)^{\frac{2}{2+d}} (\log \log n)^{\frac{d}{2+d}}. \quad (1.15)$$

Our upper bound in (1.15) can be slightly improved, see (6.4) and Remark 6.2 below. The lower bound in (i) can be shown to hold under weaker conditions; that is also the case for some cases of (ii); see (6.7) and Remark 6.7 below. If the noise distribution is in \mathcal{L}_α for some $\alpha \geq 1$, then our lower and upper bounds to μ_n coincide to leading order in the case that $d \geq 3$, $1 \leq \alpha < 1 + d/2$ and also in the case that $d = 2$, $\alpha = 1$.

Theorems 1.1 and 1.2 catch the effect of the “entropic repulsion” in a stochastically moving surface interacting with a wall by exclusion.

Many papers deal with the problem of entropic repulsion in Equilibrium Statistical Mechanics. The role of the entropic repulsion in the Gaussian free field was studied by Lebowitz and Maes (1987), Bolthausen, Deuschel and Zeitouni (1995), Deuschel (1996), Deuschel and Giacomin (1999) and Bolthausen, Deuschel and Giacomin (2001). In the Ising, SOS and related models the matter was discussed in Bricmont, El Mellouki and Fröhlich (1986), Bricmont (1990), Holický and Zahradník (1993), Cesi and Martinelli (1996), Dinaburg and Mazel (1994) and Ferrari and Martínez (1998).

The exponent $1/4$ for dynamic entropic repulsion in $d = 1$ was predicted by Lipowsky (1985) using scaling arguments. This exponent was then found numerically by Mon, Binder, Landau (1987), Binder (1990), De Coninck, Dunlop and Menu (1993). Dunlop, Ferrari and Fontes (2001) proved bounds (slightly worse than) (1.12) for a one dimensional interface related to the phase separation line in the two dimensional Ising model at zero temperature. Funaki and Olla (2001) studied a one dimensional model in a finite box rescaled as the square of the time.

The strategy to show part of Theorem 1.2 is to compare the wall process with a “free process” — in our case the serial harness — as proposed by Dunlop, Ferrari and Fontes (2001). The following lemmas are the basic ingredients in this approach. The first two concern moderate deviations of the serial harness Y_n ; they are then extended to the wall process W_n in the last one.

Lemma 1.3 *If the distribution of ε is in \mathcal{L}_1^- , then in $d \leq 2$ there exist constants $k, c, c' > 0$ such that for all $K > 0$ and $0 \leq l \leq n$,*

$$\mathbb{P}[Y_l(0) \geq K \sqrt{s(n) \log n}] \leq kn^{c-c'K}. \quad (1.16)$$

Lemma 1.4 *If the distribution of ε is in \mathcal{L}_α^- for some $\alpha \geq 1$, then in $d \geq 3$ there exist constants $k, c, c' > 0$ such that, for all $K > 0$ and $0 \leq l \leq n$,*

(i) *if $\alpha \neq 1 + d/2$, then*

$$\mathbb{P}[Y_l(0) \geq K(\log n)^{\frac{1}{\alpha} \vee \frac{2}{2+d}}] \leq kn^{c-c'K}; \quad (1.17)$$

(ii) *and if $\alpha = 1 + d/2$, then*

$$\mathbb{P}[Y_l(0) \geq KL_n(1 + 2/d)] \leq kn^{c-c'K}, \quad (1.18)$$

where $L_n(\cdot)$ is defined in (6.1) below.

Lemma 1.5 *The bounds of Lemmas 1.4 and 1.3 hold for $l = n$ if we replace Y_n with W_n , possibly with worse constants k, c .*

We conclude this introduction with a remark concerning the form (1.6) of the interaction with the wall. Two other choices are also natural. First, if the noise would push the process below zero, simply do nothing. Or, in the same case, only take the convex combination without a noise. Formally, these two cases are, respectively

$$W'_0(i) = W''_0(i) \equiv 0,$$

and for $n \geq 1$

$$W'_n(i) = \begin{cases} \sum_{j \in \mathbb{Z}^d} p(i, j) W'_{n-1}(j) + \varepsilon_n(i), & \text{if this is positive,} \\ W'_{n-1}(i), & \text{otherwise;} \end{cases} \quad (1.19)$$

and

$$W''_n(i) = \begin{cases} \sum_{j \in \mathbb{Z}^d} p(i, j) W''_{n-1}(j) + \varepsilon_n(i), & \text{if this is positive,} \\ \sum_{j \in \mathbb{Z}^d} p(i, j) W''_{n-1}(j), & \text{otherwise.} \end{cases} \quad (1.20)$$

Coupling W, W', W'' by the same realization of the noise variables, one sees that, stochastically, both $W' \geq W$ and $W'' \geq W$. This implies immediately that any lower bound for μ_n (in particular the ones in this paper) hold for $\mu'_n := \mathbb{E}W'_n(0)$ and $\mu''_n := \mathbb{E}W''_n(0)$ as well. These dominations also imply immediately the validity of the results of Theorem 1.1 (a-c) for W' and W'' . For the analogue of Theorem 1.1 (d), domination does not help (it goes in the wrong direction). An argument along the same lines as the one for W can be made for W'' straightforwardly; see paragraphs containing (2.27) and (3.2). Under the assumption that $\mathcal{P}(0, 0) > 0$, one can also make a similar argument for W' ; otherwise, the matter is more delicate, and we do not have an argument.

As for upper bounds for μ'_n, μ''_n , the ones we get for μ_n also hold for both of them, since the proof only relies on the free process started at some height r dominating stochastically the wall process started at the same height, and this holds for all three choices.

2 Delocalization

In this section we show Theorem 1.1. The wall process is *attractive*, that is,

$$\text{if } W \leq W' \text{ then } (\mathcal{P}W + \varepsilon_0)^+ \leq (\mathcal{P}W' + \varepsilon_0)^+ \quad a.s. \quad (2.1)$$

coordinatewise, which implies

$$\text{if } W_n \leq W'_n \text{ stochastically, then } W_{n+1} \leq W'_{n+1} \text{ stochastically.} \quad (2.2)$$

and since for the process with initial flat surface $0 = W_0 \leq W_1$ *a.s.* this implies (1.8).

Theorem 1.1 is a consequence of the following three lemmas.

Lemma 2.1 *There is no invariant measure for (W_n) with finite mean.*

Proof We argue by contradiction. Suppose there exists an invariant measure ν_o with finite mean. We shall argue that in this case there is positive probability for the wall process started from equilibrium to interact with the wall, which yields a positive drift, contradicting stationarity.

Suppose first that the mean $m = \nu_o(W(0))$ is strictly contained in the support $I = [-c, c]$ of the distribution of ε (c may be infinity). That is,

$$\mathbb{P}(\varepsilon > m) > 0 \quad (2.3)$$

We claim that

$$\nu_o \left[\sum_j p(j)W(j) \leq m \right] > 0. \quad (2.4)$$

Otherwise,

$$\begin{aligned} \int \nu_o(dW) \mathbb{E}[W_1(0) | W_0 = W] &= \nu_o \mathbb{E} \left[\left(\sum_{j \in \mathbb{Z}^d} p(j)W(j) + \varepsilon_1(0) \right)^+ \right] \\ &\geq \nu_o \mathbb{E} \left[\sum_{j \in \mathbb{Z}^d} p(j)W(j) + \varepsilon_1(0) \right] \\ &= \nu_o \left[\sum_{j \in \mathbb{Z}^d} p(j)W(j) \right] > m = \nu_o(W(0)), \end{aligned} \quad (2.5)$$

contradicting stationarity. We have used that ν_o is stationary, $\varepsilon_1(0)$ is symmetric, and in the second inequality $\nu_o[\sum_j p(j)W(j) > m] = 1$ (the negation of (2.4)). Denoting $u := \sum_j p(j)W(j) + \varepsilon_1(0)$, we have

$$\int \nu_o(dW) \mathbb{E}[W_1(0) | W_0 = W] - \nu_o(W(0)) = \nu_o \mathbb{E} \left[\left(\sum_j p(j)W(j) + \varepsilon_1(0) \right)^+ \right] - m$$

$$\begin{aligned}
&= m + \nu_o \mathbb{E}[-u \mathbf{1}\{u \leq 0\}] - m \\
&\geq \nu_o \mathbb{E} \left[-u \mathbf{1} \left\{ \sum_j p(j) W(j) \leq m \right\} \mathbf{1}\{\varepsilon < -m\} \right],
\end{aligned} \tag{2.6}$$

By (2.4), the first indicator is 1 on a set of positive measure, and the same holds for the second one by (2.3) and the symmetry of ε . Since the measures ν_o and \mathbb{P} are independent, (2.6) implies that $\int \nu_o(dW) \mathbb{E}[W_1(0) | W_0 = W] - \nu_o(W(0)) > 0$ which contradicts stationarity.

Finally, suppose that $|c| < \infty$ and that the mean m of ν_o is not contained in the support $[-c, c]$ of ε . We show that the process started from equilibrium has positive probability of hitting the wall. Define $n^* = \lceil \frac{4m}{c} \rceil$. Under ν_o , the event $\mathcal{A} = \{W \in (\mathbb{R}^+)^{\mathbb{Z}^d} : \sum_j p_{n^*}(j) W(j) \leq 2m\}$ has probability larger than $1/2$. Indeed, by Markov's inequality,

$$\nu_o(\mathcal{A}^c) \leq \frac{1}{2m} \sum_{j \in \mathbb{Z}^d} p_{n^*}(j) \nu_o[W(j)] = \frac{1}{2}. \tag{2.7}$$

Suppose that $\varepsilon_n(j) < -\frac{c}{2}$, for all $0 \leq n \leq n^*$ and all j such that $|j| \leq (n^* - n)v$ (where $v = \max_i(p(0, i) > 0)$), then any configuration $W \in \mathcal{A}$ evolves in such a way that $W_{n^*}(0) = 0$. The above configuration of $\varepsilon_n(j)$'s has probability larger than

$$\left(\mathbb{P}[\varepsilon < c/2] \right)^{(n^*v)^d} > 0. \tag{2.8}$$

The bounds (2.7) and (2.8) imply that $\nu_o[W(0) = 0] = \int \nu_o(dW) \mathbb{P}[W_{n^*}(0) = 0 | W_0 = W] > 0$. With a similar argument one shows that

$$\nu_o[W(j) = 0, |j| \leq v] > 0. \tag{2.9}$$

Then we can bound (2.6) from below putting 0 instead of m :

$$\nu_o \mathbb{E} \left[-u \mathbf{1} \left\{ \sum_j p(j) W(j) = 0 \right\} \mathbf{1}\{\varepsilon < 0\} \right] > 0 \tag{2.10}$$

by (2.9) and the non-triviality of ε . As before this is in contradiction with stationarity. Hence, there is no invariant measure with finite mean. \blacksquare

Lemma 2.2 *Every invariant measure for (W_n) dominates stochastically $\lim_n \mathbb{P}(W_n \in \cdot)$.*

Proof Consider the family of processes $((W_n^k, n \geq k), k \in \mathbb{Z})$ defined by

$$W_n^k = \begin{cases} 0, & \text{if } n = k, \\ (\mathcal{P}W_{n-1}^k + \varepsilon_n)^+, & \text{if } n \geq k + 1, \end{cases} \tag{2.11}$$

$(W_n^k, n \geq k)$ is the wall process evolving from time k on, having flat configuration at initial time k . It is clear that for $k \geq 0$,

$$W_0^{-k} \stackrel{d}{=} W_k^0 (= W_k). \tag{2.12}$$

Since $0 = W_k^k \leq W_k^{k-1}$, by attractiveness (2.1), $W_n^k \leq W_n^{k-1}$ for all $n \geq k$, and in particular:

$$W_0^k \leq W_0^{k-1} \quad (2.13)$$

Immediately (2.12) and (2.13) imply (1.8), that is, W_n is stochastically non decreasing.

Suppose that there exists an invariant surface Z and define

$$Z_n^k = \begin{cases} Z, & \text{if } n = k, \\ (\mathcal{P}Z_{n-1}^k + \varepsilon_n)^+, & \text{if } n \geq k + 1, \end{cases} \quad (2.14)$$

Then, since $0 = W_k^k \leq Z_k^k = Z$, (2.1) implies

$$W_n^k \leq Z_n^k \stackrel{d}{=} Z \quad (2.15)$$

for all $k \leq n$. That is, Z dominates W_n^k stochastically for all $k \leq n$. Hence the limit

$$W_n^\infty := \lim_{k \rightarrow -\infty} W_n^k \quad (2.16)$$

is *a.s.* well defined for all n . It is invariant by construction: W_n^∞ is a deterministic function of $(\varepsilon_k, k \leq n)$ which is time-translation invariant: $W_n^\infty \stackrel{d}{=} W_{n-1}^\infty$ for all n . Furthermore, since $W_n^\infty = (\mathcal{P}W_{n-1}^\infty + \varepsilon_n)^+$, the distribution of W_n^∞ is invariant for the wall process. Hence, by Lemma 2.1, $\mathbb{E}(W_n^\infty) = \infty$. By (2.15), any invariant surface Z dominates stochastically W_n^∞ . By (2.12) and (2.16),

$$\text{the law of } W_0^\infty \text{ is } \lim_n \mathbb{P}(W_n \in \cdot). \quad (2.17)$$

This concludes the proof. ■

Lemma 2.3 W_0^∞ (and hence W_n^∞ for all n) is almost surely identically infinity.

Proof The event $\{W_0^\infty = \infty\}$ belongs to the tail σ -algebra of $\{\varepsilon_k : k \leq 0\}$, and is thus trivial. Write

$$W_0^\infty = (\varepsilon_0 + \mathcal{P}W_{-1}^\infty)^+ = \dots \quad (2.18)$$

$$= (\varepsilon_0 + \mathcal{P}(\varepsilon_{-1} + \dots \mathcal{P}(\varepsilon_{-k+1} + \mathcal{P}W_{-k}^\infty)^+ \dots)^+)^+ \quad (2.19)$$

$$\geq U_k + \mathcal{P}^k W_{-k}^\infty, \quad (2.20)$$

for $k > 0$, where $U_k = \sum_{i=0}^{k-1} \mathcal{P}^i \varepsilon_{-i}$. Notice that U_k is symmetric and that U_k and W_{-k}^∞ are independent: U_k is a function of $(\varepsilon_i : -k + 1 \leq i \leq 0)$ while W_{-k}^∞ is function of $(\varepsilon_i : i \leq -k)$. Since $W_{-k}^\infty \stackrel{d}{=} W_0^\infty$, for all $k \geq 0$

$$W_0^\infty \geq V_k + \mathcal{P}^k W_0^\infty, \text{ stochastically} \quad (2.21)$$

with $V_k \stackrel{d}{=} U_k$, V_k and W_0^∞ independent.

A key observation is that W_0^∞ is ergodic for spatial shifts. This follows from the fact that W_0^∞ is a function of $\varepsilon_n(i)$'s for a cone of indices (n, i) in $-\mathbb{N} \times \mathbb{Z}^d$ with vertex in $(0, x)$. Now, $\mathbb{E}(W_0^\infty) = \infty$, the Ergodic Theorem implies that $\mathcal{P}^k W_0^\infty \rightarrow \infty$ almost surely as $k \rightarrow \infty$. Indeed,

$$\mathcal{P}^k W_0^\infty(0) = \sum_{i \in \mathbb{Z}^d} p_k(i) W_0^\infty(i) \geq \frac{c}{k^{d/2}} \sum_{|i| \leq k} W_0^\infty(i) \rightarrow \infty, \quad (2.22)$$

as $k \rightarrow \infty$, by the Ergodic Theorem. We have used the positivity of W_0^∞ and the well known Local Central Limit Theorem estimate to the effect that $\inf_{|i| \leq k} p_k(i) \geq c/k^{d/2}$ for some $c > 0$. For this estimate, aperiodicity is required; we leave the necessary and straightforward adaptations for the periodic case to the reader.

Now, (2.21), (2.22) and the symmetry of V_k imply that for arbitrary $M > 0$

$$\mathbb{P}(W_0^\infty > M) \geq \liminf_{k \rightarrow \infty} \mathbb{P}(V_k + \mathcal{P}^k W_0^\infty > M) \quad (2.23)$$

$$\geq \liminf_{k \rightarrow \infty} \mathbb{P}(V_k \geq 0) \mathbb{P}(\mathcal{P}^k W_0^\infty > M) \quad (2.24)$$

$$\geq \frac{1}{2} \liminf_{k \rightarrow \infty} \mathbb{P}(\mathcal{P}^k W_0^\infty > M) = \frac{1}{2}. \quad (2.25)$$

Thus $\mathbb{P}(W_0^\infty = \infty) \geq 1/2$ and triviality implies $\mathbb{P}(W_0^\infty = \infty) = 1$. ■

Proof of Theorem 1.1 (a) is immediate consequence of Lemmas 2.2 and 2.3: any invariant surface dominates stochastically W_0^∞ and W_0^∞ is almost surely identically infinity.

Lemma 2.3 and (2.17) imply that $W_n(0)$ converges to infinity in probability. This shows (b).

For $n \geq 0$, $\mu_n = \mathbb{E}W_n(0) = \mathbb{E}W_0^{-n}$ (by (2.12)). Taking limits,

$$\lim_n \mu_n = \lim_n \mathbb{E}W_0^{-n} = \mathbb{E}(\lim_n W_0^{-n}) = \infty \quad (2.26)$$

by the monotone convergence theorem. This shows (c).

Finally, in (3.2) below it is shown that

$$\mu_n - \mu_{n-1} = \mathbb{E} \int_{\mathcal{P}W_{n-1}}^{\infty} \mathbb{P}(\varepsilon > x) dx \quad (2.27)$$

Since ε is integrable and $\mathcal{P}W_{n-1}$ increases to infinity in probability, (2.27) converges to zero, and we get (d). ■

3 A generic lower bound

From (1.7),

$$W_n(i) = (\mathcal{P}W_{n-1}(i) + \varepsilon_n(i))^+ = \mathcal{P}W_{n-1}(i) + \varepsilon_n(i) + (-\mathcal{P}W_{n-1}(i) - \varepsilon_n(i))^+ \quad (3.1)$$

Taking expectations, since ε is symmetric,

$$\mu_n = \mu_{n-1} + \mathbb{E} \int_{\mathcal{P}W_{n-1}} \mathbb{P}(\varepsilon > x) dx. \quad (3.2)$$

As $\int_y^\infty \mathbb{P}(\varepsilon > x) dx$ is a convex function of y ,

$$\mu_n \geq \mu_{n-1} + \int_{\mathbb{E}(\mathcal{P}W_{n-1})}^\infty \mathbb{P}(\varepsilon > x) dx = \mu_{n-1} + \mathbb{E}(\varepsilon - \mu_{n-1})^+. \quad (3.3)$$

For $s \geq 0$, let $G(s) = \mathbb{E}(\varepsilon - s)^+$, $H(s) = \mu + G(s)$, and $\nu(t)$ be such that $\int_0^{\nu(t)} [G(s)]^{-1} ds = t$.

Theorem 3.1 $\mu_n \geq \nu(n)$ for all $n \geq 0$.

Remark 3.2 This general lower bound does not depend on the dimension.

Corollary 3.3 If the distribution of ε belongs to \mathcal{L}_α^+ for some $\alpha > 0$, then there exists $c_2 = c_2(\alpha) > 0$ such that

$$\mu_n \geq c_2(\log n)^{\frac{1}{\alpha}}. \quad (3.4)$$

Corollary 3.4 Suppose that the distribution of ε decays at most polynomially, i.e. $\mathbb{P}(\varepsilon > x) \geq c_0 x^{-\alpha}$ for all $x > 1$ and some positive constants c_0 and $\alpha > 1$. Then there exists $c_1 = c_1(\alpha) > 0$ such that

$$\mu_n \geq c_1 n^{\frac{1}{\alpha}}. \quad (3.5)$$

Proof of Theorem 3.1. Notice first that $\nu(t)$ is a solution of

$$\nu(t) = \int_0^t G(\nu(s)) ds$$

and thus satisfies

$$\nu(n) = \nu(n-1) + \int_{n-1}^n G(\nu(s)) ds.$$

Notice also that $G(x)$ is decreasing and $H(x)$ is increasing. We prove the lemma by induction. First, $\mu_0 = \nu(0) = 0$. Suppose that $\mu_{n-1} \geq \nu(n-1)$. Then,

$$\begin{aligned} \nu(n) &= \nu(n-1) + \int_{n-1}^n G(\nu(s)) ds \leq \nu(n-1) + G(\nu(n-1)) \\ &= H(\nu(n-1)) \leq H(\mu_{n-1}) \leq \mu_n, \end{aligned}$$

where the last inequality is (3.3). ■

Proof of Corollary 3.4. Note that

$$G(x) = \mathbb{E}(\varepsilon - x)^+ = - \int_x^{+\infty} (y - x) d\mathbb{P}[\varepsilon \geq y] = \int_x^{+\infty} \mathbb{P}[\varepsilon \geq y] dy$$

and thus

$$g(s) := \int_0^t \frac{ds}{G(s)} = \int_0^t \frac{ds}{\int_s^{+\infty} \mathbb{P}[\varepsilon \geq y] dy}. \quad (3.6)$$

Thus, from the assumption in the statement of Corollary 3.4,

$$g(t) \leq \frac{1}{c_0} \int_0^t \frac{ds}{\frac{1}{\alpha-1} s^{1-\alpha}} = \frac{\alpha-1}{c_0 \alpha} t^\alpha \quad (3.7)$$

and

$$\nu(t) \geq c_1 t^{\frac{1}{\alpha}}$$

follows immediately. ■

Proof of Corollary 3.3. As above, we have

$$g(t) \leq \frac{1}{c} \int_0^t \frac{ds}{\int_s^{+\infty} e^{-c'y^\alpha} dy} \leq c_1 \int_0^t e^{c_2 s^\alpha} ds \leq c_3 e^{c_4 t^\alpha} \quad (3.8)$$

and the result follows. ■

4 Moderate deviations for the serial harness

The proofs of Lemmas 1.3 and 1.4 are based on the behavior of $\mathbb{E}(e^{lY_n(0)})$ for small and large l , established in Lemmas 4.1 and 4.3 below.

Lemma 4.1 *Let l_n be a sequence of positive numbers such that*

$$\bar{l}_n := l_n / \sqrt{s(n)} \leq 1. \quad (4.1)$$

Then there exists a constant c such that for all $0 \leq l \leq n$

$$\mathbb{E}[e^{\bar{l}_n Y_i(0)}] \leq e^{c l_n^2}. \quad (4.2)$$

Proof For all $0 \leq l \leq n$

$$\begin{aligned} \mathbb{E}\left[e^{\bar{l}_n Y_l(0)}\right] &= \prod_{r=0}^l \prod_{j \in \mathbb{Z}^d} \mathbb{E}\left[e^{\bar{l}_n p_r(j) \varepsilon}\right] \leq \prod_{r=0}^l \prod_{j \in \mathbb{Z}^d} e^{c \bar{l}_n^2 p_r(j)^2} \\ &= \exp\{c l_n^2 s(n)^{-1} s(l)\} \leq e^{c l_n^2}, \end{aligned}$$

where $c = \mathbb{E}(e^\varepsilon)$ and we have used that for a symmetric random variable W , if $|\lambda| \leq 1$, then

$$\mathbb{E}(e^{\lambda W}) \leq 1 + \mathbb{E}(e^W) \lambda^2 \leq e^{\mathbb{E}(e^W) \lambda^2} \quad (4.3)$$

and the fact that $s(\cdot)$ is nondecreasing. \blacksquare

Proof of Lemma 1.3.

$$\mathbb{P}[Y_l(0) \geq K \sqrt{s(n) \log n}] = \mathbb{P}[\bar{l}_n Y_n(0) \geq \log n^{c'K}] \leq n^{-c'K} \mathbb{E}[e^{\bar{l}_n Y_n(0)}],$$

where $l_n = c'' \sqrt{\log n}$, for an appropriate constant c'' , and Lemma 4.1 yields the result. \blacksquare

For the proof of Lemma 1.4, we will use that in $d \geq 3$

$$s := \lim_{n \rightarrow \infty} s(n) < \infty. \quad (4.4)$$

We will also need the following converse of (4.3).

Lemma 4.2 *If the distribution of W is in \mathcal{L}_α^- for some $\alpha > 1$, then there exists a constant c such that*

$$\mathbb{E}(e^{\lambda W}) \leq e^{c \lambda^\beta}, \quad (4.5)$$

for all $l \geq 1$, where $\beta = \alpha/(\alpha - 1)$.

Proof We have that

$$\mathbb{E}e^{\lambda W} \leq 1 + c \int_0^\infty e^{lx} e^{-c'x^\alpha} dx = 1 + c_1 \int_0^\infty e^{\tilde{\lambda}x} e^{-x^\alpha} dx, \quad (4.6)$$

where $\tilde{\lambda} = l/c^{1/\alpha}$. Now, we write the integral in (4.6) as

$$\int_0^{(2\tilde{\lambda})^{\beta-1}} e^{\tilde{\lambda}x} dx + \int_{(2\tilde{\lambda})^{\beta-1}}^\infty e^{\tilde{\lambda}x-x^\alpha} dx.$$

The former integral is bounded above by $e^{c''\lambda^\beta}$. The latter one is bounded above by a uniform constant. \blacksquare

Lemma 4.3 *In $d \geq 3$, if the distribution of ε is in \mathcal{L}_α^- for some $\alpha > 1$, then there exist a constant c such that for all large q*

$$\mathbb{E}(e^{qY_n(0)}) \leq \begin{cases} e^{cq^{\beta \vee (1+2/d)}}, & \text{if } \alpha \neq 1 + d/2; \\ e^{cq^{1+2/d} \log q}, & \text{if } \alpha = 1 + d/2, \end{cases} \quad (4.7)$$

where $\beta = \alpha/(\alpha - 1)$ as before.

Proof

$$\begin{aligned} \mathbb{E}(e^{qY_n(0)}) &\leq \prod_{k=0}^{\infty} \prod_{x \in \mathbb{Z}^d} \mathbb{E}(e^{qp_k(x)\varepsilon}) \leq \prod_{k,x:qp_k(x)>1} e^{c(qp_k(x))^\beta} \prod_{k,x:qp_k(x)\leq 1} e^{c(qp_k(x))^2} \\ &= \exp \left\{ c \left[\sum_{k,x:qp_k(x)>1} (qp_k(x))^\beta + \sum_{k,x:qp_k(x)\leq 1} (qp_k(x))^2 \right] \right\}. \end{aligned} \quad (4.8)$$

We now estimate the expression within square brackets in (4.8). If $\beta \geq 2$ or, equivalently, $1 < \alpha \leq 2$, then that expression is bounded above by

$$q^\beta \sum_{k,x} p_k^2(x) = q^\beta s. \quad (4.9)$$

For the case $1 < \beta < 2$ (equivalently, $\alpha > 2$), we use the well known estimate on $p_k := \sup_{x \in \mathbb{Z}^d} p_k(x)$: there exists a constant C such that for all $k \geq 1$

$$p_k \leq Ck^{-d/2} \quad (4.10)$$

(see e.g. Spitzer (1976)) to conclude that the expression within square brackets in (4.8) is bounded above by

$$q^\beta \sum_{k=0}^{(Cq)^{2/d}} p_k^{\beta-1} + q^2 \sum_{k=(Cq)^{2/d}}^{\infty} p_k \leq C' q^\beta \sum_{k=1}^{(Cq)^{2/d}} k^{-d(\beta-1)/2} + C'' q^{1+2/d} \quad (4.11)$$

for some constants C', C'' . The result follows. \blacksquare

Proof of Lemma 1.4. Let Q_n be a sequence of positive numbers such that $Q_n = o(\log n)$ and $q_n = (\log n)/Q_n$. Then

$$\mathbb{P}[Y_l(0) \geq KQ_n] \leq \mathbb{P}[q_n Y_l(0) \geq K(\log n)] \leq n^{-K} \mathbb{E}(e^{q_n Y_l(0)}). \quad (4.12)$$

We can thus use Lemma 4.3 for q_n . Therefore, if $1 < \alpha \neq 1 + d/2$, making $Q_n = (\log n)^{\frac{1}{\alpha} \vee \frac{2}{2+d}}$, we have $q_n = (\log n)^{1 - (\frac{1}{\alpha} \vee \frac{2}{2+d})} = (\log n)^{\frac{1}{\beta} \wedge \frac{d}{d+2}}$ and thus, from (4.7)

$$\mathbb{P}[Y_l(0) \geq K(\log n)^{\frac{1}{\alpha} \vee \frac{2}{2+d}}] \leq n^{c-K}. \quad (4.13)$$

If $\alpha = 1 + d/2$, we make $Q_n = L_n(1 + 2/d)$, and thus $q_n = (\log n)/L_n(1 + 2/d) = \ell_n(1 + 2/d)$. From (4.7) and the definition of $\ell_n(1 + 2/d)$ (above (6.1) below)

$$\mathbb{P}[Y_l(0) \geq KL_n(1 + 2/d)] \leq n^{c-K}. \quad (4.14)$$

For $\alpha = 1$, we have

$$\mathbb{E}e^{Y_n(0)} = \prod_{k,x} \mathbb{E}e^{p_k(x)\varepsilon} \leq e^{c \sum_{k,x} p_k^2(x)} = e^{cs}, \quad (4.15)$$

where we have used (4.3). Thus, we obtain that

$$\mathbb{P}[Y_n(0) > K \log n] \leq Cn^{-K}.$$

■

5 Moderate deviations for the wall process

In this section we show Lemma 1.5. Introduce new processes $W_n^{0,r}$ and $Y_n^{0,r}$, which have the same evolution as W_n , respectively Y_n , but are started at time zero at height $r \in \mathbb{N}$. That is, $W_0^{0,r}(i) = Y_0^{0,r}(i) = r$, for all $i \in \mathbb{Z}^d$.

Let

$$a_n = \begin{cases} 2K(\log n)^{\frac{1}{\alpha} \vee \frac{2}{2+d}}, & \text{for the extension of (1.17);} \\ 2KL_n(1 + 2/d), & \text{for the extension of (1.18);} \\ 2K\sqrt{s(n) \log n}, & \text{for the extension of (1.16).} \end{cases} \quad (5.1)$$

Then,

$$\begin{aligned} \mathbb{P}[W_n(0) \geq a_n] &\leq \mathbb{P}[W_n^{0,r}(0) \geq a_n] \\ &= \mathbb{P}[W_n^{0,r}(0) \geq a_n, W_n^{0,r}(0) = Y_n^{0,r}(0)] + \mathbb{P}[W_n^{0,r}(0) \geq a_n, W_n^{0,r}(0) \neq Y_n^{0,r}(0)] \\ &\leq \mathbb{P}[Y_n^{0,r}(0) \geq a_n] \end{aligned} \quad (5.2)$$

$$+ \mathbb{P}[W_n^{0,r}(0) \neq Y_n^{0,r}(0)] \quad (5.3)$$

To get a bound for the probability in (5.2) of the form (1.16-1.18), we take $r = a_n/2$ and use (1.16-1.18).

The probability in (5.3) is treated as follows. Note that $W_n^{0,r}(0)$ and $Y_n^{0,r}(0)$ differ if a discrepancy occurs in the cone (v is the maximal speed of a discrepancy)

$$\{(l, j) \in \mathbb{N}_0 \times \mathbb{Z}^d : l \leq n, \quad |j| \leq v(n-l)\}, \quad (5.4)$$

that is,

$$\{Y_n^{0,r}(0) \neq W_n^{0,r}(0)\} = \{Y_l^{0,r}(j) < 0 \text{ for some } (l, j) \text{ with } l \leq n, |j| \leq v(n-l)\}.$$

Since $Y_n^{0,r}(0)$ has the same law as $Y_n(0) + r$ and by symmetry, we have

$$\mathbb{P}[Y_l^{0,r}(j) < 0] = \mathbb{P}[Y_l(j) < -r] = \mathbb{P}[Y_l(j) > r]. \quad (5.5)$$

Hence,

$$\begin{aligned} \mathbb{P}[Y_n^{0,r}(0) \leq W_n^{0,r}(0)] &= \mathbb{P}[Y_l(j) > r \text{ for some } (l, j) \text{ with } l \leq n, |j| \leq v(n-l)] \\ &\leq \sum_{l=0}^n \sum_{|j| \leq v(n-l)} \mathbb{P}[Y_l(j) > r]. \end{aligned}$$

Taking $r = a_n/2$ as before and using (1.16-1.18), we obtain

$$\mathbb{P}[Y_n^{0,r}(0) \neq W_n^{0,r}(0)] \leq kn^{c-c'K} \sum_{l=0}^n \sum_{|j| \leq v(n-l)} 1 \leq k'n^{c''-c'K}, \quad (5.6)$$

for some k', c'' . ■

6 Bounds for the wall process

For $\gamma > 1$, define $\ell_n(\gamma)$ as the solution of $x^\gamma \log x = \log n$, and let

$$L_n(\gamma) = (\log n)/\ell_n(\gamma). \quad (6.1)$$

Note that

$$(\log n)^{1-\frac{1}{\gamma}} \leq L_n(\gamma) \leq (\log n)^{1-\frac{1}{\gamma}} (\log \log n)^{\frac{1}{\gamma}} \text{ for all } n. \quad (6.2)$$

Theorem 6.1 *Suppose that the distribution of ε belongs to \mathcal{L}_α^- for some $\alpha \geq 1$. If $d \geq 3$, then there exists $c_3 = c_3(\alpha, d) > 0$ such that*

(i) *if $1 \leq \alpha \neq 1 + \frac{d}{2}$, then*

$$\mu_n \leq c_3 (\log n)^{\frac{1}{\alpha} \vee \frac{2}{2+d}}; \quad (6.3)$$

(ii) *if $\alpha = 1 + \frac{d}{2}$, then for all $\delta > 0$ we have*

$$\mu_n \leq c_3 L_n(1 + 2/d); \quad (6.4)$$

If $d = 2$, then there exists c_3 such that

$$\mu_n \leq c_3 \log n. \quad (6.5)$$

Remark 6.2 From (6.4) and (6.2), a slightly weaker alternative to (6.4) is

$$\mu_n \leq c_3 (\log n)^{\frac{2}{2+d}} (\log \log n)^{\frac{d}{2+d}}. \quad (6.6)$$

We now restrict attention to the class of exponentially decaying noise distributions. When the noise distribution is in \mathcal{L}_α , $\alpha \geq 1$, the results in Corollary 3.3 and Theorem 6.1 are our best explicit bounds (to leading order) for $d \geq 3$ and $d = 2$, $1 \leq \alpha \leq 2$. For $d = 1$, $\alpha \geq 1$ and $d = 2$, $\alpha > 2$, we have better bounds, which we discuss now.

Theorem 6.3 *If the distribution of ε is in \mathcal{L}_1^- , then for $d \leq 2$, there exist constants $c, C > 0$ such that*

$$c\sqrt{s(n)} \leq \mu_n \leq C\sqrt{s(n) \log n}, \quad (6.7)$$

where $s(n)$ is defined in (1.5). In particular

(i) for $d = 1$,

$$cn^{1/4} \leq \mu_n \leq Cn^{1/4} \sqrt{\log n}; \quad (6.8)$$

(ii) and for $d = 2$,

$$c\sqrt{\log n} \leq \mu_n \leq C \log n \quad (6.9)$$

Remark 6.4 The lower bound in (6.7) actually holds under the weaker assumption that $\mathbb{E}(\varepsilon^2) < \infty$. See Remark 6.7 below.

We prove first the lower bound (6.7). The first step is to calculate the variance of the serial harness, which will give us the proper scaling. From (1.3) we get (this is already contained in Hammersley (1967))

$$\mathbb{E} Y_n(0) = \sum_{r=0}^{n-1} \sum_{j \in \mathbb{Z}^d} p_r(j) \mathbb{E} \varepsilon = 0, \quad (6.10)$$

and

$$\begin{aligned} \mathbb{E} Y_n(0)^2 &= \sum_{r,r'=0}^{n-1} \sum_{j,j' \in \mathbb{Z}^d} p_r(j) p_{r'}(j') \mathbb{E} [\varepsilon_r(j) \varepsilon_{r'}(j')] \\ &= \sigma^2 \sum_{r=0}^{n-1} \sum_{j \in \mathbb{Z}^d} p_r(j)^2 = \sigma^2 s(n). \end{aligned} \quad (6.11)$$

The correct scaling for the serial harness is therefore $s(n)^{1/2}$, and we define accordingly

$$\tilde{Y}_n(0) \equiv s(n)^{-\frac{1}{2}} Y_n(0). \quad (6.12)$$

Analogously we define $\tilde{W}_n(0)$ for the wall process. We now show that $\tilde{Y}_n(0)$ is uniformly integrable (with respect to n).

Lemma 6.5 *The process $(\tilde{Y}_n(0))_n$ satisfies $\sup_n \mathbb{E}(e^{|\tilde{Y}_n(0)|}) < \infty$.*

Proof By symmetry of the ε , $\mathbb{E}(e^{|\tilde{Y}_n(0)|}) \leq 2\mathbb{E}(e^{\tilde{Y}_n(0)}) \leq 2e^c$, where the last inequality follows from Lemma 4.1 with $l_n \equiv 1$. \blacksquare

From Lemma (6.5) it follows immediately that $s(n)^{-1}Y_n(0)^2$ is uniformly integrable.

Lemma 6.6 *There exists a constant $c > 0$ such that for all n*

$$\mathbb{E}|\tilde{Y}_n(0)| > c. \quad (6.13)$$

Proof Clearly, for any positive M ,

$$\begin{aligned} \mathbb{E}[\tilde{Y}_n(0)^2] &= \mathbb{E} \left[\tilde{Y}_n(0)^2 \mathbf{1}\{|\tilde{Y}_n(0)| > M\} \right] + \mathbb{E} \left[\tilde{Y}_n(0)^2 \mathbf{1}\{|\tilde{Y}_n(0)| \leq M\} \right] \\ &\leq \mathbb{E} \left[\tilde{Y}_n(0)^2 \mathbf{1}\{|\tilde{Y}_n(0)| > M\} \right] + M\mathbb{E}[|\tilde{Y}_n(0)|]. \end{aligned} \quad (6.14)$$

Since $\tilde{Y}_n(0)^2$ is uniformly integrable, for each $\delta > 0$ we can choose $M > 0$ such that

$$\mathbb{E} \left[\tilde{Y}_n(0)^2 \mathbf{1}\{|\tilde{Y}_n(0)| > M\} \right] < \delta, \quad (6.15)$$

uniformly in n . Thus

$$\mathbb{E}[|\tilde{Y}_n(0)|] \geq \frac{\mathbb{E}[\tilde{Y}_n(0)^2] - \delta}{M} = \frac{\sigma^2 - \delta}{M} > c > 0, \quad (6.16)$$

for some $\delta > 0$. \blacksquare

We finally prove the result about the wall process by coupling it with the serial harness using the same disorder variables \mathcal{E} . By symmetry,

$$\mathbb{E}[|\tilde{Y}_n(0)|] = \mathbb{E}[(\tilde{Y}_n(0))^+] + \mathbb{E}[(-\tilde{Y}_n(0))^+] = 2\mathbb{E}[(\tilde{Y}_n(0))^+]. \quad (6.17)$$

On the other hand, by construction, $\tilde{W}_n(0) \geq (\tilde{Y}_n(0))^+$, and therefore,

$$\mathbb{E}[\tilde{W}_n(0)] \geq \mathbb{E}[(\tilde{Y}_n(0))^+] \geq \frac{1}{2}\mathbb{E}[|\tilde{Y}_n(0)|] \geq c' > 0. \quad (6.18)$$

This proves the lower bound (6.7).

The upper bounds (6.3-6.5) and (6.9) follow from Lemma 1.5 in the same, following way. Let a_n be as in (5.1) and $b_n = a_n/(2K)$. Then

$$\begin{aligned} \mu_n/b_n &= \mathbb{E}[W_n(0)/b_n] = \int_0^\infty \mathbb{P}(W_n(0) > Kb_n) dK \\ &\leq c/c' + k \int_{c/c'}^\infty n^{c-c'K} dK \leq C, \end{aligned}$$

for some constant C . \blacksquare

Remark 6.7 The lower bound in (6.7) actually holds under the weaker assumption that $\mathbb{E}(\varepsilon^2) < \infty$, since this is enough to have $\tilde{Y}_n(0)^2$ uniformly integrable.

Acknowledgments.

We thank Servet Martínez for many discussions on harnesses, in particular for pointing out the minimality of W_0^∞ . We thank François Dunlop for references to the physical literature.

This paper is supported by Fundação de Apoio à Pesquisa do Estado de São Paulo (FAPESP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Programa Núcleos de Excelência (PRONEX).

References

- [1] K. Binder (1990) Growth kinetics of wetting layers at surfaces. pp 31–44 in *Kinetics of Ordering and Growth at Surfaces*. Edited by M.G. Lagally. Plenum Press, New-York.
- [2] E. Bolthausen, J.-D. Deuschel, G. Giacomin (2001) Entropic repulsion and the maximum of the two-dimensional harmonic crystal. *Ann. Probab.* **29** (4) 1670–1692.
- [3] E. Bolthausen, J.-D. Deuschel, O. Zeitouni (1995) Entropic repulsion of the lattice free field. *Comm. Math. Phys.* **170** (2), 417–443.
- [4] J. Bricmont (1990) Random surfaces in statistical mechanics in Wetting phenomena. *Proceedings of the Second Workshop held at the University of Mons, Mons, October 17–19, 1988*. Edited by J. De Coninck and F. Dunlop. *Lecture Notes in Physics* **354**. Springer-Verlag, Berlin.
- [5] J. Bricmont, A. El Mellouki, J. Fröhlich (1986) Random surfaces in Statistical Mechanics - roughening, rounding, wetting *J. Stat. Phys.* **42** (5/6), 743–798
- [6] F. Cesi, F. Martinelli (1996) On the layering transition of an SOS surface interacting with a wall. I. Equilibrium results. *J. Stat. Phys.* **82** (3/4), 823–913.
- [7] J. De Coninck, F. Dunlop, F. Menu (1993) Spreading of a Solid-On-Solid drop. *Phys. Rev. E* **47**(3), 1820–1823.
- [8] J.-D. Deuschel (1996) Entropic repulsion of the lattice free field. II. The 0-boundary case. *Comm. Math. Phys.* **181** (3), 647–665.
- [9] F. M. Dunlop, P. A. Ferrari, L. R. G. Fontes (2002) A dynamic one-dimensional interface interacting with a wall. *J. Statist. Phys.* **107** (3-4), 705–727.
- [10] J.-D. Deuschel, G. Giacomin (1999) Entropic repulsion for the free field: pathwise characterization in $d \geq 3$. *Comm. Math. Phys.* **206** (2), 447–462.
- [11] E. Dinaburg, A.E. Mazel (1994) Layering transition in SOS model with external magnetic-field. *J. Stat. Phys.* **74** (3/4), 533–563.

- [12] P. A. Ferrari, S. Martínez (1998) Hamiltonians on random walk trajectories. *Stochastic Process. Appl.* **78** (1), 47–68.
- [13] J. M. Hammersley (1965/66) Harnesses. *Proc. Fifth Berkeley Sympos. Mathematical Statistics and Probability*, Vol. III, 89–117.
- [14] P. Holický, M. Zahradník (1993) On entropic repulsion in low temperature Ising models. *Cellular automata and cooperative systems* (Les Houches, 1992), 275–287, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 396, Kluwer Acad. Publ., Dordrecht.
- [15] J.L. Lebowitz, C. Maes (1987) The effect of an external field on an interface, entropic repulsion. *J. Stat. Phys.* **46** (1/2), 39–49.
- [16] R. Lipowsky (1985) Nonlinear growth of wetting layers. *J. Phys. A* **18**, L585–L590.
- [17] K. K. Mon, K. Binder, and D. P. Landau (1987) Monte Carlo simulation of the growth of wetting layers *Phys. Rev. B* **35** (7), 3683–3685.
- [18] F. Spitzer (1976) *Principles of random walk*. Springer-Verlag, New York.
- [19] Toom, A. (1997) Tails in harnesses. *J. Statist. Phys.* **88** (1-2) 347–364.