

THE ASYMPTOTIC BEHAVIOUR OF HEEGAARD GENUS

MARC LACKENBY

1. INTRODUCTION

Heegaard splittings have recently been shown to be related to a number of important conjectures in 3-manifold theory: the virtually Haken conjecture, the positive virtual b_1 conjecture and the virtually fibred conjecture [3]. Of particular importance is the rate at which the Heegaard genus of finite-sheeted covering spaces grows as a function of their degree. This was encoded in the following definitions.

Let M be a compact orientable 3-manifold. Let $\chi_-^h(M)$ be the negative of the maximal Euler characteristic of a Heegaard surface for M . Let $\chi_-^{sh}(M)$ be the negative of the maximal Euler characteristic of a strongly irreducible Heegaard surface for M , or infinity if such a surface does not exist. Define the *infimal Heegaard gradient* of M to be

$$\inf_i \left\{ \frac{\chi_-^h(M_i)}{d_i} : M_i \text{ is a degree } d_i \text{ cover of } M \right\}.$$

The *infimal strong Heegaard gradient* of M is

$$\liminf_i \left\{ \frac{\chi_-^{sh}(M_i)}{d_i} : M_i \text{ is a degree } d_i \text{ cover of } M \right\}.$$

The following conjectures were put forward in [3]. According to Theorem 1.7 of [3], either of these conjectures, together with a conjecture of Lubotzky and Sarnak [4] about the failure of Property (τ) for hyperbolic 3-manifolds, would imply the virtually Haken conjecture for hyperbolic 3-manifolds.

Heegaard gradient conjecture. *A compact orientable hyperbolic 3-manifold has zero infimal Heegaard gradient if and only if it virtually fibres over the circle.*

Strong Heegaard gradient conjecture. *Any closed orientable hyperbolic 3-manifold has positive infimal strong Heegaard gradient.*

Some evidence for these conjectures was presented in [3]. More precisely, suitably phrased versions of these conjectures were shown to be true when one

restricts attention to cyclic covers dual to a non-trivial element of $H_2(M, \partial M)$, to reducible manifolds and (in the case of the strong Heegaard gradient conjecture) to congruence covers of arithmetic hyperbolic 3-manifolds.

A less quantitative version of the conjectures is simply that $\chi_-^{sh}(M_i)$ cannot grow too slowly as a function of d_i , and that if $\chi_-^h(M_i)$ does grow sufficiently slowly, then M is virtually fibred. These expectations are confirmed in the following result, which is the main theorem of this paper.

Main Theorem. *Let M be a closed orientable 3-manifold that admits a negatively curved Riemannian metric. Let $\{M_i \rightarrow M\}$ be a collection of finite regular covers with degree d_i .*

- (1) *If $\chi_-^h(M_i)/\sqrt{d_i} \rightarrow 0$, then $b_1(M_i) > 0$ for all sufficiently large i .*
- (2) *$\chi_-^{sh}(M_i)/\sqrt{d_i}$ is bounded away from zero.*
- (3) *If $\chi_-^h(M_i)/\sqrt[3]{d_i} \rightarrow 0$, then M_i fibres over the circle for all sufficiently large i .*

A slightly weaker form of Theorem (1) appeared in [3] as Corollary 1.4, with essentially the same proof. It is included here in order to emphasise its connection to the other two results. In fact, Corollary 1.4 of [3] followed from a stronger result, Theorem 1.3 of [3], which dealt with possibly irregular covers and gave a characterisation of having positive virtual b_1 in terms of generalised Heegaard splittings. Analogously, one can construct versions of Theorems (2) and (3) above for irregular covers, in the spirit of Theorems 1.1 and 1.3 of [3]. However, we will not do so, since they are a little technical. We will content ourselves with the following corollary of the Main Theorem, which gives a necessary and sufficient condition for M to be virtually fibred in terms of the Heegaard genus of its finite covers.

Corollary. *Let M be a closed orientable 3-manifold with a negatively curved Riemannian metric. Let $\{M_i \rightarrow M\}$ be its finite-sheeted covers, and let $N(\pi_1 M_i)$ be the normaliser of $\pi_1 M_i$ in $\pi_1 M$. Then the following are equivalent:*

- (1) *M_i is fibred for infinitely many i ;*
- (2) *$\chi_-^h(M_i)[\pi_1 M : N(\pi_1 M_i)]$ has a bounded subsequence.*

Proof. (1) \Rightarrow (2). If some finite-sheeted cover \tilde{M} is fibred, then so is any finite

cyclic cover M_i of \tilde{M} dual to the fibre. The normaliser $N(\pi_1 M_i)$ contains $\pi_1 \tilde{M}$, and so $[\pi_1 M : N(\pi_1 M_i)]$ is bounded. Also, $\chi_-^h(M_i)$ is bounded, and hence, so is their product.

(2) \Rightarrow (1). Since $\chi_-^h(M_i)[\pi_1 M : N(\pi_1 M_i)]$ has a bounded subsequence, we may pass to a subsequence where $\chi_-^h(M_i)$ is bounded and $N(\pi_1 M_i)$ is constant. Let \tilde{M} be the finite-sheeted cover of M corresponding to this subgroup. Then, $\{M_i \rightarrow \tilde{M}\}$ is a collection of finite regular covers with bounded $\chi_-^h(M_i)$. Hence, by Theorem (3), M_i is fibred for all sufficiently large i . \square

2. THE PROOF

We start with a closed orientable 3-manifold M admitting a negatively curved Riemannian metric. We may assume that it is bumpy in the sense of [7]. Let $\kappa < 0$ be the supremum of its sectional curvatures. Pick a 1-vertex triangulation T of M . The edges of T , when oriented in some way, form a set \mathcal{X} of generators for $\pi_1(M)$. Let K be the 2-skeleton of the complex dual to T .

We will consider a collection $\{M_i \rightarrow M\}$ of finite regular covers of M , having the properties of the Main Theorem. In particular, we will assume (at least) that $\chi_-^h(M_i)/\sqrt{d_i} \rightarrow 0$. The triangulation T and 2-complex K lift to T_i and K_i , say, in M_i . The 1-skeleton of T_i forms the Cayley graph X_i of $G_i = \pi_1 M / \pi_1 M_i$ with respect to \mathcal{X} .

Let x be $\chi_-^h(M_i)$ when proving Theorems (1) and (3), and let x be $\chi_-^{sh}(M_i)$ when proving Theorem (2). In the proof of Theorem 3.1 in [3], the following inequality can be found:

$$h(M_i) \leq \frac{4\pi}{|\kappa|} \frac{x}{\text{Volume}(M_i)} = \frac{4\pi}{|\kappa| \text{Volume}(M)} \frac{x}{d_i},$$

where $h(M_i)$ is the Cheeger constant of M_i . By the argument of Lemma 2 in [1], there is a constant $k_1 > 0$ independent of i such that the Cheeger constant $h(X_i)$ of the graph X_i satisfies $h(X_i) \leq k_1 h(M_i)$. Setting

$$k_2 = \frac{4\pi k_1}{|\kappa| \text{Volume}(M)},$$

we deduce that

$$h(X_i) \leq k_2 x / d_i.$$

Let $V(X_i)$ be the vertex set of X_i . Let A be a non-empty subset of $V(X_i)$ such that $|\partial A|/|A| = h(X_i)$ and $|A| \leq |V(X_i)|/2 = d_i/2$. Lemma 2.4 in [3] gives that, when $h(X_i) \leq \sqrt{2/3d_i}$, T_i admits a 1-cocycle c that is not a coboundary. Since $h(X_i) \leq k_2x/d_i$, and we are assuming (at least) that $x/\sqrt{d_i} \rightarrow 0$, then such a cocycle exists for all sufficiently large i . This establishes (1) of the Main Theorem. From the proof of Lemma 2.4 in [3], c takes values in $\{-1, 0, 1\}$, and its support is a subset of the edges of ∂A . Dual to this cocycle is a transversely oriented normal surface S in T_i which is homologically non-trivial. Remove any 2-sphere components from S . This is still homologically non-trivial, since all 2-spheres in M_i are inessential, as M_i is negatively curved. The intersection of S with the 2-skeleton of T_i is a graph in S whose complementary regions are triangles and squares. Let $V(S)$ and $E(S)$ be its vertices and edges. Its vertices are in one-one correspondence with the edges of T_i in the support of c . So,

$$|V(S)| \leq |\partial A| = |A|h(X_i) \leq d_i h(X_i)/2 \leq k_2x/2.$$

The valence of each vertex is at most the maximal valence of an edge in T , k_3 , say. So,

$$|\chi(S)| < |E(S)| \leq |V(S)|k_3/2 \leq k_2k_3x/4.$$

Setting $k_4 = k_2k_3/4$, we have deduced the existence of a homologically non-trivial transversely oriented properly embedded surface S with $|\chi(S)| \leq k_4x$ and with no 2-sphere components. By compressing S and removing components if necessary, we may assume that S is also incompressible and connected. By a theorem of Freedman, Hass and Scott [2], there is an ambient isotopy taking S to a minimal surface.

Set $\epsilon/2$ to be the injectivity radius of M . This is then a lower bound for the injectivity radius of M_i . Set

$$k_5 = \epsilon + \epsilon/2\pi + 1/2|\kappa|^{1/2}.$$

Let \tilde{K} be the lift of the 2-complex K to the universal cover of M . Let k_6 be the maximum number of complementary regions of \tilde{K} within a distance k_5 of any point. Let

$$k_7 = 1/(\cosh(|\kappa|^{1/2}\epsilon/2) - 1) + 3/2,$$

$$k_8 = k_6k_7.$$

Lemma 1. *Let S be a minimal surface in M_i . Then S intersects at most $k_8|\chi(S)|$ complementary regions of K_i . Hence, running through any such region, there are at most $k_8|\chi(S)|$ translates of S under the action of G_i .*

Proof. In the proof of Proposition 5.1 in [3], it is shown that the number of k_5 -balls required to cover S is at most $k_7|\chi(S)|$. The centre of each of these balls has at most k_6 complementary regions of K_i within a distance k_5 . So, S intersects at most $k_8|\chi(S)|$ complementary regions of K_i . Each such region corresponds to an element of G_i . To prove the second half of the lemma, we may concentrate on the region corresponding to the identity. Then a translate Sg runs through here, for some g in G_i , if and only if S runs through the region corresponding to g^{-1} . Thus, there can be at most $k_8|\chi(S)|$ such g . \square

Proof of Theorem (2). Let F be a strongly irreducible Heegaard surface in M_i with $|\chi(F)| = \chi_-^{sh}(M_i) = x$. By a theorem of Pitts and Rubinstein [5], there is an ambient isotopy taking it either to a minimal surface or to the double cover of a minimal non-orientable surface, with a small tube attached. So, by Lemma 1, F intersects at most k_8x complementary regions of K_i . Hence, the number of copies of S that F intersects is at most $(k_8x)(k_8k_4x)$. This is less than d_i if $\chi_-^{sh}(M_i)/\sqrt{d_i}$ is sufficiently small. So there is a translate of S which misses F . It then lies in a complementary handlebody of F . But this is impossible, since S is incompressible. So, $\chi_-^{sh}(M_i)/\sqrt{d_i}$ is bounded away from zero. \square

Proof of Theorem (3). Let $\{F_1, \dots, F_n\}$ be a generalised Heegaard splitting for M_i , obtained by untelescoping a minimal genus Heegaard splitting. (See [6] for more details.) We may assume that the even surfaces are incompressible and have no 2-sphere components. They may therefore be realized as a minimal surface in M_i , by a theorem of Freedman, Hass and Scott [2]. We may also guarantee that the odd surfaces are strongly irreducible and may each therefore be isotoped to a minimal surface (or a double cover of a minimal non-orientable surface with a small tube attached), by a theorem of Pitts and Rubinstein [5]. After a small perturbation, we may assume that the surfaces F_1, \dots, F_n are disjoint. The number of such surfaces is at most $x + 1$, and, for each j , $|\chi(F_j)| \leq x$. Thus, $|\chi(F_1 \cup \dots \cup F_n)| \leq x(x + 1) < 2x^2$. By Lemma 1, the number of complementary regions of K_i that can intersect $F_1 \cup \dots \cup F_n$ is at most $2k_8x^2$. Let D be the corresponding subset of G_i .

Similarly, let C be the subset of G_i that corresponds to those complementary regions of K_i which intersect S . By Lemma 1, $|C| \leq k_8|\chi(S)| \leq k_8k_4x$.

We claim that, when i is sufficiently large, there are at least $3x(x+2)/2$ disjoint translates of S under G_i that are also disjoint from $F_1 \cup \dots \cup F_n$. Let $m = 3x(x+2)/2$. If the claim is not true, then for any m -tuple (Sg_1, \dots, Sg_m) of copies of S (where $g_j \in G_i$ for each j), either at least two must intersect or one copy intersects $F_1 \cup \dots \cup F_n$. In the former case, $c_1g_j = c_2g_k$, for some c_1 and c_2 in C , and for $1 \leq j < k \leq m$. Hence, $g_jg_k^{-1} \in C^{-1}C$. In the latter case, $c_1g_j = d$ for some c_1 in C and d in D , and so $g_j \in C^{-1}D$. Thus, the sets $q_{jk}^{-1}(C^{-1}C)$ and $p_j^{-1}(C^{-1}D)$ cover G_i^m , where q_{jk} and p_j are the maps

$$\begin{aligned} q_{jk}: G_i^m &\rightarrow G_i \\ (g_1, \dots, g_m) &\mapsto g_jg_k^{-1} \end{aligned}$$

$$\begin{aligned} p_j: G_i^m &\rightarrow G_i \\ (g_1, \dots, g_m) &\mapsto g_j, \end{aligned}$$

for $1 \leq j < k \leq m$. The former sets each have size $|G_i|^{m-1}|C^{-1}C|$, and the latter have size $|G_i|^{m-1}|C^{-1}D|$. So,

$$|G_i|^m \leq \binom{m}{2}|G_i|^{m-1}|C|^2 + m|G_i|^{m-1}|C||D|.$$

This implies that

$$d_i = |G_i| \leq \binom{m}{2}(k_8k_4x)^2 + m(k_8k_4x)(2k_8x^2).$$

The right-hand side has order x^6 as $i \rightarrow \infty$. However, $x/\sqrt[6]{d_i} \rightarrow 0$, which is a contradiction, proving the claim.

Consider these $3x(x+2)/2$ copies of S . At least $3x/2$ of these must lie in the same complementary region of $F_1 \cup \dots \cup F_n$, which is a collection of compression bodies. Since S is incompressible and connected, each copy of S must be parallel to a negative boundary component of one of these compression bodies. There are at most $x/2$ boundary components. So, at least 3 copies of S are parallel, and at least 2 of these are coherently oriented. The proof is now completed by the following lemma. \square

Lemma 2. *Let S be a connected properly embedded oriented incompressible surface in a closed orientable 3-manifold M_i . Suppose that the image of S under some finite order homeomorphism h of M_i is disjoint from S , parallel to it and coherently oriented. Then M_i fibres over the circle with fibre S .*

Proof. Let Y be the manifold lying between S and $h(S)$. It is copy of $S \times I$, with S and $h(S)$ corresponding to $S \times \{0\}$ and $S \times \{1\}$. Take a countable collection $\{Y_n : n \in \mathbb{Z}\}$ of copies of this manifold. Glue $S \times \{1\}$ in Y_n to $S \times \{0\}$ in Y_{n+1} , via h^{-1} . The resulting space Y_∞ is a copy of $S \times \mathbb{R}$. Let H be the automorphism of this space taking Y_n to Y_{n+1} for each n , via the ‘identity’. Let $p: Y_0 \rightarrow Y$ be the identification homeomorphism. Extend this to a map $p: Y_\infty \rightarrow M_i$ by defining $p|_{Y_n}$ to be $h^n p H^{-n}$.

We claim that this is a covering map. It may expressed as a composition $Y_\infty \rightarrow Y_\infty / \langle H^N \rangle \rightarrow M_i$, where N is the order of h . The first of these maps is obviously a covering map. The second is also, since it is a local homeomorphism and $Y_\infty / \langle H^N \rangle$ is compact. Hence, p is a covering map.

By construction, $h^n(S)$ lifts homeomorphically to $Y_n \cap Y_{n+1}$, for each n . The inverse image of S in Y_∞ includes all translates of $Y_0 \cap Y_1$ under $\langle H^N \rangle$. Since this inverse image is incompressible, we deduce that it divides Y_∞ into a collection of copies of $S \times I$. The restriction of p to one of these components is a covering map to a component of $M_i - S$. By examining this map near S , we see that it is degree one and hence a homeomorphism, and that it surjects onto $M_i - S$. Therefore, M_i is obtained by taking $S \times I$ and gluing its boundary components homeomorphically. So, it fibres over the circle with fibre S . \square

REFERENCES

1. R. BROOKS, *The spectral geometry of a tower of coverings*, J. Differential Geom. **23** (1986) 97–107.
2. M. FREEDMAN, J. HASS AND P. SCOTT, *Least area incompressible surfaces in 3-manifolds*, Invent. Math. **71** (1983) 609–642.
3. M. LACKENBY, *Heegaard splittings, the virtually Haken conjecture and Property (τ)* , Preprint.

4. A. LUBOTZKY, *Eigenvalues of the Laplacian, the first Betti number and the congruence subgroup problem*, Ann. Math. (2) **144** (1996) 441–452.
5. J. PITTS AND J. H. RUBINSTEIN, *Existence of minimal surfaces of bounded topological type in three-manifolds*. Miniconference on geometry and partial differential equations (Canberra, 1985), 163–176.
6. M. SCHARLEMANN AND A. THOMPSON, *Thin position for 3-manifolds*, Geometric topology (Haifa, 1992), 231–238, Contemp. Math., 164.
7. B. WHITE, *The space of minimal submanifolds for varying Riemannian metrics*, Indiana Math. Journal **40** (1991) 161–200.

Mathematical Institute, Oxford University,
24-29 St Giles', Oxford OX1 3LB, UK.