# ABSOLUTE CONVERGENCE OF THE SPECTRAL SIDE OF THE ARTHUR TRACE FORMULA FOR $GL_n$

WERNER MÜLLER AND BIRGIT SPEH<sup>†</sup>

#### 0. INTRODUCTION

Let E be a number field and let G be the group  $\operatorname{GL}_n$  considered as reductive algebraic group over E. Let  $\mathbb{A}$  be the ring of adèles of Eand let  $G(\mathbb{A})$  be the group of points of G with values in  $\mathbb{A}$ . Let  $G(\mathbb{A})^1$ be the intersection of the kernels of the maps  $x \mapsto |\chi(x)|, x \in G(\mathbb{A})$ , where  $\chi$  ranges over the group  $X(G)_E$  of characters of G defined over E. Then the (noninvariant) trace formula of Arthur is an identity

$$\sum_{\mathfrak{o}\in\mathfrak{O}}J_{\mathfrak{o}}(f)=\sum_{\chi\in\mathfrak{X}}J_{\chi}(f),\quad f\in C_0^\infty(G(\mathbb{A})^1),$$

between distributions on  $G(\mathbb{A})^1$ . The left hand side is the *geometric* side and the right hand side the spectral side of the trace formula.

In this paper we are concerned with the spectral side of the trace formula. The distributions  $J_{\chi}$  are defined in terms of truncated Eisenstein series. They are parametrized by the set of cuspidal data  $\mathfrak{X}$  which consists of the Weyl group orbits of pairs  $(M_B, r_B)$ , where  $M_B$  is the Levi component of a standard parabolic subgroup and  $r_B$  is an irreducible cuspidal automorphic representation of  $M_B(\mathbb{A})^1$ .

Let K be a maximal compact subgroup of  $G(\mathbb{A})$ . In this paper we prove that for all K-finite functions  $f \in C_0^{\infty}(G(\mathbb{A})^1)$ , the spectral side of the trace formula is absolutely convergent. To make this statement more explicit, we have to introduce some notation.

Let  $P_0$  be the subgroup of upper triangular matrices of G. This is the minimal standard parabolic subgroup of G. Its Levi subgroup  $M_0$  is the group of diagonal matrices. Let P be a parabolic subgroup of G, defined over E, with unipotent radical  $N_P$  such that its Levi component  $M_P$  contains  $M_0$ . We denote the split component of the center of  $M_P$ 

<sup>&</sup>lt;sup>†</sup>partially supported by NSF grant DMS-0070561

Date: June 30, 2019.

by  $A_P$  and its Lie algebra by  $\mathfrak{a}_P$ . For parabolic groups  $P \subset Q$  there is a natural surjective map  $\mathfrak{a}_P \to \mathfrak{a}_Q$  whose kernel we will denote by  $\mathfrak{a}_P^Q$ . Let  $\mathcal{A}^2(P)$  be the space of square integrable automorphic forms on  $N_P(\mathbb{A})M_P(E)\backslash G(\mathbb{A})$ . Let Q be another parabolic subgroup of G, defined over E, with Levi component  $M_Q$ , split component  $A_Q$  and corresponding Lie algebra  $\mathfrak{a}_Q$ . Let  $W(\mathfrak{a}_P,\mathfrak{a}_Q)$  be the set of all linear isomorphisms from  $\mathfrak{a}_P$  to  $\mathfrak{a}_Q$  which are restrictions of elements of the Weyl group  $W(A_0)$ . The theory of Eisenstein series associates to each  $s \in W(\mathfrak{a}_P,\mathfrak{a}_Q)$  an intertwining operator

$$M_{Q|P}(s,\lambda): \mathcal{A}^2(P) \to \mathcal{A}^2(Q), \quad \lambda \in \mathfrak{a}_{P,\mathbb{C}}^*,$$

which for  $\operatorname{Re}(\lambda)$  in a certain chamber, can be defined by an absolutely convergent integral and admits an analytic continuation to a meromorphic function of  $\lambda \in \mathfrak{a}_{P,\mathbb{C}}^*$ . Set

$$M_{Q|P}(\lambda) := M_{Q|P}(1,\lambda).$$

Let  $\Pi(M_P(\mathbb{A})^1)$  be the set of equivalence classes of irreducible unitary representations of  $M_P(\mathbb{A})^1$ . Let  $\chi \in \mathfrak{X}$  and  $\pi \in \Pi(M_P(\mathbb{A})^1)$ . Then  $(\chi, \pi)$  singles out a certain subspace  $\mathcal{A}^2_{\chi,\pi}(P)$  of  $\mathcal{A}^2(P)$  [A3, p.1249]. Let  $\overline{\mathcal{A}}^2_{\chi,\pi}(P)$  be the Hilbert space completion of  $\mathcal{A}^2_{\chi,\pi}(P)$  with respect to the canonical inner product. For each  $\lambda \in \mathfrak{a}^*_{P,\mathbb{C}}$  we have an induced representation  $\rho_{\chi,\pi}(P,\lambda)$  of  $G(\mathbb{A})$  in  $\overline{\mathcal{A}}^2_{\chi,\pi}(P)$ .

For each Levi subgroup L let  $\mathcal{P}(L)$  be the set of all parabolic subgroups with Levi component L. If P is a parabolic subgroup, let  $\Delta_P$  denote the set of simple roots of  $(P, A_P)$ . Let L be a Levi subgroup which contains  $M_P$ . Set

$$\mathfrak{M}_{L}(P,\lambda) = \lim_{\Lambda \to 0} \left( \sum_{Q_{1} \in \mathcal{P}(L)} \operatorname{vol}(\mathfrak{a}_{Q_{1}}^{G} / \mathbb{Z}(\Delta_{Q_{1}}^{\vee})) M_{Q|P}(\lambda)^{-1} \frac{M_{Q|P}(\lambda + \Lambda)}{\prod_{\alpha \in \Delta_{Q_{1}}} \Lambda(\alpha^{\vee})} \right),$$

where  $\lambda$  and  $\Lambda$  are constrained to lie in  $i\mathfrak{a}_L^*$ , and for each  $Q_1 \in \mathcal{P}(L)$ , Q is a group in  $\mathcal{P}(M_P)$  which is contained in  $Q_1$ . Then  $\mathfrak{M}_L(P,\lambda)$  is an unbounded operator which acts on the Hilbert space  $\overline{\mathcal{A}}_{\chi,\pi}^2(P)$ . In the special case that L = M and  $\dim \mathfrak{a}_L^G = 1$ , the operator  $\mathfrak{M}_L(P,\lambda)$  has a simple description. Let P be a parabolic subgroup with Levi component M. Let  $\alpha$  be the unique simple root of  $(P, A_P)$  and let  $\tilde{\omega}$ be the element in  $(\mathfrak{a}_M^G)^*$  such that  $\tilde{\omega}(\alpha^{\vee}) = 1$ . Let  $\overline{P}$  be the opposite parabolic group of P. Then

$$\mathfrak{M}_{L}(P, z\tilde{\omega}) = -\operatorname{vol}(\mathfrak{a}_{M}^{G}/\mathbb{Z}\alpha^{\vee})M_{\overline{P}|P}(z\tilde{\omega})^{-1} \cdot \frac{d}{dz}M_{\overline{P}|P}(z\tilde{\omega}).$$

Let  $f \in C_c^{\infty}(G(\mathbb{A})^1)$ . Then Arthur [A4, Theorem 8.2] proved that  $J_{\chi}(f)$  equals the sum over Levi subgroups M containing  $M_0$ , over L containing M, over  $\pi \in \Pi(M(\mathbb{A})^1)$ , and over  $s \in W^L(\mathfrak{a}_M)_{\mathrm{reg}}$ , a certain subset of the Weyl group, of the product of

$$|W_0^M||W_0|^{-1}|\det(s-1)_{\mathfrak{a}_M^L}|^{-1}|\mathcal{P}(M)|^{-1}$$

a factor to which we need not pay to much attention, and of

(0.1) 
$$\int_{i\mathfrak{a}_L^*/i\mathfrak{a}_G^*} \sum_{P \in \mathcal{P}(M)} \operatorname{tr}(\mathfrak{M}_L(P,\lambda) M_{P|P}(s,0) \rho_{\chi,\pi}(P,\lambda,f)) \, d\lambda.$$

So far, it is only known that  $\sum_{\chi \in \mathfrak{X}} |J_{\chi}(f)| < \infty$  and the goal is to show that the integral-sum obtained by summing (0.1) over  $\chi \in \mathfrak{X}$  and  $\pi \in \Pi(M(\mathbb{A})^1)$  is absolutely convergent with respect to the trace norm. For a given Levi subgroup M let  $\mathcal{L}(M)$  be the set of all Levi subgroups L with  $M \subset L$ . Put  $M(P,s) = M_{P|P}(s,0)$ . Denote by  $|| T ||_1$  the trace norm of a trace class operator T. Let  $\mathcal{C}^1(G(\mathbb{A})^1)$  be the space of integrable rapidly decreasing functions on  $G(\mathbb{A})^1$  (see §1.4 for its definition). Then our main result is the following theorem.

**Theorem 0.1.** Let  $f \in \mathcal{C}^1(G(\mathbb{A})^1)$  be K-finite. Then the sum over all  $M \in \mathcal{L}(M_0)$ ,  $L \in \mathcal{L}(M)$ ,  $\chi \in \mathfrak{X}$ ,  $\pi \in \Pi(M(\mathbb{A})^1)$ , and  $s \in W^L(\mathfrak{a}_M)_{\text{reg}}$  of the product of

$$|W_0^M||W_0|^{-1}|\det(s-1)_{\mathfrak{a}_M^L}|^{-1}$$

with

$$\int_{i\mathfrak{a}_{L}^{*}/i\mathfrak{a}_{G}^{*}} |\mathcal{P}(M)|^{-1} \sum_{P \in \mathcal{P}(M)} \|\mathfrak{M}_{L}(P,\lambda)M(P,s)\rho_{\chi,\pi}(P,\lambda,f)\|_{1} d\lambda$$

is convergent.

We think that the K-finiteness assumption can be removed. This depends on a problem in local harmonic analysis which we shall describe below.

By Theorem 0.1, the spectral side can now be rewritten in the following way. Denote by  $\Pi_{\text{disc}}(M(\mathbb{A})^1)$  the set of all  $\pi \in \Pi(M(\mathbb{A})^1)$  which are equivalent to an irreducible subrepresentation of the regular representation of  $M(\mathbb{A})^1$  in  $L^2(M(E)\backslash M(\mathbb{A})^1)$ . As in Section 7 of [A3], we shall identify any representation of  $M(\mathbb{A})^1$  with a representation of  $M(\mathbb{A})$  which is trivial on  $A_{M,\mathbb{Q}}(\mathbb{R})^0$ , where  $A_{M,\mathbb{Q}}$  is the split component of the center of the group  $\operatorname{Res}_{E/\mathbb{Q}} \operatorname{GL}_n$  obtained from Gby restricting scalars from E to  $\mathbb{Q}$ . For any parabolic group P, let  $\mathcal{A}^2_{\pi}(P) = \bigoplus_{\chi} \mathcal{A}^2_{\chi,\pi}(P)$  and for  $\lambda \in \mathfrak{a}^*_{P,\mathbb{C}}$ , let  $\rho_{\pi}(P,\lambda)$  be the induced representation of  $G(\mathbb{A})$  in  $\overline{\mathcal{A}}^2_{\pi}(P)$ , the Hilbert space completion of  $\mathcal{A}^2_{\pi}(P)$ . Given  $M \in \mathcal{L}, L \in \mathcal{L}(M), P \in \mathcal{P}(M), s \in W^L(\mathfrak{a}_M)_{\text{reg}}$  and a K-finite function  $f \in \mathcal{C}^1(G(\mathbb{A})^1)$ , let

$$J_{M,P}^{L}(f,s) = \sum_{\pi \in \Pi_{\operatorname{disc}}(M(\mathbb{A})^{1})} \int_{i\mathfrak{a}_{L}^{*}/i\mathfrak{a}_{G}^{*}} \operatorname{tr}(\mathfrak{M}_{L}(P,\lambda)M_{P|P}(s,0)\rho_{\pi}(P,\lambda,f)) \, d\lambda.$$

By Theorem 0.1 this integral-series is absolutely convergent with respect to the trace norm. Furthermore for  $M \in \mathcal{L}$  and  $s \in W^L(\mathfrak{a}_M)_{\text{reg}}$ set

$$a_{M,s} = |\mathcal{P}(M)|^{-1} |W_0^M| |W_0|^{-1} |\det(s-1)_{\mathfrak{a}_M^L}|^{-1}.$$

Then for all K-finite functions f in  $\mathcal{C}^1(G(\mathbb{A})^1)$ , the spectral side of the Arthur trace formula equals

$$\sum_{M \in \mathcal{L}} \sum_{L \in \mathcal{L}(M)} \sum_{P \in \mathcal{P}(M)} \sum_{s \in W^L(\mathfrak{a}_M)_{\text{reg}}} a_{M,s} J^L_{M,P}(f,s).$$

Note that all sums in this expression are finite.

We shall now explain the main steps of the proof of Theorem 0.1. The proof relies on Theorem 0.2 of [Mu4]. In this theorem the absolute convergence of the spectral side of the trace formula has been reduced to a problem about local components of automorphic representations. So the main issue of the present paper is to verify that for  $GL_n$ , the assumptions of Theorem 0.2 of [Mu4] are satisfied.

Let  $M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$  be a standard Levi subgroup and let  $P, Q \in \mathcal{P}(M)$ . We shall identify  $\mathfrak{a}_M^*$  with  $\mathbb{R}^r$ . Given a place v of E and an irreducible unitary representation  $\pi_v = \bigotimes_{i=1}^r \pi_{v,i}$  of  $M(E_v)$ , let  $J_{Q|P}(\pi_v, \mathbf{s}), \mathbf{s} \in \mathbb{C}^r$ , be the the local intertwining operator between the induced representations  $I_P^G(\pi_v[\mathbf{s}])$  and  $I_Q^G(\pi_v[\mathbf{s}])$ , where  $\mathbf{s} = (s_1, \dots, s_r)$  and  $\pi_v[\mathbf{s}] = \bigotimes_i (\pi_{v,i} |\det|^{s_i})$ . It follows from results of Shahidi [Sh5] that there exist normalizing factors  $r_{Q|P}(\pi_v, \mathbf{s})$  such that the normalized intertwining operators

$$R_{Q|P}(\pi_v, \mathbf{s}) = r_{Q|P}(\pi_v, \mathbf{s})^{-1} J_{Q|P}(\pi_v, \mathbf{s})$$

satisfy the properties of Theorem 2.1 of [A7]. If  $v < \infty$  and  $K_v$  is an open compact subgroup of  $G(E_v)$ , denote by  $R_{Q|P}(\pi_v, \mathbf{s})_{K_v}$  the restriction of  $R_{Q|P}(\pi_v, \mathbf{s})$  to the subspace  $\mathcal{H}_P(\pi_v)^{K_v}$  of  $K_v$ -invariant vectors in the Hilbert space  $\mathcal{H}_P(\pi_v)$  of the induced representation. If  $v|\infty$ , let  $K_v \subset G(E_v)$  be the standard maximal compact subgroup. Given  $\pi_v \in \Pi(G(E_v))$  and  $\sigma_v \in \Pi(K_v)$ , let  $R_{Q|P}(\pi_v, \mathbf{s})_{\sigma_v}$  be the restriction of  $R_{Q|P}(\pi_v, \mathbf{s})$  to the  $\sigma_v$ -isotypical subspace of  $\mathcal{H}_P(\pi_v)$ . Let  $\lambda_{\pi_v}$  be the Casimir eigenvalue of  $\pi_v$ . Finally let  $\Pi_{\text{disc}}(M(E_v))$  be the subspace of all  $\pi_v$  in  $\Pi(M(E_v))$  such that there exists an automorphic representation  $\pi$  in the discrete spectrum of  $M(\mathbb{A})$  whose local component at vis equivalent to  $\pi_v$ . Then our key result is the following proposition.

**Proposition 0.2.** Let v be a place of E. For all  $M \in \mathcal{L}$  and  $P, Q \in \mathcal{P}(M)$  the following holds.

1) If  $v < \infty$ , then for every open compact subgroup  $K_v$  of  $\operatorname{GL}_n(E_v)$  and every multi-index  $\alpha \in \mathbb{N}^r$  there exists C > 0 such that

$$(0.2) \| D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_v, i\mathbf{u})_{K_v} \| \le C$$

for all  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$  and  $\mathbf{u} \in \mathbb{R}^r$ .

2) If  $v|\infty$ , then for every  $\sigma_v \in \Pi(K_v)$  and every multi-index  $\alpha \in \mathbb{N}^r$ there exist C > 0 and  $N \in \mathbb{N}$  such that

(0.3) 
$$|| D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_{v}, i\mathbf{u})_{\sigma_{v}} || \leq C(1 + |\lambda_{\pi_{v}}| + || \mathbf{u} ||)^{N}$$

for all  $\mathbf{u} \in \mathbb{R}^r$  and all  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$ .

Together with Theorem 0.2 of [Mu4], this implies Theorem 0.1.

In order to remove the K-finiteness assumption in Theorem 0.1, one has to establish a stronger version of condition (0.3) which takes into account the dependence on  $\sigma_v$ . Let  $\lambda_{\sigma_v}$  denote the Casimir eigenvalue of a given  $\sigma_v \in \Pi(K_v)$ . Suppose that for all  $M \in \mathcal{L}$  in place of condition (0.3) the following condition holds: For every  $v \mid \infty$  and multi-index  $\alpha \in \mathbb{N}^r$  there exist C > 0 and  $N \in \mathbb{N}$  such that

(0.4) 
$$|| D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_{v}, i\mathbf{u})_{\sigma_{v}} || \leq C(1 + |\lambda_{\pi_{v}}| + |\lambda_{\sigma_{v}}| + || \mathbf{u} ||)^{N}$$

for all  $\mathbf{u} \in \mathbb{R}^r$ ,  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$  and  $\sigma_v \in \Pi(K_v)$ . Then it follows from Theorem 0.1 of [Mu4] that the spectral side of the trace formula is absolutely convergent for all  $f \in \mathcal{C}^1(G(\mathbb{A})^1)$ . At the moment, however, we don't know how to prove the stronger estimate (0.4).

The existence of estimates of the form (0.2) and (0.3) follows from nontrivial bounds on the continuous parts of the Langlands parameters of  $\pi_v \in \prod_{\text{disc}} (M(E_v))$ . To prove that appropriate bounds exist, we have to study local components of cuspidal automorphic representations of  $\operatorname{GL}_m(\mathbb{A})$ . First note that any local component  $\pi_v$  of a cuspidal automorphic representation  $\pi$  of  $\operatorname{GL}_m(\mathbb{A})$  is generic [Sk]. This implies that  $\pi_v$  is equivalent to a fully induced representation [JS3], i.e.,

$$\pi_v \cong I_{P(E_v)}^{G(E_v)}(\tau_1[t_1], ..., \tau_r[t_r]),$$

where P is a standard parabolic subgroup of type  $(n_1, ..., n_r)$ ,  $\tau_i$  are tempered representations of  $\operatorname{GL}_{n_i}(E_v)$  and the  $t_i$ 's are real numbers satisfying

$$t_1 > t_2 > \cdots > t_r.$$

Here  $\tau_i[t_i]$  is the representation  $g \mapsto \tau_i(g) |\det(g)|^{t_i}$ . For a unitary generic representation  $\pi_v$  the parameters  $t_i$  satisfy  $|t_i| < 1/2$ . In [LRS], Luo, Rudnick and Sarnak proved that for an unramified  $\pi_v$  which is the local component of a cuspidal automorphic representation of  $\operatorname{GL}_m(\mathbb{A})$ , one has

(0.5) 
$$\max_{i} |t_{i}| < \frac{1}{2} - \frac{1}{m^{2} + 1}$$

First we extend this result of Luo, Rudnick and Sarnak to all local components of cuspidal automorphic representations of  $\operatorname{GL}_m(\mathbb{A})$ . Then we use the description of the residual spectrum of  $\operatorname{GL}_m(\mathbb{A})$ , given by Mœglin and Waldspurger [MW], to prove similar bounds for the local components of all automorphic representations in the residual spectrum of  $\operatorname{GL}_m(\mathbb{A})$  (cf. Proposition 3.4 for the precise statement). Then Proposition 0.2 is a consequence of these bounds and Proposition I.10 of [MW].

Especially, it implies that there exists  $\delta > 0$  such that for every local component  $\pi_v$  of an automorphic representation  $\pi$  in the discrete spectrum of  $M(\mathbb{A})^1$  the normalized intertwining operator  $R_{Q|P}(\pi_v, \mathbf{s})$ is holomorphic in the domain  $|\operatorname{Re}(s_i - s_j)| < \delta, 1 \leq i, j \leq r$ . As the example in section 4 shows, this is a necessary condition for (0.2) and (0.3) to hold.

Finally we make some remarks about the compatability with the assumptions made in [Mu4]. First of all, the normalization of local intertwining operators used in the present paper differs from the one used in [Mu4]. The latter one relies on Langlands' general construction [CLL, §15] of normalizing factors at the finite places. So in order to be able to apply Theorem 0.2 of [Mu4], we have to compare the two different normalizations. It turns out that they differ essentially by the  $\varepsilon$ -factors appearing in the functional equation of the Rankin-Selberg L-functions. Using some estimates of conductors, it follows that Theorem 0.2 of [Mu4] also holds with respect to the normalizations given by L-functions.

Furthermore, the reductive algebraic groups G considered in [Mu4] are supposed to be defined over  $\mathbb{Q}$ . However, passing to the group  $G' = \operatorname{Res}_{E/\mathbb{Q}} G$  which is obtained from G by restriction of scalars, it follows immediately that the results of [Mu4] can also be applied to reductive algebraic groups defined over a number field.

The paper is organized as follows. In section 2 we compare the two different normalizations of intertwining operators and we prove some estimate for conductors. In section 3 we estimate the continuous parameters of local components of cuspidal automorphic representations of  $GL_m$  which generalizes results of Luo, Rudnick and Sarnak [LRS] to the case of ramified representations. Then we use the description of the residual spectrum of  $GL_m$  by Mœglin and Waldspurger [MW] to obtain estimations for the continuous parameters of all local components of automorphic representations in the discrete spectrum of  $GL_m$ . We use these results in section 4 to prove Proposition 0.2. In section 5 we consider the global normalizing factors and we show that Theorem 0.2 of [Mu4] also holds with respect to the normalizing factors defined by *L*-functions. Combined with Proposition 0.2 the proof of Theorem 0.1 follows.

Acknowledgment. The authors would like to thank G. Henniart, Ch. Kaiser and W. Zink for some very helpful discussions. Especially the last part of the proof of Lemma 2.2 was suggested by Ch. Kaiser.

## 1. Preliminaries

1.1. Let E be a number field and let A denote the ring of adèles of E. Fix a positive integer n and let G be the group  $\operatorname{GL}_n$  considered as algebraic group over E. By a parabolic subgroup of G we will always mean a parabolic subgroup which is defined over E. Let  $P_0$  be the subgroup of upper triangular matrices of G. The Levi subgroup  $M_0$  of  $P_0$  is the group of diagonal matrices in G. A parabolic subgroup P of G is called standard, if  $P \supset P_0$ . By a Levi subgroup we will mean a subgroup of G which contains  $M_0$  and is the Levi component of a parabolic subgroup of G. If  $M \subset L$  are Levi subgroups, we denote the set of Levi subgroups of L which contain M by  $\mathcal{L}^L(M)$ . Furthermore, let  $\mathcal{F}^L(M)$  denote the set of parabolic subgroups of L defined over E which contain M, and let  $\mathcal{P}^L(M)$  be the set of groups in  $\mathcal{F}^L(M)$ 

for which M is a Levi component. If L = G, we shall denote these sets by  $\mathcal{L}(M)$ ,  $\mathcal{F}(M)$  and  $\mathcal{P}(M)$ . Write  $\mathcal{L} = \mathcal{L}(M_0)$ . Suppose that  $P \in \mathcal{F}^L(M)$ . Then

$$P = N_P M_P,$$

where  $N_P$  is the unipotent radical of P and  $M_P$  is the unique Levi component of P which contains M.

Suppose that  $M \subset M_1 \subset L$  are Levi subgroups of G. If  $Q \in \mathcal{P}^L(M_1)$ and  $R \in \mathcal{P}^{M_1}(M)$ , there is a unique group  $Q(R) \in \mathcal{P}^L(M)$  which is contained in Q and whose intersection with  $M_1$  is R.

Let  $M \in \mathcal{L}$  and denote by A the split component of the center of M. Then A is defined over E. Let  $X(M)_E$  be the group of characters of M defined over E and set

$$\mathfrak{a}_M = \operatorname{Hom}(X(M)_E, \mathbb{R}).$$

Then  $\mathfrak{a}_M$  is a real vector space whose dimension equals that of A. Its dual space is

$$\mathfrak{a}_M^* = X(M)_E \otimes \mathbb{R}.$$

For any  $M \in \mathcal{L}$  there exists a partition  $(n_1, ..., n_r)$  of n such that

$$M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$$

Then  $\mathfrak{a}_M^*$  can be canonically identified with  $(\mathbb{R}^r)^*$  and  $W(\mathfrak{a}_M)$  coincides with the group  $S_r$  of permutations of the set  $\{1, ..., r\}$ .

1.2. Let H be a reductive algebraic group defined over  $\mathbb{Q}$ , let F be a local field of characteristic 0 and let K be an open compact subgroup of H(F). We shall denote by  $\Pi(H(\mathbb{A}))$  (resp.  $\Pi(H(F))$ ,  $\Pi(K)$ , etc.) the set of equivalence classes of irreducible unitary representations of  $H(\mathbb{A})$  (resp. H(F), K, etc.).

1.3. Let F be a local field of characteristic zero. If  $\pi$  is an admissible representation of  $\operatorname{GL}_m(F)$ , we shall denote by  $\tilde{\pi}$  the contragredient representation to  $\pi$ . Let  $\pi_i$ , i = 1, ..., r, be irreducible admissible representations of the group  $\operatorname{GL}_{n_i}(F)$ . Then  $\pi = \pi_1 \otimes \cdots \otimes \pi_r$  is an irreducible admissible representation of

$$M(F) = \operatorname{GL}_{n_1}(F) \times \cdots \times \operatorname{GL}_{n_r}(F).$$

For  $\mathbf{s} \in \mathbb{C}^r$  let  $\pi_i[s_i]$  be the representation of  $\operatorname{GL}_{n_i}(F)$  which is defined by

$$\pi_i[s_i](g) = |\det(g)|^{s_i} \pi_i(g), \quad g \in \operatorname{GL}_{n_i}(F).$$

Let

$$I_P^G(\pi, \mathbf{s}) = \operatorname{Ind}_{P(F)}^{G(F)}(\pi_1[s_1] \otimes \cdots \otimes \pi_r[s_r])$$

be the induced representation and denote by  $\mathcal{H}_P(\pi)$  the Hilbert space of the representation  $I_P^G(\pi, \mathbf{s})$ . We refer to  $\mathbf{s}$  as the continuous parameter of  $I_P^G(\pi, \mathbf{s})$ . Sometimes we will write  $I_P^G(\pi_1[s_1], ..., \pi_r[s_r])$  in place of  $I_P^G(\pi, \mathbf{s})$ .

1.4. Let  $G(\mathbb{A}_{\infty}) = \prod_{v \mid \infty} G(E_v)$  and denote by  $G(\mathbb{A}_{\infty})^1$  the intersection of  $G(\mathbb{A}_{\infty})$  with  $G(\mathbb{A})^1$ . Let  $\mathcal{U}(\mathfrak{g}(\mathbb{A}_{\infty})^1 \otimes \mathbb{C})$  be the universal enveloping algebra of the complexification of the Lie algebra of  $G(\mathbb{A}_{\infty})^1$ . Let  $K_f$  be an open compact subgroup of  $G(\mathbb{A}_f)^1$ . Then the double coset space  $K_f \setminus G(\mathbb{A})^1/K_f$  is a discrete union of countably many copies of  $G(\mathbb{A}_{\infty})^1$ . In particular it is a differentiable manifold. Let  $K_{\infty}$  be the standard maximal compact subgroup of  $G(\mathbb{A}_{\infty})$ . Let  $\Xi$  be the zonal spherical function on  $G(\mathbb{A}_{\infty})$  defined by  $\Xi(g) = \int_{K_{\infty}} e^{-\rho(H(kg))} dk$  and for  $g \in G(\mathbb{A}_{\infty})$ , let  $\kappa(g)$  denote the geodesic distance of the cosets  $K_{\infty}$  and  $gK_{\infty}$  in  $G(\mathbb{A}_{\infty})/K_{\infty}$ . We extend  $\Xi$  and  $\kappa$  to functions on  $G(\mathbb{A})$  in the obvious way. Suppose that  $\Omega$  is a subset of  $G(\mathbb{A})^1$  such that  $K_f \cdot \Omega \cdot K_f = \Omega$  and  $K_f \setminus \Omega/K_f$  is the disjoint union of finitely many copies of  $G(\mathbb{A}_{\infty})^1$ . Let  $\mathcal{C}^1(G(\mathbb{A})^1; \Omega, K_f)$  be the space of all functions  $h: G(\mathbb{A})^1 \to \mathbb{C}$  satisfying the following conditions:

- (1) h is biinvariant under  $K_f$ , supp  $h \subset \Omega$ , and  $h: K_f \setminus \Omega/K_f \to \mathbb{C}$  is a smooth function.
- (2) For all  $D_1, D_2 \in \mathcal{U}(\mathfrak{g}(\mathbb{A}_{\infty})^1 \otimes \mathbb{C})$  and all  $r \in \mathbb{N}$ , we have
  - $\|h\|_{D_1,D_2,r} := \sup_{x \in G(\mathbb{A})^1} \left( (1+\kappa(x))^r \Xi^{-2}(x) |D_1 * h * D_2(x)| \right) < \infty.$

Let  $\mathcal{C}^1(G(\mathbb{A})^1; \Omega, K_f)$  be equipped with the topology defined by the semi-norms  $\|\cdot\|_{D_1, D_2, r}$ . Then we denote by  $\mathcal{C}^1(G(\mathbb{A})^1)$  the topological direct limit over all pairs  $(\Omega, K_f)$  of the spaces  $\mathcal{C}^1(G(\mathbb{A})^1; \Omega, K_f)$ .

#### 2. Normalizing factors for local intertwining operators

Let F be a local field of characteristic 0. If F is non-Archimedean, let  $\mathcal{O}$  be the ring of integers of F and let  $\mathfrak{P}$  be the unique maximal ideal of  $\mathcal{O}$ . Let q be the number of elements of the residue field  $\mathcal{O}/\mathfrak{P}$ . Let  $K = \operatorname{GL}_n(\mathcal{O})$ . If F is Archimedean, let K be the standard maximal compact subgroup of  $\operatorname{GL}_n(F)$ , i.e.,  $K = \operatorname{O}(n)$ , if  $F = \mathbb{R}$ , and  $K = \operatorname{U}(n)$ , if  $F = \mathbb{C}$ .

Let  $M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$  be a standard Levi subgroup. We identify  $\mathfrak{a}_M$  with  $\mathbb{R}^r$ . Let  $P_1, P_2 \in \mathcal{P}(M)$ . Given  $\pi \in \Pi(M(F))$ , let

$$J_{P_2|P_1}(\pi, \mathbf{s}), \quad \mathbf{s} \in \mathbb{C}^r,$$

be the intertwining operator which intertwines the induced representations  $I_{P_1}^G(\pi, \mathbf{s})$  and  $I_{P_2}^G(\pi, \mathbf{s})$ . The intertwining operator  $J_{P_2|P_1}(\pi, \mathbf{s})$  is defined by an integral over  $N_{P_1}(F) \cap N_{\overline{P}_2}(F)$  which converges for Re( $\mathbf{s}$ ) in a certain chamber of  $\mathfrak{a}_M^*$ . It follows from [A7] and [CLL, §15] that the intertwining operators can be normalized in a suitable way. This means that there exist scalar valued meromorphic functions  $r_{P_2|P_1}(\pi, \mathbf{s})$ of  $\mathbf{s} \in \mathbb{C}^r$  such that the normalized intertwining operators

$$R_{P_2|P_1}(\pi, \mathbf{s}) = r_{P_2|P_1}(\pi, \mathbf{s})^{-1} J_{P_2|P_1}(\pi, \mathbf{s})$$

satisfy the properties of Theorem 2.1 of [A7]. The method used in [A7] works for every reductive group G. For  $GL_n$ , however, it follows from results of Shahidi [Sh1], [Sh5] that local intertwining operators can be normalized by L-functions.

The normalizing factors defined by the Rankin-Selberg *L*-functions can be described as follows. Fix a nontrivial continuous character  $\psi$  of the additive group  $F^+$  of F and equip F with the Haar measure which is selfdual with respect to  $\psi$ . First assume that P is a standard maximal parabolic subgroup with Levi component  $M = \operatorname{GL}_{n_1} \times \operatorname{GL}_{n_2}$ . Let  $\pi_i \in \Pi(\operatorname{GL}_{n_i}(F)), i = 1, 2$ . If F is non-Archimedean, let  $L(s, \pi_1 \times \pi_2)$ and  $\epsilon(s, \pi_1 \times \pi_2, \psi)$  be the Rankin-Selberg *L*-function and the  $\epsilon$ -factor, respectively, as defined in [JPS]. If F is Archimedean, let the *L*-function and the  $\epsilon$ -factor be defined by using the Langlands parametrization (cf. [A7], [Sh2]). Then the normalizing factor can be regarded as a function  $r_{\overline{P}|P}(\pi_1 \otimes \pi_2, s)$  of one complex variable which is given by

(2.1) 
$$r_{\overline{P}|P}(\pi_1 \otimes \pi_2, s) = \frac{L(s, \pi_1 \times \widetilde{\pi}_2)}{L(1+s, \pi_1 \times \widetilde{\pi}_2)\epsilon(s, \pi_1 \times \widetilde{\pi}_2, \psi)}.$$

For arbitrary rank, the normalizing factors are products of normalizing factors associated to rank one groups in M. Let  $e_i$ ,  $i = 1, \ldots, r$ , denote the standard basis of  $(\mathbb{R}^r)^*$ . Then there exist  $\sigma_1, \sigma_2 \in S_r$  such that the set of roots of  $P_1$  and  $P_2$ , respectively, are given by

(2.2) 
$$\Sigma_{P_k} = \{e_i - e_j \mid 1 \le i, j \le r, \sigma_k(i) < \sigma_k(j)\}, \quad k = 1, 2.$$

Put

$$I(\sigma_1, \sigma_2) = \{(i, j) \mid 1 \le i, j \le r, \, \sigma_1(i) < \sigma_1(j), \, \sigma_2(i) > \sigma_2(j)\}.$$

Then

$$\Sigma_{P_1} \cap \Sigma_{\overline{P}_2} = \{ e_i - e_j \mid (i, j) \in \mathcal{I}(\sigma_1, \sigma_2) \}$$

Let  $\pi = \pi_1 \otimes \cdots \otimes \pi_r$  where  $\pi_i \in \Pi(\operatorname{GL}_{n_i}(F)), i = 1, ..., r$ . For  $\mathbf{s} = (s_1, ..., s_r) \in \mathbb{C}^r$  set

(2.3) 
$$\prod_{\substack{(i,j)\in \mathbf{I}(\sigma_1,\sigma_2)}} \frac{L(s_i - s_j, \pi_i \times \widetilde{\pi}_j)}{L(1 + s_i - s_j, \pi_i \times \widetilde{\pi}_j)\epsilon(s_i - s_j, \pi_i \times \widetilde{\pi}_j, \psi)}.$$

Since the Rankin-Selberg *L*-factors are meromorphic functions, it follows that  $r_{P_2|P_1}(\pi, \mathbf{s})$  are meromorphic functions of  $\mathbf{s} \in \mathbb{C}^r$  and as explained in [A7, §4] and [AC, p.87], they satisfy all properties that are requested for normalizing factors.

In order to be able to apply the results of [Mu4] we have to compare the normalizing factors  $r_{Q|P}(\pi, \mathbf{s})$  with those used in [Mu4] which we denote by  $\tilde{r}_{Q|P}(\pi, \mathbf{s})$ . If F is Archimedean, the normalizing factors  $\tilde{r}_{Q|P}(\pi, \mathbf{s})$  are defined as the Artin *L*-factors and therefore, coincide with the  $r_{Q|P}(\pi, \mathbf{s})$ . Assume that F is non-Archimedean. By the construction of the normalizing factors it suffices to consider the case where P is maximal,  $Q = \overline{P}$  and  $\pi$  is square integrable. Let P be a standard maximal parabolic subgroup of  $\operatorname{GL}_m$  with Levi component

$$M = \operatorname{GL}_{m_1} \times \operatorname{GL}_{m_2}$$
.

Then the normalizing factor may be regarded as a function  $\tilde{r}_{\overline{P}|P}(\pi, s)$ of one complex variable s. We recall the construction of  $\tilde{r}_{\overline{P}|P}(\pi, s)$  for square integrable representations  $\pi$  [CLL]. It follows from [Si1], [Si2] that for every  $\pi \in \Pi_2(M(F))$  there exists a rational function  $U_P(\pi, z)$ such that the Plancherel measure  $\mu(\pi, s)$  is given by

$$\mu(\pi, s) = U_P(\pi, q^{-s}).$$

The rational function  $U_P(\pi, z)$  is of the form

$$U_P(\pi, z) = a \prod_{i=1}^r \frac{(1 - \alpha_i z)(1 - \overline{\alpha}_i^{-1} z)}{(1 - \beta_i z)(1 - \overline{\beta}_i^{-1} z)},$$

where  $|\alpha_i| \leq 1, |\beta_i| \leq 1, i = 1, ..., r$ , and  $a \in \mathbb{C}$  is a constant such that

$$a\prod_{i=1}^r \frac{\alpha_i}{\beta_i} > 0.$$

Let  $b \in \mathbb{C}$  be such that

$$|b|^2 \prod_{i=1}^r \frac{\overline{\alpha}_i}{\overline{\beta}_i} = a$$

and set

(2.4) 
$$V_P(\pi, z) = b \prod_{i=1}^r \frac{(1 - \alpha_i z)}{(1 - \beta_i z)}.$$

Then the normalizing factor  $\widetilde{r}_{\overline{P}|P}(\pi,s)$  is defined by

$$\widetilde{r}_{\overline{P}|P}(\pi,s) = V_P(\pi,q^{-s})^{-1}.$$

By definition we have

$$\mu_P(\pi, s) = \left(\overline{\widetilde{r_{P|P}}(\pi, -\overline{s})}\widetilde{r_{P|P}}(\pi, s)\right)^{-1},$$

which is one of the main conditions that the normalizing factors have to satisfy.

Let  $\pi_1$  and  $\pi_2$  be tempered representations of  $\operatorname{GL}_{m_1}(F)$  and  $\operatorname{GL}_{m_2}(F)$ , respectively. By Corollary 6.1.2 of [Sh1] the Plancherel measure is given by

$$\mu(\pi_1 \otimes \pi_2, s) = q^{f(\widetilde{\pi}_1 \times \pi_2)} \frac{L(1+s, \pi_1 \times \widetilde{\pi}_2)}{L(s, \pi_1 \times \widetilde{\pi}_2)} \frac{L(1-s, \widetilde{\pi}_1 \times \pi_2)}{L(-s, \widetilde{\pi}_1 \times \pi_2)},$$

where  $f(\tilde{\pi}_1 \times \pi_2) \in \mathbb{Z}$  is the conductor of  $\tilde{\pi}_1 \times \pi_2$ . Using the description of the Rankin-Selberg *L*-functions for tempered representations [JPS] (see also section 3), it follows that

(2.5) 
$$L(s, \pi_1 \times \widetilde{\pi}_2) = \prod_{i=1}^{m_1} \prod_{j=1}^{m_2} \left(1 - a_{ij}q^{-s}\right)^{-1},$$

with complex numbers  $a_{ij}$  satisfying  $|a_{ij}| < 1$ . Furthermore by [JPS], the  $\epsilon$ -factor  $\epsilon(s, \pi_1 \times \pi_2, \psi)$  has the following form

(2.6) 
$$\epsilon(s, \pi_1 \times \pi_2, \psi) = c(\pi_1 \times \pi_2, \psi)q^{-f(\pi_1 \times \pi_2, \psi)s},$$

with 
$$c(\pi_1 \times \pi_2, \psi) \in \mathbb{C} - \{0\}$$
 and  $f(\pi_1 \times \pi_2, \psi) \in \mathbb{Z}$ . Let  
 $c(\psi) = \max\{r \mid \mathfrak{P}^{-r} \subset \ker \psi\}$ 

Then

(2.7) 
$$f(\pi_1 \times \pi_2, \psi) = n_1 n_2 c(\psi) + f(\pi_1 \times \pi_2)$$

with  $f(\pi_1 \times \pi_2) \in \mathbb{Z}$  independent of  $\psi$ . For simplicity assume that  $c(\psi) = 0$ . By Lemma 6.1 of [Sh1] we have

(2.8) 
$$|c(\pi_1 \times \pi_2)| = q^{f(\tilde{\pi}_1 \times \pi_2)/2}.$$

Thus the  $\epsilon$ -factor can be written as

$$\epsilon(s, \pi_1 \times \pi_2, \psi) = W(\pi_1 \times \pi_2)q^{(1/2-s)f(\pi_1 \times \pi_2)},$$

where the root number  $W(\pi_1 \times \pi_2)$  satisfies  $|W(\pi_1 \times \pi_2)| = 1$ . Finally, observe that  $f(\pi_1 \times \tilde{\pi}_2) = f(\tilde{\pi}_1 \times \pi_2)$ . Using (2.5), (2.6) and (2.8) it follows that the constant b in (2.4) can be chosen to be  $\epsilon(0, \pi_1 \times \tilde{\pi}_2, \psi)$  and

(2.9) 
$$\widetilde{r}_{\overline{P}|P}(\pi_1 \otimes \pi_2, s) = \frac{L(s, \pi_1 \times \widetilde{\pi}_2)}{L(1+s, \pi_1 \times \widetilde{\pi}_2)\epsilon(0, \pi_1 \times \widetilde{\pi}_2, \psi)}.$$

Comparing (2.1) and (2.9), it follows that

(2.10) 
$$r_{\overline{P}|P}(\pi_1 \otimes \pi_2, s) = \frac{\epsilon(0, \pi_1 \times \widetilde{\pi}_2, \psi)}{\epsilon(s, \pi_1 \times \widetilde{\pi}_2, \psi)} \widetilde{r}_{\overline{P}|P}(\pi_1 \otimes \pi_2, s).$$

This can be extended to parabolic groups of arbitrary rank in the usual way. Let M be a standard Levi subgroup of  $\operatorname{GL}_n$  of type  $(n_1, \ldots, n_r)$ and let  $P_1, P_2 \in \mathcal{P}(M)$ . Using the product formula for  $\tilde{r}_{P_2|P_1}(\pi, \mathbf{s})$ [A7, p.29] and the corresponding product formula for to  $r_{P_2|P_1}(\pi, \mathbf{s})$ , we extend (2.10) to all tempered representations  $\pi$  of M(F). Finally, if  $\pi$  is any irreducible unitary representation of M(F), it can be written as a Langlands quotient  $\pi = J_R^M(\tau, \mu)$ , where R is a parabolic subgroup of  $M, \tau$  is a tempered representation of  $M_R(F)$  and  $\mu$  is a point in the chamber of  $\mathfrak{a}_R^*/\mathfrak{a}_M^*$  attached to R. Then

$$r_{P_2|P_1}(\pi, \mathbf{s}) = r_{P_2(R)|P_1(R)}(\tau, \mathbf{s} + \mu)$$

and a similar formula holds for  $\tilde{r}_{P_2|P_1}(\pi, \mathbf{s})$  [A7, (2.3)]. Let  $\sigma_1, \sigma_2 \in S_r$  be attached to  $P_1, P_2$  such that  $\Sigma_{P_i}$  is given by (2.2). Then we get

**Lemma 2.1.** For all irreducible unitary representation  $\pi = \bigotimes_{i=1}^{r} \pi_i$  of M(F) we have

$$r_{P_2|P_1}(\pi, \mathbf{s}) = \prod_{(i,j)\in I(\sigma_1, \sigma_2)} \frac{\epsilon(0, \pi_i \times \widetilde{\pi}_j, \psi)}{\epsilon(s_i - s_j, \pi_i \times \widetilde{\pi}_j, \psi)} \widetilde{r}_{P_2|P_1}(\pi, \mathbf{s}), \quad \mathbf{s} \in \mathbb{C}^r.$$

Since we will be concerned with logarithmic derivatives of normalizing factors, we need estimates for  $f(\pi_1 \times \pi_2)$ . Let  $f(\pi_i)$  be the conductor of  $\pi_i$ , i = 1, 2. Then by Theorem 1 of [BH] and Corollary (6.5) of [BHK] we have

(2.11) 
$$0 \le f(\pi_1 \times \pi_2) \le n_1 f(\pi_1) + n_2 f(\pi_2) - \inf\{f(\pi_1), f(\pi_2)\}$$

for all admissible smooth representations  $\pi_i$  of  $\operatorname{GL}_{n_i}(F)$ , i = 1, 2. Furthermore, by Corollary (6.5) of [BHK], we have  $f(\pi_1 \times \pi_2) = 0$  if and only if there exists a quasicharacter  $\chi$  of  $F^{\times}$  such that both  $\pi_1 \otimes \chi \circ \det$  and  $\pi_2 \otimes \chi^{-1} \circ \det$  are unramified principal series representations.

By (2.11) it suffices to estimate the conductors  $f(\pi_i)$ , i = 1, 2.

Given an open compact subgroup  $K \subset \operatorname{GL}_m(F)$ , let

$$\Pi(\operatorname{GL}_m(F);K) = \left\{ \pi \in \Pi(\operatorname{GL}_m(F)) \mid \pi^K \neq \{0\} \right\}.$$

**Lemma 2.2.** For every open compact subgroup K of  $GL_m(F)$  there exists C > 0 such that  $f(\pi) \leq C$  for all  $\pi \in \Pi(GL_m(F); K)$ .

*Proof.* First we reduce the proof to the case of supercuspidal representations. Let  $\pi \in \Pi(\operatorname{GL}_m(F))$ . Then there exist a parabolic subgroup Pof  $\operatorname{GL}_m$  of type  $(m_1, ..., m_r)$ , tempered representations  $\tau_j$  of  $\operatorname{GL}_{m_j}(F)$ and real numbers  $t_i$  with  $t_1 > \cdots > t_r$  such that  $\pi$  is isomorphic to the Langlands quotient  $J_P^{\operatorname{GL}_m}(\tau_1[t_1], ..., \tau_r[t_r])$ . Then by Theorem 3.4 of [J] it follows that

$$f(\pi) = f\left(I_P^{\operatorname{GL}_m}(\tau, \mathbf{s})\right) = \sum_j f(\tau_j).$$

Thus we may assume that  $\pi$  is a tempered representation of  $\operatorname{GL}_m(F)$ . A tempered representation is full induced  $\pi = I_P^G(\sigma_1, ..., \sigma_r)$  where  $\sigma_i$  is a square-integrable representation of  $\operatorname{GL}_{m_i}(F)$ . By (3.2.3) of [J] we get

$$f(\pi) = \sum_{j} f(\sigma_j),$$

which reduces the problem to the square-integrable case. Let  $\pi$  be square-integrable. By [BZ] there is a divisor d|m, a standard parabolic subgroup P of type (d, ..., d) and a unitary supercuspidal representation  $\rho$  of  $\operatorname{GL}_d(F)$  so that  $\pi$  is the unique square-integrable constituent of  $\xi = I_P^{GL_m}(\rho_1, ..., \rho_r)$ , where r = m/d and  $\rho_j = \rho \otimes |\det|^{j-(r+1)/2}$ , j =1, ..., r. By [J, (2.7.4)] there exist polynomials  $P, \widetilde{P} \in \mathbb{C}[X]$  such that

(2.12) 
$$L(s,\pi) = P(q^{-s})L(s,\xi), \quad L(s,\widetilde{\pi}) = \widetilde{P}(q^{-s})L(s,\widetilde{\xi}).$$

Moreover by (2.7.5) of [J] we have

(2.13) 
$$\epsilon(s,\pi,\psi) = \epsilon(s,\xi,\psi) \frac{P(q^{-s})}{\widetilde{P}(q^{-1+s})}.$$

Assume that  $L(s, \rho) = 1$ . Then by Proposition 3.1.3 of [J] we have  $L(s, \pi) = 1$  and  $L(s, \xi) = 1$ . Thus

$$\epsilon(s,\pi,\psi) = \epsilon(s,\xi,\psi) = \prod_{i=1}^r \epsilon(s,\rho_i,\psi).$$

Hence

$$f(\pi) = f(\xi) = rf(\rho).$$

If  $L(s, \rho) \neq 1$ , then d = 1 and  $\rho = |\det|^t$ . Then  $L(s, \rho) = (1 - q^{-s-t})^{-1}$ and it follows from the first equality of (2.12) that

$$P(X) = \prod_{i=1}^{m-1} (1 - q^{-1/2(m-1)-t+i}X)$$

Using (2.7.6) of [J], it follows that  $P(q^{-s})/\tilde{P}(q^{-1+s}) = Cq^{-(m-1)s}$ . By (2.13) we get

$$f(\pi) = mf(\rho) + m - 1.$$

It remains to relate the K-invariant subspaces. We may assume that  $K \subset \operatorname{GL}_m(\mathcal{O}_F)$  is a congruence subgroup. Set  $\sigma = \bigotimes_j \rho_j$ . Using induction by stages, we can assume that  $\pi$  is a subquotient of  $I_P^{\operatorname{GL}_m}(\sigma)$ . Hence, if  $\pi^K \neq \{0\}$ , then  $I_P^{\operatorname{GL}_m}(\sigma)^K \neq \{0\}$ . Furthermore we have

$$I_P^{\mathrm{GL}_m}(\sigma)^K = \left(I_{\mathrm{GL}_m(\mathcal{O}_F)\cap P}^{\mathrm{GL}_m(\mathcal{O}_F)}(\sigma)\right)^K \\ \hookrightarrow \bigoplus_{\mathrm{GL}_m(\mathcal{O}_F)/K} I_{K\cap P}^K(\sigma)^K \\ \cong \bigoplus_{\mathrm{GL}_m(\mathcal{O}_F)/K} \sigma^{K\cap P}.$$

Now observe that

$$K \cap P = \prod_{i=1}^{r} K_i,$$

where  $K_i \subset \operatorname{GL}_{m_i}(\mathcal{O}_F)$  are congruence subgroups. Then

$$\sigma^{K\cap P} \cong \otimes_{i=1}^r \rho_i^{K_i}.$$

Thus if  $\pi^{K} \neq \{0\}$ , then  $\rho_{i}^{K_{i}} \neq \{0\}$  for all  $i, 1 \leq i \leq r$ . Combined with the above computation of the conductors, if follows that it suffices to consider supercuspidal representations.

Let  $L = K \cdot Z(F)$ . Then L is an open subgroup of  $\operatorname{GL}_m(F)$  which is compact modulo Z(F). By the Corollary to Theorem 2 in [Bu] there exist only finitely many irreducible supercuspidal representations  $\pi$  of  $\operatorname{GL}_m(F)$  such that  $\pi^L \neq 0$ . In other words, modulo twists with unramified characters, there exist only finitely many irreducible supercuspidal representations  $\pi$  of  $\operatorname{GL}_m(F)$  such that  $\pi^K \neq 0$ . Since the conductor remains unchanged under twists with unramified characters, the lemma follows.  $\Box$ 

Finally we observe that at an infinite place v, the  $\epsilon$ -factors are defined in terms of the  $\epsilon$ -factors attached to the representation of the Weil group  $W_{E_v}$  by means of the Langlands correspondence. The  $\epsilon$ -factors of a semisimple representation of the Weil groups  $W_{\mathbb{C}}$  and  $W_{\mathbb{R}}$  are described in [Sh4, p.990]. In all cases they are of the form  $i^a$  for some integer *a*. Therefore, in the Archimedean case there exists  $b \in \mathbb{N}_0$  such that

$$\epsilon(s, \pi_1 \times \pi_2, \psi) = i^b$$

Especially, the  $\epsilon$ -factors are independent of s.

#### 3. Estimation of the continuous parameter

For the unramified places, the continuous parameters of local components of cuspidal automorphic representations of  $\operatorname{GL}_n(\mathbb{A})$  have been estimated by Luo, Rudnick and Sarnak [LRS]. The main purpose of this section is to extend the estimations of [LRS] first to ramified places and then to local components of automorphic representations in the residual spectrum of  $\operatorname{GL}_n(\mathbb{A})$ . To deal with cuspidal automorphic representations we follow the method of [LRS] which uses properties of the Rankin-Selberg *L*-functions. To this end we have to describe the local Rankin-Selberg *L*-factors. First note that any local component of a cuspidal automorphic representation of  $\operatorname{GL}_n$  is generic [Sk]. Let *F* be a local field. By [JS3], any irreducible unitary generic representation  $\pi$ of  $\operatorname{GL}_n(F)$  is equivalent to a fully induced representation

$$\pi = I_P^G(\sigma, \mathbf{s}),$$

where P is a standard parabolic subgroup of type  $(n_1, ..., n_r)$ ,  $\mathbf{s} \in \mathbb{C}^r$ and  $\sigma$  is a square integrable representation of  $M_P(F)$ . To describe the Rankin-Selberg *L*-factors, we distinguish three cases according to the type of the field F.

## **1.** $F = \mathbb{R}$ .

We note that  $\operatorname{GL}_m(\mathbb{R})$  does not have square-integrable representations if  $m \geq 3$ . Therefore we can assume that  $\pi$  is isomorphic to a fully induced representation  $I_P^G(\sigma, \mathbf{s})$ , where P is of type (2, ..., 2, 1..., 1) with 2 occurring r times, and  $\sigma$  is a product of r discrete series representations  $D_{k_i}$  of  $\operatorname{GL}_2(\mathbb{R})$  and n - 2r characters  $\chi_i$  of  $\mathbb{Z}/2\mathbb{Z}$ . Here  $D_{k_i}$  denotes the discrete series representation with the same infinitesimal character as the  $k_i$  dimensional representation. Next note that an irreducible induced representation  $I_P^G(\sigma, \mathbf{s})$  is unitary, if it is equivalent to its hermitian dual representation  $I_P^G(\sigma, \bar{\mathbf{s}})$ , i.e if there exists an intertwining operator

$$J_{\bar{P}|P}(\sigma,\bar{s}): I_P^G(\sigma,\mathbf{s}) \to I_{\bar{P}}^G(\sigma,\bar{\mathbf{s}}).$$

By [KZ] this implies that there exists a  $w \in W(a_M)$  of order 2 so that

$$(D_{k_1}[s_1] \otimes \dots D_{k_r}[s_r] \otimes \chi_{r+1}[s_{r+1}] \otimes \dots \chi_{n-r}[s_{n-r}])^w = = D_{k_1}[-\bar{s_1}] \otimes \dots D_{k_r}[-\bar{s_r}] \otimes \chi_{r+1}[-\bar{s_{r+1}}] \otimes \dots \chi_{n-r}[-\bar{s_{n-r}}]$$

So either

- a)  $s_i$  is imaginary, or
- b)  $\operatorname{Re}(s_j) > 0, \ j \leq r$ , and there exists  $k \leq r$  so that  $D_j = D_k$  and  $s_j = -\bar{s}_k$ , or
- c)  $\operatorname{Re}(s_j) > 0$ , j > r, and there exists k > 0 so that  $\chi_j = \chi_k$  and  $s_j = -\bar{s}_k$ .

If  $\chi_j$  is a nontrivial character, then  $\operatorname{Re}(s_j) = 0$  since in this case there are no complementary series representations.

The Rankin-Selberg local *L*-factors are defined in terms of *L*-factors attached to semisimple representations of the Weil group  $W_{\mathbb{R}}$  by means of the Langlands correspondence [L3]. If  $\tau$  is a semisimple representation of  $W_{\mathbb{R}}$  of degree *m* and  $\pi(\tau)$  is the associated irreducible admissible representation of  $\mathrm{GL}_m(\mathbb{R})$ , then

$$L(s,\pi(\tau)) = L(s,\tau).$$

Furthermore, if

$$\tau = \bigoplus_{1 \le j \le m} \tau_j$$

is the decomposition into irreducible representations of  $W_{\mathbb{R}}$ , then

$$L(s,\tau) = \prod_{j} L(s,\tau_j)$$

If  $\tau'$  is another semisimple representation of  $W_{\mathbb{R}}$  of degree m' and  $\pi(\tau')$  is the associated irreducible admissible representation of  $\operatorname{GL}_{m'}(\mathbb{R})$ , then the Rankin-Selberg local *L*-factor is given by

$$L(s, \pi(\tau) \times \pi(\tau')) = L(s, \tau \otimes \tau').$$

This reduces the computation of the L-factors to the case of irreducible representations of the Weil group.

The irreducible representations of the Weil group  $W_{\mathbb{R}}$  of  $\mathbb{R}$  are either 1 or 2 dimensional. To describe the *L*-factors, we define Gamma factors by

$$\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma(s/2), \quad \Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s).$$

Suppose that  $\tau$  is a two-dimensional irreducible representation of  $W_{\mathbb{R}}$ . Then

$$\tau = \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}} \theta,$$

where  $\theta$  is a (not necessarily unitary) character of  $W_{\mathbb{C}} = \mathbb{C}^*$ . Thus there exist  $t \in \mathbb{C}$  and  $n \in \mathbb{Z}$  such that

$$\theta(z) = |z|^t (z/\overline{z})^{n/2}, \quad z \in \mathbb{C}^*.$$

Then the L-factor is defined as

$$L(s,\tau) = \Gamma_{\mathbb{C}}(s+t+|n|/2).$$

The one-dimensional irreducible representations of  $W_{\mathbb{R}}$  are of the form

$$\psi_{\epsilon,t}: (z,\sigma) \in W_{\mathbb{R}} \to \operatorname{sign}^{\epsilon}(\sigma)|z|^t$$

where  $\epsilon = 0, 1$  and "sign" is the sign character of the Galois group. The *L*-factor of  $\psi_{\epsilon,t}$  is given by

$$L(s, \psi_{\epsilon,t}) = \Gamma_{\mathbb{R}}(s+t+\epsilon).$$

Next we have to consider the tensor products of irreducible representations.

If  $\tau = \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}} \theta$  and  $\tau' = \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}} \theta'$  are two-dimensional representations of  $W_{\mathbb{R}}$ , then we have

$$\begin{aligned} \tau \otimes \tau' &= \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}(\theta \otimes \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}\theta'|_{\mathbb{C}^*}) \\ &= \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}(\theta \otimes \theta') \oplus \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}(\theta \otimes \theta'^{\sigma}), \end{aligned}$$

where  $\sigma$  is the nontrivial element of the Galois group. Suppose that  $\theta'(z) = |z|^{t'} (z/\overline{z})^{n'/2}$ . Then we get

$$L(s, \tau \otimes \tau') = \Gamma_{\mathbb{C}}(s + t + t' + |n + n'|/2)\Gamma_{\mathbb{C}}(s + t + t' + |n - n'|/2).$$

Similarly, if  $\psi = \psi_{\epsilon,t'}$  is a one-dimensional representation, then we have

$$\tau \otimes \psi = \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}(\theta \otimes \psi|_{\mathbb{C}^*}),$$

and therefore, we get

$$L(s, \tau \otimes \psi) = \Gamma_{\mathbb{C}}(s + t + t' + |n|/2).$$

Finally, if  $\psi = \psi_{\epsilon,t}$  and  $\psi_{\epsilon',t'}$  are two one-dimensional representations of  $W_{\mathbb{R}}$ , then

$$L(s,\psi\otimes\psi')=\Gamma_{\mathbb{R}}(s+t+t'+\widetilde{\epsilon}),$$

where  $0 \leq \tilde{\epsilon} \leq 1$  and  $\tilde{\epsilon} = \epsilon + \epsilon' \mod 2$ .

Let  $\pi = I_P^G(\sigma, \mathbf{s})$  be an irreducible unitary principal series representation with  $\sigma$  as above, i.e.,

$$\sigma = \otimes_{i=r}^r D_{k_i} \otimes \otimes_{j=1}^{n-2r} \chi_j,$$

where the  $D_{k_i}$ 's are discrete series representations of  $\operatorname{GL}_2(\mathbb{R})$  and the  $\chi_j$ 's are unitary characters of  $\mathbb{R}^*$ . For i = 1, ..., r let  $\theta_i$  be the character of  $\mathbb{C}^*$  defined by  $\theta_i(z) = |z|^{s_i} (z/\overline{z})^{k_i/2}$  and set

$$\tau_i = \operatorname{ind}_{\mathbb{C}^*}^{W_{\mathbb{R}}}(\theta_i).$$

Put

$$\tau_{\pi} = \bigoplus_{i=1}^{r} \tau_i \oplus \bigoplus_{j=r+1}^{n-r} \psi_{\epsilon_j, s_j}.$$

Then  $\tau_{\pi}$  is the representation of the Weil group  $W_{\mathbb{R}}$  which is associated to  $\pi$ . Note that if  $\epsilon_k = 1$  then  $\operatorname{Re}(s_k) = 0$ . Then by the definition of the Rankin-Selberg convolution,  $L(s, \pi \times \tilde{\pi})$  is equal to

$$\prod_{i,j} L(s,\tau_i \otimes \tilde{\tau}_j) \cdot \prod_{i,l} L(s,\tau_i \otimes \tilde{\psi}_{\epsilon_l,s_l}) \cdot \prod_{i,l} L(s,\tilde{\tau}_i \otimes \psi_{\epsilon_l,s_l}) \\ \cdot \prod_{l,k} L(s,\psi_{\epsilon_l,s_l} \otimes \tilde{\psi}_{\epsilon_k,s_k})$$

Using the above description of the *L*-factors, it follows that  $L(s, \pi \times \widetilde{\pi})$  is a product of Gamma factors of the form

$$\Gamma_{\mathbb{C}}(s+s_i-s_j+\frac{|k_i-k_j|}{2})\cdot\Gamma_{\mathbb{C}}(s+s_i-s_j+\frac{|k_i+k_j|}{2})$$

or

$$\Gamma_{\mathbb{C}}(s+s_l-s_j+|k_j|/2)\Gamma_{\mathbb{C}}(s+s_j-s_l+|k_j|/2)$$

or

$$\Gamma_{\mathbb{R}}((s+s_l-s_k+\epsilon_{l,k})),$$

where  $0 \leq \epsilon_{l,k} \leq 1$  with  $\epsilon_{k,l} = \epsilon_l + \epsilon_k \mod 2$  and  $1 \leq i, j \leq r, r+1 \leq l, k \leq n-r$ . We conclude that the local *L*-factor  $L(s, \pi \times \tilde{\pi})$  contains for every index  $i \leq r$  with  $\operatorname{Re}(s_i) \neq 0$  a Gamma factor

 $\Gamma(s+2\operatorname{Re}(s_i)),$ 

and for every index k > r with  $\operatorname{Re}(s_k) \neq 0$  a Gamma factor

$$\Gamma((s+2\operatorname{Re}(s_k))/2).$$

Since the Gamma function  $\Gamma(s)$  has no zeros, it follows that  $L(s, \pi \times \tilde{\pi})$  has its first pole at  $s = 2 \max |\operatorname{Re}(s_i)|$ .

**2.**  $F = \mathbb{C}$ .

Suppose that  $\pi = I_P^G(\sigma, \mathbf{s})$  is a irreducible unitary representation of  $\operatorname{GL}_n(\mathbb{C})$  with a Whittaker model. Then P is the Borel subgroup and  $\sigma$  is a product of characters  $\chi_i$  where  $\chi_i(z) = (\frac{z}{|z|})^{r_i}$ ,  $r_i \in \mathbb{Z}$ . As in the real case we see that the assumption that  $\pi$  is unitary implies that the

real part of the coordinate  $s_i$  of the continuous parameter **s** is smaller than 1/2 and that for every *i* with  $\operatorname{Re}(s_i) \neq 0$  there exists a *k* so that  $\chi_i = \chi_k$  and  $s_i = -\bar{s}_k$ .

The Weil group  $W_{\mathbb{C}}$  is equal to  $\mathbb{C}^*$  and the Rankin-Selberg local L-factor can be determined in the same way as in the real case. It follows that

$$L(s, \pi \times \widetilde{\pi}) = \prod_{i,j=1}^{n} \Gamma_{\mathbb{C}} \left( s + s_i - s_j + \frac{|r_i - r_j|}{2} \right).$$

As in the real case we deduce that if  $\pi = I_P^G(\sigma, \mathbf{s})$  is an irreducible unitary principal series representation of  $\operatorname{GL}_n(\mathbb{C})$ , then the *L*-factor  $L(s, \pi \times \tilde{\pi})$  contains for every index *i* with  $\operatorname{Re}(s_i) \neq 0$  the function

$$\Gamma(s+2\operatorname{Re}(s_i))$$

as a factor.

#### **3.** F non-Archimedean.

Any irreducible unitary generic representation  $\pi$  of  $\operatorname{GL}_n(F)$  is isomorphic to an induced representation

$$\pi \cong I_P^G(\sigma, \mathbf{s}),$$

where P is defined by a partition  $(n_1, n_2, \ldots, n_h)$  with  $n_{i+1} \leq n_i$  and  $\sigma = \bigotimes_i \sigma_i$  for discrete series representations  $\sigma_i$  of  $GL_{n_i}(F)$ . By [JPS] the Rankin-Selberg *L*-factor is given by

$$L(s, \pi \times \widetilde{\pi}) = \prod_{i,j=1}^{\prime} L(s + s_i - s_j, \sigma_i \times \widetilde{\sigma}_j).$$

As in the real case we conclude that if  $\operatorname{Re}(s_i) > 0$ , then there exists k so that  $\sigma_i = \sigma_k$  and  $s_i = -\bar{s}_k$ . Thus if  $\operatorname{Re}(s_i) > 0$  then the Rankin-Selberg L-factor  $L(s, \pi \times \tilde{\pi})$  contains  $L(s - 2\operatorname{Re}(s_i), \sigma_i \times \tilde{\sigma}_i)$  as a factor.

It remains to deal with the *L*-functions for square-integrable representations. Let  $\pi$  be a square-integrable representation of  $\operatorname{GL}_m(F)$ . By [BZ] there is a divisor d|m, a standard parabolic subgroup P of  $\operatorname{GL}_m(F)$  of type  $(d, \ldots, d)$ , and an irreducible supercuspidal representation  $\rho$  of  $\operatorname{GL}_d(F)$  so that  $\pi$  is the unique quasi-square-integrable component of the induced representation  $I_P^G(\rho_1, \ldots, \rho_r)$ , where r = m/d and  $\rho_j = \rho \otimes |\det|^{j-(n+1)/2}$ . We denote this representation by  $\pi = \Delta(r, \rho)$ . The representation  $\pi$  is unitary (or equivalently square-integrable) if only if  $\rho$  in unitary. Moreover the contragredient of  $\Delta(r, \rho)$  is given by

 $\Delta(r,\rho) = \Delta(r,\tilde{\rho})$ . Let  $\sigma = \Delta(r,\rho)$  be a square integrable representation of  $\operatorname{GL}_m(F)$ . Then by Theorem 8.2 of [JPS] we have

$$L(s, \sigma \times \widetilde{\sigma}) = \prod_{j=1}^{r} L(s+r-j, \rho \times \widetilde{\rho}).$$

Thus the description of the Rankin-Selberg *L*-functions is reduced to the case of two supercuspidal representations. Let  $\rho_i$ , i = 1, 2, be supercuspidal representations of  $\operatorname{GL}_{k_i}(F)$ . By Proposition (8.1) of [JPS] we have  $L(s, \rho_1 \times \rho_2) = 1$  if  $k_1 > k_2$ . Since  $L(s, \rho_1 \times \rho_2) = L(s, \rho_2 \times \rho_1)$ , the same holds for  $k_1 < k_2$ . Let  $k_1 = k_2$ . Then by Proposition (8.1) of [JPS],  $L(s, \rho_1 \times \rho_2) = 1$ , unless  $\rho_1$  and  $\rho_2$  are in the same twist class, i.e., there exists  $t \in \mathbb{C}$  such that  $\rho_2 \cong \rho_1[t]$ . In this case we have

$$L(s,\rho_1\times\widetilde{\rho}_1[t]) = L(s+t,\rho_1\times\widetilde{\rho}_1) = \left(1-q^{-a(t+s)}\right)^{-1},$$

where  $a|k_1$  is the order of the cyclic group of unramified characters  $\chi = |\det|^u$  such that  $\rho_1 \otimes \chi \cong \rho_1$ . Let  $\sigma_i = \Delta(r_i, \rho_i)$ . Then we get

$$L(s+2\operatorname{Re}(s_i), \sigma_i \times \widetilde{\sigma}_i) = \prod_{j=1}^{r_i} L(s+2\operatorname{Re}(s_i)+r_i-j, \rho \times \widetilde{\rho})$$
$$= \prod_{j=1}^{r_i} \left(1-q^{-a_i(s+2\operatorname{Re}(s_i)+r_i-j)}\right)^{-1},$$

where  $a_i$  is the order of the cyclic group of unramified characters  $\chi$  so that  $\rho_i \otimes \chi$  is isomorphic to  $\rho_i$ . Thus we conclude that for an irreducible unitary principal series representation  $\pi = I_P^G(\sigma, \mathbf{s})$  of  $GL_n(F)$  and for every index i with  $\operatorname{Re} s_i > 0$  there exists a factor of the *L*-function  $L(s, \pi \times \tilde{\pi})$  with a pole at  $s = -2 \operatorname{Re}(s_i)$ .

Next we recall some facts about ray class characters. Let E be a number field. Let  $\mathfrak{q}$  be a nonzero integral ideal of E and denote by  $C(\mathfrak{q})$  the wide ray class group of E modulo  $\mathfrak{q}$ . We note that the term "wide" means that no positivity condition has been imposed at the real places of E. Then a character of  $C(\mathfrak{q})$  is unramified at all infinite places. Now recall that any character  $\chi$  of  $C(\mathfrak{q})$  can be identified with a character of the idele class group  $C_E = I_E/E^{\times}$  which is trivial on the congruence subgroup  $C_E^{\mathfrak{q}} = I_E^{\mathfrak{q}} E^{\times}/E^{\times}$  [Neu]. Then  $\chi = \bigotimes_v \chi_v$  where  $\chi_v$  is a character of  $E_v^{\times}$  which is unramified at all places  $v \mid \infty$  and all finite places  $\mathfrak{p} \nmid \mathfrak{q}$ . Furthermore for a finite place  $\mathfrak{p} \nmid \mathfrak{q}$ , the character  $\chi_{\mathfrak{p}}$ is given by

(3.1) 
$$\chi_{\mathfrak{p}}(\alpha) = \chi(\mathfrak{p})^{v_{\mathfrak{p}}(\alpha)}, \quad \alpha \in E_{\mathfrak{p}}^{\times},$$

where  $v_{\mathfrak{p}}: E_{\mathfrak{p}}^{\times} \to \mathbb{Z}$  is the  $\mathfrak{p}$ -adic valuation.

Let S be any finite set of finite places of E. For an integral ideal  $\mathfrak{q}$  of E let  $X_{\mathfrak{q}}$  denote the set of all wide ray class characters of conductor  $\mathfrak{q}$ such that  $\chi(\mathfrak{p}) = 1$  for all  $\mathfrak{p} \in S$ . Therefore we have  $(\mathfrak{p}, \mathfrak{q}) = 1$  for all  $\mathfrak{p} \in S$ . By (3.1) it follows that

(3.2) 
$$\chi_v = 1 \quad \text{for all } v \in S \cup S_\infty.$$

Let  $X_{\mathfrak{q}}^*$  be the subset of  $X_{\mathfrak{q}}$  consisting of all primitive characters. Set  $X = \bigcup_{\mathfrak{q}} X_{\mathfrak{q}}$  and  $X^* = \bigcup_{\mathfrak{q}} X_{\mathfrak{q}}^*$ . For  $\chi \in X_{\mathfrak{q}}^*$  and a cuspidal automorphic representation  $\pi$  of  $\operatorname{GL}_n$ , the partial Rankin-Selberg *L*-function  $L_S(s, (\pi \otimes \chi) \times \widetilde{\pi})$  is defined to be

$$L_S(s, (\pi \otimes \chi) \times \widetilde{\pi}) = \prod_{v \notin S} L(s, (\pi_v \otimes \chi_v) \times \widetilde{\pi}_v).$$

We shall use the following result of Luo, Rudnick and Sarnak which is the main result of [LRS].

**Theorem 3.1.** Given  $n, \pi, S$  as the above and any  $\beta > 1 - 2/(n^2 + 1)$ , there are infinitely many  $\chi \in X^*$  such that

$$L_S(\beta, (\pi \otimes \chi) \times \widetilde{\pi}) \neq 0.$$

Now we can establish our extension of Theorem 2 of [LRS].

**Proposition 3.2.** Suppose that  $\pi = \bigotimes_v \pi_v$  is a cuspidal automorphic representation of  $GL_n(\mathbb{A})$ . Let  $s_v = (s_{1,v}, \ldots, s_{k,v})$  be the continuous parameters of the representation  $\pi_v$ . Then we have

$$\max_{j} |\operatorname{Re}(s_{j,v})| < \frac{1}{2} - \frac{1}{n^2 + 1}.$$

*Proof.* We follow the proof of Theorem 2 in [LRS]. Let v be a place of E and set  $S = \{v\}$ . Let  $\chi \in X^*$ , where  $X^*$  is the set of ray class characters with respect to S which we defined above. The Rankin-Selberg L-function

$$L(s, (\pi \otimes \chi) \times \widetilde{\pi}) = L(s, (\pi_v \otimes \chi_v) \times \widetilde{\pi}_v) L_S(s, (\pi \otimes \chi) \times \widetilde{\pi}).$$

is holomorphic in the whole complex plane except for simple poles at s = 1 and s = 0 if  $\chi = 1$ . This follows from the work of Jacquet, Piateski-Shapiro, Shalika, Shahidi, Mœglin and Waldspurger [JPS], [JS1], [JS2], [MW], [Sh2]. By Theorem 3.1 we may choose  $\chi \in X^*$  such that  $L(s, (\pi \otimes \chi) \times \tilde{\pi})$  is an entire function and by (3.2) we have  $\chi_v = 1$ . Suppose that  $s_0$ ,  $\operatorname{Re}(s_0) > 0$ , is a pole of

$$L(s, \pi_v \times \widetilde{\pi}_v) = L(s, (\pi_v \otimes \chi_v) \times \widetilde{\pi}_v).$$

Then  $s_0$  must be a zero of  $L_S(s, (\pi \otimes \chi) \times \tilde{\pi})$ . Now suppose that  $\operatorname{Re}(s_0) > 1 - 2(1 + n^2)^{-1}$ , then by Theorem 3.1 there exists a non-trivial  $\chi \in X^*$  with  $L_S(s_0, (\pi \otimes \chi) \times \tilde{\pi}) \neq 0$  and  $\chi_v = 1$ . Hence  $L(s, \pi_v \times \tilde{\pi}_v)$  is holomorphic for  $s \geq 1 - 2(1 + n^2)^{-1}$  and therefore we get  $\operatorname{Re}(s_0) < 1 - 2(1 + n^2)^{-1}$ .

Suppose first that v is a real place and let  $\pi_v$  be the local component of  $\pi$  at v. Then  $\pi_v$  is generic and therefore, isomorphic to a fully induced representation  $I_P^G(\sigma_v, \mathbf{s})$  for some discrete series representation  $\sigma_v$  of  $M_P(\mathbb{R})$ . Suppose that  $\operatorname{Re}(s_i) < 0$ . Using the above results on the local Rankin-Selberg *L*-factors, we conclude that  $L(s, \pi_v \times \tilde{\pi}_v)$  contains as a factor  $\Gamma(s + 2\operatorname{Re}(s_i))$  or  $\Gamma((s + 2\operatorname{Re}(s_i))/2)$  and the other factors have no zeros. Thus  $s_0 = -2\operatorname{Re}(s_i)$  is a pole of this *L*-factor. Since  $\operatorname{Re}(s_i) < 0$  we deduce that  $s_0 \leq 1 - 2(n^2 + 1)^{-1}$  and thus  $-\operatorname{Re}(s_i) < 1/2 - 1(n^2 + 1)^{-1}$ . Since  $\mathbf{s}$  is the continuous parameter of a unitary representation,  $-\bar{s_i}$  is also a coordinate of  $\mathbf{s}$ . Since  $\Gamma(s - 2\operatorname{Re}(s_i))$  or  $\Gamma((s - 2\operatorname{Re}(s_i))/2)$  is also a factor of the *L*-factor of the Rankin *L*-function by the same argument  $\operatorname{Re}(s_i) < 1/2 - (n^2 + 1)^{-1}$ . Thus we conclude that  $|\operatorname{Re}(s_i)| < 1/2 - (n^2 + 1)^{-1}$ .

Next suppose that v is a complex place and  $\pi_v = I_P^G(\sigma_v, \mathbf{s})$ . Suppose that  $\operatorname{Re}(s_i) < 0$ . The local Rankin-Selberg *L*-function contains factors  $\Gamma(s+2\operatorname{Re}(s_i))$  and  $\Gamma(s-2\operatorname{Re}(s_i))$ . Thus the previous argument shows that  $|\operatorname{Re}(s_i)| < 1/2 - (n^2 + 1)^{-1}$  holds in this case too.

Finally consider the case of a finite place v. As we have seen above, for every coordinate  $s_i$  of the continuous parameter of  $\pi_v = I_P^G(\tau_v, \mathbf{s})$  the local Rankin-Selberg *L*-function contains a factor with a pole at  $-2 \operatorname{Re}(s_i)$  and the remaining factors have no zeros. Thus as before we conclude that  $|\operatorname{Re}(s_i)| < 1/2 - (n^2 + 1)^{-1}$ .

We shall now establish a similar result for the local components of residual automorphic representations of  $\operatorname{GL}_n(\mathbb{A})$ . First we recall some facts about representations of  $\operatorname{GL}_n$  over a local field F. Any irreducible unitary representation  $\pi$  of  $\operatorname{GL}_n(F)$  is equivalent to a Langlands quotient  $J_R^G(\tau,\mu)$ . This is the unique irreducible quotient of an induced representation  $I_R^G(\tau,\mu)$  where  $\tau$  is a tempered representation of  $M_R(F)$  and  $\mu$  is a point in the positive chamber attached to R. A slight variant of this description is as follows. Recall that a tempered representation  $\tau$  of  $\operatorname{GL}_m(E_v)$  can be described as follows. There exist a standard parabolic subgroup Q of type  $(m_1, ..., m_p)$  and square-integrable representations  $\delta_j$  of  $\operatorname{GL}_{m_j}(F)$  so that  $\tau$  is isomorphic to the full induced representation  $I_P^{\operatorname{GL}_m}(\delta_1, ..., \delta_p)$ . Hence by induction in stages, there exist a standard parabolic subgroup P of G of type  $(n_1, ..., n_r)$ , discrete series representations  $\delta_i$  of  $\operatorname{GL}_{n_i}(F)$  and real numbers  $s_1 \geq s_2 \geq \cdots \geq s_r$  such that  $\pi$ is equivalent to the unique irreducible quotient

$$J_P^G(\delta_1[s_1] \otimes \cdots \otimes \delta_r[s_r]),$$

of the induced representation  $I_P^G(\delta_1[s_1] \otimes \cdots \otimes \delta_r[s_r])$  [MW, I.2]. We call  $s_1, \ldots, s_r$  the (continuous) Langlands parameters of  $\pi$ .

The residual spectrum for  $\operatorname{GL}_n(\mathbb{A})$  has been determined by Mæglin and Waldspurger [MW]. Let  $\pi = \bigotimes_v \pi_v$  be an irreducible automorphic representation in the residual spectrum of  $\operatorname{GL}_n(\mathbb{A})$ . By [MW], there is a divisor k|n, a standard parabolic subgroup P of type  $(k, \ldots, k)$ , and a cuspidal automorphic representation  $\xi$  of  $\operatorname{GL}_d(\mathbb{A})$ , d = n/k, so that the representation  $\pi$  is a quotient of the induced representation

$$I_{P(\mathbb{A})}^{G(\mathbb{A})}\left(\xi[(k-1)/2]\otimes\cdots\otimes\xi[-(k-1)/2]\right).$$

**Lemma 3.3.** Let  $\pi$  and  $\xi$  be as above. Let v be a place of E and let  $s_1 \geq \cdots \geq s_r$ , be the Langlands parameters of  $\xi_v$ . Then the Langlands parameters of  $\pi_v$  are given by

$$\left(\frac{k-1}{2}+s_1,...,\frac{k-1}{2}+s_r,...,-\frac{k-1}{2}+s_1,...,-\frac{k-1}{2}+s_r\right).$$

*Proof.* Since  $\xi_v$  is a local component of a cuspidal automorphic representation  $\xi$  of  $\operatorname{GL}_d(\mathbb{A})$ , it is generic. Using induction in stages, it follows that there exist a standard parabolic subgroup R of  $\operatorname{GL}_d$  of type  $(n_1, \ldots, n_r)$ , discrete series representations  $\delta_{i,v}$  of  $\operatorname{GL}_{n_i}(E_v)$  and real numbers  $(s_1, \ldots, s_r)$  satisfying

(3.3) 
$$s_1 \ge s_2 \ge \cdots \ge s_r; \quad |s_j| < 1/2, \ j = 1, \dots, r$$

such that  $\xi_v$  is isomorphic to the full induced representation

$$\xi_v \cong I_R^{\mathrm{GL}_d}(\delta_{1,v}[s_1] \otimes \cdots \otimes \delta_{r,v}[s_r]).$$

Let  $Q = M_Q N_Q$  be the standard parabolic subgroup of  $GL_n$  whose Levi component  $M_Q$  is a product of k copies of  $M_R$  and let

$$\delta_v = (\delta_{1,v} \otimes \cdots \otimes \delta_{r,v}) \otimes \cdots \otimes (\delta_{1,v} \otimes \cdots \otimes \delta_{r,v}),$$

where  $(\delta_{1,v} \otimes \cdots \otimes \delta_{r,v})$  occurs k times. Define

$$\mu(k,\mathbf{s}) = \left(\frac{k-1}{2} + s_1, \dots, \frac{k-1}{2} + s_r, \frac{k-3}{2} + s_1, \dots, -\frac{k-1}{2} + s_r\right).$$

By induction in stages we have

$$I_P^G(\xi_v[(k-1)/2] \otimes \cdots \otimes \xi_v[-(k-1)/2]) = I_Q^G(\delta_v, \mu(k, \mathbf{s})).$$

Furthermore, by (3.3) the coordinates of  $\mu(k, \mathbf{s})$  are decreasing. Thus the induced representation  $I_Q^G(\delta_v, \mu(k, \mathbf{s}))$  has a unique irreducible quotient which must be isomorphic to  $\pi_v$ .

Next we recall a different method to parametrize irreducible unitary representations of  $\operatorname{GL}_n(F)$ . Let d|n and k = n/d. Let P be the standard parabolic subgroup of type (k, ..., k). Let  $\delta$  be a discrete series representation of  $\operatorname{GL}_d(F)$  and let  $a, b \in \mathbb{R}$  be such that  $b-a \in \mathbb{N}$ . Then the induced representation

$$I_P^G(\delta[b] \otimes \delta[b-1] \otimes \cdots \otimes \delta[a])$$

has a unique irreducible quotient which we denote by  $J(\delta, a, b)$ . Especially, if a = -(k-1)/2 and b = (k-1)/2, then we put

(3.4) 
$$J(\delta, k) := J(\delta, a, b).$$

By Theorem D of [Ta] and [Vo], for every irreducible unitary representation of  $\operatorname{GL}_n(F)$  there exist a standard parabolic subgroup P of type  $(n_1, \ldots, n_r)$ ,  $k_i | n_i$ , discrete series representations  $\delta_i$  of  $\operatorname{GL}_{d_i}(F)$ ,  $d_i = n_i/k_i$ , and real numbers  $s_1, \ldots, s_r$  with  $|s_i| < 1/2$ ,  $i = 1, \ldots, r$ , such that  $\pi$  is isomorphic to the fully induced representation:

(3.5) 
$$\pi \cong I_P^G(J(\delta_1, k_1)[s_1] \otimes \cdots \otimes J(\delta_r, k_r)[s_r]).$$

Using this parametrization, we get the following analogue to Proposition 3.2 for local components of automorphic representations in the residual spectrum of  $\operatorname{GL}_n(\mathbb{A})$ .

**Proposition 3.4.** Let  $\pi_v$  be a local component of an automorphic representation in the residual spectrum of  $\operatorname{GL}_n(\mathbb{A})$ . There exist k|n, a parabolic subgroup P of type  $(kn_1, \ldots, kn_r)$ , discrete series representations  $\delta_{i,v}$  of  $\operatorname{GL}_{n_i}(E_v)$  and real numbers  $s_1, \ldots, s_r$  satisfying

$$s_1 \ge s_2 \ge \dots \ge s_r, \quad |s_i| < 1/2 - (1+n^2)^{-1}, \quad i = 1, \dots, r,$$

such that

$$\pi_v \cong I_P^G(J(\delta_{1,v},k)[s_1] \otimes \cdots \otimes J(\delta_{r,v},k)[s_r]).$$

*Proof.* By the proof of Lemma 3.3,  $\pi_v$  is equivalent to a Langlands quotient of the form

$$\pi_v \cong J_Q^G(\delta_v, \mu(k, \mathbf{s})),$$

where the parameters  $\mathbf{s}$  satisfy (3.3). Set

$$b_i = \frac{k-1}{2} + s_i, \qquad a_i = -\frac{k-1}{2} + s_i, \quad i = 1, ..., r.$$

Suppose there exist  $1 \leq i < j \leq r$  such that the triples  $(\delta_{i,v}, a_i, b_i)$  and  $(\delta_{j,v}, a_j, b_j)$  are linked in the sense of I.6.3 or I.7 in [MW]. Suppose that  $s_i \geq s_j$ . Then it follows from (2) and (3)(i) on p.622 or from (1) and (2) on p.624 of [MW] that  $a_i \geq a_j + 1$  and  $b_i \geq b_j + 1$ . This implies  $1 \leq |s_i - s_j|$  which contradicts (3.3). Hence the triples  $(\delta_{i,v}, a_i, b_i)$  are pairwise not linked. Now observe that

$$J(\delta_{i,v}, a_i, b_i) = J(\delta_{i,v}, k)[s_i].$$

Let P be the standard parabolic subgroup with Levi component

$$\operatorname{GL}_{kn_1} \times \cdots \times \operatorname{GL}_{kn_r}$$
.

Let  $\widetilde{\delta}_v = \otimes_{i=1}^r \delta_{i,v}$  and set

$$J(\delta_v, k) := \bigotimes_{i=1}^r J(\delta_{i,v}, k)$$

Then it follows from Proposition I.9 of [MW] that the induced representation

$$I_P^G(J(\widetilde{\delta}_v,k),\mathbf{s}) := I_P^G(J(\delta_{1,v},k)[s_1] \otimes \cdots \otimes J(\delta_{r,v},k)[s_r])$$

is irreducible. By Lemma I.8, (ii), of [MW],  $I_P^G(J(\tilde{\delta}_v, k), \mathbf{s})$  is a quotient of the induced representation  $I_Q^G(\delta_v, \mu(k, \mathbf{s}))$ . Since  $I_P^G(J(\tilde{\delta}_v, k), \mathbf{s})$  is irreducible, this quotient must be the Langlands quotient. Thus

$$\pi_v \cong J_Q^G(\delta_v, \mu(k, \mathbf{s})) \cong I_P^G(J(\delta_{1,v}, k)[s_1] \otimes \cdots \otimes J(\delta_{r,v}, k)[s_r]).$$

By construction, the  $s_i$ 's are the Langlands parameters of a local component of a cuspidal automorphic representation. Therefore it follows from Proposition 3.2 that they satisfy

$$|s_i| < \frac{1}{2} - \frac{1}{n^2 + 1}, \quad i = 1, ..., r.$$

### 4. Normalized local intertwining operators

The main purpose of this section is to prove Proposition 0.2. To this end we have to establish some auxiliary results. We continue with the investigation of the parametrization (3.5) of irreducible unitary representations of  $\operatorname{GL}_m(F)$ , where F is a local field of characteristic zero.

Assume that F is Archimedean. Let  $J(\delta; k)$  be as in (3.4). Let  $K \subset \operatorname{GL}_m(F)$  be the standard maximal compact subgroup, i.e., K = O(m), if  $F = \mathbb{R}$ , and K = U(m), if  $F = \mathbb{C}$ . If  $\pi$  is an irreducible representation of  $\operatorname{GL}_m(F)$ , then we denote by  $\chi_{\pi}$  its central character.

**Lemma 4.1.** Let  $\sigma \in \Pi(K)$ . The set of all representations  $J(\delta, k)$  with  $\chi_{\delta} = 1$  and  $[J(\delta, k)|_{K} : \sigma] > 0$  is finite.

Proof. Let

$$I(\delta,k) = I_P^{\mathrm{GL}_m}(\delta[-(k-1)/2] \otimes \cdots \otimes \delta[(k-1)/2]).$$

Then by definition,  $J(\delta, k)$  is a quotient of  $I(\delta, k)$ . Therefore we have

(4.1)  $[J(\delta,k)|_K:\sigma] \le [I(\delta,k)|_K:\sigma].$ 

Let

$$K_M = K \cap M_P.$$

By [Kn, p.208] we have

(4.2) 
$$[I(\delta,k)|_K:\sigma] = \sum_{\omega\in\Pi(K_M)} \left[ (\otimes_{i=0}^{k-1}\delta[(k-1)/2-i]) \Big|_{K_M}:\omega \right] \cdot [\sigma|_{K_M}:\omega]$$

Now observe that

$$K_M = \prod_{i=1}^k \widetilde{K},$$

where  $\widetilde{K} \subset \operatorname{GL}_d(F)$  is the standard maximal compact subgroup. Thus any  $\omega \in \Pi(K_M)$  has the form

$$\omega = \otimes_i \omega_i$$

with  $\omega_i \in \Pi(\widetilde{K})$ , and therefore, we get

(4.3) 
$$\left[ \left( \otimes_{i=0}^{k-1} \delta[(k-1)/2 - i] \right) \Big|_{K_M} : \omega \right] = \prod_{i=1}^m [\delta|_{\tilde{K}} : \omega_i].$$

Suppose that  $\chi_{\delta} = 1$ . Then  $\delta$  can be identified with a discrete series representation of  $\mathrm{SL}_d(F)$ . The set of all discrete series representations of  $\mathrm{SL}_d(F)$  which contain a given  $\widetilde{K}$ -type is finite [Wa2, p.398]. Together with (4.1) - (4.3) this implies the lemma.

Now we return to the case where F is an arbitrary local field of characteristic zero. Fix a maximal compact subgroup K of  $GL_m(F)$ . Let

$$M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$$
.

Let  $k_i | n_i, d_i = n_i / k_i$  and  $\delta_i$  a discrete series representation of  $\operatorname{GL}_{d_i}(F)$ . Set

$$\pi = \bigotimes_{i=1}^r J(\delta_i, k_i).$$

**Lemma 4.2.** For all  $P, Q \in \mathcal{P}(M)$ , the normalized local intertwining operator  $R_{Q|P}(\pi, \mathbf{s})$  is holomorphic for all  $\mathbf{s} \in \mathbb{C}^r$  with  $|\operatorname{Re}(s_i - s_j)| < 1$ ,  $1 \leq i, j \leq r$ .

*Proof.* Let  $P_1, ..., P_N$  be a minimal chain of parabolic subgroups in  $\mathcal{P}(M)$  connecting P and Q. Then each pair  $(P_{l+1}, P_l)$  of consecutive parabolics in this chain is associated with adjacent Weyl chambers and it determines a unique simple root  $\alpha_l \in \Delta_{P_l}$  with  $-\alpha_l \in \Delta_{P_{l+1}}$ . Applying  $(R_2)$  of Theorem 2.1 of [A7] repeatedly, we get

$$R_{Q|P}(\pi, \mathbf{s}) = R_{P_N|P_{N-1}}(\pi, \mathbf{s}) \cdots R_{P_3|P_2}(\pi, \mathbf{s}) R_{P_2|P_1}(\pi, \mathbf{s}).$$

So it suffices to consider  $R_{P_{l+1}|P_l}(\pi, \mathbf{s})$  for  $1 \leq l \leq N-1$ . Let  $\alpha_l \in \Delta_{P_l}$  be the simple root determined by this pair. Define  $M'_l$  to be the group in  $\mathcal{L}(M)$  such that

$$\mathfrak{a}_{M_l'} = \left\{ H \in \mathfrak{a}_M \mid \alpha_l(H) = 0 \right\}.$$

Then dim $(\mathfrak{a}_M/\mathfrak{a}_{M'_l}) = 1$ . Let  $R_l$  be the unique group in  $\mathcal{P}^{M'_l}(M)$  whose simple root is  $\alpha_l$ . Let  $\phi \in \mathcal{H}_{P_l}(\pi_v)$ . Then for any  $k \in K_v$ , the function

 $\phi_k(m') = \phi(m'k), \quad m' \in M'_l(E_v),$ 

belongs to  $\mathcal{H}_{R_l}(\pi_v)$  and by Theorem 2.1 of [A7] we get

$$\left(R_{P_{l+1}|P_l}(\pi, \mathbf{s})\phi\right)_k = R_{\overline{R}_l|R_l}(\pi, \mathbf{s})(\phi_k).$$

Moreover, the root  $\alpha_l \in \Sigma_P$  determines a pair (i, j) of integers  $1 \leq i < j \leq r$  so that  $R_{\overline{R}_l|R_l}(\pi_v, \mathbf{s})$  depends only on  $s_i - s_j$ . Thus we can assume that P is maximal parabolic. Then r = 2 and

$$\pi = J(\delta_1, k_1) \otimes J(\delta_2, k_2).$$

We may regard the intertwining operator as a function  $R_{\overline{P}|P}(\pi, s)$  of one complex variable s. Let  $s \in \mathbb{C}$  and set  $s_1 = s/2$ ,  $s_2 = -s/2$ . Define

(4.4) 
$$\widetilde{\delta}_{j} = \delta_{j}[i \operatorname{Im}(s_{j})], \quad b_{j} = \frac{k_{j} - 1}{2} + \operatorname{Re}(s_{j}), \\ a_{j} = -\frac{k_{j} - 1}{2} + \operatorname{Re}(s_{j}), \quad j = 1, 2.$$

Assume that  $|\operatorname{Re}(s)| < 1$ . Then as in the proof of Proposition 3.4, it follows that the triples  $(\tilde{\delta}_1, a_1, b_1)$  and  $(\tilde{\delta}_2, a_2, b_2)$  are not linked in the sense of I.6.3 (*F* non-Archimedean) or I.7 (*F* archimedean) in [MW]. Therefore by Lemma I.6.3 or Lemma I.7, in [MW] the intertwining operator

$$R_{\overline{P}|P}(J(\delta_1, a_1, b_1) \otimes J(\delta_2, a_2, b_2), u)$$

is holomorphic at u = 0. Since

$$J(\delta_j, k_j)[s_j] = J(\widetilde{\delta}_j, a_j, b_j).$$

it follows that  $R_{\overline{P}|P}(\pi_v, s)$  is holomorphic in the strip  $|\operatorname{Re}(s)| < 1$ .  $\Box$ 

Let 
$$\pi \in \Pi(\operatorname{GL}_m(F))$$
. Then by (3.5) we have  $\pi \cong I_P^G(J, \mathbf{s})$ , where  
 $J = \bigotimes_{i=1}^r J(\delta_i, k_i), \quad \mathbf{s} = (s_1, \dots, s_r)$ 

with  $|s_i| < 1/2, i = 1, ..., r$ . Given  $\sigma \in S_r$ , let

$$M^{\sigma} = \operatorname{GL}_{\sigma(1)} \times \cdots \times \operatorname{GL}_{\sigma(r)}$$

and  $P^{\sigma}$  the standard parabolic subgroup with Levi component  $M^{\sigma}$ . Let  $\sigma$  act on J and s in the obvious way. It follows from Lemma 4.2 that

$$I_P^G(J, \mathbf{s}) \cong I_{\sigma P \sigma^{-1}}^G(\sigma J, \sigma \mathbf{s}) \cong I_{P^\sigma}^G(\sigma J, \sigma \mathbf{s}).$$

So we can assume that the  $s_i$ 's satisfy  $s_1 \ge s_2 \ge \cdots \ge s_r$ . Furthermore by Theorem D of [Ta] and [Vo] we have

$$(s_1, ..., s_r) = -(s_r, ..., s_1)$$
, and  $\delta_i \cong \delta_{r-i}$ ,  $k_i = k_{r-i}$ , if  $s_i > 0$ .

Let  $w \in S_r$  be the longest element. Then it follows that

(4.5) 
$$wJ = J, \quad ws = -s, \quad wP = \overline{P}$$

By Theorem D of [Ta] and [Vo], each representation  $J(\delta_i, k_i)$  is unitarizable. Hence J is an irreducible unitary representation of M(F). By our assumption,  $I_P^G(J, \mathbf{s})$  is unitarizable and we shall now describe the corresponding inner product in the space of K-finite vectors  $\mathcal{H}_P^0(J)$  in the Hilbert space  $\mathcal{H}_P(J)$  of the induced representation. Identify w with some element in K and let  $\ell(w)$  be the left translation by w. By  $(R_4)$  of Theorem 2.1 of [A7] and (4.5) we have

$$R_{\overline{P}|P}(J, \mathbf{s})^* = J_{P|\overline{P}}(J, -s) = J_{w\overline{P}|wP}(wJ, w\mathbf{s})$$
$$= \ell(w)R_{\overline{P}|P}(J, \mathbf{s})\ell(w)^{-1}$$

Set

$$B = \ell(w) R_{\overline{P}|P}(J, \mathbf{s}).$$

Then B is Hermitian and satisfies

(4.6) 
$$I_P^G(J, -\mathbf{s})B = BI_P^G(J, \mathbf{s})$$

Now one can verify as in [KZ, p.148] that  $(B\Phi, \Phi)_{L^2(K)}$  is positive or negative semidefinite in  $\mathcal{H}^0_P(J)$ . By Lemma 4.2 both  $R_{\overline{P}|P}(J, \mathbf{s})$  and  $R_{P|\overline{P}}(J, \mathbf{s})$  are defined and

$$R_{\overline{P}|P}(J,\mathbf{s})R_{P|\overline{P}}(J,\mathbf{s}) = \mathrm{Id}, \quad R_{P|\overline{P}}(J,\mathbf{s})R_{\overline{P}|P}(J,\mathbf{s}) = \mathrm{Id}$$

hold in  $\mathcal{H}^0_P(J)$ . Hence  $R_{\overline{P}|P}(J, \mathbf{s})$  is an isomorphism on K-finite vectors so that  $(B\Phi, \Phi)_{L^2(K)}$  is positive or negative definite. Let  $\varepsilon \in \{\pm 1\}$  be such that  $\varepsilon(B\Phi, \Phi)_{L^2(K)}$  is positive definite. Then we define an inner product in  $\mathcal{H}^0_P(J)$  by

(4.7) 
$$\langle \Phi, \Psi \rangle = \varepsilon(\ell(w) R_{\overline{P}|P}(J, \mathbf{s}) \Phi, \Psi)_{L^2(K)}, \quad \Phi, \Psi \in \mathcal{H}^0_P(J).$$

It follows as in [KZ, p.148] that

$$\langle I_P^G(J,\mathbf{s})\Phi,\Psi\rangle + \langle\Phi,I_P^G(J,\mathbf{s})\Psi\rangle = 0, \quad \Phi,\Psi\in\mathcal{H}_P^0(J),$$

where  $I_P^G(J, \mathbf{s})$  denotes the infinitesimal induced representation.

Let M be a standard Levi subgroup of type  $(n_1, ..., n_r)$  and let  $P, P' \in \mathcal{P}(M)$ . Given a place v of E, let  $\Pi_{\text{disc}}(M(E_v))$  denote the set of all  $\pi_v \in \Pi(M(E_v))$  which are local components of some automorphic representation  $\pi$  in the discrete spectrum of  $M(\mathbb{A})$ . Without loss of generality, we may assume that P is a standard parabolic subgroup. Let  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$ . Then  $\pi_v = \otimes_i \pi_{i,v}$  with  $\pi_{i,v} \in \Pi_{\text{disc}}(\text{GL}_{n_i}(E_v))$ , i = 1, ..., r. By Proposition 3.4 there exist a standard parabolic subgroup  $R_i$  of  $\text{GL}_{n_i}$  with

$$M_{R_i} = \operatorname{GL}_{n_{i1}} \times \cdots \times \operatorname{GL}_{n_{im_i}},$$

 $k_{ij}|n_{ij}$ , discrete series representations  $\delta_{ij}$  of  $\operatorname{GL}_{d_{ij}}(E_v)$ ,  $d_{ij} = n_{ij}/k_{ij}$ , and  $s_{ij} \in \mathbb{R}$  satisfying

(4.8) 
$$s_{i1} \ge \dots \ge s_{im_i}, \quad |s_{ij}| < \frac{1}{2} - \frac{1}{n^2 + 1},$$

such that

$$\pi_{i,v} \cong I_{R_i}^{\operatorname{GL}_{n_i}} \left( J(\delta_{i1}, k_{i1})[s_{i1}] \otimes \cdots \otimes J(\delta_{im_i}, k_{im_i})[s_{im_i}] \right).$$

Set  $R = \prod_i R_i$ . Then R is a standard parabolic subgroup of M. Put  $m = m_1 + \cdots + m_r$ . We identify  $\{(i, j) \mid i = 1, ..., r, j = 1, ..., m_i\}$  with  $\{1, ..., m\}$  by

$$(i,j)\mapsto \sum_{k< i}m_k+j.$$

For  $1 \leq l \leq m$  let (i, j) be the pair that corresponds to l. Put

$$\delta_l = \delta_{ij}, \quad k_l = k_{ij}, \quad s_l = s_{ij}.$$

Set

(4.9) 
$$J_{\pi_v} = \bigotimes_{i=1}^m J(\delta_i, k_i), \quad \mathbf{s}_{\pi_v} = (s_1, ..., s_m).$$

Combing the above equivalences, we get

$$\pi_v \cong I_R^M(J_{\pi_v}, \mathbf{s}_{\pi_v}).$$

Let P(R) and P'(R) be the parabolic subgroups of  $\operatorname{GL}_n$  with  $P(R) \subset P$ ,  $P'(R) \subset P'$ ,  $P(R) \cap M = R$  and  $P'(R) \cap M = R$ . Then by induction in stages, the induced representation  $I_P^G(\pi_v, \mathbf{s})$  is unitarily equivalent to the induced representation  $I_{P(R)}^G(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v})$ . The unitary map  $\mu$ which provides this equivalence, is given by the evaluation map as in [KS, p.31]. Here we identify  $\mathbf{s} \in \mathbb{C}^r$  with an element in  $\mathbb{C}^m$  by

$$(s_1, ..., s_r) \mapsto (s_1, ..., s_1, s_2, ..., s_2, ..., s_r, ..., s_r),$$

where  $s_i$  is repeated  $m_i$  times. [KS, p.31].

**Lemma 4.3.** Let  $\mu$  be the unitary equivalence of the induced represntations as above. Then we have

(4.10) 
$$\mu \circ R_{P'|P}(\pi_v, \mathbf{s}) = R_{P'(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v}) \circ \mu.$$

Proof. First consider the unnormalized intertwining operators. Recall that the unnormalized intertwining operators are defined by integrals which are absolutely convergent in a certain shifted chamber [KS, Theorem 6.6], [Sh2]. If we compare these integrals in their range of convergence as in [KS, p.31], it follows immediately that (4.10) holds for the unnormalized intertwining operators. So it remains to consider the normalizing factors. If we apply Lemma 3.3 to each component  $\pi_{i,v}$  of  $\pi_v$ , it follows that  $\pi_v$  is equivalent to a Langlands quotient of the form  $J_Q^M(\delta_v, \mu(\mathbf{k}, \mathbf{s}_{\pi_v}))$ , where  $\delta_v = \bigotimes_{i=1}^m \delta_i$  and  $\mu(\mathbf{k}, \mathbf{s}_{\pi_v}) = (\mu(k_1, \mathbf{s}_{\pi_{1,v}}), ..., \mu(k_r, \mathbf{s}_{\pi_{r,v}}))$ . Hence by (2.3) in [A7] we get

$$r_{P'|P}(\pi_v, \mathbf{s}) = r_{P'(Q)|P(Q)}(\delta_v, \mathbf{s} + \mu(\mathbf{k}, \mathbf{s}_{\pi_v})).$$

Let  $S_i$  be the standard parabolic subgroup of  $M_{R_i}$  with Levi component

$$(\operatorname{GL}_{d_{i1}} \times \cdots \times \operatorname{GL}_{d_{i1}}) \times \cdots \times (\operatorname{GL}_{d_{ip_i}} \times \cdots \times \operatorname{GL}_{d_{ip_i}})$$

where each factor  $\operatorname{GL}_{d_{ij}}$  occurs  $k_i$  times. Let  $S = \prod_i S_i$ . Then  $M_S$  and  $M_Q$  are conjugate. Let  $w \in S_m$  be the element which conjugates  $M_S$  into  $M_Q$ . Referring again to (2.3) in [A7], we get

 $r_{P'(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v}) = r_{P'(R(S))|P(R(S))}(w\delta_v, \mathbf{s} + w\mu(\mathbf{k}, \mathbf{s}_{\pi_v})).$ 

By (r.6) of [A8, p.172] it follows that

 $r_{P'(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v}) = r_{P'(R(S)^w)|P(R(S)^w)}(\delta_v, \mathbf{s} + \mu(\mathbf{k}, \mathbf{s}_{\pi_v})),$ 

where  $R(S)^w = w^{-1}R(S)w$ . Finally note that P(Q), P'(Q),  $P(R(S)^w)$  and  $P'(R(S)^w)$  have the same Levi component and the reduced roots satisfy

$$\Sigma^{\underline{r}}_{\underline{P'(R(S)^w)}} \cap \Sigma^{\underline{r}}_{P(R(S)^w)} = \Sigma^{\underline{r}}_{\underline{P'(Q)}} \cap \Sigma^{\underline{r}}_{P(Q)}.$$

Using the product formula (r.1) in [A8, p.171], it follows that

$$r_{P'(R(S)^w)|P(R(S)^w)}(\delta_v, \mathbf{s} + \mu(\mathbf{k}, \mathbf{s}_{\pi_v})) = r_{P'(Q)|P(Q)}(\delta_v, \mathbf{s} + \mu(\mathbf{k}, \mathbf{s}_{\pi_v})).$$

Combining the above equations, we get

$$r_{P'|P}(\pi_v, \mathbf{s}) = r_{P'(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v}),$$

and this finishes the proof of the lemma.

## Proof of Proposition 0.2:

Let M be a standard Levi subgroup of type  $(n_1, ..., n_r)$  and let  $P, Q \in \mathcal{P}(M)$ . We distinguish between the Archimedean and non-Archimedean case.

Case 1:  $v \mid \infty$ .

Let  $\pi_v \in \Pi(M(E_v))$ . Let  $K_v$  be the standard maximal compact subgroup of  $\operatorname{GL}_n(E_v)$ . Given  $\sigma_v \in \Pi(K_v)$ , denote by  $R_{Q|P}(\pi_v, \mathbf{s})_{\sigma_v}$  the restriction of  $R_{Q|P}(\pi_v, \mathbf{s})$  to the  $\sigma_v$ -isotypical subspace  $\mathcal{H}_P(\pi_v)_{\sigma_v}$  of the Hilbert space  $\mathcal{H}_P(\pi_v)$  of the induced representation and define  $R_{Q(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v})_{\sigma_v}$  similarly. Our goal is to estimate the norm of the derivatives of the operator  $R_{Q|P}(\pi_v, \mathbf{s})_{\sigma_v}$ . By (4.10) we have

$$\mu \circ R_{Q|P}(\pi_v, \mathbf{s})_{\sigma_v} = R_{Q(R)|P(R)}(J_{\pi_v}, \mathbf{s} + \mathbf{s}_{\pi_v})_{\sigma_v} \circ \mu,$$

so that it suffices to estimate the norm of the derivatives of the operator on the right hand side. To this end we have to compare the inner products on the spaces of  $K_v$ -finite vectors  $\mathcal{H}^0_P(\pi_v)$  and  $\mathcal{H}^0_{P(R)}(J_{\pi_v})$ . Let  $(\cdot, \cdot)$ denote the inner product in  $\mathcal{H}^0_{P(R)}(J)$  of the unitarily induced representation. By (4.7) it follows that the infinitesimal action of the induced representation  $I^G_P(\pi_v)$  on  $\mathcal{H}^0_P(\pi_v)$  is unitarily equivalent to the infinitesimal action of the induced representation  $I^G_{P(R)}(J_{\pi_v}, \mathbf{s}_{\pi_v})$  on  $\mathcal{H}^0_{P(R)}(J_{\pi_v})$ under the new inner product

$$\langle \Phi, \Psi \rangle = \varepsilon(\ell(w) R_{P(\overline{R})|P(R)}(J_{\pi_v}, \mathbf{s}_{\pi_v}) \Phi, \Psi)_{L^2(K)}, \quad \Phi, \Psi \in \mathcal{H}^0_{P(R)}(J_{\pi_v}).$$

Let  $\| \cdot \|$  denote the norm in  $\mathcal{H}^0_{P(R)}(J)$  with respect to  $\langle \cdot, \cdot \rangle$  and  $\| \cdot \|$ the norm with respect to  $(\cdot, \cdot)$ . Then for  $\Phi \in \mathcal{H}_{P(R)}(J)_{\sigma_v}$  and  $\alpha \in \mathbb{N}^r_0$ we have

$$\| D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_{v}, i\mathbf{u})_{\sigma_{v}} \Phi \|^{2} \leq \| R_{Q(\overline{R})|Q(R)}(J_{\pi_{v}}, \mathbf{s}_{\pi})_{\sigma_{v}} \|$$
  
 
$$\cdot \| D_{\mathbf{u}}^{\alpha} R_{Q(R)|P(R)}(J_{\pi_{v}}, i\mathbf{u} + \mathbf{s}_{\pi_{v}})_{\sigma_{v}} \|^{2} \| \Phi \|^{2} .$$

Let  $B = \varepsilon \ell(w) R_{P(\overline{R})|P(R)}(J_{\pi_v}, \mathbf{s}_{\pi_v})$ . As we have seen above, B is Hermitian and positive definite. Hence  $\|| \Phi |\|^2 = (B\Phi, \Phi) = \| B^{1/2}\Phi \|^2$ and therefore, we get  $\| \Phi \|^2 \le \| B^{-1/2} \|^2 \| \| \Phi \|\|^2$ . Now observe that by Lemma 4.2 and  $(R_2)$  of Theorem 2.1 of [A7], we have

$$B^{-1} = \varepsilon R_{P(R)|P(\overline{R})}(J_{\pi_v}, \mathbf{s}_{\pi_v})\ell(w)^{-1}.$$

This implies

(4.11) 
$$\begin{split} \| D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_{v}, i\mathbf{u})_{\sigma_{v}} \|^{2} \\ \leq \| R_{Q(\overline{R})|Q(R)}(J_{\pi_{v}}, \mathbf{s}_{\pi})_{\sigma_{v}} \| \| R_{P(R)|P(\overline{R})}(J_{\pi_{v}}, \mathbf{s}_{\pi})_{\sigma_{v}} \| \\ \cdot \| D_{\mathbf{u}}^{\alpha} R_{Q(R)|P(R)}(J_{\pi_{v}}, i\mathbf{u} + \mathbf{s}_{\pi_{v}})_{\sigma_{v}} \|^{2} . \end{split}$$

Note that by Lemma 4.2 the intertwining operators on the right hand side are all well defined. This leads to the first reduction of the problem. We can assume that  $\pi_v$  is of the form

(4.12) 
$$\pi_v = \bigotimes_{i=1}^r J(\delta_i, k_i),$$

but we have to estimate derivatives of  $R_{Q|P}(\pi_v, \mathbf{s})_{\sigma_v}$  for  $\mathbf{s}$  in the domain

(4.13) 
$$\Omega = \left\{ \mathbf{s} \in \mathbb{C}^r \mid |\operatorname{Re}(s_i)| \le 1/2 - (n^2 + 1)^{-1}, \ i = 1, ..., r \right\}.$$

Furthermore, since  $J(\delta_j[s], k_j) = J(\delta_j, k_j)[s]$ , we can also assume that the  $\delta_j$ 's have trivial central character. Let

$$K_{M,v} = K_v \cap M(E_v).$$

Then  $K_{M,v}$  is a maximal compact subgroup of  $M(E_v)$ . It follows from Frobenius reciprocity [Kn, p.208] that

(4.14) 
$$[I_P^G(\pi_v, \mathbf{s})|_{K_v} : \sigma_v] = \sum_{\tau \in \Pi(K_{M,v})} [\pi_v|_{K_{M,v}} : \tau] \cdot [\sigma_v|_{K_{M,v}} : \tau].$$

Let

$$U(\sigma_v) = \{ \tau \in \Pi(K_{M,v}) \mid [\sigma_v|_{K_{M,v}} : \tau] > 0 \}.$$

This is a finite set. It follows from (4.14) that  $[I_P^G(\pi_v, \mathbf{s})|_{K_v} : \sigma_v] > 0$ , implies that there exists  $\tau \in U(\sigma_v)$  such that  $[\pi_v|_{K_{M,v}} : \tau] > 0$ .

Now observe that

$$K_{M,v} = \prod_{i=1}^{\prime} K_{i,v},$$

where  $K_{i,v} \subset \operatorname{GL}_{n_i}(E_v)$  is the standard maximal compact subgroup. Let  $\tau \in \Pi(K_{M,v})$ . Then  $\tau = \bigotimes_i \tau_i$  with  $\tau_i \in \Pi(K_{i,v})$  and

$$[\pi_v|_{K_{M,v}}:\tau] = \prod_{i=1}^r [J(\delta_i, k_i)|_{K_{i,v}}:\tau_i]$$

By Lemma 4.1 it follows that there are only finitely many  $\pi_v$  of the form (4.12) such that

$$[\pi_v|_{K_{M,v}}:\tau]>0 \text{ for some } \tau \in U(\sigma_v).$$

So we may fix  $\pi_v$ . Next we apply the product formula for normalized intertwining operators in the same way as in the proof of Lemma 4.2

to reduce the problem to the case where P is maximal parabolic. Then the intertwining operator may be regarded as a function  $R_{\overline{P}|P}(\pi_v, s)$  of one complex variable s and by Lemma 4.2,  $R_{\overline{P}|P}(\pi_v, s)$  is holomorphic in the strip  $|\operatorname{Re}(s)| < 1$ . Furthermore by  $(R_6)$  of Theorem 2.1 of [A7],  $R_{Q|P}(\pi_v, s)_{\sigma_v}$  is a rational function of s. This implies immediately that for every  $k \in \mathbb{N}$  there exist C > 0 and  $N \in \mathbb{N}$  such that

$$\left\|\frac{d^k}{ds^k}R_{Q|P}(\pi_v,s)_{\sigma_v}\right\| \le C(1+|s|)^N$$

for all s in the strip  $|\operatorname{Re}(s)| < 1 - 2/(n^2 + 1)$ .

Let  $\xi_{\pi_v}$  denote the infinitesimal character of  $\pi_v$ . Combining the various steps of the reduction, it follows that for every  $\alpha \in \mathbb{N}^r$  there exist C > 0 and  $N \in \mathbb{N}$  such that

(4.15) 
$$\| D_{\mathbf{w}}^{\alpha} R_{Q|P}(\pi_{v}, i\mathbf{w})_{\sigma_{v}} \| \leq C(1+\|\xi_{\pi_{v}}\|+\|\mathbf{w}\|)^{N}$$

for all  $\mathbf{w} \in \mathbb{R}^r$  and  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$ .

It remains to estimate  $\| \xi_{\pi_v} \|$  in terms of  $|\lambda_{\pi_v}|$ . To this end note that it suffices to establish the corresponding estimate for  $GL_m(E_v)$ . Let  $\pi \in \Pi(\operatorname{GL}_m(E_v))$ . There exist a parabolic subgroup R of  $\operatorname{GL}_m$ of type  $(m_1, ..., m_r)$ , discrete series representations  $\tau_i$  of  $\operatorname{GL}_{m_i}(E_v)$  and  $s_1, \ldots, s_r \in \mathbb{C}$  satisfying  $\operatorname{Re}(s_1) \leq \operatorname{Re}(s_2) \leq \cdots \leq \operatorname{Re}(s_r)$  such that  $\pi$  is equivalent to the unique irreducible subrepresentation of the induced representation  $I_R^{\operatorname{GL}_m}(\tau_1[s_1], ..., \tau_r[s_r])$  [MW, p.609]. So it suffices to consider the induced representation  $I_R^{\operatorname{GL}_m}(\tau, \mathbf{s})$ , where  $\tau = \bigotimes_i \tau_i$  and  $\mathbf{s} = (s_1, ..., s_r)$ . Furthermore, we may assume that the central character of  $\tau$  is trivial and hence, we may regard  $\tau$  as a discrete series representation of  $M_R(E_v)^1$ . Let  $\mathfrak{m}_R^1$  denote the Lie algebra of  $M_R(E_v)^1$ and let  $\mathfrak{t} \subset \mathfrak{m}_R^1$  be a compact Cartan subalgebra. Then  $\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{a}_R$ is a Cartan subalgebra of  $\mathfrak{gl}_m(E_v)$ . Let  $\gamma: Z(\mathfrak{gl}_m(\mathbb{C})) \to I(\mathfrak{h})$  be the Harish-Chandra homorphism. Let  $\Lambda_{\tau} \in i\mathfrak{t}^*$  be the Harish-Chandra parameter of  $\tau$ . Then the infinitesimal character of  $I_R^{\operatorname{GL}_m}(\tau, \mathbf{s})$  corresponds to  $\Lambda_{\tau} + \mathbf{s} \in (\mathfrak{t} \oplus \mathfrak{a}_R)^*_{\mathbb{C}}$  [Kn, Proposition 8.22], and is given by  $\chi(Z) = (\Lambda_{\tau} + \mathbf{s})(\gamma(Z)), Z \in Z(\mathfrak{gl}_m(\mathbb{C}))$  [Kn, (8.32)]. Since  $\pi$  is an irreducible subrepresentation of  $I_R^{\operatorname{GL}_m}(\tau, \mathbf{s})$ , it has the same infinitesimal character. Therefore

$$\| \xi_{\pi} \|^2 = \| \Lambda_{\tau} \|^2 + \| \mathbf{s} \|^2.$$

Let  $H_1, ..., H_r$  be an orthonormal basis of  $\mathfrak{a}_R$  and  $H_{r+1}, ..., H_m$  an orthonormal basis of  $\mathfrak{t}$ . Then

$$\gamma(\Omega) = \sum_{i=1}^{r} H_i^2 - \sum_{j=r+1}^{m} H_j^2 - \| \rho \|^2$$

[Wa1, p.168]. Hence, the Casimir eigenvalue  $\lambda_{\pi}$  of  $\pi$  is given by

$$\lambda_{\pi} = (\Lambda_{\tau} + \mathbf{s})(\gamma(\Omega)) = \sum_{i=1}^{r} s_i^2 + \parallel \Lambda_{\tau} \parallel^2 - \parallel \rho \parallel^2$$

Since  $\pi$  is unitary, it follows from Theorem 3.3 of Chapter XI of [BW] that there exists C > 0, independent of  $\pi$ , such that  $\| \operatorname{Re}(\mathbf{s}) \| \leq C$ . Let  $K_v(m)$  be the standard maximal compact subgroup of  $\operatorname{GL}_m(E_v)$ and let  $\delta \in \Pi(K_v(m))$ . Suppose that  $\pi \in \Pi(\operatorname{GL}_m(E_v))$  is such that  $[\pi|_{K_v(m)} : \delta] > 0$ . Since  $\delta$  occurs in  $\pi$ , it also occurs in  $I_R^{\operatorname{GL}_m}(\tau, \mathbf{s})$ . Using Frobenius reciprocity as in [Kn, p.208], it follows that there exists  $\omega \in \Pi(K_v(m) \cap M_R(E_v))$  such that

$$[\tau|_{K_v(m)\cap M_R(E_v)}:\omega] > 0$$
 and  $[\delta|_{K_v(m)\cap M_R(E_v)}:\omega] > 0.$ 

Let  $\lambda_{\delta}$  and  $\lambda_{\omega}$  denote the Casimir eigenvalues of  $\delta$  and  $\omega$ , respectively. By [Mu2, (5.15)], the second inequality implies  $\lambda_{\omega} \leq \lambda_{\delta}$ . On the other hand, by [Wa2, p.398], the first inequality implies

$$\|\Lambda_{\tau}\|^{2} \leq \lambda_{\omega} + \|\rho_{R}\|.$$

Combining our estimations it follows that for every  $\delta \in \Pi(K_v(m))$  there exists C > 0 such that  $\| \xi_{\pi} \|^2 \leq C(1 + |\lambda_{\pi}|)$  for all  $\pi \in \Pi(\operatorname{GL}_m(E_v))$ satisfying  $[\pi|_{K_v(m)} : \delta] > 0$ . This implies that a similar result holds for  $M(E_v)$ , i.e., for every  $\delta_v \in \Pi(K_{M,v})$  there exists C > 0 such that

$$\| \xi_{\pi_v} \|^2 \leq C(1 + |\lambda_{\pi_v}|)$$

for all  $\pi_v \in \Pi(M(E_v))$  with  $[\pi_v|_{K_{M,v}} : \delta_v] > 0$ . Combined with (4.15) this completes the case of an Archimedean place.

## Case 2: $v < \infty$ .

Let  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$ . Let  $K_v$  be an open compact subgroup of  $\operatorname{GL}_n(E_v)$ . We may assume that  $K_v \subset \operatorname{GL}_n(\mathcal{O}_v)$  is a congruence subgroup. Let  $R_{Q|P}(\pi_v, \mathbf{s})_{K_v}$  denote the restriction of  $R_{Q|P}(\pi_v, \mathbf{s})$  to the  $K_v$ -invariant subspace in  $\mathcal{H}_P(\pi_v)$  and define  $R_{Q(R)|P(R)}(J_{\pi_v}, \mathbf{s})_{K_v}$  similarly. Now we proceed in exactly the same way as in the Archimedean case. Using (4.10) and the description (4.7) of the inner product, we get an estimation which is analogous to (4.11). From this estimation it follows immediately that it suffices to estimate the norm of derivatives of the operator  $R_{Q(R)|P(R)}(J_{\pi_v}, \mathbf{s})_{K_v}$  for  $\mathbf{s}$  in the domain (4.13). Thus we can assume that  $\pi_v$  is of the form

$$\pi_v = \otimes_{i=1}^r J(\delta_i, k_i),$$

but we have to estimate derivatives of  $R_{Q|P}(\pi_v, \mathbf{s})_{K_v}$  in the domain (4.13). Let **1** denote the trivial representation of  $K_v$ . Let

$$K_{M,v} = K_v \cap M(E_v).$$

This is again a congruence subgroup of  $M(\mathcal{O}_v)$ . As in the proof of Lemma 2.2 it follows that

(4.16) 
$$[I_P^G(\pi_v, \lambda)|_{K_v} : \mathbf{1}] \le \# [M(\mathcal{O}_v)/K_{M,v}] \cdot [\pi_v|_{K_{M,v}} : \mathbf{1}]$$

Now observe that

$$K_{M,v} = \prod_{r=1}^{l} K_{i,v},$$

where  $K_{i,v} \subset \operatorname{GL}_{n_i}(\mathcal{O}_v)$  is a congruence subgroup. Then

$$[\pi_v|_{K_{M,v}}:\mathbf{1}] = \prod_{i=1}^r [J(\delta_i, k_i)|_{K_{i,v}}:\mathbf{1}].$$

Consider an irreducible representation of  $\operatorname{GL}_m(E_v)$  of the form  $J(\delta, k)$ , , where k|m and  $\delta$  is a discrete series representation of  $\operatorname{GL}_d(E_v)$ , d = m/k. Let  $\widetilde{K}_v \subset \operatorname{GL}_m(\mathcal{O}_v)$  be a congruence subgroup. By the same reasoning as above we get

(4.17) 
$$\left[J(\delta,k)|_{\widetilde{K}_v}:\mathbf{1}\right] \leq [\delta|_{\widetilde{K}_v \cap \mathrm{GL}_d(E_v)}:\mathbf{1}]^k.$$

Again,  $\widetilde{K}_v \cap \operatorname{GL}_d(E_v)$  is a congruence subgroup of  $\operatorname{GL}_d(\mathcal{O}_v)$ , and therefore, it is an open compact subgroup of  $\operatorname{GL}_d(E_v)$ . Let  $\Pi_2(\operatorname{GL}_d(E_v), \widetilde{K}_v)$ denote the set of all discrete series representations  $\delta$  of  $\operatorname{GL}_d(E_v)$  such that  $[\delta|_{\widetilde{K}_v \cap \operatorname{GL}_d(E_v)} : 1] \geq 1$ . Then  $\Pi_2(\operatorname{GL}_d(E_v), \widetilde{K}_v)$  is a compact subset of  $\Pi_2(\operatorname{GL}_d(E_v))$  [HC2, Theorem 10].

Since v is finite, the stabilizer of  $\delta$  with respect to the action  $r \in \mathbb{R} \mapsto \delta[ir]$  in  $\Pi(\operatorname{GL}_d(E_v))$  is a lattice L and therefore, the orbit  $\mathfrak{o}_{\delta}$  of  $\delta$  is isomorphic to the circle  $\mathbb{R}/L$ . The orbit  $\mathfrak{o}_{\delta}$  is also open in  $\Pi_2(\operatorname{GL}_d(E_v))$ . Hence there exist  $\sigma_1, ..., \sigma_p \in \Pi_2(\operatorname{GL}_d(E_v), \widetilde{K}_v)$  such that

$$\Pi_2(\mathrm{GL}_d(E_v), K_v) = \mathfrak{o}_{\sigma_1} \sqcup \cdots \sqcup \mathfrak{o}_{\sigma_p}.$$

Next observe that for  $r \in \mathbb{R}$  we have

$$J(\delta[ir], k) = J(\delta, k)[ir].$$

Given  $\mathbf{u} = (u_1, ..., u_r) \in \mathbb{C}^r$ , let  $\pi_v[\mathbf{u}] = \bigotimes_i \pi_{i,v}[u_i]$ . Recall that the following equality holds

$$R_{Q|P}(\pi_v[\mathbf{u}], \mathbf{s}) = R_{Q|P}(\pi_v, \mathbf{s} + i\mathbf{u}), \quad \mathbf{u} \in \mathbb{C}^r.$$

Combining these observations it follows that it suffices to prove the lemma for fixed  $\pi_v$ . Next we apply the product formula for intertwining

operators in the same way as in the Archimedian case to reduce the problem further to the case where P is maximal parabolic and

$$\pi_v = J(\delta_1, k_1) \otimes J(\delta_2, k_2).$$

Then we may consider the intertwining operator  $R_{\overline{P}|P}(\pi_v, s)$  as a function of a single complex variable s. By Lemma 4.2,  $R_{\overline{P}|P}(\pi_v, s)$  is holomorphic in the strip  $|\operatorname{Re}(s)| < 1$ . Furthermore by Theorem 2.1 of [A7],  $R_{\overline{P}|P}(\pi_v, s)_{K_v}$  is a finite rank matrix whose coefficients are rational functions of  $q^{-s}$ . Therefore  $R_{\overline{P}|P}(\pi_v, u+iw)_{K_v}$  is a periodic function of w with period  $2\pi/\log q$ . Hence for every  $k \in \mathbb{N}$  there exists C > 0such that

$$\left|\frac{d^k}{ds^k}R_{\overline{P}|P}(\pi_v,s)_{K_v}\right| \le C$$

for all s in the strip  $|\operatorname{Re}(s) < 1 - 2/(n^2 + 1)$ . Combining all steps of the reduction, it follows that for every  $\alpha \in \mathbb{N}^r$  there exists C > 0 such that

$$\parallel D_{\mathbf{u}}^{\alpha} R_{Q|P}(\pi_v, i\mathbf{u})_{K_v} \parallel \leq C$$

for all  $\mathbf{u} \in \mathbb{R}^r$  and all  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$ . This completes the proof of Proposition 0.2.

**Remark.** As the above proof shows, the estimations hold for all generic representations  $\pi_v$  of  $M(E_v)$  whose continuous parameters  $s_1, ..., s_r$  satisfy a non-trivial bound of the form  $|s_i| < 1/2 - \varepsilon, \varepsilon > 0$ . In particular, they hold for tempered representations. For tempered representations and the real place v, Arthur [A8, Lemma 2.1] has established a refined estimation of the form (0.4). It holds for any reductive group G and is uniform in  $\sigma_v \in \Pi(K_v)$  and  $\pi_v \in \Pi_{\text{temp}}(M(\mathbb{R})/A_M(\mathbb{R}))$ . The right hand side contains in addition the Casimir eigenvalue of  $\sigma_v$ . Although we do not know how to prove it, we expect that the estimations can be made to be uniform in  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$  and  $\sigma_v \in \Pi(K_v)$ . On the other hand, as the following example shows, the estimations can not be expected to be uniform in all  $\pi_v \in \Pi(M(E_v))$ .

# Example.

Let  $G = \operatorname{GL}_4(\mathbb{R})$  and P the standard parabolic subgroup with  $M_P = \operatorname{GL}_2 \times \operatorname{GL}_2$ . Consider the representation  $I_P^G(\sigma \times \sigma, \mathbf{s})$  where  $\sigma$  is the spherical principal series representation induced from the character  $\mu = (\mu, -\mu), \mu$  real and  $0 \le \mu \le 1/2$  of the Borel subgroup of  $\operatorname{GL}_2(\mathbb{R})$ . We may assume that  $\mathbf{s} = (s, -s)$  with s real. Then

$$I_P^G(\sigma \times \sigma, \mathbf{s}) = I_B^G(\mu + s, -\mu + s, \mu - s, \mu - s).$$

For fixed  $0 \le \mu < 1/2$  this representation is irreducible for  $0 \le |s| < 1/2 - \mu$  and reducible for  $|s| = 1/2 - \mu$  [Sp]. The intertwining operator  $R_{\bar{P}|P}(\sigma \times \sigma, \mathbf{s})$  is therefore well defined on the interval  $0 \le |s| < 1/2 - \mu$  and has a pole for  $-s = 1/2 - \mu$ .

As the example shows, uniform estimates in  $\pi_v \in \Pi_{\text{disc}}(M(E_v))$  and  $\sigma \in \Pi(K_v)$  can only hold if the poles of the corresponding intertwining operators  $R_{Q|P}(\pi, \mathbf{s})$  are uniformly bounded away from the imaginary axis. We shall now show that this is indeed the case.

Let  $P = N_P M_P$  be a standard parabolic subgroup and let  $Q \in \mathcal{P}(M_P)$ . There exists  $w \in G(E)$  in the normalizer of  $M_P(E)$  so that  $wQw^{-1}$  is a standard parabolic subgroup with Levi subgroup  $M_P$ . Then for each place v of E and every  $\pi_v \in \Pi(M_P(E_v))$ , the map  $R_w : f(g) \to f(gw)$ defines an isomorphism

$$R_w: I_P^G(\pi_v, \mathbf{s}) \to I_Q^G(\pi_v^w, w(\mathbf{s})).$$

We refer to the operator

$$N(\pi_v, \mathbf{s}, w) = R_{P|Q}(\pi_v, \mathbf{s}) \circ R_w : I_P^G(\pi_v, \mathbf{s}) \to I_{wQw^{-1}}^G(\pi_v^w, w(\mathbf{s}))$$

also as a normalized intertwining operator. Since  $R_w$  is an isomorphism  $N(\pi_v, \mathbf{s}, w)$  has the same properties as  $R_{P|Q}(\pi_v, \mathbf{s})$ . We say that  $R_{Q|P}(\pi_v, \mathbf{s})$  has a pole at  $\mathbf{s}_0 \in \mathbb{C}^r$ , if  $R_{Q|P}(\pi_v, \mathbf{s})$  has a matrix coefficient with a pole at  $\mathbf{s}_0$ . Otherwise,  $R_{Q|P}(\pi_v, \mathbf{s})$  is called holomorphic in  $\mathbf{s}_0$ .

**Proposition 4.4.** Let  $M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$  be a standard Levi subgroup and let  $P, Q \in \mathcal{P}(M)$ . For every place v of E, there exists a constant  $B_v > 0$  such that for all  $\pi_v \in \prod_{\operatorname{disc}}(M(E_v))$ , the normalized intertwining operator  $R_{Q|P}(\pi_v, \mathbf{s})$  is holomorphic in the domain

$$\{\mathbf{s} \in \mathbb{C}^r \mid |\operatorname{Re}(s_i)| < B_v, \ i = 1, ..., r\}.$$

*Proof.* By the product formula for intertwining operators and induction by stages it suffices to prove the theorem for a maximal standard parabolic subgroup P of some  $\operatorname{GL}_m$  with m|n. Thus we may assume that P is a maximal standard parabolic subgroup of  $GL_m$  of type  $(m_1, m_2)$ ,  $Q = \overline{P}, \pi_v = \pi_{1,v} \otimes \pi_{2,v}$  and that  $\mathbf{s} = (s_1, s_2)$ . By Proposition 3.4,  $\pi_{i,v}$ is induced from a representation of the form

$$J(\tau_{i,1,v})[t_{i,1}] \otimes \cdots \otimes J(\tau_{i,r,v})[t_{i,r}],$$

where  $\tau_{i,j,v}$  is square integrable and  $|t_{i,j}| < 1/2 - \frac{1}{r_i^2 + 1}$ .

Let w be in the normalizer of  $M_P$  so that  $wQw^{-1}$  is a standard parabolic subgroup. Proposition I.10 in [MW] implies that the normalized

intertwining operator  $N(\tau_{1,v} \otimes \tau_{2,v}, \mathbf{s}, w)$  is holomorphic if

$$\operatorname{Re}(s_1 - s_2) > -2(\inf_{i,j}(1/2 - |t_{i,j}|)).$$

Since  $\inf_{i,j}(1/2 - |t_i|) > \frac{1}{1+n^2}$  the operator  $R_{\bar{P}|P}(\tau_{1,v} \otimes \tau_{2,v}, \mathbf{s})$  has no pole in

$$\{\mathbf{s} \in \mathbb{C}^2 \mid |\operatorname{Re}(s_i)| < (1+n^2)^{-1}, \ i=1,2\}.$$

#### 5. Absolute convergence of the spectral side

In this section we prove Theorem 0.1. First we consider the global Rankin-Selberg L-functions and the corresponding global normalizing factors defined by L-functions. Then we compare these normalizing factors with the global normalizing factors of [Mu4]. Let

$$\pi_i = \otimes_v \pi_{i,v}, \quad i = 1, 2,$$

be an irreducible unitary representation of  $\operatorname{GL}_{m_i}(\mathbb{A})$ . Then the Euler product

$$L(s,\pi_1\times\pi_2)=\prod_v L(s,\pi_{1,v}\times\pi_{2,v})$$

is absolutely convergent in some half-plane  $\operatorname{Re}(s) > c$ . To define the global  $\epsilon$ -factor we have to pick a non-trivial continuous character  $\psi$ :  $\mathbb{A}^+ \to \mathbb{C}^\times$  of the additive group  $\mathbb{A}^+$  of  $\mathbb{A}$ . Then  $\psi = \otimes_v \psi_v$  and there exists a finite set of places S, including the Archimedean ones, such that  $c(\psi_v) = 0$  and  $\pi_{1,v}$  and  $\pi_{2,v}$  are unramified for  $v \notin S$ . Hence  $f(\pi_{1,v} \times \pi_{2,v}) = 0$  for all  $v \notin S$ , and therefore  $\epsilon(s, \pi_{1,v} \times \pi_{2,v}, \psi_v) = 1$  for all  $v \notin S$ . This implies that the product

$$\epsilon(s, \pi_1 \times \pi_2, \psi) = \prod_v \epsilon(s, \pi_{1,v} \times \pi_{2,v}, \psi_v)$$

exists for all  $s \in \mathbb{C}$  and defines an entire function. We note that the global  $\epsilon$ -factor is in fact independent of  $\psi$  and therefore, will be denoted by  $\epsilon(s, \pi_1 \times \pi_2)$ .

The Rankin-Selberg *L*-functions attached to automorphic representations in the discrete spectrum have the following important properties.

**Proposition 5.1.** Let  $\pi_1$  and  $\pi_2$  be two unitary automorphic representations in the discrete spectrum of  $\operatorname{GL}_{m_1}(\mathbb{A})$  and  $\operatorname{GL}_{m_2}(\mathbb{A})$ , respectively. Let  $m = \max\{m_1, m_2\}$ . Then we have

(1) The Euler product for  $L(s, \pi_1 \times \pi_2)$  converges absolutely in the half-plane  $\operatorname{Re}(s) > m$ .

#### WERNER MÜLLER AND BIRGIT SPEH

(2)  $L(s, \pi_1 \times \pi_2)$  has a meromorphic continuation to the entire complex plane and satisfies the functional equation

(5.1) 
$$L(s,\pi_1\times\pi_2) = \epsilon(s,\pi_1\times\pi_2)L(1-s,\widetilde{\pi}_1\times\widetilde{\pi}_2).$$

(3) The function  $s \in \mathbb{C} \mapsto \prod_{j=-m}^{m} (s-j)(1-s+j)L(s,\pi_1 \times \pi_2)$  is an entire function of order one.

*Proof.* First suppose that  $\pi_1$  and  $\pi_2$  are unitary cuspidal automorphic representation. Then by Theorem 5.3 of [JS1], the Euler product is convergent in the half-plane Re(s) > 1. The functional equation is established in [Sh1, Th. 4.1] combined with [Sh3, Prop. 3.1] and [Sh3, Th. 1]. See also [Sh5] for the general case. The location of the poles of  $L(s, \pi_1 \times \pi_2)$  has been determined in the appendix of [MW], from which the first part of 3) follows. The fact that  $s(1-s)L(s, \pi_1 \times \pi_2)$  is of order one and bounded in vertical strips was proved in [RS, p.280].

Now suppose that  $\pi_1$  and  $\pi_2$  are unitary automorphic representations in the residual spectrum of  $\operatorname{GL}_{m_1}(\mathbb{A})$  and  $\operatorname{GL}_{m_2}(\mathbb{A})$ , respectively. Then by the main result of [MW] there exist  $k_i \in \mathbb{N}$  with  $k_i | m_i$ , parabolic subgroups  $P_i$  of  $\operatorname{GL}_{m_i}$  of type  $(d_i, ..., d_i)$ ,  $d_i = m_i/k_i$ , and cuspidal automorphic representations  $\xi_i$  of  $\operatorname{GL}_{d_i}(\mathbb{A})$  such that

$$\pi_i \cong J_{P_i}^{\operatorname{GL}_{m_i}}(\xi_i[(k_i-1)/2], ..., \xi_i[(1-k_i)/2]), \quad i=1,2.$$

Since  $\xi_i$  is generic, it follows from (9.4) of [JPS] that

$$L(s, \pi_1 \times \pi_2) = \prod_{i=0}^{k_1} \prod_{j=0}^{k_2} L(s + (k_1 + k_2)/2 - i - j, \xi_1 \times \xi_2).$$

Using the properties of the cuspidal *L*-functions, the proposition follows.  $\hfill \Box$ 

Now let M be a standard Levi subgroup of type  $(m_1, ..., m_r), Q, P \in \mathcal{P}(M)$  and let  $\pi$  be an automorphic representation occurring in the discrete spectrum of  $M(\mathbb{A})$ . Let  $\pi = \bigotimes_v \pi_v$  and set

(5.2) 
$$r_{Q|P}(\pi, \mathbf{s}) = \prod_{v} r_{Q|P}(\pi_{v}, \mathbf{s}), \quad \mathbf{s} \in \mathbb{C}^{r},$$

where  $r_{Q|P}(\pi_v, \mathbf{s})$  is defined by (2.3). Then it follows from (2.3) and Proposition 5.1 that the infinite product is absolutely convergent in the domain  $\operatorname{Re}(s_i - s_j) > \max\{m_1, ..., m_r\}$  and admits a meromorphic extension to the entire complex plane. Furthermore we have

$$r_{Q|P}(\pi, \mathbf{s}) := \prod_{(i,j)\in \mathrm{I}(\sigma_1, \sigma_2)} \frac{L(s_i - s_j, \pi_i \times \widetilde{\pi}_j)}{L(1 + s_i - s_j, \pi_i \times \widetilde{\pi}_j)\epsilon(s_i - s_j, \pi_i \times \widetilde{\pi}_j)}$$

where  $I(\sigma_1, \sigma_2)$  is the same as in (2.3). Let

$$\widetilde{r}_{Q|P}(\pi, \mathbf{s}) = \prod_{v} \widetilde{r}_{Q|P}(\pi_{v}, \mathbf{s})$$

be the corresponding global normalizing factor used in [Mu4]. Then  $\widetilde{r}_{Q|P}(\pi, \mathbf{s})$  is also a meromorphic function of  $\mathbf{s} \in \mathbb{C}^r$  and it follows from Lemma 2.1 that

(5.3) 
$$r_{Q|P}(\pi, \mathbf{s}) = \prod_{(i,j)\in I(\sigma_1, \sigma_2)} \frac{\epsilon(0, \pi_i \times \widetilde{\pi}_j)}{\epsilon(s_i - s_j, \pi_i \times \widetilde{\pi}_j)} \widetilde{r}_{Q|P}(\pi, \mathbf{s}), \quad \mathbf{s} \in \mathbb{C}^r.$$

Now recall that the global  $\epsilon$ -factor is of the form

$$\epsilon(s, \pi_1 \times \pi_2) = W(\pi_1 \times \pi_2) (D_{E/\mathbb{Q}}^{m_1 m_2} N(\pi_1 \times \pi_2))^{1/2-s},$$

where  $D_{E/\mathbb{Q}}$  is the discriminant of the field E,  $W(\pi_1 \times \pi_2)$  is a complex number of absolute value 1, and  $N(\pi_1 \times \pi_2)$  is an integer which is determined by the local  $\epsilon$ -factors as follows. Given a place  $v < \infty$ , let  $f(\pi_{1,v} \times \pi_{2,v})$  be the conductor of  $\pi_{1,v} \otimes \pi_{2,v}$  defined by (2.6) and (2.7). Then  $f(\pi_{1,v} \times \pi_{2,v}) = 0$  for almost all v and

$$N(\pi_1 \times \pi_2) = \prod_{v < \infty} q_v^{f(\pi_{1,v} \times \pi_{2,v})}.$$

Together with (5.3) it follows that in order to compare the normalizing factors  $r_{Q|P}(\pi, \mathbf{s})$  and  $\tilde{r}_{Q|P}(\pi, \mathbf{s})$ , we have to estimate  $N(\pi_1 \times \pi_2)$ . To this end, let  $\pi \in \Pi(\operatorname{GL}_m(\mathbb{A}))$ . Then with respect to the canonical decomposition  $\operatorname{GL}_m(\mathbb{A}) = \operatorname{GL}(\mathbb{A}_\infty) \times \operatorname{GL}_m(\mathbb{A}_f)$ , the representation  $\pi$ factorizes as  $\pi = \pi_\infty \otimes \pi_f$ . Given an open compact subgroup  $K_f \subset$  $\operatorname{GL}_m(\mathbb{A}_f)$ , set

$$\Pi(\operatorname{GL}_m(\mathbb{A}), K_f) = \left\{ \pi \in \Pi(\operatorname{GL}_m(\mathbb{A})) \mid \pi_f^{K_f} \neq 0 \right\}.$$

Let  $K_{f,i} \subset \operatorname{GL}_{n_i}(\mathbb{A}_f)$ , i = 1, 2, be two open compact subgroups and let  $K_{f,i} = \prod_{v < \infty} K_{v,i}$ . Then there exists a finite set of places S, including the Archimedean places, such that  $K_{v,i} = \operatorname{GL}_{n_i}(\mathcal{O}_{E_v})$ , i = 1, 2, and  $c(\psi_v) = 0$  for all  $v \notin S$ . Let  $\pi_i \in \Pi(\operatorname{GL}_{n_i}(\mathbb{A}), K_{f,i})$ , i = 1, 2, and let  $f(\pi_{1,v} \times \pi_{2,v})$  be the conductor of  $\pi_{1,v} \otimes \pi_{2,v}$ . By definition,  $\pi_{1,v}$  and  $\pi_{2,v}$  are unramified for all  $v \notin S$ . Hence  $f(\pi_{1,v} \times \pi_{2,v}) = 0$  for  $v \notin S$ . Let  $S_f = \{v \in S \mid v < \infty\}$ . Then  $N(\pi_1 \times \pi_2)$  is of the form

$$N(\pi_1 \times \pi_2) = \prod_{v \in S_f} q_v^{f(\pi_{1,v} \times \pi_{2,v})}$$

for all  $\pi_i \in \Pi(\operatorname{GL}_{n_i}(\mathbb{A}), K_{f,i})$ , i = 1, 2. By Lemma 2.2 and (2.11) it follows that there exists C > 0 such that for all  $\pi_i \in \Pi(\operatorname{GL}_{n_i}(\mathbb{A}), K_{f,i})$ ,

i = 1, 2, one has

$$N(\pi_1 \times \pi_2) \le C.$$

Now suppose that  $K_f \subset \operatorname{GL}_m(\mathbb{A}_f)$  is an open compact subgroup. Set

$$K_{M,f} = K_f \cap M(\mathbb{A}_f).$$

As above set

$$\Pi(M(\mathbb{A}), K_f) = \left\{ \pi \in \Pi(M(\mathbb{A})) \mid \pi_f^{K_{M,f}} \neq 0 \right\}.$$

Suppose that  $M = \operatorname{GL}_{n_1} \times \cdots \times \operatorname{GL}_{n_r}$ . Then

$$K_{M,f} = \prod_{i=1}^{\prime} K_{f,i}$$

where  $K_{f,i} \subset \operatorname{GL}_{n_i}(\mathbb{A}_f)$  is an open compact subgroup, and it follows from the above observations that there exists C > 0 such that for all representations  $\pi = \pi_1 \otimes \cdots \otimes \pi_r$  in  $\Pi(M(\mathbb{A}); K_f)$ 

(5.4) 
$$N(\pi_i \times \pi_j) \le C, \quad i, j = 1, ..., r.$$

By (5.3) this implies that the estimations of the logarithmic derivatives of normalizing factors established in [Mu4] also hold for the normalizing factors  $r_{Q|P}(\pi, \mathbf{s})$ . More precisely, assume that P is a maximal parabolic subgroup with Levi component M. Let  $A_{M,\mathbb{Q}}$  denote the split component of the center of the group  $\operatorname{Res}_{E/\mathbb{Q}} \operatorname{GL}_n$  obtained by restricting scalars from E to  $\mathbb{Q}$ . Given a unitary character  $\xi$  of  $A_{M,\mathbb{Q}}(\mathbb{R})^0$ , let  $L^2(M(E) \setminus M(\mathbb{A}))_{\xi}$  be the space of all measurable functions  $\phi$  on  $M(E) \setminus M(\mathbb{A})$  such that  $\phi(xm) = \xi(x)\phi(m)$  for all  $x \in$  $A_{M,\mathbb{Q}}(\mathbb{R})^0$ ,  $m \in M(\mathbb{A})$ , and  $\phi$  is square integrable on  $M(E) \setminus M(\mathbb{A})^1$ . Let  $\Pi_{\operatorname{disc}}(M(\mathbb{A}))_{\xi}$  denote the subspace of all  $\pi \in \Pi(M(\mathbb{A}))$  which are equivalent to a subrepresentation of the regular representation of  $M(\mathbb{A})$ in  $L^2(M(E) \setminus M(\mathbb{A}))_{\xi}$ . Set

$$\Pi_{\operatorname{disc}}(M(\mathbb{A})) := \bigcup_{\xi \in \Pi(A_{M,\mathbb{Q}}(\mathbb{R})^{0})} \Pi_{\operatorname{disc}}(M(\mathbb{A}))_{\xi}.$$

Let

$$\Pi_{\text{disc}}(M(\mathbb{A}), K_f) = \Pi_{\text{disc}}(M(\mathbb{A})) \cap \Pi(M(\mathbb{A}), K_f).$$

It follows from (5.3) and (5.4) that there exists C > 0 such that

$$\left| r_{\overline{P}|P}(\pi, iu)^{-1} \frac{d}{du} r_{\overline{P}|P}(\pi, iu) - \widetilde{r}_{\overline{P}|P}(\pi, iu)^{-1} \frac{d}{du} \widetilde{r}_{\overline{P}|P}(\pi, iu) \right| \le C$$

for all  $u \in \mathbb{R}$  and all  $\pi \in \Pi_{\text{disc}}(M(\mathbb{A}), K_f)$ . Given  $\pi \in \Pi(M(\mathbb{A}))$ , let  $\lambda_{\pi_{\infty}}$  denote the Casimir eigenvalue of  $\pi_{\infty}$ . Together with Theorem 5.3 of [Mu4] we obtain the following result.

**Proposition 5.2.** Let P be a maximal parabolic subgroup with Levi component M. Set  $m = [E : \mathbb{Q}]n^2$ . Let  $\sigma \in \Pi(K_{\infty})$ . For every  $N \ge 8m + 2$  there exists  $C_N > 0$  such that

$$\int_{\mathbb{R}} \left| r_{\overline{P}|P}(\pi, iu)^{-1} \frac{d}{du} r_{\overline{P}|P}(\pi, iu) \right| (1+u^2)^{-N} du \leq C_N (1+\lambda_{\pi_{\infty}}^2)^{18m}$$
  
for all  $\pi \in \prod_{\text{disc}} (M(\mathbb{A}), K_f)$  with  $[I_P^G(\pi_{\infty})]_{K_{\infty}} : \sigma] > 0.$ 

In general, one has to work with the multidimensional logarithmic derivatives of the normalizing factors considered in [A4]. Repeating the proof of Theorem 5.4 of [Mu4] it follows immediately that this theorem also holds for the normalizing factors  $r_{Q|P}(\pi, \mathbf{s})$ . Therefore Theorem 0.2 of [Mu4] holds with respect to the normalization of local intertwining operators used in the present paper. Combined with Proposition 0.2, this completes the proof of Theorem 0.1.

#### References

- [A1] J. Arthur, Eisenstein series and the trace formula, Proc. Sympos. Pure Math., 33, Part I, Amer. Math. Soc., Providence, R.I. (1979), 253–274.
- [A2] J. Arthur, A trace formula for reductive groups I: terms associated to classes in  $G(\mathbb{Q})$ , Duke. Math. J. 45 (1978), 911–952.
- [A3] J. Arthur, On a family of distributions obtained from Eisenstein series I: Applications of the Paley-Wiener theorem, Amer. J. Math. 104 (1982), 1243–1288.
- [A4] J. Arthur, On a family of distributions obtained from Eisenstein series II: Explicit formulas, Amer. J. Math. 104 (1982), 1289–1336.
- [A5] J. Arthur, The trace formula in invariant form, Annals of Math. 114 (1981), 1–74.
- [A6] J. Arthur, The invariant trace formula. II. Global theory, J. Amer. Math. Society 1 (1988), 501–554.
- [A7] J. Arthur, Intertwining operators and residues. I. Weighted characters, J. Funct. Analysis 84 (1989), 19–84.
- [A8] J. Arthur, On the Fourier transforms of weighted orbital integrals, J. reine angew. Math. 452 (1994), 163–217.
- [AC] J. Arthur and L. Clozel, Simple algebras, base change, and the advanced theory of the trace formula, Annals Math. Studies 120, Princeton Univ. Press, Princeton, NJ, 1989.
- [Bu] C.J. Bushnell, Induced representations of locally profinite groups, Journal of Algebra 134 (1990), 104–114.
- [BH] C.J. Bushnell and G. Henniart, An upper bound on conductors for pairs, J. Number Theory 65 (1997), 183–196.
- [BHK] C.J. Bushnell, G.M. Henniart, and P.C. Kutzko, Local Rankin-Selberg convolutions for GL<sub>n</sub>: Explicit conductor formula, J. Amer. Math. Soc. 11 (1998), 703–730.

- [BW] A. Borel and N. Wallach, Continuous cohomology, discrete subgroups, and representations of reductive groups, Annals of Math. Stud. 94, Princeton Univ. Press, 1980.
- [BZ] I.N. Bernstein and A.V. Zelevinsky, Induced representations of reductive p-adic groups, I. Ann. Scient. Éc. Norm. Sup. 10 (1977), 441–472.
- [CLL] L. Clozel, J.P. Labesse, and R. Langlands, Morning seminar on the trace formula, mimeoraphed notes, IAS, Princeton, 1984.
- [GS] S. Gelbard and F. Shahidi, Boundedness of automorphic L-functions in vertical strips, Preprint, 2000.
- [GJ] R. Godement and H. Jacquet, Zeta functions of simple algebras. Lecture Notes in Math. 260, Springer, Berlin-Heidelberg-New York, 1972.
- [HC1] Harish-Chandra, Automorphic forms on semi-simple Lie groups, Lecture Notes in Math. 62, Springer-Verlag, Berlin-Heidelberg-New York, 1968.
- [HC2] Harish-Chandra, The Plancherel formula for reductive p-adic groups, In: Collected papers, IV, pp. 353-367, Springer-Verlag, New York-Berlin-Heidelberg, 1984.
- [J] H. Jacquet, Principal L-functions of the linear group, Proc. Symp. Pure Math. 33 (1979), 63-86.
- [JPS] H. Jacquet, I.I. Piatetski-Shapiro, and J.A. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math. **105** (1983), 367-464.
- [JS1] H. Jacquet and J.A. Shalika, Euler products and classification of automorphic representations I, Amer. J. Math. 103 (1981), 499-558.
- [JS2] H. Jacquet and J.A. Shalika, Rankin-Selberg convolutions: Archimedean theory, in: Festschrift in Honor of I.I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Israel Math. Conf. Proc. 2, Weizmann, Jerusalem, 1990, 125-208.
- [JS3] H. Jacquet, J.A. Shalika, The Whittaker models of induced representations, Pacific J. Math. 109 (1983), 107–120.
- [Kn] A.W. Knapp, Representation theory of semisimple groups, Princeton University Press, Princeton, New Jersey 1986.
- [KS] A.W. Knapp and E.M. Stein, Intertwining operators for semisimple Lie groups, II, Inventiones math. 60 (1980), 9-84.
- [KZ] A.W. Knapp and G. Zuckerman, Classification theorems for representations of semisimple Lie groups, in Lecture Notes Math. Vol. 587, pp.138-159, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [Lb] J.-P. Labesse, Noninvariant base change identities, Mém. Soc. Math. France (N.S.), 61.
- [L1] R.P. Langlands, On the functional equations satisfied by Eisenstein series, Lecture Notes in Math. 544, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [L2] R.P. Langlands, Eisenstein series, In: Proc. Symp. Pure Math. Vol. 9, A. M. S., Providence, R.I., 1966.
- [L3] R.P. Langlands, On the classification of irreducible representations of real algebraic groups, Mathematical Surveys and Monographs 31, Amer. Math. Soc. 1989.
- [LRS] W. Luo, Z. Rudnick, and P. Sarnak, On the generalized Ramanujan conjecture for GL(n). Proc. Symp. Pure Math. 66, Vol. II, pp. 301–310, Amer. Math. Soc., Providence, Rhode Island, 1999.

- [MW] C. Moeglin et J.-L. Waldspurger, Le spectre résiduel de GL(n), Ann. scient. Éc. Norm. Sup.,  $4^e$  série, t. **22** (1989), 605–674.
- [Mu1] W. Müller, The trace class conjecture in the theory of automorphic forms, Annals of Math. **130** (1989), 473–529.
- [Mu2] W. Müller, The trace class conjecture in the theory of automorphic forms. II, Geom. Funct. Analysis 8 (1998), 315–355.
- [Mu3] W. Müller, On the singularities of residual intertwining operators, Geom. funct. anal. 10 (2000), 1118-1170.
- [Mu4] W. Müller, On the spectral side of the Arthur trace formula, Geom. funct. anal. **12** (2002), 669-722.
- [Neu] J. Neukirch, Algebraische Zahlentheorie, Springer-Verlag, Berlin-Heidelberg-New York, 1992.
- [RS] Z. Rudnick and P. Sarnak, Zeros of principal L-functions and random matrix theory. Duke Math. J. 81 (1996), 269–322.
- [Se] A. Selberg, Harmonic analysis, in "Collected Papers", Vol. I, Springer-Verlagf, Berlin-Heidelberg-New York (1989), 626–674.
- [Sh1] F. Shahidi, Fourier transforms of intertwining operators and Plancherel measures for  $GL_n$ , Amer. J. Math **106** (1984), 67–111.
- [Sh2] F. Shahidi, On certain L-functions, Amer. J. Math. 103 (1981), 297–356.
- [Sh3] F. Shahidi, Local coefficients and normalization of intertwining operators for  $GL_n$ , Comp. Math. 48 (1983), 271–295.
- [Sh4] F. Shahidi, Local coefficients as Artin factors for real groups, Duke Math. J. 52 (1985), 973–1007.
- [Sh5] F. Shahidi, A proof of Langlands' conjecture on Plancherel measures; Complementary series for p-adic groups, Annals Math. 132 (1990), 273–330.
- [Sk] J.A. Shalika, The multiplicity one theorem for  $GL_n$ , Annals Math. 100 (1974), 171-193.
- [Si1] A. Silberger, Special representations of reductive p-adic groups are not integrable, Annals Math. 111 (1980), 571–587.
- [Si2] A. Silberger, On Harish-Chandra's μ-function for p-adic groups, Trans. Amer. Math. Soc. 260 (1980), 113–121.
- [Si3] A. Silberger, The Langlands quotient theorem for p-adic groups, Math. Ann. 236 (1978), 95–104.
- [Sp] B. Speh , The unitary dual of  $GL(3,\mathbb{R})$  and  $GL(4,\mathbb{R})$ , Math. Ann 258, (1981/82), 113-133.
- [Ta] M. Tadić, Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case). Ann. Sci. cole Norm. Sup. (4) 19 (1986), 335–382.
- [Vo] D.B. Vogan, The unitary dual of GL(n) over an Archimedean field, Inventiones math. 83 (1986), 449–505.
- [VW] D.B. Vogan, Jr. and N.R. Wallach, Intertwining operators for real reductive Lie groups, Advances in Math. 82 (1990), 203–243.
- [Wa1] G. Warner, Harmonic analysis on semi-simple Lie groups I, Springer-Verlag, Berlin-Heidelberg-New York, 1972.
- [Wa2] G. Warner, Harmonic analysis on semi-simple Lie groups II, Springer-Verlag, Berlin-Heidelberg-New York, 1972.
- [Wh] E.T. Whitaker and G.N. Watson, A course of modern analysis, Cambridge, University Press, 4th edition, 1940.

# WERNER MÜLLER AND BIRGIT SPEH

[Z] A.V. Zelevinsky, Induced representations of reductive p-adic groups II. On irreducible representations of  $GL_n$ , Ann. Scient. Éc. Norm. Sup. 13 (1980), 165–210.

UNIVERSITÄT BONN, MATHEMATISCHES INSTITUT, BERINGSTRASSE 1, D-53115 BONN, GERMANY

*E-mail address:* mueller@math.uni-bonn.de

Department of Mathematics, 310 Malott Hall, Cornell University, Ithaca, NY 14853-4201, U.S.A.

*E-mail address*: speh@math.cornell.edu