ON THE ASYMPTOTIC LINEARITY OF CASTELNUOVO-MUMFORD REGULARITY

NGÔ VIÊT TRUNG AND HSIN-JU WANG

Dedicated to Wolmer Vasconcelos on the occasion of his sixtyfifth birthday

INTRODUCTION

Let A be a commutative Noetherian ring with unity. Let R be a standard graded algebra over A, where "standard" means $R_0 = A$ and R is generated by elements of R_1 . Let R_+ be the ideal generated by the elements of positive degree. For any finitely generated graded R-module M and $i \ge 0$ we denote by $H^i_{R_+}(M)$ the *i*th local cohomology module of M with respect to R_+ . The Castelnuovo-Mumford regularity of M is the invariant

 $\operatorname{reg}(M) := \max\{a(H_{R_{+}}^{i}(M)) + i | i \ge 0\},\$

where a(H) denotes the maximal non-vanishing degree of a graded *R*-module *H* with the convention $a(H) = -\infty$ if H = 0. It is a natural extension of the usual definition of the Castelnuovo-Mumford regularity in the case *R* is a graded algebra over a field. If *R* is a polynomial ring over a field *k*, then *M* has a finite graded minimal free resolution: $0 \longrightarrow F_r \longrightarrow \cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M$ and $\operatorname{reg}(M) = \max\{b_i(M) - i | i \ge 0\}$, where $b_i(M)$ denotes the maximal degree of the generators of F_i . Hence one may view $\operatorname{reg}(M)$ as a measure for the complexity of the structure of *M* (see e.g. [BaM93] and [EG84]).

There have been a surge of interest on the behavior of the function $\operatorname{reg}(I^n)$ where I is a homogeneous ideal in a polynomial ring over a field. It was first discovered by Bertram, Ein, and Lazarsfeld [BEL91] that if I is the defining ideal of a smooth complex variety, $\operatorname{reg}(I^n)$ is bounded by a linear function. Later, Swanson [Swa97] proved this for all homogeneous ideals. Recently, Cutkosky et al [CuHT99] and Kodiyalam [Kod00] found out that $\operatorname{reg}(I^n)$ is asymptotically a linear function. See also [Cha97], [Cut00], [CuEL], [GGP95] for related results.

The aim of this paper is to prove the above phenomenon for the general case, namely, when I is a graded ideal of a standard graded algebra R over an arbitrary ring A. That is not a simple task because the proof in the case R is a polynomial ring over a field is based strongly on the fact that finitely generated modules over R have finite minimal free resolutions, whereas that fact does not hold in general. Moreover, as far as we can see, the general case can not be reduced to this case. Our approach uses the characterization of the Castelnuovo-Mumford regularity by means of filterregular sequences combined with some results originally due to [CuHT99] and [Kod00].

¹⁹⁹¹ Mathematics Subject Classification. Primary 13C99; Secondary 13F17.

Key words and phrases. Castelnuovo regularity, filter-regular sequence, reduction, bigraded ring.

The first author is partially supported by the National Basic Research Program of Vietnam.

Even these results need to be proved differently since the original proofs are based on techniques which are not available in general such as Gröbner basis and Nakayama's lemma. More general, we will study the asymptotic regularity of the *I*-adic filtration of an arbitrary finitely generated graded *R*-module *M*. This generalization allows us to prove that $\operatorname{reg}(\overline{I^n})$, where $\overline{I^n}$ denotes the integral closure of I^n , is asymptotically a linear function with the same slope as $\operatorname{reg}(I^n)$ if *R* is a domain over an affine ring. The latter fact was not known when *R* is a polynomial ring over a field.

We call a graded ideal $J \subseteq I$ an *M*-reduction of *I* if $I^{n+1}M = JI^nM$ for some $n \ge 0$. Define

 $\rho_M(I) := \min\{d(J) \mid J \text{ is an } M \text{-reduction of } I\},\$

where for a graded module N, d(N) denotes the minimum number d such that there is a homogeneous generating set for N with elements of degree $\leq d$. Moreover, let $\varepsilon(M)$ denote the smallest degree of the homogeneous elements of M. Then our main result can be formulated as follows.

Theorem 3.2. Let R be a standard graded ring over a commutative Noetherian ring with unity and I a graded ideal of R. Let M be a finitely generated graded R-module. Then there exists an integer $e \ge \varepsilon(M)$ such that for all large n,

$$\operatorname{reg}(I^n M) = \rho_M(I)n + e.$$

1. FILTER-REGULAR SEQUENCE AND REGULARITY

Let A be an arbitrary commutative Noetherian ring with unity. Let R be a standard graded algebra over A and M a finitely generated graded R-module.

Let z_1, \ldots, z_s be linear forms in R. We call z_1, \ldots, z_s an *M*-filter-regular sequence if $z_i \notin P$ for any associated prime $P \not\supseteq R_+$ of $(z_1, \ldots, z_{i-1})M$ for $i = 1, \ldots, s$. Filterregular elements have their origin in the theory of Buchsbaum rings [STC78].

The Castelnuovo-Mumford regularity can be characterized by means of filter-regular sequences. The following characterization was proved implicitly in the proof of [Tru98, Theorem 2.4] for the case M = R. But that proof also holds for any M.

Proposition 1.1. Let z_1, \ldots, z_s be an *M*-filter-regular sequence of linear forms which generate an *M*-reduction of R_+ . Then

$$\operatorname{reg}(M) = \max \left\{ a \big((z_1, \dots, z_i) M : R_+ / (z_1, \dots, z_i) M \big) | \ i = 0, \dots, s \right\}.$$

Note that a homogeneous ideal $Q \subseteq R_+$ is an *M*-reduction of R_+ if and only if $(M/QM)_n = 0$ for all large *n*.

We shall see that any M-reduction of R_+ can be generated by an M-filter-regular sequence in a flat extension of A.

Lemma 1.2. Let Q be an M-reduction generated by the linear forms x_1, \ldots, x_s . For $i = 1, \ldots, s$ put $z_i = \sum_{j=1}^s u_{ij}x_j$, where $U = \{u_{ij} | i, j = 1, \ldots, s\}$ is a matrix of indeterminates. Put

$$A' = A[U, \det(U)^{-1}], \ R' = R \otimes_A A', \ M' = M \otimes_A A'.$$

If we view R' as a standard graded algebra over A' and M' as a graded R'-module, then z_1, \ldots, z_s is an M'-filter-regular sequence.

Proof. Since the elements z_i are defined by independent sets of indeterminates, it suffices to show that $z_1 \notin P$ for any associated prime $P \not\supseteq R'_+$ of M'. By the definition of R', such a prime P must have the form $\mathfrak{p}R'$ for some associated prime $\mathfrak{p} \not\supseteq R_+$ of M. If $Q \subseteq \mathfrak{p}$, then $(M/\mathfrak{p}M)_n = 0$ for all large n since $M/\mathfrak{p}M$ is a quotient module of M/QM. From this it follows that there is a number t such that $R^t_+M \subseteq \mathfrak{p}M$. Since $\operatorname{ann}(M) \subseteq \mathfrak{p}$, this implies $R_+ \subseteq \mathfrak{p}$, a contradiction. So we get $Q \not\subseteq \mathfrak{p}$. Since $Q = (x_1, \ldots, x_s)$, this implies $z_1 = u_{11}x_1 + \cdots + u_{1s}x_s \notin \mathfrak{p}R' = P$.

Lemma 1.2 allows us to use Proposition 1.1 for the computation of $\operatorname{reg}(M)$. Indeed, since A' =is a flat extension of A, we have $H^i_{R'_+}(M')_n \cong H^i_{R_+}(M)_n \otimes_A A'$ for all n and $i \ge 0$, whence

$$\operatorname{reg}(M) = \operatorname{reg}(M').$$

From the characterization of the regularity by means of filter-regular sequences we can deduce the following relationship between d(M) and reg(M). This was done in [Tru87, Proposition 4.1] for the case R is a polynomial ring over a local ring and M is a graded ideal of R. But the proof there also holds for the general case.

Proposition 1.3. $d(M) \leq \operatorname{reg}(M)$.

2. BIGRADED MODULE AND REGULARITY

Let $S = A[X_1, \ldots, X_s, Y_1, \ldots, Y_v]$ be a polynomial ring over a commutative Noetherian ring A with unity. Given a sequence of non-negative integers d_1, \ldots, d_v , we can view S as a bigraded ring with deg $X_i = (1, 0), i = 1, \ldots, s$, and deg $Y_j = (d_j, 1),$ $j = 1, \ldots, v$. For convenience we will assume that $d_v = \max\{d_i | i = 1, \ldots, v\}$.

Let \mathcal{M} be a finitely generated bigraded module over $A[X_1, \ldots, X_s, Y_1, \ldots, Y_v]$. For a fixed number n put

$$\mathcal{M}_n := \oplus_{a \ge 0} \mathcal{M}_{(a,n)}.$$

Then \mathcal{M}_n is a finitely generated graded module over the naturally graded polynomial ring $A[X_1, \ldots, X_s]$. We will show that $\operatorname{reg}(\mathcal{M}_n)$ is asymptotically a linear function.

If s = 0, then $\operatorname{reg}(\mathcal{M}_n)$ is the invariant

$$a(\mathcal{M}_n) = \max\{a \mid \mathcal{M}_{(a,n)} \neq 0\}.$$

This case is settled by the following result which was already known in the case A is a field [CuHT99, Theorem 3.3]. However, the proof there used Gröbner basis (and integer programming) which is not available in the general case.

Proposition 2.1. Let \mathcal{M} be a finitely generated bigraded module over the bigraded polynomial ring $A[Y_1, \ldots, Y_v]$. Then $a(\mathcal{M}_n)$ is asymptotically a linear function with $slope \leq d_v$.

Proof. The case v = 0 is trivial. Assume that $v \ge 1$. Consider the exact sequence of bigraded $A[Y_1, \ldots, Y_v]$ -modules:

$$0 \longrightarrow [0_{\mathcal{M}}: Y_{v}]_{(a,n)} \longrightarrow \mathcal{M}_{(a,n)} \xrightarrow{Y_{v}} \mathcal{M}_{(a+d_{v},n+1)} \longrightarrow [\mathcal{M}/Y_{v}\mathcal{M}]_{(a+d_{v},n+1)} \longrightarrow 0.$$

Since $0_{\mathcal{M}} : Y_v$ and $\mathcal{M}/Y_v\mathcal{M}$ can be viewed as bigraded modules over $A[Y_1, \ldots, Y_{v-1}]$, using induction we may assume that $a([0_{\mathcal{M}} : Y_v]_n)$ and $a([\mathcal{M}/Y_v\mathcal{M}]_n)$ are asymptotically linear functions with slopes $\leq d_v$. As a consequence,

$$a([0_{\mathcal{M}}: Y_v]_n) + d_v \ge a([0_{\mathcal{M}}: Y_v]_{n+1})$$
$$a(\mathcal{M}/Y_v\mathcal{M})_n) + d_v \ge a([\mathcal{M}/Y_v\mathcal{M}]_{n+1})$$

for all large n. Further, we are done if $a(\mathcal{M}_n) = a([0_{\mathcal{M}} : Y_v]_n)$ for all large n.

Since $a(\mathcal{M}_n) \geq a([0_{\mathcal{M}} : Y_v]_n)$ for all n, it remains to consider the case that there exists an infinite sequence of integers m for which $a(\mathcal{M}_m) > a([0_{\mathcal{M}} : Y_v]_m)$. Putting this condition into the above exact sequence we get

$$a(\mathcal{M}_{m+1}) = \max\{a(\mathcal{M}_m) + d_v, a([\mathcal{M}/Y_v\mathcal{M}]_{m+1})\}.$$

On the other hand, we have

$$a(\mathcal{M}_m) + d_v \ge a(\mathcal{M}/Y_v\mathcal{M})_m + d_v \ge a([\mathcal{M}/Y_v\mathcal{M}]_{m+1})$$

for m large enough. So we can find infinitely many integers m such that

$$a(\mathcal{M}_{m+1}) = a(\mathcal{M}_m) + d_v > a([0_{\mathcal{M}} : Y_v]_m) + d_v \ge a([0_{\mathcal{M}} : Y_v]_{m+1}).$$

Fix such an integer m. Similarly, we can prove that for all $n \ge m$, $a(\mathcal{M}_n) > a([0_{\mathcal{M}} : Y_v]_n)$ and therefore $a(\mathcal{M}_{n+1}) = a(\mathcal{M}_n) + d_v$. So $a(\mathcal{M}_n)$ is asymptotically a linear function with slope d_v in this case.

Now we are going to consider the general case.

Theorem 2.2. Let \mathcal{M} be a finitely generated bigraded module over the above bigraded polynomial ring $A[X_1, \ldots, X_s, Y_1, \ldots, Y_v]$. Then $\operatorname{reg}(\mathcal{M}_n)$ is asymptotically a linear function with slope $\leq d_v$.

Proof. By Proposition 2.1 we may assume that $s \ge 1$. For $i = 1, \ldots, s$ put $z_i = \sum_{j=1}^{s} u_{ij}X_j$, where $U = \{u_{ij} | i, j = 1, \ldots, s\}$ is a matrix of indeterminates. Put $A' = A[U, \det(U)^{-1}]$ and $\mathcal{M}' = \mathcal{M} \otimes_A A'$. Then \mathcal{M}' is a finitely generated bigraded module over $A'[X_1, \ldots, X_s, Y_1, \ldots, Y_v]$. Since $\mathcal{M}'_n = \mathcal{M}_n \otimes_A A'$, we have

$$\operatorname{reg}(\mathcal{M}_n) = \operatorname{reg}(\mathcal{M}'_n)$$

for all $n \ge 0$. By Lemma 1.2, z_1, \ldots, z_s is an \mathcal{M}'_n -filter-regular sequence. Put $R = A'[X_1, \ldots, X_s]$. Since $(z_1, \ldots, z_s) = (X_1, \ldots, X_s) = R_+$, applying Proposition 1.1 we get

$$\operatorname{reg}(\mathcal{M}'_n) = \max \left\{ a \Big((z_1, \ldots, z_i) \mathcal{M}'_n : R_+ / (z_1, \ldots, z_i) \mathcal{M}'_n \Big) | \ i = 0, \ldots, s \right\}.$$

It is easy to check that

 $(z_1,\ldots,z_i)\mathcal{M}'_n:R_+/(z_1,\ldots,z_i)\mathcal{M}'_n=[(z_1,\ldots,z_i)\mathcal{M}':R_+/(z_1,\ldots,z_i)\mathcal{M}']_n.$

Moreover, we may consider $(z_1, \ldots, z_i)\mathcal{M}' : R_+/(z_1, \ldots, z_i)\mathcal{M}'$ as a graded module over the bigraded polynomial ring $A'[Y_1, \ldots, Y_v]$. By Proposition 2.1, the invariant $a([(z_1, \ldots, z_i)\mathcal{M}' : R_+/(z_1, \ldots, z_i)\mathcal{M}']_n)$ is asymptotically a linear function with slope $\leq d_v$ for $i = 0, \ldots, s$. Therefore, we can conclude that $\operatorname{reg}(\mathcal{M}_n)$ is asymptotically a linear function with slope $\leq d_v$.

3. Regularity of adic filtrations

Let A be a commutative Noetherian ring with unity. Let R be a standard graded algebra over A and I a graded ideal of R. We will apply the results of the preceding sections to study the function $reg(I^nM)$ for any finitely generated graded R-module M.

First we will establish a relationship between $d(I^n M)$ and the invariants $\rho_M(I)$, $\varepsilon(M)$ introduced in the introduction. This relationship was already known for the case M = R and R is a polynomial ring over a field [Kod00, Proposition 4]. However the original proof does not hold in the general case since it uses Nakayama's lemma.

Lemma 3.1. $d(I^n M) \ge \rho_M(I)n + \varepsilon(M)$ for all $n \ge 0$.

Proof. We may write I = J + K, where J and K denote the ideals generated by the homogeneous elements of I of degree $\langle \rho_M(I) \rangle$ and $\geq \rho_M(I)$, respectively. Then

$$I^n M = J I^{n-1} M + K^n M.$$

Note that $K^n M$ is generated by homogeneous elements of degree $\geq \rho_M(I)n + \varepsilon(M)$. If $d(I^n M) < \rho_M(I)n + \varepsilon(M)$, then $I^n M = JI^{n-1}M$. Hence J is an M-reduction of I. Since by definition of J, $d(J) < \rho_M(I)$, this gives a contradiction to the definition of $\rho_M(I)$.

Now we are going to prove the main result of this paper.

Theorem 3.2. Let R be a standard graded ring over a commutative Noetherian ring with unity and I a graded ideal of R. Let M be a finitely generated graded R-module. Then there exists an integer $e \ge \varepsilon(M)$ such that for all large n,

$$\operatorname{reg}(I^n M) = \rho_M(I)n + e.$$

Proof. Let J be an M-reduction of I with $d(J) = \rho_M(I)$. Let $R[Jt] = \bigoplus_{n\geq 0} J^n t^n$ be the Rees algebra of R with respect to J and $\mathcal{M} = \bigoplus_{n\geq 0} I^n M$. Since $I^{n+1}M = JI^n M$ for all large n, we may consider \mathcal{M} as a finitely generated graded module over R[Jt]. Assume that R_1 is generated by the linear forms x_1, \ldots, x_s and J is generated by the forms y_1, \ldots, y_v with deg $y_j = d_j, j = 1, \ldots, v$. Represent R[Jt] as a quotient ring of the bigraded polynomial ring $A[X_1, \ldots, X_s, Y_1, \ldots, Y_v]$ with deg $X_i = (1, 0), i = 1, \ldots, s$, and deg $Y_j = (d_j, 1), j = 1, \ldots, v$. Then \mathcal{M} is a finitely generated bigraded module over A[X, Y] with $\mathcal{M}_n = I^n M$. By Theorem 2.2, $\operatorname{reg}(I^n M)$ is asymptotically a linear function with slope $\leq \max\{d_1, \ldots, d_v\} = d(J)$. Let dn + e be this linear function. By Proposition 1.3 and Lemma 3.1 we have

$$\operatorname{reg}(I^n M) \ge d(I^n M) \ge \rho_M(I)n + \varepsilon(M)$$

for all *n*. Therefore, $d \ge \rho_M(I)$. Since $d \le d(J) = \rho_M(I)$, we can conclude that $d = \rho_M(I)$ and $e \ge \varepsilon(M)$.

Setting M = R we immediately obtain the following consequence (see [CuHT99, Theorem 3.1] and [Kod00, Theorem 5] for the case when R is a polynomial ring over a field).

Corollary 3.3. Let R be a standard graded ring over a commutative Noetherian ring with unity and I a graded ideal of R. Then there exists an integer $e \ge 0$ such that for all large n,

$$\operatorname{reg}(I^n) = \rho_R(I)n + e.$$

In particular, we may apply Theorem 3.2 to study the asymptotic regularity of the integral closures $\overline{I^n}$ of I^n (see [CuHT99, Corollary 3.5] for the case R is a polynomial ring over a field where the slope could not be determined).

Corollary 3.4. Let R be a standard graded domain over over a commutative Noetherian ring with unity and I a graded ideal of R such that $\overline{I^{n+1}} = I\overline{I^n}$ for n sufficiently large. Then reg $(\overline{I^n})$ is asymptotically a linear function with slope $\rho_R(I)$.

Proof. By assumption, there exists an integer n_0 such that $\overline{I^{n+1}} = I\overline{I^n}$ for all $n \ge n_0$. Put $M = \overline{I^{n_0}}$. Then $I^n M = \overline{I^{n+n_0}}$ for all $n \ge 0$. As a consequence, a graded ideal $J \subseteq I$ is an *M*-reduction if and only if $\overline{J} = \overline{I}$ or, equivalently, J is an *R*-reduction of *I*. Therefore, $\rho_M(I) = \rho_R(I)$. Hence the conclusion follows from Theorem 3.2.

It is well-known that the assumption of Corollary 3.4 is satisfied for any ideal I if R is an affine domain.

References

- [BaM93] D. Bayer and D. Mumford. What can be computed in algebraic geometry? in: D. Eisenbud and L. Robbiano (eds.): Computational Algebra Geometry and Commutative Algebra, Proceedings, Cortona 1991. *Cambridge University Press*, pages 1–48, 1993.
- [BEL91] A. Bertram, L. Ein, and R. Lazarsfeld. Vanishing theorems, a theorem of Severi, and the equations defining projective varieties. J. Amer. Math. Soc., 4:587–602, 1991.
- [Cha97] K. A. Chandler. Regularity of the powers of an ideal. Comm. Algebra, 25:3773–3776, 1997.
- [Cut00] S. D. Cutkosky. Irrational asymptotic behaviour of Castelnuovo-Mumford regularity. J. Reine Angew. Math., 522:93–103, 2000.
- [CuEL] S. D. Cutkosky, L. Ein, and R. Lazarsfeld. Positivity and complexity of ideal sheaves. Preprint.
- [CuHT99] S. Cutkosky, J. Herzog, and N. V. Trung. Asymptotic behaviour of the Castelnuovo-Mumford regularity. *Compositio Mathematics*, 118:243–261, 1999.
- [EG84] D. Eisenbud and S. Goto. Linear free resolutions and minimal multiplicities. J. Algebra, 88:89–133, 1984.
- [GGP95] A. V. Geramita, A. Gimigliano, and Y. Pitteloud. Graded Betti numbers of some embedded rational n-folds. Math. Ann., 301:363–380, 1995.
- [Kod00] V. Kodiyalam. Asymptotic behaviour of Castelnuovo-Mumford regularity. Proc. Amer. Math. Soc., 128:407–411, 2000.
- [STC78] P. Schenzel, N. V. Trung, and N. T. Cuong. Verallgemeinerte Cohen-Macaulay-Moduln. Math. Nachr., 85:57–73, 1978.
- [Swa97] I. Swanson. Powers of ideals. Primary decompositions, Artin-Rees lemma and regularity. Math. Ann., 307:299–313, 1997.
- [Tru87] N. V. Trung. Reduction exponent and degree bound for the defining equations of graded rings. Proc. Amer. Math. Soc., 101:229–236, 1987.
- [Tru98] N. V. Trung. The Castelnuovo regularity of the Rees algebra and the associated graded ring. Trans. Amer. Math. Soc., 350:2813–2832, 1998.

INSTITUTE OF MATHEMATICS, BOX 631, BÒ HÔ, 10000 HANOI, VIETNAM *E-mail address*: nvtrung@thevinh.vnn.vn DEPARTMENT OF MATHEMATICS, NATIONAL CHUNG CHENG UNIVERSITY, CHIAYI 621, TAIWAN *E-mail address*: hjwang@math.ccu.edu.tw