ENTIRE INVARIANT SOLUTIONS TO MONGE-AMPÈRE EQUATIONS

ROGER BIELAWSKI

ABSTRACT. We prove existence and regularity of entire solutions to Monge-Ampère equations invariant under an irreducible action of a compact Lie group.

We consider Monge-Ampère equations of the form

$$f(\nabla\phi) \det D_{ij}\phi = g(x) \tag{0.1}$$

where f and g are nonnegative measurable functions on \mathbb{R}^n . We recall first the concept of a weak solution of (0.1). Let ϕ be a convex function. Then $\nabla \phi$ is a well-defined multi-valued mapping: $(\nabla \phi)(x)$ is the set of slopes of all supporting hyperplanes to the graph of ϕ at $(x, \phi(x))$. If B is a subset of \mathbb{R}^n , let $\nabla \phi(B)$ be its image in the multi-valued sense. Then ϕ is a weak solution of (0.1) if

$$\int_{B} g(x)dx = \int_{\nabla\phi(B)} f(y)dy \tag{0.2}$$

for every Borel set B. Let us denote the right-hand side by $\omega(B, \phi, f)$. It can be shown that it is a Borel measure on \mathbb{R}^n . A basic result [1] is that if $u_k \to u$ compactly and $f_k \to f$ uniformly, then $\omega(\cdot, u_k, f_k)$ converges to $\omega(\cdot, u, f)$ weakly, i.e. as functionals on the space of compactly supported continuous functions.

Existence of weak solutions to (0.1) defined on all of \mathbb{R}^n have been shown in [1] under the assumption that $g \in L^1(\mathbb{R}^n)$ and in [9] in the case when f = 1 and g, 1/g are bounded. In this note we wish to give a simple proof of existence and regularity of entire solutions to (0.1) under a different type of assumption: f and g are invariant under an irreducible action of a compact (Lie) group. More precisely we shall prove:

Theorem 0.1. Let $K \subset O(n)$ be a compact subgroup acting irreducibly on \mathbb{R}^n and let f, g be two nonnegative K-invariant measurable functions on \mathbb{R}^n . Furthermore, assume that f and g are locally bounded and that

$$\int_{\mathbb{R}^n} f = +\infty.$$

Then there exists a (weak) convex K-invariant solution $\phi : \mathbb{R}^n \to \mathbb{R}$ of the Monge-Ampère equation (0.1).

Theorem 0.2. In the situation of Theorem 0.1 suppose, in addition, that f and g are strictly positive and of class $C^{p,\alpha}$, $p \ge 0$, $\alpha > 0$. Then any entire K-invariant convex weak solution ϕ of (0.1) is $C^{p+2,\alpha}$.

²⁰⁰⁰ Mathematics Subject Classification. 35J60.

Research supported by an EPSRC advanced fellowship.

ROGER BIELAWSKI

Well-known examples show that this regularity result does not hold without the assumption of K-invariance, even when f = g = 1.

The above theorems have been proved before in [2] in the case when K is a reflection group. They can be applied to show existence of Kähler metrics with prescribed Ricci curvature on complexified symmetric spaces.

1. Proofs

We begin with a simple lemma:

Lemma 1.1. Let $K \subset O(n)$ be a compact subgroup acting irreducibly on \mathbb{R}^n . For any $x \in \mathbb{R}^n$, let C_x be the convex hull of the orbit $K \cdot x$. There exists an $\epsilon > 0$ such that the ball $B(0, \epsilon)$ is contained in C_x for any x with |x| = 1.

Proof. Suppose that the conclusion does not hold. Then there exists a sequence of points x_k with $|x_k| = 1$ such that the boundary of C_{x_k} contains a point y_k with $y_k \to 0$. By choosing a convergent subsequence of x_k we find a point a point x_{∞} on the unit sphere such that 0 lies on the boundary of $C_{x_{\infty}}$. As $C_{x_{\infty}}$ is convex, it lies to one side of any of its supporting hyperplanes at 0. Hence the orbit $K \cdot x_{\infty}$ lies to one side of a hyperplane H passing through 0. Therefore the center of mass of the orbit

$$m = \frac{1}{|K|} \int_K g \cdot x_\infty dg$$

cannot be the origin unless the whole orbit lies on H. On the other hand m has to be the origin as it is a K-invariant point and the representation is irreducible. It follows that the orbit $K \cdot x_{\infty}$ is contained in a hyperplane and from this the subspace generated linearly by $K \cdot x_{\infty}$ is a proper K-invariant subspace of \mathbb{R}^n , a contradiction.

We now prove Theorem 0.1. Put $f_k = f + 1/k$ and $g_k = g + 1/k$, $k \in \mathbb{N}_+$. Let B_r denote the ball of radius r centred at the origin. Let R_k be a number defined by

$$\int_{B_{R_k}} f_k = \int_{B_k} g_k$$

According to [3, 7] there exists a unique (up to a constant) strictly convex solution ϕ_k of

$$f_k(\nabla \phi_k) \det D_{ij} \phi_k = g_k(x) \tag{1.1}$$

which is of class $C^{1,\beta}$, for some $\beta > 0$, and such that $\nabla \phi_k$ maps B_k onto B_{R_k} . Moreover, as $K \subset O(n)$ and all the data are K-invariant, ϕ_k is K-invariant (this follows from uniqueness, since $\phi_k \circ g$, $g \in K$, is also a solution of (1.1)). To prove the existence of a weak solution to (0.1), defined on all of \mathbb{R}^n , it is enough to show that the functions ϕ_k are uniformly (in k) bounded on any ball B_R , $k \geq R$. Indeed, a bounded sequence of convex functions on a bounded open convex domain has a convergent subsequence. This follows from the elementary fact, which we will use repeatedly, that the slopes of supporting hyperplanes of a convex function, bounded by R on a domain G, are bounded by $2R/\delta$ on any subdomain G' such that dist $(G', \partial G) \geq \delta$.

Let us show that the functions ϕ_k , $k \ge R$, $\phi_k(0) = 0$, are bounded on B_R uniformly in k. It is enough to show that $\nabla \phi_k$ are bounded uniformly on B_R . Suppose that there exists a sequence of points x_{k_j} in B_R such that $|\nabla \phi_{k_j}(x_{k_j})| \ge j$. Let $y_{k_j} = \nabla \phi_{k_j}(x_{k_j})$. Consider the unique solution ψ_k , $\psi_k(0) = 0$, to the equation

$$g_k(\nabla\psi_k) \det D_{ij}\psi_k = f_k(y)$$

mapping $B_{R_{k_j}}$ onto B_{k_j} . According to [3], $\nabla \psi_k$ is the inverse of $\nabla \phi_k$. Thus $\nabla \psi_{k_j}(y_{k_j}) = x_{k_j}$ and hence $|\nabla \psi_{k_j}(y_{k_j})| \leq R$. Observe that $\nabla \psi_k(0) = 0$ since $\nabla \psi_k$ is *K*-equivariant and the representation is irreducible. Let v_{k_j} be the unit vector in the direction y_{k_j} . Since ψ_k is convex, $\langle \nabla \psi_{k_j}(y), v_{k_j} \rangle \leq R$ for all $y \in [0, y_{k_j}]$ and therefore $|\psi_{k_j}(y)| \leq R|y|$ for such y. Using now Lemma 1.1 we conclude that for any r > 0, ψ_{k_j} is bounded by $\frac{R}{\epsilon}r$ on B_r for large j. It follows that there is a subsequence of ψ_{k_j} convergent to a convex solution $\psi : \mathbb{R}^n \to \mathbb{R}$ of

$$g(\nabla \psi) \det D_{ij} \psi = f.$$

The function ψ is bounded by Rr/ϵ on any B_r . Since ψ is convex, $\nabla \psi$ is bounded by $4R/\epsilon$ on $B_{r/2}$. Thus $U = \nabla \psi(\mathbb{R}^n)$ is contained in $B_{4R/\epsilon}$. However

$$+\infty = \int_{\mathbb{R}^n} f = \int_U g,$$

which leads to a contradiction. Theorem 0.1 is proved.

Proposition 1.2. Suppose, in addition, that $\int_{\mathbb{R}^n} g > 0$. Then any entire K-invariant convex solution ϕ of (0.1) is a proper function.

Proof. Suppose that there exists an unbounded sequence of points x_k such that $\phi(x_k)$ is bounded. Using convexity of ϕ and Lemma 1.1 we conclude that ϕ is a bounded function. This however implies that ϕ , being convex and defined on all of \mathbb{R}^n , is constant, which contradicts the assumption on g.

This proposition allows us to prove the regularity results of Theorem 0.2. The proof is essentially the same as the one given by Caffarelli and Viaclovsky [8]; we give it here for completeness.

Consider the convex (and bounded by the last proposition) sets $\Omega_c = \{x; \phi(x) \leq c\}, c \in \mathbb{R}$. For every c, we can find R(c) > c such that dist $(\Omega_c, \partial \Omega_{R(c)}) \geq 1$. By the argument used before, the slopes $\nabla \phi$ of ϕ are bounded by 2R(c) on Ω_c . If f and g are strictly positive and continuous, then equation (0.2) implies that there are constants λ_1, λ_2 , depending on c, such that

$$\lambda_1|B| \le |\nabla\phi(B)| \le \lambda_2|B| \tag{1.2}$$

for any Borel subset $B \subset \Omega_c$. We can apply Corollary 2 in [4] to Ω_{c+1} and conclude that ϕ is strictly convex at any point of Ω_c . Since c was arbitrary, ϕ is strictly convex everywhere. Now the main result of [6] implies that ϕ is locally of class $C^{1,\beta}$, where "locally" means that β can be chosen uniformly only on bounded subsets of \mathbb{R}^n . We observe that the only additional properties of f and g we have used so far is the boundedness of 1/f and 1/g. Therefore we have proved:

Proposition 1.3. In the situation of Theorem 0.1 suppose, in addition, that 1/f and 1/g are locally bounded functions and let ϕ be an entire K-invariant convex weak solution of (0.1). Then ϕ is of class $C^{1,\beta}$ on any compact subset P, where β may depend on P.

ROGER BIELAWSKI

To finish the proof of Theorem 0.2, assume that f and g are of class $C^{0,\alpha}$. Now, ϕ is a solution of det $D_{ij}\phi = \tilde{g}$, where $\tilde{g} = g/f(\nabla \phi)$. Since, by the last proposition, $\nabla \phi$ is Hölder continuous, \tilde{g} is (locally) of class $C^{0,\gamma}$. Using (1.2), we can apply Theorem 2 in [5] to $u(x) = \phi(x) - c$, $c \in \mathbb{R}$ and the convex and bounded set $\Omega_c = \{x; u(x) \leq 0\}$ to conclude that ϕ is locally of class $C^{2,\gamma}$ and hence globally C^2 . Therefore $\tilde{g} = g/f(\nabla \phi)$ is of class $C^{0,\alpha}$ everywhere and, repeating the argument, we conclude that ϕ is $C^{2,\alpha}$. Higher regularity is standard.

References

- I.J. Bakelman Convex analysis and nonlinear geometric elliptic equations, Springer-Verlag, Berlin, 1994.
- [2] R.Bielawski 'Kähler metrics on $G^{\mathbb{C}}$ ', J. Reine Angew. Math., to appear.
- [3] Y. Brenier 'Polar factorization and monotone rearrangement of vector-valued functions', Comm. Pure Appl. Math. 44 (1991), 375–417.
- [4] L.A. Caffarelli 'A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity', Ann. of Math. (2) 131 (1990), 129–134.
- [5] L.A. Caffarelli 'Interior W^{2,p} estimates for solutions of the Monge-Ampère equation', Ann. of Math. (2) 131 (1990), 135–150.
- [6] L.A. Caffarelli 'Some regularity properties of solutions of Monge-Ampère equation', Comm. Pure Appl. Math. 44 (1991), 965–969.
- [7] L.A. Caffarelli 'The regularity of mappings with a convex potential', J. Amer. Math. Soc. 5 (1992), 99–104.
- [8] L.A. Caffarelli and J.A. Viaclovsky 'On the regularity of solutions to Monge-Ampre equations on Hessian manifolds', Comm. Partial Differential Equations 26 (2001), no. 11-12, 2339–2351.
- [9] K.-S. Chou and X.-J. Wang 'Entire solutions of the Monge-Ampère equation', Comm. Pure Appl. Math. 49 (1996), 529–539.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, GLASGOW G12 8QW, UK *E-mail address*: R.Bielawski@maths.gla.ac.uk