

On complex and noncommutative torus

Igor Nikolaev
 Department of Mathematics
 2500 University Drive N.W.
 Calgary T2N 1N4 Canada
 nikolaev@math.ucalgary.ca

June 18, 2019

Abstract

To every non-singular elliptic curve (complex torus) we assign a C^* -algebra $T_\theta = \{u, v \mid vu = e^{2\pi i\theta} uv\}$ known as noncommutative torus. It is shown that morphisms of elliptic curves generate Morita equivalence of the corresponding noncommutative tori. Real number θ we call projective curvature attached to the elliptic curve. It is proved that projective curvatures of isomorphic elliptic curves are modular equivalent numbers. We prove that projective curvature of elliptic curve with complex multiplication is a quadratic irrationality given by periodic continued fraction.

Key words and phrases: elliptic curve, noncommutative torus

AMS (MOS) Subj. Class.: 14H52, 46L85

Introduction

It is recognized that noncommutative geometry studies algebraic varieties over the noncommutative rings modulo their Morita equivalence. The origin of the subject goes back to F. Klein, who introduced noncommutative torus in a paper devoted to lattices and continued fractions ([5], see also [6]). Murray and von Neumann ([11]) discovered continuous dimensions in the noncommutative rings. This phenomenon found a spectacular application in the works of V. F. R. Jones on the classification of knots and links in the 3-dimensional space ([4]). The fundamental book of A. Connes ([1]) gives an account of the current state and perspective of the subject.

It was noticed a long time ago that Riemann surfaces is a discrete object. Let Δ be the Laplace-Beltrami operator and consider the eigenvalue problem

$\Delta u + \lambda u = 0$ on a compact Riemann surface X . It is known that $\text{Spec } X$ is discrete and consists of the eigenvalues

$$0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots, \quad \lim_{n \rightarrow \infty} \lambda_n = \infty. \quad (1)$$

One can “hear the shape of drum” in the sense that $\text{Spec } X$ determines the conformal structure of X . From this point, $\text{Spec } X$ can be viewed as a moduli set of surface X .

Let $Sp X$ be length spectrum of X , i.e. the set of lengths of all simple closed geodesics of X counting their multiplicities. Length spectrum is a discrete subset of positive real line \mathbb{R}^+ . The Selberg trace formula indicates that $Sp X$ also determines the conformal structure of X . In fact, it was shown by Wolpert that mapping $X \rightarrow Sp X$ is finite-to-one and generically one-to-one ([16]). In the same time, it is clear that total number of independent lengths of $Sp X$ cannot exceed $6g - 6$.

It is known that each homotopy class of closed curves on X has a geodesic representative. One can construct a sequence of geodesics of growing length convergent to a measured lamination on X : $\lim_{l \rightarrow \infty} Sp X = \Lambda$. The real dimension of space of measured laminations is equal to $6g - 6$ ([15]). Thus Λ can be viewed as moduli set of Riemann surface X . Irrational number θ equal to the average slope (“rotation number”) of geodesics in Λ is called *projective curvature* of X .

The present note concerns the case $X = E$, where E is a Riemann surface of genus one. In this case $E = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ is a complex torus and $\text{Spec } E, Sp E$ can be expressed through the complex modulus τ (Poisson summation formula). It can be shown that projective curvature $\theta = \theta(\tau)$ unfolds into an infinite continued fraction

$$\theta = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}} \quad (2)$$

whose positive integers a_i define noncommutative torus T_θ , i.e. a C^* -algebra on two generators u, v and relation $vu = \exp(2\pi i\theta)uv$. Under this equivalence, isomorphic complex tori correspond to Morita equivalent noncommutative tori.

When E admits complex multiplication, continued fraction for θ becomes periodic and θ real quadratic. The problem of calculation of the maximal abelian extensions of number fields was posed by L. Kronecker (Kronecker’s *Jugendtraum*). It was generalized later as Hilbert’s 12th problem. Theory of elliptic curves with complex multiplication solves the problem for the imaginary quadratic fields, cf ([8], p. 202). It was conjectured (Shimura and Stark) that for extensions of real number fields, there might exist similar theory based on the idea of “real multiplication”. Manin suggested to use “pseudolattices”

(a.k.a. dimension groups of noncommutative tori) to settle the real multiplication problem ([7]).

Present study was inspired by this bunch of problems, noncommutative geometry, and also works of Huber ([3]) and Selberg ([13]) on ζ -functions describing the asymptotic of closed geodesics on Riemann surfaces. We prove the following results.

Theorem 1 *To every non-singular elliptic curve (complex torus) there corresponds a noncommutative torus T_θ such that isomorphic curves $E \sim E'$ generate Morita equivalent tori $T_\theta \sim T_{\theta'}$. Irrational parameter θ is called projective curvature of elliptic curve and two such curves E, E' are isomorphic if and only if*

$$\theta' = \frac{a\theta + b}{c\theta + d} \quad ad - bc = \pm 1, \quad a, b, c, d \in \mathbb{Z}. \quad (3)$$

Projective curvature $\theta = \theta(\tau)$ is a continuous function of the complex modulus τ of elliptic curve E .

Theorem 2 *Projective curvature of an elliptic curve with the complex multiplication is a quadratic irrationality.*

Contents

1	Proofs	3
1.1	Proof of Theorem 1	3
1.2	Proof of Theorem 2	7
2	Background information	9
2.1	Noncommutative torus	9
2.2	Complex torus	11
2.3	Elliptic curves with complex multiplication	12

1 Proofs

The proof of both theorems is based on the rigidity of length spectrum of complex torus, cf Wolpert ([16]). The preliminary information on complex and noncommutative tori can be found in Section 2.

1.1 Proof of Theorem 1

Let us review main steps of the proof. By the rigidity lemma (Lemma 1) the length spectrum $Sp E$ defines conformal structure of E . In fact, this correspondence is a bijection. Under isomorphisms of E the length spectrum can acquire a real multiple or get a “cut of finite tail” (Lemma 2). One can relate

an infinite continued fraction to $Sp E$ using Minkowski's decomposition of the automorphisms $SL(2, \mathbb{Z})$ of lattice $L = \mathbb{Z} + \tau\mathbb{Z}$ (Lemma 3). Then isomorphic complex tori \mathbb{C}/L will have continued fractions which differ only in a finite number of terms. In other words, one can attach a Morita equivalence class of noncommutative tori to every isomorphism class of complex tori.

Lemma 1 (Rigidity of length spectrum) *Let $Sp E$ be length spectrum of a complex torus $E = \mathbb{C}/L$. Then there exists a unique complex torus with the spectrum. This correspondence is a bijection.*

Proof. Selberg ([13]) and Huber ([3]) proved that length spectrum $Sp X$ determines the eigenvalue spectrum $Spec E$ of any compact Riemann surface X (and *vice versa*). Let complex torus $E = \mathbb{C}/L$ be given by its complex modulus $\tau = \frac{\omega_1}{\omega_2} = a + ib$. One can apply the Poisson summation formula for the eigenvalues of Laplacian Δ :

$$\sum_{L'} e^{-4\pi^2 |\omega'|^2 t} = \frac{Area(E)}{4\pi t} \sum_L e^{-\frac{|\omega|^2}{4t}}, \quad (4)$$

where L' is the lattice dual to L and $\omega' \in L'$. This allows to calculate $a = Area E$ and b through unfolding the Poisson formula into the power series (McKean [9]). \square

Let $Sp X = \{l_1, l_2, \dots\}$ be length spectrum of a Riemann surface X . Let $a > 0$ be a real number. By $aSp X$ we understand the length spectrum $\{al_1, al_2, \dots\}$. Similarly, for any $m \in \mathbb{N}$ we denote by $Sp_m X$ the length spectrum $\{l_m, l_{m+1}, \dots\}$, i.e. the one obtained by deleting the first $(m-1)$ -geodesics in $Sp X$.

Lemma 2 *Let $E \sim E'$ be isomorphic complex tori. Then either:*

- (i) $Sp E' = |\alpha| Sp E$ for an $\alpha \in \mathbb{C}^\times$, or
- (ii) $Sp E' = Sp_m E$ for a $m \in \mathbb{N}$.

Proof. (i) The complex tori $E = \mathbb{C}/L, E' = \mathbb{C}/M$ are isomorphic if and only if $M = \alpha L$ for a complex number $\alpha \in \mathbb{C}^\times$. It is not hard to see that closed geodesic of E are bijective with the points of the lattice $L = \omega_1\mathbb{Z} + \omega_2\mathbb{Z}$ in the following way. Take a segment of straight line through points 0 and ω of lattice L which contains no other points of L . This segment represents a homotopy class of curves through 0 and a closed geodesic of E . Evidently, this geodesic will be the shortest in its homotopy class with the length $|\omega|$ equal to absolute value of complex number ω . Thus, $|\omega|$ belongs to the length spectrum of E .

Let now $Sp E = \{|\omega_1|, |\omega_2|, \dots\}$ with $\omega_i \in L$. Since $M = \alpha L$, one gets $Sp E' = \{|\alpha||\omega_1|, |\alpha||\omega_2|, \dots\}$ and $Sp E' = |\alpha| Sp E$. Item (i) follows.

(ii) Note that according to (i) the length spectrum $Sp X = \{l_0, l_1, l_2, \dots\}$ can be written as $Sp X = \{1, l_1, l_2, \dots\}$ after multiplication on $1/l_0$, where l_0

is the length of shortest geodesic. Note also that shortest geodesic of complex torus has homotopy type $(1, 0)$ or $(0, 1)$ (standard generators for $\pi_1 E$).

Let a, b, c, d be integers such that $ad - bc = \pm 1$ and let

$$\begin{aligned}\omega'_1 &= a\omega_1 + b\omega_2, \\ \omega'_2 &= c\omega_1 + d\omega_2,\end{aligned}\tag{5}$$

be an automorphism of the lattice $L = \omega_1\mathbb{Z} + \omega_2\mathbb{Z}$. This automorphism maps standard generators $(1, 0)$ and $(0, 1)$ of L to the vectors $\omega_1 = (a, b), \omega_2 = (c, d)$. Let their lengths be l_m, l_{m+1} , respectively.

As we showed earlier, $l_m, l_{m+1} \in Sp E$ and it is not hard to see that there are no geodesics of the intermediate length. (This gives a justification for the notation chosen.) Note that ω_1, ω_2 are standard generators for the complex torus $E' \sim E$ and therefore one of them is the shortest closed geodesics of E' . One can normalize it to the length 1.

On the other hand, there are only finite number of closed geodesics of length smaller than l_n (McKean [9]). Thus $Sp E \cap Sp E' = \{l_m, l_{m+1}, \dots\}$ for a finite number m and since (5) is automorphism of the lattice L . In other words, $Sp E' = Sp_m E$. Item (ii) follows. \square

Lemma 3 *Let $E = E(\tau)$ be a complex torus. There exists a uniquely defined continued fraction θ of form (2) corresponding to E . When $E \sim E'$ are isomorphic tori (i.e. $\tau' = \frac{a\tau+b}{c\tau+d}$, $a, b, c, d \in \mathbb{Z}$, $ad - bc = \pm 1$), then fractions θ and θ' differ only in a finite number of terms.*

Proof. Let $E = \mathbb{C}/L$ be complex torus and $Sp E = \{l_1, l_2, \dots\}$ be its length spectrum, i.e. a set of non-decreasing lengths of closed geodesics $\{\gamma_1, \gamma_2, \dots\}$. To prove the lemma, one needs to introduce a real number, θ , which measures an asymptotic slope (or direction) of the “ordered bunch” of the geodesics $\{\gamma_1, \gamma_2, \dots\}$. We wish to stress the word *ordered* here, since non-ordered sets of periodic geodesics are directionally equi-distributed (by transitivity of the geodesic flow). We shall see later that the order is critical to the value of θ . Morally, θ plays the role similar to Selberg’s zeta function for the periodic geodesics, see Huber ([3]) and Selberg ([13]).

If $\{\gamma_1, \gamma_2, \dots\}$ are closed geodesics specified by $Sp E$ as explained above, we let $\{\omega_1, \omega_2, \dots\}$ be lattice points $\omega_i \in L$ which represent closed geodesics γ_i . Two consequent points $\omega_k, \omega_{k+1} \in L$ define (uniquely) an automorphism of L as follows. Take the 1-dimensional subset $\Omega_k \subset L$ consisting of points collinear to ω_k , i.e. points lying on the line $t\omega_k, t \in \mathbb{R}$. The same notation applies to Ω_{k+1} . Set an automorphism $g_k \in SL(2, \mathbb{Z})$ such that:

$$\Omega_{k+1} = g_k(\Omega_k).\tag{6}$$

The above automorphism is unique, up to a flip. To avoid flips, one can consider orientation preserving automorphisms, e.g. this can be accomplished by fixing $\det g_{k+1} = +1$.

It is well known that every ordered sequence of $g_i \in SL(2, \mathbb{Z})$ unfolds into a continued fraction, cf Minkowski ([10]). The convergent of this fraction can be regarded as “directional averaging” of the geodesic sequence $\{\gamma_1, \gamma_2, \dots\}$. Namely, every matrix $(g_k) \in SL(2, \mathbb{Z})$ admits a unique decomposition

$$\begin{pmatrix} -q & \mp s \\ \pm p & r \end{pmatrix} = \begin{pmatrix} 0 & \mp 1 \\ \pm 1 & a_k^{(1)} \end{pmatrix} \begin{pmatrix} 0 & \pm 1 \\ \mp 1 & a_k^{(2)} \end{pmatrix} \cdots \begin{pmatrix} 0 & \mp (-1)^{m_k} \\ \pm (-1)^{m_k} & a_k^{(m_k)} \end{pmatrix}, \quad (7)$$

cf. Minkowski ([10], p. 160). Consider the limit

$$\theta = \lim_{N \rightarrow \infty} \prod_{k=1}^N (g_k) = a_1^{(1)} + \frac{1}{a_1^{(2)} + \frac{1}{a_1^{(3)} + \dots}}, \quad (8)$$

where $(a_k^{(1)}, a_k^{(2)}, \dots, a_k^{(m_k)})$ is the Minkowski decomposition of the automorphism g_k . Note that entries $a_k^{(i)}$ are positive integers, so that continued fraction (8) is regular. Such a fraction converges to an irrational number θ , which we call a *projective curvature* of complex torus E . The first part of Lemma 3 is proved.

The second part of Lemma 3 follows from item (ii) of Lemma 2 and Lemma 1. Indeed, if $E \sim E'$ are isomorphic complex tori, then by Lemma 2, (i) $Sp E' = |\alpha| Sp E$ or (ii) $Sp E' = Sp_m E$ for an $\alpha \in \mathbb{C}^\times$ and $m \in \mathbb{N}$. Note that since multiplication of L by complex number α does not affect the one-dimensional subspaces Ω_i (see above for the notation), the ordered sequence of automorphisms g_i will be the same. Thus the possibility (i) will give the same number θ .

In case (ii) it can be easily seen that the ordered sequence of automorphisms $(g_1, g_2, \dots, g_m, g_{m+1}, \dots)$ corresponding to $Sp E$ will become (g_m, g_{m+1}, \dots) for $Sp E'$. Since there is a bijection between length spectra and complex tori (Lemma 1), we get that $E \sim E'$ would correspond to continued fractions which differ only in the first m -terms. In other words, projective curvatures θ and θ' are modular equivalent irrational numbers. The second part of Lemma 3 follows. \square

To finish the proof of the first part of Theorem 1, one needs to combine Lemmas 1-3 with the fact that two noncommutative tori $T_\theta, T_{\theta'}$ are Morita equivalent if and only if their continued fractions differ only in a finite number of terms (Section 2.1).

Let us prove continuity of function $\theta = \theta(\tau)$. We wish to give a geometric proof of this fact, an analytic argument will be published elsewhere later on. Let T_1 be the Teichmüller space of complex torus. Suppose that $\phi : T \rightarrow T$ a quasiconformal map of complex torus T . It is known that ϕ acts continuously on the lamination $\Lambda = \lim_{t \rightarrow \infty} Sp T$ ([15]). Since θ is an “average slope” of Λ

on T , θ is continuous on ϕ . On the other hand, every two points (conformal structures) of T_1 can be joined by an orbit of the Teichmüller flow, ϕ^t , generated by map ϕ . Thus $\theta = \theta(\phi^t(T_1)) = \theta(\tau)$ is a continuous function. Theorem 1 is proven. \square

1.2 Proof of Theorem 2

Let us outline the idea of the proof. If E admits complex multiplication, then its complex modulus τ lies in an imaginary quadratic field K . In fact, up to an isogeny, the ring of endomorphisms $\text{End } E = \mathcal{O}_K$, where \mathcal{O}_K is the ring of integers of field K . It can be shown that L is an ideal in \mathcal{O}_K (Section 2.3). The length spectrum $Sp E$ of elliptic curve with complex multiplication is a “geometric progression” with the growth rate $|\alpha|$, where $\alpha \in \text{End } E$ (Lemma 4). One can use Klein’s lemma (Lemma 5) to characterize length spectra in terms of continued fractions. In particular, length spectrum with asymptotically geometric growth correspond to periodic continued fractions. Thus, projective curvature converges to quadratic irrationality.

Definition 1 *Length spectrum $Sp E$ of an elliptic curve $E = \mathbb{C}/L$ is called α -multiplicative, if there exists a complex number $\alpha \in \mathbb{C}^\times$ with $|\alpha| > 1$ such that*

$$Sp E = \{l_1, \dots, l_N, |\alpha|l_1, \dots, |\alpha|l_N, \dots, |\alpha|^n l_1, \dots, |\alpha|^n l_N, \dots\}, \quad (9)$$

for a $N \in \mathbb{N}$.

Lemma 4 *Let E be an elliptic curve with complex multiplication. Then its length spectrum $Sp E$ is α -multiplicative for an $\alpha \in \mathbb{C}^\times$.*

Proof. Let $E = \mathbb{C}/L$ be complex torus which admits non-trivial endomorphisms $z \mapsto \alpha z, \alpha \in K = \mathbb{Q}(\sqrt{-d})$. It is known that $\text{End } E$ is an order in the field K . In fact, up to an isogeny of E , $\text{End } E \simeq \mathcal{O}_K$, where \mathcal{O}_K is the ring of integers of imaginary quadratic field K (Section 2.3). Lattice L in this case corresponds to an ideal in \mathcal{O}_K .

Let l_1 be minimal length of closed geodesic of E . For an endomorphism $\alpha : E \rightarrow E, \alpha \in \mathbb{C}^\times$, consider the set of geodesics whose lengths are less than $|\alpha|l_1$. By the properties of $Sp E$ mentioned in Section 1.1, such a set will be finite. Let us denote the lengths of geodesics in this set by l_1, \dots, l_N . Since every geodesic in $Sp E$ is a complex number ω_i lying in the ring $L \subseteq \mathcal{O}_K$, one can consider the set of geodesics $\alpha\omega_1, \dots, \alpha\omega_N$. The length of these geodesics will be $|\alpha|l_1, \dots, |\alpha|l_N$, respectively. It is not hard to see that by the choice of number N , the first $2N$ elements of $Sp E$ are presented by the following growing sequence of geodesics: $l_1, \dots, l_N, |\alpha|l_1, \dots, |\alpha|l_N$. We proceed by iterations of α , until all closed geodesics of E are exhausted. The conclusion of Lemma 4 follows. \square

We shall need the following statement regarding geometry of the regular continued fractions, see F. Klein ([5],[6]). It is valid for any regular fraction, not necessarily periodic.

Lemma 5 (F. Klein) *Let*

$$\omega = \mu_1 + \frac{1}{\mu_2 + \frac{1}{\mu_3 + \dots}} \quad (10)$$

be a regular continued fraction. Let us denote the convergents of ω by:

$$\frac{p_{-1}}{q_{-1}} = \frac{0}{1}, \quad \frac{p_0}{q_0} = \frac{1}{0}, \quad \frac{p_1}{q_1} = \frac{\mu_1}{1}, \dots, \frac{p_\nu}{q_\nu} = \frac{\mu_\nu p_{\nu-1} + p_{\nu-2}}{\mu_\nu q_{\nu-1} + q_{\nu-2}}. \quad (11)$$

For any lattice L in \mathbb{C} , consider a segment I with ends in the points $(p_{\nu-2}, q_{\nu-2})$ and (p_ν, q_ν) . Then the segment J which joins 0 with the point $p_{\nu-1}, q_{\nu-1}$ is parallel to I and

$$|I| = \mu_\nu |J|, \quad (12)$$

where $|\bullet|$ denotes the length of the segment.

Proof. For the original formulation of this lemma, we refer the reader to ([6]). Let us outline the idea of the proof. Draw the lines through (p_ν, q_ν) , $(p_{\nu-1}, q_{\nu-1})$ and $(0, 0)$, $(p_{\nu-2}, q_{\nu-2})$. They intersect in a point A , which is common vertex of two congruent triangles with bases I and J . By formula (11) the proportion of the above triangles is equal to μ_ν . Lemma 5 follows. \square

Corollary 1 *Let $\omega_\nu = (p_\nu, q_\nu)$ be lattice points mentioned in Lemma 5. Then the length of vector ω_ν can be evaluated with the help of the following asymptotic formula:*

$$|\omega_\nu| \approx |\omega_{\nu-2}| + \mu_\nu |\omega_{\nu-1}|. \quad (13)$$

Proof. Indeed, using notation of Lemma 5, one can write $|(p_\nu, q_\nu)| \approx |(p_{\nu-2}, q_{\nu-2})| + |I|$. But according to equation (12), $|I| = \mu_\nu |(p_{\nu-1}, q_{\nu-1})|$. Corollary 1 follows. \square

Note that according to the recurrent formula (13) the length spectrum $\{|\omega_\nu|\}$ coming from continued fraction (10) is completely determined by the first two values: $|\omega_1|$ and $|\omega_2|$. Using (13), one can easily deduce the following asymptotic formula for $|\omega_\nu|$ as function of $|\omega_1|, |\omega_2|$:

$$|\omega_\nu| \approx |\omega_2| \prod_{k=3}^{\nu} \mu_k + |\omega_1| \prod_{k=4}^{\nu} \mu_k + O(\nu). \quad (14)$$

Fix N a positive integer. It follows from equation (14) that:

$$\begin{aligned} \lim_{\nu \rightarrow \infty} \frac{|\omega_{\nu+N}|}{|\omega_\nu|} &= \mu_{\nu+1} \dots \mu_{\nu+N} \lim_{\nu \rightarrow \infty} \left(\frac{\mu_\nu \dots \mu_3 |\omega_2| + \mu_\nu \dots \mu_4 |\omega_1| + O(\nu)}{\mu_\nu \dots \mu_3 |\omega_2| + \mu_\nu \dots \mu_4 |\omega_1| + O(\nu)} \right) = \\ &= \mu_{\nu+1} \dots \mu_{\nu+N}. \end{aligned} \quad (15)$$

Let E be an elliptic curve with complex multiplication. Then by Lemma 4 its length spectrum $Sp\ E$ is α -multiplicative. In other words,

$$\frac{l_{\nu+N}}{l_{\nu}} = |\alpha| = \text{Const}, \quad (16)$$

for an $N \in \mathbb{N}$ and any $\nu \bmod N$. Note that $|\alpha|$ is a rational integer. Thus, by formula (15) we have $\mu_{\nu+1} \dots \mu_{\nu+N} = \text{Const}$, for any $\nu \bmod N$. The last requirement can be satisfied if and only if continued fraction (10) is N -periodic. Theorem 2 is proven. \square

2 Background information

In present section we briefly review noncommutative and complex tori. The excellent source of information on noncommutative torus are the article of Effros and Shen ([2]) and the monograph of Rørdam, Larsen and Laustsen ([12]). The literature on complex torus is fairly vast. We would recommend for the reference Chapter VI of Silverman's book ([14]).

2.1 Noncommutative torus

By the C^* -algebra one understands a noncommutative Banach algebra with an involution ([12]). Namely, a C^* -algebra A is an algebra over \mathbb{C} with a norm $a \mapsto \|a\|$ and an involution $a \mapsto a^*$, $a \in A$, such that A is complete with respect to the norm, and such that $\|ab\| \leq \|a\| \|b\|$ and $\|a^*a\| = \|a\|^2$ for every $a, b \in A$. If A is commutative, then the Gelfand theorem says that A is isometrically $*$ -isomorphic to the C^* -algebra $C_0(X)$ of continuous complex-valued functions on a locally compact Hausdorff space X . For otherwise, A represents a “noncommutative” topological space X .

K_0 and dimension groups. Given a C^* -algebra, A , consider new C^* -algebra $M_n(A)$, i.e. the matrix algebra over A . There exists a remarkable semi-group, A^+ , connected to the set of projections in algebra $M_{\infty} = \bigcup_{n=1}^{\infty} M_n(A)$. Namely, projections $p, q \in M_{\infty}(A)$ are Murray-von Neumann equivalent $p \sim q$ if they can be presented as $p = v^*v$ and $q = vv^*$ for an element $v \in M_{\infty}(A)$. The equivalence class of projections is denoted by $[p]$. The semi-group A^+ is defined to be the set of all equivalence classes of projections in $M_{\infty}(A)$ with the binary operation $[p] + [q] = [p \oplus q]$. The Grothendieck completion of A^+ to an abelian group is called a K_0 -group of A . The functor $A \rightarrow K_0(A)$ maps the unital C^* -algebras into the category of abelian groups so that the semi-group $A^+ \subset A$ corresponds to a “positive cone” $K_0^+ \subset K_0(A)$ and the unit element $1 \in A$ corresponds to the “order unit” $[1] \in K_0(A)$. The ordered abelian group $(K_0, K_0^+, [1])$ with the order unit is called a *dimension (Elliott) group* of A . The dimension (Elliott) group is complete invariant of the AF C^* -algebras to be considered in the next paragraph.

AF C^* -algebras. An *AF* (approximately finite-dimensional) algebra is defined to be a norm closure of an ascending sequence of the finite dimensional algebras M_n 's, where M_n is an algebra of $n \times n$ matrices with the entries in \mathbb{C} . Here the index $n = (n_1, \dots, n_k)$ represents a *multi-matrix* algebra $M_n = M_{n_1} \oplus \dots \oplus M_{n_k}$. Let

$$M_1 \xrightarrow{\varphi_1} M_2 \xrightarrow{\varphi_2} \dots, \quad (17)$$

be a chain of algebras and their homomorphisms. A set-theoretic limit $A = \lim M_n$ has a natural algebraic structure given by the formula $a_m + b_k \rightarrow a + b$; here $a_m \rightarrow a, b_k \rightarrow b$ for the sequences $a_m \in M_m, b_k \in M_k$. The homeomorphisms of the above (multi-matrix) algebras admit a canonical description ([12]). Suppose that $p, q \in \mathbb{N}$ and $k \in \mathbb{Z}^+$ are such numbers that $kq \leq p$. Let us define a homomorphism $\varphi : M_q \rightarrow M_p$ by the formula

$$a \mapsto \underbrace{a \oplus \dots \oplus a}_k \oplus 0_h, \quad (18)$$

where $p = kq + h$. More generally, if $q = (q_1, \dots, q_s), p = (p_1, \dots, p_r)$ are vectors in $\mathbb{N}^s, \mathbb{N}^r$, respectively, and $\Phi = (\phi_{kl})$ is a $r \times s$ matrix with the entries in \mathbb{Z}^+ such that $\Phi(q) \leq p$, then the homomorphism φ is defined by the formula:

$$\begin{aligned} a_1 \oplus \dots \oplus a_s &\longrightarrow \underbrace{(a_1 \oplus a_1 \oplus \dots)}_{\phi_{11}} \oplus \underbrace{(a_2 \oplus a_2 \oplus \dots)}_{\phi_{12}} \oplus \dots \oplus 0_{h_1} \\ &\oplus \underbrace{(a_1 \oplus a_1 \oplus \dots)}_{\phi_{21}} \oplus \underbrace{(a_2 \oplus a_2 \oplus \dots)}_{\phi_{22}} \oplus \dots \oplus 0_{h_2} \oplus \dots \end{aligned} \quad (19)$$

where $\Phi(q) + h = p$. We say that φ is a *canonical homomorphism* between M_p and M_q . Any homomorphism $\varphi : M_q \rightarrow M_p$ can be rendered canonical ([12]).

Noncommutative torus. Fix θ irrational and consider a linear flow $\dot{x} = \theta, \dot{y} = 1$ on the torus. Let S^1 be a closed transversal to our flow. The *noncommutative torus* T_θ is a norm-closed C^* -algebra generated by the unitary operators in the Hilbert space $L^2(S^1)$:

$$Uf(t) = z(t)f(t), \quad Vf(t) = f(t - \alpha),$$

which are multiplication by a unimodular function $z(t)$ and rotation operators. It could be easily verified that $UV = e^{2\pi i \alpha} VU$. As an “abstract” algebra, T_θ is a crossed product C^* -algebra $C(S^1) \rtimes_\phi \mathbb{Z}$ of (commutative) C^* -algebra of complex-valued continuous functions on S^1 by the action of powers of ϕ , where ϕ is a rotation of S^1 through the angle $2\pi\alpha$. T_θ is not *AF*, but can be embedded into an *AF*-algebra whose dimension group is P_θ (to be specified below); the latter is known to be intimately connected with the arithmetic of the irrational numbers θ 's. The following beautiful result is due to the efforts

of many mathematicians ¹ (Effros, Elliott, Pimsner, Rieffel, Shen, Voiculescu, etc):

Theorem 3 (Classification of noncommutative tori) *Let T_θ be a noncommutative torus. Suppose that the θ has a continued fraction expansion*

$$\theta = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}} \stackrel{\text{def}}{=} [a_0, a_1, a_2, \dots].$$

Let φ_n be a composition of isometries of the lattice $\mathbb{Z}^2 \subset \mathbb{R}^2$: $\varphi_n = \begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \dots \begin{pmatrix} a_n & 1 \\ 1 & 0 \end{pmatrix}$. Then T_θ can be embedded into an AF-algebra whose dimension group is a direct limit of the ordered abelian groups: $P_\theta = \lim_{n \rightarrow \infty} (\mathbb{Z}^2, \varphi_n)$. Moreover, if $\theta = [a_0, a_1, \dots]$ and $\theta' = [b_0, b_1, \dots]$ are two irrational numbers, then P_θ and $P_{\theta'}$ are isomorphic (i.e. noncommutative tori T_θ and $T_{\theta'}$ are Morita equivalent) if and only if $a_{m+k} = b_m$ for an integer number $k \in \mathbb{Z}$. In other words, the irrational numbers θ and θ' are modular equivalent: $\theta' = \frac{a\theta+b}{c\theta+d}$, $ad-bc = \pm 1$, where $a, b, c, d \in \mathbb{Z}$ are integer numbers.

Proof. An algebraic proof of this fact can be found in ([2]). \square

2.2 Complex torus

Let L denote a lattice in the complex plane \mathbb{C} . Attached to L , there are following classic Weierstrass function $\wp(z; L)$ and Eisenstein series $G_k(L)$:

$$\wp(z; L) = \frac{1}{z^2} + \sum_{\omega \in L^\times} \left\{ \frac{1}{(z + \omega)^2} - \frac{1}{\omega^2} \right\}, \quad (20)$$

$$G_k(L) = \sum_{\omega \in L^\times} \frac{1}{\omega^{2k}}. \quad (21)$$

$\wp(z; L)$ is analytic and $G_k(L)$ is convergent for any lattice L ([14]). There exists a duality between lattices L and cubic curves E given by the following theorem.

Theorem 4 *Let L be a lattice in \mathbb{C} . Then the map $z \mapsto (\wp(z; L), \frac{1}{2}\wp'(z; L))$ is an analytic isomorphism from complex torus \mathbb{C}/L to elliptic cubic $E = E(\mathbb{C})$:*

$$E(\mathbb{C}) = \{(x, y) \in \mathbb{C}^2 \mid y^2 = x^3 - 15G_4(L)x - 35G_6(L)\}. \quad (22)$$

Conversely, to any cubic in the Weierstrass normal form $y^2 = x^3 + ax + b$ there corresponds a unique lattice L such that $a = -15G_4(L)$ and $b = -35G_6(L)$.

¹The author apologizes for possible erroneous credits regarding history of the problem. Classification of noncommutative tori seems to be an old problem; early results in this direction can be found in the works of Felix Klein ([5],[6]).

Proof. We refer the reader to ([14]) for a detailed proof of this fact. \square

Isomorphism of complex tori. Let L be a lattice in \mathbb{C} . The Riemann surface \mathbb{C}/L is called a *complex torus*. Let $f : \mathbb{C}/L \rightarrow \mathbb{C}/M$ be holomorphic and invertible map (isomorphism) between two complex tori. Since f is covered by a linear map $z \rightarrow \alpha z$ on \mathbb{C} , one can easily conclude that $\alpha L = M$ for an $\alpha \in \mathbb{C}^\times$. On the other hand, lattice L can always be written as $L = \omega_1 \mathbb{Z} + \omega_2 \mathbb{Z}$, where $\omega_1, \omega_2 \in \mathbb{C}^\times$ and $\omega_2 \neq k\omega_1$ for a $k \in \mathbb{R}$. The complex number $\tau = \frac{\omega_2}{\omega_1}$ is called a *complex modulus* of lattice L .

Lemma 6 *Two complex tori are isomorphic if and only if their complex moduli τ and τ' satisfy the equation:*

$$\tau' = \frac{a\tau + b}{c\tau + d} \quad ad - bc = \pm 1, \quad a, b, c, d \in \mathbb{Z}. \quad (23)$$

Proof. Let $L = \omega_1 \mathbb{Z} + \omega_2 \mathbb{Z}$ be a lattice. Notice that if $\omega_1 \neq 1$, one can pick up an isomorphic lattice $L' = \frac{1}{\omega_1} L$ and therefore we assume $\omega_1 = 1$ and $\omega_2 = \tau$. Let $g : L \rightarrow L$ be an automorphism of lattice L . The new coordinate system (ω'_1, ω'_2) of lattice L satisfy the equations:

$$\begin{aligned} \omega'_1 &= a\omega_1 + b\omega_2, \\ \omega'_2 &= c\omega_1 + d\omega_2, \end{aligned} \quad (24)$$

where $ad - bc = \pm 1$, $a, b, c, d \in \mathbb{Z}$. It is clear in the same time that isomorphism class of complex torus shouldn't depend on the coordinate system chosen in L . We divide the second equation by first equation in (23) and the conclusion of Lemma 6 follows. (It is not hard to see that this statement works in the opposite direction as well.) \square

One may wish to express the isomorphism criterion (23) in terms of the coefficients a, b of the elliptic cubic $y^2 = x^3 + ax + b$. Using the duality between lattices and curves (Theorem 4), the following lemma can be proved.

Lemma 7 *Let $y^2 = x^3 + ax + b$ be an elliptic cubic E . Set*

$$j(E) = -\frac{1728 \times 4a^3}{4a^3 + 27b^2}. \quad (25)$$

Then elliptic cubics E, E' are isomorphic if and only if $j(E) = j(E')$.

Proof. The proof of this fact can be found in ([14]). \square

2.3 Elliptic curves with complex multiplication

Let $E = \mathbb{C}/L$ be an elliptic curve. Consider the set $\text{End } E$ of analytic self-mappings of E . Each $f \in \text{End } E$ is covered on the complex plane by map

$z \mapsto \alpha z$ for an $\alpha \in \mathbb{C}$. It is not hard to see that $\text{End } E$ has the structure of a ring under the pointwise addition and multiplication of functions. The set $\text{End } E$ is called an *endomorphism ring* of elliptic curve E . By the remarks above, $\text{End } E$ can be thought of as a subring of complex numbers:

$$\text{End } E = \{\alpha \in \mathbb{C} \mid \alpha L \subseteq L\}. \quad (26)$$

There exists a fairly complete algebraic description of such rings. Roughly speaking, they are either “rational integers” \mathbb{Z} or integers \mathcal{O}_K of an algebraic number field K . The following lemma is true.

Lemma 8 *Let $\alpha \in \text{End } E$ be a complex number. Then either:*

- (i) α is a rational integer, or
- (ii) α is an algebraic integer in an imaginary quadratic number field $K = \mathbb{Q}(\sqrt{-d})$.

Proof. Let $L = \omega_1\mathbb{Z} + \omega_2\mathbb{Z}$ be lattice corresponding to E . If $\alpha \in \text{End } E$, then $\alpha L \subseteq L$. The latter inclusion means that there exists integers a, b, c, d , (not necessarily with $ad - bc = \pm 1$) such that:

$$\begin{aligned} \alpha\omega_1 &= a\omega_1 + b\omega_2, \\ \alpha\omega_2 &= c\omega_1 + d\omega_2. \end{aligned} \quad (27)$$

(i) Let $\alpha \in \mathbb{R}$. Then one gets a homogeneous linear system with respect to ω_1, ω_2 : $(a - \alpha)\omega_1 + b\omega_2 = 0$, $c\omega_1 + (d - \alpha)\omega_2 = 0$. Since $\omega_2 \neq k\omega_1$ for a $k \in \mathbb{R}$, one concludes that the only solution is $a - \alpha = b = c = d - \alpha = 0$. Therefore $\alpha = a = d$ is a rational integer. Item (i) of Lemma 8 follows.

(ii) Let $\alpha \notin \mathbb{R}$. We exclude α dividing second equation of (27) by the first:

$$\tau = \frac{c + d\tau}{a + b\tau}. \quad (28)$$

Therefore $b\tau^2 + (a - d)\tau - c = 0$. Note that $b \neq 0$ since τ cannot be a real number. Thus $\tau \in \mathbb{Q}(\sqrt{-d})$. Since $\alpha = a + b\tau$, $\alpha \in \mathbb{Q}(\sqrt{-d})$. Item (ii) of Lemma 8 follows. \square

Complex multiplication. If $\text{End } E$ is different from \mathbb{Z} , E is said to be *elliptic curve with complex multiplication*. If E admits complex multiplication, then its ring $\text{End } E$ is an order in an imaginary quadratic field K . In fact, E admits an isogeny (analytic homomorphism) to a curve E' such that $\text{End } E' \simeq \mathcal{O}_K$, where \mathcal{O}_K is the ring of integers of field K ([14]). Thus, by property $\alpha L \subseteq L$, lattice L is an ideal in \mathcal{O}_K . Denote by h_K the class number of field K . It is well known, that there exist h_K non-isomorphic ideals in \mathcal{O}_K . Therefore, elliptic curves $E_1 = \mathbb{C}/L_1, \dots, E_{h_K} = \mathbb{C}/L_{h_K}$ are pairwise non-isomorphic, but their endomorphism ring is the same ([14]).

References

- [1] A. Connes, Noncommutative Geometry, Academic Press, 1994.
- [2] E. G. Effros and C. L. Shen, Approximately finite C^* -algebras and continued fractions, Indiana Univ. Math. J. 29 (1980), 191-204.
- [3] H. Huber, Zur analytischen Theorie hyperbolischer Raumformen und Bewegungsgruppen, Math. Annalen 138 (1959), 1-26.
- [4] V. F. R. Jones, Subfactors and Knots, CBMS Series in Mathematics 80, Amer. Math. Soc. 1991.
- [5] F. Klein, Über eine geometrische Auffassung der gewöhnlichen Kettenbruchentwicklung, Nachrichten der Kgl. Gesellschaft der Wissenschaften zu Göttingen, Math.-phys. Klasse (1895), Heft 3.
- [6] F. Klein, Ausgewählte Kapitel der Zahlentheorie, Math. Annalen 48 (1896/97).
- [7] Yu. I. Manin, Real multiplication and noncommutative geometry, arXiv:math.AG/0202109 v1, Preprint, 46 p.
- [8] Yu. I. Manin and A. A. Panchishkin, Introduction to Number Theory, EMS, Number Theory I, Springer 1995.
- [9] H. P. McKean, Selberg's trace formula as applied to a compact Riemann surface, Comm. Pure Appl. Math. 25, (1972), 225-246.
- [10] H. Minkowski, Geometrie der Zahlen, Leipzig, 1910.
- [11] F. J. Murray and J. von Neumann, On rings of operators, Annals of Math. 37 (1936), 116-229.
- [12] M. Rørdam, F. Larsen and N. Laustsen, An introduction to K -theory for C^* -algebras, LMS Student Texts, 49, Cambridge University Press, 2000.
- [13] A. Selberg, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series, J. Indian Math. Soc. 20 (1956), 47-87.
- [14] J. H. Silverman, The Arithmetic of Elliptic Curves, Springer, 1986.
- [15] W. P. Thurston, The Geometry and Topology of Three-Manifolds, MSRI 1997, electronic edition of 1980 Princeton Univ. notes, available at <http://www.msri.org/gt3m/>; alternative reference: Three-Dimensional Geometry and Topology, ed. by Silvio Levy, vol.1, Princeton Univ. Press, 1997.
- [16] S. Wolpert, The length spectra as moduli for compact Riemann surfaces, Annals of Math. 109 (1979), 323-351.