# Morita Contexts for Corings and Equivalences<sup>\*</sup>

Jawad Y. Abuhlail<sup> $\dagger$ </sup>

Mathematics Department, Birzeit University Birzeit - Palestine

#### Abstract

In this note we study Morita contexts and Galois extensions for corings. For a coring C over a (not necessarily commutative) ground ring A we give equivalent conditions for  $\mathcal{M}^{\mathcal{C}}$  to satisfy the weak. resp. the strong structure theorem. We also characterize the so called *cleft C-Galois extensions* over commutative rings. Our approach is similar to that of Y. Doi and A. Masuoka in their work on (cleft) H-Galois extensions (e.g. [Doi94], [DM92]).

## Introduction

Let  $\mathcal{C}$  be a coring over a not necessarily commutative ring A and assume A to be a right  $\mathcal{C}$ -comodule through  $\varrho_A : A \longrightarrow A \otimes_A \mathcal{C} \simeq \mathcal{C}$ ,  $a \mapsto \mathbf{x}a$  for some group-like element  $\mathbf{x} \in \mathcal{C}$  (see [Brz02, Lemma 5.1]). In the first section we study from the viewpoint of Morita theory the relationship between A and its subring of coinvariants  $B := A^{co\mathcal{C}} := \{b \in A \mid \varrho(b) = b\mathbf{x}\}$ . We consider the A-ring  $*\mathcal{C} := \text{Hom}_{A-}(\mathcal{C}, A)$  and its left ideal  $Q := \{q \in *\mathcal{C} \mid \sum c_1 q(c_2) = q(c)\mathbf{x} \text{ for all } c \in \mathcal{C}\}$  and show that B and  $*\mathcal{C}$  are connected via a Morita context using  ${}_{B}A_{*\mathcal{C}}$  and  ${}_{*\mathcal{C}}Q_B$  as connecting bimodules. Our Morita context is in fact a generalization of Doi's Morita context presented in [Doi94].

In the second section we introduce the weak (resp. the strong) structure theorem for  $\mathcal{M}^{\mathcal{C}}$ . For the case  ${}_{A}\mathcal{C}$  is locally projective, in the sense of B. Zimmermann-Huignes, we characterize A being a generator (a progenerator) in the category of right  $\mathcal{C}$ -comodules by  $\mathcal{M}^{\mathcal{C}}$  satisfying the weak (resp. the strong) structure theorem. Here the notion of Galois corings introduced by T. Brzeziński [Brz02] plays an important role. The results and proofs are essentially module theoretic and similar to those of [MZ97] for the category  $\mathcal{M}(H)^{C}_{A}$  of Doi-Koppinen modules corresponding to a right-right Doi-Koppinen structure (H, A, C) (see also [MSTW01] for the case C = H).

The notion of a C-Galois extension A of a ring B was introduced by T. Brzeziński and S. Majid in [BM98] and is related to the so called entwining structures introduced in the

<sup>\*</sup>MSC (2000): 16D90, 16S40, 16W30, 13B02

Keywords: Morita Contexts, Hopf Algebras, (Cleft) Galois Extensions, Entwining Structures, Entwined Modules.

<sup>&</sup>lt;sup>†</sup>Current Address: Department of Mathematical Sciences, Box # 5046, KFUPM, 31261 Dhahran (Saudi Arabia), **email:** abuhlail@kfupm.edu.sa

same paper. In the third section we give equivalent conditions for a C-Galois extension A/B to be cleft. Our results generalize results of [Brz99] from the case of a base field to the case of a commutative ground ring. In the special case  $\rho(a) = \sum a_{\psi} \otimes x^{\psi}$ , for some group-like element  $x \in C$ , we get a complete generalization of [DM92, Theorem 1.5] (and [Doi94, Theorem 2.5]).

With A we denote a not necessarily commutative ring with  $1_A \neq 0_A$  and with  $\mathcal{M}_A$  (resp.  $_A\mathcal{M}, _A\mathcal{M}_A$ ) the category of *unital* right A-modules (resp. left A-modules, A-bimodules). For every right A-module W we denote by Gen( $W_A$ ) (resp.  $\sigma[W_A]$ ) the class of W-generated (resp. W-subgenerated) right A-modules. For the well developed theory of categories of type  $\sigma[W]$  the reader is referred to [Wis88, Section 15].

An A-module W is called **locally projective** (in the sense of B. Zimmermann-Huignes [Z-H76]), if for every diagram



with exact rows and F f.g.: for every A-linear map  $g: W \longrightarrow N$ , there exists an A-linear map  $g': W \longrightarrow L$ , such that the entstanding parallelogram is commutative. Note that every projective A-module is locally projective. By [Z-H76, Theorem 2.1] a left A-module W is locally projective, iff for every right A-module M the following map is injective

$$\alpha_M^W: M \otimes_A W \longrightarrow \operatorname{Hom}_{-A}(^*W, M), \ m \otimes_A w \mapsto [f \mapsto mf(w)].$$

It's easy then to see that every locally projective A-module is flat and A-cogenerated.

Let  $\mathcal{C}$  be an A-coring. We consider the canonical A-bimodule  $*\mathcal{C} := \operatorname{Hom}_{A-}(\mathcal{C}, A)$  as an A-ring with the canonical A-bimodule structure, multiplication  $(f \cdot g)(c) := \sum g(c_1 f(c_2))$ and unity  $\varepsilon_{\mathcal{C}}$ . If  $_A\mathcal{C}$  is locally projective, then we have an isomorphism of categories  $\mathcal{M}^{\mathcal{C}} \simeq \sigma[\mathcal{C}_{*\mathcal{C}}]$  (in particular  $\mathcal{M}^{\mathcal{C}} \subseteq \mathcal{M}_{*\mathcal{C}}$  is a full subcategory) and we have a left exact functor  $\operatorname{Rat}^{\mathcal{C}}(-) : \mathcal{M}_{*\mathcal{C}} \to \mathcal{M}^{\mathcal{C}}$  assigning to every right \* $\mathcal{C}$ -module its maximum  $\mathcal{C}$ -rational \* $\mathcal{C}$ -submodule, which turns to be a right  $\mathcal{C}$ -comodule. Moreover  $\mathcal{M}^{\mathcal{C}} = \mathcal{M}_{*\mathcal{C}}$  iff  $_A\mathcal{C}$  is f.g. and projective. For more investigation of the  $\mathcal{C}$ -rational \* $\mathcal{C}$ -modules see [Abu03].

After this paper was finished, it turned out that some results in this paper were discovered independently by S. Caenepeel, J. Vercruysse and S. Wang [CVW04].

### **1** Morita Contexts

In this section we fix the following: C is an A-coring with group-like element  $\mathbf{x}$  and A is a right C-comodule with structure map

$$\varrho_A: A \longrightarrow A \otimes_A \mathcal{C} \simeq \mathcal{C}, \ a \mapsto \mathbf{x} a$$

(e.g. [Brz02, Lemma 5.1]), hence  $A \in \mathcal{M}_{*\mathcal{C}}$  with  $a \leftarrow g := \sum a_{<0>}g(a_{<1>}) = g(\mathbf{x}a)$  for all  $a \in A$  and  $g \in {}^*\mathcal{C}$ . For  $M \in \mathcal{M}_{*\mathcal{C}}$  put

$$M^{\mathbf{x}} := \{ m \in M \mid mg = mg(\mathbf{x}) \text{ for all } g \in {}^{*}\mathcal{C} \}.$$

In particular  $A^{\mathbf{x}} := \{a \in A \mid a \leftarrow g = ag(\mathbf{x}) \text{ for all } g \in {}^*\mathcal{C}\} \subset A \text{ is a subring. For } M \in \mathcal{M}^{\mathcal{C}}$ we set

$$M^{co\mathcal{C}} := \{ m \in M \mid \varrho(m) = m \otimes_A \mathbf{x} \} \subseteq M^{\mathbf{x}}.$$

Obviously  $B := A^{co\mathcal{C}} = \{b \in A \mid b\mathbf{x} = \mathbf{x}b\} \subseteq A^{\mathbf{x}}$  is a subring and  $\varrho_A$  is (B, A)-bilinear. For  $M \in \mathcal{M}^{\mathcal{C}}$  we have  $M^{co\mathcal{C}} \in \mathcal{M}_B$ . Moreover we set

$$Q := \{ q \in {}^{*}\mathcal{C} \mid \sum c_1 q(c_2) = q(c)\mathbf{x} \text{ for all } c \in \mathcal{C} \} \subseteq ({}^{*}\mathcal{C})^{\mathbf{x}}.$$

**Lemma 1.1.** 1. For every right \*C-module M we have an isomorphism of right Bmodules

 $\omega_M : \operatorname{Hom}_{-*\mathcal{C}}(A, M) \longrightarrow M^{\mathbf{x}}, \ f \mapsto f(1_A)$ 

with inverse  $m \mapsto [a \mapsto ma]$ .

2. Let  ${}_{A}\mathcal{C}$  be locally projective. If  $M \in \mathcal{M}^{\mathcal{C}}$ , then  $M^{co\mathcal{C}} = M^{\mathbf{x}} \simeq \operatorname{Hom}_{-*\mathcal{C}}(A, M) = \operatorname{Hom}^{\mathcal{C}}(A, M)$ . Hence

 $\Psi_M: M^{co\mathcal{C}} \otimes_B A \longrightarrow M, \ m \otimes_B a \mapsto ma$ 

is surjective (resp. injective, bijective), iff

$$\Psi'_M$$
: Hom<sup>C</sup>(A, M)  $\otimes_B A \longrightarrow M, f \otimes_B a \mapsto f(a)$ 

is surjective (resp. injective, bijective).

- 3. We have  $\operatorname{Hom}_{*\mathcal{C}}(A, {}^{*}\mathcal{C}) \simeq ({}^{*}\mathcal{C})^{\mathbf{x}}$ . If moreover  ${}_{A}\mathcal{C}$  is A-cogenerated (resp. locally projective and  ${}^{\Box}\mathcal{C} := \operatorname{Rat}^{\mathcal{C}}({}^{*}\mathcal{C}_{*\mathcal{C}}))$ , then  $Q = ({}^{*}\mathcal{C})^{\mathbf{x}}$  (resp.  $Q = ({}^{\Box}\mathcal{C})^{co\mathcal{C}}$ ).
- 4. For every  $M \in \mathcal{M}_{*\mathcal{C}}$  (resp.  $M \in \mathcal{M}^{\mathcal{C}}$ ) and all  $m \in M$ ,  $q \in Q$  we have  $mq \in M^{\mathbf{x}}$  (resp.  $mq \in M^{co\mathcal{C}}$ ).

#### **Proof.** 1. Obvious.

2. Trivial.

3. Considering \* $\mathcal{C}$  as a right \* $\mathcal{C}$ -module via right multiplication we get Hom<sub>\* $\mathcal{C}$ </sub>( $A, \mathcal{C}$ )  $\simeq$  (\* $\mathcal{C}$ )<sup>**x**</sup> by (1). If  $q \in (\mathcal{C})^{$ **x** $}$ , then we have for all  $g \in \mathcal{C}$  and  $c \in \mathcal{C}$ :

$$g(\sum c_1 q(c_2)) = \sum g(c_1 q(c_2)) = (q \cdot g)(c) = (qg(\mathbf{x}))(c) = q(c)g(\mathbf{x}) = g(q(c)\mathbf{x}),$$

i.e.  $\sum c_1 q(c_2) - q(c)\mathbf{x} \in \operatorname{Re}(\mathcal{C}, A) := \bigcap \{\operatorname{Ke}(g) \mid g \in \operatorname{Hom}_{A-}(\mathcal{C}, A)\}$ . If  $_A\mathcal{C}$  is A-cogenerated, then  $\operatorname{Re}(\mathcal{C}, A) = 0$ , hence  $Q = ({}^*\mathcal{C})^{\mathbf{x}}$ .

Assume  $_{A}\mathcal{C}$  to be locally projective. Then we have for all  $q \in Q, g \in {}^{*}\mathcal{C}$  and  $c \in \mathcal{C}$ :

$$(q \cdot g)(c) = \sum g(c_1 q(c_2)) = g(q(c)\mathbf{x}) = q(c)g(\mathbf{x}) = (qg(\mathbf{x}))(c),$$

hence  $q \in {}^{\Box}\mathcal{C}$ , with  $\varrho(q) = q \otimes_A \mathbf{x}$ , i.e.  $q \in ({}^{\Box}\mathcal{C})^{co\mathcal{C}}$ . On the other hand, if  $q \in ({}^{\Box}\mathcal{C})^{co\mathcal{C}}$ , then for all  $g \in {}^*\mathcal{C}$  we have  $q \cdot g = qg(\mathbf{x})$ , i.e.  $q \in ({}^*\mathcal{C})^{\mathbf{x}} = Q$ .

4. Let  $M \in \mathcal{M}_{*\mathcal{C}}$ . Then we have for all  $q \in Q$ ,  $g \in {}^*\mathcal{C}$  and  $m \in M$ :

$$(mq)g = m(q \cdot g) = m(qg(\mathbf{x})) = (mq)g(\mathbf{x}),$$

i.e.  $mq \in M^{\mathbf{x}}$ . If  $M \in \mathcal{M}^{\mathcal{C}}$ , then we have for all  $m \in M$  and  $q \in Q$ :

i.e.  $mq \in M^{co\mathcal{C}}$ .

**Lemma 1.2.** 1. With the canonical actions A is a  $(B, {}^*C)$ -bimodule.

2. Q is a (\*C, B)-bimodule.

**Proof.** 1. By assumption  $A \in \mathcal{M}^{\mathcal{C}} \subseteq \mathcal{M}_{*\mathcal{C}}$ . For all  $b \in B$ ,  $a \in A$  and  $g \in {}^{*\mathcal{C}}$  we have

$$b(a \leftarrow g) = bg(\mathbf{x}a) = g(b(\mathbf{x}a)) = g(\mathbf{x}(ba)) = (ba) \leftarrow g.$$

2. For all  $a \in A$ ,  $q \in Q$  and  $c \in C$  we have

$$\sum c_1(aq)(c_2) = \sum c_1q(c_2a) = \sum (ca)_1q((ca)_2) = q(ca)\mathbf{x} = (aq)(c)\mathbf{x}.$$

For all  $q \in Q$ ,  $b \in B$  and  $c \in \mathcal{C}$  we have

$$\sum c_1(qb)(c_2) = \sum c_1q(c_2)b = q(c)\mathbf{x}b = q(c)b\mathbf{x} = (qb)(c)\mathbf{x}b$$

On the other hand we have for all  $q \in Q$ ,  $g \in {}^*\mathcal{C}$  and  $c \in \mathcal{C}$ :

$$\sum c_1(g \cdot q)(c_2) = \sum c_1q(c_{21}g(c_{22})) = \sum c_{11}q(c_{12}g(c_2))$$
  
= 
$$\sum c_{11}(g(c_2)q)(c_{12}) = \sum (g(c_2)q)(c_1)\mathbf{x}$$
  
= 
$$\sum q(c_1g(c_2))\mathbf{x} = (g \cdot q)(c)\mathbf{x}.$$

Moreover we have for all  $b \in B, q \in Q, g \in {}^*\mathcal{C}$  and  $c \in \mathcal{C}$ :

$$\begin{array}{rcl} ((g \cdot q)b)(c) &=& (g \cdot q)(c)b &=& \sum q(c_1g(c_2))b \\ &=& \sum (qb)(c_1g(c_2)) &=& (g \cdot qb)(c). \blacksquare \end{array}$$

**Theorem 1.3.** Keep the notation above fixed.

1.  $(A^{\mathbf{x}}, {}^{*}\mathcal{C}, A, ({}^{*}\mathcal{C})^{\mathbf{x}}, \widetilde{F}, \widetilde{G})$  is a Morita context derived form  $A_{{}^{*}\mathcal{C}}$ , where

2.  $(B, {}^{*}C, A, Q, F, G)$  is a Morita context, where

If moreover  $_{A}\mathcal{C}$  is locally projective, then the two Morita contexts coincide.

- **Proof.** 1. By Lemma 1.1 we have  $\operatorname{End}(A_{*\mathcal{C}}) \simeq A^{\mathbf{x}}$ ,  $(*\mathcal{C})^{\mathbf{x}} \simeq \operatorname{Hom}_{-*\mathcal{C}}(A, *\mathcal{C})$  and the result follows by [Fai81, Proposition 12.6].
  - 2. By Lemma 1.2 A is a  $(B, \mathcal{C})$ -bimodule and Q is a  $(\mathcal{C}, B)$ -bimodule. For all  $q \in Q, g \in \mathcal{C}$ ,  $a \in A$  and  $c \in \mathcal{C}$  we have

$$F(g \cdot q \otimes_B a)(c) = \sum q(c_2g(c_1))a = (g \cdot qa)(c) = (g \cdot F(q \otimes_B a))(c)$$

and

$$F(q \otimes_B a \leftarrow g)(c) = q(c)(a \leftarrow g) = q(c)g(\mathbf{x}a)$$
  
$$= g(q(c)\mathbf{x}a) = \sum_{i=1}^{n} g(c_1q(c_2)a)$$
  
$$= \sum_{i=1}^{n} g(c_1(qa)(c_2)) = (F(q \otimes_B a) \cdot g)(c),$$

hence F is \*C-bilinear. Note that by Lemma 1.1 G is well defined and is obviously B-bilinear. Moreover we have for all  $a, \tilde{a} \in A$  and  $q, \tilde{q} \in Q$  the following associativity relations:

$$(F(q \otimes_B a) \cdot \widetilde{q})(c) = \sum \widetilde{q}(c_1q(c_2)a) = \widetilde{q}(q(c)\mathbf{x}a)$$
  
$$= q(c)\widetilde{q}(\mathbf{x}a) = (qG(a \otimes_{*\mathcal{C}} \widetilde{q}))(c),$$
  
$$G(a \otimes_{*\mathcal{C}} q)\widetilde{a} = q(\mathbf{x}a)\widetilde{a} = (q\widetilde{a})(\mathbf{x}a)$$
  
$$= F(q \otimes_B \widetilde{a})(\mathbf{x}a) = a \leftarrow F(q \otimes_B \widetilde{a}).$$

If  ${}_{A}\mathcal{C}$  is locally projective, then  $A^{\mathbf{x}} = A^{co\mathcal{C}}$ ,  $({}^{*}\mathcal{C})^{\mathbf{x}} = Q$  by Lemma 1.1 and the two contexts coincide.

**1.4.** [Brz02, Definition 5.3] An A-coring C is said to be **Galois**, if there exists an A-coring isomorphism  $\chi : A \otimes_B A \longrightarrow C$  such that  $\chi(1_A \otimes_B 1_A) = \mathbf{x}$ . Recall that  $A \otimes_B A$  is an A-coring with the canonical A-bimodule structure, comultiplication

$$\Delta: A \otimes_B A \longrightarrow (A \otimes_B A) \otimes_A (A \otimes_B A), \ \widetilde{a} \otimes_B a \mapsto (\widetilde{a} \otimes_B 1_A) \otimes_A (1_A \otimes_B a)$$

and counity  $\varepsilon_{A\otimes_B A} : A \otimes_B A \longrightarrow A$ ,  $\widetilde{a} \otimes_B a \mapsto \widetilde{a}a$ .

**1.5.** Consider the functors

$$(-)^{co\mathcal{C}}: \mathcal{M}^{\mathcal{C}} \longrightarrow \mathcal{M}_B \text{ and } - \otimes_B A: \mathcal{M}_B \longrightarrow \mathcal{M}^{\mathcal{C}}.$$

By [Brz02, Proposition 5.2]  $(-\otimes_B A, (-)^{co\mathcal{C}})$  is an adjoint pair of covariant functors, where the adjunctions are given by

$$\Phi_N: N \longrightarrow (N \otimes_B A)^{co\mathcal{C}}, \ n \mapsto n \otimes_B 1_A \tag{1}$$

and

$$\Psi_M: M^{co\mathcal{C}} \otimes_B A \longrightarrow M, \ m \otimes_B a \mapsto ma.$$
<sup>(2)</sup>

If  $\Psi_M$  is an isomorphism for all  $M \in \mathcal{M}^{\mathcal{C}}$ , then we say  $\mathcal{M}^{\mathcal{C}}$  satisfies the **weak structure theorem.** If in addition  $\Phi_N$  is an isomorphism for all  $N \in \mathcal{M}_B$ , then we say  $\mathcal{M}^{\mathcal{C}}$  satisfies the **strong structure theorem** (in this case  $(-)^{co\mathcal{C}}$  and  $-\otimes_B A$  give an equivalence of categories  $\mathcal{M}^{\mathcal{C}} \simeq \mathcal{M}_B$ ).

**1.6.** Let  $W \in \mathcal{M}_A$  and consider the canonical right  $\mathcal{C}$ -comodule  $W \otimes_A \mathcal{C}$ . Then  $W \simeq (W \otimes_A \mathcal{C})^{co\mathcal{C}}$  via  $w \mapsto w \otimes_A \mathbf{x}$  with inverse  $w \otimes_A c \mapsto w \varepsilon_{\mathcal{C}}(c)$  and we define

$$\beta_W := \Psi_{W \otimes_A \mathcal{C}} : W \otimes_B A \longrightarrow W \otimes_A \mathcal{C}, \ w \otimes_B a \mapsto w \otimes_A \mathbf{x}a.$$
(3)

In particular we have for W = A the morphism of A-corings

$$\beta := \Psi_{A \otimes_A \mathcal{C}} : A \otimes_B A \longrightarrow A \otimes_A \mathcal{C} \simeq \mathcal{C}, \ \widetilde{a} \otimes_B a \mapsto \widetilde{a} \mathbf{x} a.$$

$$\tag{4}$$

If  $\beta$  is bijective, then C is a Galois A-coring and we call the ring extension A/B C-Galois.

**Theorem 1.7.** For the Morita context (B, C, A, Q, F, G) the following statements are equivalent:

- 1.  $G: A \otimes_{*\mathcal{C}} Q \longrightarrow B$  is surjective (bijective and  $B = A^{\mathbf{x}}$ );
- 2. there exists  $\widehat{q} \in Q$ , such that  $\widehat{q}(\mathbf{x}) = 1_A$ ;
- 3. for every right \*C-module M we have a B-module isomorphism  $M \otimes_{*C} Q \simeq M^{\mathbf{x}}$ .
- 4. for every right C-comodule M we have  $M \otimes_{*C} Q \simeq M^{\mathbf{co}C}$  as B-modules. If moreover  $_{A}C$  is locally projective, then (1)-(4) are moreover equivalent to:
- 5.  $A_{*C}$  is (f.g.) projective.

**Proof.** (1)  $\Rightarrow$  (2). Assume G to be surjective. Then there exist  $a_1, ..., a_k$  and  $q_1, ..., q_k \in Q$ , such that  $G(\sum_{i=1}^k a_i \otimes_{*\mathcal{C}} q_i) = 1_A$ . Set  $\widehat{q} := \sum_{i=1}^k a_i q_i \in Q$ . Then we have

$$\widehat{q}(\mathbf{x}) = \left(\sum_{i=1}^{k} a_i q_i\right)(\mathbf{x}) = \sum_{i=1}^{k} q_i(\mathbf{x}a_i) = \sum_{i=1}^{k} (a_i \leftarrow q_i) = G\left(\sum_{i=1}^{k} a_i \otimes_{*\mathcal{C}} q_i\right) = 1_A.$$

 $(2) \Rightarrow (3)$ . Consider the *B*-module morphism

$$\xi_M: M \otimes_{{}^*\mathcal{C}} Q \longrightarrow M^{\mathbf{x}}, \ m \otimes_{{}^*\mathcal{C}} q \mapsto mq.$$

Let  $\widehat{q} \in Q$  with  $\widehat{q}(\mathbf{x}) = 1_A$  and define  $\widetilde{\xi}_M : M^{\mathbf{x}} \longrightarrow M \otimes_{*\mathcal{C}} Q, m \mapsto m \otimes_{*\mathcal{C}} \widehat{q}$ . For every  $n \in M^{\mathbf{x}}$  we have

$$(\xi_M \circ \widetilde{\xi}_M)(n) = \xi_M(n \otimes_{*\mathcal{C}} \widehat{q}) = n \leftharpoonup \widehat{q} = n\widehat{q}(\mathbf{x}) = n.$$

On the other hand we have for all  $m \in M$  and  $q \in Q$ :

i.e.  $\xi_M$  is bijective with inverse  $\tilde{\xi}_M$ .

(3)  $\Rightarrow$  (4). Let  $M \in \mathcal{M}^{\mathcal{C}}$ . By Lemma 1.1 we have  $\xi_M(M \otimes_{*\mathcal{C}} Q) \subseteq M^{co\mathcal{C}} \subseteq M^{\mathbf{x}}$ . By assumption  $\xi_M : A \otimes_{*\mathcal{C}} Q \longrightarrow M^{\mathbf{x}}$  is bijective. Hence  $M^{\mathbf{x}} = M^{co\mathcal{C}}$  and  $M \otimes_{*\mathcal{C}} Q \stackrel{\xi_M}{\simeq} M^{co\mathcal{C}}$ . (4)  $\Rightarrow$  (1) We are done since  $G = \xi_A$ .

Assume  ${}_{A}\mathcal{C}$  to be locally projective.

Then  $B \simeq \operatorname{End}(A_{*\mathcal{C}}), Q \simeq \operatorname{Hom}_{-*\mathcal{C}}(A, ^*\mathcal{C})$  and we get (1)  $\iff$  (5) by [Fai81, Corollary 12.8].

**Corollary 1.8.** For the Morita context  $(B, {}^*\mathcal{C}, A, Q, F, G)$  assume there exists  $\widehat{q} \in Q$  with  $\widehat{q}(\mathbf{x}) = 1_A$  (equivalently  $G : Q \otimes_B A \longrightarrow {}^*\mathcal{C}$  is surjective). Then:

- 1. For every  $N \in \mathcal{M}_B$ ,  $\Phi_N$  is an isomorphism.
- 2. B is a left B-direct summand of A.
- **Proof.** 1. Let  $N \in \mathcal{M}_B$ . Then we have by Theorem 1.7 the isomorphisms  $G : A \otimes_{*\mathcal{C}} Q \longrightarrow B$  and  $\xi_{N \otimes_B A} : (N \otimes_B A) \otimes_{*\mathcal{C}} Q \longrightarrow (N \otimes_B A)^{co\mathcal{C}}$ . Moreover  $\Phi_N$  is given by the canonical isomorphisms

$$N \simeq N \otimes_B B \simeq N \otimes_B (A \otimes_{*\mathcal{C}} Q) \simeq (N \otimes_B A) \otimes_{*\mathcal{C}} Q \simeq (N \otimes_B A)^{co\mathcal{C}}.$$

2. The map  $\operatorname{tr}_A : A \longrightarrow B$ ,  $a \mapsto a \leftarrow \widehat{q}$  is left B-linear with  $\operatorname{tr}_A(b) = b$  for all  $b \in B$ .

**Corollary 1.9.** For the Morita context  $(B, {}^*\mathcal{C}, A, Q, F, G)$  assume there exists  $\widehat{q} \in Q$  with  $\widehat{q}(\mathbf{x}) = 1_A$  (equivalently  $G : Q \otimes_B A \longrightarrow B$  is surjective). Then:

- 1.  $_{B}A$  and  $Q_{B}$  are generators.
- 2.  $A_{*C}$  and  ${}_{*C}Q$  are f.g. and projective.
- 3.  $F: Q \otimes_B A \longrightarrow {}^*\mathcal{C}$  induces bimodule isomorphisms

 $A \simeq \operatorname{Hom}_{*\mathcal{C}}(Q, {}^*\mathcal{C}) \text{ and } Q \simeq \operatorname{Hom}_{-*\mathcal{C}}(A, {}^*\mathcal{C}).$ 

4. The bimodule structures above induce ring isomorphisms

$$B \simeq \operatorname{End}(A_{*\mathcal{C}})$$
 and  $B \simeq \operatorname{End}({_*\mathcal{C}Q})^{op}$ 

**Proof.** The result follows by standard argument of Morita Theory (e.g. [Fai81, Proposition 12.7]).■

**Proposition 1.10.** Consider the Morita context  $(B, {}^*\mathcal{C}, A, Q, F, G)$  and assume that  $F : Q \otimes_{{}^*\mathcal{C}} A \longrightarrow {}^*\mathcal{C}$  is surjective. Then:

- 1.  $A_{*\mathcal{C}}$  is a generator,  $Q \simeq \operatorname{Hom}_{B-}(A, B)$  as bimodules and  $*\mathcal{C} \simeq \operatorname{End}(Q_B)$ .
- 2.  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem (in particular A/B is  $\mathcal{C}$ -Galois).
- **Proof.** 1. The result follows by standard argument of Morita Theory (e.g. [Fai81, Proposition 12.7]).
  - 2. By assumption  $\varepsilon_{\mathcal{C}} = F(\sum_{i=1}^{k} q_i \otimes_B a_i)$  for some  $\{(q_i, a_i)\}_{i=1}^k \subseteq Q \times A$ . In this case  $\Psi_M : M^{co\mathcal{C}} \otimes_B A \longrightarrow M$  is bijective with inverse  $\widetilde{\Psi}_M : M \longrightarrow M^{co\mathcal{C}} \otimes_B A, m \mapsto \sum_{i=1}^k mq_i \otimes_B a_i$ . In fact, we have for all  $m \in M, n \in M^{co\mathcal{C}}$  and  $a \in A$ :

$$(\Psi_M \circ \widetilde{\Psi}_M)(m) = \sum_{\substack{i=1\\k}}^k (mq_i)a_i = \sum_{\substack{i=1\\k}}^k (m_{<0>}q_i(m_{<1>})a_i)$$
  
= 
$$\sum_{\substack{i=1\\k}}^k m_{<0>}(q_ia_i)(m_{<1>}) = \sum_{\substack{i=1\\k}}^k m_{<0>}\varepsilon_{\mathcal{C}}(m_{<1>})$$
  
= 
$$m$$

and

$$(\widetilde{\Psi}_{M} \circ \Psi_{M})(n \otimes_{B} a) = \sum_{i=1}^{k} (na)q_{i} \otimes_{B} a_{i} = \sum_{i=1}^{k} nq_{i}(\mathbf{x}a) \otimes_{B} a_{i}$$
$$= \sum_{i=1}^{k} n \otimes_{B} q_{i}(\mathbf{x}a)a_{i} = \sum_{i=1}^{k} n \otimes_{B} (q_{i}a_{i})(\mathbf{x}a)$$
$$= n \otimes_{B} \varepsilon_{\mathcal{C}}(\mathbf{x}a) = n \otimes_{B} a.\blacksquare$$

**Theorem 1.11.** For the Morita context  $(B, \mathcal{C}, A, Q, F, G)$  the following are equivalent:

- 1.  $F: Q \otimes_B A \longrightarrow {}^*C$  is surjective (bijective);
- 2. (a) Q<sub>B</sub> is f.g. and projective;
  (b) Ω : A → Hom<sub>-B</sub>(Q, B), a → [q → a ← q] is a bimodule isomorphism;
  (c) \*<sub>C</sub>Q is faithful.
  If <sub>A</sub>C is A-cogenerated, then (1) & (2) are moreover equivalent to:
- 3. (a) <sub>B</sub>A is f.g. and projective;
  (b) Λ : \*C → End(<sub>B</sub>A)<sup>op</sup>, g → [a → a ← g] is a ring isomorphism.
- 4.  $A_{*C}$  is a generator. If moreover  $_{A}C$  is f.g. and projective, then (1)-(4) are equivalent to:
- 5.  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem.

**Proof.** The implications  $(1) \Rightarrow (2)$ , (3), (4) follow without any finiteness conditions on  $\mathcal{C}$  by standard argument of Morita Theory (e.g. [Fai81, Proposition 12.7]). Note that  ${}_{*\mathcal{C}}Q$  is faithful by the embedding  ${}^{*\mathcal{C}} \hookrightarrow \operatorname{End}(Q_B)$  (see Proposition 1.10 (1)).

 $(2) \Rightarrow (1).$  Let  $\{(q_i, p_i)\}_{i=1}^k \subset Q \times \operatorname{Hom}_{-B}(Q, B)$  be a dual basis for  $Q_B$ . By (b) there exist  $a_1, ..., a_k \in A$ , such that  $\Omega(a_i) = q_i$  for i = 1, ..., k. For every  $q \in Q$  we have then  $(\sum_{i=1}^k q_i a_i) \cdot q = \sum_{i=1}^k q_i (a_i \leftarrow q) = \sum_{i=1}^k q_i p_i(q) = q$ , hence  $\sum_{i=1}^k q_i a_i = \varepsilon_{\mathcal{C}}$  by (c) and the \* $\mathcal{C}$ -bilinear morphism  $F: Q \otimes_{*\mathcal{C}} A \longrightarrow {}^*\mathcal{C}$  is surjective.

Assume  ${}_{A}\mathcal{C}$  to be A-cogenerated.

(3)  $\Rightarrow$  (1). Let  $\{(a_i, p_i)\}_{i=1}^k \subset A \times \operatorname{Hom}_{B-}(A, B)$  be a dual basis of  ${}_BA$ . By (b), there exist  $g_1, \ldots, g_k \in {}^*\mathcal{C}$ , such that  $\Lambda(g_i) = p_i$  for  $i = 1, \ldots, k$ . Claim:  $g_1, \ldots, g_k \in Q$ . For all  $f \in {}^*\mathcal{C}$  and  $i = 1, \ldots, k$  we have

$$\Lambda(g_i \cdot f)(a) = a \leftarrow (g_i \cdot f) = (a \leftarrow g_i) \leftarrow f)$$
  
$$= p_i(a) \leftarrow f = f(\mathbf{x}p_i(a))$$
  
$$= f(p_i(a)\mathbf{x}) = p_i(a)f(\mathbf{x})$$
  
$$= (p_if(\mathbf{x}))(a) = \Lambda(g_if(\mathbf{x}))(a),$$

hence  $g_i \cdot f = g_i f(\mathbf{x})$ , i.e.  $g_i \in ({}^*\mathcal{C})^{\mathbf{x}} = Q$  (by Lemma 1.1 (2)). Moreover for every  $a \in A$  we have:  $\Lambda(\sum_{i=1}^k g_i a_i)(a) = \sum_{i=1}^k a \leftarrow g_i a_i = \sum_{i=1}^k p_i(a)a_i = a$ , i.e.  $\sum_{i=1}^k g_i a_i = \varepsilon_{\mathcal{C}}$  and the  ${}^*\mathcal{C}$ -bilinear morphism F is surjective.

 $(4) \Rightarrow (1).$  Since  $Q \simeq \operatorname{Hom}_{*\mathcal{C}}(A, {}^*\mathcal{C})$ , we have  $\operatorname{Im}(F) = \operatorname{tr}(A, {}^*\mathcal{C}) := \sum \{\operatorname{Im}(h) : h \in \operatorname{Hom}_{*\mathcal{C}}(A, {}^*\mathcal{C})\}$ , hence  $\operatorname{Im}(F) = {}^*\mathcal{C}$  iff  $A_{*\mathcal{C}}$  is a generator (e.g. [Wis88, Page 154]).

Assume  ${}_{A}\mathcal{C}$  to be f.g. and projective.

 $(1) \Rightarrow (5)$  follows without any finiteness conditions on  $\mathcal{C}$  by Proposition 1.10 (2).

(5)  $\Rightarrow$  (1). Since  ${}_{A}\mathcal{C}$  is f.g. and projective, we have  $\mathcal{M}^{\mathcal{C}} \simeq \mathcal{M}_{*\mathcal{C}}$  (e.g. [Brz02, Lemma 4.3]), hence  ${}^{*}\mathcal{C} \in \mathcal{M}^{\mathcal{C}}$ ,  $Q = ({}^{*}\mathcal{C})^{co\mathcal{C}}$  and  $F = \Psi_{*\mathcal{C}}$ .

### 2 Galois Extensions and Equivalences

The notation of the first section remains fixed. For every  $M \in \mathcal{M}^{\mathcal{C}}$  we have the  $\mathcal{C}$ -colinear morphism

$$\Psi'_M : \operatorname{Hom}^{\mathcal{C}}(A, M) \otimes_B A \longrightarrow M, \ f \otimes_B a \mapsto f(a).$$

In this section we characterize A being a generator (resp. a progenerator) in  $\mathcal{M}^{\mathcal{C}}$  under the assumption that  $_{A}\mathcal{C}$  is locally projective. Our approach is similar to that of [MZ97] and our results generalize those obtained there for the special case of the category of Doi-Koppinen modules  $\mathcal{M}(H)_{A}^{C}$ .

**Lemma 2.1.** Assume  $_{A}C$  to be locally projective. If  $_{B}A$  is flat and A/B is C-Galois, then:

- 1. A is a subgenerator in  $\mathcal{M}^{\mathcal{C}}$ , i.e.  $\sigma[A_{*\mathcal{C}}] = \sigma[\mathcal{C}_{*\mathcal{C}}]$ .
- 2. for each  $M \in \mathcal{M}^{\mathcal{C}}, \Psi'_{M}$  is injective.
- 3. for every A-generated  $M \in \mathcal{M}^{\mathcal{C}}, \Psi'_{M}$  is an isomorphism.

**Proof.** Assume  ${}_{A}\mathcal{C}$  to be locally projective.

- 1. Since A/B is  $\mathcal{C}$ -Galois,  $\beta' := \Psi'_{\mathcal{C}}$  is an isomorphism, hence  $\mathcal{C}$  is A-generated. Consequently  $\sigma[A_{*\mathcal{C}}] \subseteq \sigma[\mathcal{C}_{*\mathcal{C}}] \subseteq \sigma[A_{*\mathcal{C}}]$ , i.e.  $\sigma[A_{*\mathcal{C}}] = \sigma[\mathcal{C}_{*\mathcal{C}}]$ .
- 2. With slight modifications, the proof of [MZ97, Lemma 3.22] applies.
- 3. If  $M \in \mathcal{M}^{\mathcal{C}}$  is A-generated, then  $\Psi'_M$  is surjective, hence bijective by (2).

The following result is a generalization of [Brz99, Proposition 3.13] (which in turn generalizes [DT89, Theorem 2.11]):

**Proposition 2.2.** Assume A/B to be C-Galois.

- 1. If  ${}_{B}A$  is flat, then  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem.
- 2. Assume there exists  $\hat{q} \in Q$ , such that  $\hat{q}(\mathbf{x}) = 1_A$ . If  ${}_BA$  is flat, or for all  $b \in B$  and  $c \in \mathcal{C}$  we have  $\hat{q}(cb) = q(c)b$ , then  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong structure theorem.

**Proof.** 1. The proof is the first part of the proof of [Brz02, Theorem 5.6].

2. By assumption and Corollary 1.8,  $\Phi_N$  is an isomorphism for all  $N \in \mathcal{M}_B$ . If  ${}_BA$  is flat, then  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem by (1). On the other hand, if for all  $b \in B$  and  $c \in \mathcal{C}$  we have  $\widehat{q}(cb) = q(c)b$ , then an analog argument to that in the proof of [Brz99, Proposition 3.13] shows that  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem.

**Theorem 2.3.** Assume  $_{A}C$  to be locally projective. Then the following are equivalent:

- 1.  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem;
- 2.  $_{B}A$  is flat and A/B is C-Galois;
- 3. <sub>B</sub>A is flat and  $\beta' := \Psi'_{\mathcal{C}}$  is an isomorphism;
- 4. <sub>B</sub>A is flat and for every A-generated  $M \in \mathcal{M}^{\mathcal{C}}, \Psi'_{M}$  is bijective;
- 5. for every  $M \in \mathcal{M}^{\mathcal{C}} = \sigma[\mathcal{C}_{*\mathcal{C}}]$ , the  $\mathcal{C}$ -colinear morphism  $\Psi'_{M}$  is bijective;
- $6. \ \sigma[\mathcal{C}_{*\mathcal{C}}] = \operatorname{Gen}(A_{*\mathcal{C}});$
- 7.  $_{B}A$  is flat,  $\sigma[\mathcal{C}_{*\mathcal{C}}] = \sigma[A_{*\mathcal{C}}]$  and  $\operatorname{Hom}_{-*\mathcal{C}}(A, -) : \operatorname{Gen}(A_{*\mathcal{C}}) \longrightarrow \mathcal{M}_{B}$  is full faithful;
- 8.  $\operatorname{Hom}^{\mathcal{C}}(A, -) : \mathcal{M}^{\mathcal{C}} \longrightarrow \mathcal{M}_B$  is faithful;
- 9. A is a generator in  $\mathcal{M}^{\mathcal{C}}$ .

**Proof.** (1)  $\iff$  (5) & (2)  $\iff$  (3) follow by Lemma 1.1. The equivalences (4)  $\iff$  (5)  $\iff$  (6)  $\iff$  (7) follow by [MZ97, Theorem 2.3]. The equivalence (8)  $\iff$  (9) is evident for any category, and moreover (6)  $\iff$  (9) by the fact that  $\text{Gen}(A_{*\mathcal{C}}) \subseteq \sigma[A_{*\mathcal{C}}] \subseteq \sigma[\mathcal{C}_{*\mathcal{C}}] = \mathcal{M}^{\mathcal{C}}$ . By Lemma 2.1 we have (3)  $\Rightarrow$  (4). Now assuming (1) we conclude that A/B is  $\mathcal{C}$ -Galois and that  $_BA$  is flat (since (1)  $\iff$  (5)  $\iff$  (7)), hence (1)  $\Rightarrow$  (2) follows and we are done.

**Definition 2.4.** ([MZ97, Definition 2.4]) A left module P over a ring S is called a **weak** generator, if for any right S-module  $Y, Y \otimes_S P = 0$  implies Y = 0. A right module P over a ring  $\mathcal{R}$  is called **quasiprogenerator** (resp. **progenerator**), if  $P_{\mathcal{R}}$  is f.g. quasiprojective and generates each of its submodules (resp.  $P_{\mathcal{R}}$  is f.g., projective and a generator).  $P_{\mathcal{R}}$ is called **faithful** (resp. **balanced**), if the canonical morphism  $\mathcal{R} \longrightarrow \operatorname{End}_{(\operatorname{End}(P_{\mathcal{R}})}P)^{op}$  is injective (resp. surjective).

**Theorem 2.5.** Assume  $_{A}C$  to be flat. Then the following are equivalent:

- 1.  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong structure theorem;
- 2. <sub>B</sub>A is faithfully flat and A/B is C-Galois.
  If moreover <sub>A</sub>C is locally projective, then (1) & (2) are moreover equivalent to:
- 3. <sub>B</sub>A is faithfully flat and  $\beta' := \Psi'_{\mathcal{C}}$  is bijective;
- 4. <sub>B</sub>A is faithfully flat and for every  $M \in \sigma[A_{*C}], \Psi'_{M}$  is bijective;
- 5.  $A_{*C}$  is quasiprojective and generates each of its submodules,  ${}_{B}A$  is a weak generator and  $\sigma[\mathcal{C}_{*C}] = \sigma[A_{*C}];$
- 6.  $A_{*\mathcal{C}}$  is a quasiprogenerator and  $\sigma[\mathcal{C}_{*\mathcal{C}}] = \sigma[A_{*\mathcal{C}}];$
- 7. <sub>B</sub>A is a weak generator,  $\Psi'_M$  is an isomorphism for every  $M \in \text{Gen}(A_{*\mathcal{C}})$  and  $\sigma[A_{*\mathcal{C}}] = \sigma[\mathcal{C}_{*\mathcal{C}}];$
- 8. Hom<sup> $\mathcal{C}$ </sup>(A, -) :  $\mathcal{M}^{\mathcal{C}} \longrightarrow \mathcal{M}_B$  is an equivalence;
- 9. A is a progenerator in  $\mathcal{M}^{\mathcal{C}}$ .

**Proof.** (1)  $\iff$  (2) is [Brz02, Theorem 5.6]. Assume  ${}_{A}C$  to be locally projective. Then (2)  $\iff$  (3) follows by Lemma 1.1 and we get (1)  $\iff$  (8)  $\iff$  (9) by characterizations of progenerators in categories of type  $\sigma[M]$  (see [Wis88, 18.5, 46.2]). Moreover (4)  $\iff$ (5)  $\iff$  (6)  $\iff$  (7) follow from [MZ97, Theorem 2.5]. Obviously (3)  $\Rightarrow$  (4) (note that (3)  $\iff$  (2)  $\iff$  (1)). Assume now (4). Then  ${}_{B}A$  is faithfully flat and moreover  $\Psi'_{C}$  is bijective, since  $C \in \sigma[A_{*C}]$  by (6), i.e. (4)  $\Rightarrow$  (3) and the proof is complete.

*Remark* 2.6. Assume  ${}_{A}\mathcal{C}$  to be locally projective. Then  $\text{Im}(F) \subseteq {}^{\Box}\mathcal{C}$ . In fact we have for all  $q \in Q$ ,  $a \in A$ ,  $g \in {}^{*}\mathcal{C}$  and  $c \in C$ :

$$((qa) \cdot g)(c) = \sum g(c_1q(c_2)a) = g(q(c)\mathbf{x}a) = q(c)g(\mathbf{x}a) = (qg(\mathbf{x}a)(c),$$

hence  $qa \in \Box C$ , with  $\varrho(qa) = q \otimes_A \mathbf{x}a$ .

**Proposition 2.7.** Assume  ${}_{A}\mathcal{C}$  to be locally projective and that there exists  $\widehat{q} \in Q$  with  $\widehat{q}(\mathbf{x}) = 1_A$  (equivalently  $G : A \otimes_{*\mathcal{C}} Q \longrightarrow B$  is surjective). Then  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong equivalence theorem, iff  $\operatorname{Im}(F) = {}^{\Box}\mathcal{C}$  and the following map is surjective for every  $M \in \mathcal{M}^{\mathcal{C}}$ 

$$\varpi_M: M \otimes_{^*\mathcal{C}} \ ^{\square}\mathcal{C} \longrightarrow M, \ m \otimes_{^*\mathcal{C}} f \mapsto mf.$$

In this case  $Q \otimes_B A \stackrel{F}{\simeq} \Box \mathcal{C}$  and  $M \otimes_{*\mathcal{C}} \Box \mathcal{C} \stackrel{\varpi_M}{\simeq} M$  for every  $M \in \mathcal{M}^{\mathcal{C}}$ .

**Proof.** Consider for every  $M \in \mathcal{M}^{\mathcal{C}}$  the commutative diagram

Assume  $\operatorname{Im}(F) = {}^{\Box}\mathcal{C}$  and  $\varpi_M$  to be surjective for every  $M \in \mathcal{M}^{\mathcal{C}}$ . Then  $\Psi_M$  is obviously surjective. Let  $K = \operatorname{Ke}(\Psi_M)$ . Since  $\Psi_M$  is a morphism in  $\mathcal{M}^{\mathcal{C}} \simeq \sigma[\mathcal{C}_{*\mathcal{C}}]$  we have  $K \in \mathcal{M}^{\mathcal{C}}$ , hence  $\Psi_K : K^{co\mathcal{C}} \otimes_B A \longrightarrow K$  is surjective. By Theorem 1.7 we have  $K \otimes_{*\mathcal{C}} Q \cong K^{co\mathcal{C}}$  and  $A \otimes_{*\mathcal{C}} Q \cong B$ , hence

$$K^{co\mathcal{C}} \simeq K \otimes_{*\mathcal{C}} Q = \operatorname{Ke}(\Psi_M) \otimes_{*\mathcal{C}} Q = \operatorname{Ke}(\Psi_M \otimes_{*\mathcal{C}} id_Q) = \operatorname{Ke}(id_{M^{co\mathcal{C}}}) = 0,$$

i.e.  $\Psi_M$  is bijective. By corollary 1.8  $\Phi_N$  is bijective for every  $N \in \mathcal{M}_B$ . Consequently  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong structure theorem.

On the other hand, assume that  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong structure theorem. Note that F is the adjunction of  $\Psi_{\Box_{\mathcal{C}}}$ , hence  $Q \otimes_B A \cong^F \Box_{\mathcal{C}}$  and consequently  $\varpi_M$  is also bijective for every  $M \in \mathcal{M}^{\mathcal{C}}$  by the commutativity of the above diagram.

*Remarks* 2.8. Assume  ${}_{A}\mathcal{C}$  to be locally projective.

- 1.  $\varpi_A : A \otimes_{*\mathcal{C}} \Box \mathcal{C} \longrightarrow A$  is surjective iff there exists  $\widehat{g} \in \Box \mathcal{C}$  with  $\widehat{g}(\mathbf{x}) = 1_A$ . To prove this assume first that  $\varpi_A$  is surjective. Then there exist  $\{(a_i, g_i)\}_{i=1}^k \subset A \times \Box \mathcal{C}$ , such that  $\sum_{i=1}^k a_i \leftarrow g_i = 1_A$ . Set  $\widehat{g} := \sum_{i=1}^k a_i g_i \in \Box \mathcal{C}$ . Then  $\widehat{g}(\mathbf{x}) = (\sum_{i=1}^k a_i g_i)(\mathbf{x}) =$  $\sum_{i=1}^k g_i(\mathbf{x}a_i) = \sum_{i=1}^k a_i \leftarrow g_i = 1_A$ . On the other hand, assume there exists  $\widehat{g} \in \Box \mathcal{C}$  with  $\widehat{g}(\mathbf{x}) = 1_A$ . Then for every  $a \in A$  we have  $1_A \leftarrow (\widehat{g}a) = (\widehat{g}a)(\mathbf{x}) = \widehat{g}(\mathbf{x})a = a$ , i.e.  $\varpi_A$ is surjective.
- 2. Assume  $\varpi_A$  to be surjective. If  $\Psi_M$  is surjective for  $M \in \mathcal{M}^{\mathcal{C}}$ , then  $\varpi_M$  is surjective, since

$$\varpi_M \circ (\Psi_M \otimes_{*\mathcal{C}} id_{\Box_{\mathcal{C}}}) = \Psi_M \circ (id_{M^{\operatorname{coc}}} \otimes_B \varpi_A).$$

### **Theorem 2.9.** Assume $_{A}C$ to be f.g. and projective. Then the following are equivalent:

- 1.  $\mathcal{M}^{\mathcal{C}}$  satisfies the weak structure theorem;
- 2.  $_{B}A$  is flat and A/B is C-Galois;
- 3.  $_{B}A$  is flat and  $\beta' := \Psi'_{\mathcal{C}}$  is an isomorphism;
- 4. <sub>B</sub>A is flat and for every A-generated  $M \in \mathcal{M}^{\mathcal{C}} = \mathcal{M}_{*\mathcal{C}}$ , the C-colinear morphism  $\Psi'_{M}$  is bijective;

- 5. for every  $M \in \mathcal{M}^{\mathcal{C}}$ , the  $\mathcal{C}$ -colinear morphism  $\Psi'_M$  is bijective;
- 6. <sub>B</sub>A is flat,  $\mathcal{M}_{*\mathcal{C}} = \sigma[A_{*\mathcal{C}}]$  and  $\operatorname{Hom}_{-*\mathcal{C}}(A, -) : \operatorname{Gen}(A_{*\mathcal{C}}) \longrightarrow \mathcal{M}_B$  is full faithful;
- 7. Hom<sub>-\* $\mathcal{C}$ </sub>(A, -) :  $\mathcal{M}_{*\mathcal{C}} \longrightarrow \mathcal{M}_B$  is faithful;
- 8.  $A_{*C}$  is a generator;
- 9.  $F: Q \otimes_B A \longrightarrow {}^*\mathcal{C}$  is surjective (bijective);
- 10. (a) Q<sub>B</sub> is f.g. and projective;
  (b) Ω : A → Hom<sub>-B</sub>(Q, B), a ↦ [q ↦ a ← q] is a bimodule isomorphism;
  (c) \*<sub>C</sub>Q is faithful;
- 11. (a)  ${}_{B}A$  is f.g. and projective; (b)  $\Lambda : {}^{*}C \longrightarrow \operatorname{End}({}_{B}A)^{op}, g \mapsto [a \mapsto a \leftarrow g]$  is a ring isomorphism.

**Proof.** The result follows by Theorems 1.11, 2.3 and the fact that in case  ${}_{A}\mathcal{C}$  is f.g. and projective  $\mathcal{M}^{\mathcal{C}} = \mathcal{M}_{*\mathcal{C}} = \sigma[\mathcal{C}_{*\mathcal{C}}]$ .

**Theorem 2.10.** (Morita, e.g. [Fai81, 4.1.3, 4.3], [MZ97, 2.6]). Let  $\mathcal{R}$  be a ring, P a right  $\mathcal{R}$ -module,  $\mathcal{S} := \operatorname{End}(P_{\mathcal{R}})$  and  $P^* := \operatorname{Hom}_{\mathcal{R}}(P, \mathcal{R})$ .

- 1. The following are equivalent:
  - (a)  $P_{\mathcal{R}}$  is a generator;
  - (b)  ${}_{\mathcal{S}}P$  is f.g. projective and  $\mathcal{R} \simeq \operatorname{End}({}_{\mathcal{S}}P)^{op}$  canonically.
- 2. The following are equivalent:
  - (a)  $P_{\mathcal{R}}$  is a faithful quasiprogenerator and  $_{\mathcal{S}}P$  is finitely generated;
  - (b)  $P_{\mathcal{R}}$  is a progenerator;
  - (c)  $_{S}P$  is a progenerator and  $P_{\mathcal{R}}$  is faithfully balanced;
  - (d)  $P_{\mathcal{R}}$  and  ${}_{\mathcal{S}}P$  are generators;
  - (e)  $P_{\mathcal{R}}$  and  ${}_{\mathcal{S}}P$  are f.g. and projective;
  - (f)  $\operatorname{Hom}_{-\mathcal{R}}(P,-): \mathcal{M}_{\mathcal{R}} \longrightarrow \mathcal{M}_{\mathcal{S}}$  is an equivalence with inverse  $\operatorname{Hom}_{-\mathcal{S}}(P^*,-);$
  - $(g) \otimes_{\mathcal{R}} P^* : \mathcal{M}_{\mathcal{R}} \longrightarrow \mathcal{M}_{\mathcal{S}} \text{ is an equivalence with inverse } \otimes_{\mathcal{S}} P.$

As a consequence of Theorems 2.5 and 2.10 we get

**Theorem 2.11.** Assume  $_{A}C$  to be f.g. and projective. Then the following are equivalent:

- 1.  $\mathcal{M}^{\mathcal{C}}$  satisfies the strong structure theorem;
- 2.  $_{B}A$  is faithfully flat and A/B is C-Galois;

- 3. <sub>B</sub>A is faithfully flat and  $\beta' := \Psi'_{\mathcal{C}}$  is bijective;
- 4. <sub>B</sub>A is faithfully flat and for every  $M \in \sigma[A_{*C}]$ , the map  $\Psi'_{M}$  is bijective;
- 5.  $A_{*C}$  is quasiprojective and generates each of its submodules,  ${}_{B}A$  is a weak generator and  $\mathcal{M}_{*C} = \sigma[A_{*C}];$
- 6.  $A_{*\mathcal{C}}$  is a quasiprogenerator and  $\mathcal{M}_{*\mathcal{C}} = \sigma[A_{*\mathcal{C}}];$
- 7. <sub>B</sub>A is a weak generator,  $\Psi'_M$  is an isomorphism for every  $M \in \text{Gen}(A_{*C})$  and  $\mathcal{M}_{*C} = \sigma[A_{*C}]$ ;
- 8.  $A_{*C}$  is a faithful quasiprogenerator and  ${}_{B}A$  is finitely generated;
- 9.  $_{B}A$  is a progenerator and  $A_{*C}$  is faithfully balanced;
- 10. Hom<sub>-\*C</sub>(A, -):  $\mathcal{M}_{*C} \longrightarrow \mathcal{M}_B$  is an equivalence with inverse Hom<sub>-B</sub>(Q, -);
- 11.  $-\otimes_{*\mathcal{C}} Q: \mathcal{M}_{*\mathcal{C}} \longrightarrow \mathcal{M}_B$  is an equivalence with inverse  $-\otimes_B A$ ;
- 12.  $A_{*C}$  and  ${}_{B}A$  are generators;
- 13.  $A_{*C}$  and  ${}_{B}A$  are f.g. and projective;
- 14.  $A_{*C}$  is a progenerator.

## **3** Cleft C-Galois Extensions

In what follows R is a commutative ring with  $1_R \neq 0_R$  and  $\mathcal{M}_R$  is the category of R-(bi)modules. For an R-coalgebra  $(C, \Delta_C, \varepsilon_C)$  and an R-algebra  $(A, \mu_A, \eta_A)$  we consider  $(\operatorname{Hom}_R(C, A), \star) := \operatorname{Hom}_R(C, A)$  as an R-algebra with the usual convolution product  $(f \star g)(c) := \sum f(c_1)g(c_2)$  and unity  $\eta_A \circ \varepsilon_C$ . The unadorned  $- \otimes - \operatorname{means} - \otimes_R -$ .

**3.1. Entwined modules.** A right-right entwining structure  $(A, C, \psi)$  over R consists of an R-algebra  $(A, \mu_A, \eta_A)$ , an R-coalgebra  $(C, \Delta_C, \varepsilon_C)$  and an R-linear map

$$\psi: C \otimes_R A \longrightarrow A \otimes_R C, \ c \otimes a \mapsto \sum a_{\psi} \otimes c^{\psi},$$

such that

$$\sum (a\widetilde{a})_{\psi} \otimes c^{\psi} = \sum a_{\psi}\widetilde{a}_{\Psi} \otimes c^{\psi\Psi}, \qquad \sum (1_A)_{\psi} \otimes c^{\psi} = 1_A \otimes c,$$
  
 
$$\sum a_{\psi} \otimes \Delta_C(c^{\psi}) = \sum a_{\psi\Psi} \otimes c_1^{\Psi} \otimes c_2^{\psi}, \qquad \sum a_{\psi}\varepsilon_C(c^{\psi}) = \varepsilon_C(c)a.$$

**3.2.** Let  $(A, C, \psi)$  be a right-right entwining structure. An **entwined module** corresponding to  $(A, C, \psi)$  is a right A-module, which is also a right C-comodule through  $\rho_M$ , such that

 $\varrho_M(ma) = \sum m_{<0>} a_{\psi} \otimes m_{<1>}^{\psi}$  for all  $m \in M$  and  $a \in A$ .

The category of right-right entwined modules and A-linear C-colinear morphisms is denoted by  $\mathcal{M}_A^C(\psi)$ . For  $M, N \in \mathcal{M}_A^C(\psi)$  we denote by  $\operatorname{Hom}_A^C(M, N)$  the set of A-linear C-colinear morphisms from M to N. With  $\#_{\psi}^{op}(C, A) := \operatorname{Hom}_R(C, A)$ , we denote the A-ring with  $(af)(c) = \sum a_{\psi}f(c^{\psi}), (fa)(c) = f(c)a$ , multiplication  $(f \cdot g)(c) = \sum f(c_2)_{\psi}g(c_1^{\psi})$  and unity  $\eta_A \circ \varepsilon_C$  (see [Abu03, Lemma 3.3]).

Entwined modules were introduced by T. Brzeziński and S. Majid [BM98] as a generalization of the Doi-Koppinen modules presented in [Doi92] and [Kop95]. By a remark of M. Takeuchi (e.g. [Brz02, Proposition 2.2]), we have an A-coring structure on  $\mathcal{C} := A \otimes_R C$ , where  $\mathcal{C}$  is an A-bimodule through  $a(\tilde{a} \otimes c) := a\tilde{a} \otimes c$ ,  $(\tilde{a} \otimes c)a := \sum \tilde{a}a_{\psi} \otimes c^{\psi}$  and has comultiplication

$$\Delta_{\mathcal{C}}: A \otimes_R C \longrightarrow (A \otimes_R C) \otimes_A (A \otimes_R C), \ a \otimes c \mapsto \sum (a \otimes c_1) \otimes_A (1_A \otimes c_2)$$

and counity  $\varepsilon_{\mathcal{C}} := id_A \otimes \varepsilon_C$ . Moreover  $\mathcal{M}^C_A(\psi) \simeq \mathcal{M}^C$ ,  $\#^{op}_{\psi}(C, A) \simeq {}^*\mathcal{C}$  as A-rings and  ${}_A\mathcal{C}$  is flat (resp. f.g., projective), if  ${}_RC$  is so (e.g. [Abu03]).

Inspired by [Doi94, 3.1] we make the following definition:

**3.3.** Let  $(A, C, \psi)$  be a right-right entwining structure over R and consider the corresponding A-coring  $\mathcal{C} := A \otimes_R C$ . We say that  $(A, C, \psi)$  satisfies the **left**  $\alpha$ -condition, if for every right A-module M the following map is injective

$$\alpha_M^{\psi}: M \otimes_R C \longrightarrow \operatorname{Hom}_R(\#_{\psi}^{op}(C, A), M), \ m \otimes c \mapsto [f \mapsto mf(c)]$$

(equivalently, if  ${}_{A}\mathcal{C}$  is locally projective).

Let M be a right  $\#^{op}_{\psi}(C, A)$ -module M and consider the canonical map  $\rho_M : M \longrightarrow$ Hom<sub>R</sub> $(\#^{op}_{\psi}(C, A), M)$ . Set Rat<sup>C</sup> $(M_{\#^{op}_{\psi}(C, A)}) := (\rho^{\psi}_M)^{-1}(M \otimes_R C)$ . We call M #-rational, if Rat<sup>C</sup> $(M_{\#^{op}_{\psi}(C, A)}) = M$  and set  $\rho_M := (\alpha^{\psi}_M)^{-1} \circ \rho_M$ . The category of #-rational right  $\#^{op}_{\psi}(C, A)$ -modules will be denoted by Rat<sup>C</sup> $(\mathcal{M}_{\#^{op}_{\psi}(C, A)})$ .

**Theorem 3.4.** ([Abu03, Theorem 3.10]) Let  $(A, C, \psi)$  be a right-right entwining structure and consider the corresponding A-coring  $C := A \otimes_R C$ .

- 1. If <sub>R</sub>C is flat, then  $\mathcal{M}^{C}_{A}(\psi)$  is a Grothendieck category with enough injective objects.
- 2. If  $_{R}C$  is locally projective (resp. f.g. and projective), then

$$\mathcal{M}_{A}^{C}(\psi) \simeq \operatorname{Rat}^{C}(\mathcal{M}_{\#_{\psi}^{op}(C,A)}) \simeq \sigma[(A \otimes_{R} C)_{\#_{\psi}^{op}(C,A)}] \ (resp. \ \mathcal{M}_{A}^{C}(\psi) \simeq \mathcal{M}_{\#_{\psi}^{op}(C,A)}). \ (5)$$

In what follows we fix a right-right entwining structure  $(A, C, \psi)$  with  $\mathcal{C} := A \otimes_R C$  the corresponding A-coring and assume that  $A \in \mathcal{M}_A^C(\psi) \simeq \mathcal{M}^C$  with

$$\varrho_A: A \longrightarrow A \otimes_R C, \ a \mapsto \sum a_{<0>} \otimes a_{<1>} = \sum 1_{<0>} a_{\psi} \otimes 1_{<1>}^{\psi}.$$

Then  $\sum \mathbf{1}_{<0>} \otimes \mathbf{1}_{<1>} \in \mathcal{C}$  is a group-like element and

$$Q \simeq \{q \in \operatorname{Hom}_R(C, A) \mid \sum q(c_2)_{\psi} \otimes c_1^{\psi} = \sum q(c) 1_{<0>} \otimes 1_{<1>} \text{ for all } c \in C\}.$$

For every  $M \in \mathcal{M}_A^C(\psi)$ , we set

$$M^{co\mathcal{C}} := \{ m \in M \mid \sum m_{<0>} \otimes m_{<1>} = \sum m \mathbb{1}_{<0>} \otimes \mathbb{1}_{<1>} \}.$$

Moreover we set  $B := A^{co\mathcal{C}}$ .

Remark 3.5. Let  $x \in C$  be a group-like element. For every right *C*-comodule *M* we put  $M^{coC} := \{m \in M \mid \varrho_M(m) = m \otimes x\}$ . If  $\varrho_A(1_A) = 1_A \otimes x$ , then we have  $M^{coC} = M^{coC}$  for every  $M \in \mathcal{M}_A^C(\psi)$ .

By [Brz99, Corollaries 3.4, 3.7]  $- \otimes_R^c A : \mathcal{M}^C \longrightarrow \mathcal{M}_A^C(\psi)$  is a functor, which is left adjoint to the forgetful functor. Here, for every  $N \in \mathcal{M}^C$ , we consider the canonical right A-module  $N \otimes_R^c A := N \otimes_R A$  with the C-coaction  $n \otimes a \mapsto \sum n_{<0>} \otimes a_{\psi} \otimes n_{<1>}^{\psi}$ .

**Proposition 3.6.** Let R be a QF ring and assume C be right semiperfect. Let <sub>R</sub>C to be locally projective (projective) and put  $C^{\Box} := \operatorname{Rat}(*_{C}C^{*})$ .

- 1. The following are equivalent:
  - (a) A is a generator in  $\mathcal{M}_A^C(\psi)$ ;
  - (b) A generates  $C^{\Box} \otimes_R^c A$  in  $\mathcal{M}_A^C(\psi)$ ;
  - (c) the map  $\Psi'_{C^{\Box}\otimes^c_{P}A}$ : Hom $^{C}_{A}(A, C^{\Box}\otimes^c_{R}A)\otimes_{B}A \longrightarrow C^{\Box}\otimes^c_{R}A$  is surjective (bijective).
- 2. The following are equivalent:
  - (a) A is a progenerator in  $\mathcal{M}_A^C(\psi)$ ;
  - (b)  $\Psi'_{C^{\Box} \otimes c_{P}A}$  is surjective (bijective) and  ${}_{B}A$  is a weak generator.

**Proof.** By [MTW01, 2.6]  $C^{\Box}$  is a generator in  $\mathcal{M}^{C}$ , hence  $C^{\Box} \otimes_{R}^{c} A$  is a generator in  $\mathcal{M}^{C}_{A}(\psi)$  by the functorial isomorphism  $\operatorname{Hom}^{C}_{A}(C^{\Box} \otimes_{R}^{c} A, M) \simeq \operatorname{Hom}^{C}(C^{\Box}, M)$  for every  $M \in \mathcal{M}^{C}_{A}(\psi)$ .

- 1. The assertions follow form the note above and Theorem 2.3.
- 2. (a)  $\Rightarrow$  (b) follows by Theorem 2.5.

(b)  $\Rightarrow$  (a). By the note above  $C^{\Box} \otimes_{R}^{c} A$  is a generator in  $\mathcal{M}_{A}^{C}(\psi)$ , and the surjectivity of  $\Psi'_{C^{\Box} \otimes_{R}^{c} A}$  makes A a generator in  $\mathcal{M}_{A}^{C}(\psi)$ . So  $_{B}A$  is flat by Theorem 2.3. The weak generator property makes  $_{B}A$  faithfully flat and we are done by Theorem 2.5.

**Definition 3.7.** A (total) integral for *C* is a *C*-colinear morphism  $\lambda : C \longrightarrow A$  (with  $\sum 1_{<0>\lambda}(1_{<1>}) = 1_A$ ). We call the ring extension A/B cleft, if there exists a \*-invertible integral. We say *A* has the right normal basis property, if there exists a left *B*-linear right *C*-colinear isomorphism  $A \simeq B \otimes_R C$ .

**Lemma 3.8.** Let  $\lambda \in \text{Hom}_R(C, A)$  be  $\star$ -invertible with inverse  $\overline{\lambda}$ . Then:

- 1.  $\lambda \in \operatorname{Hom}^{C}(C, A)$  iff  $\overline{\lambda} \in Q$ .
- 2. If  $\rho(a) = \sum a_{\psi} \otimes x^{\psi}$  for some group-like element  $x \in C$ , then there exists  $\widehat{\lambda} \in Q$ , such that  $\sum 1_{<0>}\widehat{\lambda}(1_{<1>}) = \widehat{\lambda}(x) = 1_A$  (in this case C admits a total integral, namely the  $\star$ -inverse of  $\widehat{\lambda}$ ).

**Proof.** Let  $\lambda \in \operatorname{Hom}_R(C, A)$  be  $\star$ -invertible with inverse  $\overline{\lambda}$ .

1. If  $\overline{\lambda} \in Q$ , then we have for all  $c \in C$ :

$$\sum \lambda(c_1) \otimes c_2 = \sum \lambda(c_1) 1_{\psi} \otimes c_2^{\psi}$$

$$= \sum \lambda(c_1) \varepsilon(c_3) 1_{\psi} \otimes c_2^{\psi}$$

$$= \sum \lambda(c_1) (\overline{\lambda}(c_3) \lambda(c_4))_{\psi} \otimes c_2^{\psi}$$

$$= \sum \lambda(c_1) \overline{\lambda}(c_3)_{\psi} \lambda(c_4)_{\Psi} \otimes c_2^{\psi\Psi}$$

$$= \sum \lambda(c_1) \overline{\lambda}(c_{22})_{\psi} \lambda(c_3)_{\Psi} \otimes c_{21}^{\psi\Psi}$$

$$= \sum \lambda(c_1) \overline{\lambda}(c_2) 1_{<0>\lambda} \lambda(c_3)_{\Psi} \otimes 1_{<1>}^{\Psi}$$

$$= \sum 1_{<0>\lambda} \lambda(c)_{\Psi} \otimes 1_{<1>}^{\Psi}$$

$$= \sum \lambda(c)_{<0>} \otimes \lambda(c)_{<1>}$$

i.e.  $\lambda \in \text{Hom}^{C}(C, A)$ . On the other hand, if  $\lambda \in \text{Hom}^{C}(C, A)$ , then we have for all  $c \in C$ :

$$\begin{split} \sum \overline{\lambda}(c_2)_{\psi} \otimes c_1^{\psi} &= \sum \overline{\lambda}(c_1)\lambda(c_2)\overline{\lambda}(c_4)_{\psi} \otimes c_3^{\psi} \\ &= \sum \overline{\lambda}(c_1)\lambda(c_2)_{<0>}\overline{\lambda}(c_3)_{\psi} \otimes \lambda(c_2)_{<1>}^{\psi} \\ &= \sum \overline{\lambda}(c_1)1_{<0>}\lambda(c_2)_{\psi}\overline{\lambda}(c_3)_{\Psi} \otimes 1_{<1>}^{\psi\Psi} \\ &= \sum \overline{\lambda}(c_1)1_{<0>}(\lambda(c_2)\overline{\lambda}(c_3))_{\psi} \otimes 1_{<1>}^{\psi} \\ &= \sum \overline{\lambda}(c)1_{<0>}1_{\psi} \otimes 1_{<1>}^{\psi} \\ &= \sum \overline{\lambda}(c)1_{<0>} \otimes 1_{<1>}, \end{split}$$

i.e.  $\overline{\lambda} \in Q$ .

2. Assume  $\rho(a) = \sum a_{\psi} \otimes x^{\psi}$  for some group-like element  $x \in C$ . Let  $\lambda \in \text{Hom}^{C}(C, A)$ with  $\overline{\lambda} \in Q$  (see (1)). Then  $\widehat{\lambda} := \overline{\lambda}\lambda(x) \in Q$ , since  $\lambda(x) \in B$ , and moreover  $\sum 1_{<0>}\widehat{\lambda}(1_{<1>}) = \widehat{\lambda}(x) = \overline{\lambda}(x)\lambda(x) = (\overline{\lambda} \star \lambda)(x) = \varepsilon_{C}(x)1_{A} = 1_{A}$ .

**Proposition 3.9.** Assume A/B to be cleft.

- 1.  $\mathcal{M}_{A}^{C}(\psi)$  satisfies the weak structure theorem (in particular A/B is C-Galois).
- 2. For every  $M \in \mathcal{M}_A^C(\psi)$ , the C-colinear morphism

$$\gamma_M: M \longrightarrow M^{co\mathcal{C}} \otimes_R C, \ m \mapsto \sum m_{<0>} \overline{\lambda} \otimes m_{<1>}$$

is an isomorphism.

- 3. A has the right normal basis property.
- 4. If <sub>R</sub>C is faithfully flat, then  $\mathcal{M}_A^C(\psi)$  satisfies the strong structure theorem.

**Proof.** Assume there exists a  $\star$ -invertible  $\lambda \in \text{Hom}^{C}(C, A)$  with inverse  $\overline{\lambda} \in Q$  (see Lemma 3.8 (1)).

1. Let  $M \in \mathcal{M}_A^C(\psi)$  and consider

$$\widetilde{\Psi}_M: M \longrightarrow M^{co\mathcal{C}} \otimes_B A, \ m \mapsto \sum m_{<0>} \overline{\lambda} \otimes \lambda(m_{<1>}).$$

Then we have for all  $n \in M^{co\mathcal{C}}$ ,  $m \in M$  and  $a \in A$ :

$$\begin{aligned} (\widetilde{\Psi}_{M} \circ \Psi_{M})(n \otimes a) &= \widetilde{\Psi}_{M}(na) \\ &= \sum (na_{<0>})\overline{\lambda} \otimes_{B} \lambda(a_{<1>}) \\ &= \sum na_{<0><0>}\overline{\lambda}(a_{<0><1>}) \otimes_{B} \lambda(a_{<1>}) \\ &= \sum n \otimes_{B} a_{<0><0>}\overline{\lambda}(a_{<0><1>})\lambda(a_{<1>}) \\ &= \sum n \otimes_{B} a_{<0>}\overline{\lambda}(a_{<1>1})\lambda(a_{<1>2}) \\ &= n \otimes_{B} a \end{aligned}$$

and

$$(\Psi_M \circ \widetilde{\Psi}_M)(m) = \sum (m_{<0>}\overline{\lambda})\lambda(m_{<1>})$$

$$= \sum m_{<0><0>}\overline{\lambda}(m_{<0><1>})\lambda(m_{<1>})$$

$$= \sum m_{<0>}\overline{\lambda}(m_{<1>1})\lambda(m_{<1>2})$$

$$= \sum m_{<0>}\varepsilon_C(m_{<1>})1_A$$

$$= m.$$

2. For every  $M \in \mathcal{M}_A^C(\psi)$ ,  $\gamma_M$  is bijective with inverse

$$\widetilde{\gamma}_M: M^{co\mathcal{C}} \otimes_R C \longrightarrow M, \ n \otimes c \mapsto n\lambda(c).$$

In fact we have for all  $m \in M, n \in M^{co\mathcal{C}}$  and  $c \in C$ :

$$\begin{split} (\widetilde{\gamma}_{M} \circ \gamma_{M})(m) &= \sum (m_{<0>}\overline{\lambda})\lambda(m_{<1>}) \\ &= \sum m_{<0><0>}\overline{\lambda}(m_{<0><1>})\lambda(m_{<1>}) \\ &= \sum m_{<0>}\overline{\lambda}(m_{<1>1})\lambda(m_{<1>2}) \\ &= \sum m_{<0>}\varepsilon_{C}(m_{<1>}) \\ &= m \end{split}$$

and

$$\begin{aligned} (\gamma_M \circ \widetilde{\gamma}_M)(n \otimes_B c) &= \sum (n\lambda(c))_{<0>} \overline{\lambda} \otimes (n\lambda(c))_{<0>} \\ &= \sum (n\lambda(c)_{<0>}) \overline{\lambda} \otimes \lambda(c)_{<1>} \\ &= \sum (n\lambda(c_1)) \overline{\lambda} \otimes c_2 \\ &= \sum n\lambda(c_1)_{<0>} \overline{\lambda}(\lambda(c_1)_{<1>}) \otimes c_2 \\ &= \sum n\lambda(c_{11}) \overline{\lambda}(c_{12}) \otimes c_2 \\ &= n \otimes c. \end{aligned}$$

3. By (2) the left *B*-linear right *C*-colinear map

$$\gamma_A: A \longrightarrow B \otimes_R C, a \mapsto \sum a_{<0>} \leftarrow \overline{\lambda} \otimes a_{<1>}$$

is an isomorphism with inverse  $b \otimes c \mapsto b\lambda(c)$ .

4. Assume  $_{R}C$  to be faithfully flat. By (3)  $A \simeq B \otimes_{R} C$  as left *B*-modules, hence  $_{B}A$  is faithfully flat. By (1) A/B is *C*-Glaois and we are done by Theorem 2.5.

**Theorem 3.10.** The following statements are equivalent:

- 1. A/B is cleft;
- 2.  $\mathcal{M}^{C}_{A}(\psi)$  satisfies the weak structure theorem and A has the right normal basis property;
- 3. A/B is C-Galois and A has the right normal basis property;
- 4.  $\Lambda: \#^{op}_{\psi}(C, A) \simeq \operatorname{End}({}_{B}A)^{op}, g \mapsto [a \mapsto a \leftarrow g]$  is a ring isomorphism and A has the right normal basis property.

If moreover  $_{R}C$  is faithfully flat, then (1)-(4) are equivalent to

5.  $\mathcal{M}_A^C(\psi)$  satisfies the strong structure theorem and A has the right normal basis property.

**Proof.**  $(1) \Rightarrow (2)$ . This follows by Proposition 3.9.

(2)  $\Rightarrow$  (3). By assumption  $\beta := \Psi_{A \otimes_R C}$  is an isomorphism.

(3)  $\Rightarrow$  (4). By assumption  $A \otimes_B A \simeq A \otimes_R C$  as left A-modules, hence we have the canonical isomorphisms

$$\#_{\psi}^{op}(C,A) \simeq \operatorname{Hom}_{A-}(A \otimes_{R} C,A) \simeq \operatorname{Hom}_{A-}(A \otimes_{B} A,A) \simeq \operatorname{Hom}_{B-}(A,\operatorname{End}(_{A}A)) \simeq \operatorname{End}(_{B}A).$$

 $(4) \Rightarrow (1)$ . Assume  $\theta : B \otimes_R C \longrightarrow A$  to be a left *B*-linear right *C*-colinear isomorphism and consider the right *C*-colinear morphism  $\lambda : C \longrightarrow A$ ,  $c \mapsto \theta(1_A \otimes c)$  and the left *B*linear morphism  $\delta := (id \otimes \varepsilon_C) \circ \theta^{-1} : A \longrightarrow B$ . Define  $\overline{\lambda} := \Lambda^{-1}(\delta) \in \#^{op}_{\psi}(C, A)$ . Then we have for all  $c \in C$ :

$$\sum \lambda(c_1)\overline{\lambda}(c_2) = \sum \lambda(c)_{<0>}\overline{\lambda}(\lambda(c)_{<1>}) = \lambda(c) \leftarrow \overline{\lambda}$$
  
=  $\delta(\lambda(c)) = ((id \otimes \varepsilon_C) \circ \theta^{-1})(\theta(1_A \otimes c)) = \varepsilon_C(c)1_A.$ 

On the other hand we have for all  $a \in A$ :

$$\begin{split} \Lambda(\overline{\lambda}\star\lambda)(a) &= a \leftarrow (\overline{\lambda}\star\lambda) &= \sum a_{<0>}(\overline{\lambda}\star\lambda)(a_{<1>}) \\ &= \sum a_{<0>}\overline{\lambda}(a_{<1>1})\lambda(a_{<1>2}) &= \sum a_{<0><0>}\overline{\lambda}(a_{<0><1>1})\lambda(a_{<1>}) \\ &= \sum (a_{<0>} \leftarrow \overline{\lambda})\lambda(a_{<1>}) &= \sum (a_{<0>} \leftarrow \Lambda^{-1}(\delta))\lambda(a_{<1>}) \\ &= \sum \delta(a_{<0>})\lambda(a_{<1>}) &= \sum \delta(a_{<0>})\theta(1_A\otimes a_{<1>}) \\ &= \sum \theta(\delta(a_{<0>})\otimes a_{<1>}) &= \theta(\theta^{-1}(a)) = a, \end{split}$$

hence  $\overline{\lambda} \star \lambda = \eta_A \circ \varepsilon_C$ .

Now assume  $_{R}C$  to be faithfully flat. Then  $(1) \Rightarrow (5)$  follows by Proposition 3.9 (4) and we are done.

The following result deals with the special case  $\rho(a) = \sum a_{\psi} \otimes x^{\psi}$ , for some grouplike element  $x \in C$ . In this case we obtain the equivalent statements (1)-(5) in Theorem 3.10 without any assumptions on C. **Theorem 3.11.** Assume that  $\rho(a) = \sum a_{\psi} \otimes x^{\psi}$  for some group-like element  $x \in C$ . The following statements are equivalent:

- 1. A/B is cleft;
- 2.  $\mathcal{M}_{A}^{C}(\psi)$  satisfies the strong structure theorem and A has the right normal basis property;
- 3.  $\mathcal{M}^{C}_{A}(\psi)$  satisfies the weak structure theorem and A has the right normal basis property;
- 4. A/B is C-Galois and A has the right normal basis property;
- 5.  $\Lambda: \#^{op}_{\psi}(C, A) \simeq \operatorname{End}({}_{B}A)^{op}, g \mapsto [a \mapsto a \leftarrow g]$  is a ring isomorphism and A has the right normal basis property.

**Proof.** By Theorem 3.10 it remains to prove that  $\Phi_N$  is an isomorphism for every  $N \in \mathcal{M}_B$ , if A/B is cleft. But in our special case there exists by Lemma 3.8 some  $\hat{\lambda} \in Q$  with  $\sum 1_{<0>} \hat{\lambda}(1_{<1>}) = 1_A$  and we are done by Corollary 1.8 (2).

*Remark* 3.12. Let (H, A, C) be a right-right resp. a left-right Doi-Koppinen structure. Then  $(A, C, \psi)$  is a right-right entwining structure with

$$\psi: C \otimes_R A \longrightarrow A \otimes_R C, \ c \otimes a \mapsto \sum a_{<0>} \otimes ca_{<1>}$$

resp. a left-right entwining structure with

$$\psi: A \otimes_R C \longrightarrow A \otimes_R C, \ a \otimes c \mapsto \sum a_{<0>} \otimes a_{<1>}c.$$

If x is a group-like element of C, then  $A \in \mathcal{M}(H)_A^C$  with  $\varrho(a) := \sum a_{<0>} \otimes xa_{<1>}$  (resp.  $\varrho(a) = \sum a_{<0>} \otimes a_{<1>}x$ ) and we get [DM92, Theorem 1.5] (resp. [Doi94, Theorem 2.5]) as special cases of Theorem 3.11.

### References

- [Abu03] J.Y. Abuhlail, *Rational modules for corings*, Commun. Algebra **31**, 5793-5840 (2003).
- [BDR97] M. Beattie, D. Dăscălescu and Ş. Raianu, Galois extensions for co-Frobenius Hopf algebras, J. Algebra 198, 164-183 (1997).
- [BM98] T. Brzeziński and S. Majid, *Coalgebra bundles*, Comm. Math. Phys. **191**, 467-492 (1998).
- [Brz02] T. Brzeziński, The structure of corings. Induction functors, Maschke-type theorem, and Frobenius and Galois-type properties, Algebr. Represent. Theory 5, 389-410 (2002).
- [Brz99] T. Brzeziński, On modules associated to coalgebra Galois extensions, J. Algebra 215, 290-317 (1999).

- [CFM90] M. Cohen, D. Fischman and S. Montgomery, Hopf Galois extensions, smash products and Morita equivalence, J. Algebra 133, 351-372 (1990).
- [CMZ02] S. Caenepeel, G. Militaru, and S. Zhu, Frobenius and Separable Functors for Generalized Module Categories and Nonlinear Equations, Lect. Not. Math. 1787, Springer-Verlag, Berlin (2002).
- [CS69] S. Chase and M. Sweedler, *Hopf algebras and Galois theory*, Lec. Not. Math.97 Springer-Verlag Berlin (1969).
- [CVW04] S. Caenepeel, J. Vercruysse and S. Wang, Morita Theory for corings and cleft entwining structures, J. Algebra **276**, 210-235 (2004).
- [DM92] Y. Doi and A. Masuoka, *Generalization of cleft comodule algebras*, Comm. Algebra **20**, 3703-3721 (1992).
- [Doi94] Y. Doi, Generalized smash products and Morita contexts for arbitrary Hopf algebras, Bergen-Montgomery (ed.), Advances in Hopf algebras, Marcle Dekker, Lec. Notes Pure Applied Maths 158, 39-53 (1994).
- [Doi92] Y. Doi, Unifying Hopf modules, J. Algebra 153, 373-385 (1992).
- [Doi83] Y. Doi, On the structure of relative Hopf modules, Comm. Algebra 11(3), 243-255 (1983).
- [DT89] Y. Doi and M. Takeuchi, Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and the Azumaya algebras, J. Algebra, **121**, 488-516 (1989).
- [DT86] Y. Doi and M. Takeuchi, Cleft comodule algebras for a bialgebra, Comm. Algebra 14, 801-817 (1986).
- [Fai81] C. Faith, Algebra I, Rings, Modules and Categories, Springer-Verlag (1981).
- [Kop95] M. Koppinen, Variations on the smash product with applications to groupgraded rings, J. Pure Appl. Algebra **104**, 61-80 (1995).
- [KT81] H. Kreimer and M. Takeuchi, Hopf Algebras and Galois extensions of an algebra, Indiana Univ. Math. J. 30, 675-692 (1981).
- [MSTW01] C. Menini, A. Seidel, B. Torrecillas and R. Wisbauer, A-H-bimodules and equivalences, Comm. Algebra **29(10)**, 4619-4640 (2001).
- [MTW01] C. Menini, B. Torrecillas and R. Wisbauer, Strongly rational comodules and semiperfect Hopf algebras over QF Rings, J. Pure Appl. Algebra 155, 237-255 (2001).
- [MZ97] C. Menini and M. Zuccoli, Equivalence theorems and Hopf-Galois extensions, J. Algebra 194, 245-274 (1997).
- [Sch90] H-J. Schneider, Principal homogenous spaces for arbitrary Hopf algebras, Israel
   J. Math. 72(1-2) 167-195 (1990).

- [Swe75] M. Sweedler, *The Predual theorem to the Jacobson-Bourbaki theorem*, Trans. Amer. Math. Soc. **213**, 391-406 (1975).
- [Wis02] R. Wisbauer, On the category of comodules for corings, Proc. 3rd. Int. Pal. Conf.: Math. & Math. Edu., Bethlehem (Palestine), S. Elaydi et al. (ed.), World Scientific, New Jersy, ISBN 981-02-4720-6, 325-336 (2002).
- [Wis88] R. Wisbauer, Grundlagen der Modul- und Ringtheorie, München: Verlag Reinhard Fischer (1988); Foundations of Module and Ring Theory, Gordon and Breach, Reading (1991).
- [Z-H76] B. Zimmermann-Huignes, Pure submodules of direct products of free modules, Math. Ann. 224, 233-245 (1976).