

ON GRIESS ALGEBRAS

MICHAEL ROITMAN

A vertex algebra V is a linear space, endowed with infinitely many bilinear products $(n) : V \otimes V \rightarrow V$ and a unit $\mathbb{1} \in V$, satisfying certain axioms, see § 1.1. In this paper we deal with graded vertex algebras $V = \bigoplus_{i \in \mathbb{Z}} V_i$, so that $V_i(n)V_j \subseteq V_{i+j-n-1}$ and $\mathbb{1} \in V_0$. A vertex algebra is called OZ (abbreviation of “One-Zero”) if it is graded so that $\dim V_0 = 1$ and $V_i = 0$ for $i = 1$ or $i < 0$ [12]. If V is an OZ vertex algebra, then V_2 is a commutative (but not necessary associative) algebra with respect to the product $V_2(1)V_2 \rightarrow V_2$, with an invariant symmetric bilinear form, given by $V_2(3)V_2 \rightarrow V_0$. It is called *the Griess algebra of V* .

Formulation of the results. In this paper we prove the following

Theorem 1. *Let A be a commutative algebra with non-degenerate symmetric invariant bilinear form. Then there is an OZ vertex algebra V such that $A \subseteq V_2$.*

This is a special case of the following conjecture:

Conjecture. (a) *For any commutative algebra A with a non-degenerate symmetric invariant bilinear form there is an OZ vertex algebra V such that $A = V_2$.*
 (b) *If $\dim A < \infty$, then V can be chosen so that $\dim V_n < \infty$.*
 (c) *If A has a unit e , then V can be chosen so that e is a Virasoro vector of V , see § 1.1 for the definition.*
 (d) *If $G \subset \text{Aut } A$ is a finite group of automorphisms of A , then V can be chosen so that $G \subset \text{Aut } V$.*

We construct V in three steps. First we construct a vertex algebra \widehat{V} , which is generated by A with some simple relations. We prove an analogue of Poincaré-Birkhoff-Witt theorem for \widehat{V} . There is a natural filtration

$$0 = \widehat{V}^{(-1)} \subset \widehat{V}^{(0)} \subset \widehat{V}^{(1)} \subset \widehat{V}^{(2)} \subset \dots \subset \widehat{V}$$

on \widehat{V} , so that the associated graded vertex algebra $\text{gr } \widehat{V} = \bigoplus_{i \geq 0} \widehat{V}^{(i)} / \widehat{V}^{(i-1)}$ is isomorphic to a free vertex algebra (see Theorem 2 in § 3.1).

Next we construct an algebra \overline{V} , which is a homomorphic image of \widehat{V} . The algebra \overline{V} has the following properties, summarized in Theorem 3 in § 3.4: $\overline{V}_n = 0$, for $n = 1$ or $n < 0$, \overline{V}_2 contains A , and \overline{V}_0 is isomorphic to a polynomial algebra in infinitely many variables. The main technique that we use here is the Diamond lemma and Gröbner-Shirshov bases.

Finally, we observe that any invariant bilinear form $\langle \cdot | \cdot \rangle$ on \overline{V} must coincide with the given bilinear form on A , and there are forms $\langle \cdot | \cdot \rangle$ on \overline{V} such that the codimension of $\text{Rad} \langle \cdot | \cdot \rangle \cap \overline{V}_0$ in \overline{V}_0 is 1. Since the form on A is non-degenerate, the quotient vertex algebra $V = \overline{V} / \text{Rad} \langle \cdot | \cdot \rangle$ satisfies the requirements of Theorem 1.

Motivations of the problem. Probably the most famous example of OZ vertex algebras is the Moonshine module V^\natural , constructed by Frenkel, Lepowsky and Meurman in [9], see also [3, 5]. Its Griess algebra V_2^\natural is the original 196884-dimensional commutative algebra constructed by Griess [11]. The automorphism group of V^\natural and of V_2^\natural is the Fischer-Griess Monster.

Another example of OZ vertex algebra is a Virasoro vertex algebra Vir [10, 23]. It is generated by a single Virasoro element $\omega \in Vir_2$ so that the Griess algebra of Vir is $\mathbb{k}\omega$.

If A is associative, then it is well known that A can appear as a Griess algebra, see [14, 24]. Lam [15] also showed the same for a simple Jordan algebra of type A , B or C . Other interesting examples of OZ vertex algebras and their Griess algebras can be found in [12].

We remark that if the graded vertex algebra V is of conformal field theory (CFT) type, which means that $V_n = 0$ for $n < 0$ and $\dim V_0 = 1$, then V_1 is a Lie algebra with respect to the product (0), with invariant bilinear form given by product (1). The analogous problem of finding a vertex algebra V such that V_1 is a given Lie algebra has a well-known solution: to every Lie algebra L with an invariant bilinear form there corresponds an affine Lie algebra \hat{L} , so that a certain highest weight \hat{L} -module has the desired vertex algebra structure [9, 10].

Organization of manuscript. In §1 we fix the notations and give the necessary definitions. We work with vertex algebras with sl_2 structure. They form a more general class of vertex algebras than vertex operator algebras (VOA's) as defined in e.g. [9]. We define the notion of universal enveloping algebra, following [10, 21], and present a description of invariant bilinear forms on a vertex algebra [8, 17, 21]. We also recall the notion of free vertex algebra [19, 20].

In §2 we recall the theory Gröbner-Shirshov bases for associative algebras, mostly following [19]. The standard references are [1, 2, 4, 22].

Then in §3 we prove Theorem 1, modulo several technical statements that require further considerations. These statements are proved in §5 and §6 by working with rational functions, after some preparational arguments presented in §4.

1. GENERAL FACTS ABOUT VERTEX ALGEBRAS

Here we fix the notations and give some minimal definitions. For more details on vertex algebras the reader can refer to the books [9, 13, 16, 24].

1.1. Definition of vertex algebras.

Definition 1.1. A vertex algebra is a linear space V equipped with a family of bilinear products $a \otimes b \mapsto a(n)b$, indexed by integer parameter n , and with an element $\mathbb{1} \in V$, called the unit, satisfying the identities (i)–(iv) below. Let $D : V \rightarrow V$ be the map defined by $Da = a(-2)\mathbb{1}$. Then

- (i) $a(n)b = 0$ for $n \gg 0$,
- (ii) $\mathbb{1}(n)a = \delta_{n,-1}a$ and $a(n)\mathbb{1} = \frac{1}{(-n-1)!} D^{-n-1}a$,
- (iii) $D(a(n)b) = (Da)(n)b + a(n)(Db)$ and $(Da)(n)b = -n a(n-1)b$,
- (iv) $a(m)(b(n)c) - b(n)(a(m)c) = \sum_{s \geq 0} \binom{m}{s} (a(s)b)(m+n-s)c$

for all $a, b, c \in V$ and $m, n \in \mathbb{Z}$.

This is not the only known definition of vertex algebras. Often the axioms are formulated in terms of the *left adjoint action map* $Y : V \rightarrow V[[z, z^{-1}]]$ defined for $a \in V$ by $Y(a, z) = \sum_{n \in \mathbb{Z}} a(n) z^{-n-1}$, where $a(n) : V \rightarrow V$ is the operator given by $b \mapsto a(n)b$. The most important property of these maps is that they are *local*: for any $a, b \in V$ there is $N \geq 0$ such that

$$[Y(a, w), Y(b, z)](w - z)^N = 0. \quad (1.1)$$

The minimal $N = N(a, b)$ for which this identity holds is called *the locality* of a and b . In fact, $N(a, b) = \min\{n \in \mathbb{Z} \mid a(m)b = 0 \ \forall m \geq n\}$.

Remark. One could extend this definition to allow a negative locality N , see [20].

From (iii) we get $Y(Da, z) = \partial_z Y(a, z)$, which in terms of coefficients means

$$[D, a(m)] = -m a(m-1). \quad (1.2)$$

Among other identities that hold in vertex algebras are the quasi-symmetry

$$a(n)b = - \sum_{i \geq 0} (-1)^{n+i} D^{(i)}(b(n+i)a), \quad (1.3)$$

and the associativity identity

$$\begin{aligned} (a(m)b)(n)c &= \sum_{s \geq 0} (-1)^s \binom{m}{s} a(m-s)(b(n+s)c) \\ &\quad - \sum_{s \leq m} (-1)^s \binom{m}{m-s} b(n+s)(a(m-s)c). \end{aligned} \quad (1.4)$$

For $m \geq 0$ this simplifies to

$$(a(m)b)(n)c = \sum_{s=0}^m (-1)^s \binom{m}{s} [a(m-s), b(n+s)]c,$$

which is just another form of the identity (iv) of Definition 1.1.

A vertex algebra V is called *graded* (by the integers) if $V = \bigoplus_{i \in \mathbb{Z}} V_i$ is a graded space, $V_i(n)V_j \subseteq V_{i+j-n-1}$ and $\mathbb{1} \in V_0$.

It is often required that a vertex algebra V is graded and V_2 contains a special element ω such that $\omega(0) = D$, $\omega(1)|_{V_i} = i$ and the coefficients $\omega(n)$ generate a representation of the Virasoro Lie algebra:

$$[\omega(m), \omega(n)] = (m-n)\omega(m+n-1) + \delta_{m+n,2} \frac{1}{2} \binom{m-1}{2} c$$

for some constant $c \in \mathbb{k}$ called the *central charge* of V . In this case V is called *conformal vertex algebra* or *vertex operator algebra*, especially when $\dim V_i < \infty$.

Definition 1.2. [12] A vertex algebra V is called OZ (abbreviation of “One-Zero”) if it is graded so that $V = \mathbb{k}\mathbb{1} \oplus \bigoplus_{n \geq 2} V_n$.

An OZ vertex algebra V has $\dim V_0 = 1$ and $\dim V_1 = 0$, hence the name. The component V_2 is a commutative (but not necessarily associative) algebra with respect to the product $a \otimes b \mapsto ab = a(1)b$, called the *Griess algebra* of V . The commutativity follows from (1.3). The algebra V_2 has a bilinear form $\langle a | b \rangle = a(3)b$. From (1.3) it follows that this form is symmetric, and from (1.4) it follows that $\langle ab | c \rangle = \langle a | bc \rangle$. Bilinear forms with this property are called *invariant*.

Without the condition $V_1 = 0$ the vertex algebra $V = \mathbb{k}\mathbb{1} \oplus \bigoplus_{n \geq 1} V_n$ is said to be of *CFT type*.

1.2. Correlation functions. Let $K = \{k_{ij}\}$ be an integer symmetric $l \times l$ matrix. Denote by $\Phi(K)$ the space of rational functions in variables z_1, \dots, z_l of the form

$$\alpha = p(z_1, \dots, z_l) \prod_{1 \leq i \leq l} z_i^{k_{ii}} \prod_{1 \leq i < j \leq l} (z_i - z_j)^{k_{ij}}, \quad p \in \mathbb{k}[z_1, \dots, z_l]. \quad (1.5)$$

The space $\Phi(K) = \bigoplus_{d \in \mathbb{Z}} \Phi(K)_d$ is graded in the usual sense, so that $\deg z_i = 1$.

Let V be a graded vertex algebra, and let $f : V \rightarrow \mathbb{k}$ be a functional of degree d , i.e. $f(V_n) = 0$ for $n \neq d$. Take some elements $a_1, \dots, a_l \in V$ of degrees d_1, \dots, d_l respectively, formal variables z_1, \dots, z_l and a permutation $\sigma \in S_l$ of $\{1, \dots, l\}$. Consider the formal series

$$\begin{aligned} \alpha_f^\sigma(z_1, \dots, z_l) &= f\left(Y(a_{\sigma(1)}, z_{\sigma(1)}) \cdots Y(a_{\sigma(l)}, z_{\sigma(l)}) \mathbb{1}\right) \\ &= \sum_{m_1, \dots, m_l \in \mathbb{Z}} f\left(a_{\sigma(1)}(m_1) \cdots a_{\sigma(l)}(m_l) \mathbb{1}\right) z_{\sigma(1)}^{-m_1-1} \cdots z_{\sigma(l)}^{-m_l-1}. \end{aligned} \quad (1.6)$$

It can be deduced [8] from the locality identity (1.1) and from the fact that $a(n)\mathbb{1} = 0$ for $n \geq 0$ that α_f^σ is an expansion in the domain $|z_1| > \dots > |z_l|$ of a rational function $\alpha \in \Phi(K)$ for $K = \{k_{ij}\}$ given by $k_{ii} = 0$ and $k_{ij} = -N(a_i, a_j)$, called a *correlation function* of V . Since $\deg a_i(m_i) = d_i - m_i - 1$, we get $\deg \alpha = d - \sum_{i=1}^l d_i$.

Assume that a_1, \dots, a_l are linearly independent. A simultaneous permutation of $\{a_1, \dots, a_l\}$ and $\{z_1, \dots, z_l\}$ would result in a different expansion of the same rational function. This means that α is stable under the action of the group $\Gamma \subset S_l$ generated by all transpositions (i, j) for which $a_i = a_j$.

It is shown in [8] that the rationality property of correlation functions can serve as an equivalent definition of vertex algebras.

1.3. The action of sl_2 . In this paper we need to deal with vertex algebras equipped with certain additional structure.

Definition 1.3. A vertex algebra V is said to have sl_2 -structure, if $V = \bigoplus_{d \in \mathbb{Z}} V_d$ is graded, and there is a locally nilpotent operator $D^* : V \rightarrow V$ of degree -1 , such that $D^* \mathbb{1} = 0$ and

$$[D^*, a(m)] = (2d - m - 2)a(m+1) + (D^*a)(m) \quad (1.7)$$

for every $a \in V_d$.

Let $\delta : V \rightarrow V$ be the grading derivation, defined by $\delta|_{V_d} = d$. It is easy to compute that if $D^* : V \rightarrow V$ satisfies the condition (1.7), then

$$[D^*, D] = 2\delta, \quad [\delta, D] = D, \quad [\delta, D^*] = -D^*,$$

so that D^*, D and δ span a copy of sl_2 .

All vertex algebras in this paper are assumed to have sl_2 -structure, the homomorphisms of vertex algebra preserve the action of sl_2 , etc.

An element $a \in V$ such that $D^*a = 0$ is called *minimal*. It is easy to see that if V is generated by minimal elements, then any operator $D^* : V \rightarrow V$ satisfying (1.7) must be locally nilpotent.

If V has a Virasoro element ω , then we always have $D = \omega(0)$ and $\delta = \omega(1)$, and we can take $D^* = \omega(2)$. Vertex algebras with an action of sl_2 as above were called *quasi-vertex operator algebras* in [8] and minimal elements are sometimes called *quasi-primary*.

We are going to need the following fact:

Lemma 1.1. *If $d < 0$, then $V_d = (D^*)^{1-d} V_1$.*

In fact this is true for any graded sl_2 -module V on which D^* is locally nilpotent and $\delta|_{V_d} = d$ [21]. For vertex algebras the action of sl_2 was also investigated in [7]. When V is a vertex algebra generated by minimal elements, Lemma 1.1 will follow from Corollary 5.1 in §5.2.

1.4. The universal enveloping algebra. For any vertex algebra V we can construct a Lie algebra $L = \text{Coeff } V$ in the following way [3, 13, 18, 19]. Consider the linear space $\mathbb{k}[t, t^{-1}] \otimes V$, where t is a formal variable. Denote $a(n) = a \otimes t^n$ for $n \in \mathbb{Z}$. As a linear space, L the quotient of $\mathbb{k}[t, t^{-1}] \otimes V$ by the subspace spanned by the relations $(Da)(n) = -n a(n-1)$.

The brackets are given by

$$[a(m), b(n)] = \sum_{i \geq 0} \binom{m}{i} (a(i)b)(m+n-i), \quad (1.8)$$

which is precisely the identity (iv) of Definition 1.1. The spaces $L_{\pm} = \text{Span}\{a(n) \mid n \gtrless 0\} \subset L$ are Lie subalgebras of L and we have $L = L_- \oplus L_+$.

Remark. The construction of L makes use of only the products (n) for $n \geq 0$ and the map D . This means that it works for a more generic algebraic structure, known as *conformal algebra* [13].

Now assume that the vertex algebra V has the sl_2 -structure. Then the formulas (1.2) and (1.7) define derivations $D : L \rightarrow L$ and $D^* : L \rightarrow L$ so we get an action of sl_2 on L by derivations. Denote by $\widehat{L} = L \ltimes sl_2$ the corresponding semi-direct product.

The Lie algebra $\widehat{L} = \widehat{\text{Coeff } V}$ and its universal enveloping algebra $U = U(\widehat{L})$ inherit the grading from V so that $\deg a(m) = \deg a - m - 1$. The *Frenkel-Zhu topology* [10] on a homogeneous component U_d is defined by setting the neighborhoods of 0 to be the spaces $U_d^k = \sum_{i \leq k} U_{d-i} U_i$, so that

$$\dots \subset U_d^{k-1} \subset U_d^k \subset U_d^{k+1} \subset \dots \subset U_d, \quad \bigcap_{k \in \mathbb{Z}} U_d^k = 0, \quad \bigcup_{k \in \mathbb{Z}} U_d^k = U_d.$$

Let $\bar{U} = \bigoplus_{d \in \mathbb{Z}} \bar{U}_d$ be the completion of $U(\widehat{L})$ in this topology. Consider the ideal $I \subset \bar{U}$ generated by the relations

$$(a(m)b)(n) = \sum_{s \geq 0} (-1)^s \binom{m}{s} a(m-s)b(n+s) - \sum_{s \leq m} (-1)^s \binom{m}{m-s} b(n+s)a(m-s)$$

for all $a, b \in V$ and $m, n \in \mathbb{Z}$. Note that the relations above are simply the associativity identity (1.4). Denote by $W = \bar{U}/\bar{I}$ the quotient of \bar{U} by the closure of I .

For a finite ordered set of elements $\mathcal{S} = \{a_1, \dots, a_l\}$, $a_i \in V$, let $W_{\mathcal{S}}$ be the $\langle D, \delta, D^* \rangle$ -module generated by all monomials $a_1(m_1) \cdots a_l(m_l) \in W$, $m_i \in \mathbb{Z}$.

Definition 1.4. [10, 21] The universal enveloping algebra of V is

$$U(V) = \bigcup_{\mathcal{S}} \bar{W}_{\mathcal{S}} \subset W,$$

where the union is taken over all finite ordered sets $\mathcal{S} \subset V$, and $\bar{W}_{\mathcal{S}}$ is the completion of the space $W_{\mathcal{S}}$ in the Frenkel-Zhu topology.

Remark. In fact, one can show that if \mathcal{S} and \mathcal{S}' differ by a permutation, then $\bar{W}_{\mathcal{S}} = \bar{W}_{\mathcal{S}'}$.

It is proved in [21] that any module over a vertex algebra V is a continuous module over $U(V)$, in the sense that for any sequence $u_1, u_2, \dots \in U(V)$ that converges to 0 and any $v \in M$ we have $u_i v = 0$ for $i \gg 0$. Conversely, any $U(V)$ -module M , such that for any $a \in V$ and $v \in M$ one has $a(m)v = 0$ for $m \gg 0$, is a module over V .

Remark. The algebra $W = \bar{U}(\hat{L})/\bar{I}$ is also a good candidate for universal enveloping algebra of V . It has the following property [10]: consider a graded space M such that $M_d = 0$ for $d \ll 0$; then M is a W -module if and only if M is an V -module.

On the other hand, we could define an algebra $\hat{U}(V)$ such that any series of elements from $U(\hat{L})$, that make sense as an operator on any V -module, would converge in $\hat{U}(V)$. However, this algebra is too big for our purposes, for example there is no way of defining an involution in this algebra, see § 1.5 below.

1.5. Invariant bilinear forms. The key ingredient of our constructions is the notion of invariant bilinear form on vertex algebra. Here we review the results of [21], that generalize the results of Frenkel, Huang and Lepowsky [8] and Li [17].

Let V be a vertex algebra with an sl_2 -structure, as in § 1.3. It is shown in [21], using some ideas from [8], that there is an anti-involution $u \mapsto u^*$ on the universal enveloping algebra $U(V)$ such that $D \mapsto D^*$, $\delta^* = \delta$ and

$$a(m)^* = (-1)^{\deg a} \sum_{i \geq 0} \frac{1}{i!} ((D^*)^i a) (2 \deg a - m - 2 - i)$$

for a homogeneous $a \in V$ and $m \in \mathbb{Z}$.

Let K be a linear space over \mathbb{k} .

Definition 1.5. [8, 17] A K -valued bilinear form $\langle \cdot | \cdot \rangle$ on V is called *invariant* if

$$\langle a(m)b | c \rangle = \langle b | a(m)^*c \rangle \quad \text{and} \quad \langle Da | b \rangle = \langle a | D^*b \rangle$$

for all $a, b, c \in V$ and $m \in \mathbb{Z}$.

The radical $\text{Rad}\langle \cdot | \cdot \rangle = \{a \in V \mid \langle a | b \rangle = 0 \ \forall b \in V\}$ of an invariant form is an ideal of V . Also, since $\langle \delta a | b \rangle = \langle a | \delta b \rangle$, we have $\langle V_i | V_j \rangle = 0$ for $i \neq j$.

Given a K -valued invariant form $\langle \cdot | \cdot \rangle$ on V , one can consider a functional $f : V_0 \rightarrow K$ defined by $f(a) = \langle \mathbb{1} | a \rangle$. Since $f(D^*a) = \langle \mathbb{1} | D^*a \rangle = \langle D\mathbb{1} | a \rangle = 0$, we get that $f(D^*V_1) = 0$. Also, the form can be reconstructed from f by the formula $\langle a | b \rangle = f(a(-1)^*b)$.

Proposition 1.1. [21] *There is a one-to-one correspondence between invariant K -valued bilinear forms $\langle \cdot | \cdot \rangle$ on a vertex algebra V and linear functionals $f : V_0/D^*V_1 \rightarrow K$, given by $f(a) = \langle \mathbb{1} | a \rangle$, $\langle a | b \rangle = f(a(-1)^*b)$. Moreover, every invariant bilinear form on V is symmetric.*

For the case when V is a vertex operator algebra such that $V_n = 0$ for $n \ll 0$ and $\dim V_n < \infty$ this was proved in [8, 17].

Remark. We observe that a vertex algebra V such that $V_0 = \mathbb{k}\mathbb{1}$ and $D^*V_1 = 0$ is simple if and only if the invariant \mathbb{k} -valued bilinear form on V (which is unique by the above) is non-degenerate. Indeed, any homomorphism $V \rightarrow U$ of vertex algebras must be an isometry, hence its kernel must belong to the radical of the form.

1.6. Radical of a vertex algebra. Let $I = \langle D^*V_1 \rangle \subset V$ be the ideal of a vertex algebra V generated by the space D^*V_1 . Its degree 0 component $I_0 = U(V)_0 D^*V_1$ is spanned by the elements $a_1(m_1) \cdots a_l(m_l) D^*v$ such that $a_i \in V$, $m_i \in \mathbb{Z}$, $\deg a_1(m_1) \cdots a_l(m_l) = 0$ and $v \in V_1$, since we have $DV_{-1} \subset D^*V_1 \subset I_0$, using Lemma 1.1. Note that Lemma 1.1 also implies that $V_d \subset I$ for $d < 0$.

It follows from (1.3) and (1.4) that $K = V_0/I_0$ is the commutative associative algebra with respect to the product $(-)$ with unit $\mathbb{1}$. Let $f : V_0 \rightarrow K$ be the canonical projection. By Proposition 1.1, the map f corresponds to an invariant k -valued bilinear form $\langle \cdot | \cdot \rangle$ on V .

Definition 1.6. The *radical* of V is $\text{Rad } V = \text{Rad} \langle \cdot | \cdot \rangle$.

Remark. This definition has nothing to do with the radical defined in [6].

Denote $\bar{V} = V/\text{Rad } V$. The following proposition summarizes some properties of \bar{V} that we will need later.

Proposition 1.2. [21]

- (a) $\text{Rad}(\bar{V}) = 0$.
- (b) We have $\bar{V} = \bigoplus_{n \geq 0} \bar{V}_n$, so that $\bar{V}_0 = V_0/I_0 = K$, and \bar{V} is a vertex algebra over K .
- (c) Every ideal $J_0 \subset K$ canonically corresponds to an ideal $J \subset \bar{V}$, such that $J \cap \bar{V}_0 = J_0$, in the following way: let $g : K \rightarrow K/J_0$ be the canonical projection, by Proposition 1.1 it defines a K/J_0 -valued invariant bilinear form $\langle \cdot | \cdot \rangle_g$ on \bar{V} , and we set $J = \text{Rad} \langle \cdot | \cdot \rangle_g$.

Note that all the above definitions make sense when \mathbb{k} is a commutative ring containing \mathbb{Q} . So the statement (b) means that \bar{V}_0 is of CFT type as a vertex algebra over K .

We also need the following lemma:

Lemma 1.2. (a) $(u - u^*)\mathbb{1} \in D^*V_1$ for any $u \in U(V)_0$.

(b) Let $u \in U(V)_0$ be such that $u\mathbb{1} \in \text{Rad } V$. Then $uv\mathbb{1} \in \text{Rad } V$ for every $v \in U(V)_0$.

Proof. Statement (a) is proved in [21, §3]. To prove (b), we compute, using (a),

$$uv\mathbb{1} \equiv v^*u^*\mathbb{1} \equiv v^*u\mathbb{1} \equiv 0 \pmod{\text{Rad } V}.$$

□

1.7. Delta-functions. Consider the formal series

$$\delta_k(w, z) = \sum_{i \in \mathbb{Z}} \binom{i}{k} w^{-i-1} z^{i-k}$$

for $k \geq 0$. The series $\delta_k(w, z)$ is the difference between the two expansions of $(w - z)^{-k-1}$ in the power series, so it follows that $(w - z)^n \delta_k(w, z) = 0$ for $n > k$. We will also need the following identity:

$$\delta_0(w, z) g(w, z) = g(z, z) \tag{1.9}$$

for a formal power series $g(w, z)$ provided that either of the sides makes sense.

A very useful fact in the theory of vertex algebras is that the generating function of the commutator formula (1.8) is

$$[Y(a, w), Y(b, z)] = \sum_{n \geq 0} Y(a(n)b, z) \delta_n(w, z). \tag{1.10}$$

1.8. Free vertex algebras. Here we review the construction of free vertex algebras, following [19, 20]. Let \mathcal{A} be an ordered set with a symmetric integer-valued function $N : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{Z}$ such that $N(a, a) \in 2\mathbb{Z}$ for any $a \in \mathcal{A}$. Then there is a vertex algebra $F = F_N(\mathcal{A})$ generated by \mathcal{A} such that the locality of any two generators $a, b \in \mathcal{A} \subset F$ is exactly $N(a, b)$, and the following universality property holds: for any vertex algebra V generated by \mathcal{A} with the same bound on locality there is a vertex algebra homomorphism $F \rightarrow V$ that fixes \mathcal{A} .

The natural way of grading F is by setting $\deg a = N(a, a)/2$ for any $a \in \mathcal{A}$. Obviously, the algebra F is also graded by $\mathbb{Z}_+[\mathcal{A}]$. We will refer to this grading as “grading by weights”.

Denote by $F_{\lambda,d} \subset F$ the space of all elements of degree $d \in \mathbb{Z}$ and weight $\lambda \in \mathbb{Z}_+[\mathcal{A}]$, so that

$$F_{\lambda,i}(n)F_{\mu,j} \subseteq F_{\lambda+\mu,i+j-n-1}$$

One can define the sl_2 -structure on F by $D^*a = 0$ for any $a \in \mathcal{A}$, and then

$$DF_{\lambda,i} \subset F_{\lambda,d+1}, \quad D^*F_{\lambda,i} \subset F_{\lambda,d-1}.$$

A basis of F is given by all words $a_1(m_1) \cdots a_l(m_l)\mathbb{1}$, $a_i \in \mathcal{A}$, $m_i, l \in \mathbb{Z}$, $l \geq 0$, $m_l < 0$, such that

$$m_i - m_{i+1} \leq \begin{cases} \sum_{j=i+1}^l N(a_i, a_j) - \sum_{j=i+2}^l N(a_{i+1}, a_j) & \text{if } a_i \leq a_{i+1}, \\ \sum_{j=i+1}^l N(a_i, a_j) - \sum_{j=i+2}^l N(a_{i+1}, a_j) - 1 & \text{otherwise} \end{cases}$$

for $1 \leq i \leq l-1$.

For the case when $N(a, b) = N \in \mathbb{Z}$ is constant (but $N(a, a)$ is allowed to be $N-1$ if N is odd), this simplifies to

$$m_i - m_{i+1} \leq \begin{cases} N & \text{if } a_i \leq a_{i+1}, \\ N-1 & \text{otherwise} \end{cases} \quad (1.11)$$

For any weight $\lambda = a_1 + \dots + a_l \in \mathbb{Z}_+[\mathcal{A}]$ there is the minimal degree

$$d_{\min}(\lambda) = -\frac{1}{2} \sum_{i,j=1}^l N(a_i, a_j) \in \mathbb{Z}$$

such that $\dim_{\mathbb{K}} F_{\lambda, d_{\min}(\lambda)} = 1$ and $\dim_{\mathbb{K}} F_{\lambda, d} = 0$ for $d < d_{\min}(\lambda)$.

By counting elements in the above basis, we conclude that for $d > d_{\min}(\lambda)$ the dimension of $F_{\lambda, d}$ is equal to the number of partitions of $d - d_{\min}(\lambda)$ into a sum of l non-negative integers colored by a_1, \dots, a_l .

For the case when all a_i 's are different, $\dim F_{\lambda, d}$ is equal to the number of monomials of degree $d - d_{\min}(\lambda)$ in some formal variables z_1, \dots, z_l . In general, let $\Gamma \subset S_l$ be the group of permutations σ of $\{1, \dots, l\}$ such that $a_i = a_{\sigma(i)}$ for any $1 \leq i \leq l$. Then using a combinatorial argument one can show that $\dim F_{\lambda, d}$ is equal to the dimension of the space of Γ -symmetric polynomials in z_1, \dots, z_l of degree $d - d_{\min}(\lambda)$.

Let $K = \{k_{ij}\}$ be defined by $k_{ii} = 0$ and $k_{ij} = -N(a_i, a_j)$ for $i \neq j$. Then by § 1.2, for a functional $f : F_{\lambda, d} \rightarrow \mathbb{K}$ the corresponding correlation function $\alpha_f = f(Y(a_1, z_1) \cdots Y(a_l, z_l)\mathbb{1})$ belongs to $\Phi(K)$.

Proposition 1.3. *The map $f \mapsto \alpha_f$ is a linear isomorphism $F_{\lambda, d}^* \rightarrow \Phi(K)^\Gamma$.*

Here we assume that $f(v) = 0$ for any homogeneous $v \notin F_{\lambda, d}$.

Proof. If $\alpha_f = 0$, then $f = 0$, because $F_{\lambda, d}$ is spanned by the words $a_1(m_1) \cdots a_l(m_l)\mathbb{1}$. Therefore, the map $f \mapsto \alpha_f$ is injective. Since the dimensions of $F_{\lambda, d}$ and $\Phi(K)^\Gamma$ are the same, this map must also be surjective. \square

2. DIAMOND LEMMA AND REWRITING SYSTEMS

Here we give an account of Diamond lemma for associative algebras [1, 2, 19]. This technique also works for other varieties of algebras, e.g. for commutative associative [4] or Lie [22].

Let X be some alphabet. Consider the free associative algebra $\mathbb{K}\langle X \rangle$ of non-commutative polynomials with coefficients in \mathbb{K} . Denote by X^* the set of words in X , i.e. the free semigroup with 1 generated by X .

A *rule* on $\mathbb{K}\langle X \rangle$ is a pair $\rho = w \rightarrow f$, consisting of a word $w \in X^*$ and a polynomial $f \in \mathbb{K}\langle X \rangle$. The left-hand side w is called *the principal part* of rule ρ .

A rule $\rho = w \rightarrow f$ is said to be *applicable* to a word $v \in X^*$ if w is a subword of v , i.e. $v = v'wv''$. The result of application of ρ to v is, naturally, $\rho(v) = v'fv''$. If $p \in \mathbb{K}\langle X \rangle$ is a polynomial which involves a word v , such that a rule ρ is applicable to v , then we say that ρ is *applicable* to p .

Let \mathcal{R} be a set of rules on $\mathbb{k}\langle X \rangle$. For $p, q \in \mathbb{k}\langle X \rangle$ we will write $p \xrightarrow{\mathcal{R}} q$ to denote the fact that q can be obtained as a result of applying the rules from \mathcal{R} to p .

Definition 2.1. Let \mathcal{R} be a set of rules on $\mathbb{k}\langle X \rangle$.

- (a) A word $w \in X^*$ (or a polynomial in $p \in \mathbb{k}\langle X \rangle$) is called \mathcal{R} -terminal if no rule from \mathcal{R} is applicable to w (or to p).
- (b) A set or rules \mathcal{R} is a *rewriting system* on $\mathbb{k}\langle X \rangle$ if there are no infinite sequences of the form

$$p_1 \xrightarrow{\mathcal{R}} p_2 \xrightarrow{\mathcal{R}} \dots, \quad p_i \neq p_{i+1},$$

i.e. any polynomial $p \in \mathbb{k}\langle X \rangle$ can be modified only finitely many times by rules from \mathcal{R} .

- (c) A rewriting system is *confluent* if for any polynomial $p \in \mathbb{k}\langle X \rangle$ there is the unique terminal polynomial $t = \mathcal{R}(p)$ such that $p \xrightarrow{\mathcal{R}} t$.
- (d) Let $\rho = w \rightarrow f, \sigma = u \rightarrow g \in \mathcal{R}$. A word $v \in X^*$ is called an *ambiguity* if either $v = v_1 v_2 v_3$ for some $1 \neq v_1, v_2, v_3 \in X^*$ so that $w = v_1 v_2$ and $u = v_2 v_3$, or if $\rho \neq \sigma$ but $v = w = v_1 u v_2$.

Any rule $\rho = w \rightarrow f \in \mathcal{R}$ can be viewed as a polynomial $w - f \in \mathbb{k}\langle X \rangle$. Denote by $I(\mathcal{R}) \subset \mathbb{k}\langle X \rangle$ the two-sided ideal generated by the set $\{w - f \mid w \rightarrow f \in \mathcal{R}\}$.

Now we can state the Lemma.

Lemma 2.1 (Diamond Lemma). (a) A rewriting system \mathcal{R} is confluent if and only if all terminal words form a basis of $\mathbb{k}\langle X \rangle / I(\mathcal{R})$.
 (b) A rewriting system is confluent if and only if it is confluent on all the ambiguities, that is, for any ambiguity $v \in X^*$ there is the unique \mathcal{R} -terminal polynomial $\mathcal{R}(v) \in \mathbb{k}\langle X \rangle$.

Note that the condition that the terminal words form a basis of $\mathbb{k}\langle X \rangle / I(\mathcal{R})$ is equivalent to the existence of the following membership test: a polynomial $p \in \mathbb{k}\langle X \rangle$ belongs to I if and only if $p \xrightarrow{\mathcal{R}} 0$.

Now assume that the set of words X^* has a linear monomial order. For a polynomial $p = \sum_{w \in X^*} \xi_w w \in \mathbb{k}\langle X \rangle$, $\xi_w \in \mathbb{k}$, denote $\bar{p} = \max\{w \in X^* \mid \xi_w \neq 0\}$ its principal word. Then p corresponds to a rule $\rho_p = \bar{p} \rightarrow -\xi_{\bar{p}}^{-1} \sum_{w \neq \bar{p}} \xi_w w$.

Definition 2.2. Let $I \subset \mathbb{k}\langle X \rangle$ be an ideal generated by some set $R \subset I$, and let $\mathcal{R} = \{\rho_p \mid p \in R\}$. Then R or \mathcal{R} is called a *Gröbner-Shirshov basis* of I if the rules \mathcal{R} form a confluent rewriting system.

Note that unless the order on X^* has some nice properties, it is not automatic that \mathcal{R} is a rewriting system, confluent or not. We also remark that it would be more appropriate to call R a Gröbner-Shirshov set of generators rather than a basis, because there are no independence condition.

3. PROOF OF THEOREM 1

3.1. The algebra \widehat{V} . Let A be a commutative (but not necessarily associative) algebra with an non-degenerate invariant symmetric bilinear form $\langle \cdot \mid \cdot \rangle$, so that $\langle ab \mid c \rangle = \langle a \mid bc \rangle$ for all $a, b, c \in A$. In this section we construct a certain large vertex algebra \widehat{V} such that the desired OZ algebra V will be its homomorphic image.

The algebra \widehat{V} is generated by A with the locality bound $N(a, b) = 4$ for any $a, b \in A$, subject to the following relations:

$$a(1)b = ab, \quad a(2)b = 0, \quad a(3)b = \langle a \mid b \rangle \mathbb{1}. \quad (3.1)$$

Obviously, the group $\text{Aut } A$ acts on \widehat{V} by vertex algebra automorphisms.

A simple computation shows that one can define a derivation $D^* : \widehat{V} \rightarrow \widehat{V}$ of degree -1 such that $D^*a = 0$, and this derivation satisfies all the requirements of § 1.3.

Define a filtration $0 = \widehat{V}^{(-1)} \subset \widehat{V}^{(0)} \subset \widehat{V}^{(1)} \subset \widehat{V}^{(2)} \subset \dots$ on \widehat{V} by setting

$$\widehat{V}^{(l)} = \text{Span}_{\mathbb{k}} \{ a_1(m_1) \cdots a_k(m_k) \mathbb{1} \mid a_i \in A, m_i \in \mathbb{Z}, k \leq l \}.$$

It is easy to see that $\widehat{V}^{(0)} = \mathbb{k}\mathbb{1}$, $\widehat{V}^{(1)} = \mathbb{k}[D]A$ and $\bigcup_{l \geq 0} \widehat{V}^{(l)} = \widehat{V}$. We have $\widehat{V}^{(k)}(n)\widehat{V}^{(l)} \subseteq \widehat{V}^{(k+l)}$, $D\widehat{V}^{(l)} \subset \widehat{V}^{(l)}$ and $D^*\widehat{V}^{(l)} \subset \widehat{V}^{(l)}$, so the associated graded space $\text{gr } \widehat{V} = \bigoplus_{l \geq 0} \widehat{V}^{(l)} / \widehat{V}^{(l-1)}$ is naturally a vertex algebra.

Fix a linearly ordered \mathbb{k} -linear basis \mathcal{A} of A . We claim the following analogue of Poincaré-Birkhoff-Witt theorem:

Theorem 2. *The associated graded vertex algebra $\text{gr } \widehat{V}$ is isomorphic to the free vertex algebra $F = F_N(\mathcal{A})$ generated by the set \mathcal{A} with the locality bound $N : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{Z}$ given by $N(a, b) = 1$ if $a \neq b$ and $N(a, a) = 0$.*

Denote by $F^{(l)}$ the span of all words of length l in F , so that $F^{(l)} = \widehat{V}^{(l)} / \widehat{V}^{(l-1)}$ by Theorem 2.

We note that the relations (3.1) that define \widehat{V} involve only non-negative products between generators, therefore by § 1.4 the universal enveloping algebra $U(\widehat{V})$ is a completion in the Frenkel-Zhu topology of the enveloping algebra of the Lie algebra generated by the set $X = \{a(m) \mid a \in \mathcal{A}, m \in \mathbb{Z}\}$ and relations coming from (1.8):

$$[a(m), b(n)] = [a(m-1), b(n+1)] + (ab)(m+n-1) + \binom{m-1}{2} \langle a \mid b \rangle \delta_{m+n,2}. \quad (3.2a)$$

For $a = b$ the quasi-symmetry (1.3) implies that $a(0)a = \frac{1}{2}Da^2$, so (3.2a) becomes

$$[a(m), a(n)] = \frac{1}{2}a^2(m+n-1) + \binom{m}{3} \langle a \mid a \rangle \delta_{m+n,2}. \quad (3.2b)$$

Using delta functions (see § 1.7) and specifically formula (1.10), we compute the generating function of the identities (3.2):

$$(w-z)[Y(a, w), Y(b, z)] = Y(ab, z) \delta_0(w, z) + \langle a \mid b \rangle \delta_2(w, z), \quad (3.3a)$$

which for the case when $a = b$ specializes to

$$[Y(a, w), Y(a, z)] = \frac{1}{2} \partial_z Y(a^2, z) \delta_0(w, z) + Y(a^2, z) \delta_1(w, z) + \langle a \mid a \rangle \delta_3(w, z). \quad (3.3b)$$

The reason why Theorem 2 holds is that the relations (3.2) are in fact a Gröbner-Shirshov basis (see § 2) of the ideal generated by them. Let us introduce a linear order on words X^* . We set $a(m) < b(n)$ if either $m < n$ or $m = n$ and $a < b$. Then we will compare words from X^* first by length and then lexicographically. Then relations (3.2) correspond to rewriting system \mathcal{R} (see § 2) that consists of the rules

$$\begin{aligned} a(m)b(n) &\longrightarrow b(n)a(m) + [a(m-1), b(n+1)] \\ &\quad + (ab)(m+n-1) + \delta_{m+n,2} \binom{m-1}{2} \langle a \mid b \rangle, \end{aligned} \quad (3.4a)$$

where $m > n$ if $a > b$ and $m > n+1$ if $a \leq b$, and also

$$a(m+1)a(m) \longrightarrow a(m)a(m+1) + \frac{1}{2}a^2(2m). \quad (3.4b)$$

By rather lengthy calculations, similar to those in [19], one can show that \mathcal{R} is confluent. If $ab = \langle a \mid b \rangle = 0$, then the relations (3.2) define the universal enveloping algebra of the free algebra $F = F_N(\mathcal{A})$, and in [19] it was shown that the corresponding rewriting system is confluent. Then the Diamond Lemma 2.1 states that the set \mathcal{T} of \mathcal{R} -terminal words forms a basis of both \widehat{V} and of F . These terminal words are given by (1.11), which in this case gives the set of words $a_1(m_1)a_2(m_2) \cdots a_l(m_l)$ such that

$$m_i - m_{i+1} \leq \begin{cases} 1 & \text{if } a_i < a_{i+1} \\ 0 & \text{otherwise,} \end{cases} \quad \text{for } 1 \leq i \leq l-1. \quad (3.5)$$

Therefore a basis of \widehat{V} is given by all expressions $a_1(m_1)a_2(m_2) \cdots a_l(m_l)\mathbb{1}$ such that in addition to (3.5) one has $m_l < 0$.

3.2. Order on words.

We start with introducing a linear order on the set of words $w = a_1(m_1) \dots a_l(m_l)$ of degree 0, where $a_i \in \mathcal{A}$ and $m_i \in \mathbb{Z}$. We will write $|w| = l$. We will call such a word *indecomposable* if it does not have a proper tail $a_k(m_k) \dots a_l(m_l)$, $1 \leq k < l$, of degree 0.

Assume that w is an indecomposable word. Define $\tau(w)$ by

$$\tau(w) = \min\{\deg v \mid v \text{ is a proper tail of } w\}$$

if w has a proper tail of either negative degree or of degree 1, and set $\tau(w) = \infty$ otherwise. Set also $\tau(a(1)) = 0$ for $a \in \mathcal{A}$, these will be the only indecomposable words w with $\tau(w) = 0$. For a decomposable word w , let $\tau(w)$ be the minimal of $\tau(w_i)$ for w_i running over the indecomposable parts of w .

We fix a linear monomial order \prec on the set of degree 0 words as above, such that the following conditions hold:

- If $\tau(u) < \tau(v)$, then $u \prec v$.
- If u and v are indecomposable words, such that $\tau(u) = \tau(v)$, then $u \prec v$ if and only if $u < v$, the later being the order introduced in §3.1.
- Let $u_1 \prec u_2 \prec \dots$ and $v_1 \prec v_2 \prec \dots$ be indecomposable terms of the words u and v respectively. Then if $u_1 = v_1, \dots, u_i = v_i$ but $u_{i+1} \prec v_{i+1}$, then $u \prec v$.

Note that these conditions do not define the order \prec uniquely.

We will need the following easy property of the rules \mathcal{R} .

Lemma 3.1. *Let $w \xrightarrow{\mathcal{R}} \sum_{v \in \mathcal{B}} k_v v$. Then $\tau(v) \leq \tau(w)$ whenever $k_v \neq 0$.*

Recall that \mathcal{T} is the set of the words $a_1(m_1) \dots a_l(m_l) \in X^*$ such that the condition (3.5) holds. These are the terminal words with respect to the rules \mathcal{R} given by (3.4). Set $\mathcal{T}_d = \mathcal{T} \cap \widehat{V}_d$ and

$$\mathcal{T}_0^+ = \{a_1(m_1) \dots a_l(m_l) \in \mathcal{T}_0 \mid m_l < 0\}.$$

By the results of §3.1, the set $\{w\mathbb{1} \mid w \in \mathcal{T}_0^+\}$ is a basis of \widehat{V}_0 . Let $S \subset \mathcal{T}_0$ be the set of all indecomposable words of degree 0. We will deal with rules that act on words in S . We can interpret any element $u \in \mathcal{T}_0$ as a word in the alphabet S . We will identify \widehat{V}_0 with a subspace in $\mathbb{k}\langle S \rangle$, so that an element $v = \sum_{w \in \mathcal{T}_0^+} k_w w\mathbb{1} \in \widehat{V}_0$, $k_w \in \mathbb{k}$, is the same as $\sum_{w \in \mathcal{T}_0^+} k_w w \in \mathbb{k}\langle S \rangle$. The order \prec is also an order on S^* . As in §2, any element $v = \sum_{u \in S^*} k_u u\mathbb{1} \in \widehat{V}_0$ defines a rule $\rho_v = \bar{v} \longrightarrow -k_{\bar{v}}^{-1} \sum_{u \neq \bar{v}} k_u u$ on S^* , where $\bar{v} = \max\{u \mid k_u \neq 0\}$.

Let $S_1 = \{s \in S \mid \tau(s) \leq 1\}$. We have $\widehat{V}_0 = Q_{\leq 1} \oplus Q_{>1}$, where

$$Q_{\leq 1} = \text{Span}\{w\mathbb{1} \mid w \in \mathcal{T}_0, \tau(w) \leq 1\} \quad \text{and} \quad Q_{>1} = \text{Span}\{w\mathbb{1} \mid w \in \mathcal{T}_0, \tau(w) > 1\}.$$

Lemma 3.1 implies that $Q_{\leq 1} = \text{Span}\{w\mathbb{1} \mid w \in X^*, \tau(w) \leq 1\}$.

3.3. Extended radical of \widehat{V} . Consider the space $D^*\widehat{V}_1 \subset \widehat{V}_0$. It inherits the filtration $(D^*\widehat{V}_1)^{(0)} \subset (D^*\widehat{V}_1)^{(1)} \subset (D^*\widehat{V}_1)^{(2)} \subset \dots \subset D^*\widehat{V}_1$, where $(D^*\widehat{V}_1)^{(l)} = D^*\widehat{V}_1 \cap \widehat{V}_0^{(l)}$, so that by Theorem 2 we have $\text{gr } D^*\widehat{V}_1 = \bigoplus_l (D^*\widehat{V}_1)^{(l)} / (D^*\widehat{V}_1)^{(l-1)} \subset F_0$. In §5 we are going to prove the following strengthening of Theorem 2:

Proposition 3.1. $\text{gr } D^*\widehat{V}_1 = D^*F_1 \subset F_0$.

Remark. In fact, it will follow from the results of §6 that the associated graded vertex algebra of $\text{Rad } \widehat{V}$ is isomorphic to $\text{Rad } F$.

In the free algebra F the unit $\mathbb{1} \in F_0$ is not in D^*F_1 , since $F_1^{(l)} = 0$ for $l \leq 5$, so we get the following important corollary

Corollary 3.1. $\mathbb{1} \notin D^*\widehat{V}_1$.

Recall from §1.6 that $\text{Rad}_0 = U(\widehat{V})_0 D^*\widehat{V}_1$ is the degree 0 component of $\text{Rad } \widehat{V}$. Define an extension $N \subset \widehat{V}_0$ of the radical by $N = \text{Rad}_0 + Q_{\leq 1}$, and set $\bar{N} = \{u \in \mathbb{k}\langle S \rangle \mid u\mathbb{1} \in N\}$. Using Lemma 1.2(b), it is easy to see that $\bar{N} \subset \mathbb{k}\langle S \rangle$ is an ideal generated by $D^*\widehat{V}_1$ and S_1 .

We are going to construct a Gröbner-Shirshov basis of N . It will consist of the following rules:

$$\begin{aligned}\mathcal{H} &= \left\{ \rho_v \left| \begin{array}{l} v = D^* a_1(m_1) \cdots a_l(m_l) \mathbb{1} \in \widehat{V}_0, \ a_i \in \mathcal{A}, \ m_i \in \mathbb{Z} \\ \bar{v} \in S \text{ is an indecomposable word of length } l, \ \tau(\bar{v}) > 1 \end{array} \right. \right\}, \\ \mathcal{C} &= \{ s_1 s_2 \longrightarrow s_2 s_1 \mid s_1 \succ s_2 \in S \}, \\ \mathcal{S}_1 &= \{ s \longrightarrow 0 \mid \tau(s) \leq 1 \}.\end{aligned}$$

Denote $\mathcal{N} = \mathcal{H} \cup \mathcal{C} \cup \mathcal{S}_1$. The rules \mathcal{N} have the following important property: they all are of the form $s \xrightarrow{x}$, for $s \in S$.

Proposition 3.2. \mathcal{N} is a Gröbner-Shirshov basis of \bar{N} .

We are going to split the proof of Proposition 3.2 into the following lemmas:

Lemma 3.2. $I(\mathcal{N}) \subset N$.

Lemma 3.3. \mathcal{N} is a rewriting system.

Lemma 3.4. The rules \mathcal{N} are applicable to every non-zero element $x \in \bar{N}$.

Proof of Proposition 3.2. By Lemma 3.2 we get $I(\mathcal{N}) \subset \bar{N}$. On the other hand, by Lemma 3.4, an element $0 \neq x \in \bar{N}$ cannot be \mathcal{N} -terminal, yet by Lemma 3.3 every element of \bar{N} is reduced to a terminal by finitely many steps. It follows that $x \xrightarrow{\mathcal{N}} 0$, which by Lemma 2.1 (a) means that \mathcal{N} is a Gröbner-Shirshov basis of \bar{N} . \square

Proof of Lemma 3.2. We have $I(\mathcal{H}) \subset D^* \widehat{V}_1 \subset N$ by definition. Using Lemma 1.2, it is easy to see that $uv \equiv u^*v \pmod{\text{Rad}_0}$ for any $u \in U(\widehat{V})_0$ and $v \in \widehat{V}_0$. Therefore, for any $u_1, u_2 \in U(\widehat{V})_0$ and $v \in \widehat{V}_0$ we get

$$u_1 u_2 v \mathbb{1} - u_2 u_1 v \mathbb{1} \equiv u_1 u_2 v \mathbb{1} - u_2^* u_1^* v \mathbb{1} \equiv 0 \pmod{\text{Rad}}.$$

This implies that $I(\mathcal{C}) \subset \text{Rad}_0$.

We are left to show that $S_1 \subset N$. Let $w \in S_1$. If $\tau(w) = 1$, then $w \in \bar{N}$ by definition. If $\tau(w) = 0$, then $w = a(1)$ for some $a \in \mathcal{A}$, and then $w \mathbb{1} = 0 \in N$. Finally, assume that $\tau(w) < 0$. Then $w = w_1 w_2$, so that $\deg w_1 = -\deg w_2 = d > 0$. By Lemma 1.1 there is $v \in \widehat{V}_1$, such that $w_2 \mathbb{1} = (D^*)^{1+d} v$. So we get

$$w_1 w_2 \mathbb{1} = w_1 (D^*)^{1+d} v = (D^*)^d w_1 D^* v - [(D^*)^d, w_1] D^* v \in \text{Rad}_0 \subset N.$$

\square

Proof of Lemma 3.3. Let $w = a_1(m_1) \cdots a_l(m_l) \in S^*$ be a word. We will prove by induction on its length $l = |w|$ as a word in the alphabet $X = \{a(m) \mid a \in \mathcal{A}, m \in \mathbb{Z}\}$ that there is no infinite sequence of rule applications $w \xrightarrow{\mathcal{N}} x_1 \xrightarrow{\mathcal{N}} x_2 \xrightarrow{\mathcal{N}} \dots$.

If $v \longrightarrow x \in \mathcal{H}$ for some $v \in \widehat{V}_0^{(k)}$, then by Proposition 3.1 we have $v \pmod{\widehat{V}_0^{(k-1)}} \in D^* F_0^{(k)}$, and therefore $k \geq 6$, since $D^* F_0^{(k)} = 0$ for $k \leq 5$. Therefore, if $l < 6$, the rules \mathcal{H} are not applicable to w .

Now assume that $w \longrightarrow x \in \mathcal{H}$. Then any word of length l that appears in x must be of the form $a_{i_1}(n_1) \cdots a_{i_l}(n_l) \in S^*$, where (i_1, \dots, i_l) is a permutation of $(1, \dots, l)$ and $\sum n_i = \sum m_i = -2l$. Now the lemma follows from the observation that there are only finitely many such words that do not have a tail of negative degree; and if a word u has a tail of negative degree, then the only way of applying \mathcal{N} to u is $u \xrightarrow{S_1} 0$. \square

For the proof of Lemma 3.4 we will have to use two additional statements. Let $N^{(l)} = N \cap \widehat{V}^{(l)}$ and $Q_{\leq 1}^{(l)} = Q_{\leq 1}^{(l)} \cap \widehat{V}^{(l)}$. Note that $Q_{\leq 1}^{(l)} = N^{(l)} = 0$ for $l \leq 5$. First of all, we need

Proposition 3.3. $N^{(l)} = (D^* \widehat{V}_1)^{(l)} + Q_{\leq 1}^{(l)}$.

In order to formulate the second statement, we need to introduce some auxiliary constructions. Recall from § 3.1 that F is the free vertex algebra generated by the set $\mathcal{A} \subset F_2$ with respect to

locality function $N : \mathcal{A} \times \mathcal{A} \rightarrow \mathbb{Z}_+$ given by $N(a, b) = 1$ for $a \neq b$ and $N(a, a) = 0$ (see §1.8). Let X_0^* be the set of all words $w \in X^*$ if $\deg w = 0$. We define the following subspaces of F_0 :

$$\begin{aligned} E_0 &= \text{Span} \{ uv \mid 1 \neq u \in X_0^*, 1 \neq v \in F_0 \}, \\ E_1 &= \text{Span} \{ u1 \mid u \in X_0^*, \tau(u) \leq 1 \}, \\ P &= \text{Span} \{ uD^*v \mid 1 \neq u \in X_0^*, v \in F_1 \}, \\ C &= \text{Span} \{ u_1[u_2, u_3]u_41 \mid u_1, u_2, u_3, u_4 \in X_0^* \}. \end{aligned}$$

Note that in the special case when the product on form on A are not 0, we have $F = \widehat{V}$ and $Q_{\leq 1} = E_1$. So Proposition 3.3 implies that $\text{Rad}_0 F + E_1 = D^*F_1 + E_1$, in particular,

$$P \equiv \text{Span} \{ ur \mid 1 \neq u \in U(F)_0, r \in \text{Rad}_0 F \} \pmod{E_1}.$$

We are interested in the space $(E_0 + E_1) \cap (D^*F_1 + E_1) \subset F_0/E_1$. Obviously, $P, C \subset E_0$ and $P \subset \text{Rad}_0 F \subset F^*F_1 + E_1$. By the argument similar to that in the proof of Lemma 3.2, we have also $C \subset F^*F_1 + E_1$. We claim that P and C span the whole subspace $(E_0 + E_1) \cap (D^*F_1 + E_1)$ modulo E_1 :

Proposition 3.4. $(E_0 + E_1) \cap (D^*F_1 + E_1) = P + C + E_1 \subset F_0$.

We will prove Proposition 3.3 in §6.2 and Proposition 3.4 in §6.4.

Proof of Lemma 3.4. Let $0 \neq x \in N$. By Proposition 3.3 we can write $x = y + y'$, where $y \in D^*\widehat{V}_1$ and $y' \in Q_{\leq 1}$. Let $y \in (D^*\widehat{V}_1)^{(l)} \setminus (D^*\widehat{V}_1)^{(l-1)}$, and assume that the decomposition $x = y + y'$ gives the minimal possible value of l . Let $y = \sum_{w \in \mathcal{T}_0^+} k_w w$, $k_w \in \mathbb{k}$, and let $y_0 = \sum_{w \in \mathcal{T}_0^+, |w|=l} k_w w$ be the length l component of y . We are going to prove the Lemma by induction on l . As we have seen before, if $l \leq 5$, then $x = 0$.

First we claim that $y_0 \notin Q_{\leq 1}$. Indeed, otherwise $x = (y - y_0) + (y' + y_0) \in N^{(l-1)} + Q_{\leq 1}$ so by Proposition 3.3 we would have $x \in (D^*\widehat{V}_1)^{(l-1)} + Q_{\leq 1}$ which contradicts to the minimality of l .

So we have shown that the set $\Omega = \{ w \in \mathcal{T}_0^+ \mid k_w \neq 0, |w| = l, \tau(w) > 1 \}$ is not empty. Consider the following two cases:

Case 1. The set Ω contains an indecomposable word.

Case 2. All words in Ω are decomposable.

In Case 1, let $v \in \mathcal{T}_0^+$ be the maximal indecomposable word in Ω with respect to the order \prec . Then $v = \bar{y}$ and there is a rule $\rho = v \rightarrow \dots \in \mathcal{H}$ which is applicable to y , and therefore to x , since by Lemma 3.1, the word $v1$ cannot appear in the expansion of $y' \in Q_{\leq 1}$ in the basis $\{ w1 \mid w \in \mathcal{T}_0^+ \}$.

Now assume that the Case 2 takes place. By Theorem 2 there is a projection $\pi : \widehat{V}_0^{(l)} \rightarrow F_0^{(l)}$ such that $\text{Ker } \pi = \widehat{V}_0^{(l-1)}$. Since all the words w that appear in y_0 are either decomposable or have $\tau(w) \leq 1$, we get $\pi(y_0) \in E_0 + E_1$. On the other hand, since $y \in D^*\widehat{V}_1$, we obtain that $\pi(y_0) = \pi(y) \in D^*F_1$, and therefore that $\pi(y_0) \in (E_0 + E_1) \cap (D^*F_1 + E_1)$. So by Proposition 3.4 the element $\pi(y_0)$ can be decomposed as $\pi(y_0) = p + c + e_1$ so that $p \in P$, $c \in C$, $e_1 \in E_1$, and $p \notin C + E_1$, $c \notin E_1$.

Assume first that $p \neq 0$. Then p can be written as $p = \sum_i u_i r_i$, where $u_i \in S$, $|u_i| > 0$ are pairwise different indecomposable \mathcal{R} -terminal words, and $r_i \in \text{Rad}_0 F$. We can assume that $u_i \notin S_1$, since otherwise $u_i r_i \in E_1$ can be removed from the expansion of p .

By induction, there is a rule $\rho \in \mathcal{N}$ which is applicable to the leading word \bar{r}_i of r_i . If $\rho \in \mathcal{S}_1$, then $r_i \in E_1$, and then $u_i r_i \in E_1$, so we can assume that $u_i r_i$ does not appear in the expansion of p . We can also assume that $\rho \notin \mathcal{C}$, because otherwise we can replace r_i by the result $\rho(r_i)$ of applying ρ to r_i ; the difference $r_i - \rho(r_i)$ is in $C \subset \text{Rad}_0 F$ and it needs to be added to c to compensate for this change.

So we get that a rule $\rho = s \rightarrow \dots \in \mathcal{H}$, $s = \bar{\rho} \in S$, is applicable to $u_i \bar{r}_i$. We will show that ρ is applicable to $\pi(y_0)$ as well. We see that the word $u_i \bar{r}_i$ is \mathcal{R} -terminal, and appears with a non-zero coefficient in the expansion of p in the basis $\{ w1 \mid w \in \mathcal{T}_0^+ \}$ of F_0 . Therefore, there is a word w that differs from $u_i \bar{r}_i$ by an permutation of indecomposable terms, that appears with non-zero coefficient

in the expansion of $p + c$. But then the same word w must appear with non-zero coefficient in the expansion of $\pi(y_0)$, because all the words that appear in the expansion of e_1 are from S_1 and $w \notin S_1$.

It remains to consider the case when $p = 0$. But then $c \neq 0$, since we have assumed that $y_0 \notin Q_{\leq 1}$, and the rules \mathcal{C} apply to c and therefore to $\pi(y_0)$.

So we have proved that the rules \mathcal{N} apply to $\pi(y_0)$. This means that they are also applicable to y_0 , since they have the same expansions in the basis $\{w\mathbb{1} \mid w \in \mathcal{T}_0^+\}$, therefore \mathcal{N} are applicable to y and x as well. \square

3.4. The construction of V . In this section we assume that the form $\langle \cdot | \cdot \rangle_A$ on A is non-degenerate. We start with the following observation.

Lemma 3.5. *Any invariant bilinear form $\langle \cdot | \cdot \rangle_{\widehat{V}}$ on \widehat{V} , such that $\langle \mathbb{1} | \mathbb{1} \rangle_{\widehat{V}} = 1$, agrees with the form on A .*

Proof. For any $a, b \in A$ we have, using that $a(-1)^* = a(3)$,

$$\langle a | b \rangle_{\widehat{V}} = \langle a(-1)\mathbb{1} | b \rangle_{\widehat{V}} = \langle \mathbb{1} | a(3)b \rangle_{\widehat{V}} = \langle a | b \rangle_A \langle \mathbb{1} | \mathbb{1} \rangle_{\widehat{V}} = \langle a | b \rangle_A.$$

\square

We will also need the following lemma, which we will prove in §6.5. Recall that $F_d^{(l)}$ denotes the span of all words of length l in F . For a space $E \subseteq F_d$ set $E^{(l)} = E \cap F_d^{(l)}$.

Lemma 3.6. *If $\dim A \geq 2$ and $l \geq 8$, then $E_0^{(l)} + E_1^{(l)} + D^*F_1^{(l)} \subsetneq F_0^{(l)}$.*

By Proposition 1.2 there is an ideal $J \subset \widehat{V}$ such that $J \cap \widehat{V}_0 = N$. Denote $\overline{V} = \widehat{V}/J$ and $K = \widehat{V}_0/N$. The following theorem summarizes some properties of \overline{V} .

Theorem 3. (a) $\overline{V}_d = 0$ for $d < 0$ and for $d = 1$.
 (b) If $\dim A > 1$, the algebra $\overline{V}_0 = K$ is isomorphic to a polynomial algebra in infinitely many variables and \overline{V} is a vertex algebra over K . If $\dim A = 1$, then $K = \mathbb{k}$.
 (c) $A \subset \overline{V}_2$.
 (d) There are no non-zero ideals $I \subset \overline{V}$ such that $I \cap \overline{V}_0 = 0$.
 (e) The vertex algebra \overline{V} depends canonically on A , i.e. the group $\text{Aut } A$ acts on \overline{V} by vertex algebra automorphisms.

Proof. (a) If $d < 0$, then by Lemma 1.1 we have $V_d \subset \langle D^*\widehat{V}_1 \rangle \subset J$. Since $U(\widehat{V})_{-1}v \in \langle S_1 \rangle \in N \subset J$, for any $v \in \widehat{V}_1$, we get $\widehat{V}_1 \subset N$. This proves (a).

(b) Combining Proposition 3.2 with Lemma 2.1, we see that the set of all \mathcal{N} -reduced words in S^* form a basis of K . We note that the rules \mathcal{H} and \mathcal{S}_1 are all of the form $s \xrightarrow{x}$, where $s \in S$, and \mathcal{C} are the commutativity relations. This implies that K is a polynomial algebra generated by the set $S_{\mathcal{N}} \subset S$ of all \mathcal{N} -terminal words in S .

It follows from Lemma 3.6 that if $\dim A > 1$ and $l \geq 8$, then the set $S_{\mathcal{N}}^{(l)} \subset S_{\mathcal{N}}$ of \mathcal{N} -terminal words in S of length l is non-empty. Indeed, let $\phi : \widehat{V}_0^{(l)} \rightarrow F_0^{(l)}$ be the canonical projection, see Theorem 2. Then Proposition 3.1 implies that $\phi(N) \subset E_1^{(l)} + D^*F_1^{(l)}$. Also, if $w \in S^*$ is a decomposable word of length l , then $\phi(w\mathbb{1}) \in E_0^{(l)}$. Therefore, the set $\phi(S_{\mathcal{N}}^{(l)})$ spans $F_0^{(l)}$ modulo $E_0^{(l)} + E_1^{(l)} + D^*F_1^{(l)}$, and we apply Lemma 3.6.

(c) We have $\overline{V} = \widehat{V}/\text{Rad}\langle \cdot | \cdot \rangle$, where $\langle \cdot | \cdot \rangle$ is a canonical invariant K -valued bilinear form corresponding to the map $\widehat{V}_0 \rightarrow \widehat{V}_0/N$, see §1.6. By Lemma 3.5, this form $\langle \cdot | \cdot \rangle$ coincides with the bilinear form $\langle \cdot | \cdot \rangle_A$ on A . Since $\langle \cdot | \cdot \rangle_A$ is non-degenerate, we get $\text{Rad}\langle \cdot | \cdot \rangle \cap A = 0$.

(d) holds since any ideal $I \subset \widehat{V}$ such that $I \cap \widehat{V}_0 \subseteq N$ must belong to $\text{Rad}\langle \cdot | \cdot \rangle$.

(e) is clear, since \widehat{V} depends canonically on A and so does $N \subset \widehat{V}_0$. \square

Proof of Theorem 1. Let $\chi : K \rightarrow \mathbb{k}$ be a character of K . By Proposition 1.1 it defines an invariant \mathbb{k} -bilinear form $\langle \cdot | \cdot \rangle_\chi$ on \bar{V} . Set $V = \bar{V} / \text{Rad} \langle \cdot | \cdot \rangle_\chi$. Since χ is a character, we have $\text{Ker } \chi \subset \text{Rad} \langle \cdot | \cdot \rangle_\chi$, and therefore $V_0 = \mathbb{k}\mathbb{1}$. Theorem 3 (a) implies that V is an OZ vertex algebra. As in the proof of Theorem 3 (c), we deduce from Lemma 3.5 that $A \cap \text{Rad} \langle \cdot | \cdot \rangle_\chi = 0$, therefore $A \subseteq V_2$.

Remark. Let $G \subset \text{Aut } A$ be a group of automorphisms of A . By Theorem 3 (e), G acts in \bar{V} . If we can choose the character $\chi : K = \bar{V}_0 \rightarrow k$ to be G -invariant, then G will act on V as well.

4. CORRELATION FUNCTIONS ON \hat{V}

In order to prove all these numerous statements that are left unproven in § 3.3 and § 3.4, we need to develop some techniques, that utilize the notion of correlation function, see § 1.2.

4.1. Take some elements $a_1, \dots, a_l \in \mathcal{A}$. Denote by $W_d \subseteq \hat{V}^{(l)}$ the span of words $a_1(m_1) \cdots a_l(m_l) \mathbb{1} \in \hat{V}$ of degree d for all $m_i \in \mathbb{Z}$. Denote also $W'_d = W_d \cap \hat{V}^{(l-1)}$. By Theorem 2 we have $W_d / W'_d \cong F_{\lambda, d}$, where $\lambda = a_1 + \dots + a_l \in \mathbb{Z}_+[\mathcal{A}]$.

Denote by $\Gamma \subset S_l$ the group of permutations generated by all transpositions (i, j) for which $a_i = a_j$. Let $\Phi = \Phi(K)$ where $K = \{k_{ij}\}$ is given by $k_{ii} = 0$ and $k_{ij} = -4$ for $i \neq j$. Take some formal variables z_1, \dots, z_l . Recall from § 1.2 that for any functional $f : W_0 \rightarrow \mathbb{k}$ and a permutation $\sigma \in S_l$ of $\{1, \dots, l\}$ the power series $\alpha_f^\sigma(z_1, \dots, z_l)$ given by (1.6) converge in the domain $|z_{\sigma(1)}| > \dots > |z_{\sigma(l)}|$ to a Γ -symmetric rational function $\alpha_f \in \Phi_{-2l}^\Gamma$, which does not depend on the permutation σ .

The following lemma gives more information about the poles of α_f .

Lemma 4.1. *For every $1 \leq i < j \leq l$ the function $\alpha = \alpha_f$ can be expanded*

$$\alpha = \sum_{k \geq -4} (z_i - z_j)^k \alpha_k(z_1, \dots, \widehat{z_i}, \dots, z_l), \quad (4.1)$$

where $\alpha_{-3} = 0$ and

$$\begin{aligned} \alpha_{-4} &= \langle a_i | a_j \rangle f \left(Y(a_1, z_1) \cdots \widehat{Y(a_i, z_i)} \cdots \widehat{Y(a_j, z_j)} \cdots Y(a_l, z_l) \mathbb{1} \right), \\ \alpha_{-2} &= f \left(Y(a_i a_j, z_j) Y(a_1, z_1) \cdots \widehat{Y(a_i, z_i)} \cdots \widehat{Y(a_j, z_j)} \cdots Y(a_l, z_l) \mathbb{1} \right). \end{aligned}$$

If in addition $a_i = a_j$, then

$$\alpha_{-1} = \frac{1}{2} \partial_{z_j} f \left(Y(a_j^2, z_j) Y(a_1, z_1) \cdots \widehat{Y(a_i, z_i)} \cdots \widehat{Y(a_j, z_j)} \cdots Y(a_l, z_l) \mathbb{1} \right).$$

Here the hat $\widehat{}$ over a term means that the term is omitted. In particular, α_{-4} does not depend on both z_i and z_j .

Proof. Since $N(a_i, a_j) = 4$, the order of pole of α at $z_i - z_j$ is at most 4, see § 1.2. We show now that the coefficients α_k have the desired form.

Let $\sigma \in S_l$ be a permutation such that $\sigma(1) = i$ and $\sigma(2) = j$. Look at the difference $\alpha^\sigma - \alpha^{(1,2)\sigma}$ between the two power series expansions of the rational function α . On one hand, from (4.1) we get

$$(z_i - z_j)(\alpha^\sigma - \alpha^{(1,2)\sigma}) = \delta_2(z_i, z_j) \alpha_{-4} + \delta_1(z_i, z_j) \alpha_{-3} + \delta_0(z_i, z_j) \alpha_{-2}.$$

On the other hand, using (3.3a), we get

$$\begin{aligned} (z_i - z_j)(\alpha^\sigma - \alpha^{(1,2)\sigma}) &= (z_i - z_j) f \left([Y(a_i)(z_i), Y(a_j)(z_j)] Y(a_{\sigma(3)})(z_{\sigma(3)}) \cdots Y(a_{\sigma(l)})(z_{\sigma(l)}) \mathbb{1} \right) \\ &= f \left(Y(a_i a_j)(z_j) Y(a_{\sigma(3)})(z_{\sigma(3)}) \cdots Y(a_{\sigma(l)})(z_{\sigma(l)}) \mathbb{1} \right) \delta_0(z_i, z_j) \\ &\quad + \langle a_i | a_j \rangle f \left(Y(a_{\sigma(3)})(z_{\sigma(3)}) \cdots Y(a_{\sigma(l)})(z_{\sigma(l)}) \mathbb{1} \right) \delta_2(z_i, z_j), \end{aligned}$$

Comparing the coefficients with $\delta_k(z_i, z_j)$ gives the statements about α_{-4} , α_{-3} and α_{-2} . If $a_i = a_j$, then we obtain α_{-1} in the same way by taking $\alpha^\sigma - \alpha^{(1,2)\sigma}$ instead of $(z_i - z_j)(\alpha^\sigma - \alpha^{(1,2)\sigma})$ and using (3.3b) instead of (3.3a). □

Let $\Phi^s \subset \Phi$ the space of all rational functions in Φ that may have only simple poles in $z_i - z_j$.

Corollary 4.1. *For a functional $f : W_0 \rightarrow \mathbb{k}$ the condition $f(W'_0) = 0$ holds if and only if $\alpha_f \in (\Phi^s)^\Gamma$.*

Proof. By Lemma 4.1, the correlation function α_f does not have multiple poles if and only if $f(w) = 0$ for all words $w \in W'_0$ of the form

$$(a_i a_j)(m_j) a_1(m_1) \cdots \widehat{a_i(m_i)} \cdots \widehat{a_j(m_j)} \cdots a_l(m_l) \mathbb{1} \quad \text{or} \quad (4.2)$$

$$a_1(m_1) \cdots \widehat{a_i(m_i)} \cdots \widehat{a_j(m_j)} \cdots a_l(m_l) \mathbb{1}.$$

Since these words span the whole W'_0 , we get $f(W'_0) = 0$. \square

Remark. Alternatively, we could argue as follows. If $f(W') = 0$, then f can be viewed as a functional on F_λ , and hence by §1.8 we get $\alpha_f \in (\Phi^s)^\Gamma$. In the other direction, if $\alpha \in \Phi^s$, then by §1.8 there is a functional $g : F_\lambda \rightarrow \mathbb{k}$ such that its correlation function on F_λ is α . But the lifting of g to a functional on W must coincide with f , since a functional is uniquely defined by its correlation function. Note that we are making essential use of Theorem 2 here.

4.2. The operator Δ . Consider the differential operator $\Delta_z(n) = z^2 \partial_z - nz$. It is easy to compute using (1.7) that for any minimal element a of degree d in a vertex algebra one has $[D^*, Y(a, z)] = \Delta_z(-2d)Y(a, z)$. The most important case for us is when $n = -4$. Denote $\Delta = \Delta_{z_1}(-4) + \dots + \Delta_{z_l}(-4)$. Then we get

$$D^* Y(b_1, z_1) \cdots Y(b_l, z_l) \mathbb{1} = \Delta Y(b_1, z_1) \cdots Y(b_l, z_l) \mathbb{1} \quad (4.3)$$

for any $b_1, \dots, b_l \in A$. Let $f : W_0 \rightarrow \mathbb{k}$ be a functional and $\alpha_f \in \Phi^\Gamma$ be the corresponding correlation function, as in §4. The formula (4.3) implies that $f(D^*W_1) = 0$ if and only if $\Delta\alpha_f = 0$.

Fix a pair of indices $1 \leq i < j \leq l$, and let $\Delta' = \Delta - \Delta_{z_i}(-4)$ and $\Delta'' = \Delta - \Delta_{z_i}(-4) - \Delta_{z_j}(-4)$.

Lemma 4.2. *The expansion of $\Delta\alpha$ in the powers of $z_i - z_j$ has form*

$$\Delta\alpha = (z_i - z_j)^{-4} \Delta'' \alpha_{-4} + (z_i - z_j)^{-2} \Delta' \alpha_{-2} + O((z_i - z_j)^{-1}),$$

where α_k 's are the coefficients of α in the expansion (4.1).

Proof. Using the commutation relation $[\Delta, (z_i - z_j)^k] = k(z_i - z_j)^k(z_i + z_j)$, we compute

$$\begin{aligned} \Delta(z_i - z_j)^k \alpha_k &= (z_i - z_j)^k (\Delta + k(z_i + z_j)) \alpha_k \\ &= (z_i - z_j)^k (\Delta'' + \Delta_{z_i}(k+4) + \Delta_{z_j}(k+4)) \alpha_k. \end{aligned}$$

If $k = -4$, then, since $\partial_{z_i} \alpha_{-4} = \partial_{z_j} \alpha_{-4} = 0$, we get $(\Delta'' + \Delta_{z_i}(0) + \Delta_{z_j}(0)) \alpha_{-4} = \Delta'' \alpha_{-4}$, and if $k = -2$, then $(\Delta'' + \Delta_{z_i}(2) + \Delta_{z_j}(2)) \alpha_{-2} = \Delta' \alpha_{-2} + O(z_i - z_j)$, and the lemma follows. \square

Corollary 4.2. *Let $f : W_0 \rightarrow \mathbb{k}$ be a functional and $\alpha_f \in \Phi^\Gamma$ be the corresponding correlation function. Then $f(D^*W'_1) = 0$ if and only if $\Delta\alpha_f$ has only simple poles.*

Proof. By Lemma 4.1 and Lemma 4.2, the function $\Delta\alpha_f$ has only simple poles if and only if $f(D^*w) = 0$ for every word $w \in W'_1$ of the form (4.2). Since these words span W'_1 , this is equivalent to having $f(D^*W'_1) = 0$. \square

5. RATIONAL FUNCTIONS AND DIFFERENTIAL OPERATORS

5.1. Reductions of Proposition 3.1. We are going to reduce Proposition 3.1 to some elementary facts about rational functions. Our first step in this direction is the following:

Lemma 5.1. *$D^*V_1^{(l)} \cap V_0^{(l-1)} = D^*V_1^{(l-1)}$ for all $l \geq 1$.*

Note that a priori we have $D^*V_1^{(l)} \cap V_0^{(l-1)} \supseteq D^*V_1^{(l-1)}$.

Reduction of Proposition 3.1 to Lemma 5.1.

The component of length l of $\text{gr } D^*\widehat{V}_1$ is $D^*\widehat{V}_1 \cap \widehat{V}_0^{(l)} / D^*\widehat{V}_1 \cap \widehat{V}_0^{(l-1)}$, and by Theorem 2, the corresponding component of D^*F_1 is isomorphic to $D^*V_1^{(l)} / D^*V_1^{(l-1)}$. So it is enough to show that $D^*\widehat{V}_1 \cap \widehat{V}_0^{(l)} = D^*\widehat{V}_1^{(l)}$.

Clearly, we have $D^*\widehat{V}_1 \cap \widehat{V}_0^{(l)} \supset D^*\widehat{V}_1^{(l)}$. Assume on the contrary that there is $v \in \widehat{V}_1^{(k)} \setminus \widehat{V}_1^{(k-1)}$, such that $D^*v \in \widehat{V}_0^{(l)}$, where $k > l$ is minimal possible. Then $D^*v \in \widehat{V}_0^{(k-1)}$, so if Lemma 5.1 holds, then there is $u \in \widehat{V}_1^{(k-1)}$ such that $D^*u = D^*v \in \widehat{V}_0^{(l)}$, which contradicts the minimality of k . \square

As in § 4, for a weight $\lambda = a_1 + \dots + a_l \in \mathbb{Z}_+[A]$, let $W_d = W_d(\lambda) \subseteq \widehat{V}^{(l)}$ be the span of all words $a_1(m_1) \dots a_l(m_l) \mathbf{1} \in \widehat{V}$ of degree d and $W'_d = W_d \cap \widehat{V}^{(l-1)}$. We show next that Lemma 5.1 follows from the following statement.

Lemma 5.2. $D^*W_1 \cap W'_0 = D^*W'_1$.

Reduction of Lemma 5.1 to Lemma 5.2. Assume on the contrary that Lemma 5.1 fails. This means that there is $w \in \widehat{V}_1^{(l)}$ such that $D^*w \in \widehat{V}_0^{(l-1)} \setminus D^*\widehat{V}_1^{(l-1)}$. We can represent w as a sum $w = \sum_{\lambda \in \mathbb{Z}_+[A]} w_\lambda$ for $w_\lambda \in W_1(\lambda)$. Then, using Theorem 2, we see that $D^*w_\lambda \in \widehat{V}_0^{(l-1)}$ for all λ , and at least for some λ we must have $D^*w_\lambda \notin D^*\widehat{V}_1^{(l-1)}$. \square

Recall that $\Phi = \Phi(K)$ for the matrix $K = \{k_{ij}\}$ with $k_{ii} = 0$ and $k_{ij} = -4$, and $\Phi^s \subset \Phi$ are those functions that have only simple poles.

Lemma 5.3. *If $\alpha \in \Phi_{-2l}$ is such that $\Delta\alpha \in \Phi_{-2l+1}^s$, then there is a function $\beta \in \Phi_{-2l}^s$ such that $\Delta\beta = \Delta\alpha$.*

Reduction of Lemma 5.2 to Lemma 5.3. Clearly, $D^*W_1 \cap W'_0 \supseteq D^*W'_1$. Let $f : W_0 \rightarrow \mathbb{k}$ be a functional such that $f(D^*W'_1) = 0$. We show that Lemma 5.3 implies that $f(D^*W_1 \cap W'_0) = 0$.

Let $\alpha = \alpha_f$ be the correlation function corresponding to f , constructed in § 1.2. By Corollary 4.2 we know that $\Delta\alpha \in (\Phi^s)^\Gamma$. If Lemma 5.3 holds, then there is a function $\beta \in \Phi^s$ such that $\Delta\alpha = \Delta\beta$. By taking a Γ -orbit of β , we can make sure that $\beta \in (\Phi^s)^\Gamma$. But then by § 1.8 there is a functional $g : F_\lambda \rightarrow \mathbb{k}$ such that β is the correlation function of g on F_λ . Lift g to a functional $g : W_0 \rightarrow \mathbb{k}$ such that $g(W'_0) = 0$. Then look at the functional $f - g$. We have $(f - g)|_{W'_0} = f|_{W'_0}$, and $(f - g)(D^*W_1) = 0$. It follows that $f(D^*W_1 \cap W'_0) = 0$. \square

Note that Lemma 5.3 is just a statement about rational functions from $\Phi(K)$. We will prove this lemma in § 5.3, but first we need to look at such spaces of rational functions more closely.

5.2. Regular functions. Consider again the differential operator $\Delta_z(n) = z^2\partial_z - nz$. For the formal variables z_1, \dots, z_l , and for a sequence of integers n_1, \dots, n_l set

$$\Delta(n_1, \dots, n_l) = \Delta_{z_1}(n_1) + \dots + \Delta_{z_l}(n_l).$$

Using the relations

$$\begin{aligned} \Delta(n_1, \dots, n_l)(z_i - z_j) &= (z_i - z_j) \Delta(n_1, \dots, n_i - 1, \dots, n_j - 1, \dots, n_l), \\ \Delta(n_1, \dots, n_l)z_i &= z_i \Delta(n_1, \dots, n_i - 1, \dots, n_l), \end{aligned} \tag{5.1}$$

we see that $\Delta(n_1, \dots, n_l)$ is an operator on $\Phi(K)$ of degree 1.

We will call a function $\alpha \in \Phi(K)$ *regular* if $\Delta(n_1, \dots, n_l)\alpha = 0$.

Our first concern is to describe the space $\text{Ker } \Delta(n_1, \dots, n_l) \subset \Phi(K)$ of regular functions.

Take another integer symmetric $l \times l$ matrix $S = \{s_{ij}\}$. We say that $S \geq K$ if $s_{ij} \geq k_{ij}$ for every $1 \leq i, j \leq l$. For such a matrix S define

$$\pi(S) = \prod_{1 \leq i \leq l} z_i^{s_{ii}} \prod_{1 \leq i < j \leq l} (z_i - z_j)^{s_{ij}} \in \Phi(K). \tag{5.2}$$

The relations (5.1) imply that $\Delta(n_1, \dots, n_l)\pi(S) = 0$ if $\sum_{j=1}^l s_{ij} = n_i$ for all $1 \leq i \leq l$.

Lemma 5.4. *The kernel of $\Delta(n_1, \dots, n_l) : \Phi(K) \rightarrow \Phi(K)$ is*

$$\text{Span} \left\{ \pi(S) \mid S = \{s_{ij}\} \geq K, \sum_{j=1}^l s_{ij} = n_i \ \forall \ 1 \leq i \leq l \right\}.$$

Proof. Step 1. Set $k_i = \sum_{j=1}^l k_{ij}$ for $1 \leq i \leq l$. If $\alpha \in \Phi(K)$ is given by (1.5), then (5.1) implies that $\Delta(n_1, \dots, n_l) \alpha = 0$ if and only if $\Delta(n_1 - k_1, \dots, n_l - k_l) p = 0$. Therefore, it is enough to prove the Lemma for the case when $K = 0$.

Step 2. So let $p \in \mathbb{k}[z_1, \dots, z_l]$ be a polynomial such that $\Delta(n_1, \dots, n_l) p = 0$. Expand p in the powers of $z_{l-1} - z_l$:

$$p = p_0 + (z_{l-1} - z_l) p_1 + \dots + (z_{l-1} - z_l)^m p_m, \quad p_i \in \mathbb{k}[z_1, \dots, z_{l-1}]. \quad (5.3)$$

Using (5.1) we compute

$$\begin{aligned} \Delta(n_1, \dots, n_l) (z_{l-1} - z_l)^k p_k &= (z_{l-1} - z_l)^k \Delta(n_1, \dots, n_{l-2}, n_{l-1} - k, n_l - k) p_k \\ &= (z_{l-1} - z_l)^k \left(\Delta(n_1, \dots, n_{l-2}, n_{l-1} - k) p_k - (n_l - k) z_l p_k \right) \\ &= (z_{l-1} - z_l)^k \Delta(n_1, \dots, n_{l-2}, n_{l-1} + n_l - 2k) p_k + (n_l - k) (z_{l-1} - z_l)^{k+1} p_k. \end{aligned}$$

By taking $k = 0$ we see that $\Delta(n_1, \dots, n_{l-2}, n_{l-1} + n_l) p_0 = 0$. Taking $k = m$ implies that $m = n_l$, and therefore, since we could take any other pair $z_i - z_j$ instead of $z_{l-1} - z_l$, we conclude that $\deg_{z_i} p = n_i$.

Step 3. We prove the Lemma by induction on l and $\sum n_i$. By Step 2, if $n_i < 0$ for at least one i , then $\text{Ker } \Delta(n_1, \dots, n_l) = 0$, and also $\text{Ker } \Delta(0, \dots, 0) = \mathbb{k}$. On the other hand, if $l = 1$, then $\Delta(n)p(z) = 0$ means that $p(z)$ satisfies the differential equation $zp' - np = 0$, therefore $p \in \mathbb{k}z^n$. So the Lemma holds for all these cases.

If p is divisible by some $z_i - z_j$, then $p = (z_i - z_j)q$, so that $\Delta(n_1, \dots, n_i - 1, \dots, n_j - 1, \dots, n_l) q = 0$; by induction, q is a linear combination of the products $\pi(S)$, and therefore, so is p . So we can assume that $p_0 \neq 0$ in (5.3).

By Step 2, we have $\Delta(n_1, \dots, n_{l-2}, n_{l-1} + n_l) p_0 = 0$, therefore by induction,

$$p_0 = \sum_r \xi_r \pi(S^{(r)}),$$

where $\xi_r \in \mathbb{k}$, and $S^{(r)} = \{s_{ij}^{(r)}\}$, $s_{ij}^{(r)} \in \mathbb{Z}_+$, is a symmetric $l-1 \times l-1$ matrix such that

$$\sum_{j=1}^{l-1} s_{ij}^{(r)} = n_i \quad \text{for } 1 \leq i \leq l-2 \quad \text{and} \quad \sum_{j=1}^{l-1} s_{j, l-1}^{(r)} = n_{l-1} + n_l.$$

For each such matrix $S^{(r)}$ fix a symmetric $l \times l$ matrix $\hat{S}^{(r)} = \{\hat{s}_{ij}^{(r)}\}$, $\hat{s}_{ij}^{(r)} \in \mathbb{Z}_+$, such that

- $\hat{s}_{ij}^{(r)} = s_{ij}^{(r)}$ for $1 \leq i, j \leq l-2$,
- $\hat{s}_{i, l-1}^{(r)} + \hat{s}_{il}^{(r)} = s_{i, l-1}^{(r)}$ for $1 \leq i \leq l-2$,
- $\hat{s}_{l-1, l-1}^{(r)} + \hat{s}_{ll}^{(r)} = s_{l-1, l-1}^{(r)}$,
- $\hat{s}_{l-1, l}^{(r)} = 0$,
- $\sum_{j=1}^l \hat{s}_{ij}^{(r)} = n_i$ for $i = l-1, l$.

Clearly, such a choice of $\hat{S}^{(r)}$ always exists, and we have $\sum_{j=1}^l \hat{s}_{ij}^{(r)} = n_i$ for all $1 \leq i \leq l$, which implies that $\Delta(n_1, \dots, n_l) \pi(\hat{S}^{(r)}) = 0$. We also have

$$\pi(\hat{S}^{(r)})|_{z_l = z_{l-1}} = \pi(S^{(r)}).$$

Now take $q = \sum_r \xi_r \pi(\hat{S}^{(r)})$. Then $p - q$ is divisible by $z_{l-1} - z_l$, and also $\Delta(n_1, \dots, n_l) (p - q) = 0$, so $p - q$ is a linear combination of products $\pi(S)$, hence so is p . \square

We will need the following simple corollary of Lemma 5.4 for the case when the matrix K has zeros on the diagonal:

Corollary 5.1. *Let $\Delta = \Delta(n_1, \dots, n_l)$ and $K = \{k_{ij}\}$ be a symmetric integer $l \times l$ matrix with $k_{ii} = 0$. Let $\Phi = \Phi(K) = \bigoplus_{d \in \mathbb{Z}} \Phi_d$, and $(\text{Ker } \Delta)_d = \text{Ker } \Delta \cap \Phi_d$. Then $(\text{Ker } \Delta)_d = 0$ for $d < d_0 = \frac{1}{2} \sum n_i$ and if $d_0 \in \mathbb{Z}$, then*

$$(\text{Ker } \Delta)_{d_0} = \text{Span} \left\{ \pi(S) \mid S = \{s_{ij}\} \geq K, s_{ii} = 0, \sum_j s_{ij} = n_i \right\}.$$

Proof. Let $S = \{s_{ij}\}$ be a symmetric integer matrix, such that $\pi(S) \in (\text{Ker } \Delta)_d$. Then $\sum_j s_{ij} = n_i$ because $\Delta \pi(S) = 0$, and $s_{ii} \geq 0$ because Φ does not contain functions with poles at z_i . By Lemma 5.4, the space $(\text{Ker } \Delta)_d$ is spanned by the products $\pi(S)$ for such S . Denote

$$S_0 = \sum_{i=1}^l s_{ii} \geq 0, \quad S_1 = \sum_{1 \leq i < j \leq l} s_{ij}.$$

Then we have $S_0 + 2S_1 = \sum_i n_i$ and $S_0 + S_1 = d$, therefore $0 \leq S_0 = 2d - \sum_i n_i$. If $2d = \sum_i n_i$, then $S_0 = 0$, which can happen only if $s_{ii} = 0$ for all $1 \leq i \leq l$. \square

Remark. Let V be a vertex algebra generated by minimal elements. Corollary 5.1 implies that $D^*V_{d+1} = V_d$ for $d < 0$. Indeed, let $f : V_d \rightarrow \mathbb{k}$ be a linear functional, such that $f(D^*V_{d+1}) = 0$. Take some minimal elements $a_1, \dots, a_l \in V$ of degrees $\deg a_i = d_i$. Then the corresponding correlation function α_f belongs to $\Phi(K)$ for $K = \{k_{ij}\}$ given by $k_{ii} = 0$ and $k_{ij} = -N(a_i, a_j)$, see § 1.2. By Corollary 5.1, if $\alpha_f \neq 0$, then $\deg \alpha_f \geq -\sum d_i$, therefore $d = \deg \alpha_f + \sum d_i \geq 0$.

5.3. Proof of Lemma 5.3. Take a matrix $K = \{k_{ij}\}_{i,j=1}^l$ such that $k_{ij} < 0$ for $i \neq j$ and $k_{ii} = 0$. Let $K^s = \{k_{ij}^s\} \geq K$ be defined by $k_{ij}^s = -1$ for $i \neq j$ and $k_{ii}^s = 0$. Then $\Phi^s = \Phi(K^s) \subset \Phi(K)$ is the space of functions in $\Phi(K)$ that have at most simple poles at $z_i - z_j$. Denote $\Delta = \Delta(-4, \dots, -4)$ (l times). We need to show that if $\alpha \in \Phi(K)_{-2l}$ is such that $\Delta \alpha \in \Phi_{-2l+1}^s$, then there is a function $\beta \in \Phi_{-2l}^s$ such that $\Delta \beta = \Delta \alpha$.

Note that the degree of any function in Φ^s is at least $-l(l-1)/2$, therefore, if $l \leq 4$, then $\Phi_{-2l+1}^s = 0$, so we can assume that $l \geq 5$.

Using induction, we can assume that the lemma is true for any matrix $K' > K$.

Assume that $k = k_{l-1,l} < -1$. Expand α into a power series in $z_{l-1} - z_l$:

$$\alpha = \sum_{n \geq 0} (z_{l-1} - z_l)^{n+k} \alpha_n(z_1, \dots, z_{l-1}).$$

Using (5.1), we compute

$$\Delta \alpha = (z_{l-1} - z_l)^{n+k} \Delta(-4, \dots, -4, -8 - 2k) \alpha_0 + O((z_{l-1} - z_l)^{k+1}),$$

and therefore $\Delta(-4, \dots, -4, -8 - 2k) \alpha_0 = 0$.

The function α_0 belongs to $\Phi(N)$, for the matrix $N = \{n_{ij}\}_{i,j=1}^{l-1}$ given by

$$n_{ij} = k_{ij} \text{ for } 1 \leq i < j \leq l-2 \text{ and } n_{i,l-1} = k_{i,l-1} + k_{il} \text{ for } 1 \leq i \leq l-2.$$

By Corollary 5.1, α_0 is a linear combination of products $\pi(S)$ for some symmetric integer $(l-1) \times (l-1)$ matrices $S = \{s_{ij}\} \geq N$ such that

$$s_{ii} = 0 \text{ and } \sum_{j=1}^{l-1} s_{ij} = \begin{cases} -4 & \text{if } 1 \leq i \leq l-2, \\ -8 - 2k & \text{if } i = l-1. \end{cases}$$

For any such matrix S we choose a symmetric integer $l \times l$ matrix $\hat{S} = \{\hat{s}_{ij}\}$ such that

- $\hat{s}_{ii} = 0$,
- $\hat{s}_{ij} = s_{ij}$ for $1 \leq i, j \leq l-2$,
- $\hat{s}_{ij} \geq \min\{-1, k_{ij}\}$,
- $\hat{s}_{i,l-1} + \hat{s}_{il} = s_{i,l-1}$ for $1 \leq i \leq l-2$,
- $\hat{s}_{l-1,l} = k$,
- $\sum_{i=1}^{l-2} \hat{s}_{i,l-1} = \sum_{i=1}^{l-2} \hat{s}_{i,l} = -4 - k$.

This is possible due to the following statement, which we will prove at the end of this subsection:

Lemma 5.5. *For any integers s_1, \dots, s_m , k'_1, \dots, k'_m , k''_1, \dots, k''_m and n , such that*

$$m + n \geq 0, \quad k'_i, k''_i \leq -1, \quad s_i \geq k'_i + k''_i \quad \text{and} \quad s_1 + \dots + s_m = 2n,$$

it is always possible to partition $s_i = s'_i + s''_i$ so that

$$s'_i \geq k'_i, \quad s''_i \geq k''_i \quad \text{and} \quad s'_1 + \dots + s'_m = s''_1 + \dots + s''_m = n.$$

We apply Lemma 5.5 to the sequences $s_{1,l-1}, \dots, s_{l-2,l-1}$, $\min\{-1, k_{1,l-1}\}, \dots, \min\{-1, k_{l-2,l-1}\}$, $\min\{-1, k_{1l}\}, \dots, \min\{-1, k_{l-2,l}\}$, taking $m = l - 2$ and $n = -4 - k$. Since $l \geq 5$ and $k \leq -2$, we have $m + n = l - 2 - k - 4 \geq 1$. Then we set $\hat{s}_{i,l-1} = s'_i$ and $\hat{s}_{il} = s''_i$. Note that $\pi(\hat{S}) \in \Phi(K)$ and $\Delta\pi(\hat{S}) = 0$.

So if $\alpha_0 = \sum_m \xi_m \pi(S^{(m)})$ for $\xi_m \in \mathbb{k}$, then set $\gamma = \sum_m \xi_m \pi(\hat{S}^{(m)}) \in \Phi(K)$. Then $\Delta\gamma = 0$ and


$$\gamma = (z_{l-1} - z_l)^k \alpha_0 + O((z_{l-1} - z_l)^{k+1}).$$

Now consider the function $\alpha - \gamma$. We have $\Delta\alpha - \gamma = \Delta\alpha \in \Phi^s$. The order of $\alpha - \gamma$ at $z_{l-1} - z_l$ is bigger than $k = k_{l-1,l}$, and the order of $\alpha - \gamma$ at any other point $z_i - z_j$ is at least that of α , therefore $\alpha - \gamma$ belongs to a space of functions $\Phi(K')$ for some matrix $K' > K$. By induction, there is a function $\beta \in \Phi^s$ such that $\Delta\beta = \Delta\alpha$. \square

Proof of Lemma 5.5. Set $x_i = s'_i$. We are looking for a vector $(x_1, \dots, x_m) \in \mathbb{Z}^m$ that belongs to the box $k'_i \leq x_i \leq s_i - k''_i$ such that $x_1 + \dots + x_m = n$. To show that this box contains a vector with the sum of coordinates equal to n , we need to check that the sum of coordinates of the two extremal vertices bound n from above and from below. Indeed, using that $k'_i, k''_i \leq -1$ and $m \geq -n$, we get $\sum k'_i \leq -m \leq n$, and similarly, $\sum_{i=1}^m s_i - k''_i = 2n - \sum k''_i \geq 2n + m \geq n$. \square

5.4. Some combinatorics. Using Lemma 5.4 we can investigate the dimensions of the spaces $F_{\lambda,0}/D^*F_{\lambda,1}$, where F is the free vertex algebra generated by \mathcal{A} (see §3.1) and $\lambda \in \mathbb{Z}_+[\mathcal{A}]$ is a weight. We see that if $\lambda = a_1 + \dots + a_l$ is such that $a_i \neq a_j \in \mathcal{A}$, then the space $F_{\lambda,0}/D^*F_{\lambda,1}$ is in one-to-one correspondence with the span Π of products $\pi(S)$, where $S = \{s_{ij}\}$ runs over all $l \times l$ symmetric integer matrices such that $s_{ii} = 0$, $s_{ij} \geq -1$ and $\sum_i s_{ij} = -4$ for every $1 \leq j \leq l$.

To every such product $\pi(S)$ we can correspond a diagram in the following way: take l points p_1, \dots, p_l on a line, and connect points p_i and p_j with $s_{ij} + 1$ arcs placed above the line. Then one can show that these of the products that correspond to a planar diagram form a basis of Π . For example, if $l = 6$, then $\dim F_{\lambda,0}/D^*F_{\lambda,1} = 5$, since there are 5 such planar diagrams,



that correspond to the following matrices S :

$$\begin{pmatrix} 0 & 0 & -1 & -1 & -1 & -1 \\ 0 & 0 & -1 & -1 & -1 & -1 \\ -1 & -1 & 0 & 0 & -1 & -1 \\ -1 & -1 & 0 & 0 & -1 & -1 \\ -1 & -1 & -1 & -1 & 0 & 0 \\ -1 & -1 & -1 & -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 & -1 & -1 & -1 \\ 0 & 0 & -1 & -1 & -1 & -1 \\ -1 & -1 & 0 & -1 & -1 & 0 \\ -1 & -1 & -1 & 0 & 0 & -1 \\ -1 & -1 & -1 & 0 & 0 & -1 \\ -1 & -1 & 0 & -1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & -1 & 0 & -1 & -1 \\ -1 & 0 & 0 & -1 & -1 & -1 \\ -1 & 0 & 0 & -1 & -1 & -1 \\ 0 & -1 & -1 & 0 & -1 & -1 \\ -1 & -1 & -1 & -1 & 0 & 0 \\ -1 & -1 & -1 & -1 & 0 & 0 \end{pmatrix},$$

$$\begin{pmatrix} 0 & -1 & -1 & -1 & -1 & 0 \\ -1 & 0 & 0 & -1 & -1 & -1 \\ -1 & 0 & 0 & -1 & -1 & -1 \\ -1 & -1 & -1 & 0 & 0 & -1 \\ -1 & -1 & -1 & 0 & 0 & -1 \\ 0 & -1 & -1 & -1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & -1 & -1 & -1 & 0 \\ -1 & 0 & -1 & -1 & 0 & -1 \\ -1 & -1 & 0 & 0 & -1 & -1 \\ -1 & -1 & 0 & 0 & -1 & -1 \\ -1 & 0 & -1 & -1 & 0 & -1 \\ 0 & -1 & -1 & -1 & -1 & 0 \end{pmatrix}.$$

6. EXPANSIONS OF RATIONAL FUNCTIONS

6.1. Expansions of rational functions. Let $\alpha = \alpha(z_1, \dots, z_l)$ be a homogeneous rational function that has poles only at either z_i or $z_i - z_j$. In other words, $\alpha \in \Phi = \Phi(K)$ for some matrix K , see §1.2. Let $\xi = (\Xi_1, \dots, \Xi_k)$ be an ordered partition $\{1, \dots, l\} = \Xi_1 \sqcup \dots \sqcup \Xi_k$ of the set $\{1, \dots, l\}$ into a disjoint union of subspaces $\Xi_j \neq \emptyset$. Take a permutation $\sigma = (i_1, \dots, i_l)$ of $(1, \dots, l)$ such that $\Xi_1 = \{i_1, \dots, i_{|\Xi_1|}\}$, $\Xi_2 = \{i_{|\Xi_1|+1}, \dots, i_{|\Xi_1|+|\Xi_2|}\}$, etc. Consider the power series expansion α^σ of α in the domain $|z_{i_1}| > |z_{i_2}| > \dots > |z_{i_l}|$.

Take some part Ξ_j of the partition ξ . Let x be a monomial in the variables $\{z_i \mid i \notin \Xi_j\}$. Denote by α_x the coefficient of x in the series α^σ , so that we get

$$\alpha = \sum_x \alpha_x x. \quad (6.1)$$

If $\deg x = \deg \alpha - n$ for some $n \in \mathbb{Z}$, then $\deg \alpha_x = n$ and $\alpha_x \in \Phi(K_j)_n$, where K_j is the diagonal minor of K , corresponding to the set of indices Ξ_j . Therefore, the space $\text{Span}\{\alpha_x \mid \deg x = \deg \alpha - n\} \subset \Phi(K_j)_n$ is finite dimensional. Let $\alpha_{n,1}^{(j)}, \alpha_{n,2}^{(j)}, \dots$ be its basis. Now substitute every function α_x in (6.1) by its linear expansion in this basis and then apply the above expansion procedure to the coefficients of $\alpha_{n,i}^{(j)}$'s. So after several such steps our expansion of α becomes

$$\alpha = \sum_{n_1 + \dots + n_k = \deg \alpha} \sum_i \alpha_{n_1,i}^{(1)} \alpha_{n_2,i}^{(2)} \dots \alpha_{n_k,i}^{(k)}, \quad (6.2)$$

where $\alpha_{n,i}^{(j)}$ is a rational function in the variables $\{z_m \mid m \in \Xi_j\}$ of degree n_j .

The rational functions in this section are always obtained as correlation functions of linear functionals on \widehat{V} . As in §4.1, for any $a_1, \dots, a_l \in \mathcal{A}$ we set $W_0 \subset \widehat{V}_0$ to be the span of all words $a_1(m_1) \dots a_l(m_l) \mathbb{1}$ of degree 0, and then any functional $f : W_0 \rightarrow \mathbb{k}$ defines the correlation function $\alpha = \alpha_f(z_1, \dots, z_l)$ of degree $-2l$. As we have seen before, many properties of f can be expressed in terms of α . For example, $f(D^*W_1) = 0$ if and only if α is a regular function, see §5.2.

Now we are going to extend the definition of τ from §3.2 to rational functions. Recall that for $\alpha(z_1, \dots, z_l) \in \Phi$ and a permutation $\sigma \in S_l$ we have denoted by α^σ the power series expansion of α in the domain $|z_{i_1}| > |z_{i_2}| > \dots > |z_{i_l}|$. Set

$$\tau(\alpha) = \min \left\{ \tau(a_{\sigma(1)}(m_1) \dots a_{\sigma(l)}(m_l)) \mid \begin{array}{l} \sigma \in S_l, \text{ the coefficient of } z_{\sigma(1)}^{-m_1-1} \dots z_{\sigma(l)}^{-m_l-1} \\ \text{in } \alpha^\sigma \text{ is non-zero} \end{array} \right\}.$$

If $\alpha = \alpha_f$ is the correlation function of a functional $f : W_0 \rightarrow \mathbb{k}$, then, for example we have $f(Q_{\leq 1}) = 0$ if and only if $\tau(\alpha) > 1$.

Most of the rational functions α that we deal with will have $\tau(\alpha) \leq 0$. This means that in the expansion (6.2), a summand corresponding to n_1, \dots, n_k is equal to 0 whenever

$$n_j + n_{j+1} + \dots + n_k < -2|\Xi_j \sqcup \Xi_{j+1} \sqcup \dots \sqcup \Xi_k|$$

for some $1 < j \leq k$. To avoid writing cumbersome subscripts, let us omit the subscript n from $\alpha_{n,i}^{(j)}$ when $n = -2|\Xi_j|$. Using this convention, denote by

$$\alpha(\xi) = \sum_i \alpha_i^{(1)} \alpha_i^{(2)} \dots \alpha_i^{(k)} \quad (6.3)$$

the “lowest degree” component of α in the expansion (6.2).

6.2. Proof of Proposition 3.3. We are going to prove a slightly stronger statement. Let

$$\begin{aligned} Q'_{\leq 1} = & \text{Span} \{ w \mathbb{1} \mid w \in X^*, \deg w = 0, \tau(w) < 0 \} \\ & \cup \{ u a(1) w \mathbb{1} \mid u, w \in X^*, \deg u = \deg w = 0, a \in \mathcal{A} \} \\ & \cup \{ u a(2) w \mathbb{1} \mid u, w \in X^*, \deg u = -1, \deg w = 0, a \in \mathcal{A} \}. \end{aligned}$$

Lemma 1.2 (b) implies that $Q'_{\leq 1} \subset \text{Rad}_0$. Recall that we have denoted $H = D^* \widehat{V}_1$. We will prove

Lemma 6.1. $\text{Rad}_0^{(l)} = H^{(l)} + Q'_{\leq 1}^{(l)}$.

Clearly this lemma implies Proposition 3.3, because $N = \text{Rad}_0 \widehat{V} + Q_{\leq 1}$. Note that we only need to show inclusion “ \subseteq ”.

Proof. Step 0. First of all we note that it is enough to prove that $\text{Rad}_0 \cap W_0 = (H \cap W_0) + (Q'_{\leq 1} \cap W_0)$ for any space $W_0 \subset \widehat{V}_0$ corresponding to a collection $a_1, \dots, a_l \in \mathcal{A}$. This is because Rad_0 is spanned by all elements of the form

$$a_1(m_1) \dots a_j(m_j) D^* a_{j+1}(m_{j+1}) \dots a_l(m_l) \mathbb{1} \in \text{Rad}_0 \cap W_0,$$

such that $\deg a_1(m_1) \cdots a_j(m_j) = 0$, and $\deg a_{j+1}(m_{j+1}) \cdots a_l(m_l) = 1$, see § 1.6.

Step 1. We need to show that if a functional $f : W_0 \rightarrow \mathbb{k}$ satisfies $f(Q'_{\leq 1}) = f(H) = 0$, then also $f(\text{Rad}_0) = 0$.

It is easy to see what the condition $f(\text{Rad}_0) = 0$ means in terms of the corresponding correlation function $\alpha = \alpha_f$. Let $\xi = (\Xi_1, \Xi_2)$, $\Xi_1 = \{i_1, \dots, i_k\}$, $\Xi_2 = \{i_{k+1}, \dots, i_l\}$ be an ordered partition of $\{1, \dots, l\}$. For a degree 0 word $u = a_{i_1}(m_1) \cdots a_{i_k}(m_k)$ consider the correlation function

$$\alpha_u^{(2)}(z_{i_{k+1}}, \dots, z_{i_l}) = f(u Y(a_{i_{k+1}}, z_{i_{k+1}}) \cdots Y(a_{i_l}, z_{i_l}) \mathbf{1}). \quad (6.4)$$

Note that $\alpha_u^{(2)} = \alpha_x$ (see (6.1)) for $x = z_{i_1}^{-m_1-1} \cdots z_{i_k}^{-m_k-1}$. Then $f(\text{Rad}_0) = 0$ if and only if $\alpha_u^{(2)}$ is a regular function for every u . This is equivalent to the condition that $\alpha_i^{(2)}$ is regular for every i in (6.2).

Step 2. We show here that for any word $v = a_{i_{k+1}}(m_{k+1}) \cdots a_{i_l}(m_l)$ the function

$$\alpha_v^{(1)}(z_{i_1}, \dots, z_{i_k}) = f(Y(a_{i_1}, z_{i_1}) \cdots Y(a_{i_k}, z_{i_k}) v \mathbf{1})$$

is regular. Since $\alpha_v^{(1)} = \alpha_x$ for $x = z_{i_{k+1}}^{-m_{k+1}-1} \cdots z_{i_l}^{-m_l-1}$, this is equivalent to having all $\alpha_i^{(1)}$'s regular in (6.2).

Denote $\Delta^{(m)} = \sum_{j \in \Xi_m} \Delta_{z_j}(-4)$ for $m = 1, 2$ (see § 5.2). We apply Δ to (6.2) and get

$$0 = \Delta \alpha = \sum_{n \geq -2(l-k)} \sum_i (\Delta^{(1)} \alpha_{-2l-n,i}^{(1)}) \alpha_{n,i}^{(2)} + \alpha_{-2l-n,i}^{(1)} (\Delta^{(2)} \alpha_{n,i}^{(2)}).$$

In this expansion the component of lowest degree in the variables $z_{i_{k+1}}, \dots, z_{i_l}$ is

$$\sum_i (\Delta^{(1)} \alpha_i^{(1)}) \alpha_i^{(2)},$$

and it has to be equal to 0, therefore $\Delta^{(1)} \alpha_i^{(1)} = 0$ for all i .

Step 3. Now we show that for any words $u = a_{i_1}(m_1) \cdots a_{i_k}(m_k)$ and $v = a_{i_{k+1}}(m_{k+1}) \cdots a_{i_l}(m_l)$ as above we have $f((u - u^*)v \mathbf{1}) = 0$. Indeed, it follows from Lemma 1.2 (a) that $u - u^* = [D^*, u_1] + u_0$, for some $u_1 \in U(\widehat{V})_1$ and $u_0 \in U(\widehat{V})_0$ such that $u_0 \mathbf{1} = 0$. But then u_0 belongs to the left ideal in $U(\widehat{V})$ generated by the set $\{a(m) \mid a \in A, m \in \mathbb{Z}_+\}$, therefore $u_0 v \mathbf{1} \in Q'_{\leq 1}$ and hence $f(u_0 v \mathbf{1}) = 0$. So we get $f((u - u^*)v \mathbf{1}) = f([D^*, u_1]v \mathbf{1}) = 0$ by Step 2.

needs a ref
in § 1

Step 4. By Lemma 1.2 (a) and the assumptions on f , we have $f((w - w^*) \mathbf{1}) = 0$. So we compute, using Step 3,

$$\begin{aligned} \alpha_u^{(2)} &= \sum_{m_{k+1}, \dots, m_l} f(u a_{i_{k+1}}(m_{k+1}) \cdots a_{i_l}(m_l) \mathbf{1}) z_{i_{k+1}}^{-m_{k+1}-1} \cdots z_{i_l}^{-m_l-1} \\ &= \sum_{m_{k+1}, \dots, m_l} f((a_{i_{k+1}}(m_{k+1}) \cdots a_{i_l}(m_l))^* u^* \mathbf{1}) z_{i_{k+1}}^{-m_{k+1}-1} \cdots z_{i_l}^{-m_l-1} \\ &= \sum_{m_{k+1}, \dots, m_l} f(a_{i_{k+1}}(m_{k+1}) \cdots a_{i_l}(m_l) u^* \mathbf{1}) z_{i_{k+1}}^{-m_{k+1}-1} \cdots z_{i_l}^{-m_l-1}, \end{aligned}$$

and this is exactly $\alpha_{u^*}^{(1)}(z_{i_{k+1}}, \dots, z_{i_l})$ corresponding to the opposite partition $\{1, \dots, l\} = \Xi_2 \sqcup \Xi_1$, which is regular by Step 3. \square

6.3. Products of functions. Let $\xi = (\Xi_1, \dots, \Xi_k)$ and $\theta = (\Theta_1, \dots, \Theta_s)$ be two ordered partitions of $\{1, \dots, l\}$ as above. Define their product to be

$$\xi\theta = (\Xi_1 \cap \Theta_1, \Xi_1 \cap \Theta_2, \dots, \Xi_1 \cap \Theta_s, \dots, \Xi_k \cap \Theta_s).$$

Note that the partitions $\xi\theta$ and $\theta\xi$ differ only by a permutation of parts.

For the proof of Proposition 3.4 we will need the following lemma.

Lemma 6.2. *Let $\xi = (\Xi_1, \dots, \Xi_k)$ be a partition of $\{1, \dots, l\}$, let $\alpha^{(j)}$, $1 \leq j \leq k$, be a rational function as above in the variables $\{z_i \mid i \in \Xi_j\}$, and let $\alpha = \alpha^{(1)} \dots \alpha^{(k)}$.*

- (a) *Let $n = -1, 0$ or 1 . If $\tau(\alpha^{(j)}) > n$ for all $1 \leq j \leq k$, then also $\tau(\alpha) > n$.*
- (b) *Let $\theta = (\Theta_1, \dots, \Theta_s)$ be another partition, and assume that $\tau(\alpha^{(j)}) > 0$ for every $1 \leq j \leq k$. Then $\alpha(\theta) = \alpha(\theta\xi)$.*

Proof. Take a permutation $\sigma \in S_l$ and let α^σ be the corresponding power series expansion of α . For a subset $\Omega \subset \{1, \dots, l\}$ consider a monomial $x = \prod_{i \in \Omega} z_i^{-m_i-1}$. Let α_x be the coefficient of x in the series α^σ . Then

$$\alpha_x = \alpha_{x_1}^{(1)} \dots \alpha_{x_k}^{(k)}, \quad (6.5)$$

where $x_j = \prod_{i \in \Xi_j \cap \Omega} z_i^{-m_i-1}$ and $\alpha_{x_j}^{(j)}$ is the coefficient of x_j in the power series expansion of $\alpha^{(j)}$.

For a monomial $x = z_{i_1}^{-m_1-1} \dots z_{i_t}^{-m_t-1}$ let $w(x) = a_{i_1}(m_1) \dots a_{i_t}(m_t)$ be the corresponding word. Then we have

$$\deg w(x) = \sum_{j=1}^k \deg w(x_j).$$

In order to prove (a), take $x = z_{\sigma(t)}^{-m_t-1} \dots z_{\sigma(l)}^{-m_l-1}$ for some $1 \leq t \leq l$, set $w = w(x)$ and assume that the following condition holds:

$$\deg w < 0 \text{ or } (n \geq 0 \text{ and } w = a_{\sigma(t)}(1)v, \deg v = 0) \text{ or } (n = 1 \text{ and } \deg w = 1).$$

Clearly, we have $\tau(\alpha) > n$ if and only if $\alpha_x = 0$ for all such monomials x .

Assume that $\deg w < 0$. Then there must exist some $1 \leq j \leq k$ such that $\deg w(x_j) < 0$. But then $\alpha_{x_j}^{(j)} = 0$ since $\tau(\alpha_j) \geq 0$, therefore $\alpha_x = 0$ by (6.5).

If $n \geq 0$ and $w = a_{\sigma(t)}(1)v$ for $\deg v = \deg w = 0$, then either there is j such that $\deg w(x_j) < 0$ or we have $\deg w(x_j) = 0$ for all j . In the latter case, let j be such that $\sigma(t) \in \Xi_j$. Then $\alpha_{x_j}^{(j)} = \alpha_x = 0$.

Finally, if $n = 1$ and $\deg w = 1$, then there is j such that either $\deg w(x_j) < 0$ or $\deg w(x_j) = 1$. In both cases $\alpha_{x_j}^{(j)} = \alpha_x = 0$.

Now we prove (b). Take a part Θ_r of θ , and set $\Omega = \{1, \dots, l\} \setminus \Theta_r$. Let $x = x_r$ be a monomial in $\{z_i \mid i \in \Omega\}$ of degree $-2|\Omega|$, so that $\deg w(x_r) = 0$. Then by (6.5) we get

$$\alpha_{x_r} = \alpha_{x_{r1}}^{(1)} \dots \alpha_{x_{rk}}^{(k)},$$

and this proves (b), because the component $\alpha(\theta)$ is by definition spanned by the products $\alpha_{x_1} \dots \alpha_{x_s}$, see § 6.1. \square

6.4. Proof of Proposition 3.4. In this section we deal with the free vertex algebra F . Fix a weight $\lambda = a_1 + \dots + a_l \in \mathbb{Z}[\mathcal{A}]$. As in § 1.2, for a functional $f : F_{\lambda,0} \rightarrow \mathbb{k}$ there corresponds the correlating function $\alpha_f \in \Phi(\mathbf{K}) = \Phi$ of degree $-2l$, where the $l \times l$ matrix \mathbf{K} has 0 on the diagonal and -1 everywhere else. By § 1.8 this gives a one-to-one correspondence between $\text{Hom}(F_{\lambda,0}, \mathbb{k})$ and Φ_{-2l}^Γ , where Γ is the group of permutations generated by the transpositions (i, j) such that $a_i = a_j$. We will call the functions from Φ_{-2l}^Γ *admissible*.

We need to show that $(E_0 + E_1) \cap (D^*F_1 + E_1) \subseteq P + C + E_1 \subset F_0$, since the other inclusion is already established in § 3.3. All these spaces are homogeneous, therefore it is sufficient to prove this statement for the homogeneous components of weight λ . We will show that if a functional $f : F_0 \rightarrow \mathbb{k}$ satisfies $f(P + C + E_1) = 0$, then also $f((E_0 + E_1) \cap (D^*F_1 + E_1)) = 0$. Let $\alpha = \alpha_f \in \Phi(\mathbf{K})$ be the correlation function of f .

Claim A. For any partition $\xi = (\Xi_1, \dots, \Xi_k)$, the component $\alpha(\xi) \in \Phi_{-2l}$ is regular, has $\tau(\alpha(\xi)) > 1$ and satisfies

$$\sigma(\alpha(\xi)) = \sigma(\alpha)(\sigma(\xi)) = \alpha(\sigma(\xi)). \quad (6.6)$$

Proof. Let us investigate the conditions $f(P) = 0$, $f(C) = 0$ and $f(E_1) = 0$ separately.

We have $f(P) = 0$ if and only if for any partition $\xi = (\Xi_1, \Xi_2)$ all the terms $\alpha_i^{(2)}$ in $\alpha(\xi)$ (see (6.3)), are regular admissible. Indeed, clearly $\alpha_i^{(2)}$ is admissible, since its order of poles at $z_i - z_j$ or z_i cannot be bigger than that of α . And $\alpha_i^{(2)}$ is regular, since f is zero on any element of the form uD^*v for $1 \leq u \in X_0^*$ and $v \in F_1$, and therefore the function $\alpha_u^{(2)}$, given by (6.4), is regular.

It is easy to see that $f(C) = 0$ if and only if for any pair of partitions ξ and ξ' that differ one from another by a permutation of parts, we have $\alpha(\xi) = \alpha(\xi')$.

The condition $f(E_1) = 0$ means that $\tau(\alpha) > 1$. We claim that then $\tau(\alpha(\xi)) > 1$ for any partition $\xi = (\Xi_1, \dots, \Xi_k)$. Indeed, take some $\Xi_j = \{i_1, \dots, i_s\}$. By §6.1, every function $\alpha_i^{(j)}$ (see (6.2)) is a linear combination of the functions

$$\alpha_{uv} = \sum_{m_1 + \dots + m_s = s} f(u a_{i_1}(m_1) \cdots a_{i_s}(m_s) v \mathbb{1}) z_{i_1}^{-m_1-1} \cdots z_{i_s}^{-m_s-1}$$

for $u, v \in U(\widehat{V})_0$. If $\tau(a_{i_1}(m_1) \cdots a_{i_s}(m_s)) \leq 1$, then $f(u a_{i_1}(m_1) \cdots a_{i_s}(m_s) v \mathbb{1}) = 0$, therefore $\tau(f_{uv}) > 1$, and hence also $\tau(\alpha(\xi)) > 1$ by Lemma 6.2 (a).

Let $\sigma \in \Gamma$. Denote by $\sigma(\xi)$ the partition $(\sigma\Xi_1) \sqcup \dots \sqcup (\sigma\Xi_k)$ such that $\sigma\Xi_j = \{\sigma(i) \mid i \in \Xi_j\}$, and for a function $\beta(z_1, \dots, z_l)$ set $\sigma(\beta) = \beta(z_{\sigma(1)}, \dots, z_{\sigma(l)})$. This implies (6.6), since α is Γ -symmetric.

Combining all this together, we see that $f(P + C + E_1) = 0$ if and only if for any partition $\xi = (\Xi_1, \dots, \Xi_k)$ all terms $\alpha_i^{(j)}$ in (6.3) are regular admissible and have $\tau(\alpha_i^{(j)}) > 1$, so Claim A follows. \square

Let us now look at the condition $f((E_0 + E_1) \cap (D^*F_1 + E_1)) = 0$. It is equivalent to the existence of a functional $g : F_0/E_1 \rightarrow \mathbb{k}$ such that

$$g(D^*F_1 + E_1) = 0 \quad \text{and} \quad g|_{E_0+E_1} = f|_{E_0+E_1}.$$

If β is the correlation function of g , then β must be regular admissible, such that $\tau(\beta) > 1$ and $\beta(\xi) = \alpha(\xi)$ for any partition ξ . Moreover, if we find such a function β , then it should correspond to a desired functional g .

Denote by Π the set of all *unordered* partitions of $\{1, \dots, l\}$. Note that if $\xi \in \Pi$ the expression $\alpha(\xi)$ makes sense, since $f(C) = 0$. For a partition $\xi \in \Pi$ let $|\xi|$ be the number of parts in ξ and let $\Pi_k = \{\xi \in \Pi \mid |\xi| = k\}$.

We claim that the following function β satisfies all our requirements:

$$\beta = \sum_{\xi \in \Pi} (-1)^{|\xi|} (|\xi| - 1)! \alpha(\xi). \quad (6.7)$$

Using the properties of $\alpha(\xi)$ established in Claim A, we see that the only thing that remains to be shown is

Claim B. Let $\beta \in \Phi_{-2l}^\Gamma$ be given by (6.7). Then $\beta(\xi) = \alpha(\xi)$ for any partition $\xi \in \Pi$.

Proof. For $\xi, \theta \in \Pi$ write $\xi < \theta$ if θ is obtained from ξ by dividing parts of ξ into smaller subsets. It is enough to prove Claim B for $\xi \in \Pi_2$, since if $|\theta| > 2$, then $\theta > \xi$ for some $\xi \in \Pi_2$ and then $\alpha(\theta) = \alpha(\xi)(\theta) = \beta(\xi)(\theta) = \beta(\theta)$.

So let $\xi = \{\Xi_1, \Xi_2\} \in \Pi_2$. Denote by $\Pi_\xi(m_1, m_2) \subset \Pi_{m_1+m_2}$ the set of all partitions that can be obtained from ξ by dividing Ξ_i into m_i parts, $i = 1, 2$. Note that $\Pi_\xi(1, 1) = \{\xi\}$. Fix a partition $\theta \in \Pi_\xi(m_1, m_2)$ and an integer $k \geq 2$ and set $n = m_1 + m_2 - k$. We claim that

$$|\{\pi \in \Pi_k \mid \pi\xi = \theta\}| = \begin{cases} n! \binom{m_1}{n} \binom{m_2}{n} & \text{if } n \geq 0, \\ 0 & \text{if } n < 0. \end{cases}$$

Indeed, let $\theta = \{\Theta_1, \dots, \Theta_{m_1+m_2}\}$. Set $\theta_i = \{\Theta_j \mid \Theta_j \subseteq \Xi_i\}$, so that $|\theta_i| = m_i$ for $i = 1, 2$. Then the set $\{\pi \in \Pi_k \mid \pi\xi = \theta\}$ is in one-to-one correspondence with the set of all n -tuples of pairs

$$(\Theta_{i_1}, \Theta_{j_1}), \dots, (\Theta_{i_n}, \Theta_{j_n}),$$

such that $\Theta_{i_s} \in \theta_1$, $\Theta_{j_s} \in \theta_2$ and all Θ_{i_s} and Θ_{j_s} are pairwise different. This is because to every such n -tuple we can correspond the partition

$$\pi = (\theta \setminus \{\Theta_{i_s}, \Theta_{j_s}\}_{s=1}^n) \cup \{\Theta_{i_s} \cup \Theta_{j_s}\}_{s=1}^n \in \Pi_k,$$

obtained by joining together the sets Θ_{i_s} and Θ_{j_s} for $s = 1, \dots, n$. Clearly, the set of of these n -tuples has the required cardinality.

Now we compute, using Lemma 6.2 (b),

$$\begin{aligned} \beta(\xi) &= \sum_{k \geq 2} (-1)^k (k-1)! \left(\sum_{\pi \in \Pi_k} \alpha(\pi) \right) (\xi) \\ &= \sum_{k \geq 2} (-1)^k (k-1)! \sum_{\pi \in \Pi_k} \alpha(\pi\xi) \\ &= \sum_{\substack{m_1, m_2 \geq 1 \\ k \geq 2}} (-1)^k (k-1)! (m_1 + m_2 - k)! \binom{m_1}{k-m_2} \binom{m_2}{k-m_1} \sum_{\theta \in \Pi_\xi(m_1, m_2)} \alpha(\theta) \\ &= \alpha(\xi) + \sum_{\substack{m_1+m_2 \geq 3 \\ k \geq 2}} (-1)^k (k-1)! (m_1 + m_2 - k)! \binom{m_1}{k-m_2} \binom{m_2}{k-m_1} \sum_{\theta \in \Pi_\xi(m_1, m_2)} \alpha(\theta) \end{aligned}$$

This is equal to $\alpha(\xi)$ if we show that

$$\sum_{k=\max(m_1, m_2)}^{m_1+m_2} (-1)^k (k-1)! (m_1 + m_2 - k)! \binom{m_1}{k-m_2} \binom{m_2}{k-m_1} = 0$$

for any $m_1, m_2 \geq 0$. But

$$(k-1)! (m_1 + m_2 - k)! \binom{m_1}{k-m_2} \binom{m_2}{k-m_1} = \frac{(m_1!)^2 (m_2!)^2}{(m_1 + m_2)!} \frac{1}{k} \binom{k}{m_1} \binom{k}{m_2} \binom{m_1 + m_2}{k},$$

and $k^{-1} \binom{k}{m_1} \binom{k}{m_2}$ is a polynomial in k of degree $m_1 + m_2 - 1$, so the statement follows from the fact that $\sum_k (-1)^k k^s \binom{n}{k} = 0$ for any $s < n$. \square

6.5. Proof of Lemma 3.6. Let $\lambda = la + lb \in \mathbb{Z}_+[\mathcal{A}]$ for some $a \neq b \in \mathcal{A}$ and $l \geq 4$. Consider the $2l \times 2l$ matrix $S = \begin{pmatrix} 0 & R \\ R^T & 0 \end{pmatrix}$, where $R = \{r_{ij}\}_{i,j=1}^l$ is given by

$$r_{ij} = \begin{cases} -1 & \text{if } i \leq j \leq i+3 \pmod{l}, \\ 0 & \text{otherwise.} \end{cases}$$

We can view S as an adjacency matrix of a bipartite graph G drawn on Figure 1, that has l “even” vertices v_1, \dots, v_l drawn on the bottom row, and l “odd” vertices v_{l+1}, \dots, v_{2l} , drawn on the top row. An edge connects v_i with v_j in G if and only if $s_{ij} = -1$.

The group $\Gamma \subset S_{2l}$ in this case is equal to $S_l \times S_l$. Let $\alpha = \alpha(z_1, \dots, z_{2l}) = \sum_{\sigma \in \Gamma} \sigma \pi(S)$, where $\pi(S)$ is defined in (5.2). Need to prove the following fact:

Claim.

- (a) $0 \neq \alpha \in \Phi_{-4l}^\Gamma$ is a regular admissible function.
- (b) For any partition $\xi = (\Xi_1, \Xi_2)$ of $\{1, \dots, 2l\}$ in two parts, we have $\alpha_{n,i}^{(2)} = 0$ in (6.2) for $n < -2|\Xi_2| + 2$ and all i .

Indeed, it would follow then that the non-zero functional $f : F_{\lambda,0} \rightarrow \mathbb{k}$, corresponding to α is equal to zero on $D^*F_{\lambda,1}$ (since α is regular) and on $E_0 + E_1$, since it will be zero on every word w that has a tail of degree less than 2.

Proof. The function α is regular by § 5.2. Note that for any $\sigma \in S_{2l}$, we have $\sigma\pi(S) = \pi(\sigma S)$, where σ acts on S by permuting both rows and columns. Therefore, for any $\sigma \in \Gamma$ we have

$$\sigma\pi(S)|_{z_{l+1}=\dots=z_{2l}=0} = z_1^{-4} \cdots z_l^{-4},$$

hence $\alpha(z_1, \dots, z_l, 0, \dots, 0) = |\Gamma| z_1^{-4} \cdots z_l^{-4} \neq 0$. This proves (a).

To prove (b), it is enough to show that this property holds for $\alpha = \sigma\pi(S)$ for every $\sigma \in \Gamma$. Since $\sigma\pi(S) = \pi(\sigma S)$ for any $\sigma \in \Gamma$, it is sufficient to show that the property (b) holds for $\alpha = \pi(S)$.

For a subset $\Xi \subset \{1, \dots, 2l\}$ let S_Ξ be the minor of S consisting of the rows and columns with indices in Ξ . Suppose that $\Xi_1 = \{i_1, \dots, i_m\}$. Then the term of minimal degree in the decomposition (6.2) of $\pi(S)$ is

$$z_{i_1}^{k_1} \cdots z_{i_m}^{k_m} \pi(S_{\Xi_1}) \pi(S_{\Xi_2}).$$

So we need to show that $\deg \pi(S_\Xi) \geq -2|\Xi| + 2$ for any $\Xi \subset \{1, \dots, 2l\}$.

This follows from the following property of the corresponding graph G : for every partition of vertices into two subsets, there are at least 4 edges that have vertices in different subsets. This is true because G has two Hamiltonian paths with no common edges. They are marked by the solid and the dashed lines on Figure 1.

□




FIGURE 1. The graph G . The upper and lower rows of vertices are periodic with period l .

REFERENCES

- [1] G. M. Bergman. The Diamond lemma for ring theory. *Adv. in Math.*, 29(2):178–218, 1978.
- [2] L. A. Bokut'. Imbeddings into simple associative algebras. *Algebra i Logika*, 15(2):117–142, 245, 1976.
- [3] R. E. Borcherds. Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. U.S.A.*, 83(10):3068–3071, 1986.
- [4] B. Buchberger. Ein algorithmisches Kriterium für die Lösbarkeit eines algebraischen Gleichungssystems. *Aequationes Math.*, 4:374–383, 1970. English translation in *Gröbner bases and applications (Linz, 1998)*, London Math. Soc. Lecture Note Ser., 251:535–545, Cambridge Univ. Press, Cambridge, 1998.
- [5] C. Dong. Representations of the Moonshine module vertex operator algebra. In *Mathematical aspects of conformal and topological field theories and quantum groups (South Hadley, MA, 1992)*, volume 175 of *Contemp. Math.*, pages 27–36. Amer. Math. Soc., Providence, RI, 1994.
- [6] C. Dong, H. Li, G. Mason, and P. S. Montague. The radical of a vertex operator algebra. In *The Monster and Lie algebras (Columbus, OH, 1996)*, volume 7 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 17–25. de Gruyter, Berlin, 1998.
- [7] C. Dong, Z. Lin, and G. Mason. On vertex operator algebras as \mathfrak{sl}_2 -modules. In *Groups, difference sets, and the Monster (Columbus, OH, 1993)*, volume 4 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 349–362. de Gruyter, Berlin, 1996.
- [8] I. B. Frenkel, Y.-Z. Huang, and J. Lepowsky. On axiomatic approaches to vertex operator algebras and modules. *Mem. Amer. Math. Soc.*, 104(494):viii+64, 1993.
- [9] I. B. Frenkel, J. Lepowsky, and A. Meurman. *Vertex Operator Algebras and the Monster*. Academic Press, Boston, MA, 1988.
- [10] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66(1):123–168, 1992.
- [11] R. L. Griess, Jr. The Friendly Giant. *Invent. Math.*, 69(1):1–102, 1982.
- [12] R. L. Griess, Jr. GNAVOA. I. Studies in groups, nonassociative algebras and vertex operator algebras. In *Vertex operator algebras in mathematics and physics (Toronto, ON, 2000)*, volume 39 of *Fields Inst. Commun.*, pages 71–88. Amer. Math. Soc., Providence, RI, 2003.
- [13] V. G. Kac. *Vertex Algebras for Beginners*, volume 10 of *University Lecture Series*. AMS, Providence, RI, second edition, 1998.

- [14] C. H. Lam. Construction of vertex operator algebras from commutative associative algebras. *Comm. Algebra*, 24(14):4339–4360, 1996.
- [15] C. H. Lam. On VOA associated with special Jordan algebras. *Comm. Algebra*, 27(4):1665–1681, 1999.
- [16] J. Lepowsky and H. Li. *Introduction to vertex operator algebras and their representations*, volume 227 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2004.
- [17] H. Li. Symmetric invariant bilinear forms on vertex operator algebras. *J. Pure Appl. Algebra*, 96(3):279–297, 1994.
- [18] H. Li. An analogue of the Hom functor and a generalized nuclear democracy theorem. *Duke Math. J.*, 93(1):73–114, 1998.
- [19] M. Roitman. On free conformal and vertex algebras. *J. Algebra*, 217(2):496–527, 1999.
- [20] M. Roitman. Combinatorics of free vertex algebras. *J. Algebra*, 255(2):297–323, 2002.
- [21] M. Roitman. Invariant bilinear forms on a vertex algebra, 2003. To appear.
- [22] A. I. Shirshov. Some algorithmic problems for Lie algebras. *Sibirsk. Mat. Ž.*, 3:292–296, 1962. English translation in *ACM SIGSAM Bull.*, 33(2):3–6, 1999.
- [23] W. Wang. Rationality of Virasoro vertex operator algebras. *Internat. Math. Res. Notices*, no. 7:197–211, 1993.
- [24] X. Xu. *Introduction to vertex operator superalgebras and their modules*, volume 456 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 1998.

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, ANN ARBOR, MI 48109-1109
E-mail address: roitman@umich.edu