

Selection Criterion for Log-Linear Models Using Statistical Learning Theory

Daniel J. L. Herrmann

Max-Planck-Institut für biologische Kybernetik
Spemannstr. 38, D-72076 Tübingen, Germany
daniel.herrmann@tuebingen.mpg.de

Dominik Janzing

Institut für Algorithmen und Kognitive Systeme
Universität Karlsruhe, Am Fasanengarten 5
D-76131 Karlsruhe, Germany
janzing@ira.uka.de

February 7, 2003

Abstract

Log-linear models are a well-established method for describing statistical dependencies among a set of n random variables. The observed frequencies of the n -tuples are explained by a joint probability such that its logarithm is a sum of functions, where each function depends on as few variables as possible. We obtain for this class a new model selection criterion using nonasymptotic concepts of statistical learning theory. We calculate the VC dimension for the class of k -factor log-linear models. In this way we are not only able to select the model with the appropriate complexity, but obtain also statements on the reliability of the estimated probability distribution. Furthermore we show that the selection of the best model among a set of models with the same complexity can be written as a convex optimization problem.

Key words: Markov network, AIC, non-asymptotic statistic, VC dimension structural risk minimization, convex optimization

1 INTRODUCTION

Suppose a scientist is interested in the relation between n features described by the random variables X_1, X_2, \dots, X_n . Observing dependencies among these features is an important part of all scientific disciplines. The dependencies may be deterministic, e.g.,

$$X_n = f(X_1, \dots, X_{n-1})$$

or (in the generic case) statistical, i.e., the probability for the event $X_1 = x_1, X_2 = x_2, \dots, X_n = x_n$, for short $\mathbf{X} = \mathbf{x}$, is not the product of the probabilities of the n events $X_j = x_j$. All information about these dependencies is contained in the joint probability distribution P that assigns the probability $P(\mathbf{x}) = P(x_1, x_2, \dots, x_n)$ to each n -tuple. Notice that for large n no “reasonable” sample size is sufficient to determine the probabilities of all $P(x_1, x_2, \dots, x_n)$ with good reliability since the size of $\Omega := \times_j \Omega_j$ (where each Ω_j denotes the set of possible values x_j) is exponentially large in n . This shows that we should not try to learn the joint probability distribution, we rather have to develop inference rules for learning the statistical dependencies *in a weaker sense*. The task is therefore to estimate some properties of the joint distribution described by empirical data of a given size.

In the last years methods to judge scientific theories by their predicted models on real world data sets become more and more important (Hagenaars 1994, Ishii-Kuntz 1997, Pitt and Myung 2002) and since the accessible computational power is increasing strongly this trend will proceed in future.

It is a well-known fact, see e.g. (Akaike 1973, Hansen and Yu 2001), that criteria describing the goodness of fit for the model to the data set only are not enough to judge the scientific relevance of the model. Additionally one has to take the complexity of the model into account. Usually the number of free parameters are treated as a measure for the model complexity and are incorporated in criteria like in Akaike’s Information Criterion (AIC) (Akaike 1973), or Bayesian Information Criterion of Schwarz (1978). Bayesian Model Selection (Kass and Raftery 1995) and Minimum Description Length (Rissanen 1996, Hansen and Yu 2001) take also the functional form of the model

into account. However, these criteria do not give any statements on the reliability of the estimation.

In this article we derive reliability statements from statistical learning theory (Vapnik 1998) and show that the trade-off between goodness of fit and model complexity can be treated with structural risk minimization.

Let us describe roughly the link between the ideas presented here and the question of minimal description length in information theory, see (Barron 1991, Rissanen 1996) and references therein. In the latter approach a risk functional is minimized over a set of probability distributions where each probability distribution is penalized according to its description length. More explicitly, the risk functional consists of two terms: the “empirical risk” evaluates the “goodness of fit” and a regularization term penalizes the code length. Barron and Cover (1991) obtain bounds on the convergence rate of the risk functional in the Hellinger distance. The difference to our approach lies in the fact the regularization function can in principle be chosen arbitrarily, only Kraft’s inequality must hold; whereas we use a penalty term coming directly from the complexity of the considered class of log-linear models and is based on statistical learning theory. Vapnik and Chervonenkis (1971) addressed the question of model selection from empirical data in a probabilistic setting (see also Vapnik (1999) and references therein). They introduced the so-called VC dimension of a function set and showed that the VC dimension of the function set which the model can implement is a crucial quantity to describe its complexity. In general the VC dimension does not agree with the usual dimension or the number of free parameters.

From the statistical learning point of view, a scientific theory delivers a prior on the class of models under consideration. Then the model is chosen according to some criterion measuring the fit on the empirical data. In this work we consider the class of all log-linear models and prior models with few statistical dependencies among the random variables, i.e. we prefer less complex models in terms of statistical independency as long as they describe the data well enough. With this assumption we estimate the model (complexity) directly from the empirical data rather than testing one model against another using test methods like the Pearson or likelihood ratio chi-square criterion, see e.g. Christensen 1997, Goodman 1978. Akaike (1973) already formulated this view point, however he did not have the conceptual tools of learning theory (Vapnik 1995).

The article is organized as follows. In Section 2 we introduce log-linear models and explain the prior for model selection leading to a risk functional on the data. We discuss the commonly used model selection criteria and explain their shortcomings. We present the idea of Markov networks and relate them to log-linear models. In Section 3.1 we calculate the VC dimension of the models under consideration and give the central result about the estimation of the risk functional for the true probability distribution. In Section 3.2 we use the idea of the structural risk minimization to formulate the model estimation for log-linear models. In Section 3.3 we show that the optimal model is the solution of a convex optimization problem.

2 LOG-LINEAR MODELS

2.1 The Empirical Risk Functional

In the work presented here we restrict our attention to discrete random variables and assume that \mathbf{X} takes values in a finite set $\Omega = \times_{j=1}^n \Omega_j$. Assume a scientist tells us that he has found that the true joint probability distribution P is given by P_t and we would like to test whether he is right or not. Suppose that his model is given by the distribution that all n -tuples in Ω occur with equal probability. Assume, in contrast, that the *true* distribution P assigns the probability $1/k$ to k specific n -tuples $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in \Omega$ and that this k n -tuples are chosen without any simple law. Assume furthermore that k is large compared to the available sample size but small compared to the size of Ω . Then we have only little chance to recognize that his model is wrong. Only if the k n -tuples are selected by a simple law we would mistrust the scientist's hypothesis. This example illustrates that we need a quality criterion for models which can be tested on a reasonable sample size. We suggest to use following criterion. For each n tuple \mathbf{x} in our data set we give penalty points depending on how likely its appearance was according to the hypothetical joint distribution. A good choice, for instance, is the negative logarithm of the hypothetical probability $P_t(\mathbf{x})$. Let T be the training set, then we obtain

$$-\frac{1}{\#T} \sum_{\mathbf{x} \in T} \ln P_t(\mathbf{x})$$

as the “penalty” for the model $P_t(\mathbf{x})$. Notice that this sum converges in the large sample size limit to

$$R(P_t) := - \sum_{\mathbf{x} \in \Omega} P(\mathbf{x}) \ln P_t(\mathbf{x}). \quad (1)$$

Note that $R(P_t)$ is closely related to the Kullback-Leibler distance (Cover and Thomas 1991) and can never be smaller than the entropy

$$S(P) := - \sum_{\mathbf{x} \in \Omega} P(\mathbf{x}) \ln P(\mathbf{x}).$$

Hence, even if the scientist has predicted the true measure exactly, he will never get less than $S(P)$ penalty points. If $S(P)$ is high, he is not punished for building *wrong* models but for selecting the *wrong* features, i.e. features with too few dependencies. If the scientist considers additional random variables (“features influencing the considered ones”) more specific statements might be possible. Reasonable research consists not only in observing dependencies among a given set of features, it consists also in *selecting* statistically relevant features yielding a low entropy of the joint distribution. Therefore the “empirical risk functional”

$$R_{emp}(P_t) := - \frac{1}{\#T} \sum_{\mathbf{x} \in T} \ln P_t(\mathbf{x}) \quad (2)$$

is a good measure to *test* the quality of the distribution P_t for large sample size. This risk functional is well-known for estimating probabilities (Vapnik 1998).

If the space of all probability distributions is too large compared to the sample size, we should not minimize the empirical risk over all joint distributions. In order to infer a probability distribution from the statistical data we give the space a class structure. We use log-linear models (Christensen 1997, Goodman 1978) to define suitable classes. The simplest class of log-linear models is given in the case when all random variables are statistically independent. Then the logarithm of the joint probability distribution can be written as a sum of functions, each function depends on one variable only

$$\ln P_t(x_1, x_2, \dots, x_n) = \sum_{j=1}^n f_j(x_j),$$

with $f_j(x_j) = \ln P(x_j)$. The next higher class is given by allowing two-variable interaction terms

$$\ln P_t(x_1, \dots, x_n) = \sum_{j \leq n} f_j(x_j) + \sum_{i, j \leq n} g_{i,j}(x_i, x_j).$$

The next higher class contains terms with three-variable interactions, etc.. Probability distributions with the property that their logarithm can be expressed as sum of functions with k variables are called k -factor distributions. In other words the data is described by a k -factor *log-linear model*. The idea of log-linear models come from the fact that some variables influence each other *directly*, expressed by the interaction terms, whereas the other dependencies are caused *indirectly* by influencing intermediate variables. The hope that real world data are well explained by log-linear models can be backed up by the idea that each variable is influenced only by a few other variables directly. The graph of the variables with edges representing the direct influences is simple. This idea can be formulated using *Markov networks*, see Section 2.3.

2.2 Model Selection and AIC

A simple way of choosing the degree of the interaction terms in a log-linear model goes as follows. We begin with the simplest model class and decide on the basis of some significance test whether the data suggest to reject the model or not. If so then we include two-variable interactions. The same significance test is applied to the new model. The procedure ends if frequencies observed in the data and the probabilities given by the model do not differ significantly according to the significance test. Let us describe two popular significance tests (Christensen 1997, Goodman 1978).

Let $j \in \{1, \dots, k\}$ be the probability space consisting of k possible events. Let p_1, \dots, p_k be the probabilities of the model and m_1, \dots, m_k the observed frequencies. The sum $l := \sum_j m_j$ is the sample size. Then we calculate

1. Pearson chi-square Test:

$$X^2 := \sum_j \frac{(m_j - lp_j)^2}{lp_j}$$

2. Deviance Test:

$$G^2 := 2 \sum_j m_j \ln \frac{m_j}{lp_j}.$$

Remark: $G^2/(2l)$ is the Kullback-Leibler distance (Cover and Thomas 1991) between the probability distribution $(p_j)_{j \leq k}$ and the relative frequencies $(m_j/l)_{j \leq k}$. Furthermore the minimizer of G^2 and R_{emp} are the same, i.e., $G^2/(2l) - R_{emp}$ does not depend on $(p_j)_{j \leq k}$.

Under the assumption that $(p_j)_{j \leq k}$ is the true probability distribution the values of the random variables X^2 and G^2 are in the large sample limit χ^2 -distributed (Christensen 1997). Using this argument the model is rejected if the X^2 or G^2 values are too large. The range of acceptable values for X^2 and G^2 is based on the following rule:

First we have to calculate the number of degrees of freedom (df) of the model. (df) is given by the number of possible events minus 1 minus the number of free parameters in the model class. Given df one can look up in a table with χ^2 distribution whether the probability that X^2 or G^2 , respectively, is outside a certain confidence interval.

However, the idea of rejecting a simple model only if the data contradicts the externally given significance level leads to conservative models. Suppose the outcome of the chi-squared test of a model yields values of χ^2 or G^2 within a certain confidence interval. Nevertheless there might be a model with only one interaction term more such that the χ^2 or G^2 value is considerably decreased. In this case we will certainly prefer the model that is *a little bit* more complex. This example illustrates that the externally given significance level implicitly determines the model selection. Therefore it is crucial to find a more systematic way for model selection and prevent in this way over- and underfitting.

Let $(P_\theta)_{\theta \in \mathbb{R}^k}$ be a set of probability distributions with k parameters and assume that the true probability distribution P lies in $(P_\theta)_{\theta \in \mathbb{R}^k}$. The true probability distribution P is the minimizer of R and for a sample size which is large compared to the size of $(P_\theta)_{\theta \in \mathbb{R}^k}$, the empirical risk functional R_{emp} is a good approximation for R . In this case we can search for the minimizer of R_{emp} instead. This leads to the Deviance test above. However we are interested in the case where we cannot assure that the data set is sufficiently large. Then the minimizer of R_{emp} is in general not a good estimation for the minimizer of R .

Akaike (1973) suggested to modify the empirical risk functional according to the number of free parameters of the model under consideration. He showed under some regularity conditions on the mapping $\theta \mapsto P_\theta$, that

$$R_{emp} + df/l$$

is a consistent estimator of R , i.e., it has the same expectation value as R . However minimizing $R_{emp} + df/l$ for a finite data set give us no information about the true risk R . In Section 3.1 we obtain some bounds on the true risk and suggest to minimize this *guaranteed* risk rather than the *expected* risk, see Section 3.1.

2.3 Why Log-linear models

In Section 2 we have introduced log-linear model classes without giving any justification for it. In this section we explain why one should expect that probability distributions generating real world data can be described by k -factor models with small k . The key idea is that the distribution of a random variable X_j is only influenced by the values a few others variables. This kind of statistical relations can be encoded by a Markov network.

The presentation of Markov networks in this work follows Pearl (1988). For each random variable X_i we define the Markov boundary $\mathcal{B}_i \subset \{X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n\}$ as the smallest set fulfilling

$$P(x_i | x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) = P(x_i | \mathcal{B}_i). \quad (3)$$

Observe that this definition does not take any (temporal or causal) order of the random variables into account. It will therefore lead to an undirected graph. This is in contrast to the directed graphs in the context of Bayesian networks (Pearl 1985) which are for instance useful to formalize causal structure (Pearl 2000).

In case of strict positive probability distributions the Markov boundary is unique. Actually, strict positivity of the probability distributions is also a crucial condition for our approach to learn probabilities as we will see in Section 3.1. We now define the Markov network as the graph \mathcal{G} with n nodes also denoted by X_1, \dots, X_n and undirected edges between X_i and X_j if and only if $X_j \in \mathcal{B}_i$ (or equivalently $X_i \in \mathcal{B}_j$). In other words, we can generate the Markov network for a distribution P if we connect each of the n nodes with the element of its Markov boundary. As a consequence we obtain that two random

variables X_i and X_j , $i \neq j$ are conditionally independent with respect to a set $\mathbf{Z} \subset \mathbf{X} \setminus \{X_i, X_j\}$, i.e.

$$P(X_i|X_j, \mathbf{Z}) = P(X_i|\mathbf{Z}),$$

if and only if every path from X_i to X_j has one node in \mathbf{Z} . In other words, if \mathbf{Z} *blocks* every path between X_i and X_j then X_i and X_j are conditionally independent (with respect to \mathbf{Z}). A *clique* in the graph \mathcal{G} is a subgraph in which all nodes are connected to each other. We have an (partial) order on the cliques given by the inclusion as sets.

Let \mathcal{G} be a Markov graph for a strictly positive probability distribution P and let C be the set of maximal cliques in \mathcal{G} . Then one can show that P factorizes as follows

$$P(\mathbf{x}) = \frac{1}{N} \prod_{c \in C} \phi_c(\mathbf{x}_c), \quad (4)$$

where $\mathbf{X}_c \subset \mathbf{X}$ is the set of random variables in the cliques c . N is the normalization constant and ϕ_c are the *compatibility functions* w.r.t. the clique c , also called the potential functions.

Suppose two distributions have the same Markov graph, then they describe the same (in)dependency in \mathbf{X} . In Section 3.1 we will see that the maximal number in a clique $\deg(\mathcal{G}) = \max_{c \in C} \#c$ are appropriate parameters to subdivide the set of all distributions $\mathcal{P}(\mathbf{X})$ into *small* model classes.

Consider the case that every clique has less than $k + 1$ elements. Then the distribution can be described by a k -factor model due to the factorization in eq. (4). The Markov boundary can be understood in a rather literary sense if the variables represent some quantities that are measured at different positions in the real space. Then the belief to obtain probability distributions that correspond to simple Markov networks stems from the locality principle: distant variables influence each other only indirectly via other variables.

In statistical physics log-linear models can be justified even more directly. Assume the variable X_j describes the physical state of particle j and that the total energy of the system in the state $\mathbf{x} = (x_1, \dots, x_n)$ is a sum of functions depending on at most k particles, i.e.,

$$E(\mathbf{x}) = \sum_{\mathbf{j}} f_{\mathbf{j}}(x_{j_1}, \dots, x_{j_k}),$$

where \mathbf{j} runs over all k -subsets in $\{1, \dots, n\}$. Then the thermodynamic equilibrium state, the so-called Gibbs distribution, is up to a

normalization factor directly given by the exponential of the negative energy function. Hence the equilibrium distribution is exactly described by a k -factor model. In statistical physics one usually considers two particle interactions only, therefore 2-factor models describe the equilibrium state.

Now we have justified why the class structure of log-linear models is natural. However, we have to refine our hierarchy in some respect. So far, we did not formulate any restrictions on the range of the probability distribution of the models. Intuitively it is clear that we can assign a small probability to an event $\mathbf{x} \in \Omega$ only if we have a very large data set. This means that reasonable model selection should allow small probabilities only for large data sets. However, the empirical risk functional R_{emp} penalizes for small probabilities only if they are contained in the data set. Therefore the value of R_{emp} depends strongly on the specific data set if the probability distribution is not bounded from below by a strict positive constant. Actually, for deriving a bound for the risk functional R in Section 3.1 we precisely need to bound the probability in this way.

Definition 1 *Let \mathcal{P}_λ be the set of all strict positive functions greater than $\lambda > 0$. Let \mathcal{P}^k be the set of probability distributions P corresponding to a k -factor model, i.e.,*

$$P(x_1, x_2, \dots, x_n) = \prod_{\mathbf{j}} q_{\mathbf{j}}(x_{j_1}, x_{j_2}, \dots, x_{j_s}),$$

where each $q_{\mathbf{j}}$ is a positive function and $\mathbf{j} := (j_1, j_2, \dots, j_s)$ runs over all possible subsets of $\{1, 2, \dots, n\}$ with s elements. Then we define the model class of degree k by

$$\mathcal{H}_k^\lambda = \mathcal{P}_\lambda(\mathbf{X}) \cap \mathcal{P}^k. \quad (5)$$

Remark: The values $q_{\mathbf{j}}(x_{j_1}, x_{j_2}, \dots, x_{j_s})$ cannot be interpreted as probabilities since they can be greater than 1.

the

3 LEARNING LOG-LINEAR MODELS

It is a well-known problem in any mathematical theory of learning that a model can explain the training data well but does not fit the

data observed in the future if the model class is too large. If the model class is small enough, one knows with high confidence that the model which minimizes the error on the training data will also explain the future data almost optimal within this model set.

3.1 Risk estimation

For a set of real-valued functions on an arbitrary set Ω a relevant measure for the size of this set is the so-called VC dimension (Vapnik-Chervonenkis). First, let us define the VC dimension (Vapnik 1998) of a set of indicator functions (f_α) , i.e. $f_\alpha : \Omega \rightarrow \{0, 1\}$.

Definition 2 *Let Λ be an index set of arbitrary cardinality. Let $(f_\alpha)_{\alpha \in \Lambda}$ be a set of indicator functions on Ω . Then the VC dimension of $(f_\alpha)_{\alpha \in \Lambda}$ is the largest number h such that there exist h points $x_1, x_2, \dots, x_h \in \Omega$ with the property that for every function $\chi : \{x_1, \dots, x_h\} \rightarrow \{0, 1\}$ there exists a function f_α such that its restriction to $\{x_1, x_2, \dots, x_h\}$ coincides with χ .*

The VC dimension for a set of real-valued functions is defined by the VC dimension for the set of corresponding indicator functions.

Definition 3 *Let $(f_\alpha)_{\alpha \in \Lambda}$ be a family of real-valued functions on a set Ω . Then the VC dimension of $(f_\alpha)_{\alpha \in \Lambda}$ is the VC dimension of the family of the indicator functions $(\chi_\mu \circ f_\alpha)_{\mu \in \mathbb{R}, \alpha \in \Lambda}$, where χ_μ is $\chi_\mu(x) = 0$ for $x < \mu$ and $\chi_\mu(x) = 1$ for $x \geq \mu$.*

The notion of VC dimension plays a crucial role in statistical learning theory. It is a measure which indicates whether a learning machine overfits or not. Roughly speaking, if a family of functions which can be implemented by a learning machine has small VC dimension then every function which fits well the training data will fit with high probability the test data as well. The following theorem is an Corollary of Theorem 5.1 pp.192 in (Vapnik 1998).

Theorem 4 *Let $(f_\alpha)_{\alpha \in \Lambda}$ be a measurable set of bounded real-valued functions on Ω , $A \leq f_\alpha(z) \leq B$, and let the set of indicator functions have finite VC dimension h . Let μ be a probability distribution on $\Omega \times \mathbb{R}$ and let the data $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_l$ be drawn according to μ (independent*

and identically distributed). Then we have with probability at least $1 - \eta$

$$\tilde{R}(f_\alpha) \leq \tilde{R}_{emp}(f_\alpha) + (B - A) \sqrt{\frac{h - \ln h + \ln 16l - \ln \eta}{l}},$$

where $\tilde{R}_{emp}(f_\alpha) := \frac{1}{l} \sum_{j \leq l} f_\alpha(\mathbf{x}_j)$ is the empirical risk and $\tilde{R}(f_\alpha) := \int f_\alpha(\mathbf{x}) d\mu(\mathbf{x})$ the true risk.

Remark: We have $R(P_\alpha) = \tilde{R}(\ln P_\alpha)$ and $R_{emp}(P_\alpha) = \tilde{R}_{emp}(\ln P_\alpha)$ when (P_α) is a set of probability distributions. Therefore we can use this theorem to calculate the generalization error of equation (1) for the family of probability distributions defined in equation (5).

First, let us interpret the generalization error in this context. Suppose we have found a probability distribution P_t in some model class that minimizes R then we know that in this space P_t is the closest distribution to the true distribution in the Kullback-Leibler distance. Furthermore, we can conclude with high confidence that future data will fit to this distribution as well.

Next we estimate the VC dimension of product distributions since this will be the leading intuition for estimating the VC dimension of the model class \mathcal{H}_k^λ .

Lemma 1 *Let X_1, X_2, \dots, X_n be discrete random variables where X_i takes m_i different values in Ω_i for $i = 1, \dots, n$. Then the VC dimension of the family of all product distributions $\mathcal{P}^1 = \{P(x_1, \dots, x_n) = P(x_1) \cdot \dots \cdot P(x_n) : P \text{ is a probability distribution}\}$ is given by $N = 1 + \sum_{i=1}^n m_i - n$.*

Remark: For the families $(\ln P)_{P \in \mathcal{P}^1}$ and $(P)_{P \in \mathcal{P}^1}$ the set of indicator functions coincides. Hence $(\ln P)_{P \in \mathcal{P}^1}$ and $(P)_{P \in \mathcal{P}^1}$ have the same VC dimension.

Proof: Let $x_{j;0}$ be an arbitrary element of the set Ω_j . Due to $P(\mathbf{x}) = P(x_1) \cdot \dots \cdot P(x_n)$ we can write the logarithm of the joint probability as

$$\ln P(\mathbf{x}) = \sum_j \ln(P(x_j)/P(x_{j;0})) + \sum_j \ln P(x_{j;0})$$

We can characterize an n -tuple $\mathbf{x} = (x_1, x_2, \dots, x_n)$ uniquely by an $N = \sum_j (m_j - 1)$ dimensional vector with entries 0 and 1 as follows:

The vector consists of n blocks of dimension $(m_j - 1)$ characterizing x_j . Let $x_{j;0}, x_{j;1}, \dots, x_{j;m_j-1}$ be the elements of Ω_j in an arbitrary ordering. For $x_j = x_{j;i}$ with $i \neq 0$ let the i -th entry of the j -th block be 1 and the other entries of the j -th block be zero. For $x_j = x_{j;0}$ let all entries of the j -th block be zero.

For each n -tuple the logarithm of its probability is given as follows. Let $c^{\mathbf{x}} \in \mathbb{R}^N$ be the vector corresponding to $\mathbf{x} := (x_1, x_2, \dots, x_n)$. Then we have

$$\ln P(\mathbf{x}) = \sum_l c_l^{\mathbf{x}} f_l + d = \langle c^{\mathbf{x}}, f \rangle + d$$

where the coordinates f_l of f are given by the values $\ln(P(x_{j;i})/P(x_{j;0}))$ with $j \leq n$ and $1 \leq i \leq m_j - 1$. The constant term d is defined as $d := \sum_j \ln P(x_{j;0})$. This shows that $\ln P(\mathbf{x})$ can be written as an affine functional in \mathbb{R}^N . Hence the VC dimension of \mathcal{P} is at most $N + 1$ (see (Vapnik 1998), Example in Chapter 5.2.3). To show that it is not smaller than $N + 1$ note that there is no restriction at all to the set of all possible vectors f if d is chosen appropriately. This can be seen as follows. Let $g_1, g_2, \dots, g_{m_j-1}$ be the entries of f in the j -th block. Then set

$$P(x_{j;i}) := \frac{\exp(g_i)}{\sum_{1 \leq i \leq m_j-1} \exp(g_i) + 1}$$

and

$$P(x_{j;0}) := \frac{1}{\sum_{1 \leq i \leq m_j-1} \exp(g_i) + 1}.$$

Now we have to show that one can find $N + 1$ vectors that can be classified in all 2^{N+1} possible ways such that the vectors correspond to n -tuples. Consider the vectors e_j having 1 at the position j and 0 elsewhere. Obviously, each vector e_j corresponds to a possible n -tuple. As $N + 1$ -th vector we choose the origin $(0, 0, \dots, 0)$. It corresponds to the n -tuple $(x_{1;0}, x_{2;0}, \dots, x_{n;0})$. These vectors can be classified in all 2^{N+1} possibilities as follows. Choose an arbitrary indicator function χ on these $N + 1$ points and set $f := (\pm 1, \pm 1, \dots, \pm 1)$ with positive sign at position j if and only if $\chi(e_j) = 1$. Define d as above in order to ensure that

$$\mathbf{x} \mapsto \langle c^{\mathbf{x}}, f \rangle + d$$

defines the logarithm of a probability function. Set $a := d \pm 1/2$ and choose the sign positive if and only if $\chi((0, 0, \dots, 0)) = 1$. Then the indicator function $\chi_a \circ w$ with $w(\mathbf{x}) := \langle c^{\mathbf{x}}, f \rangle + d$ coincides with the

desired classifier χ . Hence the VC dimension of product distributions is $N + 1$. \square

Now we give a bound for the VC dimension of the model classes introduced in Section 2.3.

Lemma 2 *The VC dimension of \mathcal{P}^k is at most*

$$h_k := \sum_{\mathbf{j}} m_{j_1} m_{j_2} \dots m_{j_k},$$

where m_i is the size of the set Ω_i .

Remark: h_s is at most $O(n^k)$ if m_i is uniformly bounded.

Proof: Let $P \in \mathcal{P}^k$. Since P factorizes we can characterize the expression $\ln P$ uniquely by a vector $f \in \mathbb{R}^h$ as follows: the vector f consists of $\binom{n}{k}$ blocks where the blocks are indexed by \mathbf{j} given in the Lemma. The block \mathbf{j} has size $\prod_i m_{j_i}$ and the entries within the block have the values

$$\ln q_{\mathbf{j}}(x_{j_1}, x_{j_2}, \dots, x_{j_k}),$$

where $(x_{j_1}, x_{j_2}, \dots, x_{j_k})$ runs over all elements of $\Omega_{j_1} \times \Omega_{j_2} \times \dots \times \Omega_{j_k}$. Each n -tuple $\mathbf{x} = (x_1, x_2, \dots, x_n)$ can be characterized by a vector $c^{\mathbf{x}} \in \mathbb{R}^h$ as follows: In each block \mathbf{j} there is exact one coordinate that corresponds to the s -tuple $(x_{j_1}, x_{j_2}, \dots, x_{j_k})$. Set this entry to 1 and the other entries of the block to 0. Then $\ln P(\mathbf{x})$ is given by the inner product of the vectors f and $c^{\mathbf{x}}$. We obtain the lemma since the VC dimension of the set of linear functionals on \mathbb{R}^h is h , see (Vapnik 1998). \square

Obviously, the model class \mathcal{H}_k^λ is a subset of \mathcal{P}^k and so we can give an upper bound the VC dimension of \mathcal{H}_k^λ with h_k which is independent of λ . Therefore Lemma 2 and Theorem 4 yields to the following theorem.

Theorem 5 *Let \mathcal{H}_k^λ be the model class of degree k and h_k be the upper bound on the VC dimension of \mathcal{H}_k^λ given in Lemma 2. Then for a training data $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_l$ we have for $P_\alpha \in \mathcal{H}_k^\lambda$ with probability at least $1 - \eta$*

$$R(P_\alpha) \leq R_{emp}(P_\alpha) + \phi(k, \lambda, \eta), \quad (6)$$

with $\phi(k, \lambda, \eta) := -\ln \lambda \sqrt{\frac{h_k - \ln h_k + \ln 16 + \ln l - \ln \eta}{l}}$.

The task to find the minimizer of R_{emp} in \mathcal{H}_k^λ even for fixed λ and k is computationally expensive. We will not treat this question here.

Theorem 5 tells us how to estimate the risk $R(P_\alpha)$ for a a priori chosen model class \mathcal{H}_k^λ . However we do not know which class to select. This question is addressed in the next section.

3.2 Estimating Log-linear Models from empirical data

We do not want to exclude k -factor models with large k a priori if the data strongly indicates that such a complex model is appropriate. Similarly, we do not want to exclude distributions with small λ . We rather want to avoid such models if the data does not give us strong enough reason for choosing them. To solve this problem we use a generalized version of the structural risk minimization principle (Vapnik 1998, Vapnik 1995). In this way the result of Theorem 5 can be extended such that k and λ need not to be chosen in advanced. For doing so we have to define a prior probability measure ν on the set of possible pairs (k, λ) in advance in order to get a bound on the true risk.

We obtain from Theorem 5 with a standard union bound argument, see e.g. Lemma 4.1 in (Herbrich 2002), following corollary.

Corollary 1 *Let $(\lambda_n)_{n \in \mathbb{N}}$ be a strict monotone decreasing sequence converging to zero. Let ν be a strict positive probability measure on \mathbb{N}^2 and $A_{kn} = \{1, \dots, k\} \times \{1, \dots, n\}$. Then for training data $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_l$ we have with probability at least $1 - \eta$*

$$R(P) \leq R_{emp}(P) + \phi(k, \lambda_n, \eta \nu(A_{kn})),$$

for $P \in \mathcal{H}_k^{\lambda_n}$ and ϕ defined above.

The probability measure ν in Corollary 1 has to be chosen before seeing the training data. Usually one chooses ν to be inversely proportional to ϕ . However, it should be emphasized that the statement in Corollary 1 does not assume that the measure ν is chosen according to any prior probabilities for the possible probability measures as used in Bayesian learning.

Now we can formulate structural risk minimization as follows:

$$\begin{aligned} & \underset{(k,n) \in \mathbb{N}^2}{\text{Minimize}} && t + \phi(k, \lambda, \eta \nu(A_{kn})) \\ & \text{subject to} && t = \min_{P \in \mathcal{H}_k^{\lambda_n}} R_{emp}(P) \end{aligned}$$

3.3 Model Computation by Convex Optimization

In this section we show that the minimization of the empirical risk R_{emp} is a convex optimization problem in the class of all k -factor models for a fixed k and λ . We formulate the optimization problem for $\ln P$ rather than for P . Let $\mathbf{x}^1, \dots, \mathbf{x}^l$ be the training data. For each set $\mathbf{j} = \{j_1, \dots, j_k\} \subset \{1, \dots, n\}$ let $V_{\mathbf{j}}$ be the vector space of real-valued functions on the set $\Omega_{\mathbf{j}} := \Omega_{j_1} \times \dots \times \Omega_{j_k}$. Define

$$V^k := \oplus_{\mathbf{j}} V_{\mathbf{j}}$$

where \mathbf{j} runs over all subsets of $\{1, \dots, n\}$ of size k . Now fix k and \mathbf{j} of size k . The set $\Omega_{\mathbf{j}}$ is discrete and so the functions

$$e_{w_1, w_2, \dots, w_k}(v_1, v_2, \dots, v_k) = \begin{cases} 1 & \text{if } w_1 = v_1, \dots, w_k = v_k \\ 0 & \text{else} \end{cases}$$

form a canonical basis of $V_{\mathbf{j}}$. Further let $\mathbf{x}_{\mathbf{j}}$ be the k -tuple given by the restriction of $\mathbf{x} \in \Omega$ to the variables x_{j_1}, \dots, x_{j_k} and $\mathbf{x}_{\mathbf{j}}^i$ the restriction of the training data \mathbf{x}^i to the variables of $\mathbf{x}_{\mathbf{j}}$.

Now, let us formulate the optimization problem of the empirical risk. Find a vector $f = \oplus_{\mathbf{j}} f_{\mathbf{j}} \in V^k$ with $f_{\mathbf{j}} \in V_{\mathbf{j}}$ such that

$$\tilde{R}_{emp}(f) := \frac{1}{l} \sum_{i=1}^l \sum_{\mathbf{j}} f_{\mathbf{j}}(\mathbf{x}_{\mathbf{j}}^i) \quad (7)$$

is minimal subject to the constraints

$$Z(f) := \sum_{\mathbf{x} \in \Omega} \exp\left(\sum_{\mathbf{j}} f_{\mathbf{j}}(\mathbf{x}_{\mathbf{j}})\right) = 1$$

and

$$G_{\mathbf{x}}(f) := \sum_{\mathbf{j}} f_{\mathbf{j}}(\mathbf{x}_{\mathbf{j}}) \geq \log \lambda \quad \text{for all } \mathbf{x} \in \Omega.$$

Then the probability measure $P \in \mathcal{H}_k^\lambda$ is given by $\ln f$. The functions \tilde{R}_{emp} and $G_{\mathbf{x}}$ are obviously linear in f . The function Z is convex in f since the exponent is linear in f and the exponential function is convex. Clearly, the sum of the convex functions

$$f \mapsto \exp\left(\sum_{\mathbf{j}} f_{\mathbf{j}}(\mathbf{x}_{\mathbf{j}})\right)$$

over all \mathbf{x} is convex. Therefore the problem can be solved with standard convex optimization methods (Pallaschke and Rolewicz 1997). However, for a large number of variables the solution becomes computationally expensive. Notice that the number of constraints in (7) increases exponentially with n . Furthermore the calculation of $\tilde{R}_{emp}(f)$ involves the summation over exponentially many possible n -tuples \mathbf{x} . This problem is similar to the calculation of the partition function in statistical mechanics which requires a summation over exponentially many states of a many-particle system. Those computational problems are not different from those appearing in conventional approaches to log-linear model selection.

4 CONCLUSION

We used structural risk minimization of statistical learning theory to obtain a new selection criterion for log-linear models. In this way we have not only a criterion to choose the suitable model complexity, but we found also a bound on the actual risk, i.e. the key feature of our approach is that it provides statements of the form “if a log-linear model of certain simplicity fits well to the observed data then we are guaranteed with high probability that it will fit future observations as well”. This kind of nonasymptotic statistics become more and more important for in research areas like cognitive and social science, where the number of possibly relevant features for the system under investigation is large. It offers a new way of developing models, namely, inferring statistical dependencies among a large number of features from data.

We structured the class of log-linear models according to the degree of the interaction terms. Using other measures of complexity than VC dimension like Gaussian complexity (Bartlett and Mendelson 2001) we believe that the obtained bounds can be improved. The structure of the model classes can be refined taking for example prior knowledge into account. In future this model selection criterion need to be tested on real-world data and the results compared to existing ones.

References

- Cover, T. and Thomas, J. (1991), “Elements of Information Theory,” Wileys Series in Telecommunications, New York.

- Vapnik, V. (1998), “Statistical learning theory.” John Wileys & Sons, New York.
- Vapnik, V. (1995), “The Nature of Statistical learning theory,” Springer-Verlag, New York.
- Christensen, R. (1997), “Log-linear Models and Logistic Regression,” Springer Verlag.
- Goodman, L. (1978), “Analyzing Qualitative/Categorical Data,” Addison Wesley, Cambridge MA, 1978.
- Herbrich, R. (2002) “Learning kernel classifiers,” MIT Press, Cambridge, MA, 2002.
- Vapnik, V. N. and Chervonenkis, A. J. (1971), “On the uniform convergence of relative frequencies of events to their probabilities,” *Theory Probab. Apl.*, **16** pp. 264–280.
- Pearl, J. (1985) “Bayesian networks: A model of self-activated memory for evidential reasoning,” In *Proceedings, Cognitive Science Society*, pp. 329–334, Greenwich, Albex.
- Pearl, J. “Causality. Models, Reasoning, and Inference”. Cambridge University Press, Cambridge 2000.
- Pearl, J. (1988) *Probabilistic Reasoning in Intelligent Systems*. San Mateo, CA: Morgan Kaufmann.
- Schwarz, G. (1978), “Estimating the Dimension of a Model,” *Annals of Statistics*, **6**, pp. 461-464.
- Kass, R. E. and Raftery, A. E. (1995) “Bayes factors,” *JASA*, **90**, pp. 773-795.
- Akaike, H. (1973) “Extension of the maximum likelihood principle,” In B. Petrov and F. Csáki, ed., *2nd International Symposium on Information theory*, pages 267–281. Akadémiai Kiado.
- Barron, A. and Cover, T. (1991) “Minimum Complexity Density Estimation,” *IEEE Transaction on Information Theory*, **37**, 4, pp. 1034–1054.

- Hagenaars, J. A. (1994), "Loglinear models with latent variables," *Sage University Paper series on Quantitative Applications in Social Sciences* **07-094**. Newbury, CA:Sage.
- Ishii-Kuntz, M. (1997). "Ordinal loglinear models," *Sage University Paper series on Quantitative Applications in Social Sciences* **07-097**. Newbury, CA:Sage.
- Pitt, M. and Myung, J. (2002), "When a good fit can be bad," *Trends Cognitive Sciences*, **6**, 10, pp. 421-425.
- Rissanen, J. (1996) "Fisher Information and Stochastic Complexity," *IEEE Transaction on Information Theory*, **42**.
- Hansen, M. H. and Yu, B. (2001), "Model Selection and the Principle of Minimum Description Length," *JASA*, **96**, 454, pp. 746-774.
- Hansen, M. and Yu, B. (2002) "Minimum description length model selection criteria for generalized models," <http://cm-labs.com/who/cocteau/papers/index.html>, 2002.
- Wocjan, P. and Janzing, D. and Beth, T. (2002), "Required sample size for learning sparse Bayesian networks with many variables," *LANL-preprint* <http://xxx.lanl.gov/abs/cs.LG/0204052>.
- Pallaschke, D. and Rolewicz, S. (1997), "Foundations of Mathematical Optimization," Vol. 388, Kluwer Academic Publishers, Dordrecht.
- Bartlett, P.L. and Mendelson, S. Rademacher and gaussian complexities: risk bounds and structural results, Proceedings of the 14th annual conference on Computational Learning Theory, 224-240, 2001.