LIFTS OF C_{∞} AND L_{∞} -MORPHISMS TO G_{∞} -MORPHISMS

Grégory Ginot^(a), Gilles Halbout^(b)

^(a) Laboratoire Analyse Géométrie et Applications, Université

Paris 13 et Ecole Normale Supèrieure de Cachan, France

^(b) Institut de Recherche Mathématique Avancée, Université

Louis Pasteur et CNRS, Strasbourg, France

ABSTRACT. Let \mathfrak{g}_2 be the Hochschild complex of cochains on $C^{\infty}(\mathbb{R}^n)$ and \mathfrak{g}_1 be the space of multivector fields on \mathbb{R}^n . In this paper we prove that given any G_{∞} -structure (i.e. Gerstenhaber algebra up to homotopy structure) on \mathfrak{g}_2 , and any C_{∞} -morphism φ (i.e. morphism of commutative, associative algebra up to homotopy) between \mathfrak{g}_1 and \mathfrak{g}_2 , there exists a G_{∞} morphism Φ between \mathfrak{g}_1 and \mathfrak{g}_2 that restricts to φ . We also show that any L_{∞} -morphism (i.e. morphism of Lie algebra up to homotopy), in particular the one constructed by Kontsevich, can be deformed into a G_{∞} -morphism, using Tamarkin's method for any G_{∞} -structure on \mathfrak{g}_2 . We also show that any two of such G_{∞} -morphisms are homotopic.

0-Introduction

Let M be a differential manifold and $\mathfrak{g}_2 = (C^{\bullet}(A, A), b)$ be the Hochschild cochain complex on $A = C^{\infty}(M)$. The classical Hochschild-Kostant-Rosenberg theorem states that the cohomology of \mathfrak{g}_2 is the graded Lie algebra $\mathfrak{g}_1 = \Gamma(M, \wedge^{\bullet}TM)$ of multivector fields on M. There is also a graded Lie algebra structure on \mathfrak{g}_2 given by the Gerstenhaber bracket. In particular \mathfrak{g}_1 and \mathfrak{g}_2 are also Lie algebras up to homotopy $(L_{\infty}$ -algebra for short). In the case $M = \mathbb{R}^n$, using different methods, Kontsevich ([Ko1] and [Ko2]) and Tamarkin ([Ta]) have proved the existence of Lie homomorphisms "up to homotopy" $(L_{\infty}$ -morphisms) from \mathfrak{g}_1 to \mathfrak{g}_2 . Kontsevich's proof uses graph complex and is related to multizeta functions whereas Tamarkin's construction uses the existence of Drinfeld's associators. In fact Tamarkin's L_{∞} -morphism comes from the restriction of a Gerstenhaber algebra up to homotopy homomorphism $(G_{\infty}$ -morphism) from \mathfrak{g}_1 to \mathfrak{g}_2 . The G_{∞} -algebra structure on \mathfrak{g}_1 is induced by its classical Gerstenhaber algebra structure and a far less trivial G_{∞} -structure on \mathfrak{g}_2 was proved to exist by Tamarkin [Ta] and relies on a Drinfeld's associator. Tamarkin's G_{∞} -morphism also restricts into a commutative, associative up to homotopy morphism (C_{∞} -morphism for short). The C_{∞} -structure on \mathfrak{g}_2 (given by

Keywords : Deformation quantization, star-product, homotopy formulas, homological methods. **AMS Classification :** Primary 16E40, 53D55, Secondary 18D50, 16S80.

restriction of the G_{∞} -one) highly depends on Drinfeld's associator, and any two choices of a Drinfeld associator yields a priori different C_{∞} -structures. When M is a Poisson manifold, Kontsevich and Tamarkin homomorphisms imply the existence of a star-product (see [BFFLS1] and [BFFLS2] for a definition). A connection between the two approaches has been given in [KS] but the morphisms given by Kontsevich and Tamarkin are not the same. The aim of this paper is to show that, given any G_{∞} -structure on \mathfrak{g}_2 and any C_{∞} -morphism φ between \mathfrak{g}_1 and \mathfrak{g}_2 , there exists a G_{∞} -morphism Φ between \mathfrak{g}_1 and \mathfrak{g}_2 that restricts to φ . We also show that any L_{∞} -morphism can be deformed into a G_{∞} -one. In the first section, we fix notation and recall the definitions of L_{∞} and G_{∞} -structures. In the second section we state and prove the main theorem. In the last section we show that any two G_{∞} -morphisms given by Tamarkin's method are homotopic.

Remark : In the sequel, unless otherwise is stated, the manifold M is supposed to be \mathbb{R}^n for some $n \ge 1$. Most results could be generalized to other manifolds using techniques of Kontsevich [Ko1] (also see [TS], [CFT]).

1- C_{∞} , L_{∞} and G_{∞} -structures

For any graded vector space \mathfrak{g} , we choose the following degree on $\wedge^{\bullet}\mathfrak{g}$: if X_1, \ldots, X_k are homogeneous elements of respective degree $|X_1|, \ldots, |X_k|$, then

$$|X_1 \wedge \dots \wedge X_k| = |X_1| + \dots + |X_k| - k$$

In particular the component $\mathfrak{g} = \wedge^1 \mathfrak{g} \subset \wedge^{\bullet} \mathfrak{g}$ is the same as the space \mathfrak{g} with degree shifted by one. The space $\wedge^{\bullet} \mathfrak{g}$ with the deconcatenation cobracket is the cofree cocommutative coalgebra on \mathfrak{g} with degree shifted by one (see [LS], Section 2). Any degree one map $d^k : \wedge^k \mathfrak{g} \to \mathfrak{g}$ $(k \ge 1)$ extends into a derivation $d^k : \wedge^{\bullet} \mathfrak{g} \to \wedge^{\bullet} \mathfrak{g}$ of the coalgebra $\wedge^{\bullet} \mathfrak{g}$ by cofreeness property.

Definition 1.1. A vector space \mathfrak{g} is endowed with a L_{∞} -algebra (Lie algebras "up to homotopy") structure if there are degree one linear maps $m^{1,\dots,1}$, with k ones : $\wedge^k \mathfrak{g} \to \mathfrak{g}$ such that if we extend them to maps $\wedge^{\bullet} \mathfrak{g} \to \wedge^{\bullet} \mathfrak{g}$, then $d \circ d = 0$ where d is the derivation

$$d = m^1 + m^{1,1} + \dots + m^{1,\dots,1} + \dots$$

For more details on L_{∞} -structures, see [LS]. It follows from the definition that a L_{∞} -algebra structure induces a differential coalgebra structure on $\wedge^{\bullet}\mathfrak{g}$ and that the map $m^{1}:\mathfrak{g} \to \mathfrak{g}$ is a differential. If $m^{1,\dots,1}: \wedge^{k}\mathfrak{g} \to \mathfrak{g}$ are 0 for $k \geq 3$, we get the usual definition of (differential if $m^{1} \neq 0$) graded Lie algebras.

For any graded vector space \mathfrak{g} , we denote $\underline{\mathfrak{g}}^{\otimes n}$ the quotient of $\mathfrak{g}^{\otimes n}$ by the image of all shuffles of length n (see [GK] or [GH] for details). The graded vector space $\bigoplus_{n\geq 0} \underline{\mathfrak{g}}^{\otimes n}$ is a quotient coalgebra of the tensor coalgebra $\bigoplus_{n\geq 0} \mathfrak{g}^{\otimes n}$. It is well known that this coalgebra $\bigoplus_{n\geq 0} \underline{\mathfrak{g}}^{\otimes n}$ is the cofree Lie coalgebra on the vector space \mathfrak{g} (with degree shifted by minus one).

Definition 1.2. A C_{∞} -algebra (commutative and associative "up to homotopy" algebra) structure on a vector space \mathfrak{g} is given by a collection of degree one linear maps m^k : $\underline{\mathfrak{g}}^{\otimes k} \to \mathfrak{g}$ such that if we extend them to maps $\oplus \underline{\mathfrak{g}}^{\otimes \bullet} \to \oplus \underline{\mathfrak{g}}^{\otimes \bullet}$, then $d \circ d = 0$ where d is the derivation

$$d = m^1 + m^2 + m^3 + \cdots$$

In particular a C_{∞} -algebra is an A_{∞} -algebra.

For any space \mathfrak{g} , we denote $\wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$ the graded space

$$\wedge^{\bullet}\underline{\mathfrak{g}}^{\otimes \bullet} = \bigoplus_{m \ge 1, \ p_1 + \dots + p_n = m} \underline{\mathfrak{g}}^{\otimes p_1} \wedge \dots \wedge \underline{\mathfrak{g}}^{\otimes p_n}.$$

We use the following grading on $\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}$: for $x_1^1, \cdots, x_n^{p_n} \in \mathfrak{g}$, we define

$$|\underline{x_1^1 \otimes \cdots \otimes x_1^{p_1}} \wedge \cdots \wedge \underline{x_n^1 \otimes \cdots \otimes x_n^{p_n}}| = \sum_{i_1}^{p_1} |x_1^{i_1}| + \cdots + \sum_{i_n}^{p_1} |x_n^{i_n}| - n.$$

Notice that the induced grading on $\wedge^{\bullet}\mathfrak{g} \subset \wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$ is the same than the one introduced above. The cobracket on $\oplus \mathfrak{g}^{\otimes \bullet}$ and the coproduct on $\wedge^{\bullet}\mathfrak{g}$ extend to a cobracket and a coproduct on $\wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$ which yield a Gerstenhaber coalgebra structure on $\wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$. It is well known that this coalgebra structure is cofree (see [Gi], Section 3 for example).

Definition 1.3. A G_{∞} -algebra (Gerstenhaber algebra "up to homotopy") structure on a graded vector space \mathfrak{g} is given by a collection of degree one maps

$$m^{p_1,\ldots,p_n}$$
 : $\underline{\mathfrak{g}^{\otimes p_1}}\wedge\cdots\wedge\underline{\mathfrak{g}^{\otimes p_n}}\to\mathfrak{g}$

indexed by $p_1, \ldots p_n \ge 1$ such that their canonical extension: $\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet} \to \wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}$ satisfies $d \circ d = 0$ where

$$d = \sum_{m \ge 1, p_1 + \dots + p_n = m} m^{p_1, \dots, p_n}.$$

Again, as the coalgebra structure of $\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}$ is cofree, the map d makes $\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}$ into a differential coalgebra. If the maps m^{p_1,\ldots,p_n} are 0 for $(p_1,p_2,\ldots) \neq (1,0,\ldots), (1,1,0,\ldots)$ or $(2,0,\ldots)$, we get the usual definition of (differential if $m^1 \neq 0$) Gerstenhaber algebra.

The space of multivector fields \mathfrak{g}_1 is endowed with a graded Lie bracket $[-, -]_S$ called the Schouten bracket (see [Kos]). This Lie algebra can be extended into a Gerstenhaber algebra, with commutative structure given by the exterior product: $(\alpha, \beta) \mapsto \alpha \wedge \beta$

Setting $d_1 = m_1^{1,1} + m_1^2$, where $m_1^{1,1} : \wedge^2 \mathfrak{g}_1 \to \mathfrak{g}_1$, and $m_1^2 : \underline{\mathfrak{g}_1}^{\otimes 2} \to \mathfrak{g}_1$ are the extension of the Schouten bracket and the exterior product, we find that (\mathfrak{g}_1, d_1) is a G_{∞} -algebra.

In the same way, one can define a differential Lie algebra structure on the vector space $\mathfrak{g}_2 = C(A, A) = \bigoplus_{k\geq 0} C^k(A, A)$, the space of Hochschild cochains (generated by differential klinear maps from A^k to A), where $A = C^{\infty}(M)$ is the algebra of smooth differential functions over M. Its bracket $[-, -]_G$, called the Gerstenhaber bracket, is defined, for $D, E \in \mathfrak{g}_2$, by

$$[D, E]_G = \{D|E\} - (-1)^{|E||D|} \{E|D\},\$$

where

$$\{D|E\}(x_1,...,x_{d+e-1}) = \sum_{i\geq 0} (-1)^{|E|\cdot i} D(x_1,...,x_i,E(x_{i+1},...,x_{i+e}),...).$$

The space \mathfrak{g}_2 has a grading defined by $|D| = k \Leftrightarrow D \in C^{k+1}(A, A)$ and its differential is $b = [m, -]_G$, where $m \in C^2(A, A)$ is the commutative multiplication on A.

Tamarkin (see [Ta] or also [GH]) stated the existence of a G_{∞} -structure on \mathfrak{g}_2 (depending on a choice of a Drinfeld associator) given by a differential $d_2 = m_2^1 + m_2^{1,1} + m_2^2 + \cdots + m_2^{p_1,\dots,p_n} + \cdots$, on $\wedge \mathfrak{g}_2^{\otimes \mathfrak{o}}$ satisfying $d_2 \circ d_2 = 0$. Although this structure is non-explicit, it satisfies the following three properties :

- (a) m_2^1 is the extension of the differential b
- (b) $m_2^{1,1}$ is the extension of the Gerstenhaber bracket $[-,-]_G$ and $m_2^{1,1,\dots,1} = 0$
- (c) m_2^2 induces the exterior product in cohomology and the collection of the $(m^k)_{k>1}$ defines a C_{∞} -structure on $\mathfrak{g}_2.(1.1)$

Definition 1.4. A L_{∞} -morphism between two L_{∞} -algebras $(\mathfrak{g}_1, d_1 = m_1^1 + ...)$ and $(\mathfrak{g}_2, d_2 = m_2^1 + ...)$ is a morphism of differential coalgebras

$$\varphi : (\wedge^{\bullet} \mathfrak{g}_1, d_1) \to (\wedge^{\bullet} \mathfrak{g}_2, d_2).$$
(1.2)

Such a map φ is uniquely determined by a collection of maps $\varphi^n : \wedge^n \mathfrak{g}_1 \to \mathfrak{g}_2$ (again by cofreeness properties). In the case \mathfrak{g}_1 and \mathfrak{g}_2 are respectively the graded Lie algebra $(\Gamma(M, \wedge TM), [-, -]_S)$ and the differential graded Lie algebra $(C(A, A), [-, -]_G)$, the formality theorems of Kontsevich and Tamarkin state the existence of a L_{∞} -morphism between \mathfrak{g}_1 and \mathfrak{g}_2 such that φ^1 is the Hochschild-Kostant-Rosenberg quasi-isomorphism.

Definition 1.5. A morphism of C_{∞} -algebras between two C_{∞} -algebras (\mathfrak{g}_1, d_1) and (\mathfrak{g}_2, d_2) is a map $\phi : (\oplus \mathfrak{g}_1^{\otimes \bullet}, d_1) \to (\oplus \mathfrak{g}_2^{\otimes \bullet}, d_2)$ of codifferential coalgebras.

A C_{∞} -morphism is in particular a morphism of A_{∞} -algebras and is uniquely determined by maps $\partial^k : \mathfrak{g}^{\otimes k} \to \mathfrak{g}$.

Definition 1.6. A morphism of G_{∞} -algebras between two G_{∞} -algebras (\mathfrak{g}_1, d_1) and (\mathfrak{g}_2, d_2) is a map $\phi : (\wedge^{\bullet}\mathfrak{g}_1^{\otimes \bullet}, d_1) \to (\wedge^{\bullet}\mathfrak{g}_2^{\otimes \bullet}, d_2)$ of codifferential coalgebras.

There are coalgebras inclusions $\wedge^{\bullet}\mathfrak{g} \to \wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$, $\oplus \mathfrak{g}^{\otimes \bullet} \to \wedge^{\bullet}\mathfrak{g}^{\otimes \bullet}$ and it is easy to check that any G_{∞} -morphism between two G_{∞} -algebras $(\mathfrak{g}, \sum m^{p_1,\dots,p_n})$, $(\mathfrak{g}', \sum m'^{p_1,\dots,p_n})$ restricts to a L_{∞} -morphism $(\wedge^{\bullet}\mathfrak{g}, \sum m^{1,\dots,1}) \to (\wedge^{\bullet}\mathfrak{g}', \sum m'^{1,\dots,1})$ and a C_{∞} -morphism $(\oplus \mathfrak{g}^{\otimes \bullet}, \sum m^k) \to (\oplus \mathfrak{g}'^{\otimes \bullet}, \sum m'^k)$. In the case \mathfrak{g}_1 and \mathfrak{g}_2 are as above, Tamarkin's theorem states that there exists a G_{∞} -morphism between the two G_{∞} algebras \mathfrak{g}_1 and \mathfrak{g}_2 (with the G_{∞} structure he built) that restricts to a C_{∞} and a L_{∞} -morphism.

2-Main theorem

We keep the notations of the previous section, in particular \mathfrak{g}_2 is the Hochschild complex of cochains on $C^{\infty}(M)$ and \mathfrak{g}_1 its cohomology. Here is our main theorem.

Theorem 2.1. Given any G_{∞} -structure d_2 on \mathfrak{g}_2 satisfying the three properties of (1.1), and any C_{∞} -morphism φ between \mathfrak{g}_1 and \mathfrak{g}_2 such that φ^1 is the Hochschild-Kostant-Rosenberg map, there exists a G_{∞} -morphism $\Phi : (\mathfrak{g}_1, d_1) \to (\mathfrak{g}_2, d_2)$ that restricts to φ .

Also, given any L_{∞} -morphism γ between \mathfrak{g}_1 and \mathfrak{g}_2 such that γ^1 is the Hochschild-Kostant-Rosenberg map, there exists a G_{∞} -structure (\mathfrak{g}_1, d'_1) on \mathfrak{g}_1 and G_{∞} -morphism Γ : $(\mathfrak{g}_1, d'_1) \rightarrow (\mathfrak{g}_2, d_2)$ that restricts to γ . Moreover there exists a G_{∞} -morphism Γ' : $(\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_1, d_1)$.

In particular, Theorem 2.1 applies to the formality map of Kontsevich and also to any C_{∞} -map derived (see [Ta], [GH]) from any B_{∞} -structure on \mathfrak{g}_2 lifting the Gerstenhaber structure of \mathfrak{g}_1 .

Let us first recall the proof of Tamarkin's formality theorem (see [GH] for more details):

- 1. First one proves there exists a G_{∞} -structure on \mathfrak{g}_2 , with differential d_2 , as in (1.1).
- 2. Then, one constructs a G_{∞} -structure on \mathfrak{g}_1 given by a differential d'_1 together with a G_{∞} -morphism Φ between (\mathfrak{g}_1, d'_1) and (\mathfrak{g}_2, d_2) .
- 3. Finally, one constructs a G_{∞} -morphism Φ' between (\mathfrak{g}_1, d_1) and (\mathfrak{g}_1, d'_1) .

The composition $\Phi \circ \Phi'$ is then a G_{∞} -morphism between (\mathfrak{g}_1, d_1) and (\mathfrak{g}_2, d_2) , thus restricts to a L_{∞} -morphism between the differential graded Lie algebras \mathfrak{g}_1 and \mathfrak{g}_2 .

We suppose now that, in the first step, we take any G_{∞} -structure on \mathfrak{g}_2 given by a differential d_2 and we suppose we are given a C_{∞} -morphism φ and a L_{∞} -morphism γ between \mathfrak{g}_1 and \mathfrak{g}_2 satisfying $\gamma^1 = \varphi^1 = \varphi_{\text{HKR}}$ the Hochschild-Kostant-Rosenberg quasi-isomorphism. *Proof of Theorem 2.1:*

The Theorem will follow if we prove that steps 2 and 3 of Tamarkin's construction are still true with the extra conditions that the restriction of the G_{∞} -morphism Φ (resp. Φ') on the C_{∞} -structures is the C_{∞} -morphism $\varphi : \mathfrak{g}_1 \to \mathfrak{g}_2$ (resp. id).

Let us recall (see [GH]) that the constructions of Φ and d'_1 can be made by induction. For i = 1, 2 and $n \ge 0$, let us set

$$V_i^{[n]} = \bigoplus_{p_1 + \dots + p_k = n} \underline{\mathfrak{g}_i^{\otimes p_1}} \wedge \dots \wedge \underline{\mathfrak{g}_i^{\otimes p_k}}$$

and $V_i^{[\leq n]} = \sum_{k \leq n} V_i^{[k]}$. Let $d_2^{[n]}$ and $d_2^{[\leq n]}$ be the sums

$$d_2^{[n]} = \sum_{p_1 + \dots + p_k = n} d_2^{p_1, \dots, p_k}$$
 and $d_2^{[\leq n]} = \sum_{p \leq n} d_2^{[p]}$

Clearly, $d_2 = \sum_{n \ge 1} d_2^{[n]}$. In the same way, we denote $d'_1 = \sum_{n \ge 1} d'_1^{[n]}$ with

$$d'_{1}^{[n]} = \sum_{p_{1}+\dots+p_{k}=n} d'_{1}^{p_{1},\dots,p_{k}}$$
 and $d'_{1}^{[\leq n]} = \sum_{1 \leq k \leq n} d'_{1}^{[k]}$

We know from Section 1 that a morphism $\Phi : (\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}, d'_1) \to (\wedge \overset{\bullet}{\mathfrak{g}_2^{\otimes \bullet}}, d_2)$ is uniquely determined by its components $\Phi^{p_1, \dots, p_k} : \underline{\mathfrak{g}_1^{\otimes p_1}} \wedge \dots \wedge \underline{\mathfrak{g}_1^{\otimes p_k}} \to \mathfrak{g}_2$. Again, we have $\Phi = \sum_{n \geq 1} \Phi^{[n]}$ with

$$\Phi^{[n]} = \sum_{p_1 + \dots + p_k = n} \Phi^{p_1, \dots, p_k}$$
 and $\Phi^{[\leq n]} = \sum_{1 \leq k \leq n} \Phi^{[k]}$.

We want to construct the maps $d'_{1}^{[n]}$ and $\Phi^{[n]}$ by induction with the initial condition

$$d'_{1}^{[1]} = 0$$
 and $\Phi^{[1]} = \varphi_{\text{HKR}},$

where φ_{HKR} : $(\mathfrak{g}_1, 0) \to (\mathfrak{g}_2, b)$ is the Hochschild-Kostant-Rosenberg quasi-isomorphism (see [HKR]) defined, for $\alpha \in \mathfrak{g}_1, f_1, \cdots, f_n \in A$, by

$$\varphi_{\text{HKR}} : \alpha \mapsto ((f_1, \dots, f_n) \mapsto \langle \alpha, df_1 \wedge \dots \wedge df_n \rangle).$$

Moreover, we want the following extra conditions to be true:

$$\Phi^{k \ge 2} = \varphi^k, \qquad {d'}_1^2 = d_1^2, \qquad {d'}_1^{k \ge 3} = 0.$$
(2.3)

Now suppose the construction is done for n-1 $(n \ge 2)$, i.e., we have built maps $(d'_1^{[i]})_{i \le n-1}$ and $(\Phi^{[i]})_{i \le n-1}$ satisfying conditions (2.3) and

$$\Phi^{[\leq n-1]} \circ d'_{1}^{[\leq n-1]} = d_{2}^{[\leq n-1]} \circ \Phi^{[\leq n-1]} \text{ on } V_{1}^{[\leq n-1]} \text{ and } d'_{1}^{[\leq n-1]} \circ d'_{1}^{[\leq n-1]} = 0 \text{ on } V_{1}^{[\leq n]}.$$
(2.4)

In [GH], we prove that for any such $(d'_1^{[i]})_{i \leq n-1}$ and $(\Phi^{[i]})_{i \leq n-1}$, one can construct $d'_1^{[n]}$ and $\Phi^{[n]}$ such that condition (2.4) is true for n instead of n-1. To complete the proof of Theorem 2.1 (step 2), we have to show that $d'_1^{[n]}$ and $\Phi^{[n]}$ can be chosen to satisfy conditions (2.3). In the equation 2.4, the terms d'_1^k and Φ^k only act on V_1^k . So one can replace Φ^n with φ^n , d'_1^2 with d_1^2 (or $d'_1^i, i \geq 3$ with 0) provided conditions (2.4) are still satisfied on V_1^n . The other terms acting on V_1^n in the equation (2.4) only involve terms $\Phi^m = \varphi^m$ and d'_1^m . Then conditions (2.4) on $V_1^{1,\dots,1}$ are the equations that should be satisfied by a C_∞ -morphism between the C_∞ -algebras $(\mathfrak{g}_1, d'_1^{1,1} = d_1^{1,1})$ and $(\mathfrak{g}_2, \sum_{k\geq 1} d_2^k)$ restricted to V_1^n . Hence by hypothesis on φ the conditions hold.

Similarly the construction of Φ' can be made by induction. Let us recall the proof given in [GH]. Again a morphism $\Phi' : (\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}, d_1) \to (\wedge \overset{\bullet}{\mathfrak{g}_2^{\otimes \bullet}}, d'_1)$ is uniquely determined by its components $\Phi'^{p_1, \dots, p_k} : \underline{\mathfrak{g}_1^{\otimes p_1}} \wedge \dots \wedge \underline{\mathfrak{g}_1^{\otimes p_k}} \to \mathfrak{g}_1$. We write $\Phi' = \sum_{n \ge 1} \Phi'^{[n]}$ with

$$\Phi'^{[n]} = \sum_{p_1 + \dots + p_k = n} \Phi'^{p_1, \dots, p_k}$$
 and $\Phi'^{[\leq n]} = \sum_{1 \leq k \leq n} \Phi'^{[k]}$.

We construct the maps ${\Phi'}^{[n]}$ by induction with the initial condition ${\Phi'}^{[1]} = id$. Moreover, we want the following extra conditions to be true:

$$\Phi'^n = 0 \text{ for } n \ge 2. \tag{2.5}$$

Now suppose the construction is done for n-1 $(n \ge 2)$, i.e., we have built maps $(\Phi'^{[i]})_{i\le n-1}$ satisfying conditions (2.5) and

$$\Phi^{\prime [\leq n-1]} d_1^{[\leq n]} = d_1^{\prime [\leq n]} \Phi^{\prime [\leq n-1]} \text{ on } V_1^{[\leq n]}.$$
(2.6)

In [GH], we prove that for any such $(\Phi'^{[i]})_{i \leq n-1}$, one can construct $\Phi'^{[n]}$ such that condition (2.6) is true for *n* instead of n-1 in the following way : making the equation $\Phi' d_1 = d'_1 \Phi'$ on $V_1^{[n+1]}$ explicit, we get

$$\Phi^{\prime [\leq n]} d_1^{[\leq n+1]} = d_1^{\prime [\leq n+1]} \Phi^{\prime [\leq n]}.$$
(2.7)

If we now take into account that $d_1^{[i]} = 0$ for $i \neq 2$, $d_1'^{[1]} = 0$ and that on $V_1^{[n+1]}$ we have $\Phi'^{[k]}d_1^{[l]} = d_1'^{[\leq k]}\Phi'^{[l]} = 0$ for k+l > n+2, the identity (2.7) becomes

$$\Phi'^{[\leq n]} d_1^{[2]} = \sum_{k=2}^{n+1} d_1'^{[k]} \Phi'^{[\leq n-k+2]}.$$

As $d'_1{}^{[2]} = d_1{}^{[2]}$, (2.7) is equivalent to

$$d_1^{[2]}\Psi'^{[\leq n]} - \Phi'^{[\leq n]}d_1^{[2]} = \left[d_1^{[2]}, \Phi'^{[\leq n]}\right] = -\sum_{k=3}^{n+1} d_1'^{[k]}\Phi'^{[\leq n-k+2]}$$

Notice that $d_1^{[2]} = m_1^{1,1} + m_1^2$. Then the construction will be possible when the term $\sum_{k=3}^{n+1} d_1'^{[k]} \psi'^{[\leq n-k+2]}$ is a couboundary in the subcomplex of $(\operatorname{End}(\wedge^{\bullet}\mathfrak{g}_1^{\otimes \bullet}), [d_1^{[2]}, -])$ consisting of maps which restrict to zero on $\bigoplus_{n\geq 2}\mathfrak{g}_1^{\otimes n}$. It is always a cocycle by straightforward computation (see [GH]) and the subcomplex is acyclic because both $(\operatorname{End}(\wedge^{\bullet}\mathfrak{g}_1^{\otimes \bullet}), [d_1^{[2]}, -])$ and the Harrison cohomology of \mathfrak{g}_1 are trivial according to Tamarkin [Ta] (see also [GH] Proposition 5.1 and [Hi] 5.4).

In the case of the L_{∞} -morphism γ , the first step is similar: the fact that γ is a L_{∞} -map enables us to build a G_{∞} -structure (\mathfrak{g}_1, d'_1) on \mathfrak{g}_1 and a G_{∞} -morphism $\Gamma : (\mathfrak{g}_1, d'_1) \rightarrow (\mathfrak{g}_2, d_2)$ such that:

$$\Gamma^{1,\dots,1} = \gamma^{1,\dots,1}, \qquad d'_{1}^{1,1} = d_{1}^{1,1}, \qquad d'_{1}^{1,1,\dots,1} = 0.$$
 (2.8)

For the second step, we have to build a map Γ' satisfying the equation

$$d_1^{[2]} \Gamma'^{[\leq n]} - \Gamma'^{[\leq n]} d_1^{[2]} = \left[d_1^{[2]}, \Gamma'^{[\leq n]} \right] = -\sum_{k=3}^{n+1} d_1'^{[k]} \Gamma'^{[\leq n-k+2]}$$

on $V_1^{[n+1]}$ for any $n \geq 1$. Again, because Tamarkin has prooved that the complex $(\operatorname{End}(\wedge \mathfrak{g}_1^{\otimes \mathfrak{o}}), [d_1^{[2]}, -])$ is acyclic (we are in the case $M = \mathbb{R}^n$), the result follows from the fact that $\sum_{k=3}^{n+1} d_1'^{[k]} \Gamma'^{[\leq n-k+2]}$ is a cocycle. The difference with the C_{∞} -case is that the $\Gamma'^{1,\ldots,1}$ could be non zero.

3-The difference between two G_{∞} -maps

In this section we investigate the difference between two differents G_{∞} -formality maps. We fix once for all a G_{∞} -structure on \mathfrak{g}_2 (given by a differential d_2) satisfying the conditions (1.1) and a morphism of G_{∞} -algebras $T : (\mathfrak{g}_1, d_1) \to (\mathfrak{g}_2, d_2)$ such that $T^1 : \mathfrak{g}_1 \to \mathfrak{g}_2$ is φ_{HKR} . Let $K : (\mathfrak{g}_1, d_1) \to (\mathfrak{g}_2, d_2)$ be any other G_{∞} -morphism with $K^1 = \varphi_{\text{HKR}}$ (for example any lift of a Kontsevich formality map or any G_{∞} -maps lifting another C_{∞} morphism). **Theorem 3.1.** There exists a map $h : \wedge \overset{\bullet}{\underline{g}_1^{\otimes \bullet}} \to \wedge \overset{\bullet}{\underline{g}_2^{\otimes \bullet}}$ such that

$$T - K = h \circ d_1 + d_2 \circ h.$$

In other words the formality G_{∞} -morphisms K and T are homotopic.

The maps T and K are elements of the cochain complex $\left(\operatorname{Hom}(\wedge^{\bullet}\underline{\mathfrak{g}_{1}^{\otimes \bullet}}, \wedge^{\bullet}\underline{\mathfrak{g}_{2}^{\otimes \bullet}}), \delta\right)$ with differential given, for all $f \in \operatorname{Hom}(\wedge^{\bullet}\underline{\mathfrak{g}_{1}^{\otimes \bullet}}, \wedge^{\bullet}\underline{\mathfrak{g}_{2}^{\otimes \bullet}}), |f| = k$, by

$$\delta(f) = d_2 \circ f - (-1)^k f \circ d_1.$$

We first compare this cochain complex with the complexes $\left(\operatorname{End}(\wedge \underline{\mathfrak{g}_1^{\otimes \bullet}}), [d_1; -]\right)$ and $\left(\operatorname{End}(\wedge \underline{\mathfrak{g}_2^{\otimes \bullet}}), [d_2; -]\right)$ (where [-; -] is the graded commutator of morphisms). There are morphisms

$$T_*: \operatorname{End}(\wedge \underline{\bullet}\underline{\mathfrak{g}_1^{\otimes \bullet}}) \to \operatorname{Hom}(\wedge \underline{\bullet}\underline{\mathfrak{g}_1^{\otimes \bullet}}, \wedge \underline{\bullet}\underline{\mathfrak{g}_2^{\otimes \bullet}}), \ T^*: \operatorname{End}(\wedge \underline{\bullet}\underline{\mathfrak{g}_2^{\otimes \bullet}}) \to \operatorname{Hom}(\wedge \underline{\bullet}\underline{\mathfrak{g}_1^{\otimes \bullet}}, \wedge \underline{\bullet}\underline{\mathfrak{g}_2^{\otimes \bullet}})$$

defined, for $f \in \operatorname{End}(\wedge \underline{\mathfrak{g}_2^{\otimes \bullet}})$ and $g \in \operatorname{Hom}(\wedge \underline{\mathfrak{g}_1^{\otimes \bullet}}, \wedge \underline{\mathfrak{g}_2^{\otimes \bullet}})$, by

$$T_*(f) = T \circ f, \ T^*(g) = g \circ T.$$

Lemma 3.2. The morphisms

$$T_*: \left(\operatorname{End}(\wedge^{\bullet} \underline{\mathfrak{g}_1^{\otimes \bullet}}), [d_1; -] \right) \to \left(\operatorname{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}_2^{\otimes \bullet}}, \wedge^{\bullet} \underline{\mathfrak{g}_2^{\otimes \bullet}}), \delta \right) \leftarrow \left(\operatorname{End}(\wedge^{\bullet} \underline{\mathfrak{g}_2^{\otimes \bullet}}), [d_2; -] \right) : T^*$$

of cochain complexes are quasi-isomorphisms.

Remark: This lemma holds for every manifold M and any G_{∞} -morphism $T : (\mathfrak{g}_1, d_1) \rightarrow (g_2, d_2)$.

Proof: First we show that T_* is a morphism of complexes. Let $f \in \text{End}(\wedge \underline{\mathfrak{g}_2^{\otimes \bullet}})$ with |f| = k, then

$$T_*([d_1; f]) = T \circ d_1 \circ f - (-1)^k T \circ f \circ d_1$$

= $d_2 \circ (T \circ f) - (-1)^k (T \circ f) \circ d_1$
= $\delta(T_*(f)).$

Let us prove now that T_* is a quasi-isomorphism. For any graded vector space \mathfrak{g} , the space $\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes \bullet}{\mathfrak{g}}$ has the structure of a filtered space where the *m*-level of the filtration is $F^m(\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes \bullet}{\mathfrak{g}}) = \bigoplus_{p_1 + \dots + p_n - 1 \leq m} \overset{\bullet}{\mathfrak{g}} \overset{\otimes p_1}{\mathfrak{g}} \wedge \dots \overset{\bullet}{\mathfrak{g}} \overset{\otimes p_n}{\mathfrak{g}}$. Clearly the differential d_1 and d_2 are compatible with the filtrations on $\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}} \overset{\bullet}{\mathfrak{g}}$ and $\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}} \overset{\bullet}{\mathfrak{g}}$, hence $\operatorname{End}(\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}})$ and $\operatorname{Hom}(\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}}, \wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}})$ are filtered cochain complex. This yields two spectral sequences (lying in the first quadrant) $E_{\bullet}^{\bullet,\bullet}$ and $\widetilde{E}_{\bullet}^{\bullet,\bullet}$ which converge respectively toward the cohomology $H^{\bullet}(\operatorname{End}(\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}}))$ and $H^{\bullet}(\operatorname{Hom}(\wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}}, \wedge \overset{\bullet}{\mathfrak{g}} \overset{\otimes}{\mathfrak{g}}))$. By standard spectral sequence techniques it is enough to

prove that the map $T^0_* : E_0^{\bullet, \bullet} \to \widetilde{E}_0^{\bullet, \bullet}$ induced by T_* on the associated graded is a quasi-isomorphism.

The induced differentials on $E_0^{\bullet,\bullet}$ and $\widetilde{E}_0^{\bullet,\bullet}$ are respectively $[d_1^1, -] = 0$ and $d_2^1 \circ (-) - (-) \circ d_1^1 = b \circ (-)$ where b is the Hochschild coboundary. By cofreeness property we have the following two isomorphisms

$$E_0^{\bullet,\bullet} \cong \operatorname{End}(\mathfrak{g}_1), \qquad \widetilde{E}_0^{\bullet,\bullet} \cong \operatorname{Hom}(\mathfrak{g}_1,\mathfrak{g}_2).$$

The map $T^0_*: E^{\bullet,\bullet}_0 \to \widetilde{E}^0_{\bullet\bullet}$ induced by T_* is $\varphi_{\text{HKR}} \circ (-)$. Let $p: \mathfrak{g}_2 \to \mathfrak{g}_1$ be the projection onto the cohomology, *i.e.* $p \circ \varphi_{\text{HKR}} = \text{id.}$ Let $u: \mathfrak{g}_1 \to \mathfrak{g}_2$ be any map satisfying b(u) = 0and set $v = p \circ u \in \text{End}(\mathfrak{g}_1)$. One can choose a map $w: \mathfrak{g}_1 \to \mathfrak{g}_2$ which satisfies for any $x \in \mathfrak{g}_1$ the following identity

$$\varphi_{\mathbf{HKB}} \circ p \circ u(x) - u(x) = b \circ w(x).$$

It follows that $\varphi_{\text{HKR}}(v)$ has the same class of homology as u which proves the surjectivity of T^0_* in cohomology. The identity $p \circ \varphi_{\text{HKR}} = \text{id}$ implies easily that T^0_* is also injective in cohomology which finish the proof of the lemma for T_* .

The proof that T^* is also a quasi-isomorphism is analogous.

Proof of Theorem 3.1:.

It is easy to check that T - K is a cocycle in $\left(\operatorname{Hom}(\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}, \wedge \overset{\bullet}{\mathfrak{g}_2^{\otimes \bullet}}), \delta\right)$. The complex of cochain $\left(\operatorname{End}(\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}), [d_1, -]\right) \cong \left(\operatorname{Hom}(\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}, \mathfrak{g}_1), [d_1, -]\right)$ is trigraded with $||_1$ being the degree coming from the graduation of \mathfrak{g}_1 and any element x lying in $\underline{\mathfrak{g}_1^{\otimes p_1}} \wedge \cdots \wedge \underline{\mathfrak{g}_1^{\otimes p_q}}$ satisfies $|x|_2 = q - 1, |x|_3 = p_1 + \ldots p_q - q$. In the case $M = \mathbb{R}^n$, the cohomology $H^{\bullet}\left(\operatorname{End}(\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}), [d_1, -]\right)$ is concentrated in bidegree $(||_2, ||_3) = (0, 0)$ (see [Ta], [Hi]). By Lemma 3.2, this is also the case for the cochain complex $\left(\operatorname{Hom}(\wedge \overset{\bullet}{\mathfrak{g}_1^{\otimes \bullet}}, \wedge \overset{\bullet}{\mathfrak{g}_2^{\otimes \bullet}}), \delta\right)$. Thus, its cohomology classes are determined by complex morphisms $(\mathfrak{g}_1, 0) \to (\mathfrak{g}_2, d_2^1)$ and it is enough to prove that T and K determine the same complex morphism $(\mathfrak{g}_1, 0) \to (\mathfrak{g}_2, d_2^1 = b)$ which is clear because T^1 and K^1 are both equal to the Hochschild-Kostant-Rosenberg map.

Remark. It is possible to have an explicit formula for the map h in Theorem .3.1. In fact the quasi-isomorphism coming from Lemma 3.2 can be made explicit using explicit homotopy formulae for the Hochschild-Kostant-Rosenberg map (see [Ha] for example) and deformation retract techniques (instead of spectral sequences) as in [Ka]. The same techniques also apply to give explicit formulae for the quasi-isomorphism giving the acyclicity of $\left(\operatorname{End}(\wedge^{\bullet}\mathfrak{g}_{1}^{\otimes \bullet}), [d_{1}; -]\right)$ in the proof of theorem 3.1 (see [GH] for example)

References

- [BFFLS1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, Quantum mechanics as a deformation of classical mechanics, Lett. Math. Phys. 1 (1975), 521–530.
- [BFFLS2] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, Deformation theory and quantization, I and II, Ann. Phys. 111 (1977), 61–151.
- [CFT] A. S. Cattaneo, G. Felder, L. Tomassini, From local to global deformation quantization of Poisson manifolds, Duke Math. J. 115 (2002), 329–352.
- [Gi] G. Ginot, Homologie et modèle minimal des algèbres de Gerstenhaber, Ann. Math. Blaise Pascal 11 (2004), 91–127.
- [GH] G. Ginot, G. Halbout, A formality theorem for Poisson manifold, Lett. Math. Phys. 66 (2003), 37–64.
- [GK] V. Ginzburg, M. Kapranov, Koszul duality for operads, Duke Math. J. 76 (1994), 203–272.
- [Ha] G. Halbout, Formule d'homotopie entre les complexes de Hochschild et de de Rham, Compositio Math. 126 (2001), 123–145.
- [Hi] V. Hinich, Tamarkin's proof of Kontsevich's formality theorem, Forum Math. 15 (2003), 591–614.
- [HKR] G. Hochschild, B. Kostant and A. Rosenberg, *Differential forms on regular affine algebras*, Transactions AMS **102** (1962), 383-408.
- [Ka] C. Kassel, Homologie cyclique, caractère de Chern et lemme de perturbation, J. Reine Angew. Math. 408 (1990), 159–180.
- [Ko1] M. Kontsevich, Formality conjecture. Deformation theory and symplectic geometry, Math. Phys. Stud. 20 (1996), 139–156.
- [Ko2] M. Kontsevich, Deformation quantization of Poisson manifolds, Lett. Math. Phys. 66 (2003), 157–216.
- [KS] M. Kontsevich, Y. Soibelman, Deformations of algebras over operads and the Deligne conjecture, Math. Phys. Stud. 21 (2000), 255–307.
- [Kos] J. L. Koszul, Crochet de Schouten-Nijenhuis et cohomologie in "Elie Cartan et les mathématiques d'aujourd'hui", Astérisque (1985), 257-271.
- [LS] T. Lada, J. D. Stasheff, Introduction to SH Lie algebras for physicists, nternat. J. Theoret. Phys. 32 (1993), 1087-1103.
- [Ta] D. Tamarkin, Another proof of M. Kontsevich's formality theorem, math. QA/9803025.
- [TS] D. Tamarkin, B Tsygan, Noncommutative differential calculus, homotopy BV algebras and formality conjectures, Methods Funct. Anal. Topology 6 (2000), 85–97.

(a) LABORATOIRE ANALYSE GÉOMÉTRIE ET APPLICATIONS, UNIVERSITÉ PARIS 13, CENTRE DES MATHÉMATIQUES ET DE LEURS APPLICATIONS, ENS CACHAN, 61, AV. DU PRÉSIDENT WILSON, 94230 CACHAN, FRANCE
E-MAIL:GINOT@CMLA.ENS-CACHAN.FR

 ^(b) Institut de Recherche Mathématique Avancée, Université Louis Pasteur et CNRS
 7, rue René Descartes, 67084 Strasbourg, France E-Mail:halbout@math.u-strasbg.fr