

PROBABILISTIC ASPECTS OF AL-SALAM-CHIHARA POLYNOMIALS

WŁODZIMIERZ BRYC, WOJCIECH MATYSIAK, AND PAWEŁ J. SZABŁOWSKI

ABSTRACT. We solve the connection coefficient problem between the Al-Salam–Chihara polynomials and the q -Hermite polynomials, and we use the resulting identity to answer a question from probability theory. We also derive the distribution of some Al-Salam–Chihara polynomials, and compute determinants of related Hankel matrices.

1. INTRODUCTION AND MAIN IDENTITY

The aim of the paper is to point out the connection of Al-Salam–Chihara polynomials with a regression problem in probability, and to use it to give a new simple derivation of their density. Our approach exploits identity (1.8) below, which connects the Al-Salam–Chihara polynomials to the continuous q -Hermite polynomials. This connection is more direct and elementary but less general than the technique of attachment exploited in [BI96, Section 2]. We also compute determinants of Hankel matrices with entries which are linear combinations of the q -Hermite polynomials.

The Al-Salam–Chihara polynomials were introduced in [ASC76]. We are interested in the renormalized Al-Salam–Chihara polynomials $\{p_n(x|q, a, b)\}$, which are defined by the following three term recurrence relation

$$(1.1) \quad p_{n+1}(x) = (x - aq^n) p_n(x) - (1 - bq^{n-1}) [n]_q p_{n-1}(x) \quad (n \geq 0),$$

with the usual initial conditions $p_{-1}(x) = 0$, $p_0(x) = 1$. Here, we use the standard notation

$$\begin{aligned} [n]_q &= 1 + q + \cdots + q^{n-1}, \\ [n]_q! &= [1]_q [2]_q \cdots [n]_q, \\ \begin{bmatrix} n \\ k \end{bmatrix}_q &= \frac{[n]_q!}{[n-k]_q! [k]_q!} \end{aligned}$$

with the usual conventions $[0]_q = 0$, $[0]_q! = 1$.

For $|q| < 1$, their generating function

$$f(t, x|q, a, b) = \sum_{n=0}^{\infty} \frac{t^n}{[n]_q!} p_n(x|q, a, b)$$

Date: April 11, 2003 .

1991 Mathematics Subject Classification. Primary: 33D45 Secondary: 05A30, 15A15, 42C05.

Key words and phrases. q -Hermite polynomials, matrix of moments, orthogonal polynomials, determinants, polynomial regression.

is given by

$$(1.2) \quad f(t, x|q, a, b) = \prod_{k=0}^{\infty} \frac{1 - (1-q)atq^k + (1-q)bt^2q^{2k}}{1 - (1-q)xtq^k + (1-q)t^2q^{2k}};$$

compare [AI84, (3.6) and (3.10)].

The corresponding (renormalized) continuous q -Hermite polynomials $H_n(x|q) = p_n(x|q, 0, 0)$ satisfy the three term recurrence relation

$$(1.3) \quad H_{n+1}(x) = xH_n(x) - [n]_q H_{n-1}(x).$$

For $|q| < 1$ their generating function $\phi(t, x|q) = \sum_{n=0}^{\infty} \frac{t^n}{[n]_q!} H_n(x|q)$ is

$$(1.4) \quad \phi(t, x|q) = \prod_{k=0}^{\infty} (1 - (1-q)xtq^k + (1-q)t^2q^{2k})^{-1}.$$

Of course, these are well known special cases of (1.1) and (1.2), see [ISV87, (2.11) and (2.12)] which we state here for further reference.

We will also use polynomials $\{B_n(x|q)\}$ defined by the three term recurrence relation

$$(1.5) \quad B_{n+1}(x) = -q^n x B_n(x) + q^{n-1} [n]_q B_{n-1}(x) \quad (n \geq 0)$$

with the usual initial conditions $B_{-1} = 0, B_0 = 1$. These polynomials are related to the q -Hermite polynomials by

$$(1.6) \quad B_n(x|q) = \begin{cases} i^n q^{n(n-2)/2} H_n(i\sqrt{q}x|\frac{1}{q}) & \text{if } q > 0 \\ (-1)^{n(n-1)/2} |q|^{n(n-2)/2} H_n(-\sqrt{|q}|x|\frac{1}{q}) & \text{if } q < 0 \end{cases}.$$

Indeed, both sides of (1.6) are equal at $n = -1, 0$ and satisfy (1.5) for $n \geq 0$.

Their generating function $\psi(t, x|q) = \sum_{n=0}^{\infty} \frac{t^n}{[n]_q!} B_n(x|q)$ is given by

$$(1.7) \quad \psi(t, x|q) = \prod_{k=0}^{\infty} (1 - (1-q)xtq^k + (1-q)t^2q^{2k}).$$

To see this, we compute the q -derivative for $t \neq 0, q \neq 1$ and use (1.5) to get

$$\begin{aligned} \frac{\psi(t, x|q) - \psi(qt, x|q)}{(1-q)t} &= \sum_{n=0}^{\infty} \frac{t^n}{[n]_q!} B_{n+1}(x|q) = \\ &= -x \sum_{n=0}^{\infty} \frac{(qt)^n}{[n]_q!} B_n(x|q) + t \sum_{n=0}^{\infty} \frac{(qt)^n}{[n]_q!} B_n(x|q) = (-x + t)\psi(qt, x|q). \end{aligned}$$

Thus

$$\begin{aligned} \psi(t, x|q) &= \psi(qt, x|q) (1 - (1-q)xt + (1-q)t^2) = \\ &= \psi(q^n t, x|q) \prod_{k=0}^{n-1} (1 - (1-q)xtq^k + (1-q)t^2q^{2k}), \end{aligned}$$

and (1.7) follows since $\psi(q^n t, x|q) \rightarrow \psi(0, x|q) = 1$ as $n \rightarrow \infty$ when $|q| < 1$.

We now point out the mutual relationship between the Al-Salam–Chihara polynomials $\{p_n(x|q, a, b)\}$ and the polynomials $\{H_n(x|q)\}_{n \geq 0}$ and $\{B_n(x|q)\}_{n \geq 0}$.

Theorem 1. *For all $a, b, q \in \mathbb{C}$, $b \neq 0$, and $n \geq 1$ we have*

$$(1.8) \quad p_n(x|q, a, b^2) = \sum_{k=1}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_q b^{n-k} B_{n-k}(\frac{a}{b}|q) \left(H_k(x|q) - b^k H_k(\frac{a}{b}|q) \right)$$

Proof. From recurrence relations (1.1), (1.3), and (1.5), it is clear that $p_n(x|q, a, b)$, $H_n(x|q)$, and $B_n(x|q)$ are given by polynomial expressions in variable q . The q -binomial coefficient $\begin{bmatrix} n \\ k \end{bmatrix}_q$ is also a polynomial in q . Therefore, we see that identity (1.8) is equivalent to a polynomial identity in variable $q \in \mathbb{C}$. Hence it is enough to prove that (1.8) holds true for all $|q| < 1$. When $|q| < 1$, inspecting (1.2), (1.4), and (1.7) we notice that

$$(1.9) \quad f(t, x|q, a, b^2) = \psi(bt, a/b|q)\phi(t, x|q),$$

and

$$(1.10) \quad \psi(t, x|q)\phi(t, x|q) = 1.$$

Therefore,

$$f(t, x|q, a, b^2) = 1 + \psi(bt, a/b|q) (\phi(t, x|q) - \phi(bt, a/b|q)),$$

which is valid for all small enough $|t|$. Comparing the coefficients at t^n for $n \geq 1$ and taking into account that $H_k(x|q) - b^k H_k(\frac{a}{b}|q) = 0$ for $k = 0$, we get (1.8). ■

Remark 1. One could split (1.8) into the following two separate identities, which are implied by (1.9) and (1.10) respectively:

$$(1.11) \quad \forall n \geq 0 : p_n(x|q, a, b^2) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q b^{n-k} B_{n-k}(\frac{a}{b}|q) H_k(x|q),$$

$$(1.12) \quad \forall n \geq 1 : \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q B_{n-k}(x) H_k(x|q) = 0.$$

Formula (1.11) is a renormalized inverse of formula [IRS99, (4.7)], who express the q -Hermite polynomials as linear combinations of Al-Salam-Chihara polynomials. Formula (1.12) resembles [Car56, (2.28)], who considers q -Hermite polynomials of the form $h_n(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q x^k$, paired with $b_n(x|q) = h_n(x|1/q)$.

2. PROBABILISTIC ASPECTS

Quadratic regression questions in the paper [Bry01] lead to the problem of determining all probability distributions μ which are defined indirectly by the relationships

$$(2.1) \quad \int H_n(x|q)\mu(dx) = \rho^n H_n(y|q) \quad n = 1, 2, \dots,$$

where $y, \rho, q \in \mathbb{R}$ are fixed parameters, and $\{H_n\}_{n \geq 0}$ is the family of the q -Hermite polynomials.

Our next result shows that this problem can be solved using the Al-Salam-Chihara polynomials.

Theorem 2. If $\mu = \mu(dx|\rho, y)$ satisfies (2.1), then its orthogonal polynomials are Al-Salam-Chihara polynomials $\{p_n(x|q, a, b)\}$ with $a = \rho y$, $b = \rho^2$.

Proof. Recall that $H_n(x|q) = p_n(x|q, 0, 0)$. Thus if $\rho = 0$ then (2.1) implies that $\int p_n(x|q, a, b)\mu(dx) = 0$ for all $n = 1, 2, \dots$. Suppose now that $\rho \neq 0$. Combining (1.8) with (2.1) we get

$$\int p_n(x|q, a, b)\mu(dx) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \rho^{n-k} B_{n-k}(y|q) \int (H_k(x|q) - \rho^k H_k(y|q)) \mu(dx) = 0$$

for all $n = 1, 2, \dots$. Since $\{p_n\}$ satisfy a three step recurrence, this implies $\int p_k(x)p_n(x)\mu(dx) = 0$ for all $0 \leq k < n$. ■

Next we answer an unresolved case from [Bry01].

Corollary 1. *If $\rho, q, y \in \mathbb{R}$ are such that $q > 1$ and $\rho \neq 0$ then (2.1) has no probabilistic solution μ .*

Proof. Suppose that μ is positive and solves (2.1). Then its monic orthogonal polynomials satisfy the three term recurrence relation (1.1)

$$(2.2) \quad p_{n+1}(x) = (x - \rho y q^n) p_n(x) - (1 - \rho^2 q^{n-1}) [n]_q p_{n-1}(x).$$

For a positive measure, the coefficients at the third term in the recurrence must be non-negative, $1 - \rho^2 q^{n-1} \geq 0$ for all n , which is false. ■

From Theorem 2 it follows that if the solution of (2.1) exists, then it is given by the distribution of the Al-Salam-Chihara polynomials. The distribution of the Al-Salam-Chihara polynomials is derived in [AI84, Chapter 3]. However, in [Bry01, Proposition 8.1] we found the solution of (2.1) which relies solely on the facts about the q -Hermite polynomials. We repeat the latter argument here, and then use it to re-derive the distribution of the corresponding Al-Salam-Chihara polynomials.

Corollary 2. *If $\rho, q, y \in \mathbb{R}$ are such that $|\rho| < 1, |q| < 1$, and $y^2(1 - q) < 4$, then the probabilistic solution of (2.1) is given by the absolutely continuous measure μ with the density on $x^2 < 4/(1 - q)$ given by*

$$\frac{\sqrt{1 - q}}{2\pi\sqrt{4 - (1 - q)x^2}} \prod_{k=0}^{\infty} \frac{(1 - \rho^2 q^k) (1 - q^{k+1}) ((1 + q^k)^2 - (1 - q)x^2 q^k)}{(1 - \rho^2 q^{2k})^2 - (1 - q)\rho q^k(1 + \rho^2 q^{2k})xy + (1 - q)\rho^2(x^2 + y^2)q^{2k}}$$

Proof. The distribution of the q -Hermite polynomials $H_n(x|q)$ is supported on $x^2 < 4/(1 - q)$ with the density

$$f_H(x) = \frac{\sqrt{1 - q}}{2\pi\sqrt{4 - (1 - q)x^2}} \prod_{k=0}^{\infty} ((1 + q^k)^2 - (1 - q)x^2 q^k) \prod_{k=0}^{\infty} (1 - q^{k+1}),$$

see [ISV87, (2.15)]. Moreover, since $|H_n(x)| \leq C_q(n + 1)(1 - q)^{-n/2}$ when $x^2, y^2 \leq 4/(1 - q)$, the series

$$(2.3) \quad g_H(x, y, \rho) = \sum_{n=0}^{\infty} \frac{\rho^n}{[n]_q!} H_n(x) H_n(y)$$

converges uniformly and defines the Poisson-Mehler kernel which is given by

$$(2.4) \quad g_H(x, y, \rho) = \prod_{k=0}^{\infty} \frac{(1 - \rho^2 q^k)}{(1 - \rho^2 q^{2k})^2 - (1 - q)\rho q^k(1 + \rho^2 q^{2k})xy + (1 - q)\rho^2(x^2 + y^2)q^{2k}};$$

this is the renormalized version of the well known result, see eg. [IS88, (2.2)] who consider the q -Hermite polynomials given by $\{(1-q)^{n/2}H_n(2x/\sqrt{1-q}|q)\}$ instead of our $\{H_n(x|q)\}$.

Since (1.3) implies that $\int H_n^2(x|q)f_H(x)dx = [n]_q!$, it follows from (2.3) that

$$\int_{-2/\sqrt{1-q}}^{2/\sqrt{1-q}} H_n(x|q)g_H(x, y, \rho)f_H(x)dx = \rho^n H_n(y).$$

■

Corollary 3. *If $q, a, b \in \mathbb{R}$ are such that $|q| < 1$, $0 < b < 1$, and $a^2(1-q) < 4b$, then the distribution of the Al-Salam-Chihara polynomials $\{p_n(x|q, a, b)\}$ is absolutely continuous with the density on $x^2 < 4/(1-q)$ given by*

$$\frac{\sqrt{1-q}}{2\pi\sqrt{4-(1-q)x^2}} \prod_{k=0}^{\infty} \frac{(1-bq^k)(1-q^{k+1})((1+q^k)^2 - (1-q)x^2q^k)}{(1-bq^{2k})^2 - (1-q)aq^k(1+bq^{2k})x + (1-q)(bx^2 + a^2)q^{2k}}.$$

Proof. By Theorem 2, the distribution of polynomials p_n solves (2.1) with $\rho = \sqrt{b}$, $y = a/\rho$. Thus the formula follows from Corollary 2. ■

3. DETERMINANTS OF HANKEL MATRICES

In this section we are interested in calculating the determinants of the Hankel matrices

$$M_n = [m_{i+j}]_{i,j=0,\dots,n-1},$$

where $m_i = \int x^i \mu(dx)$ are the moments of certain (perhaps signed) measure μ . It is well known that for positive measures we must have $\det M_n \geq 0$, and that these determinants can be read out from the three term recurrence for the corresponding monic orthogonal polynomials.

Consider first the moments $m_k(y) = \int x^k \mu(dx)$ of the (perhaps signed) measure $\mu = \mu_{y,\rho}$, which solves (2.1). Then $m_k(y)$ are polynomials of degree k in variable y , and can be written as follows. Let $a_{n,2i}$, $i \leq \lfloor n/2 \rfloor$ be the coefficients in the expansion of monomial x^n into the q -Hermite polynomials,

$$x^n = \sum_{i=0}^{\lfloor n/2 \rfloor} a_{n,2i} H_{n-2i}(x|q), \quad n \geq 0.$$

Then

$$m_n(y) = \int x^n d\mu(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \rho^{n-2k} a_{n,2k} H_{n-2k}(y|q).$$

Let S_n be the Hankel matrix of moments $m_k(y)$,

$$S_n(y|q, \rho) = \begin{bmatrix} m_0(y) & m_1(y) & \dots & m_{n-1}(y) \\ m_1(y) & m_2(y) & & \\ \vdots & & \ddots & \vdots \\ m_{n-1}(y) & & \dots & m_{2n-2}(y) \end{bmatrix}.$$

It is well known that $\det S_n$ is the product of the coefficients at the third term of (2.2), which implies the following.

Corollary 4. $\det S_{n+1} / \det S_n = [n]_q! \prod_{i=1}^n (1 - \rho^2 q^{i-1})$.

Our second Hankel matrix has an even simpler form. As indicated in [IS97], [IS02] the q -Hermite polynomials can be viewed as moments of a signed measure, $H_n(x|q) = \int u^n \mu(du|x, q)$. It turns out that measure $\mu(du|x, q)$ cannot be positive even for a single value of x . To see this, consider the following $n \times n$ matrices

$$M_n(x|q) = \begin{bmatrix} H_0(x|q) & H_1(x|q) & H_2(x|q) & \dots & H_{n-1}(x|q) \\ H_1(x|q) & H_2(x|q) & & & H_n(x|q) \\ H_2(x|q) & & \ddots & & H_{n+1}(x|q) \\ \vdots & & & \ddots & \vdots \\ H_{n-1}(x|q) & H_n(x|q) & & \dots & H_{2n-2}(x|q) \end{bmatrix}$$

The following q -generalization of [Kra99, (3.55)] shows that the determinants $\det M_n(x|q)$ are free of the variable x , and take negative values.

Theorem 3.

$$\frac{\det M_{n+1}}{\det M_n} = (-1)^n q^{n(n-1)/2} [n]_q!$$

Proof. Using (1.3), we row-reduce the first column of the matrix. Namely, from the second row of M_{n+1} , we subtract the first one multiplied by x . Similarly, for $i \geq 3$, we subtract x times row $i-1$ and add the $i-2$ -th row multiplied by $[i-1]_q$. Taking (1.3) into account, $\det M_{n+1}(x|q)$ becomes

$$\det \begin{bmatrix} H_0 & H_1 & H_2 & \dots & H_n \\ 0 & ([0] - [1])H_0 & ([0] - [2])H_1 & \dots & ([0] - [n])H_{n-1} \\ 0 & ([1] - [2])H_1 & ([1] - [3])H_2 & \dots & ([1] - [n+1])H_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & ([n-1] - [n])H_{n-1} & ([n-1] - [n+1])H_n & \dots & ([n-1] - [2n-1])H_{2n} \end{bmatrix}$$

Now, we use the fact that for $m \leq n$ we have

$$[n]_q - [m]_q = q^m [n-m]_q.$$

Thus $\det M_{n+1}(x|q)$ becomes

$$\det \begin{bmatrix} H_0 & H_1 & H_2 & \dots & H_{n-1} \\ 0 & -[1]H_0 & -[2]H_1 & \dots & -[n]H_{n-1} \\ 0 & -q[1]H_1 & -q[2]H_2 & \dots & -q[n]H_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & -q^{n-1}[1]H_{n-1} & -q^{n-1}[2]H_n & \dots & -q^{n-1}[n]H_{2n} \end{bmatrix}$$

Expanding $\det M_{n+1}$ with respect to the first column, and factoring out the common factors $-q^{i-1}$ from the i -th row and $[j]_q$ from the j -th column of the resulting matrix, we get

$$\begin{aligned} \det M_{n+1} &= (-1)^n q^{\sum_{i=1}^{n-1} i} \prod_{j=1}^n [j]_q \det M_n \\ &= (-1)^n q^{n(n-1)/2} [n]_q! \det M_n. \end{aligned}$$

■

Formula stated in Corollary 4 was originally discovered through symbolic computations and motivated this paper. We were unable to find a direct algebraic proof along the lines of the proof of Theorem 3, and our search for the explanation of

why $\det S_n(y)$ does not depend on y lead us to Al-Salam–Chihara polynomials and identity (1.8).

The fact that Hankel determinants formed of certain linear combinations of the q -Hermite polynomials do not depend on the argument of these polynomials as exposed in Theorem 3 and Corollary 4 is striking and unexpected to us. A natural question arises whether other linear combinations have this property.

Acknowledgement Part of the research of WB was conducted while visiting Warsaw University of Technology. The authors thank Jacek Wołowski for several helpful discussions.

REFERENCES

- [AI84] Richard Askey and Mourad Ismail. Recurrence relations, continued fractions, and orthogonal polynomials. *Mem. Amer. Math. Soc.*, 49(300):iv+108, 1984.
- [ASC76] W. A. Al-Salam and T. S. Chihara. Convolutions of orthonormal polynomials. *SIAM J. Math. Anal.*, 7(1):16–28, 1976.
- [BI96] Christian Berg and Mourad E. H. Ismail. q -Hermite polynomials and classical orthogonal polynomials. *Canad. J. Math.*, 48(1):43–63, 1996.
- [Bry01] Włodzimierz Bryc. Stationary random fields with linear regressions. *Annals of Probability*, 29:504–519, 2001.
- [Car56] L. Carlitz. Some polynomials related to theta functions. *Ann. Mat. Pura Appl. (4)*, 41:359–373, 1956.
- [IRS99] Mourad E. H. Ismail, Mizan Rahman, and Dennis Stanton. Quadratic q -exponentials and connection coefficient problems. *Proc. Amer. Math. Soc.*, 127(10):2931–2941, 1999.
- [IS88] Mourad E. H. Ismail and Dennis Stanton. On the Askey-Wilson and Rogers polynomials. *Canad. J. Math.*, 40(5):1025–1045, 1988.
- [IS97] Mourad E. H. Ismail and Dennis Stanton. Classical orthogonal polynomials as moments. *Can. J. Math.*, 49(3):520–542, 1997.
- [IS02] Mourad E. H. Ismail and Dennis Stanton. q -integral and moment representations for q -orthogonal polynomials. *Canad. J. Math.*, 54(4):709–735, 2002.
- [ISV87] M. E. H. Ismail, D. Stanton, and G. Viennot. The combinatorics of q -Hermite polynomials and the Askey-Wilson integral. *Europ. J. Combinatorics*, 8:379–392, 1987.
- [Kra99] Christian Krattenthaler. Advanced determinant calculus. *Sém. Lothar. Combin.*, 42:Art. B42q, 67 pp. (electronic), 1999.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CINCINNATI, P.O. BOX 210025, CINCINNATI, OH 45221–0025

E-mail address: Wlodzimierz.Bryc@UC.edu

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, PL. POLITECHNIKI 1, 00-661 WARSZAWA, POLAND

E-mail address: wmatysiak@elka.pw.edu.pl

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, PL. POLITECHNIKI 1, 00-661 WARSZAWA, POLAND

E-mail address: pszablowski@elka.pw.edu.pl