

# EXT VANISHING AND INFINITE AUSLANDER-BUCHSBAUM

PETER JØRGENSEN

ABSTRACT. A vanishing theorem is proved for Ext groups over non-commutative graded algebras. Along the way, an “infinite” version is proved of the non-commutative Auslander-Buchsbaum theorem.

## 0. INTRODUCTION

Let  $R$  be a noetherian local commutative ring, and let  $X$  be a finitely generated  $R$ -module of finite projective dimension. The classical Auslander-Buchsbaum theorem states

$$\mathrm{pd} X = \mathrm{depth} R - \mathrm{depth} X.$$

This can also be phrased as an Ext vanishing theorem, namely, if  $M$  is any  $R$ -module, then

$$\mathrm{Ext}_R^i(X, M) = 0 \quad \text{for } i > \mathrm{depth} R - \mathrm{depth} X. \quad (1)$$

In [1] is proved a surprising variation of this: Suppose that  $R$  is complete in the  $\mathfrak{m}$ -adic topology. Then equation (1) remains true if  $X$  is *any*  $R$ -module of finite projective dimension, provided  $M$  is finitely generated. In other words, the condition of being finitely generated is shifted from  $X$  to  $M$ .

In theorem 2.4 below, this result will be generalized to the situation of a non-commutative noetherian  $\mathbb{N}$ -graded connected algebra.

The route goes through an “infinite” version of the non-commutative Auslander-Buchsbaum theorem, given in theorem 1.4. This result is a substantial improvement of the original non-commutative Auslander-Buchsbaum theorem, as given in [3, thm. 3.2], in that the condition of dealing only with finitely generated modules is dropped.

The notation of this paper is standard, and is already on record in several places such as [2] or [3]. So I will not say much, except that throughout,  $k$  is a field, and  $A$  is a noetherian  $\mathbb{N}$ -graded connected

---

2000 *Mathematics Subject Classification.* 16E30, 16W50.

*Key words and phrases.*  $k$ -flat dimension, depth, infinite non-commutative Auslander-Buchsbaum theorem, Ext groups, vanishing theorem.

$k$ -algebra. However, let me give one important word of caution: Everything in sight is graded. So for instance,  $D(A)$  stands for  $D(\mathbf{Gr} A)$ , the derived category of the abelian category  $\mathbf{Gr}(A)$  of  $\mathbb{Z}$ -graded  $A$ -left-modules and graded homomorphisms of degree zero.

# 1. AUSLANDER-BUCHSBAUM

**Definition 1.1.** For  $X$  in  $D(A)$ , define the  $k$ -flat dimension by

$$\mathrm{k.f.d} X = -\inf k \overset{\mathrm{L}}{\otimes}_A X.$$

**Remark 1.2.** Using a minimal free resolution, it is easy to see that if the cohomology of  $X$  is bounded and finitely generated, then

$$\mathrm{k.f.d} X = \mathrm{fd} X = \mathrm{pd} X,$$

where  $\mathrm{fd}$  stands for flat dimension and  $\mathrm{pd}$  stands for projective dimension.

In the following lemma is used  $D^b(A)$ , the full subcategory of  $D(A)$  consisting of complexes with bounded cohomology, and  $D^+(A^{\mathrm{op}})$ , the full subcategory of  $D(A^{\mathrm{op}})$  consisting of complexes with cohomology vanishing in low degrees.

**Lemma 1.3.** *Let  $X$  in  $D^b(A)$  have  $\mathrm{fd} X < \infty$ , and let  $S$  in  $D^+(A^{\mathrm{op}})$  have  $\dim_k h^i(S) < \infty$  for each  $i$ . Then*

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = \inf S - \mathrm{k.f.d} X.$$

*Proof.* Observe that  $\mathrm{fd} X < \infty$  implies

$$\inf k \overset{\mathrm{L}}{\otimes}_A X > -\infty,$$

hence

$$\mathrm{k.f.d} X < \infty. \tag{2}$$

If

$$\inf S = \infty \tag{3}$$

then  $S$  is zero. Then  $S \overset{\mathrm{L}}{\otimes}_A X$  is also zero, so

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = \infty. \tag{4}$$

Combining equations (2), (3), and (4) shows that for  $\inf S = \infty$ , the lemma's equation reads

$$\infty = \infty - (\text{something smaller than } \infty)$$

which is true.

So for the rest of the proof, I can assume  $\inf S < \infty$ , that is,  $S$  is non-zero. Note that  $S$  is in  $D^+(A^{\text{op}})$ , so  $\inf S > -\infty$ , so  $\inf S$  is a finite number.

Let me first do the case where  $S$  is concentrated in degree zero. Here  $S$  is just a graded  $A$ -right-module which is finite dimensional over  $k$ , and the lemma claims

$$\inf S \overset{\text{L}}{\otimes}_A X = -\text{k.fd } X = \inf k \overset{\text{L}}{\otimes}_A X. \quad (5)$$

To prove this, note that  $\dim_k S < \infty$  gives that there is a short exact sequence  $0 \rightarrow k(\ell) \rightarrow S \rightarrow \tilde{S} \rightarrow 0$  of graded  $A$ -right-modules. This gives a distinguished triangle  $k(\ell) \rightarrow S \rightarrow \tilde{S} \rightarrow$  in  $D(A^{\text{op}})$ , and tensoring with  $X$  and taking the cohomology long exact sequence gives a sequence consisting of pieces

$$h^i(k(\ell) \overset{\text{L}}{\otimes}_A X) \rightarrow h^i(S \overset{\text{L}}{\otimes}_A X) \rightarrow h^i(\tilde{S} \overset{\text{L}}{\otimes}_A X).$$

Induction on  $\dim_k S$  now shows that the lowest  $i$  with

$$h^i(k \overset{\text{L}}{\otimes}_A X) \neq 0$$

equals the lowest  $i$  with

$$h^i(S \overset{\text{L}}{\otimes}_A X) \neq 0,$$

and equation (5) follows. The induction works because  $\inf k \overset{\text{L}}{\otimes}_A X > -\infty$ , and even works for  $\inf k \overset{\text{L}}{\otimes}_A X = \infty$ .

Let me next do the case where  $S$  is general. There is a standard spectral sequence

$$E_2^{pq} = h^p(h^q(S) \overset{\text{L}}{\otimes}_A X) \Rightarrow h^{p+q}(S \overset{\text{L}}{\otimes}_A X).$$

Since  $\text{fd } X < \infty$ , the spectral sequence is first quadrant up to shift, hence converges. Now,  $\dim_k h^q(S) < \infty$  holds for each  $q$ , so if  $h^q(S)$  is non-zero, then the special case of the lemma dealt with above applies to  $h^q(S) \overset{\text{L}}{\otimes}_A X$  and shows

$$\inf h^q(S) \overset{\text{L}}{\otimes}_A X = -\text{k.fd } X. \quad (6)$$

There are now two cases: First, if

$$\text{k.fd } X = -\infty \quad (7)$$

then equation (6) gives that if  $h^q(S)$  is non-zero, then  $\inf h^q(S) \overset{\text{L}}{\otimes}_A X = \infty$ , that is,  $h^p(h^q(S) \overset{\text{L}}{\otimes}_A X)$  is zero for each  $p$ . Of course this also holds for  $h^q(S)$  equal to zero, and so in the spectral sequence,  $E_2^{pq}$

is identically zero. Therefore the limit  $\mathrm{h}^{p+q}(S \overset{\mathrm{L}}{\otimes}_A X)$  of the spectral sequence is also zero, so  $S \overset{\mathrm{L}}{\otimes}_A X$  is zero, so

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = \infty. \quad (8)$$

But  $\inf S$  is a finite number, and combining this with equations (7) and (8) says that the lemma's equation reads

$$\infty = (\text{a finite number}) - (-\infty)$$

which is true.

Secondly, if

$$\mathrm{k.f.d} X > -\infty,$$

then equation (6) says that if  $\mathrm{h}^q(S)$  is non-zero, then  $\mathrm{h}^p(\mathrm{h}^q(S) \overset{\mathrm{L}}{\otimes}_A M)$  is non-zero for  $p = -\mathrm{k.f.d} X$ , but zero for  $p < -\mathrm{k.f.d} X$ . And of course, if  $\mathrm{h}^q(S)$  is zero, then  $\mathrm{h}^p(\mathrm{h}^q(S) \overset{\mathrm{L}}{\otimes}_A X)$  is zero for each  $p$ . So in the spectral sequence,  $E_2^{pq}$  is non-zero for  $p = -\mathrm{k.f.d} X$  and  $q = \inf S$ , but zero for lower  $p$  or  $q$ . Hence  $E_2^{-\mathrm{k.f.d} X, \inf S}$  can be used in a standard corner argument which shows that the lowest non-zero term in the limit  $\mathrm{h}^{p+q}(S \overset{\mathrm{L}}{\otimes}_A X)$  of the spectral sequence has degree  $p + q = -\mathrm{k.f.d} X + \inf S$ . Hence

$$\inf S \overset{\mathrm{L}}{\otimes}_A X = -\mathrm{k.f.d} X + \inf S,$$

proving the lemma's equation.  $\square$

**Theorem 1.4** (Infinite Auslander-Buchsbaum). *Assume that  $A$  satisfies  $\dim_k \mathrm{Ext}_A^i(k, A) < \infty$  for each  $i$ . Let  $X$  in  $\mathrm{D}^b(A)$  have  $\mathrm{fd} X < \infty$ . Then*

$$\mathrm{depth} X = \mathrm{depth} A - \mathrm{k.f.d} X.$$

*Proof.* I have

$$\mathrm{RHom}_A(k, X) \cong \mathrm{RHom}_A(k, A \overset{\mathrm{L}}{\otimes}_A X) \cong \mathrm{RHom}_A(k, A) \overset{\mathrm{L}}{\otimes}_A X,$$

where the second  $\cong$  holds because it holds if  $X$  is just a graded flat module, so also holds if  $X$  is quasi-isomorphic to a bounded complex of graded flat modules, and this is the case since  $X$  is in  $\mathrm{D}^b(A)$  and has  $\mathrm{fd} X < \infty$ .

Thus

$$\begin{aligned}
\text{depth } X &= \inf \text{RHom}_A(k, X) \\
&= \inf \text{RHom}_A(k, A) \overset{\text{L}}{\otimes}_A X \\
&\stackrel{(a)}{=} \inf \text{RHom}_A(k, A) - \text{k.fd } X \\
&= \text{depth } A - \text{k.fd } X,
\end{aligned}$$

where (a) is by lemma 1.3. The lemma applies because  $\text{RHom}_A(k, A)$  is in  $\mathbf{D}^+(A^{\text{op}})$ , and has  $\dim_k \text{h}^i \text{RHom}_A(k, A) = \dim_k \text{Ext}_A^i(k, A) < \infty$  for each  $i$  by assumption.  $\square$

**Remark 1.5.** Theorem 1.4 even holds for  $\text{depth } A = \infty$ , where the theorem states  $\text{depth } X = \infty$ .

On the other hand, suppose  $\text{depth } A < \infty$ . Then it is easy to see that it makes sense to rearrange the equation in theorem 1.4 as

$$\text{k.fd } X = \text{depth } A - \text{depth } X.$$

If the cohomology of  $X$  is bounded and finitely generated, then this equation reads

$$\text{pd } X = \text{depth } A - \text{depth } X$$

by remark 1.2. This is the original non-commutative Auslander-Buchsbaum theorem, as proved in [3, thm. 3.2].

## 2. EXT VANISHING

**Lemma 2.1.** *Let  $X$  in  $\mathbf{D}^b(A)$  have  $\text{fd } X < \infty$ , and let  $T$  in  $\mathbf{D}^+(A^{\text{op}})$  be so that  $\text{h}^i(T)$  is a graded torsion module for each  $i$ . Then*

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T - \text{k.fd } X.$$

*Proof.* As in the proof of lemma 1.3, it is easy to see that for  $\inf T = \infty$ , the lemma's inequality trivially reads  $\infty \geq \infty$ . So for the rest of the proof I can assume  $\inf T < \infty$ . Note that  $T$  is in  $\mathbf{D}^+(A^{\text{op}})$ , so  $\inf T > -\infty$ , so  $\inf T$  is a finite number.

Let me first do the case where  $T$  is concentrated in degree zero. Here  $T$  is just a graded torsion module, and the lemma claims

$$\inf T \overset{\text{L}}{\otimes}_A X \geq -\text{k.fd } X = \inf k \overset{\text{L}}{\otimes}_A X.$$

If  $F$  is a flat resolution of  $X$ , then this amounts to

$$\inf T \otimes_A F \geq \inf k \otimes_A F. \tag{9}$$

To prove this, note that as  $T$  is a graded torsion module, it is the colimit of the system

$$T\langle 1 \rangle \subseteq T\langle 2 \rangle \subseteq \cdots$$

where

$$T\langle j \rangle = \{ t \in T \mid A_{\geq j} t = 0 \}.$$

Each quotient  $T\langle j \rangle / T\langle j-1 \rangle$  is annihilated by  $A_{\geq 1}$  so has the form  $\coprod_{\alpha} k(\ell_{\alpha})$ , so there are short exact sequences of the form

$$0 \rightarrow T\langle j-1 \rangle \longrightarrow T\langle j \rangle \longrightarrow \coprod_{\alpha} k(\ell_{\alpha}) \rightarrow 0.$$

Tensoring such a sequence with  $F$  gives a short exact sequence of complexes because  $F$  consists of graded flat modules. The corresponding cohomology long exact sequence consists of pieces

$$h^i(T\langle j-1 \rangle \otimes_A F) \longrightarrow h^i(T\langle j \rangle \otimes_A F) \longrightarrow \coprod_{\alpha} h^i(k \otimes_A F)(\ell_{\alpha}).$$

Induction on  $j$  now makes clear that

$$h^i(k \otimes_A F) = 0$$

implies

$$h^i(T\langle j \rangle \otimes_A F) = 0 \text{ for each } j,$$

and this further gives

$$h^i(T \otimes_A F) \cong h^i(\operatorname{colim} T\langle j \rangle \otimes_A F) \cong \operatorname{colim} h^i(T\langle j \rangle \otimes_A F) = 0,$$

so the inequality (9) follows.

Let me next do the case where  $T$  is general. There is a standard spectral sequence

$$E_2^{pq} = h^p(h^q(T) \overset{L}{\otimes}_A X) \Rightarrow h^{p+q}(T \overset{L}{\otimes}_A X).$$

Since  $\operatorname{fd} X < \infty$ , the spectral sequence is first quadrant up to shift, hence converges. Now, each  $h^q(T)$  is a graded torsion module, so if  $h^q(T)$  is non-zero, then the special case of the lemma dealt with above applies to  $h^q(T) \overset{L}{\otimes}_A X$  and shows

$$\inf h^q(T) \overset{L}{\otimes}_A X \geq -\operatorname{k.f.d.} X.$$

Hence if  $h^q(T)$  is non-zero, then  $h^p(h^q(T) \overset{L}{\otimes}_A X)$  is zero for  $p < -\operatorname{k.f.d.} X$ . And of course, if  $h^q(T)$  is zero, then  $h^p(h^q(T) \overset{L}{\otimes}_A X)$  is zero for each  $p$ . So in the spectral sequence,  $E_2^{pq}$  is zero for  $p < -\operatorname{k.f.d.} X$  or  $q < \inf T$ . Therefore the limit  $h^{p+q}(T \overset{L}{\otimes}_A X)$  of the spectral sequence is zero for  $p+q < -\operatorname{k.f.d.} X + \inf T$ , and this says

$$\inf T \overset{L}{\otimes}_A X \geq -\operatorname{k.f.d.} X + \inf T,$$

proving the lemma's inequality.  $\square$

**Lemma 2.2.** *Let  $A$  satisfy  $\text{depth } A < \infty$  and  $\dim_k \text{Ext}_A^i(k, A) < \infty$  for each  $i$ .*

*Let  $X$  in  $\mathbf{D}^b(A)$  have  $\text{fd } X < \infty$ , and let  $T$  in  $\mathbf{D}^+(A^{\text{op}})$  be so that  $h^i(T)$  is a graded torsion module for each  $i$ . Then*

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T + \text{depth } X - \text{depth } A.$$

*Proof.* Using lemma 2.1 and remark 1.5 gives  $\geq$  and  $=$  in

$$\inf T \overset{\text{L}}{\otimes}_A X \geq \inf T - \text{k.fd } X = \inf T + \text{depth } X - \text{depth } A.$$

□

In the following theorem is used  $\mathbf{D}_{\text{fg}}^-(A)$ , the full subcategory of  $\mathbf{D}(A)$  consisting of complexes whose cohomology vanishes in high degrees and consists of finitely generated graded modules.

**Theorem 2.3.** *Let  $A$  satisfy  $\text{depth } A < \infty$  and  $\dim_k \text{Ext}_A^i(k, A) < \infty$  for each  $i$ .*

*Let  $X$  in  $\mathbf{D}^b(A)$  have  $\text{fd } X < \infty$ , and let  $M$  be in  $\mathbf{D}_{\text{fg}}^-(A)$ . Then*

$$\sup \text{RHom}_A(X, M) \leq \sup M - \text{depth } X + \text{depth } A.$$

*Proof.* It is easy to see that since  $M$  is in  $\mathbf{D}_{\text{fg}}^-(A)$ , the Matlis dual  $M'$  is in  $\mathbf{D}^+(A^{\text{op}})$  and has  $h^i(M')$  a graded torsion module for each  $i$ . So

$$\begin{aligned} \sup \text{RHom}_A(X, M) &= \sup \text{RHom}_A(X, M'') \\ &\stackrel{(a)}{=} \sup((M' \overset{\text{L}}{\otimes}_A X)') \\ &= -\inf M' \overset{\text{L}}{\otimes}_A X \\ &\stackrel{(b)}{\leq} -\inf M' - \text{depth } X + \text{depth } A \\ &= \sup M - \text{depth } X + \text{depth } A, \end{aligned}$$

where (a) is by adjunction and (b) is by lemma 2.2. □

The following is the special case of theorem 2.3 where  $X$  and  $M$  are concentrated in degree zero, that is, where  $X$  and  $M$  are graded modules.

**Theorem 2.4** (Ext vanishing). *Assume that  $A$  satisfies  $\text{depth } A < \infty$  and  $\dim_k \text{Ext}_A^i(k, A) < \infty$  for each  $i$ .*

*Let  $X$  in  $\text{Gr}(A)$  have  $\text{fd } X < \infty$ , and let  $M$  be in  $\text{gr}(A)$ . Then*

$$\text{Ext}_A^i(X, M) = 0 \text{ for } i > \text{depth } A - \text{depth } X.$$

This says that for  $\text{fd } X < \infty$ , the number  $\text{depth } A - \text{depth } X$  plays the role of projective dimension of  $X$ , but only with respect to finitely generated graded modules  $M$ .

Of course, this fails when  $M$  is general, as illustrated by the following example.

**Example 2.5.** Let  $A$  be the polynomial algebra  $k[x]$ . Then the conditions of theorem 2.4 are satisfied, and it is classical that  $\text{depth } A$  is 1.

Let  $X$  be  $k[x, x^{-1}]$ . Then  $\text{depth } X \geq 1$  because  $X$  is a graded torsion free module, so  $\text{depth } A - \text{depth } X \leq 0$  and theorem 2.4 gives

$$\text{Ext}_A^i(X, M) = 0 \quad \text{for } i > 0$$

for  $M$  in  $\text{gr}(A)$ .

However, this must fail when  $M$  is general, for otherwise  $X$  would be a projective object of  $\text{Gr}(A)$  which it is certainly not.

**Acknowledgement.** I thank Anders Frankild for showing me [1].

#### REFERENCES

- [1] A. Frankild, *Vanishing of local homology*, to appear in Math. Z.
- [2] P. Jørgensen, *Gorenstein homomorphisms of non-commutative rings*, J. Algebra **211** (1999), 240–267.
- [3] P. Jørgensen, *Non-commutative graded homological identities*, J. London Math. Soc. (2) **57** (1998), 336–350.

DANISH NATIONAL LIBRARY OF SCIENCE AND MEDICINE, NØRRE ALLÉ 49,  
2200 KØBENHAVN N, DK-DENMARK

*E-mail address:* pej@dnlib.dk, [www.geocities.com/popjoerg](http://www.geocities.com/popjoerg)