

q -Multiple Zeta Functions and q -Multiple Polylogarithms*

Jianqiang Zhao[†]

Department of Mathematics, University of Pennsylvania, PA 19104, USA

Abstract. We shall define the q -analogs of multiple zeta functions and multiple polylogarithms in this paper and study their properties, based on the work of Kaneko et al. and Schlesinger, respectively.

1 Introduction and definitions

Let $0 < q < 1$ and for any positive integer k define its q -analogue $[k] = [k]_q = (1 - q^k)/(1 - q)$. In [5] Kaneko et al. define a function of two complex variables $f_q(s; t) = \sum_{k=1}^{\infty} q^{kt}/[k]^s$ such that the q -analogue of Riemann zeta function is realized as

$$\zeta_q(s) := f_q(s; s - 1).$$

After analytically continuing $f_q(s; t)$ to \mathbb{C}^2 as a meromorphic function they prove the following main result

Theorem 1.1. ([5, Theorem 2]) *For any $s \in \mathbb{C} \setminus \{1 + \frac{2\pi i}{\log q} \mathbb{Z}\}$ we have*

$$\lim_{q \uparrow 1} \zeta_q(s) = \zeta(s).$$

They also study the special values of $\zeta_q(s)$ at non-negative integers. In this paper we shall generalize these to the (Euler-Zagier) multiple zeta functions, which are defined as nested generalizations of Riemann zeta function $\zeta(s)$:

$$\zeta(s_1, \dots, s_d) = \sum_{0 < k_1 < \dots < k_d} k_1^{-s_1} \dots k_d^{-s_d} \quad (1)$$

for complex variables s_1, \dots, s_d satisfying $\sigma_j + \dots + \sigma_d > d - j + 1$ for all $j = 1, \dots, d$. Here and in what follows, whenever $s \in \mathbb{C}$ we always write $\sigma = \Re(s)$, the real part of s . The analytic continuation of multiple zeta functions has been studied independently in [1] and [11]. We know that $\zeta(s_1, \dots, s_d)$ can be extended to a meromorphic function on $\mathbb{C}^d \setminus \mathfrak{S}_d$ where

$$\mathfrak{S}_d = \left\{ (s_1, \dots, s_d) \in \mathbb{C}^d \left| \begin{array}{l} s_d = 1, s_{d-1} + s_d = 2, 1, -2m \ \forall m \in \mathbb{Z}_{\geq 0}, \\ \text{and } s_j + \dots + s_d \in \mathbb{Z}_{\leq d-j+1} \ \forall j \leq d-2. \end{array} \right. \right\} \quad (2)$$

To find the q -analogue of multiple zeta functions we first define an auxiliary function of $2d$ complex variables $s_1, \dots, s_d, t_1, \dots, t_d \in \mathbb{C}$

$$f_q(s_1, \dots, s_d; t_1, \dots, t_d) = \sum_{0 < k_1 < \dots < k_d} \frac{q^{k_1 t_1 + \dots + k_d t_d}}{[k_1]^{s_1} \dots [k_d]^{s_d}}$$

which converges if $\Re(t_j + \dots + t_d) > 0$ for all $j = 1, \dots, d$ (see Proposition 2.2). In the next section we are going to analytically continue this function to \mathbb{C}^{2d} as a meromorphic function with explicitly defined poles.

*2000 Mathematics Subject Classification: Primary: 11M41, 81R50; Secondary: 11B68, 05A30, 11R42.

[†]Partially supported by NSF grant DMS0139813

We now define the q -multiple zeta function by specialization of f_q :

$$\zeta_q(s_1, \dots, s_d) := f_q(s_1, \dots, s_d; s_1 - 1, \dots, s_d - 1)$$

which will be shown to be the correct q -analog of multiple zeta functions. When $\sigma_j > 1$ for all j we can express this by the series

$$\zeta_q(s_1, \dots, s_d) = \sum_{0 < k_1 < \dots < k_d} \frac{q^{k_1(s_1-1) + \dots + k_d(s_d-1)}}{[k_1]^{s_1} \dots [k_d]^{s_d}}. \quad (3)$$

Note that when $d = 1$ this is the same as the q -analog of the Riemann zeta function defined in [5]. Put

$$\mathfrak{S}'_d = \left\{ (s_1, \dots, s_d) \in \mathbb{C}^d : \begin{array}{l} s_d \in 1 + \frac{2\pi i}{\log q} \mathbb{Z}, \text{ or } s_d \in \mathbb{Z}_{\leq 0} + \frac{2\pi i}{\log q} \mathbb{Z}_{\neq 0}, \\ \text{or } s_j + \dots + s_d \in \mathbb{Z}_{\leq d-j+1} + \frac{2\pi i}{\log q} \mathbb{Z}, \ j < d \end{array} \right\} \supset \mathfrak{S}_d.$$

Here the last part in \mathfrak{S}'_d is vacuous if $d = 1$. The primary goal of this paper is to prove

Main Theorem. *The q -multiple zeta function $\zeta_q(s_1, \dots, s_d)$ can be extended to a meromorphic function on $\mathbb{C}^d \setminus \mathfrak{S}'_d$ with simple poles along \mathfrak{S}'_d . Further, for all $(s_1, \dots, s_d) \in \mathbb{C}^d \setminus \mathfrak{S}'_d$*

$$\lim_{q \uparrow 1} \zeta_q(s_1, \dots, s_d) = \zeta(s_1, \dots, s_d).$$

In the last section of this paper, we propose a new definition of the q -multiple polylogarithms and briefly study their properties.

2 Analytic continuations of f_q and ζ_q

The purpose of this section is two-fold: first we will use the auxiliary functions f_q introduced in the first section to give a quick analytic continuation of q -multiple zeta functions $\zeta_q(s_1, \dots, s_d)$, though this is not enough to show it's the right q -analog of the multiple zeta functions. Second, we write down these expressions involving binomial coefficients explicitly which will be used to study special values of $\zeta_q(s_1, \dots, s_d)$ in section 6.

We need a simple lemma first.

Lemma 2.1. *Let k be a positive integer. For all $1 > q > 1/2$ we have*

$$1 \leq \frac{1 - q^k}{1 - q} < 2k.$$

Proof. The inequality $1 \leq \frac{1 - q^k}{1 - q}$ is obvious. Let $f(q) = 2k(1 - q) - (1 - q^k)$. We show that $f(q) \geq 0$ for all $0 < q < 1$. Indeed, for all such q

$$f'(q) = -2k + kq^{k-1} \leq -k < 0.$$

So f is strictly decreasing and the positivity of f follows from $f(1) = 0$. This implies that $2k > (1 - q^k)/(1 - q)$ as desired. \square

Proposition 2.2. *The function*

$$f_q(s_1, \dots, s_d; t_1, \dots, t_d) = \sum_{0 < k_1 < \dots < k_d} \frac{q^{k_1 t_1 + \dots + k_d t_d}}{[k_1]^{s_1} \dots [k_d]^{s_d}}$$

converges if $\Re(t_j + \dots + t_d) > 0$ for all $j = 1, \dots, d$. It can be analytically continued to a meromorphic function over \mathbb{C}^{2d} via the series expansion

$$f_q(s_1, \dots, s_d; t_1, \dots, t_d) = (1 - q)^{\text{wt}(\vec{s})} \sum_{r_1, \dots, r_d=0}^{+\infty} \prod_{j=1}^d \left[\binom{s_j + r_j - 1}{r_j} \frac{q^{j(r_j + t_j)}}{1 - q^{r_j + t_j + \dots + r_d + t_d}} \right], \quad (4)$$

where $\text{wt}(\vec{s}) = s_1 + \dots + s_d$. It has the following (simple) poles: $t_j + \dots + t_d \in \mathbb{Z}_{\leq 0} + \frac{2\pi i}{\log q} \mathbb{Z}$.

Proof. Assume $|\Re(s_j)| < N_j$ and let $\tau_j = \Re(t_j)$ for all $j = 1, \dots, d$. By Lemma 2.2

$$\left| f_q(s_1, \dots, s_d; t_1, \dots, t_d) \right| < \sum_{0 < k_1 < \dots < k_d} \prod_{j=1}^d (2k_j)^{N_j} q^{k_j \tau_j}. \quad (5)$$

Let $k = k_{d-1}$ (when $d = 1$ take $k_0 = 0$), $n = k_d$, $N = N_d$, and $\tau = \tau_d$. Then by root test $\sum_{n > k} n^N q^{n\tau}$ converges and moreover

$$\sum_{n > k} n^N q^{n\tau} = \left(q \frac{d}{dq} \right)^N \sum_{n > k} q^{n\tau} = (1 - q^\tau)^{-N} f_N(k; q, \tau) q^{k\tau} \quad (6)$$

where $f_N(x; q, \tau) = \sum_{l=0}^N c_l x^l$ is a polynomial of degree N whose coefficients depend only on the constants N , q and τ . This proves the first part of the lemma when $d = 1$. In the general case it follows from (5) and (6) that

$$\left| f_q(s_1, \dots, s_d; t_1, \dots, t_d) \right| < \frac{2^{N_1 + \dots + N_d}}{(1 - q^\tau)^N} \sum_{l=0}^N c_l \sum_{0 < k_1 < \dots < k_{d-1} < k} \left(\prod_{j=1}^{d-2} (k_j)^{N_j} q^{k_j \tau_j} \right) k^{N_{d-1} + l} q^{k(\tau_{d-1} + \tau_d)}.$$

Hence the first part of the lemma follows from an easy induction on d .

By binomial expansion $(1 - x)^{-s} = \sum_{r=0}^{\infty} \binom{s+r-1}{r} x^r$ we get

$$f_q(s_1, \dots, s_d; t_1, \dots, t_d) = (1 - q)^{\text{wt}(\vec{s})} \sum_{0 < k_1 < \dots < k_d} \sum_{r_1, \dots, r_d=0}^{+\infty} \left(\prod_{j=1}^d \binom{s_j + r_j - 1}{r_j} q^{k_j(r_j + t_j)} \right).$$

As $0 < q < 1$ the series converges absolutely by Stirling's formula so we can exchange the summations. The proposition follows immediately from the next lemma by taking $x_j = q^{t_j + r_j}$ for $j = 1, \dots, d$. \square

Lemma 2.3. *Let $x_j \in \mathbb{C}$ such that $|x_j| < 1$ for $j = 1, \dots, d$. Then*

$$\sum_{0 < k_1 < \dots < k_d} \prod_{j=1}^d x_j^{k_j} = \prod_{j=1}^d \frac{x_j^j}{1 - x_j} = \prod_{j=1}^d \frac{x_j \cdots x_d}{1 - x_j \cdots x_d}. \quad (7)$$

Proof. When $n = 1$ this is clear. For $n \geq 2$ we denote the right hand side of (7) by $F_d(x_1, \dots, x_d)$. Then

$$\sum_{0 < k_1 < \dots < k_d} \prod_{j=1}^d x_j^{k_j} = \sum_{0 < k_1 < \dots < k_{d-1}} \prod_{j=1}^{d-1} x_j^{k_j} \sum_{k_d > k_{d-1}} x_d^{k_d} = \frac{x_d}{1 - x_d} F_{d-1}(x_1, \dots, x_{d-2}, x_{d-1} x_d).$$

The lemma follows immediately by induction. \square

Recall that $\zeta_q(s_1, \dots, s_d) = f_q(s_1, \dots, s_d; s_1 - 1, \dots, s_d - 1)$. Hence we have the following immediate consequence.

Theorem 2.4. *The q -multiple zeta function $\zeta_q(s_1, \dots, s_d)$ can be extended to a meromorphic function with simple poles lying along \mathfrak{S}'_d :*

$$\zeta_q(s_1, \dots, s_d) = (1 - q)^{\text{wt}(\vec{s})} \sum_{r_1, \dots, r_d=0}^{+\infty} \prod_{j=1}^d \left[\binom{s_j + r_j - 1}{r_j} \frac{q^{j(r_j + s_j - 1)}}{1 - q^{r_j + s_j + \dots + r_d + s_d - d + j - 1}} \right].$$

To see the effect of taking different specializations of t_j in f_q we define the shifting operators \mathcal{S}_j ($1 \leq j \leq d$) on the multiple zeta functions as follows:

$$\mathcal{S}_j \zeta(s_1, \dots, s_d) = \zeta(s_1, \dots, s_d) + (1 - q) \zeta(s_1, \dots, s_j - 1, \dots, s_d). \quad (8)$$

It is obvious that these operators are commutative.

Proposition 2.5. *Let n_1, \dots, n_d be non-negative integers. Then we have*

$$\begin{aligned} & f_q(s_1, \dots, s_d; s_1 - 1 - n_1, \dots, s_d - 1 - n_d) \\ &= \mathcal{S}_1^{n_1} \circ \dots \circ \mathcal{S}_d^{n_d} \zeta(s_1, \dots, s_d) = \sum_{r_1=0}^{n_1} \dots \sum_{r_d=0}^{n_d} \left(\prod_{j=1}^d \binom{n_j}{r_j} (1-q)^{r_j} \right) \zeta_q(s_1 - r_1, \dots, s_d - r_d). \end{aligned}$$

Proof. We only sketch the proof in the case $n_1 = \dots = n_{d-1} = 0$. The general case is completely similar. In the rest of the paper we always let \mathcal{S} be the shifting operator on the last variable. Suppose $n_d = n = 1$. Then

$$\begin{aligned} & f_q(s_1, \dots, s_d; s_1 - 1, \dots, s_{d-1} - 1, s_d - 2) \\ &= \sum_{0 < k_1 < \dots < k_d} \frac{q^{k_1(s_1-1) + \dots + k_{d-1}(s_{d-1}-1) + k_d(s_d-2)}}{[k_1]^{s_1} \dots [k_d]^{s_d}} \\ &= \sum_{0 < k_1 < \dots < k_d} \frac{q^{k_1(s_1-1) + \dots + k_{d-1}(s_{d-1}-1)}}{[k_1]^{s_1} \dots [k_{d-1}]^{s_{d-1}}} \cdot \frac{q^{k_d(s_d-2)}(1 - q^{k_d}) + q^{k_d(s_d-1)}}{[k_d]^{s_d}} \\ &= \mathcal{S} \zeta(s_1, \dots, s_d). \end{aligned}$$

The rest follows easily by induction. □

The next corollary answers an implicit question in [5].

Corollary 2.6. *Let n be a positive integer. The specialization of t in $f_q(s; t)$ to $s - 1 - n$ is*

$$f_q(s; s - 1 - n) = \mathcal{S}^n \zeta_q(s) = \sum_{r=0}^n \binom{n}{r} (1-q)^r \zeta_q(s - r).$$

We observe that one effect of the shifting operator is to bring in more poles. Essentially, \mathcal{S}^n shifts all the poles of $\zeta_q(s)$ by n to the right on the complex plane.

3 Analytic continuation of multiple zeta functions

Let's begin with a review of some classical results on Bernoulli polynomials $B_k(x)$ which are defined by the generating function

$$\frac{te^{xt}}{e^t - 1} = \sum_{k=0}^{\infty} B_k(x) \frac{t^k}{k!}.$$

Let $\tilde{B}_k(x)$ be the “periodic Bernoulli polynomial”

$$\tilde{B}_k(x) = B_k(\{x\}), \quad x \geq 1,$$

where $\{x\}$ is the fractional part of x . Then we have ([10, Ch. IX, Misc. Ex. 12])

$$\tilde{B}_k(x) = -k! \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{e^{2\pi i n x}}{(2\pi i n)^k}. \quad (9)$$

Recall that the Bernoulli numbers satisfy $B_k = B_k(1)$ if $k \geq 2$ while $B_0 = 1$ and $B_1 = -1/2$.

Lemma 3.1. *For every positive integer $M \geq 2$ and $x > 1$ we have*

$$\left| \tilde{B}_M(x) \right| \leq \frac{4M!}{(2\pi)^M}.$$

Proof. It follows from the fact that $\zeta(M) \leq \zeta(2) = \pi^2/6 < 2$ for $M \geq 2$. □

We know that one can analytically continue the multiple zeta functions as independently presented in [11] and [1] by different methods. Moreover, $\zeta(s_1, \dots, s_d)$ has singularities on the hyperplanes in \mathfrak{S}_d defined by (2). However, neither approach is suitable for our purpose here. So we follow the idea in [5] to provide a third approach in the rest of this section. The same idea will also be used to deal with the q -multiple zeta functions.

Let's recall the classical Euler-Maclaurin summation formula [10, 7.21]. Let $f(x)$ be any (complex-valued) C^∞ function on $[1, \infty)$ and let m and M be two positive integers. Then we have

$$\begin{aligned} \sum_{n=1}^m f(n) &= \int_1^m f(x) dx + \frac{1}{2}(f(1) + f(m)) + \sum_{r=1}^M \frac{B_{r+1}}{(r+1)!} (f^{(r)}(m) - f^{(r)}(1)) \\ &\quad - \frac{(-1)^{M+1}}{(M+1)!} \int_1^m \tilde{B}_{M+1}(x) f^{(M+1)}(x) dx. \end{aligned} \quad (10)$$

To simplify our notation, in definition (1) we replace s_d, k_{d-1}, k_d by s, k and n , respectively. Taking $f(x) = 1/x^s$ and $m = k$ and ∞ in (10) we have:

$$\begin{aligned} \sum_{n>k} \frac{1}{n^s} &= \sum_{n=1}^{\infty} \frac{1}{n^s} - \sum_{n=1}^k \frac{1}{n^s} \\ &= \int_k^{\infty} f(x) dx - \frac{1}{2}f(k) - \sum_{r=1}^M \frac{B_{r+1}}{(r+1)!} f^{(r)}(k) - \frac{(-1)^{M+1}}{(M+1)!} \int_k^{\infty} \tilde{B}_{M+1}(x) f^{(M+1)}(x) dx \\ &= \frac{1}{(s-1)k^{s-1}} - \frac{1}{2k^s} + \sum_{r=1}^M \frac{B_{r+1}}{(r+1)!} \frac{(s)_r}{k^{s+r}} - \frac{(s)_{M+1}}{(M+1)!} \int_k^{\infty} \frac{\tilde{B}_{M+1}(x)}{x^{s+M+1}} dx. \end{aligned}$$

Here we have used the fact that $B_k = 0$ if $k \geq 3$ is odd. By definition (1) we have

Theorem 3.2. *For all $(s_1, \dots, s_d) \in \mathbb{C}^d \setminus \mathfrak{S}_d$ and $M > 1 + |\sigma_d| + |\sigma_{d-1}|$ we have*

$$\begin{aligned} \zeta(s_1, \dots, s_d) &= \sum_{r=0}^{M+1} \frac{B_r}{r!} (s_d)_{r-1} \cdot \zeta(s_1, \dots, s_{d-1} + s_d + r - 1) \\ &\quad - \frac{(s_d)_{M+1}}{(M+1)!} \sum_{0 < k_1 < \dots < k_d} \frac{1}{k_1^{s_1} \dots k_{d-1}^{s_{d-1}}} \int_{k_{d-1}}^{\infty} \frac{\tilde{B}_{M+1}(x)}{x^{s_d+M+1}} dx. \end{aligned} \quad (11)$$

where we set $(s)_0 = 1$ and $(s)_{-1} = 1/(s-1)$. This provides an analytic continuation of $\zeta(s_1, \dots, s_d)$ to $\mathbb{C}^d \setminus \mathfrak{S}_d$.

Proof. We only need to show that the series in (11) converges. Lemma 3.1 implies (if $d = 2$ then take $k_0 = 1$)

$$\sum_{k_{d-1}=k_{d-2}}^{\infty} \left| \frac{1}{k_{d-1}^{s_{d-1}}} \int_{k_{d-1}}^{\infty} \frac{\tilde{B}_M(x)}{x^{M+s_d+1}} dx \right| \leq \frac{4M!}{(2\pi)^M (M - |\sigma_d|)} \sum_{k_{d-1}=k_{d-2}}^{\infty} \frac{1}{k_{d-1}^{M-|\sigma_{d-1}|-|\sigma_d|}}$$

which converges absolutely whenever $M > 1 + |\sigma_d| + |\sigma_{d-1}|$. \square

4 Proof of Main Theorem

Fix $(s_1, \dots, s_d) \in \mathbb{C}^d$ such that $\sigma_j + \dots + \sigma_d < d - j + 1$ for all $j = 1, \dots, d$. When $d = 1$ this is Theorem 1.1 due to Kaneko et al. We now assume $d \geq 2$ and proceed by induction. The key is a recursive formula for $\zeta_q(s_1, \dots, s_d)$ similar to (11) for $\zeta(s_1, \dots, s_d)$. To derive this formula we appeal to the Euler-Maclaurin summation formula (10) again. Hence we set

$$F(x) = \frac{q^{x(s-1)}}{(1-q^x)^s}$$

as in [5]. Then

$$F'(x) = (\log q) q^{x(s-1)} \frac{s-1+q^x}{(1-q^x)^{s+1}},$$

$$F''(x) = (\log q)^2 q^{x(s-1)} \frac{s(s+1) - 3s(1-q^x) + (1-q^x)^2}{(1-q^x)^{s+2}}.$$

In definition (3) we replace s_d , k_{d-1} , and k_d by s , k and n , respectively. We now take $M = 1$, $f(x) = F(x+k-1)$ and let $m \rightarrow \infty$ in (10) and get

$$\begin{aligned} \sum_{n>k} \frac{q^{n(s-1)}}{[n]^s} &= (1-q)^s \left(-F(k) + \sum_{n=1}^{\infty} f(n) \right) \\ &= (1-q)^s \left(\int_k^{\infty} F(x) dx - \frac{1}{2} F(k) - \frac{1}{12} F'(k) - \frac{1}{2} \int_1^{\infty} \tilde{B}_2(x) f''(x) dx \right) \\ &= \frac{(q-1)}{(s-1) \log q} \frac{q^{k(s-1)}}{[k]^{s-1}} - \frac{1}{2} \frac{q^{k(s-1)}}{[k]^s} + \frac{1}{12} \frac{\log q}{q-1} \frac{q^{k(s-1)}(s+q^k-1)}{[k]^{s+1}} \\ &\quad - \frac{(1-q)^s (\log q)^2}{2} \int_k^{\infty} \tilde{B}_2(x) q^{x(s-1)} \frac{s(s+1) - 3s(1-q^x) + (1-q^x)^2}{(1-q^x)^{s+2}} dx \end{aligned} \quad (12)$$

because $\tilde{B}_2(x+k-1) = \tilde{B}_2(x)$ by periodicity. By the same argument as in [5], setting the incomplete beta integrals

$$b_t(\alpha, \beta) = \int_0^t u^{\alpha-1} (1-u)^{\beta-1} du,$$

we can obtain from (9) the following expression for the last term in (12):

$$(12) = - \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{(1-q)^s \log q}{(2\pi i n)^2} \sum_{\nu=\pm 1, 0} a_{\nu}(s) b_{q^k}(s-1+\delta n, -s+\nu) \quad (13)$$

where $\delta = 2\pi i / \log q$, $a_{-1}(s) = s(s+1)$, $a_0(s) = -3s$, and $a_1(s) = 1$. Repeatedly applying integration by parts on these incomplete beta integrals we get for $\nu = \pm 1, 0$ and positive integer $M \geq 2$

$$\begin{aligned} b_{q^k}(s-1+\delta n, -s+\nu) &= \sum_{r=1}^{M-1} (-1)^{r-1} \frac{(s+1-\nu)_{r-1}}{(s-1+\delta n)_r} \frac{q^{k(s+r-2)}}{(1-q^k)^{s+r-\nu}} \\ &\quad + (-1)^{M-1} \frac{(s+1-\nu)_{M-1}}{(s-1+\delta n)_{M-1}} b_{q^k}(s-2+M+\delta n, -s-M+1+\nu). \end{aligned} \quad (14)$$

Set $\vec{s}' = (s_1, \dots, s_{d-2})$ if $d \geq 3$ and $\vec{s}' = \emptyset$ if $d = 2$. Putting (3), (12), (13) and (14) together and applying Proposition 2.5 we get

$$\begin{aligned} \zeta_q(\vec{s}', s_{d-1}, s_d) &= \frac{(q-1)}{(s-1) \log q} \zeta_q(\vec{s}', s_{d-1} + s_d - 1) - \frac{1}{2} \mathcal{S} \zeta_q(\vec{s}', s_{d-1} + s_d) \\ &\quad + \frac{s}{12} \frac{\log q}{q-1} \mathcal{S}^2 \zeta_q(\vec{s}', s_{d-1} + s_d + 1) + \frac{\log q}{12} \mathcal{S} \zeta_q(\vec{s}', s_{d-1} + s_d) - \sum_{\nu=\pm 1, 0} (C_{\nu} + D_{\nu}) \end{aligned} \quad (15)$$

where C_{ν} and D_{ν} are the contributions from the sum involving $b_{q^k}(\dots, -s+\nu)$. Explicitly they are computed as follows. Write

$$T(q, s, n, r) = \prod_{j=0}^{r-1} (2\pi i n + (s-1+j) \log q)^{-1}. \quad (16)$$

Then

$$\begin{aligned}
C_{-1} &= \sum_{r=1}^{M-1} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, r)}{(2\pi i n)^2} \left(\frac{\log q}{q-1} \right)^{r+1} (s)_{r+1} \cdot \mathcal{S}^3 \zeta_q(\vec{s}', s_{d-1} + s_d + r + 1), \\
D_{-1} &= - \sum_{0 < k_1 < \dots < k_{d-2} < k_{d-1}} \frac{q^{k_1(s_1-1) + \dots + k_{d-1}(s_{d-1}-1)}}{[k_1]^{s_1} \dots [k_{d-1}]^{s_{d-1}}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, M-1)}{(2\pi i n)^2} \\
&\quad \cdot \left(\frac{\log q}{q-1} \right)^{M+1} (s)_{M+1} \cdot \int_{k_{d-1}}^{\infty} e^{2\pi i n x} q^{x(s-2+M)} \left(\frac{1-q^x}{1-q} \right)^{-s-M-1} dx \\
&= \sum_{0 < k_1 < \dots < k_{d-2}} \frac{q^{k_1(s_1-1) + \dots + k_{d-2}(s_{d-2}-1)}}{[k_1]^{s_1} \dots [k_{d-2}]^{s_{d-2}}} \sum_{k=k_{d-2}+1}^{\infty} R(M, q, k, s_{d-1}, s).
\end{aligned}$$

where we replace the index k_{d-1} by k . Similarly,

$$\begin{aligned}
C_0 &= 3 \log q \sum_{r=1}^{M-1} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, r)}{(2\pi i n)^2} \left(\frac{\log q}{q-1} \right)^r (s)_r \cdot \mathcal{S}^2 \zeta_q(\vec{s}', s_{d-1} + s_d + r), \\
D_0 &= -3 \log q \sum_{0 < k_1 < \dots < k_{d-2} < k_{d-1}} \frac{q^{k_1(s_1-1) + \dots + k_{d-1}(s_{d-1}-1)}}{[k_1]^{s_1} \dots [k_{d-1}]^{s_{d-1}}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, M-1)}{(2\pi i n)^2} \\
&\quad \cdot \left(\frac{\log q}{q-1} \right)^M (s)_M \cdot \int_{k_{d-1}}^{\infty} e^{2\pi i n x} q^{x(s-2+M)} \left(\frac{1-q^x}{1-q} \right)^{-s-M} dx,
\end{aligned}$$

and

$$\begin{aligned}
C_1 &= (\log q)^2 \sum_{r=1}^{M-1} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, r)}{(2\pi i n)^2} \left(\frac{\log q}{q-1} \right)^{r-1} (s)_{r-1} \cdot \mathcal{S} \zeta_q(\vec{s}', s_{d-1} + s_d + r - 1), \\
D_1 &= -(\log q)^2 \sum_{0 < k_1 < \dots < k_{d-2} < k_{d-1}} \frac{q^{k_1(s_1-1) + \dots + k_{d-1}(s_{d-1}-1)}}{[k_1]^{s_1} \dots [k_{d-1}]^{s_{d-1}}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{T(q, s, n, M-1)}{(2\pi i n)^2} \\
&\quad \cdot \left(\frac{\log q}{q-1} \right)^{M-1} (s)_{M-1} \cdot \int_{k_{d-1}}^{\infty} e^{2\pi i n x} q^{x(s-2+M)} \left(\frac{1-q^x}{1-q} \right)^{-s-M+1} dx.
\end{aligned}$$

The crucial step next is to control the summations over k_{d-1} and show that they converge uniformly with respect to q . When $0 < q \leq 1/2$ this is clear. The only non-trivial part is when $q \uparrow 1$. So we assume $1/2 < q < 1$. Note that

$$\lim_{q \uparrow 1} T(q, s, n, r) = \frac{1}{(2\pi i n)^r}, \quad \lim_{q \uparrow 1} \frac{\log q}{q-1} = 1.$$

Lemma 4.1. *Let $s_d = s = \sigma + i\tau$. Let $q_0 = \max\{1/2, e^{(6-2\pi)/\tau}\}$ if $\tau > 0$ and let $q_0 = 1/2$ if $\tau \leq 0$. Then for all $1 > q > q_0$ and positive integer k we have*

$$\left| \frac{\log q}{q-1} \right| < 2, \quad \text{and} \quad |T(q, s, n, r)| < \frac{1}{(6n)^r}.$$

Proof. Let $f(q) = 2(1-q) + \log q$. Then $f'(q) = -2 + 1/q < 0$ whenever $q > q_0$. So $f(q) > f(1) = 0$ whenever $1 > q > q_0$. This implies that $2(1-q) > -\log q$ whence $\log q/(q-1) < 2$.

To bound $T(q, s, n, r)$ we consider each of its factors in definition (16). For each $0 \leq j < r$ we have

$$|2\pi i n + (s-1+j) \log q|^2 = ((\sigma-1+j) \log q)^2 + (2\pi n + \tau \log q)^2 \geq (2\pi n + \tau \log q)^2$$

which is independent j . If $\tau \leq 0$ then clearly $|2\pi i n + (s-1+j) \log q| > 6n$. If $\tau > 0$ then it follows from $q > e^{(6-2\pi)\tau}$ that

$$2\pi n + \tau \log q > 2\pi n + 6 - 2\pi \geq 6n,$$

as desired. \square

Next we want to bound the integral terms in D_{-1} . Let $|\sigma_d| < N$ and $|\sigma_{d-1}| < N'$ for some positive integers N and N' . Fix an arbitrary $x > k$ and a positive integer $M > 16 + 2N + 6 \sum_{j=1}^{d-1} (N_j + 1)$. Then

$$q^{-k(M/6-N'-1)} \left| q^{k(s'-1)} q^{x(s-2+M)} \left(\frac{1-q^x}{1-q} \right)^{-s-M-1} \right| < q^{x(M-N-2)-kM/6} \left(\frac{1-q}{1-q^x} \right)^{M-N+1}.$$

Denote by $g(q)$ the right hand side of the above inequality.

Lemma 4.2. *Let $1/2 < q < 1$. Then $g(q)$ is increasing as a function of q so that*

$$g(q) \leq \lim_{q \uparrow 1} g(q) = \frac{1}{x^{M-N+1}}.$$

Proof. Taking the logarithmic derivative of $g(q)$ we have

$$\begin{aligned} \frac{g'(q)}{g(q)} &= \frac{x(M-N-2)-kM/6}{q} + (M-N+1) \frac{xq^{x-1}(1-q) - (1-q^x)}{(1-q)(1-q^x)} \\ &= \frac{(1-q)(1-q^x)(x(M-N-2)-kM/6) + (M-N+1)(xq^x(1-q) - q + q^{x+1})}{q(1-q)(1-q^x)} \end{aligned}$$

whose numerator is denoted by $h(q)$. Then

$$\begin{aligned} h'(q) &= ((x+1)q^x - xq^{x-1} - 1)(x(M-N-2) - kM/6) \\ &\quad + (M-N+1)(x(xq^{x-1} - (x+1)q^x) - 1 + (x+1)q^x). \end{aligned}$$

Clearly $h'(1) = 0$ and moreover

$$\begin{aligned} q^{2-x} h''(q) &= (x(x+1)q - x(x-1))(x(M-N-2) - kM/6) \\ &\quad + (M-N+1)(x(x(x-1) - x(x+1)q) + x(x+1)q) \\ &= x(x-1)(kM/6 + 3x) + qx(x+1)(M-N+1 - (kM/6 + 3x)) \\ &= (kM/6 + 3x)(x^2(1-q) - x(1+q)) + qx(x+1)(M-N+1) \\ &> qx(x+1)(M-N+1) - x(1+q)(kM/6 + 3x) \quad (\text{since } 1 > q) \\ &\geq qx^2 \left\{ M-N+1 - \frac{1+q}{q}(3 + M/6) \right\} \quad (\text{since } k < x) \\ &> qx^2(M/2 - N - 8) > 0 \end{aligned}$$

where we used the fact that if $q > 1/2$ then $(1+q)/q < 3$. This implies that $h'(q)$ is increasing so that $h'(q) < 0$ for all $1 > q > 1/2$ (recall that $h'(1) = 0$). It follows that $h(q)$ is decreasing. But $h(1) = 0$ so we know $h(q) > 0$ for all such q . Thus $g'(q) > 0$ and therefore $g(q)$ is increasing. This completes the proof of the lemma. \square

We now can bound the innermost sum of D_{-1} . From Lemma 2.1, Lemma 4.1 and Lemma 4.2 we have (if $d = 2$ then take $k_0 = 1$)

$$\begin{aligned} \sum_{k=1+k_{d-2}}^{\infty} |R(M, q, k, s', s)| &< \sum_{k=1+k_{d-2}}^{\infty} (2k)^{N'} \frac{2\zeta(M+1)}{4\pi^2 6^{M-1}} 2^{M+1} (N)_{M+1} q^{k(M/6-N'-1)} \int_k^{\infty} \frac{dx}{x^{M-N+1}} \\ &< \frac{(M+1)!}{M-N} \binom{M+N}{M+1} \sum_{k=1+k_{d-2}}^{\infty} \frac{q^{k(M/6-N'-1)}}{k^{M-N-N'}} \\ &< (M+1)!(2M)^{M+1} \sum_{k=1+k_{d-2}}^{\infty} q^{k(M/6-N'-1)} \end{aligned}$$

since $2^{N'+M+2}\zeta(M+1) < 4\pi^2 6^{M-1}$ and $M - N > 2$. Therefore by Lemma 2.1

$$|D_{-1}| < (2M)^{2M} \sum_{0 < k_1 < \dots < k_{d-2}} \left(\prod_{l=1}^{d-2} k_l^{N_l} q^{k_l(-N_l-1)} \right) q^{k_{d-2}(M/6 - N_{d-1} - 1)}$$

which converges as proved in Proposition 2.2.

Exactly the same argument applies to the integral terms in D_0 and D_1 which we leave to the interested readers. These convergence results imply two things. First we can show by induction on d that (15) gives rise to an analytic continuation of $\zeta_q(s_1, \dots, s_d)$ as a meromorphic function on $\mathbb{C}^d \setminus \mathfrak{S}'_d$. Second, also by induction on d , we now can conclude that it's legitimate to take the limit $q \uparrow 1$ inside the sums of C_ν and D_ν to get (note that $\lim_{q \uparrow 1} \mathcal{S}^n \zeta_q(\vec{s}) = \zeta(\vec{s})$ for any $\vec{s} \in \mathbb{C}^{d-1} \setminus \mathfrak{S}'_{d-1}$ and any positive integer n)

$$\begin{aligned} \lim_{q \uparrow 1} \zeta_q(\vec{s}', s_{d-1}, s_d) &= \frac{1}{s-1} \zeta(\vec{s}', s_{d-1} + s_d - 1) - \frac{1}{2} \zeta(\vec{s}', s_{d-1} + s_d) + \frac{s}{12} \zeta(\vec{s}', s_{d-1} + s_d + 1) \\ &\quad - \sum_{r=1}^{M-1} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{(2\pi i n)^{r+2}} (s)_{r+1} \cdot \zeta(\vec{s}', s_{d-1} + s_d + r + 1) \\ &\quad + \sum_{0 < k_1 < \dots < k_{d-1}} \frac{1}{k_1^{s_1} \dots k_{d-1}^{s_{d-1}}} \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{(s)_{M+1}}{(2\pi i n)^{M+1}} \int_{k_{d-1}}^{\infty} e^{2\pi i n x} x^{-s-M-1} dx \\ &= \sum_{r=0}^{M+1} \frac{B_r}{r!} (s_d)_{r-1} \cdot \zeta(s_1, \dots, s_{d-1} + s_d + r - 1) \\ &\quad - \frac{1}{(M+1)!} \sum_{0 < k_1 < \dots < k_d} \frac{1}{k_1^{s_1} \dots k_d^{s_d}} \int_{k_{d-1}}^{\infty} \tilde{B}_{M+1}(x) \frac{(s)_{M+1}}{x^{s+M+1}} dx \end{aligned}$$

by (9) and its specialization with $x = 1$

$$\sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{(2\pi i n)^{r+2}} = -\frac{\tilde{B}_{r+2}(1)}{(r+2)!} = -\frac{B_{r+2}}{(r+2)!}.$$

The main theorem now mostly follows from Theorem 3.2. The poles at $s_d = m - \frac{2\pi i}{\log q} n$ are given by the first term in formula (15) when $m = 1$ and $n = 0$ and by the terms $T(q, s_d, n, r)$ as defined in (16) if $m \leq 1$ and $n \neq 0$. The location of the other poles are obtained by induction using those poles of the q -Riemann zeta function presented in Theorem 1.1 for the initial step. This completes the proof of our main theorem.

5 q -shuffle relations

The classical multiple zeta functions satisfy shuffle relations originating from their series representations. For example,

$$\zeta(s_1)\zeta(s_2) = \zeta(s_1, s_2) + \zeta(s_2, s_1) + \zeta(s_1 + s_2). \quad (17)$$

In general we can define a shuffle operation $*$ on finite words so that for any complex numbers a, b (regarded as letters) and words w_1 and w_2 of complex numbers

$$aw_1 * bw_2 = a(w_1 * bw_2) + b(aw_1 * w_2) + (a+b)(w_1 * w_2)$$

so that treating complex variables as words we have

$$\zeta(w_1)\zeta(w_2) = \zeta(w_1 * w_2).$$

To generalize these relations we first define shifting operators on words of complex variables:

$$(s_1, \dots, s_{i-1}, \mathcal{S}(s_i), s_{i+1}, \dots, s_d) = \mathcal{S}_i(s_1, \dots, s_d)$$

and set $\zeta(\mathcal{S}_i(s_1, \dots, s_d)) = \mathcal{S}_i \zeta(s_1, \dots, s_d)$. Now we define the q -shuffle operator $*_q$ on words by

$$aw_1 *_q bw_2 = a(w_1 *_q bw_2) + b(aw_1 *_q w_2) + \mathcal{S}(a+b)(w_1 *_q w_2).$$

Theorem 5.1. *For any two words of complex variables w_1 and w_2 we have*

$$\zeta_q(w_1)\zeta_q(w_2) = \zeta_q(w_1 *_q w_2).$$

Proof. Induction on the length of $w_1 w_2$. □

For example

$$\zeta_q(s_1)\zeta_q(s_2) = \zeta_q(s_1, s_2) + \zeta_q(s_2, s_1) + \zeta_q(s_1 + s_2) + (q-1)\zeta_q(s_1 + s_2 - 1). \quad (18)$$

We can recover the shuffle relation of (17) by taking $\lim_{q \uparrow 1}$.

6 Special values of $\zeta_q(s_1, s_2)$

For integers n_1, \dots, n_d we set

$$\begin{aligned} \zeta_q(n_1, \dots, n_d) &= \lim_{s_1 \rightarrow n_1} \cdots \lim_{s_d \rightarrow n_d} \zeta_q(s_1, \dots, s_d), \\ \zeta_q^R(n_1, \dots, n_d) &= \lim_{s_d \rightarrow n_d} \cdots \lim_{s_1 \rightarrow n_1} \zeta_q(s_1, \dots, s_d) \end{aligned}$$

if the limits exist. Interesting phenomena occur already in the case $d = 2$ and these should be generalized to arbitrary depth. By Theorem 2.4 we get

$$\begin{aligned} \zeta_q(s_1, s_2) &= (1-q)^{s_1+s_2} \sum_{r_1, r_2=0}^{+\infty} \binom{s_1+r_1-1}{r_1} \binom{s_2+r_2-1}{r_2} \frac{q^{2s_2+2r_2+s_1+r_1-3}}{(1-q^{s_2+r_2-1})(1-q^{s_2+r_2+s_1+r_1-2})} \\ &= (1-q)^{s_1+s_2} \left\{ \frac{q^{2s_2+s_1-3}}{(1-q^{s_2-1})(1-q^{s_2+s_1-2})} + \frac{s_1 q^{2s_2+s_1-2}}{(1-q^{s_2-1})(1-q^{s_2+s_1-1})} \right. \\ &\quad + \frac{s_2 q^{2s_2+s_1-1}}{(1-q^{s_2})(1-q^{s_2+s_1-1})} + \frac{s_1 s_2 q^{2s_2+s_1}}{(1-q^{s_2})(1-q^{s_2+s_1})} \\ &\quad \left. + \frac{s_1(s_1+1)q^{2s_2+s_1-1}}{2(1-q^{s_2-1})(1-q^{s_2+s_1})} + \frac{s_2(s_2+1)q^{2s_2+s_1+1}}{2(1-q^{s_2+1})(1-q^{s_2+s_1})} + \cdots \right\} \end{aligned}$$

Clearly we have

$$\begin{aligned} \zeta_q(0, 0) &= \lim_{s_1 \rightarrow 0} \lim_{s_2 \rightarrow 0} \zeta_q(s_1, s_2) = \frac{1}{(q^2-1)(q-1)} - \frac{3}{2(q-1)\log q} + \frac{1}{\log^2 q}, \\ \zeta_q^R(0, 0) &= \lim_{s_2 \rightarrow 0} \lim_{s_1 \rightarrow 0} \zeta_q(s_1, s_2) = \frac{1}{(q^2-1)(q-1)} - \frac{1}{(q-1)\log q} + \frac{q}{2(q-1)\log q}. \end{aligned}$$

It is not too hard to find that

$$\lim_{q \uparrow 1} \zeta_q(0, 0) = \frac{1}{3}, \quad \lim_{q \uparrow 1} \zeta_q^R(0, 0) = \frac{5}{12}.$$

This is consistent with what we found in [11] by using generalized functions (distributions). See also equations (22) below. In [11] we further showed that near $(0, 0)$ the double zeta function has the following asymptotic expansion:

$$\zeta(s_1, s_2) = \frac{4s_1 + 5s_2}{12(s_1 + s_2)} + R(s_1, s_2)$$

where $R(s_1, s_2)$ is analytic at $(0, 0)$ and $\lim_{(s_1, s_2) \rightarrow (0, 0)} R(s_1, s_2) = 0$.

Let n, k be two non-negative integers, and $m = k - n - 2$. We now consider the double zeta function around $(s_1, s_2) = (-m, -n)$ which has the following expression by Theorem 3.2:

$$\begin{aligned} & \frac{-1}{n+1} \zeta(s_1 + s_2 - 1) - \frac{1}{2} \zeta(s_1 + s_2) - \sum_{r=2}^{n+1} \frac{B_r}{r} \binom{n}{r-1} \zeta(s_1 + s_2 + r - 1) \\ & + \alpha(m)(-1)^n \frac{B_k}{k!} n! (k - n - 2)! (s_2 + n) \zeta(s_2 + s_1 + k - 1) \end{aligned} \quad (19)$$

where $\alpha(m) = 0$ if $m \leq -1$ and $\alpha(m) = 1$ if $m \geq 0$. Note that the last term is zero when computing

$$\zeta(-m, -n) = \lim_{s_1 \rightarrow -m} \lim_{s_2 \rightarrow -n} \zeta(s_1, s_2)$$

while it has possibly nontrivial contribution for $\zeta^R(-m, -n)$ since

$$\lim_{s_2 \rightarrow -n} \lim_{s_1 \rightarrow -m} (s_2 + n) \zeta(s_2 + s_1 + k - 1) = 1.$$

We get

k, n	pole, residue	indeterminacy, $\zeta = \zeta^R(n + 2 - k, -n)$
$k = 0$	$-1/(n+1)$	NO
$k = 1$	$-1/2$	NO
$2 \nmid k, 3 \leq k \leq n+1$	NO	$\frac{B_{k-1}}{2(k-1)} + \sum_{r=k-1}^{n+1} \frac{B_r}{r} \binom{n}{r-1} \zeta(r+1-k)$
$2 \nmid k, k = n+2$	NO	$B_{k-1}/(k-1)$
$2 \nmid k, k > n+2$	NO	$B_{k-1}/2(k-1)$
$2 k, 2 \leq k \leq n+1$	$(-1)^{k+1} \frac{B_k}{k} \binom{n}{k-1}$	NO

Table 1: Poles and indeterminacy of double zeta function.

When $m \geq 0$ and $2|k$ the values of $\zeta(-m, -n)$ and $\zeta^R(-m, -n)$ are different in general:

$$\zeta(-m, -n) = \frac{B_k}{k(n+1)} + \sum_{r=1}^{n+1} \frac{B_r}{r} \binom{n}{r-1} \frac{B_{k-r}}{k-r} \quad (20)$$

$$\zeta^R(-m, -n) = \zeta(-m, -n) + (-1)^n \frac{B_k}{k!} n! (k - n - 2)!. \quad (21)$$

Note that the term corresponding to $r = 1$ is non-zero if and only if $k = 2$ (and $n = 0$). From this observation we again recover that

$$\zeta(0, 0) = \frac{1}{3}, \quad \zeta^R(0, 0) = \frac{5}{12}. \quad (22)$$

We now consider the q -double zeta function.

Theorem 6.1. *Let k, n be two non-negative integers, and $m = k - n - 2$. If $m \leq -1$ then the q -double zeta function $\zeta_q(s_1, s_2)$ has a pole at $(-m, -n)$ with residue given by:*

$$\frac{\text{Res}_{(s_1, s_2) = (-m, -n)} \zeta_q(s_1, s_2)}{-(1-q)^{2-k} (\log q)^{-1}} = \begin{cases} \sum_{r=0}^k (-1)^r \binom{n+1-r}{k-r} \binom{n}{r} / (q^{n+1-r} - 1) & \text{if } k \leq n, \\ \sum_{r=0}^n (-1)^r \binom{n}{r} / (q^{n+1-r} - 1) - \frac{(-1)^n}{(n+1) \log q} & \text{if } k = n+1. \end{cases}$$

Proof. Use Theorem 2.4. □

Corollary 6.2. *Let n be a non-negative integer. Then*

$$\operatorname{Res}_{(s_1, s_2)=(1, -n)} \zeta_q(s_1, s_2) = \frac{q-1}{\log q} \zeta_q(-n) \quad (23)$$

and

$$\lim_{q \uparrow 1} \zeta_q(-n) = -\frac{B_{n+1}}{n+1} = \operatorname{Res}_{(s_1, s_2)=(1, -n)} \zeta(s_1, s_2). \quad (24)$$

Proof. Equation (23) follows from the case $m = -1$ in the above theorem and [5, (6)]:

$$\zeta_q(-n) = (1-q)^{-n} \left\{ \sum_{r=0}^n (-1)^r \binom{n}{r} / (q^{n+1-r} - 1) - \frac{(-1)^n}{(n+1) \log q} \right\}.$$

The first equality in (24) is [5, Thm. 1] and the second equality follows from Table 1. \square

Corollary 6.3. *Let k, n be two non-negative integers, $m = k - n - 2 \leq -2$. Then*

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2)=(-m, -n)} \zeta_q(s_1, s_2) = \operatorname{Res}_{(s_1, s_2)=(-m, -n)} \zeta(s_1, s_2). \quad (25)$$

Proof. By Theorem 6.1 and Table 1 we only need to prove

$$\begin{aligned} \lim_{q \uparrow 1} (1-q)^{1-k} \sum_{r=0}^k (-1)^r \binom{n+1-r}{k-r} \binom{n}{r} / (q^{n+1-r} - 1) \\ = \begin{cases} \frac{-1}{n+1} & \text{if } k = 0, \\ (-1)^{1-k} \frac{B_k}{k} \binom{n}{k-1} & \text{if } 1 \leq k \leq n. \end{cases} \end{aligned} \quad (26)$$

First by generating function of the Bernoulli numbers

$$\frac{1}{q^{n+1-r} - 1} = \frac{1}{e^{(n+1-r) \log q} - 1} = \sum_{l=0}^{\infty} \frac{B_l}{l!} ((n+1-r) \log q)^{l-1}.$$

Plugging this into the left hand side of equation (26), replacing $1-q$ by $-\log q$, and exchanging the summation we get

$$\begin{aligned} (-\log q)^{1-k} \sum_{r=0}^k (-1)^r \binom{n+1-r}{k-r} \binom{n}{r} / (q^{n+1-r} - 1) \\ = (-1)^{1-k} \sum_{l=0}^{\infty} \frac{B_l}{l!} (\log q)^{l-k} \sum_{r=0}^k (-1)^r \binom{n+1-r}{k-r} \binom{n}{r} (n+1-r)^{l-1}. \end{aligned} \quad (27)$$

Then the inner sum over r is the coefficient of x^k of the following polynomial

$$\begin{aligned} f_l(x) &= \sum_{k=0}^{n+1} \sum_{r=0}^k (-1)^r \binom{n+1-r}{k-r} \binom{n}{r} (n+1-r)^{l-1} x^k \\ &= \sum_{r=0}^n (-1)^r \binom{n}{r} (n+1-r)^{l-1} \sum_{k=r}^{n+1} \binom{n+1-r}{k-r} x^k \\ &= (x+1)^{n+1} \sum_{r=0}^n (-y)^r \binom{n}{r} (n+1-r)^{l-1}, \end{aligned} \quad (28)$$

where $y = x/(x+1)$. When $l = 0$ this expression becomes

$$f_0(x) = \frac{(x+1)^{n+1}}{n+1} \sum_{r=0}^n (-y)^r \binom{n+1}{r} = \frac{(x+1)^{n+1}}{n+1} \left[(1-y)^{n+1} - (-y)^{n+1} \right] = \frac{1}{n+1} \left[1 - (-x)^{n+1} \right].$$

Note that $k \leq n$ we see the coefficient of x^k in $f_0(x)$ is 0 if $k > 0$ and it's $1/(n+1)$ if $k = 0$. If $k = 0$ then only the constant term $-1/(n+1)$ in (27) remains when $q \uparrow 1$ which proves the corollary in this case. So we can assume $l, k > 0$. Then

$$\begin{aligned} f_l(x) &= \left(z \frac{d}{dz} \right)^{l-1} \left\{ (x+1)^{n+1} \sum_{r=0}^n (-y)^r \binom{n}{r} z^{n+1-r} \right\} \Big|_{z=1} \\ &= \left(z \frac{d}{dz} \right)^{l-1} \left\{ (x+1)^{n+1} z(z-y)^n \right\} \Big|_{z=1}. \end{aligned}$$

Note that highest degree term in $f_l(x)$ is contained in

$$\begin{aligned} (x+1)^{n+1} \left(\frac{d}{dz} \right)^{l-1} (z-y)^n \Big|_{z=1} &= n(n-1) \cdots (n-l+2) (x+1)^{n+1} (1-y)^{n-l+1} \\ &= n(n-1) \cdots (n-l+2) (x+1)^l. \end{aligned}$$

If $l = 1$ one can easily modify this to get just $x+1$. If $l < k$ then the coefficient of x^k in $f_l(x)$ is 0. If $l = k$ it is equal to

$$n(n-1) \cdots (n-k+2) = (k-1)! \binom{n}{k-1}.$$

The last express is valid even for $k = l = 1$. Thus the range of l in the outer sum of (27) starts from k . Moreover, the first term of (27) is

$$(-1)^{1-k} \frac{B_k}{k} \binom{n}{k-1}$$

as desired. This completes the proof of the corollary. \square

Proposition 6.4. *Let k, n be two non-negative integers such that $n \geq k$ and k is even. Let $m = k - n - 2$. Then*

$$\operatorname{Res}_{(s_1, s_2) = (-m, -n)} \zeta_q(s_1, s_2) = \frac{-f(q)(q-1)/\log q}{D(q)}, \quad D(q) = \prod_{j=n+1-k}^{n+1} F(q, j)^{\epsilon_j}$$

where $f(q) \in \mathbb{Z}[q]$ is a palindrome with leading coefficient $\binom{n}{k}$, $F(q, j) \in \mathbb{Z}[q]$ is a factor of $(q^j - 1)/(q - 1)$, $\epsilon = 0$ or 1 , and $\deg_q D(q) = n + \deg_q f(q)$, such that

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2) = (-m, -n)} \zeta_q(s_1, s_2) = \operatorname{Res}_{(s_1, s_2) = (-m, -n)} \zeta(s_1, s_2).$$

Proof. The computational proof is left as an exercise for the interested readers. \square

Example 6.5. By Theorem 6.1 we find with the help of Maple

$$\operatorname{Res}_{(s_1, s_2) = (4, -4)} \zeta_q(s_1, s_2) = \frac{-2q^3(3q^2 + 4q + 3)(q-1)/\log q}{P_1(q, 2)P_1(q, 3)P_1(q, 4)}.$$

where $P_a(q, m) = \sum_{j=0}^m q^{aj}$. Moreover we can check that

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2) = (4, -4)} \zeta_q(s_1, s_2) = \operatorname{Res}_{(s_1, s_2) = (4, -4)} \zeta(s_1, s_2) = -\frac{1}{3}$$

by Table 1 with $k = 2$ and $n = 4$.

Example 6.6. From Theorem 6.1 we get

$$\operatorname{Res}_{(s_1, s_2) = (6, -8)} \zeta_q(s_1, s_2) = \frac{-14q^5 g(q)(q-1)/\log q}{P_1(q, 4)P_1(q, 5)P_1(q, 6)P_2(q, 3)P_3(q, 2)}$$

where $g(q)$ is a polynomial in q of degree 14 satisfying

$$q^{14}f(1/q) = f(q) = 5q^{14} + 6q^{13} + 8q^{12} + 7q^{11} - q^{10} - 20q^9 - 30q^8 - 34q^7 - \dots$$

Then we can compute with Maple

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2) = (6, -8)} \zeta_q(s_1, s_2) = \operatorname{Res}_{(s_1, s_2) = (6, -8)} \zeta(s_1, s_2) = \frac{7}{15}$$

by Table 1 with $k = 4$ and $n = 8$.

Example 6.7. Consider the point $(s_1, s_2) = (5, -9)$. We have

$$\operatorname{Res}_{(s_1, s_2) = (5, -9)} \zeta_q(s_1, s_2) = \frac{-42q^4 g(q)(q-1)/\log q}{P_1(q, 4)P_1(q, 6)P_1(q, 7)P_2(q, 2)P_3(q, 2)A(q, 4)}$$

where $A(q, m) = \sum_{j=0}^m (-1)^j q^j$ and $g(q)$ is a polynomial in q of degree 18 satisfying

$$q^{18}g(1/q) = g(q) = 2q^{18} - q^{17} - 7q^{15} - 11q^{14} - 16q^{13} - 4q^{12} + 9q^{11} + 28q^{10} + 30q^9 + \dots$$

so that we again have the equality

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2) = (5, -9)} \zeta_q(s_1, s_2) = \operatorname{Res}_{(s_1, s_2) = (5, -9)} \zeta(s_1, s_2) = -\frac{1}{2}$$

by Table 1 with $k = 6$ and $n = 9$.

Theorem 6.8. Let m, n be two non-negative integers and $k = m + n + 2$. Then $\zeta_q(s_1, s_2)$ has indeterminacy at $(-m, -n)$ such that

$$\begin{aligned} \zeta_q(-m, -n) = & (1-q)^{2-k} \left\{ \frac{(-1)^k}{(m+1)(n+1)(\log q)^2} \right. \\ & + \sum_{r=0}^m \frac{(-1)^{r+n+1}}{(n+1)\log q} \binom{m}{r} \frac{1}{q^{m+1-r}-1} \\ & + \sum_{r=0}^n \frac{(-1)^{r+m+1}}{\log q} \frac{m!(n+1-r)!}{(k-r)!} \binom{n}{r} \frac{1}{q^{n+1-r}-1} \\ & \left. + \sum_{r_1=0}^m \sum_{r_2=0}^n (-1)^{r_1+r_2} \binom{m}{r_1} \binom{n}{r_2} \frac{1}{q^{n+1-r_2}-1} \frac{1}{q^{k-r_1-r_2}-1} \right\}, \end{aligned}$$

and

$$\begin{aligned} \zeta_q^R(-m, -n) = & (1-q)^{2-k} \left\{ \sum_{r=0}^m \frac{(-1)^{r+n+1}}{(n+1)\log q} \binom{m}{r} \frac{1}{q^{m+1-r}-1} \right. \\ & + \sum_{r=0}^m \frac{(-1)^{r+n}}{\log q} \binom{k-n-2}{r} \frac{n!(m+1-r)!}{(k-r)!} \frac{q^{m+1-r}}{q^{m+1-r}-1} \\ & \left. + \sum_{r_1=0}^m \sum_{r_2=0}^n (-1)^{r_1+r_2} \binom{m}{r_1} \binom{n}{r_2} \frac{1}{q^{n+1-r_2}-1} \frac{1}{q^{k-r_1-r_2}-1} \right\}. \end{aligned}$$

Proof. Use Theorem 2.4. □

Similar to Cor. 6.2 and Cor. 6.3 we have

Corollary 6.9. Let m and n be two non-negative integers. Then

$$\lim_{q \uparrow 1} \zeta_q(-m, -n) = \zeta(s_1, s_2), \quad \lim_{q \uparrow 1} \zeta_q^R(-m, -n) = \zeta^R(s_1, s_2). \quad (29)$$

Proof. Set $k = m + n + 2$. We consider $\zeta_q^R(-m, -n)$ first. From Theorem 6.8 we get

$$\zeta_q^R(-m, -n) = (q-1)^{-k} \left(\frac{q-1}{\log q} \right)^2 (A + B + C),$$

where

$$\begin{aligned} A &= \sum_{j=0}^{\infty} \frac{B_j}{j!} (\log q)^j \sum_{r=0}^m \frac{(-1)^{m+r+1}}{n+1} \binom{m}{r} (m+1-r)^{j-1}, \\ B &= \sum_{i=0}^{\infty} \frac{B_i}{i!} (\log q)^i \sum_{r=0}^m (-1)^{r+m+i} \binom{m}{r} \frac{n!(m+1-r)!}{(k-r)!} (m+1-r)^{i-1}, \\ C &= \sum_{i,j=0}^{\infty} \frac{B_i B_j}{i! j!} (\log q)^{i+j} \sum_{r_1=0}^m \sum_{r_2=0}^n (-1)^{k+r_1+r_2} \binom{m}{r_1} \binom{n}{r_2} (n+1-r_2)^{i-1} (k-r_1-r_2)^{j-1}. \end{aligned}$$

We first compute B as follows. Write

$$B = \sum_{i=0}^{\infty} \frac{B_i}{i!} (\log q)^i W(m, n, i),$$

where

$$W(m, n, i) = \sum_{r=0}^m (-1)^{r+k+n+i} \frac{m!n!}{r!(k-r)!} (m+1-r)^i.$$

If $i = 0$ then we can prove by decreasing induction on n that

$$W(m, n, 0) = \sum_{r=0}^m (-1)^{r+k+n} \frac{m!n!}{r!(k-r)!} = \frac{1}{k(n+1)}. \quad (30)$$

This is trivial if $n = k - 2$. Suppose (30) is true for $n \geq 1$ then we have

$$W(m, n-1, 0) = -\frac{k-n-1}{n} W(m, n, 0) + \frac{(n-1)!}{(n+1)!} = \frac{1}{kn},$$

as desired.

Similarly, we can compute C as follows. Put

$$C = \sum_{i,j=0}^{\infty} \frac{B_i B_j (\log q)^{i+j}}{i! j! (n+1)} \sum_{r_1=0}^m \sum_{r_2=0}^n (-1)^{k+r_1+r_2} \binom{m}{r_1} \binom{n+1}{r_2} (n+1-r_2)^i (k-r_1-r_2)^{j-1}.$$

We now change the upper limit of r_2 from n to $n+1$ in the above. The extra terms correspond to those by setting $i = 0$ and $r_2 = n+1$, which produce exactly A . Therefore,

$$C = \sum_{i,j=0}^{\infty} \frac{B_i B_j (\log q)^{i+j}}{i! j!} V(m, n, i, j) - A$$

where

$$V(m, n, i, j) = \sum_{r_1=0}^m \sum_{r_2=0}^{n+1} \frac{(-1)^{k+r_1+r_2}}{n+1} \binom{m}{r_1} \binom{n+1}{r_2} (n+1-r_2)^i (k-r_1-r_2)^{j-1}.$$

For $j \geq 1$ we have

$$\begin{aligned}
V(m, n, i, j) &= \left(x \frac{d}{dx}\right)^{j-1} \left\{ \left(y \frac{d}{dy}\right)^i \left\{ \sum_{r_1=0}^m \sum_{r_2=0}^{n+1} \frac{(-1)^{k+r_1+r_2}}{n+1} \right. \right. \\
&\quad \left. \left. \binom{m}{r_1} \binom{n+1}{r_2} y^{n+1-r_2} x^{k-r_1-r_2} \right\} \Big|_{y=1} \right\} \Big|_{x=1} \\
&= \frac{(-1)^k}{n+1} \left(x \frac{d}{dx}\right)^{j-1} \left\{ \left(y \frac{d}{dy}\right)^i \left\{ x(xy-1)^{n+1} (x-1)^{k-n-2} \right\} \Big|_{y=1} \right\} \Big|_{x=1} \quad (31) \\
&= \begin{cases} 0 & \text{if } i+j < k, \\ \frac{(-1)^k n!}{(n+1-i)!} (k-i-1)! & \text{if } i+j = k, i \leq n+1. \end{cases}
\end{aligned}$$

We have used the fact that if $i+j = k$ and $i > n+1$ then $j < k-n-1$ and by exchanging the two operators $x(d/dx)$ and $y(d/dy)$ we can easily show that (31) is zero. So if $l < k$ the total contribution to the coefficient of $(\log q)^l$ from $V(m, n, i, j)$ with $j > 0$ is trivial and if $l = k$ it is equal to

$$\sum_{i=0}^{n+1} B_i B_{k-i} \frac{(-1)^k n! (k-i-1)!}{i! (k-i)! (n+1-i)!} = \begin{cases} \frac{B_{k-1}}{2(k-1)} & \text{if } 2 \nmid k, \\ \frac{B_k}{k(n+1)} + \sum_{i=1}^{n+1} \frac{B_i B_{k-i}}{i(k-i)} \binom{n}{i-1} & \text{if } 2 \mid k, \end{cases} \quad (32)$$

because $k \geq n+2 \geq 2$ and $B_k = 0$ if k is odd.

To deal with $V(m, n, i, 0)$ note that equation (31) still makes sense if we interpret the operator $x(d/dx)^{-1}$ as follows:

$$\left(x \frac{d}{dx}\right)^{-1} \left\{ F(x) \right\} \Big|_{x=1} = \int_0^1 \frac{F(x)}{x} dx$$

whenever $F(0) = 0$. Thus we get

$$V(m, n, i, 0) = \frac{(-1)^k}{n+1} \left(y \frac{d}{dy}\right)^i \left\{ \int_0^1 (xy-1)^{n+1} (x-1)^m dx \right\} \Big|_{y=1}.$$

Therefore if $i = 0$ then we get

$$V(m, n, 0, 0) = \frac{(-1)^k}{n+1} \int_0^1 (x-1)^{k-1} dx = -\frac{1}{k(n+1)} = -W(m, n, 0) \quad (33)$$

from equation (30). If $i \geq 1$ then integrating by parts we get

$$\begin{aligned}
&\int_0^1 (xy-1)^{n+1} (x-1)^m dx \\
&= \frac{(xy-1)^{n+2}}{y(n+2)} (x-1)^m \Big|_0^1 - \frac{m}{y(n+2)} \int_0^1 (xy-1)^{n+2} (x-1)^{m-1} dx \\
&= \dots \\
&= (-1)^{k+1} \left(\frac{1}{y(n+2)} - \frac{m}{y^2(n+2)(n+3)} + \dots + (-1)^{m-1} \frac{m!(n+1)!}{y^m(m+n+1)!} \right. \\
&\quad \left. + (-1)^m \frac{m!(n+1)!}{y^m(m+n+1)!} \int_0^1 (xy-1)^{m+n+1} dx, \right) \\
&= (-1)^{k+n} \frac{m!(n+1)!}{k!} \frac{(y-1)^k}{y^{m+1}} + \sum_{r=0}^m (-1)^{r+k+1} \frac{m!(n+1)!}{y^{r+1}(m-r)!(n+2+r)!}.
\end{aligned}$$

It follows from changing the index r to $m - r$ that

$$\begin{aligned} V(m, n, i, 0) &= (-1)^{k+n} \frac{m!n!}{k!} \left(y \frac{d}{dy} \right)^i \left\{ \frac{(y-1)^k}{y^{m+1}} \right\} \Big|_{y=1} + \sum_{r=0}^m (-1)^{r+n+i+1} \frac{m!n!}{r!(k-r)!} (m+1-r)^i \\ &= \begin{cases} (-1)^{k+1} W(m, n, 0) & \text{if } 0 < i < k, \\ (-1)^{k+1} W(m, n, 0) + (-1)^{k+n} n! (k-n-2)! & \text{if } i = k. \end{cases} \end{aligned} \quad (34)$$

Thus when $0 < i < k$ and k is even we have $V(m, n, i, 0) = -W(m, n, i)$. It follows from (32), (33) and (34) that

$$\lim_{q \uparrow 1} \zeta_q^R(-m, -n) = \zeta^R(s_1, s_2)$$

since $B_k = 0$ if $k > 2$ is odd.

Let's turn to prove the first equality in (29). By Theorem 6.8 we have

$$\zeta_q(-m, -n) = (q-1)^{-k} \left(\frac{q-1}{\log q} \right)^2 (D + A + E + C)$$

where A and C are as above and

$$D = \frac{1}{(m+1)(n+1)}, \quad E = \sum_{i=0}^{\infty} \frac{B_i}{i!} (\log q)^i U(m, n, i),$$

where

$$U(m, n, i) = - \sum_{r=0}^n (-1)^{r+n} \frac{m!n!}{r!(k-r)!} (n+1-r)^i.$$

Hence

$$U(m, n, 0) = W(n, m, 0) = \frac{-1}{k(m+1)} = \frac{-1}{k(k-n-1)} = \frac{1}{k(n+1)} - D. \quad (35)$$

We only need to show that

$$U(m, n, i) = \begin{cases} W(m, n, i) & \text{if } 0 < i < k, \\ W(m, n, i) - (-1)^n n! (k-n-2)! & \text{if } i = k. \end{cases} \quad (36)$$

Indeed when $i > 0$ we have

$$\begin{aligned} U(m, n, i) &= (-1)^{n+1} \left(y \frac{d}{dy} \right)^i \left\{ \sum_{r=0}^n (-1)^r \frac{m!n!}{k!} \binom{k}{r} y^{n+1-r} \right\} \Big|_{y=1} \\ &= (-1)^{n+1} \frac{m!n!}{k!} \left(y \frac{d}{dy} \right)^i \left\{ y^{n+1-k} (y-1)^k - \sum_{r=n+1}^k (-1)^r \binom{k}{r} y^{n+1-r} \right\} \Big|_{y=1} \\ &= (-1)^{n+1} \frac{m!n!}{k!} \left(y \frac{d}{dy} \right)^i \left\{ \frac{(y-1)^k}{y^{m+1}} \right\} + \sum_{r=n+1}^k (-1)^{r+n} \frac{m!n!}{r!(k-r)!} (n+1-r)^i. \end{aligned}$$

Then first term is 0 if $i < k$ and it's $(-1)^{n+1} m!n!$ if $i = k$. When $r = n+1$ the summand in the second term is zero since $i > 0$. So we can let r range only from $n+2$ to k . Then change the index r to $k-r$ (and let r run from 0 to m) we can see immediately that the second term is exactly $W(m, n, i)$. This proves equation (36) which together with (35) implies the first equation in (29). We thus finish the proof of the corollary. \square

We conclude this section by remarking that by shuffle relation (18) we can also analyze $\zeta_q(s_1, s_2)$ at $(-n, n+2-k)$ for any non-negative integers k and n . For example, it's easy to compute directly

that

$$\begin{aligned} \operatorname{Res}_{(s_1, s_2)=(-3, 2)} \zeta_q(s_1, s_2) &= -\frac{1}{(1-q)\log q} \sum_{r=0}^3 (-1)^r \binom{3}{r} (r+1) \frac{q^{r+1}}{1-q^{r+1}} \\ &= \frac{-q(q-1)^2}{(q+1)(q^2+1)(q^2+q+1)\log q}, \end{aligned}$$

which can be obtained also by the shuffle relation (18) and the expression

$$\operatorname{Res}_{(s_1, s_2)=(2, -3)} \zeta_q(s_1, s_2) = \frac{q(q-1)^2}{(q+1)(q^2+1)(q^2+q+1)\log q}$$

by taking $k = n = 3$ in Theorem 6.1. Thus $(-3, 2)$ is a simple pole of $\zeta_q(s_1, s_2)$. On the other hand $\zeta(s_1, s_2)$ does not have a pole along $s_1 + s_2 = -1$. Indeed we find that

$$\lim_{q \uparrow 1} \operatorname{Res}_{(s_1, s_2)=(-3, 2)} \{\zeta_q(s_1, s_2)\} = 0.$$

7 q -multiple polylogarithms

It is well known that special values of the multiple zeta function $\zeta(s_1, \dots, s_d)$ at positive integers (n_1, \dots, n_d) can be regarded as single-valued version of multiple polylogarithm $Li_{n_1, \dots, n_d}(z_1, \dots, z_d)$ evaluated at $z_1 = \dots = z_d = 1$. For $|z_j| < 1$ these functions can be defined as

$$Li_{n_1, \dots, n_d}(z_1, \dots, z_d) = \sum_{0 < k_1 < \dots < k_d} \frac{z_1^{k_1} \dots z_d^{k_d}}{k_1^{n_1} \dots k_d^{n_d}}.$$

By Chen's iterated integral

$$(-1)^d \int_0^1 \frac{dt_1}{t_1 - a_1} \circ \left(\frac{dt_1}{t_1} \right)^{\circ(n_1-1)} \circ \dots \circ \frac{dt_d}{t_d - a_d} \circ \left(\frac{dt_d}{t_d} \right)^{\circ(n_d-1)}, \quad (37)$$

where $a_j = 1/\prod_{i=j}^d z_i$ for all $j = 1, \dots, d$, we can obtain the analytic continuation of this function as a multi-valued function on $\mathbb{C}^d \setminus \mathfrak{D}_d$ where

$$\mathfrak{D}_d = \left\{ (z_1, \dots, z_d) \in \mathbb{C}^d : \prod_{i=j}^d z_i = 1, j = 1, \dots, d \right\}.$$

When $|z_j| < 1$ we define its q -analog ($0 < q < 1$) by

$$Li_{q; n_1, \dots, n_d}(z_1, \dots, z_d) = \sum_{0 < k_1 < \dots < k_d} \frac{z_1^{k_1} \dots z_d^{k_d}}{[k_1]^{n_1} \dots [k_d]^{n_d}}.$$

Clearly when $q \uparrow 1$ we recover the ordinary multiple polylogarithm. Moreover, the special value of q -multiple zeta function at positive integers

$$\zeta_q(n_1, \dots, n_d) = Li_{q; n_1, \dots, n_d}(q^{n_1-1}, \dots, q^{n_d-1}).$$

Note that our definition of the q -multiple polylogarithms is different from that of [8]. In case of logarithm and dilogarithm our definitions are different from that of [7]. We want to convince the readers that ours are also good analogs of the ordinary ones.

We can mimic the method in section 2 to get the analytic continuation of $Li_{q; n_1, \dots, n_d}(z_1, \dots, z_d)$.

Theorem 7.1. *The q -multiple polylogarithm function $Li_{q; n_1, \dots, n_d}(z_1, \dots, z_d)$ converges if $|z_j| < 1$ for all $j = 1, \dots, d$. It can be analytically continued to a multi-valued function over $\mathbb{C}^d \setminus \mathfrak{D}_{q; d}$ via the series expansion*

$$Li_{q; n_1, \dots, n_d}(z_1, \dots, z_d) = (1-q)^{n_1 + \dots + n_d} \sum_{r_1, \dots, r_d=0}^{+\infty} \prod_{j=1}^d \left[\binom{n_j + r_j - 1}{r_j} \frac{z_j^{r_j} q^{j r_j}}{1 - (z_j \dots z_d) q^{r_j + \dots + r_d}} \right]. \quad (38)$$

Proof. The first part of the lemma is obvious. Let's concentrate on the analytic continuation. By binomial expansion $(1-x)^{-n} = \sum_{r=0}^{\infty} \binom{n+r-1}{r} x^r$ we get

$$Li_{q;n_1, \dots, n_d}(z_1, \dots, z_d) = (1-q)^{n_1 + \dots + n_d} \sum_{0 < k_1 < \dots < k_d} \sum_{r_1, \dots, r_d=0}^{+\infty} \prod_{j=1}^d \binom{n_j + r_j - 1}{r_j} (z_j q^{r_j})^{k_j}.$$

As $0 < q < 1$ the series converges absolutely by Stirling's formula so we can exchange the summations. The theorem follows immediately from Lemma 2.3 by taking $x_j = z_j q^{r_j}$. \square

However, this analytic continuation is not suitable for comparing with its ordinary counterpart. Then we define Jackson's q -differential operator (cf. [3]) by

$$D_{q;z} f(z) = \frac{f(z) - f(qz)}{(1-q)z}.$$

Lemma 7.2. *Let d, n_1, \dots, n_d be positive integers. If $n_j \geq 2$ then we have*

$$D_{q;z_j} Li_{q;n_1, \dots, n_d}(z_1, \dots, z_d) = \frac{1}{z_j} Li_{q;n_1, \dots, n_j-1, \dots, n_d}(z_1, \dots, z_d);$$

if $d \geq 2$ and $n_j = 1$ then

$$\begin{aligned} D_{q;z_j} Li_{q;n_1, \dots, n_d}(z_1, \dots, z_d) &= \frac{1}{1-z_j} Li_{q;n_1, \dots, \widehat{n_j}, \dots, n_d}(z_1, \dots, z_{j-1} z_j, \dots, z_d) \\ &\quad - \frac{1}{z_j(1-z_j)} Li_{q;n_1, \dots, \widehat{n_j}, \dots, n_d}(z_1, \dots, z_j z_{j+1}, \dots, z_d). \end{aligned}$$

Here the second term does not appear if $j = d$. If $d = n_1 = 1$ then

$$D_{q;z} Li_{q;1}(z) = \frac{1}{1-z}.$$

Proof. Clear. \square

The same properties listed in the lemma are satisfied by the ordinary multiple polylogarithms. We note that the first equation in [8, Lemma 1] is valid only for $n_j \geq 2$.

Recall that for any continuous function $f(x)$ on $[a, b]$ Jackson's q -integral (cf. [4]) is defined by

$$\int_a^b f(x) d_q x := \sum_{i=0}^{\infty} f(a + q^i(b-a))(q^i - q^{i+q})(b-a).$$

Then for every $x \in [a, b]$ we have

$$\int_a^x D_{q;t} f(t) d_q t = f(x) - f(a), \quad \text{and} \quad \lim_{q \uparrow 1} \int_a^x f(t) d_q t = \int_a^x f(t) dt. \quad (39)$$

Similar to Chen's iterated integrals we can define the q -iterated integrals as follows:

$$\int_a^b \frac{d_q t_1}{t_1 - a_1} \circ \dots \circ \frac{d_q t_r}{t_r - a_r} := \int_a^b \left(\int_a^{t_r} \dots \int_a^{t_3} \left(\int_a^{t_2} \frac{d_q t_1}{t_1 - a_1} \right) \frac{d_q t_2}{t_2 - a_2} \dots \frac{d_q t_{r-1}}{t_{r-1} - a_{r-1}} \right) \frac{d_q t_r}{t_r - a_r}.$$

Define

$$\mathfrak{D}_{q;d} =: \left\{ (z_1, \dots, z_d) \in \mathbb{C}^d : \prod_{i=j}^d z_i = q^{-m}, m \in \mathbb{Z}_{\geq 0}, j = 1, \dots, d \right\}.$$

Corollary 7.3. *We can analytically continue $Li_{q;n_1,\dots,n_d}(z_1,\dots,z_d)$ to $\mathbb{C}^d \setminus \mathfrak{D}_{q;d}$ by the q -iterated integral*

$$(-1)^d \int_0^1 \frac{d_q t_1}{t_1 - a_1} \circ \left(\frac{d_q t_1}{t_1} \right)^{\circ(n_1-1)} \circ \dots \circ \frac{d_q t_d}{t_d - a_d} \circ \left(\frac{d_q t_d}{t_d} \right)^{\circ(n_d-1)},$$

where $a_j = 1/\prod_{i=j}^d z_i$ for all $j = 1, \dots, d$. Further, for all $(z_1, \dots, z_d) \in \mathbb{C}^d$ such that $\prod_{i=j}^d z_i \notin [1, +\infty)$ for $1 \leq j \leq d$ we have

$$\lim_{q \uparrow 1} Li_{q;n_1,\dots,n_d}(z_1, \dots, z_d) = Li_{n_1,\dots,n_d}(z_1, \dots, z_d),$$

where $Li_{n_1,\dots,n_d}(z_1, \dots, z_d)$ is defined by (37) with the path being the straight line segment from 0 to 1 in \mathbb{C}^1 .

Proof. It follows from Lemma 7.2 and equation (39). The singular set $\mathfrak{D}_{q;d}$ is determined by Theorem 7.1 so that for each $j = 1, \dots, d$ the function $1/(t - a_j)$ is continuous on $[0, 1]$. \square

References

- [1] S. Akiyama, S. Egami, and Y. Tanigawa, *Analytic continuation of multiple zeta-functions and their values at non-positive integers*, Acta Arith, **98** (2001), pp. 107–116.
- [2] S. Akiyama and Y. Tanigawa, *Multiple zeta values at non-positive integers*. The Ramanujan J., **5** (2001), pp. 327–351.
- [3] F.H. Jackson, *On q -functions and a certain difference operator*, Trans. Roy. Soc. Edin. **46** (1908), pp. 253–281.
- [4] F.H. Jackson, *On q -definite integrals*, Quart. J. Pure and Appl. Math. **41** (1908), pp. 193–203.
- [5] M. Kaneko, N. Kurokawa, and M. Wakayama, *A variation of Euler’s approach to values of the Riemann zeta function*, xxx.lanl.gov/abs/math.QA/0206171.
- [6] A. Klimyk and K. Schmüdgen, *Quantum groups and their representations*, Berlin, New York, Springer, 1997.
- [7] T. H. Koornwinder, *Special functions and q -commuting variables*, math.QA/9608008
- [8] K.-G. Schlesinger, *Some remarks on q -deformed multiple polylogarithms*, math.QA/0111022.
- [9] G. Gasper and M. Rahman, *Basic hypergeometric series*, Cambridge (England); New York, Cambridge University Press, 1990.
- [10] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge University Press.
- [11] J. Zhao, *Analytic continuation of multiple zeta functions*, Proc. of AMS, **128**(1999), pp. 1275–1283. MR 2000j:11132

Email: jqz@math.upenn.edu