

A note on Abelian varieties embedded in quadrics

Luis Fuentes García *

Abstract: We show that if A is a d -dimensional abelian variety in a smooth quadric of dimension $2d$ then $d = 1$ and A is an elliptic curve of bidegree $(2, 2)$ on a quadric. This extends a result of Van de Ven which says that A only can be embedded in $\mathbf{P}^{2d}$ when $d = 1$ or 2 .

Mathematics Subject Classifications (1991): Primary, 14K05; secondary, 14E25, 14C99.

Key Words: Abelian varieties, quadrics.

1 Introduction.

Let A be a d -dimensional abelian variety embedded in $\mathbf{P}^N$. It is well known that $2d \leq N$. Moreover, in [7] Van de Ven proved that the equality holds only when $d = 1$ or 2 .

It is a natural question to study the possibilities for d when the abelian variety A is embedded in any other smooth $2d$ -dimensional variety V . In particular, here we study the embedding in smooth quadrics. We obtain the following result:

Theorem 1.1 *If A is a d -dimensional abelian variety in a smooth quadric of dimension $2d$ then $d = 1$ and A is an elliptic curve of bidegree $(2, 2)$ on a quadric.*

We will use similar methods to Van de Ven's proof. The calculation of the self intersection of A in the quadric and the Riemann-Roch theorem for abelian varieties allow only the cases $d = 1, 2, 3$.

The case $d = 1$ is the classical elliptic curve of type $(2, 2)$ contained in the smooth quadric of $\mathbf{P}^3$.

When $d = 2$, A is an abelian surface in $\mathbf{P}^5$. We see that is the projection of an abelian surface $A' \subset \mathbf{P}^6$ given by a $(1, 7)$ polarization. By a result [6] due

*Supported by EAGER.

to R. Lazarsfeld, this is projectively normal and it is not contained in quadrics. Therefore, A is not contained in quadrics either.

Finally, two results [4], [5] of J.N.Iyer allow us to discard the case $d = 3$.

2 Proof of the Theorem.

Let $j : A \hookrightarrow Q$ be an embedding of a d -dimensional abelian variety into a $2d$ -dimensional smooth quadric, with $d > 1$. The Chow ring of the smooth quadric in codimension d is generated by cocycles α and β with the relations $\alpha^2 = \beta^2 = 1$, $\alpha\beta = 0$. Thus, A will be equivalent to $a\alpha + b\beta$ and

$$A.A = a^2 + b^2. \quad (1)$$

On the other hand, by the self-intersection formula ([3], pag 431) we have $A.A = j_*c_d(N_{A,Q})$. To obtain $c_d(N_{A,Q})$, let us consider the normal bundle sequence:

$$0 \longrightarrow T_A \longrightarrow j^*T_Q \longrightarrow N_{A,Q} \longrightarrow 0$$

Since the tangent bundle of an abelian variety is trivial, we see that $c(N_{A,Q}) = j^*(c(T_Q))$. We compute the class of the tangent bundle of a quadric in the following lemma:

Lemma 2.1 *Let $i : Q \hookrightarrow \mathbf{P}^{n+1}$ be an n -dimensional smooth quadric in $\mathbf{P}^{n+1}$. Then*

$$c(T_Q) = (1 + \overline{H})^{n+2}(1 + 2\overline{H})^{-1}$$

where $\overline{H} = i^*H$ and H is a hyperplane in $\mathbf{P}^{n+1}$.

Proof: We have an exact sequence:

$$0 \longrightarrow T_Q \longrightarrow i^*T_{\mathbf{P}^{n+1}} \longrightarrow N_{Q,\mathbf{P}^{n+1}} \longrightarrow 0$$

Since Q is a hypersurface $N_{Q,\mathbf{P}^{n+1}} \cong \mathcal{O}_Q(Q) \cong \mathcal{O}_Q(2\overline{H})$ and the total class of the normal bundle is $c(N_{Q,\mathbf{P}^{n+1}}) = 1 + 2\overline{H}$. On the other hand, it is well known that $c(T_{\mathbf{P}^{n+1}}) = (1 + H)^{n+2}$. Now, from the splitting principle the claim follows. ■

Let us apply this lemma to the previous situation. We obtain

$$c(N_{A,Q}) = (1 + h)^{2d+2}(1 + 2h)^{-1} = \sum_{k=0}^{2d+2} \binom{2d+2}{k} h^k \sum_{l=0}^{\infty} (-2h)^{-l}$$

where $h = j^*\overline{H}$. In particular, the top class is

$$c_d = F_d h^d, \text{ with } F_d = \sum_{k=0}^d \binom{2d+2}{k} (-2)^{(d-k)}.$$

Substituting this into the self-intersection formula, we have:

$$A.A = F_d j_* (j^* \overline{H}^d) = F_d \overline{H}^d . j_* A = F_d (a\alpha + b\beta) . \overline{H}^d = F_d (a + b).$$

Combining this expression with (1) we obtain the following relation

$$a^2 + b^2 = F_d (a + b) \quad (2)$$

or equivalently,

$$(a - \frac{F_d}{2})^2 + (b - \frac{F_d}{2})^2 = \frac{F_d^2}{2}.$$

We are interested in bounding the degree of A , when (a, b) satisfy this equation. Note that this is a circle of center $(\frac{F_d}{2}, \frac{F_d}{2})$ and radius $\frac{F_d}{\sqrt{2}}$. Since $\deg(A) = a + b$, it is clear that the maximal degree is reached when $(a, b) = (F_d, F_d)$, that is,

$$\deg(A) \leq 2F_d.$$

On the other hand, the abelian variety is embedded in $Q \subset \mathbf{P}^{2d+1}$. When $d > 2$, by Van de Ven's Theorem, it spans $\mathbf{P}^{2d+1}$. Furthermore, by the Riemann-Roch theorem for abelian varieties, we know that $h^0(\mathcal{O}_A(h)) = \frac{\deg(A)}{d!}$. Thus, we have the following inequality:

$$\deg(A) \geq 2(d+1)!$$

Comparing the two bounds we see that a sufficient condition for the non-existence of the embedding j is $F_d < (d+1)!$. Now,

$$F_d = \sum_{k=0}^d \binom{2d+2}{k} (-2)^{(d-k)} \leq \sum_{k=0}^d \binom{2d+2}{k} (2)^d \leq 2^d 2^{2d+1} = 2^{3d+1}.$$

We see that $(d+1)! > 2^{3d+1} \geq F_d$ when $d = 17$. A simple inductive argument shows that this holds if $d \geq 17$.

If $d \leq 17$, using the exact value of F_d , we see that $(d+1)! > F_d$ for any $d > 3$.

We conclude that the unique possibilities are $d = 2$ or $d = 3$.

First, suppose that A is an abelian surface contained in a quadric. $F_2 = 7$ and we can check that the unique positive integer solution of the equation (2) is $a = b = 7$. Thus A must be an abelian surface of degree 14 given by the polarization $(1, 7)$. Note that $A \subset Q \subset \mathbf{P}^5$ is not linearly normal, that is, it is the projection of a linearly normal abelian surface $A' \subset \mathbf{P}^6$. The quadric Q can be lifted to a quadric containing the surface A' .

Lazarsfeld proved in [6] that a very ample divisor of type $(1, d)$ with $d \geq 13$ or $d = 7, 8, 9$ is projectively normal. From this the following sequence is exact:

$$0 \longrightarrow H^0(I_{A', \mathbf{P}^6}(2)) \longrightarrow H^0(\mathcal{O}_{\mathbf{P}^6}(2)) \longrightarrow H^0(\mathcal{O}_{A'}(2)) \longrightarrow 0$$

Since $h^0(\mathcal{O}_{\mathbf{P}^6}(2)) = h^0(\mathcal{O}_{A'}(2)) = 28$, there are not quadrics containing the abelian surface A' and we obtain a contradiction.

Finally, suppose that $d = 3$. Now, $F_3 = 24 = (3 + 1)!$, so the degree of the abelian variety is exactly $2F_3 = 48$. The line bundle $\mathcal{O}_A(h)$ corresponds to a divisor of type $(1, 1, 8)$ or $(1, 2, 4)$. But J.N.Iyer prove in [4] that a line bundle of type $(1, \dots, 1, 2d + 1)$ is never very ample. Moreover, in [5] she studies the map defined by a line bundle of type $(1, 2, 4)$ in a generic abelian threefold. She obtains that it is birational but not an isomorphism onto its image. Note that the very ampleness is an open condition for polarized abelian varieties (see [1]). It follows that a linear system of type $(1, 2, 4)$ cannot be very ample on any abelian threefold and this completes the proof.

Remark 2.2 *The sequence $F_d = \sum_{k=0}^d \binom{2d+2}{k} (-2)^{(d-k)}$ is related to the Fine numbers. For a reference see [2].*

Acknowledgement I thank K. Hulek for suggesting me this problem and for his advice and interest. I am grateful to the Institut für Mathematik of Hannover for its hospitality and especially to E. Schellhammer for his patience. I want also thank E. Deutsch for his remark about bounding F_d and its relation with the Fine numbers.

References

- [1] DEBARRE, O.; HULEK, K.; SPANDAW, J.; *Very ample linear systems on abelian varieties.* Math. Ann., **300**, 181-202 (1994).
- [2] DEUTSCH, E.; SHAPIRO, L. *A survey of the Fine numbers.* Discrete Math., **241**, 241-265 (2001).
- [3] HARTSHORNE, R. *Algebraic Geometry.* GTM, **52**. Springer-Verlag, 1977.
- [4] IYER, J.N. *Linear systems on abelian varieties of dimension $2g + 1$.* Proc. Am. Math. Soc. **130**, No.4, 959-962 (2002).
- [5] IYER, J.N. *Line bundles of type $(1, \dots, 1, 2, \dots, 2, 4, \dots, 4)$ on abelian varieties.* Int. J. Math. **12**, No.1, 125-142 (2001)
- [6] LAZARSFELD, R. *Projectivité normale des surfaces abéliennes.* Rédigé par O. Debarre. Prepublication No. **14**, Europroj- C.I.M.P.A., Nice (1990).
- [7] VAN DE VEN, A. *On the embedding of Abelian varieties in projective spaces.* Ann. Mat. Pura Appl., IV. Ser. **103**, 127-129 (1975).

E-mail: luisfg@usc.es

Luis Fuentes García. Institut für Mathematik. 30167 Hannover, Germany.