

# ON THE CACHAZO-DOUGLAS-SEIBERG-WITTEN CONJECTURE FOR SIMPLE LIE ALGEBRAS

PAVEL ETINGOF AND VICTOR KAC

ABSTRACT. Recently, motivated by supersymmetric gauge theory, Cachazo, Douglas, Seiberg, and Witten proposed a conjecture about finite dimensional simple Lie algebras, and checked it in the classical cases. We prove the conjecture for type  $G_2$ , and also verify a consequence of the conjecture in the general case.

## 1. THE CDSW CONJECTURE

Let  $\mathfrak{g}$  be a simple finite dimensional Lie algebra over  $\mathbb{C}$ . We fix an invariant form on  $\mathfrak{g}$  and do not distinguish  $\mathfrak{g}$  and  $\mathfrak{g}^*$ . Let  $g$  be the dual Coxeter number of  $\mathfrak{g}$ . Consider the associative algebra  $R = \wedge(\mathfrak{g} \oplus \mathfrak{g})$ . This algebra is naturally  $\mathbb{Z}_+$ -bigraded (with the two copies of  $\mathfrak{g}$  sitting in degrees  $(1,0)$  and  $(0,1)$ , respectively). The degree  $(2,0)$ ,  $(1,1)$ , and  $(0,2)$  components of  $R$  are  $\wedge^2 \mathfrak{g}$ ,  $\mathfrak{g} \otimes \mathfrak{g}$ ,  $\wedge^2 \mathfrak{g}$ , respectively, hence each of them canonically contains a copy of  $\mathfrak{g}$ . Let  $I$  be the ideal in  $R$  generated by these three copies of  $\mathfrak{g}$ , and  $A = R/I$ .

The associative algebra  $A$  may be interpreted as follows. Let  $\Pi\mathfrak{g}$  denote  $\mathfrak{g}$  regarded as an odd vector space. Then  $R$  may be thought of as the algebra of regular functions on  $\Pi\mathfrak{g} \times \Pi\mathfrak{g}$ . We have the supercommutator map  $\{, \} : \Pi\mathfrak{g} \times \Pi\mathfrak{g} \rightarrow \mathfrak{g}$  given by the formula  $\{X, Y\} = XY + YX$ , where the products are taken in the universal enveloping algebra (this is a morphism of supermanifolds). The ideal  $I$  in  $R$  is then defined by the relations  $\{X, X\} = 0$ ,  $\{X, Y\} = 0$ ,  $\{Y, Y\} = 0$ , so  $A$  is the algebra of functions on the superscheme defined by these equations.

In [CDSW,W], the following conjecture is proposed, and proved for classical  $\mathfrak{g}$ :

**Conjecture 1.1.** (i) The algebra  $A^{\mathfrak{g}}$  of  $\mathfrak{g}$ -invariants in the algebra  $A$  is generated by the unique invariant element  $S$  of  $A$  of degree  $(1,1)$  (namely,  $S = \text{Tr}|_V(XY)$ , where  $V$  is a non-trivial irreducible finite dimensional representation of  $\mathfrak{g}$ ).

(ii)  $S^g = 0$ .

(iii)  $S^{g-1} \neq 0$ .

Here we prove the conjecture for  $\mathfrak{g}$  of type  $G_2$ . We believe that the method of the proof should be relevant in the general case.

We also prove in general the following result, which is a consequence of the conjecture.

**Proposition 1.2.** *Any homogeneous element of  $A^{\mathfrak{g}}$  is of degree  $(d, d)$  for some  $d$ ; therefore, the algebra  $A^{\mathfrak{g}}$  is purely even, and the natural action of  $sl(2)$  on it (by linear transformations of  $X, Y$ ) is trivial.*

**Remark.** Conjecture 1.1 has the following cohomological interpretation. Let  $\mathfrak{g}[x, y]$  denote the Lie algebra of polynomials of  $x, y$  with values in  $\mathfrak{g}$ . Consider the algebra of relative cohomology  $H^*(\mathfrak{g}[x, y], \mathfrak{g}, \mathbb{C})$ . It is graded by the cohomological

degree  $D$  and the  $x, y$ -degree  $d$ . It is clear that for any nonzero cochain, one has  $D \leq d$ . Thus the homogeneous elements of the algebra  $H^*(\mathfrak{g}[x, y], \mathfrak{g}, \mathbb{C})$  for which  $D = d$  form a subalgebra  $E$ . Conjecture 1.1 then states that  $E$  is generated by an element  $S$  of degree 2 with the defining relation  $S^g = 0$ .

Note that in this formulation, one can obviously replace  $\mathfrak{g}[x, y]$  with its quotient by any ideal spanned by homogeneous elements of  $x, y$ -degree  $d \geq 3$ .

## 2. PROOF

The main idea of the proof is to consider first the algebra  $B$  of functions on the superscheme of  $X \in \Pi\mathfrak{g}$  such that  $\{X, X\} = 0$ . This algebra was intensively studied by Kostant, Peterson, and others (see [K]) and is rather well understood. In particular, it is known that as a  $\mathfrak{g}$ -module,  $B$  is a direct sum of  $2^{\text{rk}(\mathfrak{g})}$  non-isomorphic simple  $\mathfrak{g}$ -modules  $V_{\mathfrak{a}}$  parametrized by abelian ideals  $\mathfrak{a}$  in a Borel subalgebra  $\mathfrak{b}$  of  $\mathfrak{g}$ . Namely, if  $\mathfrak{a}$  is such an ideal of dimension  $d$  then it defines (by taking its top exterior power) a nonzero vector  $v_{\mathfrak{a}}$  in  $\wedge^d \mathfrak{g}$  (defined up to scaling). This vector generates an irreducible submodule in  $\wedge^d \mathfrak{g}$  (with highest weight vector  $v_{\mathfrak{a}}$ ), and the sum of these submodules is a complement to the kernel of the projection  $\wedge \mathfrak{g} \rightarrow B$ . In fact, this is the unique invariant complement, because in each degree  $d$  it is the eigenspace of the quadratic Casimir  $C$  with eigenvalue  $d$  (the largest possible eigenvalue on  $\wedge^d \mathfrak{g}$ ; here the Casimir is normalized so that  $C|_{\mathfrak{g}} = 1$ ). Thus the (direct) sum of  $V_{\mathfrak{a}}$  is canonically identified with  $B$  as a  $\mathfrak{g}$ -module.

The algebra  $A$  is obtained from  $B \otimes B$  by taking a quotient by the additional relation  $\{X, Y\} = 0$  and then taking invariants. In other words,  $A = (B \otimes B)^{\mathfrak{g}}/L$ , where  $L$  is the space of invariants in the ideal  $\tilde{L}$  in  $B \otimes B$  given by the relation  $\{X, Y\} = 0$ .

Thus, let us look more carefully at the algebra  $(B \otimes B)^{\mathfrak{g}}$ . From the above we see that a basis of  $(B \otimes B)^{\mathfrak{g}}$  is given by the elements  $z_{\mathfrak{a}}$ , the canonical elements in  $V_{\mathfrak{a}} \otimes V_{\mathfrak{a}}^*$ . The element  $z_{\mathfrak{a}}$  sits in bidegree  $(d, d)$ , where  $d$  is the dimension of  $\mathfrak{a}$ . This implies Proposition 1.2 (the  $sl(2)$  action on  $A^{\mathfrak{g}}$  is trivial because the  $sl(2)$ -weight of a vector in  $A$  of degree  $(d_1, d_2)$  is  $d_1 - d_2$ ).

Now we proceed to prove the conjecture for  $G_2$ . Let  $\omega_1$  and  $\omega_2 = \theta$  be the fundamental weights of  $G_2$  ( $\theta$  is the highest root), and  $\alpha_1, \alpha_2$  the corresponding simple roots (so that  $\theta = 3\alpha_1 + 2\alpha_2$ ). The abelian ideals in the corresponding Borel subalgebra are  $\mathfrak{a}_0 = 0$ ,  $\mathfrak{a}_1 = \mathbb{C}e_{\theta}$ ,  $\mathfrak{a}_2 = \mathbb{C}e_{\theta} \oplus \mathbb{C}e_{3\alpha_1 + \alpha_2}$ ,  $\mathfrak{a}_3 = \mathbb{C}e_{\theta} \oplus \mathbb{C}e_{3\alpha_1 + \alpha_2} \oplus \mathbb{C}e_{2\alpha_1 + \alpha_2}$ . Thus the  $\mathfrak{g}$ -module  $B$  has 4 irreducible components:  $V_0 = \mathbb{C}$  in degree 0,  $V_1 = \mathfrak{g}$  in degree 1,  $V_2 = V(3\omega_1)$  in degree 2, and  $V_3 = V(\omega_1 + 2\omega_2)$  in degree 3.

Let  $S$  be the invariant element of degree  $(1, 1)$  in  $\wedge \mathfrak{g} \otimes \wedge \mathfrak{g}$ . Clearly the projection of powers of  $S$  onto  $B \otimes B$  is nonzero in degree 0, 1, 2, 3 (this follows from the fact that for each degree  $d$  one has a canonical decomposition  $\wedge^d \mathfrak{g} = V_{\mathfrak{a}_d} \oplus \text{Ker}(\wedge^d \mathfrak{g} \rightarrow B[d])$ , where  $B[d]$  is the degree  $d$  component of  $B$ ).

The dual Coxeter number of  $G_2$  is 4. Thus, our job is just to show that the ideal generated by the relation  $\{X, Y\} = 0$  in  $B \otimes B$  contains no nonzero  $\mathfrak{g}$ -invariants.

A  $\mathfrak{g}$ -invariant of degree  $(d+1, d+1)$  in this ideal is a linear combination of elements of the form  $C_w := \sum f_{ijk}(x_i \otimes x_j)w(x_k)$ , where  $x_i$  is an orthonormal basis of  $\mathfrak{g}$ ,  $f_{ijk}$  are the structure constants of  $\mathfrak{g}$  in this basis, and  $w : \mathfrak{g} \rightarrow \text{End}(V_{\mathfrak{a}}) = V_{\mathfrak{a}} \otimes V_{\mathfrak{a}}^*$  is a homomorphism of representations (here for brevity  $\mathfrak{a} := \mathfrak{a}_d$ ). It is easy to check

that the only homomorphisms  $w$  relevant to our situation (i.e., for  $d = 1, 2$ ) are given by the action of  $\mathfrak{g}$  on  $V_{\mathfrak{a}}$ .<sup>1</sup> So the result follows from the following lemma:

**Lemma 2.1.** *If  $w$  is the action map then  $C_w$  is zero in  $B \otimes B$ .*

*Proof.* Let us regard  $C_w$  as an operator on  $\wedge^{d+1}(\mathfrak{g})$ . Then it has the form

$$C_w = \sum f_{ijk} W_{x_i} L_{x_k} I_{x_j},$$

where  $I, L, W$  are the operators of contraction, Lie derivative, and wedging, respectively. This can be more shortly written as

$$C_w = \sum W_{x_i} L_{[x_i, x_j]} I_{x_j}.$$

We claim that if  $v \in V_{\mathfrak{a}}$  then  $C_w v$  is zero (this would imply that  $C_w$  is zero in  $B \otimes B$ ). Set  $y = p_1 \wedge \dots \wedge p_{d+1}$ . Then we get

$$\begin{aligned} C_w y &= \sum_{i,j,r,s} (-1)^{r+s} x_i \wedge ([x_i, p_r], p_s) - ([x_i, p_s], p_r) \wedge p_1 \wedge \dots \wedge \hat{p}_r \dots \wedge \hat{p}_s \dots \wedge p_{d+1} = \\ &= \sum_{i,j,r,s} (-1)^{r+s} x_i \wedge [[p_s, p_r] x_i] \wedge p_1 \wedge \dots \wedge \hat{p}_r \dots \wedge \hat{p}_s \dots \wedge p_{d+1}. \end{aligned}$$

This implies the result, since  $V_{\mathfrak{a}}$  coincides with the kernel of the map  $\wedge^{d+1} \mathfrak{g} \rightarrow \mathfrak{g} \otimes \wedge^{d-1} \mathfrak{g}$ , given by

$$y \mapsto \sum_{r,s} (-1)^{r+s} [p_s, p_r] \otimes p_1 \wedge \dots \wedge \hat{p}_r \dots \wedge \hat{p}_s \dots \wedge p_{d+1}.$$

□

## References

[CDSV] Freddy Cachazo, Michael R. Douglas, Nathan Seiberg, Edward Witten, Chiral Rings and Anomalies in Supersymmetric Gauge Theory, hep-th/0211170.

[K] Bertram Kostant, On  $\wedge \mathfrak{g}$  for a semisimple Lie algebra  $\mathfrak{g}$ , as an equivariant module over the symmetric algebra  $S(\mathfrak{g})$ . Analysis on homogeneous spaces and representation theory of Lie groups, Okayama–Kyoto (1997), 129–144, Adv. Stud. Pure Math., 26, Math. Soc. Japan, Tokyo, 2000.

[W] Edward Witten, Chiral Ring Of  $\mathrm{Sp}(N)$  And  $\mathrm{SO}(N)$  Supersymmetric Gauge Theory In Four Dimensions hep-th/0302194.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

*E-mail address:* [etingof@math.mit.edu](mailto:etingof@math.mit.edu)

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

*E-mail address:* [kac@math.mit.edu](mailto:kac@math.mit.edu)

---

<sup>1</sup>This has to do with the fact that the highest weight of  $V_2$  does not involve  $\omega_2$ . The situation is exactly the same for type  $B_2$ , except that it is not true there that all  $w$  come from the action of  $\mathfrak{g}$ ; this results in  $S^3 = 0$  for  $B_2$ , while  $S^3 \neq 0$  for  $G_2$ .