

# Differential calculus on a novel cross-product quantum algebra

**Deepak Parashar**

Department of Mathematics, University of Wales Swansea, Singleton Park, Swansea SA2 8PP, United Kingdom

E-mail: D.Parashar@swansea.ac.uk

URL: <http://www-maths.swan.ac.uk/staff/dp/>

**Abstract.** We investigate the algebro-geometric structure of a novel two-parameter quantum deformation which exhibits the nature of a semidirect or cross-product algebra built upon  $GL(2) \otimes GL(1)$ , and is related to several other known examples of quantum groups. Following the R-matrix framework, we construct the  $L^\pm$  functionals and address the problem of duality for this quantum group. This naturally leads to the construction of a bicovariant differential calculus that depends only on one deformation parameter, respects the cross-product structure and has interesting applications. The corresponding Jordanian and hybrid deformation is also explored.

## 1. The quantum algebra $\mathcal{A}_{r,s}$

The biparametric  $q$ -deformation  $\mathcal{A}_{r,s}$  is defined [1] to be the semidirect or cross-product  $GL_r(2) \rtimes \mathbb{C}[f, f^{-1}]$  built on the vector space  $GL_r(2) \otimes \mathbb{C}[f, f^{-1}]$  where  $GL_r(2) = \mathbb{C}[a, b, c, d]$  modulo the relations

$$\begin{aligned} ab &= r^{-1}ba, & bd &= r^{-1}db \\ ac &= r^{-1}ca, & cd &= r^{-1}dc \\ bc &= cb, & [a, d] &= (r^{-1} - r)bc \end{aligned} \tag{1}$$

and  $\mathbb{C}[f, f^{-1}]$  has the cross relations

$$\begin{aligned} af &= fa, & cf &= sfc \\ bf &= s^{-1}fb, & df &= fd \end{aligned} \tag{2}$$

$\mathcal{A}_{r,s}$  can also be interpreted as a skew Laurent polynomial ring  $GL_r[f, f^{-1}; \sigma]$  where  $\sigma$  is the automorphism given by the action of element  $f$  on  $GL_r(2)$ . If we let  $A = GL_r(2)$  and  $H = \mathbb{C}[f, f^{-1}]$ , then  $A$  is a left  $H$ -module algebra and the action of  $f$  is given by

$$f \triangleright a = a, \quad f \triangleright b = sb, \quad f \triangleright c = s^{-1}c, \quad f \triangleright d = d \tag{3}$$

## 2. The dual algebra $\mathcal{U}_{r,s}$

Knowing properties of cross-product algebras [2, 3], we already know that the algebra dual to  $\mathcal{A}_{r,s}$  would be the cross-coproduct coalgebra  $\mathcal{U}_{r,s} = U_r(gl(2)) \rtimes \mathbb{C}[[\phi]]$  with  $\phi$  as an element dual to  $f$ . As a vector space, the dual is  $\mathcal{U}_{r,s} = U_r(gl(2)) \otimes U(u(1))$ . Now, the duality relation between  $\langle GL_r(2), U_r(gl(2)) \rangle$  is already well-known [4], while that between  $\langle \mathbb{C}[f, f^{-1}], U(u(1)) \rangle$  is given by  $\langle f, \phi \rangle = 1$ , i.e.,  $U(u(1)) = \mathbb{C}[[\phi]]$ . More precisely, we work

algebraically with  $\mathbb{C}[s^\phi, s^{-\phi}]$  where  $\langle f, s^\phi \rangle = s$ . This induces duality on the vector space tensor products, the left action dualises to the left coaction, and this results in the dual algebra being a cross-coproduct  $\mathcal{U}_{r,s} = U_r(gl(2)) \underset{s}{\bowtie} \mathbb{C}[[\phi]]$ . Let us recall that  $U_r(gl(2))$ , the algebra dual to  $GL_r(2)$ , is isomorphic to the tensor product  $U_r(sl(2)) \otimes \tilde{U}(u(1))$  where  $U_r(sl(2))$  has the usual generators  $\{H, X_\pm\}$  and  $\tilde{U}(u(1)) = \mathbb{C}[[\xi]] = \mathbb{C}[r^\xi, r^{-\xi}]$  with  $\xi$  central. Therefore,  $\mathcal{U}_{r,s}$  is nothing but  $U_r(sl(2))$  and two central generators  $\xi$  and  $\phi$ , where  $\xi$  is the generating element of  $\tilde{U}(u(1))$  and  $\phi$  is the generating element of  $U(u(1))$ . Also note that  $s^\phi$  is dually paired with the element  $f$  of  $\mathcal{A}_{r,s}$ .

### 3. R-matrix relations

In the quantum group language,  $\mathcal{A}_{r,s}$  is understood as a novel Hopf algebra [5, 6] generated by  $\{a, b, c, d, f\}$  arranged in the matrix form

$$T = \begin{pmatrix} f & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{pmatrix} \quad (4)$$

with the labelling 0, 1, 2, and  $\{r, s\}$  are the two deformation parameters. The  $R$ -matrix

$$R = \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & S^{-1} & 0 & 0 \\ 0 & \Lambda & S & 0 \\ 0 & 0 & 0 & R_r \end{pmatrix} \quad (5)$$

is in block form, i.e., in the order (00), (01), (02), (10), (20), (11), (12), (21), (22) (which is chosen in conjunction with the block form of the  $T$ -matrix) where

$$R_r = \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & \lambda & 1 & 0 \\ 0 & 0 & 0 & r \end{pmatrix}; \quad S = \begin{pmatrix} s & 0 \\ 0 & 1 \end{pmatrix}; \quad \Lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}; \quad \lambda = r - r^{-1}$$

The  $RTT$  relations  $RT_1T_2 = T_2T_1R$  (where  $T_1 = T \otimes \mathbf{1}$  and  $T_2 = \mathbf{1} \otimes T$ ) then yield the commutation relations (1) and (2) between the generators. The Hopf algebra structure underlying  $\mathcal{A}_{r,s}$  is  $\Delta(T) = T \dot{\otimes} T$ ,  $\varepsilon(T) = \mathbf{1}$ . The Casimir operator  $\delta = ad - r^{-1}bc$  is invertible and the antipode is

$$S(f) = f^{-1}, \quad S(a) = \delta^{-1}d, \quad S(b) = -\delta^{-1}rb, \quad S(c) = -\delta^{-1}r^{-1}c, \quad S(d) = \delta^{-1}a \quad (6)$$

The quantum determinant  $\mathcal{D} = \delta f$  is group-like but not central. There are several interesting features of this deformation [5, 6, 1] which cannot be mentioned here. In the  $R$ -matrix formulation of matrix quantum groups, a basic step is to construct functionals (matrices)  $L^+$  and  $L^-$  which are dual to the matrix of generators (4) in the fundamental representation. These functions are defined by their value on the matrix of generators  $T$

$$\langle (L^\pm)_b^a, T_d^c \rangle = (R^\pm)_{bd}^{ac} \quad (7)$$

where

$$(R^+)^{ac}_{bd} = c^+(R)^{ca}_{db} \quad (8)$$

$$(R^-)^{ac}_{bd} = c^-(R^{-1})^{ac}_{bd} \quad (9)$$

and  $c^+, c^-$  are free parameters. For  $\mathcal{A}_{r,s}$  we make the following ansatz for the  $L^\pm$  matrices:

$$L^+ = \begin{pmatrix} J & 0 & 0 \\ 0 & M & P \\ 0 & 0 & N \end{pmatrix} \quad \text{and} \quad L^- = \begin{pmatrix} J' & 0 & 0 \\ 0 & M' & 0 \\ 0 & Q & N' \end{pmatrix} \quad (10)$$

where

$$\begin{aligned} J &= s^{-\frac{1}{2}(F-A+D-1)} r^{\frac{1}{2}(F-A-D+1)}, & J' &= s^{-\frac{1}{2}(F-A+D-1)} r^{-\frac{1}{2}(F-A-D+1)} \\ M &= s^{-\frac{1}{2}(F-A-D+1)} r^{\frac{1}{2}(-F+A-D+1)}, & M' &= s^{-\frac{1}{2}(F-A-D+1)} r^{-\frac{1}{2}(-F+A-D+1)} \\ N &= s^{-\frac{1}{2}(F+A+D-1)} r^{\frac{1}{2}(-F-A+D+1)}, & N' &= s^{-\frac{1}{2}(F+A+D-1)} r^{-\frac{1}{2}(-F-A+D+1)} \\ P &= \lambda C, & Q &= -\lambda B \end{aligned}$$

and  $\{A, B, C, D, F\}$  is the set of generating elements of the dual algebra  $\mathcal{U}_{r,s}$ . This is consistent with the action on the generators of  $\mathcal{A}_{r,s}$  and gives the correct duality pairings. The commutation algebra is given by the  $RLL$  relations  $R_{12}L_2^\pm L_1^\pm = L_1^\pm L_2^\pm R_{12}$ ,  $R_{12}L_2^+ L_1^- = L_1^- L_2^+ R_{12}$ , where  $L_1^\pm = L^\pm \otimes \mathbf{1}$ ,  $L_2^\pm = \mathbf{1} \otimes L^\pm$ , and  $R_{12}$  is the same as (5). Finally, we obtain a single-parameter deformation of  $U(gl(2)) \otimes U(u(1))$  as an algebra. Including the coproduct, we again obtain  $\mathcal{U}_{r,s}$  as a semidirect product  $U_r(gl(2)) \rtimes_s U(u(1))$ .

#### 4. Differential calculus on $\mathcal{A}_{r,s}$

The  $R$ -matrix procedure [7] is known to provide a natural framework to construct differential calculus on matrix quantum groups. We note here that the  $\mathcal{A}_{r,s}$  deformation is not a full matrix quantum group, but an appropriate quotient of one (of multiparameter  $q$ -deformed  $GL(3)$ , to be precise). Nevertheless, it turns out that the constructive differential calculus methods [8] work equally well for such quotients. The bimodule  $\Gamma$  (space of quantum one-forms  $\omega$ ) is characterised by the commutation relations between  $\omega$  and  $a \in \mathcal{A}$  ( $\equiv \mathcal{A}_{r,s}$ )

$$\omega a = (\mathbf{1} \otimes g) \Delta(a) \omega \quad (11)$$

and the linear functional  $g \in \mathcal{A}' (= \text{Hom}(\mathcal{A}, \mathbb{C}))$  is defined in terms of the  $L^\pm$  matrices

$$g = S(L^+) L^- \quad (12)$$

Thus, in terms of components we have

$$\omega_{ij} a = [(\mathbf{1} \otimes S(l_{ki}^+) l_{jl}^-) \Delta(a)] \omega_{kl} \quad (13)$$

using  $L^\pm = l_{ij}^\pm$  and  $\omega = \omega_{ij}$  where  $i, j = 1..3$ . From these relations, one can obtain the commutation relations of all the left-invariant one-forms with the generating elements of  $\mathcal{A}$ . The left-invariant vector fields  $\chi_{ij}$  on  $\mathcal{A}$  are given by the expression

$$\chi_{ij} = S(l_{ik}^+) l_{kj}^- - \delta_{ij} \varepsilon \quad (14)$$

The vector fields act on the generating elements as

$$\chi_{ij} a = (S(l_{ik}^+) l_{kj}^- - \delta_{ij} \varepsilon) a \quad (15)$$

Furthermore, using the formula  $\mathbf{d}a = \sum_i (\chi_i * a)\omega^i$ , we obtain the action of the exterior derivatives ( $\mathbf{d} : \mathcal{A} \longrightarrow \Gamma$ ):

$$\mathbf{d}a = (r^{-2} - 1)a\omega^1 - \lambda b\omega^+ \quad (16)$$

$$\mathbf{d}b = \lambda^2 b\omega^1 - \lambda a\omega^- + (r^{-2} - 1)b\omega^2 \quad (17)$$

$$\mathbf{d}c = (r^{-2} - 1)c\omega^1 - \lambda d\omega^+ \quad (18)$$

$$\mathbf{d}d = \lambda^2 d\omega^1 - \lambda c\omega^- + (r^{-2} - 1)d\omega^2 \quad (19)$$

$$\mathbf{d}f = (r^{-2} - 1)f\omega^0 \quad (20)$$

where  $\omega^0 = \omega_{11}, \omega^1 = \omega_{22}, \omega^+ = \omega_{23}, \omega^- = \omega_{32}, \omega^2 = \omega_{33}$ .  $\mathbf{d}\mathcal{A}$  generates  $\Gamma$  as a left  $\mathcal{A}$ -module, and this defines a first-order differential calculus  $(\Gamma, \mathbf{d})$  on  $\mathcal{A}_{r,s}$ . The calculus is bicovariant due to the coexistence of the left ( $\Delta_L : \Gamma \longrightarrow \mathcal{A} \otimes \Gamma$ ) and the right ( $\Delta_R : \Gamma \longrightarrow \Gamma \otimes \mathcal{A}$ ) actions. Curiously, using the Leibniz rule it can be checked that

$$\mathbf{d}(af - fa) = 0, \quad \mathbf{d}(cf - sfc) = 0, \quad \mathbf{d}(bf - s^{-1}fb) = 0, \quad \mathbf{d}(df - fd) = 0, \quad (21)$$

which is consistent with cross relations (2), and so the differential calculus also respects the cross-product structure of  $\mathcal{A}_{rs}$ .

## 5. The Jordanian deformation $\mathcal{A}_{m,k}$

$\mathcal{A}_{r,s}$  can be contracted [6] (by means of singular limit of similarity transformations) to obtain a nonstandard or Jordanian analogue, say  $\mathcal{A}_{m,k}$ , with deformation parameters  $\{m, k\}$  and the associated  $R$ -matrix is triangular. In analogy with  $\mathcal{A}_{r,s}$ ,  $\mathcal{A}_{m,k}$  can also be considered as the semidirect or cross-product  $GL_m(2) \rtimes_k \mathbb{C}[f, f^{-1}]$  built upon the vector space  $GL_m(2) \otimes \mathbb{C}[f, f^{-1}]$ , where  $GL_m(2)$  is itself a Jordanian deformation of  $GL(2)$ . Thus,  $\mathcal{A}_{m,k}$  can also be interpreted as a skew Laurent polynomial ring  $GL_m[f, f^{-1}; \sigma]$  where  $\sigma$  is the automorphism given by the action of element  $f$  on  $GL_m(2)$ .

## 6. Conclusions

The  $\mathcal{A}_{r,s}$  and  $\mathcal{A}_{m,k}$  deformations provide interesting new examples of cross-product quantum algebras, both of which have  $GL(2) \otimes GL(1)$  as their classical limits. The differential calculus on  $\mathcal{A}_{r,s}$  also has an inherent cross-product structure, embeds the calculus on  $GL_q(2)$  and is also related to the calculus on  $GL_{p,q}(2)$ . It would be interesting to investigate the calculus on the Jordanian  $\mathcal{A}_{m,k}$ , and on the *hybrid/intermediate* [9] deformation obtained during the course of the contraction of  $\mathcal{A}_{r,s}$  to  $\mathcal{A}_{m,k}$ .

## References

- [1] Parashar D 2001 J. Math. Phys. 42 5431-5443
- [2] Majid S 1995 Foundations of Quantum Group Theory (Cambridge: CUP)
- [3] Klimyk A and Schmüdgen K 1997 Quantum Groups and Their Representations (Springer)
- [4] Sudbery A 1990 Proc. Workshop on Quantum Groups, Argonne (edited by Curtright T, Fairlie D and Zachos C) pp. 33-51
- [5] Basu-Mallick B 1994 hep-th/9402142
- [6] Parashar D and McDermott R J 2000 J. Math. Phys. 41 2403-2416
- [7] Faddeev L D, Reshetikhin N Y and Takhtajan L A 1990 Len. Math. J. 1 193-225
- [8] Jurčo B 1991 Lett. Math. Phys. 22 177-186; 1994 preprint CERN-TH 9417/94
- [9] Ballesteros A, Herranz F J and Parashar P 1999 J. Phys. A: Math. Gen. 32, 2369-2385