

ON THE LOCATION OF CONCENTRATION POINTS FOR SINGULARLY PERTURBED ELLIPTIC EQUATIONS

SIMONE SECCHI AND MARCO SQUASSINA

ABSTRACT. By means of a variational identity of Pohožaev-Pucci-Serrin type for solutions of class C^1 recently obtained, we give some necessary conditions for locating the concentration points for a class of quasi-linear elliptic problems in divergence form. More precisely we show that the points where the concentration occurs must be critical, either in a generalized or in the classical sense, for a suitable ground state function.

1. INTRODUCTION

Let $\varepsilon > 0$, $n \geq 3$ and $1 < p < n$. In this paper we consider the following class of singularly perturbed quasi-linear elliptic problems in divergence form

$$(P_\varepsilon) \quad \begin{cases} -\varepsilon^p \operatorname{div}(\alpha(x) \partial_\xi \beta(\nabla u)) + V(x) u^{p-1} = K(x) f(u) & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n. \end{cases}$$

We assume that the functions $\alpha, V, K: \mathbb{R}^n \rightarrow \mathbb{R}$ are positive and of class C^1 with $\alpha, K \in L^\infty(\mathbb{R}^n)$ and

$$\inf_{x \in \mathbb{R}^n} \alpha(x) > 0, \quad \inf_{x \in \mathbb{R}^n} V(x) > 0.$$

The function $\beta: \mathbb{R}^n \rightarrow \mathbb{R}$ is strictly convex and positively homogeneous of degree p , i.e. $\beta(\lambda \xi) = \lambda^p \beta(\xi)$ for every $\lambda > 0$ and $\xi \in \mathbb{R}^n$. Moreover, there exist $\nu > 0$ and $c_1, c_2 > 0$ such that

$$(1.1) \quad \nu |\xi|^p \leq \beta(\xi) \leq c_1 |\xi|^p,$$

$$(1.2) \quad |\nabla \beta(\xi)| \leq c_2 |\xi|^{p-1},$$

for every $\xi \in \mathbb{R}^n$. The nonlinearity $f: \mathbb{R}^+ \rightarrow \mathbb{R}$ is of class C^1 and such that

$$(1.3) \quad \lim_{s \rightarrow 0} \frac{f(s)}{s^{p-1}} = 0, \quad \lim_{s \rightarrow \infty} \frac{f(s)}{s^{q-1}} = 0$$

1991 *Mathematics Subject Classification.* 35J40; 58E05.

Key words and phrases. Quasi-linear elliptic equations, concentrating sequences, Pohožaev-Pucci-Serrin identity, Clarke's subdifferential.

The first author was supported by the MIUR national research project "Variational methods and nonlinear differential equations". The second author was supported by the MIUR national research project "Variational and topological methods in the study of nonlinear phenomena" and by the Istituto Nazionale di Alta Matematica "F. Severi" (INdAM).

for some $p < q < p^*$, $p^* = np/(n - p)$, and

$$(1.4) \quad 0 < \vartheta F(s) \leq f(s)s, \quad \text{for every } s > 0$$

for some $\vartheta > p$.

Let us define the space $W_V(\mathbb{R}^n)$ by setting

$$W_V(\mathbb{R}^n) := \left\{ u \in W^{1,p}(\mathbb{R}^n) : \int_{\mathbb{R}^n} V(x)|u|^p dx < \infty \right\},$$

endowed with the natural norm $\|u\|_{W_V}^p = \int_{\mathbb{R}^n} |\nabla u|^p dx + \int_{\mathbb{R}^n} V(x)|u|^p dx$. We write $H_V(\mathbb{R}^n)$ in place of $W_V(\mathbb{R}^n)$ if $p = 2$. Under the previous assumptions, if $K \equiv 1$, it has been recently proved in [8, 19] that if for some compact subset $\Lambda \subset \mathbb{R}^n$ we have

$$(1.5) \quad V(z_0) = \min_{\Lambda} V < \min_{z \in \partial\Lambda} V(z), \quad \alpha(z_0) = \min_{z \in \Lambda} \alpha(z)$$

then for every ε sufficiently small there exists a solution $u_\varepsilon \in W_V(\mathbb{R}^n)$ of (P_ε) which has a maximum point $z_\varepsilon \in \Lambda$, with

$$\lim_{\varepsilon \rightarrow 0} V(z_\varepsilon) = \min_{\Lambda} V, \quad \lim_{\varepsilon \rightarrow 0} \|u_\varepsilon\|_{L^\infty(\Omega \setminus B_\varrho(z_\varepsilon))} = 0 \quad \text{for every } \varrho > 0.$$

In the semi-linear case this kind of problems have been deeply investigated in the last decade and also stronger results can be found in the literature (see e.g. [1, 4, 5, 12, 16, 21]). In this paper, in Theorems 2.7, 2.12 and 3.6, we establish some necessary conditions for a sequence of solutions (u_{ε_h}) of (P_ε) to concentrate around a given point $z_0 \in \mathbb{R}^n$, in the sense of Definition 2.9. As a first condition, the vectors

$$\nabla \alpha(z_0), \quad \nabla V(z_0), \quad \nabla K(z_0)$$

must be linearly dependent.

If $\beta(\xi) = \xi$, we will show in Theorem 3.6 that if z_0 is a concentration point for a sequence $(u_{\varepsilon_h}) \subset H_V(\mathbb{R}^n)$ of solutions of the problem then there exists a locally Lipschitz function $\Sigma : \mathbb{R}^n \rightarrow \mathbb{R}$ (see Definition 2.3), the ground state function, which has, under suitable assumptions, a critical point in the sense of Clarke's subdifferential at z_0 , that is

$$0 \in \partial \Sigma(z_0).$$

Under more stringent assumptions, it turns out that Σ admits all the directional derivatives at z_0 and

$$\nabla \Sigma(z_0) = 0.$$

In the general case, in Theorem 2.7, we show that for a sequence $(u_{\varepsilon_h}) \subset W_V(\mathbb{R}^n)$ of solutions to (P_ε) which concentrates around a point $z_0 \in \mathbb{R}^n$ and which admits an integrable barrier, then z_0 has to be critical in a generalized sense. For instance, in the case when $\beta(\xi) = |\xi|^{p-2}\xi$, we prove in Theorem 2.12 that if $(u_{\varepsilon_h}) \subset W_V(\mathbb{R}^n)$ is a sequence of solutions of the problem

$$\begin{cases} -\varepsilon^p \Delta_p u + V(x)u^{p-1} = K(x)u^{q-1} & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n \end{cases}$$

concentrating at a point z_0 , then

$$z_0 \in \mathfrak{C}(\Sigma),$$

where $\mathfrak{C}(\Sigma)$ is called the set of formal critical points of Σ (see Definition 2.2). To the authors' knowledge, this kind of necessary conditions in terms of the generalized derivatives seems to be new. Quite interestingly, the lack of uniqueness for the limiting problem (that is the problem with frozen coefficients)

$$(P_z) \quad \begin{cases} -\alpha(z) \operatorname{div}(\partial_\xi \beta(\nabla u)) + V(z)u^{p-1} = K(z)f(u) & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n \end{cases}$$

forces a lack of regularity for Σ . Some sharp conditions ensuring uniqueness for (P_z) can be found in [2, 7, 18]. For instance, for $1 < p \leq 2$, $\beta(\xi) = |\xi|^{p-2}\xi$ and $f(u) = u^{q-1}$ with $p < q < p^*$, we have uniqueness for (P_z) and Σ admits all the directional derivatives.

Some necessary conditions for the location of concentration points were previously obtained by Ambrosetti et al. in [1] and by Wang and Zeng in [21, 22] in the case $p = 2$ and $\beta(\xi) = \xi$. Their approach is based on a repeated use of the Divergence Theorem. With respect to those papers we prove our main results by means of a locally Lipschitz variant of the celebrated Pucci-Serrin variational identity ([14]). In our possibly degenerate setting, classical C^2 solutions might not exist, the highest general regularity class being $C^{1,\beta}$ (see [20]). Therefore, the classical identity is not applicable in our framework. However, it has been recently shown in [3] that, under minimal regularity assumptions, the identity holds for locally Lipschitz solutions (see Theorem 2.6), provided that the operator is strictly convex in the gradient, which, from our viewpoint, is a very natural requirement.

This identity has also turned out to be useful in characterizing the exact energy level of the least energy solutions of the problem (P_z) . Indeed in [8, Theorem 3.2] it was proved that (P_z) admits a least energy solution $u_z \in W^{1,p}(\mathbb{R}^n)$ having the Mountain-Pass energy level c_z . This is precisely the motivation that led us to define the ground-energy function Σ also in a degenerate setting.

2. THE QUASI-LINEAR CASE

If z is fixed in \mathbb{R}^n , we consider the limiting functional $I_z : W^{1,p}(\mathbb{R}^n) \rightarrow \mathbb{R}$,

$$(2.1) \quad I_z(u) := \alpha(z) \int_{\mathbb{R}^n} \beta(\nabla u) dx + \frac{V(z)}{p} \int_{\mathbb{R}^n} |u|^p dx - K(z) \int_{\mathbb{R}^n} F(u) dx,$$

where $F(s) = \int_0^s f(t) dt$. It follows from our assumptions on β and f that I_z is a C^1 functional and its critical points are solutions of the limiting problem (P_z) . We define the minimax value c_z for I_z by setting

$$(2.2) \quad c_z := \inf_{\gamma \in \mathcal{P}_z} \sup_{t \in [0,1]} I_z(\gamma(t)),$$

$$\mathcal{P}_z := \left\{ \gamma \in C([0,1], W^{1,p}(\mathbb{R}^n)) : \gamma(0) = 0, I_z(\gamma(1)) < 0 \right\}.$$

We now introduce two functions $\partial\Gamma^-$, $\partial\Gamma^+$ that will be useful in the sequel.

Definition 2.1. For every $z, w \in \mathbb{R}^n$ we define $\partial\Gamma^-(z; w)$ and $\partial\Gamma^+(z; w)$ by setting

$$\begin{aligned}\partial\Gamma^-(z; w) &:= \sup_{v \in G(z)} \nabla_z I_z(v) \cdot w, \\ \partial\Gamma^+(z; w) &:= \inf_{v \in G(z)} \nabla_z I_z(v) \cdot w,\end{aligned}$$

where $G(z)$ is the set of all the nontrivial solutions of the limiting problem (P_z) (up to translations). Explicitly,

$$\begin{aligned}\partial\Gamma^-(z; w) &= \sup_{v \in G(z)} \left[\nabla\alpha(z) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v) dx \right. \\ &\quad \left. + \nabla V(z) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \nabla K(z) \cdot w \int_{\mathbb{R}^n} F(v) dx \right], \\ \partial\Gamma^+(z; w) &= \inf_{v \in G(z)} \left[\nabla\alpha(z) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v) dx \right. \\ &\quad \left. + \nabla V(z) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \nabla K(z) \cdot w \int_{\mathbb{R}^n} F(v) dx \right],\end{aligned}$$

for every $z, w \in \mathbb{R}^n$.

The motivation for introducing the functions $\partial\Gamma^\pm(z; w)$ will be clear later on, in view of formulas (3.2) of Lemma 3.1.

Definition 2.2. We define the set $\mathfrak{C}(\Sigma)$ of formal critical points of Σ by

$$\mathfrak{C}(\Sigma) := \left\{ z \in \mathbb{R}^n : \partial\Gamma^-(z, w) \geq 0 \text{ and } \partial\Gamma^+(z, w) \leq 0 \text{ for every } w \in \mathbb{R}^n \right\}.$$

Definition 2.3. We define the ground-energy function $\Sigma : \mathbb{R}^n \rightarrow \mathbb{R}$ by setting

$$(2.3) \quad \Sigma(z) := \min\{I_z(u) : u \in W^{1,p}(\mathbb{R}^n) \text{ is a nontrivial solution of } (P_z)\}$$

for every $z \in \mathbb{R}^n$.

Lemma 2.4. Assume that

$$(2.4) \quad s \in \mathbb{R}^+ \mapsto \frac{f(s)}{s^{p-1}} \text{ is increasing.}$$

Then the following facts hold.

(i) the map Σ is well defined and continuous and

$$\Sigma(z) = c_z, \quad \text{for all } z \in \mathbb{R}^n;$$

(ii) the map Σ can be written as

$$\Sigma(z) = \inf_{u \in W^{1,p}(\mathbb{R}^n) \setminus \{0\}} \max_{\theta \geq 0} I_z(\theta u) = \inf_{u \in \mathcal{N}_z} I_z(u), \quad \text{for all } z \in \mathbb{R}^n,$$

where the Nehari manifold \mathcal{N}_z is defined as

$$\mathcal{N}_z := \left\{ u \in W^{1,p}(\mathbb{R}^n) \setminus \{0\} : I'_z(u)[u] = 0 \right\}.$$

Proof. By [8, Theorem 3.2], for every $z \in \mathbb{R}^n$ the problem (P_z) admits a solution $v_z \in W^{1,p}(\mathbb{R}^n)$ such that

$$I_z(v_z) = \inf \{ I_z(u) : u \in W^{1,p}(\mathbb{R}^n) \text{ is a nontrivial solution of } (P_z) \}.$$

Moreover, $I_z(v_z) = c_z$, where c_z is as in (2.2). The continuity of Σ then follows from the continuity of the map $z \mapsto c_z$, which we now prove directly using an argument envisaged by Rabinowitz in [16].

For $\alpha, V, K \in \mathbb{R}$, define the functional $I_{\alpha,V,K} : W^{1,p}(\mathbb{R}^n) \rightarrow \mathbb{R}$ by

$$I_{\alpha,V,K}(u) := \alpha \int_{\mathbb{R}^n} \beta(\nabla u) dx + \frac{V}{p} \int_{\mathbb{R}^n} |u|^p dx - K \int_{\mathbb{R}^n} F(u) dx.$$

Let us set

$$c(\alpha, V, K) := \inf_{\gamma \in \mathcal{P}_{\alpha,V,K}} \max_{t \in [0,1]} I_{\alpha,V,K}(\gamma(t)),$$

$$\mathcal{P}_{\alpha,V,K} := \left\{ \gamma \in C([0,1], W^{1,p}(\mathbb{R}^n)) : \gamma(0) = 0, \quad I_{\alpha,V,K}(\gamma(1)) < 0 \right\}.$$

Claim: For every $(\alpha, V, K) \in \mathbb{R}^3$ we have

$$(2.5) \quad \lim_{\eta \rightarrow 0} c(\alpha + \eta, V + \eta, K - \eta) = c(\alpha, V, K).$$

We first observe that a simple adaptation of [16, Lemma 3.17] yields

$$(2.6) \quad \alpha_1 > \alpha_2, \quad V_1 > V_2, \quad K_1 < K_2 \implies c(\alpha_1, V_1, K_1) \geq c(\alpha_2, V_2, K_2).$$

The proof of (2.5) will be accomplished indirectly. By virtue of (2.6), we get

$$\lim_{\eta \rightarrow 0^-} c(\alpha + \eta, V + \eta, K - \eta) := c^- \leq c(\alpha, V, K).$$

Suppose that $c^- < c(\alpha, V, K)$. For the sake of brevity, we define

$$J_\eta(u) := I_{\alpha+\eta, V+\eta, K-\eta}(u).$$

Let $\eta_h \rightarrow 0^-$ as $h \rightarrow \infty$, and $\delta_j \rightarrow 0^+$ as $j \rightarrow \infty$. For each $h \in \mathbb{N}$, by Lemma 2.4, there is a sequence (u_{hj}) in $W^{1,p}(\mathbb{R}^n) \setminus \{0\}$ such that

$$(2.7) \quad \alpha \int_{\mathbb{R}^n} \beta(\nabla u_{hj}) dx + V \int_{\mathbb{R}^n} |u_{hj}|^p dx = 1$$

and

$$(2.8) \quad \max_{\theta \geq 0} J_{\eta_h}(\theta u_{hj}) \leq c(\alpha + \eta_h, V + \eta_h, K - \eta_h) + \delta_j.$$

We can choose (u_{hj}) satisfying (2.7), since the position $u \mapsto \alpha \int_{\mathbb{R}^n} \beta(\nabla u) + V \int_{\mathbb{R}^n} |u|^p$ defines on $W^{1,p}(\mathbb{R}^n)$ a norm equivalent to the natural one, as follows from (1.1). Take

now $h = j$, set $u_h = u_{hh}$. Hence

$$\begin{aligned}
c(\alpha, V, K) &\leq \max_{\theta \geq 0} I_{\alpha, V, K}(\theta u_h) = I_{\alpha, V, K}(\phi(u_h)u_h) \\
&= J_{\eta_h}(\phi(u_h)u_h) - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \frac{|u_h|^p}{p} dx - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \beta(\nabla u_h) dx \\
&\quad + \eta_h \int_{\mathbb{R}^n} F(\phi(u_h)u_h) dx \\
&\leq \max_{\theta \geq 0} J_{\eta_h}(\theta u_h) - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \frac{|u_h|^p}{p} dx - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \beta(\nabla u_h) dx \\
&\quad + \eta_h \int_{\mathbb{R}^n} F(\phi(u_h)u_h) dx \\
&\leq c(\alpha + \eta_h, V + \eta_h, K - \eta_h) + \delta_h - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \frac{|u_h|^p}{p} dx \\
&\quad - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \beta(\nabla u_h) dx + \eta_h \int_{\mathbb{R}^n} F(\phi(u_h)u_h) dx \\
&\leq c^- + \delta_h - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \frac{|u_h|^p}{p} dx - \eta_h \phi(u_h)^p \int_{\mathbb{R}^n} \beta(\nabla u_h) dx \\
&\quad + \eta_h \int_{\mathbb{R}^n} F(\phi(u_h)u_h) dx.
\end{aligned}$$

Since $\|u_h\| = 1$, the sequence (u_h) is bounded in $W_V(\mathbb{R}^n)$. Recalling the properties of F and the Sobolev embeddings, if the sequence $(\phi(u_h))$ is bounded with respect to h , then the last chain of inequalities contradicts $c^- < c(\alpha, V, K)$ at least for h sufficiently large.

At this point, one can show exactly as in [16, pp. 281–282] that there exists a constant $C > 0$ such that

$$\phi(u_h) \leq C \quad \text{for all } h \text{ sufficiently large.}$$

We conclude that $c^- < c(\alpha, V, K)$ is impossible. In a completely similar fashion one can prove that the inequality

$$c(\alpha, V, K) < \lim_{\eta \rightarrow 0^+} c(\alpha + \eta, V + \eta, K - \eta)$$

leads to a contradiction. Therefore (2.5) holds true and the proof of the claim is complete.

Let now (z_h) be a sequence in \mathbb{R}^n such that $z_h \rightarrow z$ as $h \rightarrow \infty$. Observe that, given $\eta > 0$, for large $h \in \mathbb{N}$ we have

$$\begin{aligned}
V(z) + \eta &\geq V(z) + |V(z_h) - V(z)| \\
&\geq V(z) \geq V(z) - |V(z_h) - V(z)| \\
&\geq V(z) - \eta
\end{aligned}$$

and similar relations for α and K . Therefore the continuity of $z \mapsto c_z$ follows from the previous claim, applied with $\alpha = \alpha(z)$, $V = V(z)$, $K = K(z)$. This completes the proof of (i).

To prove (ii), it suffices to argue as in [13, Proposition 2.5]. \square

Remark 2.5. We point out that the regularity of Σ for $p \neq 2$ is, in general, a hard matter owing to the lack of regularity of the solutions of the limiting problem. If instead $\beta(\xi) = \xi$ the map Σ is locally Lipschitz, as shown in Lemma 3.1.

Let $\mathcal{L} : \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a function of class C^1 such that the function

$$\xi \mapsto \mathcal{L}(x, s, \xi)$$

is strictly convex for every $(x, s) \in \mathbb{R}^n \times \mathbb{R}$, and let $\varphi \in L_{\text{loc}}^\infty(\mathbb{R}^n)$.

We now recall a Pucci-Serrin variational identity for locally Lipschitz continuous solutions of a general class of Euler equations, proved in [3]. As we have already pointed out in the introduction, the classical identity [14] is not applicable here, since it requires the C^2 regularity of the solutions while the maximal regularity for degenerate equations is $C^{1,\beta}$ (see e.g. [20]).

Theorem 2.6. *Let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally Lipschitz solution of*

$$-\operatorname{div}(\partial_\xi \mathcal{L}(x, u, \nabla u)) + \partial_s \mathcal{L}(x, u, \nabla u) = \varphi \quad \text{in } \mathcal{D}'(\mathbb{R}^n).$$

Then

$$(2.9) \quad \sum_{i,j=1}^n \int_{\mathbb{R}^n} \partial_i h^j \partial_{\xi_i} \mathcal{L}(x, u, \nabla u) \partial_j u + \\ - \int_{\mathbb{R}^n} [(\operatorname{div} h) \mathcal{L}(x, u, \nabla u) + h \cdot \partial_x \mathcal{L}(x, u, \nabla u)] = \int_{\mathbb{R}^n} (h \cdot \nabla u) \varphi$$

for every $h \in C_c^1(\mathbb{R}^n, \mathbb{R}^n)$.

The following is the main result of this section.

Theorem 2.7. *Let $z_0 \in \mathbb{R}^n$ and assume that (u_{ε_h}) is a sequence of solutions of problem (P_ε) such that*

$$(2.10) \quad u_{\varepsilon_h} = v_0 \left(\frac{\cdot - z_0}{\varepsilon_h} \right) + o(1), \quad \text{strongly in } W_V(\mathbb{R}^n)$$

for some $v_0 \in W_V(\mathbb{R}^n) \setminus \{0\}$. Then the following facts hold:

(a) *the vectors*

$$\nabla \alpha(z_0), \quad \nabla V(z_0), \quad \nabla K(z_0)$$

are linearly dependent;

(b) *$z_0 \in \mathfrak{C}(\Sigma)$, that is z_0 is a formal critical point of Σ ;*

(c) *if $G(z_0) = \{v_0\}$, then all the partial derivatives of Σ at z_0 exist and*

$$\nabla \Sigma(z_0) = 0,$$

that is z_0 is a critical point of Σ .

Proof. We write u_h instead of u_{ε_h} and we define

$$(2.11) \quad v_h(x) := u_h(z_0 + \varepsilon_h x).$$

Therefore v_h satisfies the rescaled equation

$$-\operatorname{div}(\alpha(z_0 + \varepsilon_h x) \partial_\xi \beta(\nabla v_h)) + V(z_0 + \varepsilon_h x) v_h^{p-1} = K(z_0 + \varepsilon_h x) f(v_h) \quad \text{in } \mathbb{R}^n.$$

By (2.10), we have $v_h \rightarrow v_0$ strongly in $W_V(\mathbb{R}^n)$. Clearly (v_h) is bounded in $W_V(\mathbb{R}^n)$. We now prove that $v_h \rightarrow v_0$ in the C^1 sense over the compact sets of \mathbb{R}^n and that v_0 is a nontrivial positive solution of the equation

$$(2.12) \quad -\alpha(z_0) \operatorname{div}(\partial_\xi \beta(\nabla v)) + V(z_0) v^{p-1} = K(z_0) f(v) \quad \text{in } \mathbb{R}^n.$$

Let us set

$$\begin{aligned} d_h(x) &:= \begin{cases} V(z_0 + \varepsilon_h x) - K(z_0 + \varepsilon_h x) \frac{f(v_h(x))}{v_h^{p-1}(x)} & \text{if } v_h(x) \neq 0 \\ 0 & \text{if } v_h(x) = 0, \end{cases} \\ A(x, s, \xi) &:= \alpha(z_0 + \varepsilon_h x) \nabla \beta(\xi), \\ B(x, s, \xi) &:= d_h(x) s^{p-1}, \end{aligned}$$

for every $x \in \mathbb{R}^n$, $s \in \mathbb{R}^+$ and $\xi \in \mathbb{R}^n$. Taking into account the growth condition on $\nabla \beta$, the strict convexity of β in ξ , we get

$$A(x, s, \xi) \cdot \xi \geq \nu |\xi|^p, \quad |A(x, s, \xi)| \leq c_2 |\xi|^{p-1}.$$

Now, there exists $\delta > 0$ sufficiently small such that $d_h \in L^{n/(p-\delta)}(B_{2\varrho})$ for every $\varrho > 0$ and

$$S = \sup_h \|d_h\|_{L^{n/(p-\delta)}(B_{2\varrho})} \leq D_\varrho \left(1 + \sup_{h \in \mathbb{N}} \|v_h\|_{L^{p^*}(B_{2\varrho})} \right) < \infty$$

for some $D_\varrho > 0$. Since we have $\operatorname{div}(A(x, v_h, \nabla v_h)) = B(x, v_h, \nabla v_h)$ for every $h \in \mathbb{N}$, by virtue of [17, Theorem 1] there exists a radius $\varrho > 0$ and a positive constant $M = M(\nu, c_2, S\varrho^\delta)$ such that

$$\sup_{h \in \mathbb{N}} \max_{x \in B_\varrho} |v_h(x)| \leq M(2\varrho)^{-N/p} \sup_{h \in \mathbb{N}} \|v_h\|_{L^p(B_{2\varrho})} < \infty$$

so that (v_h) is uniformly bounded in B_ϱ . Then, by [17, Theorem 8], up to a subsequence (v_h) converges uniformly to v_0 in a small neighborhood of zero. Similarly one shows that $v_h \rightarrow v_0$ in $C_{\text{loc}}^1(\mathbb{R}^n)$. Therefore, it is easily seen that v_0 is a nontrivial positive solution of (2.12). Since the map β is strictly convex, we can use Theorem 2.6 by choosing in (2.9) $\varphi = 0$ and

$$\begin{aligned} \mathcal{L}(s, \xi) &:= \alpha(z_0 + \varepsilon_h x) \beta(\xi) + V(z_0 + \varepsilon_h x) \frac{s^p}{p} - K(z_0 + \varepsilon_h x) F(s), \\ h(x) = h_{\varepsilon, k}(x) &:= (\underbrace{0, \dots, 0}_{k-1}, T(\varepsilon x), \underbrace{0, \dots, 0}_{n-k}), \end{aligned}$$

for every $x \in \mathbb{R}^n$, $s \in \mathbb{R}^+$ and $\xi \in \mathbb{R}^n$, the function $T \in C_c^1(\mathbb{R}^n)$ being chosen so that $T(x) = 1$ if $|x| \leq 1$ and $T(x) = 0$ if $|x| \geq 2$. In particular, $h_{\varepsilon, k} \in C_c^1(\mathbb{R}^n, \mathbb{R}^n)$ and

$$\partial_i h_{\varepsilon, k}^j(x) = \varepsilon \partial_i T(\varepsilon x) \delta_{kj} \quad \text{for every } x \in \mathbb{R}^n.$$

Then, it follows from (2.9) that

$$\begin{aligned}
0 &= \sum_{i=1}^n \int_{\mathbb{R}^n} \varepsilon \partial_i T(\varepsilon x) \alpha(z_0 + \varepsilon_h x) \partial_{\xi_i} \beta(\nabla v_h) \partial_k v_h \, dx \\
&\quad - \int_{\mathbb{R}^n} \varepsilon \partial_k T(\varepsilon x) \left[\alpha(z_0 + \varepsilon_h x) \beta(\nabla v_h) + V(z_0 + \varepsilon_h x) \frac{v_h^p}{p} - K(z_0 + \varepsilon_h x) F(v_h) \right] dx \\
&\quad - \int_{\mathbb{R}^n} \varepsilon_h T(\varepsilon x) \left[\frac{\partial \alpha}{\partial x_k}(z_0 + \varepsilon_h x) \beta(\nabla v_h) + \frac{\partial V}{\partial x_k}(z_0 + \varepsilon_h x) \frac{v_h^p}{p} - \frac{\partial K}{\partial x_k}(z_0 + \varepsilon_h x) F(v_h) \right] dx
\end{aligned}$$

for every $k = 1, \dots, n$. Since the sequence (v_h) is bounded in $W_V(\mathbb{R}^n)$, by (1.1), (1.2) and the boundedness of α and K , we have

$$\begin{aligned}
&\left| \sum_{i=1}^n \int_{\mathbb{R}^n} \partial_i T(\varepsilon x) \alpha(z_0 + \varepsilon_h x) \partial_{\xi_i} \beta(\nabla v_h) \partial_k v_h \, dx \right| \leq C, \\
&\left| \int_{\mathbb{R}^n} \partial_k T(\varepsilon x) \left[\alpha(z_0 + \varepsilon_h x) \beta(\nabla v_h) + V(z_0 + \varepsilon_h x) \frac{v_h^p}{p} - K(z_0 + \varepsilon_h x) F(v_h) \right] dx \right| \leq C'
\end{aligned}$$

for some positive constants C, C' . Therefore, letting $\varepsilon \rightarrow 0$ yields

$$\begin{aligned}
(2.13) \quad &\int_{\mathbb{R}^n} \left[\frac{\partial \alpha}{\partial x_k}(z_0 + \varepsilon_h x) \beta(\nabla v_h) + \frac{\partial V}{\partial x_k}(z_0 + \varepsilon_h x) \frac{v_h^p}{p} + \right. \\
&\quad \left. - \frac{\partial K}{\partial x_k}(z_0 + \varepsilon_h x) F(v_h) \right] dx = 0
\end{aligned}$$

for every $h \in \mathbb{N}$. Letting $h \rightarrow \infty$, by (2.10), we find

$$\frac{\partial \alpha}{\partial x_k}(z_0) \int_{\mathbb{R}^n} \beta(\nabla v_0) \, dx + \frac{\partial V}{\partial x_k}(z_0) \int_{\mathbb{R}^n} \frac{v_0^p}{p} \, dx - \frac{\partial K}{\partial x_k}(z_0) \int_{\mathbb{R}^n} F(v_0) \, dx = 0$$

for every $k = 1, \dots, n$, which yields

$$\nabla \alpha(z_0) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v_0) \, dx + \nabla V(z_0) \cdot w \int_{\mathbb{R}^n} \frac{v_0^p}{p} \, dx = \nabla K(z_0) \cdot w \int_{\mathbb{R}^n} F(v_0) \, dx$$

for every $w \in \mathbb{R}^n$. Then, since $v_0 \not\equiv 0$, assertion (a) follows. Moreover, since $v_0 \in G(z_0)$, by the definition of $\partial \Gamma^-$ we obtain

$$\begin{aligned}
\partial \Gamma^-(z_0; w) &= \sup_{v \in G(z_0)} \left[\nabla \alpha(z_0) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v) \, dx \right. \\
&\quad \left. + \nabla V(z_0) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} \, dx - \nabla K(z_0) \cdot w \int_{\mathbb{R}^n} F(v) \, dx \right] \\
&\geq \nabla \alpha(z_0) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v_0) \, dx \\
&\quad + \nabla V(z_0) \cdot w \int_{\mathbb{R}^n} \frac{v_0^p}{p} \, dx - \nabla K(z_0) \cdot w \int_{\mathbb{R}^n} F(v_0) \, dx = 0
\end{aligned}$$

for every $w \in \mathbb{R}^n$. Analogously, by the definition of $\partial\Gamma^+$, we have

$$\begin{aligned} \partial\Gamma^+(z_0; w) &= \inf_{v \in G(z_0)} \left[\nabla\alpha(z_0) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v) dx \right. \\ &\quad \left. + \nabla V(z_0) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \nabla K(z_0) \cdot w \int_{\mathbb{R}^n} F(v) dx \right] \\ &\leq \nabla\alpha(z_0) \cdot w \int_{\mathbb{R}^n} \beta(\nabla v_0) dx \\ &\quad + \nabla V(z_0) \cdot w \int_{\mathbb{R}^n} \frac{v_0^p}{p} dx - \nabla K(z_0) \cdot w \int_{\mathbb{R}^n} F(v_0) dx = 0 \end{aligned}$$

for every $w \in \mathbb{R}^n$. Therefore $z_0 \in \mathfrak{C}(\Sigma)$ and assertion (b) is proved. If $G(z_0) = \{v_0\}$, then clearly Σ admits all the directional derivatives at z_0 and

$$\frac{\partial\Sigma}{\partial w}(z_0) = \partial\Gamma^-(z_0; w) = \partial\Gamma^+(z_0; w) = 0 \quad \text{for every } w \in \mathbb{R}^n,$$

by virtue of (b). This proves assertion (c). \square

The strong convergence required by (2.10) allows us to take the limit as $h \rightarrow \infty$ in equation (2.13). In the semi-linear situation one can construct uniform exponential barriers for the family (v_h) , and therefore the strong convergence of (v_h) follows easily from the Lebesgue Convergence Theorem (see [13, 21, 22]). The well-known loss of regularity for solutions of quasi-linear equations is usually an obstruction to this kind of argument. However, if the solutions belong to a suitable Sobolev space, then a pointwise concentration suffices, see Corollary 2.10.

Remark 2.8. We wish to point out that Theorem 2.7 holds true also for the more general class of quasi-linear equations

$$-\varepsilon^p \operatorname{div}(\alpha(x) \partial_\xi \beta(u, \nabla u)) + \varepsilon^p \alpha(x) \partial_s \beta(u, \nabla u) + V(x) u^{p-1} = K(x) f(u),$$

under suitable assumptions on the terms $\partial_\xi \beta(s, \xi)$ and $\partial_s \beta(s, \xi)$. On the other hand, although the ground-energy function Σ can be defined exactly as in Definition 2.3 and $\Sigma(z) = c_z$ (cf. [8, Theorem 3.2]), the presence of u itself in the function β makes the problems of the regularity of Σ and of the decay at infinity for the rescaled family of solutions very complicated, even in the nondegenerate case $p = 2$.

Definition 2.9. Let $z_0 \in \mathbb{R}^n$. We say that a sequence (u_{ε_h}) of solutions of (P_{ε_h}) concentrates at z_0 if $u_{\varepsilon_h}(z_0) \geq \ell > 0$ and for every $\eta > 0$ there exist $\varrho > 0$ and $h_0 \in \mathbb{N}$ such that

$$u_{\varepsilon_h}(x) \leq \eta \quad \text{for every } h \geq h_0 \text{ and } |x - z_0| \geq \varepsilon_h \varrho.$$

This is exactly the same notion of concentration adopted in [21, 22].

Corollary 2.10. Let (u_{ε_h}) be a family of solutions of (P_ε) which concentrates at a point $z_0 \in \mathbb{R}^n$. Suppose that, for every h sufficiently large,

$$u_{\varepsilon_h} \in C_d^1(\mathbb{R}^n) \cap W^{2,n}(\mathbb{R}^n),$$

where

$$C_d^1(\mathbb{R}^n) := \left\{ u \in C^1(\mathbb{R}^n) : \lim_{|x| \rightarrow \infty} u(x) = 0, \lim_{|x| \rightarrow \infty} \nabla u(x) = 0 \right\}.$$

Then all the conclusions of Theorem 2.7 hold true.

Proof. If $u_{\varepsilon_h} \in C_d^1(\mathbb{R}^n) \cap W^{2,n}(\mathbb{R}^n)$, then one can apply the results contained in [15] to show that u_{ε_h} decays exponentially fast at infinity, uniformly with respect to h , together with all its partial derivatives. Hence one can pass to the limit in (2.13), and complete the proof as in Theorem 2.7. \square

For the particular but important case $\alpha(x) = 1$, $\beta(\xi) = |\xi|^{p-2}\xi$ and $f(s) = s^{q-1}$, $p < q < p^*$, we can still prove a fast decay at infinity for the solutions.

Lemma 2.11. *Let (u_{ε_h}) be a sequence of solutions of the problem*

$$\begin{cases} -\varepsilon^p \Delta_p u + V(x)u^{p-1} = K(x)u^{q-1} & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n \end{cases}$$

which concentrates at $z_0 \in \mathbb{R}^n$. Then, if we set

$$v_h(x) := u_{\varepsilon_h}(z_0 + \varepsilon_h x),$$

for each $\eta > 0$ there exist $R_\eta, C_\eta > 0$ independent of h such that

$$|v_h(x)| \leq C_\eta \exp \left\{ - \left(\frac{\eta}{p-1} \right)^{1/p} |x| \right\},$$

for every $|x| \geq R_\eta$ and every $h \in \mathbb{N}$.

Proof. For every $h \in \mathbb{N}$, v_h satisfies the equation

$$-\Delta_p v_h + V(z_0 + \varepsilon_h x)v_h^{p-1} = K(z_0 + \varepsilon_h x)v_h^{q-1} \quad \text{in } \mathbb{R}^n.$$

Since u_{ε_h} is a concentrating sequence, it results that

$$\lim_{|x| \rightarrow \infty} v_h(x) = 0, \quad \text{uniformly in } h \in \mathbb{N}.$$

Set $\inf_{x \in \mathbb{R}^n} V(x) = V_0$. Then, given $\eta > 0$ there exists $R_\eta > 0$ independent of h such that

$$V(z_0 + \varepsilon_h x)v_h(x)^{p-1} - K(z_0 + \varepsilon_h x)v_h(x)^{q-1} \geq (V_0 - \eta)v_h(x)^{p-1},$$

for every $|x| \geq R_\eta$. It follows that the inequality

$$(2.14) \quad -\operatorname{div}(|\nabla v_h|^{p-2} \nabla v_h) + (V_0 - \eta)v_h^{p-1} \leq 0$$

holds true for every $h \in \mathbb{N}$ and $|x| \geq R_\eta$. Define now the function

$$\Phi(x) := C_\eta \exp \left\{ - \left(\frac{V_0 - \eta}{p-1} \right)^{1/p} |x| \right\},$$

where

$$C_\eta := \exp \left\{ \left(\frac{V_0 - \eta}{p-1} \right)^{1/p} R_\eta \right\} \max_{|x|=R_\eta} v_h(x).$$

Notice that, since v_h is uniformly bounded, we can assume that C_η is independent of h . Now, exactly the same computations as in [10, Theorem 2.8] yield

$$(2.15) \quad -\operatorname{div}(|\nabla\Phi|^{p-2}\nabla\Phi) + (V_0 - \eta)\Phi^{p-1} \geq 0.$$

Testing inequalities (2.14) and (2.15) with $\phi = (v_h - \Phi)^+$ on $\{|x| \geq R_\eta\}$ yields

$$\begin{aligned} \int_{\{|x| \geq R_\eta\} \cap \{v_h \geq \Phi\}} (|\nabla v_h|^{p-2}\nabla v_h \cdot \nabla(v_h - \Phi) + (V_0 - \eta)v_h^{p-1}(v_h - \Phi)) \, dx &\leq 0, \\ \int_{\{|x| \geq R_\eta\} \cap \{v_h \geq \Phi\}} (|\nabla\Phi|^{p-2}\nabla\Phi \cdot \nabla(v_h - \Phi) + (V_0 - \eta)\Phi^{p-1}(v_h - \Phi)) \, dx &\geq 0. \end{aligned}$$

By subtracting the previous inequality and taking into account that

$$\sum_{i=1}^n (|\xi|^{p-2}\xi_i - |\zeta|^{p-2}\zeta_i)(\xi_i - \zeta_i) > 0, \quad \text{for every } \xi \neq \zeta,$$

we get

$$\int_{\{|x| \geq R_\eta\} \cap \{v_h \geq \Phi\}} (v_h^{p-1} - \Phi^{p-1})(v_h - \Phi) \, dx \leq 0.$$

Since v_h and Φ are continuous functions, it has to be

$$\{|x| \geq R_\eta\} \cap \{v_h \geq \Phi\} = \emptyset,$$

which implies the assertion. \square

Theorem 2.12. *Let (u_{ε_h}) be a sequence of solutions of the problem*

$$(2.16) \quad \begin{cases} -\varepsilon^p \Delta_p u + V(x)u^{p-1} = K(x)u^{q-1} & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n \end{cases}$$

which concentrates at $z_0 \in \mathbb{R}^n$. Then the following facts hold:

- (a) *the vectors $\nabla V(z_0)$ and $\nabla K(z_0)$ are linearly dependent;*
- (b) *$z_0 \in \mathfrak{C}(\Sigma)$, that is z_0 is a formal critical point of Σ ;*
- (c) *if $1 < p \leq 2$ then all the partial derivatives of Σ at z_0 exist and $\nabla \Sigma(z_0) = 0$, that is z_0 is a critical point of Σ .*

Proof. By virtue of Lemma 2.11 we can pass to the limit in equation (2.13) and get assertions (a) and (b) as in Theorem 2.7. If $1 < p \leq 2$, by combining the results of [2, 7] and [18], for every $z \in \mathbb{R}^n$, problem (P_z) admits a unique positive C^1 solution (up to translations) with $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then $G(z_0) = \{v_0\}$ and assertion (c) follows by the corresponding assertion in Theorem 2.7. \square

3. THE SEMI-LINEAR CASE

The main goal of this section is that of getting, in the particular case $p = 2$ and $\beta(\xi) = \xi$, i.e. semi-linear equations, a more accurate version of Theorem 2.7. For any $z \in \mathbb{R}^n$ we consider the limiting functional $I_z : W^{1,p}(\mathbb{R}^n) \rightarrow \mathbb{R}$,

$$(3.1) \quad I_z(u) := \alpha(z) \int_{\mathbb{R}^n} |\nabla u|^2 dx + \frac{V(z)}{p} \int_{\mathbb{R}^n} |u|^p dx - K(z) \int_{\mathbb{R}^n} F(u) dx.$$

Clearly, its critical points are solutions on (P_z) . The minimax levels c_z of I_z are defined according to (2.2).

As we see in the following lemma, in this situation, the map Σ introduced in Definition 2.3 has some further regularity properties.

Lemma 3.1. *If condition (2.4) with $p = 2$ holds, then the following facts are true:*

- (i) *the map Σ is locally Lipschitz;*
- (ii) *the directional derivatives from the left and the right of Σ at z along w , which we denote respectively by $(\frac{\partial \Sigma}{\partial w})^-(z)$ and $(\frac{\partial \Sigma}{\partial w})^+(z)$, exist at every point $z \in \mathbb{R}^n$ and it holds*

$$\begin{aligned} \left(\frac{\partial \Sigma}{\partial w}\right)^-(z) &= \sup_{v \in S(z)} \left[\nabla \alpha(z) \cdot w \int_{\mathbb{R}^n} \frac{|\nabla v|^2}{2} dx \right. \\ &\quad \left. + \nabla V(z) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \nabla K(z) \cdot w \int_{\mathbb{R}^n} F(v) dx \right], \\ \left(\frac{\partial \Sigma}{\partial w}\right)^+(z) &= \inf_{v \in S(z)} \left[\nabla \alpha(z) \cdot w \int_{\mathbb{R}^n} \frac{|\nabla v|^2}{2} dx \right. \\ &\quad \left. + \nabla V(z) \cdot w \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \nabla K(z) \cdot w \int_{\mathbb{R}^n} F(v) dx \right], \end{aligned}$$

for every $z, w \in \mathbb{R}^n$, where $S(z)$ is the set of all the nontrivial solutions of (P_z) corresponding to the energy level $\Sigma(z)$ (up to translations).

In particular, if $G(z) = S(z)$, we have

$$(3.2) \quad \partial \Gamma^-(z; w) = \left(\frac{\partial \Sigma}{\partial w}\right)^-(z), \quad \partial \Gamma^+(z; w) = \left(\frac{\partial \Sigma}{\partial w}\right)^+(z),$$

for every $w \in \mathbb{R}^n$.

Proof. By the results of [22], Σ is a locally Lipschitz map. We remark here that, since z acts as a parameter, the functional I_z is invariant under orthogonal change of variables. Therefore, without loss of generality, to prove the formulas for the left

and right directional derivatives of Σ it suffices to show that

$$\begin{aligned} \left(\frac{\partial \Sigma}{\partial z_i} \right)^- (z) &= \sup_{v \in S(z)} \left[\frac{\partial \alpha}{\partial z_i}(z) \int_{\mathbb{R}^n} \frac{|\nabla v|^2}{2} dx \right. \\ &\quad \left. + \frac{\partial V}{\partial z_i}(z) \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \frac{\partial K}{\partial z_i}(z) \int_{\mathbb{R}^n} F(v) dx \right], \\ \left(\frac{\partial \Sigma}{\partial z_i} \right)^+ (z) &= \inf_{v \in S(z)} \left[\frac{\partial \alpha}{\partial z_i}(z) \int_{\mathbb{R}^n} \frac{|\nabla v|^2}{2} dx \right. \\ &\quad \left. + \frac{\partial V}{\partial z_i}(z) \int_{\mathbb{R}^n} \frac{|v|^p}{p} dx - \frac{\partial K}{\partial z_i}(z) \int_{\mathbb{R}^n} F(v) dx \right], \end{aligned}$$

for every $i = 1, \dots, n$. This again can be found in [13, 22]. Finally, formulas (3.2) follow by the definition of $\partial \Gamma^\pm(z; w)$. \square

Remark 3.2. Assume that $p = 2$ and $f(u) = u^{q-1}$, where $2 < q < 2^*$. Then Σ can be given an explicit form: there exists a constant $C > 0$ such that

$$\Sigma(z) = CV(z)^{\frac{q}{q-2} - \frac{n}{2}} \sqrt{\alpha(z)},$$

for every $z \in \mathbb{R}^n$. See also [13].

Let us now recall from [6] two definitions of non-smooth analysis that will be useful in the sequel.

Definition 3.3. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally Lipschitz function near a given point $z \in \mathbb{R}^n$. The generalized derivative of the function f at z along the direction $w \in \mathbb{R}^n$ is defined by

$$f^0(z; w) := \limsup_{\substack{\xi \rightarrow z \\ \lambda \rightarrow 0+}} \frac{f(\xi + \lambda w) - f(\xi)}{\lambda}.$$

Definition 3.4. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a locally Lipschitz function near a given point $z \in \mathbb{R}^n$. The Clarke subdifferential (or generalized gradient) of f at z is defined by

$$\partial f(z) := \left\{ \eta \in \mathbb{R}^n : f^0(z, w) \geq \eta \cdot w \text{ for every } w \in \mathbb{R}^n \right\}.$$

Here \cdot stands for the usual inner product of \mathbb{R}^n .

Proposition 3.5. For every $z \in \mathbb{R}^n$ the set $\partial f(z)$ is nonempty, convex and

$$\partial(-f)(z) = -\partial f(z).$$

Proof. See [6, Proposition 2.3.1]. \square

The following is the main result of this section.

Theorem 3.6. Assume that (u_{ε_h}) is a sequence of solutions of the problem

$$(3.3) \quad \begin{cases} -\varepsilon^2 \operatorname{div}(\alpha(x) \nabla u) + V(x)u = K(x)f(u) & \text{in } \mathbb{R}^n \\ u > 0 & \text{in } \mathbb{R}^n, \end{cases}$$

which concentrates at z_0 . Then the following facts hold:

(a) *the vectors*

$$\nabla\alpha(z_0), \nabla V(z_0), \nabla K(z_0)$$

are linearly dependent;

(b) $z_0 \in \mathfrak{C}(\Sigma)$, *that is z_0 is a formal critical point of Σ ;*

(c) *if either $G(z_0) = S(z_0)$ or*

$$\varepsilon^{-n} J_{\varepsilon_h}(u_{\varepsilon_h}) \rightarrow c_{z_0},$$

where

$$(3.4) \quad J_\varepsilon(v) = \frac{\varepsilon^2}{2} \int_{\mathbb{R}^n} \alpha(x) |\nabla v|^2 dx + \frac{1}{2} \int_{\mathbb{R}^n} V(x) |v|^2 dx - \int_{\mathbb{R}^n} K(x) F(v) dx,$$

we have

$$0 \in \partial\Sigma(z_0),$$

that is z_0 is a critical point of Σ in the sense of Clarke's subdifferential;

(d) *if $S(z_0) = \{v_0\}$, then all the partial derivatives of Σ at z_0 exist and*

$$\nabla\Sigma(z_0) = 0,$$

whence z_0 is a critical point of Σ .

Proof. For problem (3.3) it is possible to prove the existence of uniform exponentially decaying barriers. Then we can pass to the limit in equation (2.13), to get assertions (a) and (b) as in Theorem 2.7. If $S(z_0) = G(z_0)$, by combining formulas (3.2) of Lemma 3.1 with (b) of Theorem 2.7, we have

$$(3.5) \quad \left(\frac{\partial\Sigma}{\partial w} \right)^- (z_0) \geq 0, \quad \left(\frac{\partial\Sigma}{\partial w} \right)^+ (z_0) \leq 0$$

for every $w \in \mathbb{R}^n$. In particular, it holds

$$\left(\frac{\partial(-\Sigma)}{\partial w} \right)^+ (z_0) \geq 0 \quad \text{for every } w \in \mathbb{R}^n.$$

Then, by the definition of $(-\Sigma)^0(z_0; w)$ we get

$$(-\Sigma)^0(z_0; w) \geq \left(\frac{\partial(-\Sigma)}{\partial w} \right)^+ (z_0) \geq 0 \quad \text{for every } w \in \mathbb{R}^n.$$

By the definition of $\partial(-\Sigma)(z_0)$ we get

$$0 \in \partial(-\Sigma)(z_0),$$

which, together with Proposition 3.5, yields assertion (c). To prove the same conclusion when

$$\varepsilon^{-n} J_{\varepsilon_h}(u_{\varepsilon_h}) \rightarrow c_{z_0},$$

we simply remark that $c_{z_0} = \Sigma(z_0)$. Therefore, if v_0 is the limit of the sequence (v_h) defined in (2.11), then $v_0 \in S(z_0)$ because we can exploit again some exponential barrier to pass to the limit. Now the proof proceeds as in the previous case. Finally,

if $S(z_0) = \{v_0\}$, the map Σ has all the directional derivatives at z_0 and, by virtue of (3.5) they are equal to zero. This proves (d). \square

We would like to remark that a different definition of concentration for single-peak solutions has been used in [9]. We recall it here, suitably adapted to our purposes.

Definition 3.7. Let $u_\varepsilon \in C^2(\mathbb{R}^n)$ be a family of positive solutions of the equation

$$-\varepsilon^2 \operatorname{div}(\alpha(x)\nabla u) + V(x)u = K(x)f(u) \quad \text{in } \mathbb{R}^n.$$

Let $x_\varepsilon \in \mathbb{R}^n$ with $\max_{\mathbb{R}^n} u_\varepsilon = u_\varepsilon(x_\varepsilon)$. We say that u_ε concentrates at $z_0 \in \mathbb{R}^n$ if

- (i) $x_\varepsilon \rightarrow z_0$ as $\varepsilon \rightarrow 0$;
- (ii) $\varepsilon^{-n} J_\varepsilon(u_\varepsilon) \rightarrow c_{z_0}$ as $\varepsilon \rightarrow 0$,

where J_ε is as in (3.4).

It is not difficult to check that if (u_ε) is a sequence as in this definition, then (u_ε) concentrates at z_0 in the sense of Definition 2.9, at an exponential rate (see Lemma 4.2 in [9]). In particular, by Theorem 3.6, we have $0 \in \partial\Sigma(z_0)$.

Remark 3.8. We finish with an open problem. Assume that (u_h) is a sequence of solutions of (P_ε) with $p = 2$ and $\beta(\xi) = \xi$, namely the semi-linear case. Suppose that these solutions concentrate at $z_0 \in \mathbb{R}^n$, and $S(z_0) = \{v_0\}$. Is it possible to prove that z_0 is a C^1 -stable critical point of Σ , according to the definition of Yanyan Li [11]?

REFERENCES

- [1] A. Ambrosetti, M. Badiale, S. Cingolani, *Semiclassical states of nonlinear Schrödinger equations*, Arch. Ration. Mech. Anal. **140** (1997), 285–300.
- [2] L. Damascelli, M. Ramaswamy, *Symmetry of C^1 solutions of p -Laplace equations in \mathbb{R}^N* , Adv. Nonlinear Stud. **1** (2001), 40–64.
- [3] M. Degiovanni, A. Musesti, M. Squassina, *On the regularity of solutions in the Pucci-Serrin identity*, Calc. Var. Partial Differential Equations, in press.
- [4] M. Del Pino, P. Felmer, *Local mountain passes for semilinear elliptic problems in unbounded domains*, Calc. Var. Partial Differential Equations **4** (1996), 121–137.
- [5] M. Del Pino, P. Felmer, *Semi-classical states for nonlinear Schrödinger equations*, J. Functional Anal. **149** (1997), 245–265.
- [6] F.H. Clarke, *Optimization and nonsmooth analysis*, Wiley-Interscience publication, (1983).
- [7] G.B. Li, *some properties of weak solutions of nonlinear scalar field equations*, Ann. Acad. Sci. Fenn. Ser. A. Math. **15** (1990), 27–36.
- [8] A. Giacomini, M. Squassina, *Multipeak solutions for a class of degenerate elliptic problems*, Asymptotic Anal., in press.
- [9] M. Grossi, R. Molle, *On the shape of the solutions of some semilinear elliptic problems*, Commun. Contemp. Math. **5** (2003), no. 1, 85–99.
- [10] Y. Kabeya, *Multiple solutions of some quasilinear elliptic equations on \mathbb{R}^n* , Adv. Math. Sci. Appl. **4** (1994), 161–190.
- [11] Y. Y. Li, *On a singularly perturbed elliptic equation*, Adv. Diff. Equations **2** (1997), 955–980.
- [12] Y.-G. Oh, *Existence of semiclassical bound states of nonlinear Schrödinger equations with potentials of the class $(V)_a$* , Comm. Partial Differential Equations **13** (1988), 1499–1519.
- [13] A. Pomponio, S. Secchi, *On a class of singularly perturbed elliptic equations in divergence form: existence and multiplicity results*, Preprint (2003), available at <http://babbage.sissa.it/>.

- [14] P. Pucci, J. Serrin, *A general variational identity*, Indiana Univ. Math. J. **35** (1986), 681–703.
- [15] P. Rabier, C.A. Stuart, *Exponential decay of the solutions of quasilinear second-order equations and Pohožaev identities*, J. Differential Equations **165** (2000), 199–234.
- [16] P.H. Rabinowitz, *On a class of nonlinear Schrödinger equations*, Z. Angew. Math. Phys. **43** (1992), 270–291.
- [17] J. Serrin, *Local behavior of solutions of quasi-linear equations*, Acta Math. **111** (1964), 247–302.
- [18] J. Serrin, M. Tang, *Uniqueness of ground states for quasilinear elliptic equations*, Indiana Univ. Math. J. **49** (2000), 897–923.
- [19] M. Squassina, *Spike solutions for a class of singularly perturbed quasilinear elliptic equations*, Nonlinear Anal., in press.
- [20] P. Tolksdorf, *Regularity for a more general class of quasilinear elliptic equations*, J. Differential Equations **51** (1984), 126–150.
- [21] X. Wang, *On concentration of positive bound states of nonlinear Schrödinger equations*, Comm. Math. Phys. **153** (1993), 229–244.
- [22] X. Wang, B. Zeng, *On concentration of positive bound states of nonlinear Schrödinger equations with competing potential functions*, SIAM J. Math. Anal. **28** (1997), 633–655.

DIPARTIMENTO DI MATEMATICA “L.TONELLI”
UNIVERSITÀ DI PISA
VIA BUONARROTI 2, I-56127 PISA, ITALY
<http://www.dm.unipi.it/~secchi>
E-mail address: secchi@dm.unipi.it

DIPARTIMENTO DI MATEMATICA “F.BRIOSCHI”
POLITECNICO DI MILANO
VIA L. DA VINCI 32, I-20133 MILANO, ITALY
<http://www1.mate.polimi.it/~squassina>
E-mail address: squassina@mate.polimi.it