KIRWAN MAP AND MODULI SPACE OF FLAT CONNECTIONS

SÉBASTIEN RACANIÈRE

ABSTRACT. If K is a compact Lie group and g 2 an integer, the space K^{2g} is endowed with the structure of a Hamiltonian space with a Lie group valued moment map . Let be in the centre of K. The reduction 1 ()=K is homeomorphic to a moduli space of flat connections. When K is simply connected, a direct consequence of a recent paper of Bott, Tolman and Weitsman is to give a set of generators for the K-equivariant cohomology of 1 (). Another method to construct classes in H $_{\rm K}$ (1 ()) is by using the so called universal bundle. When the group is SU (n) and is a generator of the centre, these last classes are known to also generate the equivariant cohomology of 1 (). The aim of this paper is to compare the classes constructed using the result of Bott, Tolman and Weitsman and the ones using the universal bundle.

INDEX OF NOTATION

- Group of relative integers Ζ
- Field of reals R
- Integer bigger than 1 g
- F
- Free group on 2g generators $x_1; :::; x_{2g}$ Element in F given by $\begin{array}{c} g \\ j=1 \end{array} \left[x_{2j 1}; x_{2j} \right]$ Quotient of F by the relation $\begin{array}{c} g \\ g \\ j=1 \end{array} \left[x_{2j 1}; x_{2j} \right] = 1$ R Closed Riemann surface of genus q 0
 - with the interior of a disc removed
- Κ Simply connected compact Lie group
- Z (K) Centre of K
- Quotient of K by its centre Z (K) Κ_c

Acknowledgement

I would like to thank Frances Kirwan for her comments on a preliminary version of this article.

1. INTRODUCTION

All cohomology will have coefficients in R.

Let be a closed compact Riemann surface of genus g. Let $_0$ be a compact Riemann surface with boundary obtained from by removing a small disc.

Let K be a compact simply connected Lie group. We denote by K_c the quotient of K by its centre Z (K). In other words, K_c is the projectivisation of K. Let be in the centre of K. Let K act on itself by conjugation and on K^{2g} by diagonal conjugation. The point

is fixed by K. Let EK ! BK be a classifying bundle for K. If M is a topological space acted on by K, the equivariant cohomology $H_K (M)$ of M is the singular cohomology of $M_K = M_K EK$. If M is a manifold and it is acted on smoothly by K, then $H_K (M)$ is also the cohomology of the Cartan-de Rham complex_K (M).

Let be the equivariant map

Let $Y = {}^{1}$ (). The space 1 ()=K is homeomorphic to the moduli space of flat connections on the trivial bundle K ${}_{0}$! ${}_{0}$ (all K principal bundles over ${}_{0}$ are trivial) with prescribed holonomy around the boundary S^{1} of ${}_{0}$.

Call the restriction map

$$:H_{K}$$
 (K^{2g}) ! H_{K} (Y):

Let $b_1; \ldots; b_r$ be primitive elements which generate H (K) and let $c_1; \ldots; c_r$ be their respective transgressions in H (BK). One can extend $b_1; \ldots; b_r$ to equivariant classes $b_1^K ; \ldots; b_r^K$ on K. Let $pr_i : K^{2g}$! K be the projection on the ith factor and $b_{j,i}^K$ the pull-back of b_j^K by pr_i .

Because is a moment map on the quasi-Hamiltonian space K ^{2g} (see [2]), the paper [5] gives a way of constructing equivariant classes $a_1; :::; a_r$ on Y (see Equation (1) on page 6) such that f (c_j); $(b_{j,i}^K); a_j; j = 1; :::; r; i = 1; :::; 2gg$ generate the K -equivariant cohomology of Y.

Also, one can construct a K -equivariant $K_{\rm c}\mbox{-}{\rm principal}$ bundle over

Y : the so called universal bundle. The Künneth decomposition of the equivariant characteristic classes of this bundle allows us to construct classes $fc_{j}^{0};b_{j;i}^{0};a_{j}^{0};j = 1;:::;r;i = 1;:::;2gg in H_{K}$ (Y) (see Equation (2) on page 7). At least when K is the special unitary group and is a generator of the centre of K , these classes are known to generate $H_{\rm K}$ (Y) as a ring. They are the Atiyah-Bott-Biswas-Raghavendra classes¹.

The aim of this paper is to prove the following theorem.

Theorem 3.2. — Let K be a compact, connected and simply connected Lie group. Let be in the centre of K such that K acts locally freely on Y. Let $a_j^0; b_{j;i}^0; c_j^0; a_j; b_{j;i}^K; c_j$ be equivariant classes defined as above. The following relations hold

$$C_{j}^{0} = (C_{j})$$

 $b_{j;i}^{0} = (b_{j;i}^{K})$:

Also, the classes fa_jg depend on certain choices but one can choose them so that

$$a_{j}^{0} = a_{j}$$
:

This article is organised as followed. In Section 2, I recall what [5] says about the restriction map involved in the reduction of K 2g at . I also show how to use this result to construct generators for the K - equivariant cohomology of Y . In Section 3, I describe the construction of equivariant characteristic classes on Y using a universal bundle over

Y. I also outline the proof of Theorem 3.2, leaving the most technical details for Section 4.

2. Generators for the equivariant cohomology of Y

Let be the fundamental group of . Let us fix a presentation

=
$$hx_1; :::; x_{2g}; \sum_{j=1}^{Y^g} [x_{2j 1}; x_{2j}] = 1i:$$

Let F be the fundamental group of $_0$. We fix a presentation

$$F = hx_1; \ldots; x_{2\alpha}i$$

such that the map

F !

induced by the inclusion of $_0$ in $% 10^{-1}$, is given by the obvious map x_k 7 ! $x_k.$

¹Atiyah and Bott [1] were the first ones to construct generators of the cohomology of the moduli space. Their classes depended on the choice of a universal bundle which was only defined up to the tensor product with a line bundle. The universal bundle of this article is actually the projectivisation of theirs. It is well-defined and so are the generators constructed from it. This was proved by Biswas and Raghavendra in [3]

SÉBASTIEN RACANIÈRE

Let be in the centre of K. Let Y be the subset of K^{2g} defined by

$$Y = (X_{1}; ...; X_{2g}) 2 K^{2g} j_{j=1}^{Y^{g}} [X_{2j1}; X_{2j}] = :$$

The group K acts on K^{2g} by conjugation. This action restricts to Y, since is in the centre of K. The quotient M = Y = K can be identified with a moduli space of flat connections on the trivial principal bundle K $_0 !_0$ (because K is simply connected, all principal bundles are trivial over $_0$), with holonomy around the boundary Q $_0$. If K acts locally freely on Y, then the cohomology of M is isomorphic to the K-equivariant cohomology of Y (recall that cohomology is taken with coefficients in R).

Let be the restriction map

The space K $^{\rm 2g}$ has the structure of a quasi-Hamiltonian space (see [2]) with moment map

:
$$K^{2g}$$
 ! Q_{g} K
(X₁:::;X_{2g}) 7! $Q_{j=1}^{g}$ [X_{2j1};X_{2j}]:

The map is a submersion at any point of $Y = {}^1$ () if and only if K acts locally freely on Y. In this nice case, the space M is a compact symplectic orbifold since it is the regular reduction of K 2g at .

Let b_1 ;:::; b_r be primitive elements in H (K) that generate the cohomology of K as a ring

H (K) =
$$\begin{pmatrix} X^{r} \\ R \\ j=1 \end{pmatrix}$$

Each b_i is of odd degree.

Let c_1 ;:::; c_r be the transgressions in H (B K) of respectively b_1 ;:::; b_r , thus deg c_j = deg b_j + 1 and

H (BK) =
$$R[c_1; :::; c_r]$$
:

For each K-principal bundle $G \quad ! \quad X$, the classes $c_1; :::; c_r$ define characteristic classes c_1 (G); :::; c_r (G) in H (X).

Proposition 2.1. Let EK ! BK be the classifying bundle for K. The bundle $EK_{K}K^{n}$! BK is cohomologically trivial as a ring, that is

$$H_{K}$$
 (Kⁿ) ' H (BK) H (Kⁿ)

as rings.

This result is not new but I will propose here a proof which is original as far as I know.

Also, because I want to state the principal results as soon as possible, I will make use in the following proof, as well as in others, of Lemma 3.7 even if I haven't proved it yet.

Proof. We will prove the case n = 1. The general case is similar, one only has to use Proposition 3.5 instead of Lemma 3.7 (or use the fact that if X has an equivariantly formal action of K then the diagonal action of K on Xⁿ is also equivariantly formal).

Let us first prove that H_K (K) and H (BK) H (K) are isomorphic as H (BK) modules. By the Leray-Hirsch Theorem, all we have to do is prove that the homomorphism

is surjective. Let b_j be one of the generators of $H \ll 1$. Let $D ! S^1 \ltimes be$ the K-equivariant K-principal bundle of Lemma 3.7. As a K-principal bundle, its characteristic class $c_j \oplus 0$ is $dt \ b_j$ (Lemma 3.7). As an immediate corollary, we deduce that there exists b_j^K in $H_K \ll 1$ such that $c_j \oplus \kappa \in K$ = 1 $c_j + dt \ b_j^K$, where b_j^K restricts to b_j . Now, because the b_j 's generate $H \ll 1$ as a ring, the homomorphism

Н_К (К) ! Н (К)

is surjective and we have proven the existence of a H (BK) module isomorphism between H_K (K) and H (BK) H (K). The existence of a ring isomorphism follows because the cohomology H (K) of the fibre in K _K EK ! BK is generated by classes of odd degree (see [9]). 2

Remark 2.2. Let s be the section

Let n be an integer (we will need it to be either 1 or 2g). Let $B_{j,i}^{K}$ be the pull-back, under the projection on the *i*-th factor K^{n} ! K, of B_{j}^{K} .

Lemma 2.3. The K -equivariant cohomology of Kⁿ is isomorphic as a ring to γ

 H_{K} (Kⁿ) ' ($Rb_{j;i}^{K}$) R [c₁;:::;c_r]:

SÉBASTIEN RACANIÈRE

Before stating the result of Bott, Tolman and Weitsman for , we need to construct some classes in H $_{\rm K}$ (Y).

Lemma 2.4. For all j, the pull-back under of \mathbb{B}_{j}^{K} is zero

 $(b_{i}^{K}) = 0$:

In their article [5], the authors already gave this result. I give here a proof using Lemma 3.7.

Proof. Consider the diagram

where the horizontal map m_1 is induced by the moment map : K^{2g} ! K and the vertical map m_2 is given by the injection of S¹ as the boundary of $_0$. The pull-back of D by m_1 and the pull-back of F by m_2 are isomorphic. Indeed they are both equivariantly isomorphic to the bundle

(R K^{2g} K)=Z ! S¹ K^{2g}

where the group Z acts by

We then deduce the lemma from Proposition 3.5 and Lemma 3.7 by noticing that the injection of S^1 as the boundary of $_0$ induces the null map in degree 1 cohomology. 2

Let b_1 ;:::; b_r be equivariant forms representing respectively b_1^K ;:::; b_r^K .

Remark 2.5. Such forms will be chosen in Section 4 where it will be proved that their pull-backs by the section s of Remark 2.2 vanish.

The preceding lemma implies there exist equivariant forms a_1 ;:::; a_r on K ^{2g} such that

(1)
$$(b_j) = da_j; 8j:$$

The $a_1\textit{;:::;}a_r$ are closed when restricted to Y . They define classes $a_1\textit{;:::;}a_r$ in H $_K$ (Y).

Theorem 2.6 (Bott, Tolman and Weitsman). The equivariant cohomology H_K (Y) is generated as a ring by the image of and the $a_1; :::; a_r$.

As an immediate consequence, we have the following corollary.

Corollary 2.7. The equivariant cohomology H_{K} (Y) is generated as a ring by the classes (C_j); $(B_{j;i}^{K})$; a_{j} for $j = 1; \ldots; r$ and $i = 1; \ldots; 2g$.

```
3. Some classical classes in H _{\rm K} (Y )
```

Let e be the universal cover of . The group acts on e. Since we have chosen a presentation for F, we have an isomorphism

Hom (F;K) !
$$K^{2g}$$

7! $((x_1);...; (x_{2g})):$

We will use this isomorphism to identify $H \circ m$ (F;K) and K^{2g}. Let us define an action of on $\stackrel{e}{=}$ Y K_c

where **p** is any element in **F** lying above **p**. This action is well-defined and free. The quotient **E**, together with the projection **:E ! Y** define a K_c -principal bundle. The actions of **K** on **Y** by conjugation and on K_c by multiplication on the left, make **E** into a **K**-equivariant K_c -principal bundle over **Y**. We call it the universal bundle.

Example 3.1. For K = SU(n) and a generator of Z(SU(n)), this is the projectivisation of a universal bundle for the moduli space of semi-stable vector bundles of fixed rank and determinant (see [1] and [9]).

Because K is compact and simply connected, we have

H (BK) ' H (BK_c):

The classes $c_1; :::; c_r$ in H (BK_c) define equivariant characteristic classes for the bundle E (recall that these are the same as the usual characteristic classes of the K-principal bundle E _K EK ! Y _K EK). These classes live in H_K (Y)' H () H_K (Y). Let ₁;:::; _{2g} be a basis of H (), dual to $x_1;:::;x_{2g}$ in = ₁(). Let V be a volume form of volume 1 for . Using the Künneth decomposition, let us write

(2)
$$c_{j} (E K E K) = 1 c_{j}^{0} + \sum_{i=1}^{X^{2g}} b_{j;i}^{0} + V a_{j}^{0}$$
:

In Example 3.1, this classes are the known Atiyah-Bott – Biswas-Raghavendra generators of H_K (Y). A natural question is: what is the relation between these classes and the ones of Corollary 2.7. An answer is given in the following theorem.

Theorem 3.2. Let K be compact, connected and simply connected Lie group. Let be in the centre of K such that is a regular value of the moment map . Let the $fa_{j}^{0};b_{j;i}^{0};c_{j}^{0};a_{j};b_{j;i}^{K};c_{j}g$ be equivariant classes defined as above. The following relations hold

$$C_{j}^{0} = (C_{j})$$

 $b_{j;i}^{0} = (C_{j;i}^{K})$

Also, the classes $\mathtt{fa}_{\mathtt{j}}\mathtt{g}$ depend on certain choices but one can choose them so that

$$a_{j}^{0} = a_{j}$$
:

Proof. Let us fix j. The first part of the theorem is a consequence of Proposition 3.4 and Proposition 3.5. Indeed, by Proposition 3.4

$$c_j(E_K EK) j_0 Y = c_j(F_K EK) j_0 Y$$

But

(3)
$$c_{j}(E K E K) j_{0} Y = 1 c_{j}^{0} + i b_{j;i}^{0}$$

and by Proposition 3.5

(4)
$$c_{j}(F_{K} E K) j_{0} Y = (id_{c_{j}})c_{j}(F_{K} E K)$$

 $= 1 (c_{j}) + \sum_{i=1}^{\chi^{2g}} (c_{j,i}):$

The first part of the theorem follows by comparing Equation (3) and Equation (4).

The second part of Theorem 3.2 is a consequence of the following Lemma and of the definition of the classes a_1 ;:::; a_r .

Lemma 3.3. One can choose representatives b_j^{K} and a_j^{0} of respectively b_j^{K} and a_j^{0} such that a_j^{0} extends to an equivariant class on K^{2g} whose differential is (b_j^{K}) .

The proof of this Lemma, together with the proof of Lemma 3.7, is given in the next Section.

This finishes the proof of Theorem 3.2.

2

Let e_0 be the universal covering for $_0$. Let us define an action of $F = _1 (_0)$ on $e_0 - K^{2g} - K$

$$F = e_0 K^{2g} K ! e_0 K^{2g} K$$

(p; ; ;k) 7! (p; ; (p)k):

This action is free and the quotient F is a K-principal bundle on $_{0}$ K^{2g}. The action of K by conjugation on K^{2g} and by multiplication on the left on K makes F into a K-equivariant K-principal bundle. The situation here is somehow confusing since there are two different actions of K on F. In order to clarify the situation, I will call the action that makes F a K-principal bundle the principal action and call the action that makes it K-equivariant the covering action ('covering' because it covers the action of K on K^{2g}).

Proposition 3.4. The projectivisation of $F j_{0Y}$ is K -equivariantly isomorphic to $E j_{0Y}$.

Proof. The proof is easy.

Now all we have to do is compute the equivariant characteristic classes of F. The cohomologies of and $_0$ are the same in degree 0 and 1 and $_0$ has no cohomology in degree 2. We use the same basis $_1;:::; _{2g}$ for H¹() and H¹($_0$).

Proposition 3.5. Using the Künneth decomposition, the equivariant characteristic classes of F are

$$c_{j} (F K E K) = 1 c_{j} + \sum_{i=1}^{X^{2g}} b_{ji}^{K}$$

Proof. The surface $_0$ is homotopic to a wedge product of 2g circles. As is easily seen, Proposition 3.5 will follow from the computation of the characteristic classes of the restriction of F to the n-th circle in the wedge product times the m-th copy of K in K^{2g}. When $n \in m$, the restriction of F is trivial. Thus, I only have to do the computation for n = m. This computation is the object of the next two lemmas. 2

Let us call D_n the restriction of F to the n-th circle in the wedge product times the n-th copy of K in K^{2g}. I identify the circle S¹ with the quotient R=Z. I denote by t the natural coordinate in R, thus dt defines a volume form on S¹.

Lemma 3.6. All the D_n 's are isomorphic to the same bundle $D_{!}$ S¹ K. The total space D is (R K K)=Z where the action of Z is defined by

Z (R K K) ! R K K
(
$$;t;k_1;k_2$$
) 7! (t+ $;k_1;k_1k_2$)

and the projection is

(R K K)=Z ! S^{1} K ft;k₁;k₂g 7! (ftg;k₁):

Proof. Fix an integer n between 1 and 2g and let $x = x_n$ be the n-th generator of F.

Recall that F was defined as a quotient

 $F = (e_0 K^{2g} K) = (0);$

Let h be the group homomorphism

Z !
$$_{1}(_{0})$$

7! x:

This morphism makes Z into a subgroup of $_1(_0)$. Choose a point e_0 in $_0$ lying above the base point $_0$ of $_0$. Choose a loop $S^1 ! _0$ representing x. This loop lifts to a map $: \mathbb{R} ! e_0$ such that $(0) = e_0$ and for an integer (t +) = x (t).

The covering action of K (the one that covers the action of K on $S^1 ext{ K}$) on D_n is induced by the conjugation on the first factor K in R K K and by multiplication on the left on the second factor K. Let :K ! K^{2g} be the injection of the n-th factor. Define a map

 $(R \quad K \quad K) = Z \quad ! \quad (e_0 \quad K^{2g} \quad K) = _1 (_0) \\ [t; X; k] \quad 7 \; ! \qquad [(t); (X); k]:$

This map is well-defined since it sends the translated <code>t+ ;X;X k]of</code> <code>t;X;k]by</code> to

$$[(t+); (X);X k] = [x_n (t); (X);Xk] = [(t); (X);k]:$$

It is also K -equivariant because, for `in K, it sends [t;AdX; `k] to

The lemma follows.

2

Lemma 3.7. The characteristic classes of D are

$$c_j (D) = dt b_j$$
:

This last lemma is actually very useful since it can be used to prove Proposition 3.5 which leads to Theorem 3.2, and also Proposition 2.1 and Lemma 2.4.

4. Proof of Lemma 3.3 and Lemma 3.7

The aim of this section is to prove Lemma 3.3 and Lemma 3.7 starting with the last one.

The proof of Lemma 3.7 goes as follows: in a similar way to Jeffrey[7], I construct explicitly a map from $S^1 ext{K}$ to the fat realisation of $B ext{K}$ as a simplicial manifold such that the pull-back of $E ext{K}$ under this map is the principal bundle D_n . I then use a result of Shulman to prove Lemma 3.7.

Let us start by introducing the simplicial realisations of EG and BG for any group G (see [10]). Let

$${}^{m} = (t_{0}; :::; t_{m}) 2 [0; 1]^{n+1} j X^{n} t_{i} = 1$$

be the standard m-simplex. The simplicial realisation \overline{NG} , respectively NG, of EG, respectively BG, is given by the contravariant functor

f ^m; m 2 Ng ! fcategory of smooth manifoldsg

which to each $_{m}$ associates \overline{NG} (m) = G^{m+1} , respectively NG (m) = G^{m} . For ibetween 0 and m, the map $_{i}:\overline{NG}$ (m) ! \overline{NG} (m 1), respectively $_{i}:NG$ (m) ! NG (m 1), corresponding to the ith face map $^{i}: {}^{m-1}$! m is given by the omission of the ith term

$$_{i}(k_{0}; \ldots; k_{m}) = (k_{0}; \ldots; k_{i}; \ldots; k_{m});$$

respectively

$$(5)_{i}(k_{1}; \dots; k_{m}) = \begin{cases} 8 \\ < (k_{2}; \dots; k_{m}) & i = 0 \\ (k_{1}; k_{2}; \dots; k_{i}k_{i+1}; \dots; k_{m}) & i = 1; \dots; m \\ (k_{1}; \dots; k_{m-1}) & i = m : \end{cases}$$

The projection NG (m) ! NG (m) is the map

$$q_{n} (k_{0}; \ldots; k_{m}) = (k_{1}k_{2}^{1}; \ldots; k_{m-1}k_{m}^{1}):$$

We then take EG, respectively BG, to be the fat realisation of \overline{NG} , resp. NG, that is EG is the quotient of $m^m \overline{NG}$ (m) by the relation

$$(i i d)(t;x)$$
 $(i d_i)(tx)$

for (t;x) in ^{m 1} \overline{NG} (m) and BG is the quotient of ^F_m ^m NG (m) by the relation

 $(^{i} id)(t;x)$ (id $_{i})(tx)$

for (t;x) in ^{m 1} NG (m). The projection q:EG ! BG is induced by the maps q_n . The action of G on EG is induced by the diagonal multiplication of G on the right on NG $(m) = G^{m+1}$ for each m. In fact, there is a second action of G on EG, the covering action, which descends to an action of G on BG. This action on EG is induced by the diagonal multiplication on the left of G on \overline{NG} (m) = G^{m+1} , it descends to an action on BG induced by the diagonal conjugation on NG (m) = G^m .

Notice that

$$S^{1} = BZ$$
$$= B$$
$$_{0} = BF$$

In the simplicial models, the injection of $_0$ in $\,$ is given by the projections

$$\mathbf{F}^{n}$$
 ! n :

The surface $_0$ is homotopic to the wedge product of 2g circles. The inclusion of the \pm th circle S^1 ! $_0$ is given by

$$Z^{n} : F^{n}$$

7! $(x_{i};...;x_{i}):$

Let $R = {Q \atop j=1}^{q} [x_{2j1}; x_{2j}]$ be in F. The inclusion of S^1 as the boundary of 0 is the realisation of the simplicial map

The classifying map $f : S^1 \times I = BK$ for the bundle D is the realisation of the simplicial map

The homology of S^1 is the Eilenberg-Mac Lane group homology H (Z) of Z (see [8]). Denote by x a generator of Z. A generator of H₁(Z) is given by x.

Since the centre of K is finite, on the one hand K and K_c, and on the other hand BK and BK_c have naturally isomorphic cohomologies. In addition, every K-principal bundle defines a K_c-principal bundle in a natural way: one just has to compose the classifying map of the K-principal bundle with the projection BK ! BK_c. The simplicial version of this last projection is given by the maps Kⁿ ! Kⁿ_c. Moreover, the characteristic classes of the induced K_c-principal bundle are the same as the one of the K-principal bundle.

The cohomology of BK_{c} is the cohomology of a double complex $p_{\mathcal{P}_{q}} = {}^{p} (MK_{c}(q))$ with two differentials: one is the usual differential of forms d: ${}^{p} (MK_{c}(q))$! ${}^{p+1} (MK_{c}(q))$ and the other is

:
$${}^{p}(NK_{c}(q)) : {}^{p}(NK_{c}(q+1)) = {}^{q+1}_{i=0}(1)^{i}{}^{n}_{i}$$

Recall that the maps "i were defined in (4). The two differentials d and commute and the total differential is $d+(1)^{p}$ on ^{prq}. Shulman [11] (see also Bott, Shulman and Stasheff [4]) proved that each class c_{j} in H^{2rj} (BK_c) can be represented by a form $c_{j;1} + \frac{1}{2r} \dot{c}n$, with $c_{j;k} 2 = \frac{2r k_{jk}}{r}$. That this form is closed means that

$$\begin{array}{rcl} dc_{j;1} &=& 0 \\ c_{j;k} &=& (1)^k dc_{j;k+1} \mbox{ for } 1 & j < r_j \\ c_{j;r} &=& 0 \end{array}$$

Also, the closed form $c_{j;1} 2^{-2r_j - 1}$ (K $_c$) is a representative of b_j .

Everything in the above paragraph can be done equivariantly. One just has to replace cohomology by equivariant cohomology, the complex ${}^{p_{\mathcal{A}}} = {}^{p} (\mathbb{N} K_{c}(\mathbf{q}))$ by ${}^{p_{\mathcal{A}}} = {}^{p}_{K} (\mathbb{N} K_{c}(\mathbf{q}))$, the differential d by d_{K} , the forms \mathbf{c}_{j} by their respective extensions c_{j}^{K} in $H_{K}^{2r_{j}}$ (BK_c), and so on. Hence it is clear that the pairing hx; f c_{j} is b_{j}^{K} . This together with the fact that the restriction of D to S¹ is trivial proves Lemma 3.7. We see that Lemma 3.7 is almost tautological.

I chose $c_{j;l}^{K}$ as a representative b_{j}^{K} of b_{j}^{K} . The section

is obtained from the inclusion of the identity 1 into K by applying the functor, from K -spaces to spaces,

$$X 7! X _{K} EK:$$

Because the inclusion map flg ! K is constant and because $c_{j;l}^{K}$ is of odd degree, it follows that $s b_{j}^{K} = s c_{j;l}^{K} = 0$. This proves Remark 2.5.

The equivariant forms $C_{j;1}^{K}$ and $C_{j;2}^{K}$ satisfy $d_{K} C_{j;2}^{K} = C_{j;1}^{K}$. A classifying map for the K-equivariant K_c-principal bundle E is the realisation of the equivariant simplicial map

: Y ⁿ !
$$K_{c}^{n}$$

(;p_1;:::;p_n) 7! ((p_1);:::; (p_n)):

The homology of is the Eilenberg-Mac Lane group homology H $\,$ (). A generator of H $_2$ () is (see [6, Proposition 3.9])

$$c = \frac{X^{2g}}{\lim_{i=1}^{d} \frac{QR}{QX_{i}}} \quad x_{i};$$

where $\frac{@R}{@x_i}$ is the Fox free differential calculus (see [6, Section 3]). In fact,

$$\frac{QR}{Qx_i} = \int_{j}^{0} \int_{i}^{1} \mathbf{i};$$

where $_{i} 2 F$ is

$$\begin{array}{cccc} 0 & = & Q_{i1} \\ 2i1 & = & & \\ 1 & = 1 \\ 2i1 & = & & \\ 2i1 & x_{2i1} \\ 2i & = & & \\ 2i1 & x_{2i1} \\ 2i1 & x_{2i1} \\ 1 \\ 2i & = & & \\ 2i1 & x_{2i1} \\ 1 \\ 2i1 & = & & \\ 2i1 & x_{2i1} \\ 1 \\ 2i1 & x_{2i1} \end{array}$$

A representative a_j^0 of a_j^0 is given by

where, for $\mathtt{p2}$ F, the evaluation map is

$$ev_p$$
: Hom (F;K_c) ! K_c
7! (p):

The last line of the above computation of a_j^0 clearly shows that a_j^0 extends to a form on K ^{2g}. The differential of this form is

where

$$z_{i;0} = x_i; z_{i;1} = x_i \text{ and } z_{i;2} = i$$

Because

$$\begin{array}{rcl} & & \mbox{ev}_{z_{i;0}^0} & = & \mbox{ev}_{z_{i;1}^1} \\ & & \mbox{ev}_{z_{2i;1}^1} & = & \mbox{ev}_{z_{2i-1;2}^1} \\ & \mbox{ev}_{z_{2i-1;1}^0} & = & \mbox{ev}_{z_{2i;2}^0} \\ & \mbox{ev}_{z_{2i;1}^0} & = & \mbox{ev}_{z_{2i-1;1}^1} \\ & \mbox{ev}_{z_{2i;1}^0} & = & \mbox{ev}_{z_{2i;1}^1} \\ & \mbox{ev}_{z_{2i;1}^0} & = & \mbox{ev}_{z_{2i;2}^1} \end{array}$$

many terms cancel each other in the expression (6) above. We are left with

$$da_{j}^{0} = ev_{z_{2g;2}^{1}}c_{j;1}^{K} ev_{z_{1;2}^{0}}c_{j;1}^{K}$$
:

But $z_{1;2}^0 = 1$ the identity of F and hence $ev_{z_{1;2}^0}$ is a constant map, whereas $z_{2g;2}^1 = {Q \atop j=1}^g [x_{2j \ 1}; x_{2j}]$ and hence $ev_{z_{2g;2}^1} = .$ Since $c_{j;1}^K$ is a representative of b_{i}^K , we deduce that on K^{2g}

$$da_j^0 = b_j^K$$
:

This proves Lemma 3.3.

References

- M. F. ATIYAH and R. BOTT The Yang-Mills Equations Over Riemann Surfaces, Phil. Trans. Royal Soc. Lond. A308 (1982), 523-615.
- [2] A. ALEKSEEV, A. MALKIN and E. MEINRENKEN Lie group valued moment maps, J. Differential Geom. 48 (1998), 445–495.
- [3] I. BISWAS and N. Raghavendra Canonical generators of the cohomology of moduli of parabolic bundles on curve, Math. Ann. 306 (1996), 1–14.
- [4] R. BOTT, H. SHULMAN and J. STASHEFF On the de Rham theory of certain classifying spaces, Adv. Math. 20 (1976), 43–56.
- [5] R. BOTT, S. TOLMAN and J. WEITSMAN Surjectivity for Hamiltonian Loop Group Spaces, arXiv:math.DG/0210036 v1 (Oct. 2002).
- [6] W. GOLDMAN The symplectic nature of the fundamental groups of surfaces, Adv. Math. 54 (1984), 200-225.
- [7] L. JEFFREY Group cohomology construction of the cohomology of moduli spaces of flat connections on 2 manifolds, Duke Math. J. 77 (1995), 407–429.
- [8] S. MAC LANE Homology, Grundlehren Math. Wiss. 114 (1963), Springer-Verlag, Berlin.
- [9] S. RACANIÈRE *Restriction map in a regular reduction of* SU (n)^{2g}, to appear in Commentarii Mathematica Helvetici.
- [10] G. SEGAL Classifying spaces and spectral sequences, Inst. Hautes Études Sci. Publ. Math. 34 (1968), 105–112.
- [11] H. B. SHULMAN Characteristic Classes and Foliations, Doctorate (1968), University of California, Berkeley.

MATHEMATICS DEPARTMENT, SOUTH KENSINGTON CAMPUS, IMPERIAL COL-LEGE LONDON, SW7 2AZ, UK

E-mail address: s.racaniere@ic.ac.uk *URL*: www.ma.ic.ac.uk/~racani