Spectral portraits and the resolvent growth of a model problem associated with the Orr–Sommerfeld equation

Shkalikov A. A.*

Department of Mechanics and Mathematics Moscow State University ashkalikov@yahoo.com

The Orr–Sommerfeld equation is obtained by linearization of the Navier–Stokes system in the infinite spacial layer $(x, \xi, \eta) \in \mathbb{R}^3$, where $|x| \leq 1$ and $(\xi, \eta) \in \mathbb{R}^2$. This equation has the form

$$(D^{2} - \alpha^{2})y - i\alpha R \left[q(x)(D^{2} - \alpha^{2}) - q''(x)\right]y = i\alpha R\lambda (D^{2} - \alpha^{2})y.$$
(1)

Here D = d/dx, α is the wave number ($\alpha \neq 0$) appearing after the separation of the variables $(\xi, \eta) \in \mathbb{R}^2$, R is the Reynolds number, q(x) is the stationary unperturbed velocity profile and λ is the spectral parameter. Usually the boundary conditions

$$y(\pm 1) = y'(\pm 1) = 0 \tag{2}$$

are associated with equation (1).

Our main goal is to understand the spectrum behaviour of problem (1), (2) as the Reynolds number R tends to the infinity (this means that the liquid becomes to be ideal). For these purposes we associate with (1), (2) a problem of the form

$$-i\varepsilon^2 z'' + q(x)z = \lambda z \tag{3}$$

$$z(-1) = z(1) = 0, (4)$$

assuming that q(x) is the same as in (1) and $\varepsilon^2 = (\alpha R)^{-1}$. It turns out that this problem serves as a good model for the study of the original Orr–Sommerfeld problem (1)–(2). Now the question is: for which function q(x) we can describe the spectral portraits of problem (3), (4) as $\varepsilon \to 0$? The answer is not simple even for particular functions q(x), for example $q(x) = x^2$ or $q(x) = (x - \alpha)^2$, $\alpha \in \mathbb{R}$. Only in the case q(x) = x the problem can be solved explicitly [1], [2]. At the moment we have general result only for analytic monotonous functions.

We shall say that a function q(x) belongs to the class AM if the following conditions hold: q(x) is analytic on the segment [-1, 1] and admits an analytic continuation in a domain G such that the range q(G) contains the semistrip $\Pi = \{\lambda \mid a < \operatorname{Re} \lambda < b, \operatorname{Im} \lambda < 0\}$ where a = q(-1), b = q(1). Moreover, if $D = q^{-1}(\Pi) \subset G$, then q(z) is continuous on the closure \overline{D} and the map $q(z): \overline{D} \to \overline{\Pi}$ is one to one.

^{*}Research supported by RFFI grant No 01-01-00691 and by the grant "The Universities of Russia"

Assuming that $q(x) \in AM$ we introduce the functions

$$Q^{\pm}(\lambda) = \int_{\xi_{\lambda}}^{\pm 1} \sqrt{i(q(\xi) - \lambda)} \, d\xi, \qquad \lambda \in \overline{\Pi},$$

where $\xi_{\lambda} \in \overline{D}$ is the root of the equation $q(\xi) - \lambda = 0$, and the function

$$Q(\lambda) = \int_{-1}^{1} \sqrt{i(q(x) - \lambda)} \, dx, \qquad \lambda \in \overline{\Pi}.$$

These functions are analytic in Π and continuous in $\overline{\Pi}$, provided $q(x) \in AM$.

The following theorem is a generalization of the previous author's result [3].

Theorem 1. Let $q(x) \in AM$. Then the curves $\tilde{\gamma}_{\pm}$ determined in the semistrip Π by the equations $\operatorname{Re} Q^{\pm}(\lambda) = 0$ have the only intersection point in Π , say λ_0 . The curve $\tilde{\gamma}_{\infty}$ determined in Π by the equation $\operatorname{Re} Q(\lambda) = 0$, is a function with respect to the imaginary axis and intersect the curves $\tilde{\gamma}^{\pm}$ in the only point λ_0 . Denote by γ_+ , γ_- and γ_{∞} the parts of $\tilde{\gamma}_+$, $\tilde{\gamma}_-$ and $\tilde{\gamma}_{\infty}$ connecting the intersection point λ_0 with the points a, b and $-i\infty$, respectively. Denote also

$$\Gamma = \gamma_+ \cup \gamma_- \cup \gamma_\infty$$

Then the spectrum of model problem (3), (4) is concentrated along Γ as $\varepsilon \to 0$, i. e. $\forall \delta > 0$ there are no eigenvalues outside the δ -neighbourhood of Γ , provided that $\varepsilon < \varepsilon_0 = \varepsilon_0(\delta)$. Moreover, the eigenvalues of the problem near the curves γ_+ , γ_- and γ_{∞} are determined from the conditions

$$i \int_{\xi_{\lambda}}^{1} \sqrt{i(q(\xi) - \lambda)} d\xi = \varepsilon \pi (k - 1/4), \qquad k \in \mathbb{Z},$$
$$i \int_{\xi_{\lambda}}^{-1} \sqrt{i(q(\xi) - \lambda)} d\xi = \varepsilon \pi (k - 1/4), \qquad k \in \mathbb{Z},$$
$$i \int_{-1}^{1} \sqrt{i(q(\xi) - \lambda)} d\xi = \varepsilon \pi k, \qquad k \in \mathbb{Z}.$$

More precizely, let μ_k^+ , μ_k^- and μ_k are solutions of these equations lying on the curves γ_+ , $\gamma_$ and γ_{∞} outside δ -neighbourhoods of the points a, b and λ_0 . Then there is a constant $C = C(\delta)$ such that each circle of the radius $C\varepsilon^2$ centered at the points μ_k^- , μ_k^+ and μ_k contains the only eigenvalue. The eigenvalue counting functions along the curves γ_+ , γ_- and γ_{∞} have representations

$$N_{\pm}(\lambda) = \pm \frac{1}{i\pi\varepsilon} \int_{\xi_{\lambda}}^{\pm 1} \sqrt{i(q(\xi) - \lambda)} \, d\xi + O(1),$$
$$N(\lambda) = \frac{1}{i\pi\varepsilon} \int_{-1}^{1} \sqrt{i(q(\xi) - \lambda)} \, d\xi + O(1),$$

where O(1) is uniformly bounded as $\varepsilon \to 0$ provided that λ lies outside some neighbourhoods of the points a, b and λ_0 .

Our next results clarify the connection of model problem (3), (4) with original Orr-Sommerfeld problem (1)-(2).

Theorem 2. Let $q(x) \in AM$. Denote by $\Delta_m(\lambda, \varepsilon)$ and $\Delta_{OS}(\lambda, \varepsilon)$ the characteristic determinants of the model and Orr-Sommerfeld problems, respectively. Then

$$\Delta_{OS}(\lambda,\varepsilon) = \Delta_m(\lambda,\varepsilon)R(\lambda)(1+O(\varepsilon)),$$

where $R(\lambda) \neq 0$ for $\lambda \in \Pi$ and $|O(\varepsilon)| \leq C\varepsilon$ for $\lambda \in \Pi \setminus \Gamma_{\delta}$. Here Γ_{δ} is the δ -neighborhood of the limit spectral graph Γ determined in Theorem 1, and the constant C depends only on δ .

The technique, developed in [4] allows to prove the similarity of the zero counting functions for a pair of holomorphic functions, provided that they have similar growth near the curves where the zeros are located (in our case on the boundary of Γ_{δ}). Hence, Theorem 1 leads to the following result.

Theorem 3. Let $q(x) \in AM$. Then the limit spectral graph of the Orr-Sommerfeld problem coincides with the graph Γ of the corresponding model problem, i.e. $\forall \delta > 0$ there are no eigenvalues of problem (1), (2) outside the δ -neighbourhood of Γ , provided that $R > R_0 = R_0(\delta)$. Moreover, the eigenvalue counting functions $N_{\pm}(\lambda)$ and $N_{\infty}(\lambda)$ along the curves γ_{\pm} and γ_{∞} for problem (1), (2) have representations

$$N_{\pm}(\lambda) = \pm \frac{1}{i\pi\varepsilon} \int_{\xi_{\lambda}}^{\pm 1} \sqrt{i(q(\xi) - \lambda)} \, d\xi \, (1 + o(1))$$
$$N(\lambda) = \frac{1}{i\pi\varepsilon} \int_{-1}^{1} \sqrt{i(q(\xi) - \lambda)} \, d\xi \, (1 + o(1))$$

where $o(1) \to 0$ as $\varepsilon \to 0$ uniformly for all λ lying outside some neighbourhoods of the points a, b and λ_0 .

Next our goal is to understand the resolvent behaviour of the operators corresponding to problems in question. Denote by $z_k = z_k(x, \varepsilon)$ the normed eigenfunctions of the model problems (for symplicity we assume that there is no associated functions, although they do appear for some discrete values of $\varepsilon_j \to 0$). It is known that the system $\{z_n\}_1^\infty$ forms a Riesz basis in $L_2(-1, 1)$, therefore, the Gramm matrix $\{(z_k, z_j)\}_{k,j=1}^\infty$ generates a bounded and boundely invertible operator T in l_2 . The number

$$C = C(\varepsilon) = ||T|| \cdot ||T^{-1}||$$
(5)

characterizes "the quality" of the basis $\{z_k\}_1^\infty$. It is called the Riesz constant.

Let $L = L(\varepsilon)$ be the operator in $L_2(-1, 1)$ associated with model problem (3), (4). It can be easily proved that

$$\|(L-\lambda)^{-1}\| \leq \frac{C(\varepsilon)}{\operatorname{dist}(\lambda,\sigma(L))}$$

where $C(\varepsilon)$ is defined by (5). The question is: how rapidly grows the constant $C(\varepsilon)$?

Theorem 4. Denote by Ω the domain bounded by the curves γ_+, γ_- and the segment [a, b]. Then, given compact K in Ω there are numbers $\tau > 0$ and $\varepsilon_0 > 0$ such that $||(L - \lambda)^{-1}|| \ge e^{\tau/\varepsilon}$ for all $\lambda \in K$ and $\varepsilon < \varepsilon_0$. In particular, the constant $C(\varepsilon)$ defined in (5) grows exponentially, *i.e.* there is a number $\eta > 0$ such that $C(\varepsilon) \ge e^{\eta/\varepsilon}$.

It was proved in [5] that there is no operator in $L_2(-1, 1)$ associated with Orr-Sommerfeld problem. However, there is such an operator in the Sobolev space $\mathring{W}_2^1[-1, 1]$. Denote this operator by S and consider the operator T in l_2 generated by the Gramm matrix $\{(y_k, y_j)\}_{k,j=1}^{\infty}$ assuming that y_k are the normed eigenfunctions of the operator S in the space $\mathring{W}_2^1[-1, 1]$ and the scalar product is taken in the same space.

Theorem 5. The resolvent $(S - \lambda)^{-1}$ and the constant $C(\varepsilon)$ defined by (5) admit the same estimates from below as in Theorem 4.

The estimates from below for the resolvents $(L - \lambda)^{-1}$ and $(S - \lambda)^{-1}$ confirm the following fact: for sufficiently small $\varepsilon > 0$ the pseudospectra of these operators occupies "almost whole" domain Ω . So, the last theorems prove some observations of Trefethen [6] and Reddy, Schmidt and Henningson [7] on the pseudospectra of the Orr–Sommerfeld operator. We remark also that a weaker estimate $C(\varepsilon) \ge \tau \varepsilon^{1/3}$ for the Riesz constant was obtained by Stepin [8].

References

- [1] Dyachenko A. V., Shkalikov A. A. On a model problem for the Orr–Sommerfeld equation with linear profile // Funct. Analysis and its Appl., **36**, No 3 (2002), P. 71–75.
- Shkalikov A. A. The limit behaviour of the spectrum for large parameter values in a model problem // Math. Zametki, 62 (1997), P. 950–953 (in russian); English transl.: Math. Notes, 62 (1997), P. 796–799.
- [3] Shkalikov A. A. Quasi-classical eigenvalue distribution for a non-selfadjoint Sturm-Liouville operators. In book: Spectral Analysis of Differential and Difference operators. Stephan Banach Intern. Math. Center. Warsaw, 2001, P. 37–40.
- [4] Shkalikov A. A. Theorems of Tauberian type on the zero distribution of holomorphic functions // Math. Sborn., 125, No 3 (1984), P. 317–347.
- [5] Shkalikov A. A. How to Define the Orr-Sommerfeld Operator? //Moscow University Math.Bulletin, 1998, No 4, P. 65–75.
- [6] Trefethen L. N. Pseudospectra of Linear Operators, in: ISIAM 95: Proceedings of the Third Int. Congress of Industrial and Applied Math., Academic Verlag, Berlin, 1996. P. 401–434.
- [7] Reddy S. G., Schmidt P. J. and Henningson D. S., Pseudospectra of the Orr-Sommerfeld operator // SIAM J.Appl.Math. 53 (1993) No 1, P. 15–47.
- [8] Stepin S. A. Non-selfadjoint Singular Perturbations and Spectral Properties of the Orr-Sommerfeld Problem // Math. Sbornik 188 (1997) No 1, P. 129–146.