MAXIMAL DIVISORIAL IDEALS AND t-MAXIMAL IDEALS

STEFANIA GABELLI AND MOSHE ROITMAN

ABSTRACT. We give conditions for a maximal divisorial ideal to be *t*-maximal and show with examples that, even in a completely integrally closed domain, maximal divisorial ideals need not be *t*-maximal.

INTRODUCTION

The v-operation and the t-operation are the best known and most useful star operations; mainly because the structure of certain semigroups of t-ideals reflects the multiplicative properties of an integral domain. In this context an important role is played by the prime and the maximal v- and t-ideals.

Since the t-operation is a star operation of finite type, a domain R has always t-maximal ideals. On the other hand, the set of v-maximal ideals may be empty.

In this paper we deal with the following question:

Assume that M is a v-maximal ideal of R, is M necessarily a t-maximal ideal?

We show that although the answer is positive in a large class of domains, namely in the class of v-coherent domains, it is negative in general. In fact we give two examples of a v-maximal ideal P that is not a t-maximal ideal. In the first example P is an upper to zero of a completely integrally closed polynomial ring, thus P is v-invertible. In the second example P is a strongly divisorial ideal of an integrally closed semigroup ring.

1. Preliminaries and notations

Throughout this paper R will denote an integral domain with quotient field K. We will refer to a fractional ideal as an *ideal* and will call a fractional ideal contained in R an *integral ideal*.

¹⁹⁹¹ Mathematics Subject Classification. Primary: 13C13; Secondary: 13G05. Key words and phrases. divisorial ideal, star operation, t-ideal.

We recall that a star operation is an application $I \to I^*$ from the set F(R) of nonzero ideals of R to itself such that:

(1) $R^* = R$ and $(aI)^* = aI^*$, for all $a \in K \setminus \{0\}$;

(2) $I \subseteq I^*$ and $I \subseteq J \Rightarrow I^* \subseteq J^*$;

(3) $I^{**} = I^*$.

General references for systems of ideals and star operations are [13, 16, 17, 22].

We denote by f(R) the set of nonzero finitely generated ideals of R. A star operation * is of *finite type* if, for each $I \in F(R)$, $I^* = \bigcup \{J^* \mid J \subseteq I \text{ and } J \in f(R)\}$. To any star operation *, we can associate a star operation $*_f$ of finite type by defining $I^{*_f} = \bigcup \{J^* \mid J \subseteq I \text{ and } J \in f(R)\}$. Clearly $I^{*_f} \subseteq I^*$.

The v- and the t-operations are particular star operations, defined in the following way.

For a pair of nonzero ideals I and J of a domain R we let (J : I)denote the set $\{x \in K | xI \subseteq J\}$ and $(J :_R I)$ denote the set $\{x \in R | xI \subseteq J\}$. We set $I_v = (R : (R : I))$ and $I_t = \bigcup J_v$ with the union taken over all finitely generated ideals J contained in I.

The t-operation is the finite type star operation associated to the v-operation.

A nonzero ideal I is called a *-*ideal* if $I = I^*$. Thus a nonzero ideal I is a *v*-*ideal*, or is *divisorial*, if $I = I_v$, and it is a *t*-*ideal* if $I = I_t$. Note that I is a *t*-ideal if and only if $J_v \subseteq I$ whenever J is finitely generated and $J \subseteq I$.

The set $F_*(R)$ of *-ideals of R is a semigroup with respect to the *-multiplication, defined by $(I, J) \to (IJ)^*$, with unity R.

We say that an ideal $I \in F(R)$ is *-*invertible* if I^* is a unit in the semigroup $F_*(R)$. In this case the *-*inverse* of I is (R : I). Thus I is *-invertible if and only if $(I(R : I))^* = R$. Invertible ideals are (*-invertible) *-ideals.

We have $I \subseteq I^* \subseteq I_v$, so that any divisorial ideal is a *-ideal and any *-invertible ideal is *v*-invertible. In particular a divisorial ideal is a *t*-ideal and a *t*-invertible ideal is *v*-invertible.

A nonzero ideal I is *-finite if $I^* = J^*$ for some finitely generated ideal J. Since the v- and the t-operation coincide on finitely generated ideals and since $I_t = J_t$ implies $I_v = J_v$, an ideal I is t-finite if and only if $I_v = J_v$ (equivalently (R : I) = (R : J)) for some finitely generated ideal $J \subseteq I$. It follows that the set $f_v(R)$ of the v-finite divisorial ideals coincides with the set of the t-finite t-ideals. $f_v(R)$ is a sub-semigroup of $F_v(R)$. An ideal I is t-invertible if and only if it is v-invertible and both I and (R : I) are t-finite. Hence the set of the t-invertible t-ideals of R, here denoted by T(R), is the largest subgroup of $f_v(R)$. The importance of the notion of t-invertibility is well illustrated in [28].

Denoting by Inv(R) the group of the invertible ideals of R, we have

$$Inv(R) \subseteq T(R) \subseteq f_v(R) \subseteq F_v(R) \subseteq F_t(R) \subseteq F(R)$$

and

$$Inv(R) \subseteq f(R).$$

Several important classes of domains may be characterized by the fact that some of these inclusions are equalities. For example R is a *Prüfer* domain if and only if Inv(R) = f(R) [13], it is a *Krull* domain if and only if $T(R) = F_t(R)$ [24] and it is a *Prüfer v-multiplication* domain, for short a *PvMD*, if and only if $T(R) = f_v(R)$ [16]. A *Mori* domain is a domain satisfying the ascending chain condition on integral divisorial ideals and has the property that $f_v(R) = F_t(R)$. Noetherian and Krull domains are Mori. A recent reference for Mori domains is [1]. The class of domains with the property that $F_v(R) = F(R)$ have been studied by several authors [3, 18, 26, 4, 5]. A domain such that $F_v(R) = F_t(R)$ is called in [20] a *TV*-domain. Examples of *TV*-domains are given in [20, 23, 7]. Mori and pseudovaluations domains which are not valuation domains are *TV*-domains.

2. When a maximal divisorial ideal is t-maximal

A prime *-ideal is called a *-prime. A *-maximal ideal is an ideal that is maximal in the set of the proper integral *-ideals. A v-maximal ideal is also called a maximal divisorial ideal. A *-maximal ideal is a prime ideal (if it exists).

If * is a star operation of finite type, an easy application of the Zorn Lemma shows that the set *-Max(R) of the *-maximal ideals of R is not empty. Moreover, for each $I \in F(R)$, $I^* = \bigcap_{M \in *-\text{Max}(R)} I^* R_M$ [16]. In particular the set of the *t*-maximal ideals is not empty and $I_t = \bigcap_{M \in t-\text{Max}(R)} I_t R_M$. On the contrary, the set of maximal divisorial ideals may be empty, like for example when R is a rank-one nondiscrete valuation domain.

If M is a *-maximal ideal that is not *-invertible, then $M = (M(R : M))^*$ and so (M : M) = (R : M). An ideal I with the property that (R : I) = (I : I) is called *strong*. A strong ideal is never *-invertible and we have just seen that a *-maximal ideal is either *-invertible or strong.

An ideal which is strong and divisorial is called *strongly divisorial*.

Proposition 2.1.

- (1) If M is a maximal divisorial ideal of R, then $M = x^{-1}R \cap R$, for some element $x \in K$. Hence $(R:M) = (R + xR)_v$.
- (2) If P is a prime divisorial ideal of R such that (R : P) = R + xR, for some element $x \in K$, then P is maximal divisorial.

Proof. (1) If $x \in (R:M) \setminus R$, then $M \subseteq x^{-1}R$ and $R \not\subseteq x^{-1}R$. Since an intersection of divisorial ideals is divisorial and M is v-maximal, we have $M = x^{-1}R \cap R = (R:R+xR)$.

(2) Let Q be a proper divisorial ideal containing P. Since Q is divisorial, $(R : Q) \notin R$. Since $(R : Q) \subseteq (R : P) = R + xR$, we see that there exists an element $y \in R$ such that $xy \in (R : Q) \setminus R$. Thus $y \notin P$, and $xyQ \subseteq R$. Since P = (R : R + xR), we obtain that $yQ \subseteq P$. Since P is a prime ideal, we conclude that $Q \subseteq P$. Hence P is maximal divisorial.

In a Mori domain R, all the prime divisorial ideals are of the form $x^{-1}R \cap R = (R: R + xR)$ [19, Corollary 2.5].

A domain has the property that each t-maximal ideal is divisorial if and only if every ideal I such that (R : I) = R is t-finite [20, Proposition 2.4]. A domain of this type is called an H-domain in [15]. A TV-domain is clearly an H-domain, but the converse is not true [20, 2].

The following proposition gives conditions for a divisorial prime ideal to be a *t*-maximal ideal. A proof can be found in [10].

Proposition 2.2.

- (1) A v-invertible divisorial prime is maximal divisorial;
- (2) A v-finite maximal divisorial ideal is t-maximal;
- (3) A v-finite v-invertible divisorial prime is t-invertible;
- (4) A t-invertible t-prime is t-maximal.

We remark that in general a *-invertible *-prime need not be *maximal (for example a principal prime ideal is not necessarily a maximal ideal) and that a (non-prime) v-finite v-invertible divisorial ideal need not be t-invertible [6].

Corollary 2.3. Assume that each maximal divisorial ideal of R is a t-maximal ideal. Then each v-invertible divisorial prime is a t-invertible t-maximal ideal.

Proof. Let P be a v-invertible divisorial prime. By Proposition 2.2, P is maximal divisorial and so t-maximal. Since P is not strong, then it is t-invertible.

In general, if each v-invertible divisorial prime of R is t-invertible, it is not true that each v-invertible ideal is t-invertible. This last property

is in fact equivalent to R being an H-domain [28, Proposition 4.2]. The ring of entire functions is not an H-domain, but all its divisorial primes are t-invertible (see for example [10, Section 2]).

A *v*-coherent domain is a domain R with the property that, for each finitely generated ideal J, the ideal (R : J) is *v*-finite. This class of domains was first studied (under a different name) in [25] and is very large, properly including PvMD's, Mori domains and coherent domains [25, 11]. (A domain is coherent if each finitely generated ideal is finitely presented, or, equivalently, if the intersection of each pair of finitely generated ideals is finitely generated.)

Proposition 2.4. If R is v-coherent, then each maximal divisorial ideal is t-maximal.

Proof. Let M be a maximal divisorial ideal of R. Then $M = x^{-1}R \cap R = (R : R + xR)$ for some $x \in K$ (Proposition 2.1). Since R is v-coherent, then M is v-finite and so t-maximal by Proposition 2.2. \Box

A domain R is completely integrally closed if and only if $F_v(R)$ is a group under v-multiplication [13]. If $F_v(R) = T(R)$, then R is a completely integrally closed H-domain, equivalently a Krull domain [10, 15].

A divisorial prime of a completely integrally closed domain, being v-invertible, is always maximal divisorial by Proposition 2.2. We will see in the next section that it need not be t-maximal. As a matter of fact, a divisorial prime P of a completely integrally closed domain has height one and P is t-maximal if and only if it is v-finite, if and only if it is t-invertible [10, Theorem 2.3].

A completely integrally closed v-coherent domain is a (completely integrally closed) PvMD. In this case each divisorial prime is t-maximal by Corollary 2.3.

We now turn to the case of polynomial rings.

We denote by \mathbf{X} a set of independent indeterminates over R and by $R[\mathbf{X}]$ the polynomial ring in this set of indeterminates.

It is well known that the correspondence $I \mapsto I[\mathbf{X}]$ induces inclusion preserving injective maps $t(R) \longrightarrow t(R[\mathbf{X}])$ and $D(R) \longrightarrow D(R[\mathbf{X}])$. Moreover, M is a t-maximal ideal, respectively a maximal divisorial ideal, of $R[\mathbf{X}]$ such that $M \cap R \neq (0)$, if and only if $M = (M \cap R)[\mathbf{X}]$ and $M \cap R$ is a t-maximal ideal, respectively a maximal divisorial ideal, of R (see for example [8, Lemma 2.1] and [27, Theorem 3.6]).

Thus, if each maximal divisorial ideal of $R[\mathbf{X}]$ is t-maximal, R has the same property.

On the other hand, Example 3.1 in the next section shows that if $M \cap R = (0)$, then M may be maximal divisorial but not t-maximal.

A prime ideal Q of $R[\mathbf{X}]$ such that $Q \cap R = (0)$ is called an *upper to* zero. Q is an upper to zero of height one if and only if $Q = fK[\mathbf{X}] \cap R[\mathbf{X}]$ for some polynomial $f \in R[\mathbf{X}]$, irreducible in $K[\mathbf{X}]$ [12, Lemma 2.1]. In one indeterminate, all the uppers to zero are of this form.

Recall that if R is integrally closed and f is a nonzero polynomial of $R[\mathbf{X}]$, then $fK[\mathbf{X}] \cap R[\mathbf{X}] = f(R : c(f))[\mathbf{X}]$ [13, Corollary 34.9]. (Here c(f) denotes the *content* of f, that is the fractional ideal of Rgenerated by the coefficients of f.) Hence if R is integrally closed, an upper to zero of height one is always divisorial and if R is completely integrally closed, an upper to zero of height one, being v-invertible, is always maximal divisorial.

In general, an upper to zero is t-maximal if and only if it is t-invertible; in this case it has height one [12, Section 3]. We now show that a similar result holds for the v-operation.

Proposition 2.5. A divisorial upper to zero is a maximal divisorial ideal if and only if it is v-invertible. In this case it has height one.

Proof. A divisorial v-invertible prime is always maximal divisorial (Proposition 2.2 (1)).

Conversely, let $P \subseteq R[\mathbf{X}]$ be an upper to zero that is maximal divisorial. Then $P = \frac{f}{g}R[\mathbf{X}] \cap R[\mathbf{X}] \subseteq fK[\mathbf{X}] \cap R[\mathbf{X}]$, for some $f, g \in R[\mathbf{X}]$, $g \neq 0$ (Proposition 2.1(1)). Since $P \cap R = (0)$ and $f = \frac{f}{g}g \in P$, then $f \notin R$. We may also assume that f and g are coprime in $K[\mathbf{X}]$.

Let $h = f\alpha \in fK[\mathbf{X}] \cap R[\mathbf{X}]$, with $\alpha \in K[\mathbf{X}]$. There is a nonzero $c \in R$ such that $c\alpha \in R[\mathbf{X}]$. Hence $ch = (c\alpha)f = (c\alpha g)\frac{f}{g} \in P$. Since $c \notin P$, then $h \in P$.

We conclude that $P = fK[\mathbf{X}] \cap R[\mathbf{X}]$ has height one.

In addition, $\frac{g}{f} \in (R[\mathbf{X}] : P)$, but $\frac{g}{f} \notin (P : P)$. Otherwise $g = \frac{g}{f}f \in P$ and so $g = \frac{f}{g}t$ for some $t \in R[\mathbf{X}]$. Then f divides g^2 in $K[\mathbf{X}]$, which is impossible, because f and g are coprime and $f \notin K$.

It follows that P is not strong and, being maximal divisorial, is v-invertible.

The following result was proved in [15] for one indeterminate.

Proposition 2.6. R is an H-domain if and only if $R[\mathbf{X}]$ is an H-domain.

Proof. An extended prime $P[\mathbf{X}]$ is a *t*-maximal ideal, respectively a maximal divisorial ideal, if and only if so is P, [8, Lemma 2.1] and

[27, Theorem 3.6]. A *t*-maximal upper to zero is *t*-invertible by [12, Theorem 2.3]. Hence it is divisorial. \Box

The domain R is said to be a UMT-domain if every upper to zero of R[X] is a t-maximal ideal [21]. This property is stable under polynomial extensions, in fact R is a UMT-domain if and only if $R[\mathbf{X}]$ is a UMT-domain [8, Theorem 2.4]. The integrally closed UMT-domains are exactly the PvMDs [21, Proposition 3.2].

The following proposition is immediate.

Proposition 2.7. Assume that R is an UMT-domain. Then each maximal divisorial ideal of R is t-maximal if and only if $R[\mathbf{X}]$ has the same property.

We conclude this section recalling that it is not known whether R*v*-coherent implies that R[X] is *v*-coherent. This is true under the additional hypothesis that R is integrally closed [25]. In this case, each prime of R[X] upper to zero is divisorial *v*-finite. When R is *v*-coherent and completely integrally closed (thus a completely integrally closed PvMD), each upper to zero of R[X] is *t*-maximal (and *t*-invertible).

3. Maximal divisorial ideals that are not t-maximal

In this section we give two examples of a maximal divisorial ideal P of an integral domain R that is not a *t*-maximal ideal. In the first example R is a completely integrally closed polynomial ring in one indeterminate and P is an upper to zero, thus P is *v*-invertible. In the second example R is an integrally closed semigroup ring and P is strongly divisorial.

Example 3.1. An upper to zero P of a completely integrally closed polynomial ring R[X] that is maximal divisorial but not t-maximal. P is necessarily v-invertible.

Let y, z and $\mathbf{t} = \{t_n (n \ge 1)\}$ be independent indeterminates over a field k. Let S be the semigroup of monomials f of $k[y, z, \mathbf{t}]$ satisfying the conditions $\deg_{y,z} f \ge \deg_{t_n} f$ for all $n \ge 1$, and let R = k[S] the semigroup ring over k generated by S.

Set

$$P = (y + zX)K[X] \cap R[X],$$

where K is the field of fractions of R and X is an indeterminate over R. Then R (and so also R[X]) is completely integrally closed, and P is a maximal divisorial ideal of R[X] that is not t-maximal.

Proof.

(1) R[X] is completely integrally closed.

It is enough to show that R is completely integrally closed. Since R = k[S] is a semigroup ring over the field k, by [14, Corollary 12.7 (2)] to this end it suffices to show that the semigroup S is completely integrally closed.

Let $u, v, w \in S$ so that $u(\frac{v}{w})^m \in S$ for all $m \ge 1$. Fix $n \ge 1$. Then $\deg_{y,z}(u(\frac{v}{w})^m) \ge \deg_{t_n}(u(\frac{v}{w})^m)$ for all m. Hence

$$\deg_{y,z} u + m \deg_{y,z}(\frac{v}{w}) \ge \deg_{t_n} u + m \deg_{t_n}(\frac{v}{w}).$$

Divide by m and let m go to ∞ to obtain that $\deg_{y,z}(\frac{v}{w}) \geq \deg_{t_n}(\frac{v}{w})$. The same argument shows that $\frac{v}{w}$ is a monomial, that is has a nonnegative degree in each indeterminate. It follows that $\frac{v}{w} \in S$; thus S is completely integrally closed.

(2) P is an upper to zero of R[X] that is a v-invertible maximal divisorial ideal.

P is clearly an upper to zero. Since *R* is integrally closed, then P = (y+zX)(R:(y,z))[X] by [13, Corollary 34.9], hence *P* is divisorial. But R[X] is completely integrally closed; thus *P* is *v*-invertible and so maximal divisorial (Proposition 2.2).

(3) P is not t-maximal.

Let $Q = (y, z)k[y, z, \mathbf{t}] \cap R$. Then QR[X] is a proper t-ideal of R[X] properly containing P.

To verify this, let F be a finite subset of QR[X]. Let t_n be an indeterminate that does not occur in the polynomials in the set F. Then $t_n f \in R[X]$ for all $f \in F$, so $t_n \in (R[X] : F)$. If $g \in (F)_v$, then $gt_n \in R[X]$. Hence $\deg_{y,z} gt_n \ge 1$ and $g \in Q$. It follows that $(F)_v \subseteq Q$, so Q is a t-ideal.

Example 3.2. An example of a strong maximal divisorial ideal of an integrally closed domain R that is not t-maximal.

Let k be a field and let $Y, Z, \mathbf{X} = \{X_n : n \ge 1\}, \mathbf{T} = \{T_n : n \ge 1\}$ be independent indeterminates over k. Let S be the set of monomials f in $k[Y, Z, \mathbf{X}, \mathbf{T}]$ satisfying the following two conditions:

- (a) If Z occurs in f, then some X_n occurs in f.
- (b) For all n, if T_n occurs in f, then either Y or X_i occurs in f for some $i \leq n$.

Clearly, S is a semigroup containing **X** and Y. Let R = k[S] be the semigroup ring over S and set

$$P = (\mathbf{X})k[Y, Z, \mathbf{X}, \mathbf{T}] \cap R.$$

8

Then R is integrally closed and P is a strong maximal divisorial ideal of R that is not t-maximal.

- *Proof.* We will use repeatedly that P is a monomial ideal of R.
 - (1) R is integrally closed.

By [14, Corollary 12.11 (2)], it is enough to show that the monoid S is integrally closed. If f is an element in the quotient group of S such that $f^n \in S$ for some $n \ge 1$, then f is a monomial. Since f^n satisfies conditions (a)-(b), it is clear that f also satisfies them, thus $f \in S$. We conclude that R is integrally closed.

(2) $P = RZ^{-1} \cap R$. Hence P is a divisorial ideal.

Clearly, any monomial in ZP satisfies conditions (a)-(b), hence $ZP \subseteq R$. Thus $P \subseteq RZ^{-1} \cap R$.

For the reverse inclusion, it is enough to show that any monomial $f \in RZ^{-1}$ belongs to P. Since $Zf \in R$, we see that Zfsatisfies conditions (a)-(b) and so does f, thus $f \in R$. Using again that $Zf \in R$ we see that some X_n occurs in f, hence $f \in P$.

(3) (R:P) = R[Z].

Using conditions (a)-(b), we see that $R[Z] \subseteq (R:P)$.

For the reverse inclusion, let u be a quotient of monomials in (R : P). Since $uX_1, uX_2 \in R$, we see that uX_1 and uX_2 are monomials, hence, by factoriality, u also is a monomial. Let $u = Z^k u_0$, where $k \ge 0$, u_0 is a monomial and Z does not occur in u_0 . Choose a positive integer N such that N > i for all T_i 's occurring in u. Since $Z^k u_0 X_N \in R$, we see that u_0 satisfies condition (b); hence $u_0 \in R$, so $u \in R[Z]$.

(4) P is a strong maximal divisorial ideal.

We have $(R:P) = R[Z] \subseteq (P:P)$, thus (R:P) = (P:P), that is, P is strong.

Assume that P is not maximal divisorial, so there is a divisorial ideal Q properly containing P. Let $f \in Q \setminus P$. We may assume that no X_n occurs in f, thus Z does not occur in f either by condition (a) above. Let $g \in (R : Q) \setminus R$, thus $g \in (R : P) = R[Z], g = \sum_{i=0}^{n} a_i Z^i$, where $a_0, \ldots, a_n \in R$. We may assume that $a_n Z^n \notin R$, thus $n \ge 1$. We also may assume that no X_i occurs in a_n . Thus no X_i occurs in fa_n , which implies that $fa_n Z^n \notin R$. Since R = k[S], we obtain that $fg = fa_n Z^n + \cdots \notin R$, a contradiction.

(5) The ideal M = (S)R is a maximal ideal of R properly containing P and is a t-ideal.

Clearly M is a maximal ideal containing P. Since $Y \in M \setminus P$, we have $P \subsetneq M$.

To show that M is a *t*-ideal, let F be a finite subset of Mand let N be a positive integer such that N > i for each T_i occurring in some element of F. From conditions (a)-(b) it follows that $M \subseteq (\mathbf{X}, Y)k[Y, Z, \mathbf{X}, \mathbf{T}]$. Hence $T_N F \subseteq R$. Thus $(F)_v \subseteq (R : T_N) \cap R$. Since $T_N \notin R$ and since $(R : T_N) \cap R$ is a monomial ideal, we obtain that $(R : T_N) \cap R \subseteq (S)R = M$. It follows that $(F)_v \subseteq M$ and that M is a t-ideal.

References

- V. Barucci, Mori domains, Non-Noetherian commutative ring theory, 57-74, Math. Appl., 520, Kluwer Acad. Publ., Dordrecht, 2000.
- [2] V. Barucci, S. Gabelli and M. Roitman, On semi-Krull domains, J. Algebra 145 (1992), 306-328.
- [3] H. Bass, On the ubiquity of Gorenstein rings, Math. Z. 82 (1963), 8-28.
- [4] S. Bazzoni, Divisorial domains, Forum Math. 12 (2000), 397-419.
- [5] S. Bazzoni and L. Salce, Warfield domains, J. Algebra 185 (1996), 836-868.
- [6] J. Dieudonné, Sur la théorie de la divisibilité, Bull. Soc. Math. France 69 (1941), 133-134.
- [7] Chul Ju Hwan and Gyu Whan Chang, Prüfer v-multiplication domains in which each t-ideal is divisorial, Bull. Korean Math. Soc. 35 (1998), 259-268.
- [8] M. Fontana, S. Gabelli and E. Houston, UMT-domains and domains with Prüfer integral closure, Comm. Algebra 26 (1998) 1017-1039.
- [9] M. Fontana, J. A. Huckaba and Ira J. Papick. Prüfer domains, M. Dekker, 1997.
- [10] S. Gabelli, Completely integrally closed domains and t-ideals, Bollettino U.M.I. 3-B (1989), 327-342.
- [11] S. Gabelli and H. Houston, Coherentlike conditions in pullbacks, Michigan Math. J. 44 (1997), 99-123.
- [12] S. Gabelli, H. Houston and T. Lucas, The t# property for integral domains, manuscript.
- [13] R. Gilmer, Multiplicative Ideal Theory, M. Dekker, 1972.
- [14] R. Gilmer, Commutative semigroup rings, The University of Chicago Press, 1984.
- [15] S. Glaz and W. Vasconcelos, Flat ideals II, Manuscripta Math. 22 (1977), 325-341.
- [16] M. Griffin, Some results on v-multiplication rings, Canad. J. Math. 19 (1967), 710-722.
- [17] F. Halter-Koch, Ideal systems. An introduction to multiplicative ideal theory, Monographs and Textbooks in Pure and Applied Mathematics, 211, M. Dekker, 1998.
- [18] W. Heinzer, Integral domains in which each nonzero ideal is divisorial, Mathematika 15 (1968), 164-170.

- [19] E. G. Houston, T. G. Lucas and T. M. Viswanathan, Primary decomposition of divisorial ideals in Mori domains, J. Algebra 117 (1988), 327-342.
- [20] E. G. Houston and M. Zafrullah, Integral domains in which each t-ideal is divisorial, Michigan Math. J. 35 (1988), 291-300.
- [21] E. G. Houston and M. Zafrullah, On t-invertibility II, Comm. Algebra 17 (1989), 1955-1969.
- [22] P. Jaffard, Les Systèmes d'Idéaux, Dunod, Paris, 1970.
- [23] B. G. Kang, Some questions about Prüfer v-multiplication domains. Comm. Algebra 17 (1989), 553–564.
- [24] B. G. Kang, On the converse of a well-known fact about Krull domains. J. Algebra 124 (1989), 284–299.
- [25] D. Nour el Abidine, Groupe des classes de certain anneaux intégres et idéaux transformés, Thèse de Doctorat, Lyon, 1992.
- [26] E. Matlis, Reflexive domains, J. Algebra 8 (1968), 1-33.
- [27] M. Roitman, On Mori domains and commutative rings with CC^{\perp} I, J. Pure Applied Algebra 56 (1989), 247-268.
- [28] M. Zafrullah, Putting t-invertibility to use, Non-Noetherian commutative ring theory, 429-457, Math. Appl., 520, Kluwer Acad. Publ., Dordrecht, 2000.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI ROMA TRE, LARGO S. L. MURIALDO, 1, 00146 ROMA, ITALY

E-mail address: gabelli@mat.uniroma3.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, MOUNT CARMEL, HAIFA 31905, ISRAEL

E-mail address: mroitman@math.haifa.ac.il