

**NORM ESTIMATES FOR THE DIFFERENCE BETWEEN
BOCHNER'S INTEGRAL AND THE CONVEX COMBINATION
OF FUNCTION'S VALUES**

P. CERONE, Y.J. CHO \star , S.S. DRAGOMIR, J.K. KIM, AND S.S. KIM \diamond

ABSTRACT. Norm estimates are developed between the Bochner integral of a vector-valued function in Banach spaces having the Radon-Nikodym property and the convex combination of function values taken on a division of the interval $[a, b]$.

1. INTRODUCTION

A Banach space X with the property that every absolutely continuous X -valued function is almost everywhere differentiable is said to be a *Radon-Nikodym* space [7, pp. 217–219] or [2, 13] (see also [3]). For example, every reflexive Banach space (in particular, every Hilbert space) is a Radon-Nikodym space, but the space $L_\infty[0, 1]$ of all \mathbb{K} -valued, essentially bounded functions defined on the interval $[0, 1]$, endowed with the norm

$$\|g\|_\infty := \text{ess} \sup_{t \in [0, 1]} |g(t)|,$$

is a Banach space which is not a Radon-Nikodym space.

A function $f : [a, b] \rightarrow X$ is *measurable* if there exists a sequence of simple functions (f_n) (with $f_n : [a, b] \rightarrow X$) which converges punctually a.e. on $[a, b]$.

It is well-known that a measurable function $f : [a, b] \rightarrow X$ is Bochner integrable if and only if its norm, that is, the function $t \mapsto \|f\|(t) := \|f(t)\| : [a, b] \rightarrow X$ is Lebesgue integrable on $[a, b]$, (see for example [12]). The Bochner integral of f shall be represented by $(B) \int_a^b f$.

Further, we use the integration by parts formula, which holds under the following general conditions:

Let $-\infty < a < b < \infty$ and f, g be two mappings defined on $[a, b]$ such that f is C -valued and g is X -valued, where X is a real or complex Banach space. If f, g are differentiable on $[a, b]$ and their derivatives are Bochner integrable on $[a, b]$, then

$$(B) \int_a^b f'g = f(b)g(b) - f(a)g(b) - (B) \int_a^b fg'.$$

For some results on the Ostrowski inequality for real-valued functions, see [1], [5], [10] and [11], and the references therein.

The following theorem concerning a version of Ostrowski's inequality for vector-valued functions has been obtained in [3].

Date: July 16, 2003.

1991 *Mathematics Subject Classification.* Primary 26D15; Secondary 41A55.

Key words and phrases. Bochner's Integral, Ostrowski Inequality, Quadrature Formulae
 \star, \diamond Corresponding authors.

Theorem 1. Let $(X; \|\cdot\|)$ be a Banach space with the Radon-Nikodym property and $f : [a, b] \rightarrow X$ an absolutely continuous function on $[a, b]$ with the property that $f' \in L_\infty([a, b]; X)$, i.e.,

$$\|f'\|_{[a,b],\infty} := \text{ess sup}_{t \in [a,b]} \|f'(t)\| < \infty.$$

Then we have the inequalities:

$$\begin{aligned}
 (1.1) \quad & \left\| f(s) - \frac{1}{b-a} (B) \int_a^b f(t) dt \right\| \\
 & \leq \frac{1}{b-a} \left[\int_a^s (t-a) \|f'(t)\| dt + \int_s^b (b-t) \|f'(t)\| dt \right] \\
 & \leq \frac{1}{2(b-a)} \left[(s-a)^2 \|f'\|_{[a,s],\infty} + (b-s)^2 \|f'\|_{[s,b],\infty} \right] \\
 & \leq \left[\frac{1}{4} + \left(\frac{s-\frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) \|f'\|_{[a,b],\infty} \\
 & \leq \frac{1}{2} (b-a) \|f'\|_{[a,b],\infty};
 \end{aligned}$$

for any $s \in [a, b]$, where $(B) \int_a^b f(t) dt$ is the Bochner integral of f .

Bounds involving the p -norms, $p \in [1, \infty)$, of the derivative f' , are embodied in the following theorem [3].

Theorem 2. Let $(X, \|\cdot\|)$ be a Banach space with the Radon-Nikodym property and $f : [a, b] \rightarrow X$ be an absolutely continuous function on $[a, b]$ with the property that $f' \in L_p([a, b]; X)$, $p \in [1, \infty)$, i.e.,

$$(1.2) \quad \|f'\|_{[a,b],p} := \left(\int_a^b \|f'(t)\|^p dt \right)^{\frac{1}{p}} < \infty.$$

Then we have the inequalities:

$$\begin{aligned}
 (1.3) \quad & \left\| f(s) - \frac{1}{b-a} (B) \int_a^b f(t) dt \right\| \\
 & \leq \frac{1}{b-a} \left[\int_a^s (t-a) \|f'(t)\| dt + \int_s^b (b-t) \|f'(t)\| dt \right]
 \end{aligned}$$

$$\begin{aligned}
&\leq \begin{cases} \frac{1}{b-a} \left[(s-a) \|f'\|_{[a,s],1} + (b-s) \|f'\|_{[s,b],1} \right] & \text{if } f' \in L_1([a,b];X); \\ \frac{1}{(b-a)(q+1)^{\frac{1}{q}}} \left[(s-a)^{\frac{1}{q}+1} \|f'\|_{[a,s],p} + (b-s)^{\frac{1}{q}+1} \|f'\|_{[s,b],p} \right] & \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1 \text{ and } f' \in L_p([a,b];X); \end{cases} \\
&\leq \begin{cases} \left[\frac{1}{2} + \left| \frac{s - \frac{a+b}{2}}{b-a} \right| \right] \|f'\|_{[a,b],1} & \text{if } f' \in L_1([a,b];X); \\ \frac{1}{(q+1)^{\frac{1}{q}}} \left[\left(\frac{s-a}{b-a} \right)^{q+1} + \left(\frac{b-s}{b-a} \right)^{q+1} \right]^{\frac{1}{q}} (b-a)^{\frac{1}{q}} \|f'\|_{[a,b],p} & \text{if } f' \in L_p([a,b];X). \end{cases}
\end{aligned}$$

The main aim of this paper is to point out estimates between the Bochner integral of a vector-valued function, with values in Banach spaces having the Radon-Nikodym property and a convex combination of values taken on a given division of the interval $[a, b]$. The obtained results naturally extend the Ostrowski type inequalities mentioned above. Some particular cases for two and three points rules are also given.

2. THE RESULTS

Let $a \leq b$ and $c \in \mathbb{R}$. Define the mapping

$$(2.1) \quad \mu_p(a, c, b) := \begin{cases} \int_a^b |t - c|^p dt & \text{if } p \in [1, \infty); \\ \max_{t \in [a, b]} |t - c| & \text{if } p = \infty. \end{cases}$$

We observe that:

(1) If $c < a$, then

$$\begin{aligned}
\mu_p(a, c, b) &= \int_a^b (t - c)^p dt \\
&= \frac{1}{p+1} \left[(b-c)^{p+1} - (a-c)^{p+1} \right], \quad \text{for } p \in [1, \infty)
\end{aligned}$$

and

$$\mu_\infty(a, c, b) = b - c.$$

(2) If $c \in [a, b]$, then

$$\begin{aligned}
\mu_p(a, c, b) &= \int_a^c (c-t)^p dt + \int_c^b (t-c)^p dt \\
&= \frac{1}{p+1} \left[(c-a)^{p+1} + (b-c)^{p+1} \right]
\end{aligned}$$

for $p \in [1, \infty)$ and

$$\mu_\infty(a, c, b) = \max(c-a, b-c) = \frac{1}{2}(b-a) + \left| c - \frac{a+b}{2} \right|.$$

(3) If $b < c$, then

$$\begin{aligned}\mu_p(a, c, b) &= \int_a^b (c-t)^p dt \\ &= \frac{1}{p+1} [(c-a)^{p+1} - (c-b)^{p+1}], \quad \text{for } p \in [1, \infty)\end{aligned}$$

and

$$\mu_\infty(a, c, b) = c - a.$$

Consequently, we may conclude that

$$\mu_p(a, c, b) = \begin{cases} \frac{1}{p+1} [(b-c)^{p+1} - (a-c)^{p+1}] & \text{if } c < a; \\ \frac{1}{p+1} [(c-a)^{p+1} + (b-c)^{p+1}] & \text{if } c \in [a, b]; \\ \frac{1}{p+1} [(c-a)^{p+1} - (c-b)^{p+1}] & \text{if } b < c; \end{cases}$$

for $p \in [1, \infty)$ and

$$\mu_\infty(a, c, b) = \begin{cases} b - c & \text{if } c < a; \\ \frac{1}{2}(b-a) + |c - \frac{a+b}{2}| & \text{if } c \in [a, b]; \\ c - a & \text{if } b < c; \end{cases}$$

where $\mu_s(a, c, b), s \in [1, \infty]$ is as defined in (2.1).

The following integral identity is of interest.

Lemma 1. *Let $f : [a, b] \rightarrow X$ be an absolutely continuous function on the Banach space X , X is with the property of Radon-Nikodym, $a \leq x_1 \leq \dots \leq x_{n-1} \leq x_n \leq b$ and $p_i > 0$ ($i = 1, \dots, n$) with $\sum_{i=1}^n p_i = 1$. Then we have the identity:*

$$\begin{aligned}(2.2) \quad \sum_{i=1}^n p_i f(x_i) - \frac{1}{b-a} (B) \int_a^b f(t) dt &= \frac{1}{b-a} (B) \int_a^{x_1} (t-a) f'(t) dt \\ &\quad + \frac{1}{b-a} \sum_{i=1}^{n-1} (B) \int_{x_i}^{x_{i+1}} [t - (P_i b + \bar{P}_i a)] f'(t) dt \\ &\quad + \frac{1}{b-a} (B) \int_{x_n}^b (t-b) f'(t) dt,\end{aligned}$$

where $(B) \int_a^b f(t) dt$ is the Bochner integral, $P_i := \sum_{k=1}^i p_k$ and $\bar{P}_i = 1 - P_i$.

The sum in the middle is assumed to be zero when $n = 1$.

Proof. We know that, on utilizing the integration by parts formula, for any $x \in [a, b]$, we have the representation (see for example [3])

$$(2.3) \quad f(x) = \frac{1}{b-a} (B) \int_a^b f(t) dt + \frac{1}{b-a} (B) \int_a^b k(x, t) f'(t) dt,$$

where

$$k(x, t) = \begin{cases} t - a & \text{if } a \leq t \leq x \leq b, \\ t - b & \text{if } a \leq x < t \leq b. \end{cases}$$

Putting in (2.3) $x = x_i$ ($i = 1, \dots, n$), multiplying by $p_i \geq 0$ and summing over i from 1 to n , we deduce

$$(2.4) \quad \sum_{i=1}^n p_i f(x_i) = \frac{1}{b-a}(B) \int_a^b f(t) dt + \frac{1}{b-a}(B) \int_a^b \left[\sum_{i=1}^n p_i k(x_i, t) \right] f'(t) dt.$$

However,

$$k(x_1, t) = \begin{cases} t - a & \text{if } a \leq t \leq x_1 \leq b, \\ t - b & \text{if } a \leq x_1 < t \leq b, \end{cases}$$

.....

$$k(x_n, t) = \begin{cases} t - a & \text{if } a \leq t \leq x_n \leq b, \\ t - b & \text{if } a \leq x_n < t \leq b, \end{cases}$$

then

$$(2.5) \quad S(\bar{x}, \bar{p}, t) := \sum_{i=1}^n p_i k(x_i, t)$$

$$= \begin{cases} p_1(t-a) + p_2(t-a) + \dots + p_{n-1}(t-a) + p_n(t-a), & a \leq t \leq x_1 \leq b, \\ p_1(t-b) + p_2(t-b) + \dots + p_{n-1}(t-b) + p_n(t-b), & a \leq x_1 < t \leq x_2 \leq b, \end{cases}$$

.....

$$= \begin{cases} p_1(t-b) + p_2(t-b) + \dots + p_{n-1}(t-b) + p_n(t-b), & a \leq x_{n-1} \leq t \leq x_n \leq b, \\ p_1(t-b) + p_2(t-b) + \dots + p_{n-1}(t-b) + p_n(t-b), & a \leq x_n < t \leq b, \end{cases}$$

$$= \begin{cases} t - a, & a \leq t \leq x_1 \leq b, \\ p_1(t-b) + (p_2 + \dots + p_n)(t-a), & a \leq x_1 < t \leq x_2 \leq b, \end{cases}$$

.....

$$= \begin{cases} (p_1 + \dots + p_{n-1})(t-b) + p_n(t-a), & a \leq x_{n-1} \leq t \leq x_n \leq b, \\ t - b, & a \leq x_n < t \leq b, \end{cases}$$

$$= \begin{cases} t - a, & a \leq t \leq x_1 \leq b, \\ t - [p_1 b + (p_2 + \dots + p_n) a], & a \leq x_1 < t \leq x_2 \leq b, \end{cases}$$

.....

$$= \begin{cases} t - [(p_1 + \dots + p_{n-1}) b + p_n a], & a \leq x_{n-1} \leq t \leq x_n \leq b, \\ t - b, & a \leq x_n < t \leq b, \end{cases}$$

$$= \begin{cases} t - a, & a \leq t \leq x_1 \leq b, \\ t - (P_1 b + \bar{P}_1 a), & a \leq x_1 < t \leq x_2 \leq b, \end{cases}$$

.....

$$= \begin{cases} t - (P_i b + \bar{P}_i a) & a \leq x_i \leq t \leq x_{i+1} \leq b, \\ t - (P_{n-1} b + \bar{P}_{n-1} a), & a \leq x_{n-1} \leq t \leq x_n \leq b, \\ t - b, & a \leq x_n < t \leq b. \end{cases}$$

Consequently, by (2.4) and (2.5), we have

$$\begin{aligned}
(2.6) \quad & \sum_{i=1}^n p_i f(x_i) \\
& = \frac{1}{b-a} (B) \int_a^b f(t) dt + \frac{1}{b-a} (B) \int_a^b S(\bar{x}, \bar{p}, t) f'(t) dt \quad (\text{by (2.5)}) \\
& = \frac{1}{b-a} (B) \int_a^b f(t) dt + \frac{1}{b-a} (B) \int_a^{x_1} (t-a) f'(t) dt \\
& \quad + \frac{1}{b-a} \sum_{i=1}^{n-1} (B) \int_a^b [t - (P_i b + \bar{P}_i a)] f'(t) dt \\
& \quad + \frac{1}{b-a} (B) \int_{x_n}^b (t-b) f'(t) dt,
\end{aligned}$$

and the representation (2.2) is proved. ■

The following result in approximating the Bochner integral $(B) \int_a^b f(t) dt$ in terms of the convex combination of $(f(x_i))_{i=\overline{1,n}}$ with the weights $(p_i)_{i=\overline{1,n}}$ holds.

Theorem 3. Assume that $f : [a, b] \rightarrow X$, $(x_i)_{i=\overline{1,n}}$ and $(p_i)_{i=\overline{1,n}}$ are as in Lemma 1. Then we have the inequality:

$$\begin{aligned}
(2.7) \quad & \left\| (B) \int_a^b f(t) dt - (b-a) \sum_{i=1}^n p_i f(x_i) \right\| \\
& \leq \int_a^{x_1} (t-a) \|f'(t)\| dt + \sum_{i=1}^{n-1} \int_{x_i}^{x_{i+1}} |t - (P_i b + \bar{P}_i a)| \|f'(t)\| dt \\
& \quad + \int_{x_n}^b (b-t) \|f'(t)\| dt
\end{aligned}$$

$$\begin{aligned}
& \leq \left\{ \begin{array}{l} (x_1 - a) \|f'\|_{[a, x_1], 1} + \sum_{i=1}^{n-1} \mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], 1} \\ \quad + (b - x_n) \|f'\|_{[x_n, b], 1} \\ \frac{(x_1 - a)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[a, x_1], p} + \sum_{i=1}^{n-1} [\mu_q(x_i, P_i b + \bar{P}_i a, x_{i+1})]^{\frac{1}{q}} \|f'\|_{[x_i, x_{i+1}], p} \\ \quad + \frac{(b - x_n)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[x_n, b], p} \quad \text{where } p > 1, \frac{1}{p} + \frac{1}{q} = 1 \\ \quad \text{and if } f' \in L_p([a, b]; X); \\ \frac{(x_1 - a)^2}{2} \|f'\|_{[a, x_1], \infty} + \sum_{i=1}^{n-1} \mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], \infty} \\ \quad + \frac{(b - x_n)^2}{2} \|f'\|_{[x_n, b], \infty} \quad \text{if } f' \in L_\infty([a, b]; X); \end{array} \right.
\end{aligned}$$

$$\leq \begin{cases} \max \left(x_1 - a, \max_{i=1, n-1} \{ \mu_\infty (x_i, P_i b + \bar{P}_i a, x_{i+1}) \}, b - x_n \right) \| |f'| \|_{[a, b], 1} \\ \quad \text{if } f' \in L_1 ([a, b]; X); \\ \left[\frac{(x_1 - a)^{q+1}}{q+1} + \sum_{i=1}^{n-1} \mu_q (x_i, P_i b + \bar{P}_i a, x_{i+1}) + \frac{(b - x_n)^{q+1}}{q+1} \right]^{\frac{1}{q}} \| |f'| \|_{[a, b], p} \\ \quad \text{where } p > 1, \frac{1}{p} + \frac{1}{q} = 1 \text{ and if } f' \in L_p ([a, b]; X); \\ \left[\frac{(x_1 - a)^2}{2} + \sum_{i=1}^{n-1} \mu_1 (x_i, P_i b + \bar{P}_i a, x_{i+1}) + \frac{(b - x_n)^2}{2} \right] \| |f'| \|_{[a, b], \infty} \\ \quad \text{if } f' \in L_\infty ([a, b]; X); \end{cases}$$

where $L_p ([a, b]; X)$, $p \in [1, \infty]$ are the usual vector-valued Lebesgue spaces and

$$\begin{aligned} \| |h| \|_{[\alpha, \beta], \infty} &:= \text{ess} \sup_{t \in [\alpha, \beta]} \| h(t) \|, \\ \| |h| \|_{[\alpha, \beta], p} &:= \left(\int_\alpha^\beta \| h(t) \|^p dt \right)^{\frac{1}{p}}, \quad p \geq 1, \end{aligned}$$

and the functions $\mu_q (\cdot, \cdot, \cdot)$, $q \in [1, \infty]$ were defined in (2.1).

Proof. Using the properties of the norm, we have, by (2.2), that

$$\begin{aligned} &\left\| (B) \int_a^b f(t) dt - (b-a) \sum_{i=1}^n p_i f(x_i) \right\| \\ &\leq \int_a^{x_1} (t-a) \| f'(t) \| dt + \sum_{i=1}^{n-1} \int_{x_i}^{x_{i+1}} |t - (P_i b + \bar{P}_i a)| \| f'(t) \| dt \\ &\quad + \int_{x_n}^b (b-t) \| f'(t) \| dt \end{aligned}$$

and the first inequality in (2.7) is proved.

Now, observe that

$$\begin{aligned} &\int_a^{x_1} (t-a) \| f'(t) \| dt \\ &\leq \begin{cases} (x_1 - a) \| |f'| \|_{[a, x_1], 1}; \\ \frac{(x_1 - a)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \| |f'| \|_{[a, x_1], p} \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p ([a, b]; X); \\ \frac{(x_1 - a)^2}{2} \| |f'| \|_{[a, x_1], \infty} \quad \text{if } f' \in L_\infty ([a, b]; X); \end{cases} \end{aligned}$$

and

$$\begin{aligned} & \sum_{i=1}^{n-1} \int_{x_i}^{x_{i+1}} |t - (P_i b + \bar{P}_i a)| \|f'(t)\| dt \\ & \leq \begin{cases} \sup_{t \in [x_i, x_{i+1}]} |t - (P_i b + \bar{P}_i a)| \|f'\|_{[x_i, x_{i+1}], 1}; \\ \left(\int_{x_i}^{x_{i+1}} |t - (P_i b + \bar{P}_i a)|^q dt \right)^{\frac{1}{q}} \|f'\|_{[x_i, x_{i+1}], p} \\ \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, \text{ and } f' \in L_p([a, b]; X); \\ \int_{x_i}^{x_{i+1}} |t - (P_i b + \bar{P}_i a)| dt \|f'\|_{[x_i, x_{i+1}], \infty} \quad \text{if } f' \in L_\infty([a, b]; X); \end{cases} \\ & = \begin{cases} \mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], 1}; \\ [\mu_q(x_i, P_i b + \bar{P}_i a, x_{i+1})]^{\frac{1}{q}} \|f'\|_{[x_i, x_{i+1}], p} \\ \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, \text{ and } f' \in L_p([a, b]; X); \\ \mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], \infty} \quad \text{if } f' \in L_\infty([a, b]; X); \end{cases} \end{aligned}$$

with

$$\begin{aligned} & \int_{x_n}^b (b-t) \|f'(t)\| dt \\ & \leq \begin{cases} (b-x_n) \|f'\|_{[x_n, b], 1}; \\ \frac{(b-x_n)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[x_n, b], p} \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a, b]; X); \\ \frac{(b-x_n)^2}{2} \|f'\|_{[x_n, b], \infty} \quad \text{if } f' \in L_\infty([a, b]; X); \end{cases} \end{aligned}$$

giving the second inequality in (2.7).

Finally, observe that

$$\begin{aligned} & (x_1 - a) \|f'\|_{[a, x_1], 1} + \sum_{i=1}^{n-1} \mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], 1} \\ & + (b - x_n) \|f'\|_{[x_n, b], 1} \\ & \leq \max \left\{ x_1 - a, \max_{i=1, n-1} \{ \mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) \}, b - x_n \right\} \\ & \quad \times \left[\|f'\|_{[a, x_1], 1} + \sum_{i=1}^{n-1} \|f'\|_{[x_i, x_{i+1}], 1} + \|f'\|_{[x_n, b], 1} \right] \\ & \leq \max \left\{ x_1 - a, \max_{i=1, n-1} \{ \mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) \}, b - x_n \right\} \|f'\|_{[a, b], 1}. \end{aligned}$$

Further, by the discrete Hölder's inequality we have that

$$\begin{aligned}
& \frac{(x_1 - a)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[a, x_1], p} + \sum_{i=1}^{n-1} [\mu_q(x_i, P_i b + \bar{P}_i a, x_{i+1})]^{\frac{1}{q}} \|f'\|_{[x_i, x_{i+1}], p} \\
& + \frac{(b - x_n)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[x_n, b], p} \\
\leq & \left\{ \left[\frac{(x_1 - a)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \right]^q + \sum_{i=1}^{n-1} \left([\mu_q(x_i, P_i b + \bar{P}_i a, x_{i+1})]^{\frac{1}{q}} \right)^q \right. \\
& \left. + \left[\frac{(b - x_n)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \right]^q \right\}^{\frac{1}{q}} \times \left[\|f'\|_{[x_n, b], p}^p + \|f'\|_{[a, x_1], p}^p + \sum_{i=1}^{n-1} \|f'\|_{[x_i, x_{i+1}], p}^p \right]^{\frac{1}{p}} \\
= & \left[\frac{(x_1 - a)^{q+1}}{q+1} + \sum_{i=1}^{n-1} \mu_q(x_i, P_i b + \bar{P}_i a, x_{i+1}) + \frac{(b - x_n)^{q+1}}{q+1} \right]^{\frac{1}{q}} \|f'\|_{[a, b], p}
\end{aligned}$$

and

$$\begin{aligned}
& \frac{(x_1 - a)^2}{2} \|f'\|_{[a, x_1], \infty} + \sum_{i=1}^{n-1} \mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) \|f'\|_{[x_i, x_{i+1}], \infty} \\
& + \frac{(b - x_n)^2}{2} \|f'\|_{[x_n, b], \infty} \\
\leq & \left[\frac{(x_1 - a)^2}{2} + \sum_{i=1}^{n-1} \mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) + \frac{(b - x_n)^2}{2} \right] \\
& \times \max \left\{ \|f'\|_{[a, x_1], \infty}, \max_{i=1, n-1} \|f'\|_{[x_i, x_{i+1}], \infty}, \|f'\|_{[x_n, b], \infty} \right\} \\
= & \left[\frac{(x_1 - a)^2}{2} + \sum_{i=1}^{n-1} \mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) + \frac{(b - x_n)^2}{2} \right] \|f'\|_{[a, b], \infty};
\end{aligned}$$

and the theorem is completely proved. ■

It is a natural assumption to consider the weights $p_i > 0$ ($i = 1, \dots, n$) for which $\xi_i := P_i b + \bar{P}_i a$ ($\in [a, b]$) will be in the interval $[x_i, x_{i+1}]$ ($i = 1, \dots, n$). In this case we have:

$$\mu_\infty(x_i, P_i b + \bar{P}_i a, x_{i+1}) = \frac{1}{2} h_i + \left| P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right|,$$

where $h_i := x_{i+1} - x_i$, and for $p \in [1, \infty)$

$$\mu_p(x_i, P_i b + \bar{P}_i a, x_{i+1}) = \frac{1}{p+1} \left[(P_i b + \bar{P}_i a - x_i)^{p+1} + (x_{i+1} - P_i b - \bar{P}_i a)^{p+1} \right].$$

Note that for $p = 1$, we have

$$\mu_1(x_i, P_i b + \bar{P}_i a, x_{i+1}) = \frac{1}{4} h_i^2 + \left(P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right)^2.$$

The following corollary is important for applications.

Corollary 1. *With the assumptions of Lemma 1 and if $x_i \leq P_i b + \bar{P}_i a \leq x_{i+1}$ for each $i = 1, \dots, n-1$, then we have the inequalities:*

$$(2.8) \quad \left\| (B) \int_a^b f(t) dt - (b-a) \sum_{i=1}^n p_i f(x_i) \right\| \\ \leq \begin{cases} (x_1 - a) \|f'\|_{[a, x_1], 1} + \sum_{i=1}^{n-1} \left[\frac{1}{2} h_i + \left| P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right| \right] \|f'\|_{[x_i, x_{i+1}], 1} \\ \quad + (b - x_n) \|f'\|_{[x_n, b], 1}; \\ \frac{(x_1 - a)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[a, x_1], p} \\ + \frac{1}{(q+1)^{\frac{1}{q}}} \sum_{i=1}^{n-1} \left[(P_i b + \bar{P}_i a - x_i)^{q+1} + (x_{i+1} - P_i b - \bar{P}_i a)^{q+1} \right]^{\frac{1}{q}} \|f'\|_{[x_i, x_{i+1}], p} \\ \quad + \frac{(b - x_n)^{1+\frac{1}{q}}}{(q+1)^{\frac{1}{q}}} \|f'\|_{[x_n, b], p} \quad \text{where } p > 1, \frac{1}{p} + \frac{1}{q} = 1 \\ \quad \text{and if } f' \in L_p([a, b]; X); \\ \frac{(x_1 - a)^2}{2} \|f'\|_{[a, x_1], \infty} + \sum_{i=1}^{n-1} \left[\frac{1}{4} h_i^2 + \left(P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right)^2 \right] \|f'\|_{[x_i, x_{i+1}], \infty} \\ \quad + \frac{(b - x_n)^2}{2} \|f'\|_{[x_n, b], \infty} \quad \text{if } f' \in L_\infty([a, b]; X); \\ \max \left(x_1 - a, \frac{1}{2} \max_{i=1, n-1} h_i + \max_{i=1, n-1} \left| P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right|, b - x_n \right) \|f'\|_{[a, b], 1} \\ \quad \text{if } f' \in L_1([a, b]; X); \\ \frac{1}{(q+1)^{\frac{1}{q}}} \left[(x_1 - a)^{q+1} + \sum_{i=1}^{n-1} \left[(P_i b + \bar{P}_i a - x_i)^{q+1} \right. \right. \\ \quad \left. \left. + (x_{i+1} - P_i b - \bar{P}_i a)^{q+1} \right] + (b - x_n)^{q+1} \right]^{\frac{1}{q}} \|f'\|_{[a, b], p} \\ \quad \text{if } p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a, b]; X); \\ \left[\frac{(x_1 - a)^2}{2} + \sum_{i=1}^{n-1} \left[\frac{1}{4} h_i^2 + \left(P_i b + \bar{P}_i a - \frac{x_i + x_{i+1}}{2} \right)^2 \right] + \frac{(b - x_n)^2}{2} \right] \|f'\|_{[a, b], \infty} \\ \quad \text{if } f' \in L_\infty([a, b]; X). \end{cases}$$

Remark 1. *For $n = 1$, we recapture from (2.8) the Ostrowski type inequalities incorporated in Theorems 1 and 2.*

3. THE CASE OF TWO POINTS

The following proposition is a particular case of Corollary 1 for $n = 2$ and will be considered in some details since there are important for applications.

Proposition 1. *Let $(X, \|\cdot\|)$ be a Banach space with the Radon-Nikodym property and $f : [a, b] \rightarrow X$ be an absolutely continuous function on $[a, b]$. If $a \leq x_1 \leq x_2 \leq b$ ($b > a$) and $t \in [0, 1]$ satisfies the condition*

$$(0 \leq) \frac{x_1 - a}{b - a} \leq t \leq \frac{x_2 - a}{b - a} (\leq 1),$$

then we have the inequalities

$$(3.1) \quad \left\| (B) \int_a^b f(s) ds - (b-a) [tf(x_2) + (1-t)f(x_1)] \right\| \leq \begin{cases} (x_1 - a) \|f'\|_{[a,x_1],1} + \left[\frac{1}{2} (x_2 - x_1) + |tb + (1-t)a - \frac{x_1+x_2}{2}| \right] \|f'\|_{[x_1,x_2],1} \\ + (b - x_2) \|f'\|_{[x_2,b],1}; \\ \frac{1}{(q+1)^{1/q}} \left\{ (x_1 - a)^{1+1/q} \|f'\|_{[a,x_1],p} \right. \\ + \left[(tb + (1-t)a - x_1)^{q+1} + (x_2 - tb - (1-t)a)^{q+1} \right]^{1/q} \|f'\|_{[x_1,x_2],p} \\ \left. + (b - x_2)^{1+1/q} \|f'\|_{[x_2,b],p} \right\}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{(x_1 - a)^2}{2} \|f'\|_{[a,x_1],\infty} + \left[\frac{1}{4} (x_2 - x_1)^2 + \left[tb + (1-t)a - \frac{x_1+x_2}{2} \right]^2 \right] \|f'\|_{[x_1,x_2],\infty} \\ + \frac{(b - x_2)^2}{2} \|f'\|_{[x_2,b],\infty}, f' \in L_\infty([a,b];X); \\ \max \left\{ x_1 - a, \frac{1}{2} (x_2 - x_1) + |tb + (1-t)a - \frac{x_1+x_2}{2}|, b - x_2 \right\} \|f'\|_{[a,b],1}; \\ \frac{1}{(q+1)^{1/q}} \left\{ (x_1 - a)^{q+1} + (tb + (1-t)a - x_1)^{q+1} + (x_2 - tb - (1-t)a)^{q+1} \right. \\ \left. + (b - x_2)^{q+1} \right\}^{1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \left[\frac{(x_1 - a)^2}{2} + \frac{1}{4} (x_2 - x_1)^2 + \left[tb + (1-t)a - \frac{x_1+x_2}{2} \right]^2 + \frac{(b - x_2)^2}{2} \right] \|f'\|_{[a,b],\infty}, \\ f' \in L_\infty([a,b];X). \end{cases}$$

The following particular inequalities are of interest.

1. If $x_1 = a, x_2 = b$, then for any $t \in [0, 1]$, we have the inequalities

$$(3.2) \quad \left\| (B) \int_a^b f(s) ds - (b-a) [tf(b) + (1-t)f(a)] \right\| \leq$$

$$\leq \begin{cases} \left[\frac{1}{2} (b-a) + |tb + (1-t)a - \frac{a+b}{2}| \right] \|f'\|_{[a,b],1}; \\ \frac{1}{(q+1)^{1/q}} \left[t^{q+1} + (1-t)^{q+1} \right]^{1/q} (b-a)^{1+1/q} \|f'\|_{[a,b],p}, \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \left[\frac{1}{4} (b-a)^2 + \left(tb + (1-t)a - \frac{a+b}{2} \right)^2 \right] \|f'\|_{[a,b],\infty}, f' \in L_\infty([a,b];X). \end{cases}$$

The best inequality one can get from (3.2) is for $t = \frac{1}{2}$, obtaining the trapezoidal rule

$$(3.3) \quad \left\| (B) \int_a^b f(s) ds - (b-a) \cdot \frac{f(b) + f(a)}{2} \right\| \leq \begin{cases} \frac{1}{2} (b-a) \|f'\|_{[a,b],1}; \\ \frac{1}{2(q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{1}{4} (b-a)^2 \|f'\|_{[a,b],\infty}, f' \in L_\infty([a,b];X). \end{cases}$$

2. If $x_1 = \frac{3a+b}{4}, x_2 = \frac{a+3b}{4}$, then for any $t \in [\frac{1}{4}, \frac{3}{4}]$ we have the inequalities

$$(3.4) \quad \left\| (B) \int_a^b f(s) ds - (b-a) \left[tf\left(\frac{3a+b}{4}\right) + (1-t)f\left(\frac{a+3b}{4}\right) \right] \right\| \leq \begin{cases} \frac{b-a}{4} \|f'\|_{[a,\frac{3a+b}{4}],1} + \left[\frac{1}{4}(b-a) + |tb + (1-t)a - \frac{a+b}{2}| \right] \|f'\|_{[\frac{3a+b}{4},\frac{a+3b}{4}],1} \\ + \frac{b-a}{4} \|f'\|_{[\frac{a+3b}{4},b],1}; \\ \frac{1}{4^{q+1}(q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[a,\frac{3a+b}{4}],p} \\ + \frac{1}{(q+1)^{1/q}} \left[(tb + (1-t)a - \frac{3a+b}{4})^{q+1} \right. \\ \left. + \left(\frac{a+3b}{4} - tb - (1-t)a \right)^{q+1} \right]^{1/q} \|f'\|_{[\frac{3a+b}{4},\frac{a+3b}{4}],p} \\ + \frac{1}{4^{q+1}(q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[\frac{a+3b}{4},b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{(b-a)^2}{8} \|f'\|_{[a,\frac{3a+b}{4}],\infty} + \left[\frac{1}{16}(b-a)^2 + [tb + (1-t)a - \frac{a+b}{2}]^2 \right] \|f'\|_{[\frac{3a+b}{4},\frac{a+3b}{4}],\infty} \\ + \frac{(b-a)^2}{8} \|f'\|_{[\frac{a+3b}{4},b],\infty}, f' \in L_\infty([a,b];X); \\ \left[\frac{1}{4}(b-a) + |tb + (1-t)a - \frac{a+b}{2}| \right] \|f'\|_{[a,b],1}; \\ \frac{1}{(q+1)^{1/q}} \left[\frac{2(b-a)^{q+1}}{4^{q+1}} + \left(tb + (1-t)a - \frac{3a+b}{4} \right)^{q+1} + \left(\frac{a+3b}{4} - tb - (1-t)a \right)^{q+1} \right]^{1/q} \\ \times (b-a)^{1+1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \left[\frac{1}{8}(b-a)^2 + \left(tb + (1-t)a - \frac{a+b}{2} \right)^2 \right] \|f'\|_{[a,b],\infty}, f' \in L_\infty([a,b];X). \end{cases}$$

The best inequality one can get from (3.4) is for $t = \frac{1}{2}$, obtaining

$$(3.5) \quad \left\| (B) \int_a^b f(s) ds - (b-a) \cdot \frac{f\left(\frac{a+3b}{4}\right) + f\left(\frac{3a+b}{4}\right)}{2} \right\| \leq \begin{cases} \frac{1}{4} (b-a) \|f'\|_{[a,b],1}; \\ \frac{1}{4(q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{1}{8} (b-a)^2 \|f'\|_{[a,b],\infty}, f' \in L_\infty([a,b];X). \end{cases}$$

Remark 2. One may realize that, instead of using the trapezoidal rule in approximating the Bochner integral $(B) \int_a^b f(t) dt$, that one should use the rule

$$(3.6) \quad QT(f; a, b) := (b - a) \cdot \frac{f\left(\frac{a+3b}{4}\right) + f\left(\frac{3a+b}{4}\right)}{2},$$

which provides a halving of the bound on the error.

4. THE CASE OF THREE POINTS

The case of three points is important for applications since it contains amongst others Simpson's quadrature rule.

The following proposition holds:

Proposition 2. Let $(X, \|\cdot\|)$ be a Banach space with the Radon-Nikodym property and $f : [a, b] \rightarrow X$ be an absolutely continuous function on $[a, b]$. If $a \leq x_1 \leq x_2 \leq x_3 \leq b$ ($b > a$) and $\alpha, \beta, \gamma \in [0, 1]$ with $\alpha + \beta + \gamma = 1$ satisfies the condition

$$(4.1) \quad (0 \leq) \frac{x_1 - a}{b - a} \leq \alpha \leq \frac{x_2 - a}{b - a} \leq \alpha + \beta \leq \frac{x_3 - a}{b - a} (\leq 1),$$

then we have the inequalities

$$(4.2) \quad \left\| (B) \int_a^b f(t) dt - (b - a) [\alpha f(x_1) + \beta f(x_2) + (1 - \alpha - \beta) f(x_3)] \right\|$$

$$\leq \begin{cases} (x_1 - a) \|f'\|_{[a, x_1], 1} + \left[\frac{1}{2} (x_2 - x_1) + |\alpha b + (1 - \alpha) a - \frac{x_1 + x_2}{2}| \right] \|f'\|_{[x_1, x_2], 1} \\ + \left[\frac{1}{2} (x_3 - x_2) + |(\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{x_2 + x_3}{2}| \right] \|f'\|_{[x_2, x_3], 1} \\ + (b - x_2) \|f'\|_{[x_2, b], 1}; \\ \\ \frac{1}{(q+1)^{1/q}} \left\{ (x_1 - a)^{1+1/q} \|f'\|_{[a, x_1], p} \right. \\ \left. + \left[(\alpha b + (1 - \alpha) a - x_1)^{q+1} + (x_2 - \alpha b - (1 - \alpha) a)^{q+1} \right]^{1/q} \|f'\|_{[x_1, x_2], p} \right\} \\ + \left[((\alpha + \beta) b + (1 - \alpha - \beta) a - x_2)^{q+1} + (x_3 - (\alpha + \beta) b - (1 - \alpha - \beta) a)^{q+1} \right]^{1/q} \|f'\|_{[x_2, x_3], p} \\ + (b - x_3)^{1+1/q} \|f'\|_{[x_3, b], p}, \quad p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a, b]; X); \\ \\ \frac{(x_1 - a)^2}{2} \|f'\|_{[a, x_1], \infty} + \left[\frac{1}{4} (x_2 - x_1)^2 + [\alpha b + (1 - \alpha) a - \frac{x_1 + x_2}{2}]^2 \right] \|f'\|_{[x_1, x_2], \infty} \\ \left[\frac{1}{4} (x_3 - x_2)^2 + [(\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{x_2 + x_3}{2}]^2 \right] \|f'\|_{[x_2, x_3], \infty} \\ + \frac{(b - x_3)^2}{2} \|f'\|_{[x_3, b], \infty}, \quad f' \in L_\infty([a, b]; X); \end{cases}$$

$$\leq \begin{cases} \max \left\{ x_1 - a, \frac{1}{2}(x_2 - x_1) + \left| ab + (1 - \alpha) a - \frac{x_1+x_2}{2} \right|, \right. \\ \left. \frac{1}{2}(x_2 - x_1) + \left| (\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{x_2+x_3}{2} \right|, b - x_2 \right\} \|f'\|_{[a,b],1}; \\ \frac{1}{(q+1)^{1/q}} \left\{ (x_1 - a)^{q+1} + (ab + (1 - \alpha) a - x_1)^{q+1} + (x_2 - ab - (1 - \alpha) a)^{q+1} \right. \\ \left. + ((\alpha + \beta) b + (1 - \alpha - \beta) a - x_2)^{q+1} + (x_3 - (\alpha + \beta) b - (1 - \alpha - \beta) a)^{q+1} \right. \\ \left. + (b - x_3)^{q+1} \right\} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b]; X); \\ \left[\frac{(x_1-a)^2}{2} + \frac{1}{4}(x_2-x_1)^2 + \left[ab + (1 - \alpha) a - \frac{x_1+x_2}{2} \right]^2 \right. \\ \left. + \frac{1}{4}(x_3-x_2)^2 + \left[(\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{x_2+x_3}{2} \right]^2 + \frac{(b-x_3)^2}{2} \right] \|f'\|_{[a,b],\infty}, \\ f' \in L_\infty([a,b]; X). \end{cases}$$

The following particular inequalities are of interest.

1. Assume that $x_1 = a, x_2 = \frac{a+b}{2}, x_3 = b$ and $\alpha, \beta \in [0, 1]$ so that $0 \leq \alpha \leq \frac{1}{2} \leq \alpha + \beta \leq 1$, then we have the inequalities

$$(4.3) \quad \left\| (B) \int_a^b f(t) dt - (b-a) \left[\alpha f(a) + \beta f\left(\frac{a+b}{2}\right) + (1-\alpha-\beta) f(b) \right] \right\|$$

$$\leq \begin{cases} \left[\frac{1}{4}(b-a) + \left| ab + (1 - \alpha) a - \frac{3a+b}{4} \right| \right] \|f'\|_{[a,\frac{a+b}{2}],1} \\ + \left[\frac{1}{4}(b-a) + \left| (\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{a+3b}{4} \right| \right] \|f'\|_{[\frac{a+b}{2},b],1}; \\ \frac{1}{(q+1)^{1/q}} \left\{ \left[\alpha^{q+1} (b-a)^{q+1} + \left(\frac{a+b}{2} - ab - (1 - \alpha) a \right)^{q+1} \right]^{1/q} \|f'\|_{[a,\frac{a+b}{2}],p} \right. \\ \left. + \left[((\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{a+b}{2})^{q+1} + (1 - \alpha - \beta)^{q+1} (b-a)^{q+1} \right]^{1/q} \right\} \|f'\|_{[\frac{a+b}{2},b],p} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b]; X); \\ \left[\frac{1}{16}(b-a)^2 + \left[ab + (1 - \alpha) a - \frac{3a+b}{4} \right]^2 \right] \|f'\|_{[a,\frac{a+b}{2}],\infty} \\ \left[\frac{1}{16}(b-a)^2 + \left[(\alpha + \beta) b + (1 - \alpha - \beta) a - \frac{a+3b}{4} \right]^2 \right] \|f'\|_{[\frac{a+b}{2},b],\infty} \\ f' \in L_\infty([a,b]; X); \end{cases}$$

$$\leq \begin{cases} \max \left\{ \left[\frac{1}{4} (b-a) + |\alpha b + (1-\alpha)a - \frac{3a+b}{4}| \right], \right. \\ \left. \left[\frac{1}{4} (b-a) + |(\alpha+\beta)b + (1-\alpha-\beta)a - \frac{a+3b}{4}| \right] \right\} \|f'\|_{[a,b],1} \\ \\ \frac{1}{(q+1)^{1/q}} \left\{ \alpha^{q+1} (b-a)^{q+1} + \left(\frac{a+b}{2} - \alpha b - (1-\alpha)a \right)^{q+1} \right. \\ \left. + \left((\alpha+\beta)b + (1-\alpha-\beta)a - \frac{a+b}{2} \right)^{q+1} + (1-\alpha-\beta)^{q+1} (b-a)^{q+1} \right\}^{1/q} \|f'\|_{[a,b],p} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \\ \left\{ \frac{1}{8} (b-a)^2 + \left[\alpha b + (1-\alpha)a - \frac{3a+b}{4} \right]^2 \right. \\ \left. + \left[(\alpha+\beta)b + (1-\alpha-\beta)a - \frac{a+3b}{4} \right]^2 \right\} \|f'\|_{[a,b],\infty} \\ f' \in L_\infty([a,b];X). \end{cases}$$

It is easy to see that, the best inequality one can derive from (4.3) is the one for $\alpha = \frac{1}{4}$ and $\beta = \frac{3}{4}$, getting

$$(4.4) \quad \left\| (B) \int_a^b f(t) dt - (b-a) \left[\frac{f(a) + f(b)}{4} + \frac{1}{2} f\left(\frac{a+b}{2}\right) \right] \right\|$$

$$\leq \begin{cases} \frac{1}{4} (b-a) \|f'\|_{[a,b],1}; \\ \\ \frac{1}{2^{2+1/q}(q+1)^{1/q}} (b-a)^{1+1/q} \left\{ \|f'\|_{[a,\frac{a+b}{2}],p} + \|f'\|_{[\frac{a+b}{2},b],p} \right\} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \\ \frac{1}{16} (b-a)^2 \left[\|f'\|_{[a,\frac{a+b}{2}],\infty} + \|f'\|_{[\frac{a+b}{2},b],\infty} \right], f' \in L_\infty([a,b];X); \\ \\ \frac{1}{4} (b-a) \|f'\|_{[a,b],1}; \\ \\ \frac{1}{2^{2+1/q}(q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \\ \frac{1}{8} (b-a)^2 \|f'\|_{[a,b],\infty} f' \in L_\infty([a,b];X). \end{cases}$$

The inequality (4.3) incorporates *Simpson's rule* as well. Indeed, if we choose $\alpha = \frac{1}{6}$, $\beta = \frac{4}{6}$, then we get from (4.3) the result

$$(4.5) \quad \left\| (B) \int_a^b f(t) dt - \frac{(b-a)}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] \right\|$$

$$\begin{aligned} & \leq \begin{cases} \frac{1}{3}(b-a) \|f'\|_{[a,b],1}; \\ \frac{1}{(q+1)^{1/q}} \frac{(2^{q+1}+1)^{1/q}}{6^{1+1/q}} (b-a)^{1+1/q} \left\{ \|f'\|_{[a,\frac{a+b}{2}],p} + \|f'\|_{[\frac{a+b}{2},b],p} \right\} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{5}{72}(b-a)^2 \left[\|f'\|_{[a,\frac{a+b}{2}],\infty} + \|f'\|_{[\frac{a+b}{2},b],\infty} \right], f' \in L_\infty([a,b];X); \end{cases} \\ & \leq \begin{cases} \frac{1}{3}(b-a) \|f'\|_{[a,b],1}; \\ \frac{(2^{q+1}+1)^{1/q}}{2 \cdot 3^{1+1/q} (q+1)^{1/q}} (b-a)^{1+1/q} \|f'\|_{[a,b],p}, p > 1, \frac{1}{p} + \frac{1}{q} = 1, f' \in L_p([a,b];X); \\ \frac{5}{36}(b-a)^2 \|f'\|_{[a,b],\infty} f' \in L_\infty([a,b];X). \end{cases} \end{aligned}$$

Remark 3. It is obvious that, if the values at $a, \frac{a+b}{2}$ and b of the function $f : [a,b] \rightarrow X$ are available, then one should choose the rule

$$QS(f; a, b) := (b-a) \left[\frac{f(a) + f(b)}{4} + \frac{1}{2} f\left(\frac{a+b}{2}\right) \right]$$

that provides a better approximation for the Bochner integral $(B) \int_a^b f(t) dt$ than the classical Simpson's rule.

2. If one chooses $x_1 = \frac{3a+b}{4}, x_2 = \frac{a+b}{2}, x_3 = \frac{a+3b}{4}$, then by the use of inequality (4.2), one can derive estimates for the norm of difference

$$(B) \int_a^b f(t) dt - (b-a) \left[\alpha f\left(\frac{3a+b}{4}\right) + \beta f\left(\frac{a+b}{2}\right) + (1-\alpha-\beta) f\left(\frac{a+3b}{4}\right) \right]$$

in terms of the Lebesgue norms of the derivative f' .

We omit the details.

For more scalar-valued three point quadrature rules, see [6].

Acknowledgement: S. S. Dragomir and Y. J. Cho greatly acknowledge the financial support from the Brain Pool Program (2002) of the Korean Federation of Science and Technology Societies. The research was performed under the "Memorandum of Understanding" between Victoria University and Gyeongsang National University.

REFERENCES

- [1] G.A. ANASTASSIOU, Ostrowski type inequalities, *Proc. of Amer. Math. Soc.*, **123**(12) (1995), 3775-3781.
- [2] W. ARENDT, C.J.K. BATTY, M. HIEBER and F. NEUBRANDER, *Vector-Valued Laplace Transforms and Cauchy Problems*, Birkhäuser Verlag Basel, 2001.
- [3] N.S. BARNETT, C. BUŞE, P. CERONE and S.S. DRAGOMIR, Ostrowski's inequality for vector-valued functions and applications, *Comp. & Math. Appl.*, **44**(2002), 559-572.
- [4] S. BOCHNER, Integration von Funktionen, deren Werte die Elemente eines Vektorraumes, *Fund. Math.*, **20** (1933), 262-276.
- [5] P. CERONE and S.S. DRAGOMIR, Midpoint-type rules from an inequalities point of view, *Handbook of Analytic-Computational Methods in Applied Mathematics*, Editor: G. Anastassiou, CRC Press, New York (2000), 135-200.
- [6] P. CERONE and S.S. DRAGOMIR, Three point quadrature rules, in *Ostrowski Type Inequalities and Applications in Numerical Integration*, S.S. Dragomir and Th.M Rassias (Eds), Kluwer Academic Publishers, Dordrecht, 2002, 141-250

- [7] J. DIETSEL and J.J. UHL, Jr., *Vector Measures*, Amer. Math. Society, Mathematical Surveys, **15**, Providence, Rhode Island, (1977).
- [8] S.S. DRAGOMIR and S. WANG, A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, **28** (1997), 239-244.
- [9] S.S. DRAGOMIR and S. WANG, A new inequality of Ostrowski's type in L_p -norm, *Indian J. of Math.*, **40**(3) (1998), 295-304.
- [10] S.S. DRAGOMIR and Th.M. RASSIAS (Ed.), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publishers, Dordrecht, 2002.
- [11] A.M. FINK, Bounds on the derivation of a function from its averages, *Czech. Math. J.*, **42** (1992), 289-310.
- [12] E. HILLE and R.S. PHILIPS, *Functional Analysis and Semi-Groups*, Amer. Math. Soc. Colloq. Publ., Vol. 31, Providence, R.I. (1957).
- [13] J. MIKUSIŃSKI, *The Bochner Integral*, Birkhäuser Verlag Basel, 1978.
- [14] R. NAGEL (Ed.), *One-Parameter Semigroups of Positive Operators*, Springer Lect. Notes in Math., 1184 (1986).
- [15] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer Verlag, (1983).

SCHOOL OF COMPUTER SCIENCE AND MATHEMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY,
PO Box 14428, MCMC 8001, VICTORIA, AUSTRALIA.

E-mail address: pietro.cerone@vu.edu.au

URL: <http://rgmia.vu.edu.au/cerone/index.html>

DEPARTMENT OF MATHEMATICS EDUCATION, COLLEGE OF EDUCATION, GYEONGSANG NATIONAL
UNIVERSITY, CHINJU 660-701, KOREA

E-mail address: yjcho@nongae.gsnu.ac.kr

SCHOOL OF COMPUTER SCIENCE AND MATHEMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY,
PO Box 14428, MCMC 8001, VICTORIA, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>

DEPARTMENT OF MATHEMATICS, KYUNG NAM UNIVERSITY, MASAN,, KYUNG NAM 631-701, Ko-
REA

E-mail address: jongkyuk@kyungnam.ac.kr

DEPARTMENT OF MATHEMATICS, DONGEUI UNIVERSITY, PUSAN 614-714, KOREA

E-mail address: sskim@dongeui.ac.kr