A Proof On Arnold's Chord Conjecture On Cotangent Bundles *

Renyi Ma
Department of Mathematics
Tsinghua University
Beijing, 100084
People's Republic of China
rma@math.tsinghua.edu.cn

Abstract

In this article, we prove that there exists at least one chord which is characteristic of Reeb vector field connecting a given Legendre submanifold in a contact manifolds of induced type in the cotangent bundles of any smooth open manifolds which confirms the Arnold conjecture in cotangent bundles.

Keywords J-holomorphic curves, Legendre submanifolds, Reeb chord. **2000 MR Subject Classification** 32Q65, 53D35, 53D12

1 Introduction and results

Let Σ be a smooth closed oriented manifold of dimension 2n-1. A contact form on Σ is a 1-form such that $\lambda \wedge (d\lambda)^{n-1}$ is a volume form on Σ . Associated to λ there is the so-called Reeb vectorfield X_{λ} defined by

$$i_X \lambda \equiv 1, \quad i_X d\lambda \equiv 0.$$

^{*}Project 19871044 Supported by NSF

Concerning the dynamics of Reeb flow, there is a well-known conjecture raised by Arnold in [2] which concerned the Reeb orbit and Legendre submanifold in a contact manifold. If (Σ, λ) is a contact manifold with contact form λ of dimension 2n-1, then a Legendre submanifold is a submanifold \mathcal{L} of Σ , which is (n-1)dimensional and everywhere tangent to the contact structure $\ker \lambda$. Then a characteristic chord for (λ, \mathcal{L}) is a smooth path $x:[0,T] \to \Sigma, T>0$ with $\dot{x}(t)=X_{\lambda}(x(t))$ for $t\in(0,T),x(0),x(T)\in\mathcal{L}$ Arnold raised the following conjecture:

Conjecture1(see[2]). Let λ_0 be the standard tight contact form $\lambda_0 = \frac{1}{2}(x_1dy_1 - y_1dx_1 + x_2dy_2 - y_2dx_2)$ on the three sphere $S^3 = \{(x_1, y_1, x_2, y_2) \in R^4 | x_1^2 + y_1^2 + x_2^2 + y_2^2 = 1\}$. If $f: S^3 \to (0, \infty)$ is a smooth function and \mathcal{L} is a Legendre knot in S^3 , then there is a characteristic chord for $(f\lambda_0, \mathcal{L})$.

This conjecture was completely solved in [16, 17].

In this paper we improve the Gromov's proof on that there exists at least one intersection point for the weakly exact Lagrangian submanifold under the weakly Lagrangian isotopy [10, $2.3.B_{3-4}$] to prove:

Theorem 1.1 Let (Σ, λ) be a contact manifold with contact form λ of induced type or Weinstein type in the cotangent bundles of any open smooth manifold with symplectic form $\sum_{i=1}^{n} dp_i \wedge dq_i$ induced by Liouville form $\alpha = \sum_{i=1}^{n} p_i dq_i$, i.e., there exists a transversal vector field Z to Σ such that $L_Z \omega = \omega$, $\lambda = i_Z \omega$. Let X_λ its Reeb vector field and \mathcal{L} a closed Legendre submanifold. Then $p_i dq_i - \lambda$ defines an element $[p_i dq_i - \lambda] \in H^1(\Sigma)$. If $[p_i dq_i - \lambda] = 0$, then there exists at least one characteristic chord for (X_λ, \mathcal{L}) .

Sketch of proofs: We work in the framework as in [10, 16]. In Section 2, we study the linear Cauchy-Riemann operator and sketch some basic properties. In section 3, first we construct a Lagrangian submanifold W under the assumption that there does not exist Reeb chord connecting the Legendre submanifold \mathcal{L} ; second, we study the space $\mathcal{D}(V,W)$ consisting of contractible disks in manifold V with boundary in Lagrangian submanifold V and construct a Fredholm section of tangent bundle of $\mathcal{D}(V,W)$. In section 4, following [10], we construct a non-proper Fredholm section by using a special anti-holomorphic section as in [10, 16]. In section 5, we transform the non-homogenious Cauchy-Riemann equations to J-holomorphic curves. In section 6, we finish the proof of Theorem 1.1. as in [10].

2 Linear Fredholm Theory

For $100 < k < \infty$ consider the Hilbert space V_k consisting of all maps $u \in H^{k,2}(D, \mathbb{C}^n)$, such that $u(z) \in \mathbb{R}^n \subset \mathbb{C}^n$ for almost all $z \in \partial D$. L_{k-1}

denotes the usual Hilbert L_{k-1} -space $H_{k-1}(D, \mathbb{C}^n)$. We define an operator $\bar{\partial}: V_p \mapsto L_p$ by

$$\bar{\partial}u = u_s + iu_t \tag{2.1}$$

where the coordinates on D are (s,t)=s+it, $D=\{z||z|\leq 1\}$. The following result is well known(see[4, 21]).

Proposition 2.1 $\bar{\partial}: V_p \mapsto L_p$ is a surjective real linear Fredholm operator of index n. The kernel consists of the constant real valued maps.

Let $(C^n, \sigma = -Im(\cdot, \cdot))$ be the standard symplectic space. We consider a real n-dimensional plane $R^n \subset C^n$. It is called Lagrangian if the skew-scalar product of any two vectors of R^n equals zero. For example, the plane $\{(p,q)|p=0\}$ and $\{(p,q)|q=0\}$ are Lagrangian subspaces. The manifold of all (nonoriented) Lagrangian subspaces of R^{2n} is called the Lagrangian-Grassmanian $\Lambda(n)$. One can prove that the fundamental group of $\Lambda(n)$ is free cyclic, i.e. $\pi_1(\Lambda(n)) = Z$. Next assume $(\Gamma(z))_{z \in \partial D}$ is a smooth map associating to a point $z \in \partial D$ a Lagrangian subspace $\Gamma(z)$ of C^n , i.e. $(\Gamma(z))_{z \in \partial D}$ defines a smooth curve α in the Lagrangian-Grassmanian manifold $\Lambda(n)$. Since $\pi_1(\Lambda(n)) = Z$, one have $[\alpha] = ke$, we call integer k the Maslov index of curve α and denote it by $m(\Gamma)$, see([2]).

Now let $z: S^1 \to R^n \subset C^n$ be a smooth curve. Then it defines a constant loop α in Lagrangian-Grassmanian manifold $\Lambda(n)$. This loop defines the Maslov index $m(\alpha)$ of the map z which is easily seen to be zero.

Now Let (V, ω) be a symplectic manifold and $W \subset V$ a Lagrangian submanifold. Let $u:(D,\partial D)\to (V,W)$ be a smooth map homotopic to constant map $u_0:(D,\partial D)\to p\in W$. Then u^*TV is a symplectic vector bundle and $(u|_{\partial D})^*TW$ be a Lagrangian subbundle in u^*TV . Since u is homotopic to u_0 by h(t,z) with $h(0,\cdot)=u_0$ and $h(1,\cdot)=u$, we can take a trivialization of h^*TV as

$$\Phi(h^*TV) = [0,1] \times D \times C^n$$

and

$$\Phi(h|_{\partial D})^*TW \subset [0,1] \times S^1 \times C^n$$
.

Let

$$\pi_2: [0,1] \times D \times C^n \to C^n$$

then

$$\bar{h}:(t,z)\in[0,1]\times S^1\to\pi_2\Phi(h|_{\partial D})^*TW|(t,z)\in\Lambda(n).$$

Lemma 2.1 Let $u:(D^2,\partial D^2)\to (V,W)$ be a $C^k-map\ (k\geq 1)$ as above. Then,

$$m(\bar{u}) = 0$$

Proof. Since the homotopy h(t,z) induces a homotopy \bar{h} in Lagrangian-Grassmanian manifold. Note that $m(\bar{h}(0,\cdot))=0$. By the homotopy invariance of Maslov index, we know that $m(\bar{u})=0$.

Consider the partial differential equation

$$\bar{\partial}u + A(z)u = 0 \text{ on } D \tag{2.2}$$

$$u(z) \in \Gamma(z)R^n \text{ for } z \in \partial D$$
 (2.3)

$$\Gamma(z) \in GL(2n, R) \cap Sp(2n)$$
 (2.4)

$$m(\Gamma) = 0 \tag{2.5}$$

For $100 < k < \infty$ consider the Banach space \bar{V}_k consisting of all maps $u \in H^{k,2}(D, \mathbb{C}^n)$ such that $u(z) \in \Gamma(z)$ for almost all $z \in \partial D$. Let L_{k-1} the usual Hilbert space $H_{k-1}(D, \mathbb{C}^n)$.

We define an operator $P: \overline{V}_k \to L_{k-1}$ by

$$P(u) = \bar{\partial}u + Au \tag{2.6}$$

where D as in (2.1).

Proposition 2.2 $\bar{\partial}: \bar{V}_k \to L_{k-1}$ is a real linear Fredholm operator of index n.

Proof: see [4, 10, 21].

3 Nonlinear Fredholm Theory

3.1 Construction of Lagrangian Submanifold

Let M be an open manifold and $(T^*M, p_i dq_i)$ be the cotangent bundle of open manifold with the Liouville form $p_i dq_i$. Since M is open, there exists a function $g: M \to R$ without critical point. The translation by tTdg along the fibre gives a hamiltonnian isotopy of T^*M :

$$h_t^T(q,p) = (q, p + tTdg(q))$$
(3.1)

$$h_t^{T*}(p_i dq_i) = p_i dq_i + tT dg. (3.2)$$

Lemma 3.1 For any given compact set $K \subset T^*M$, there exists $T = T_K$ such that $h_1^T(K) \cap K = \emptyset$.

Proof. Similar to [10, 15]

Let $\Sigma \subset T^*M$ be a closed hypersurface, if there exists a vector field V defined in the neighbourhood U of Σ transversal to Σ such that $L_V\omega = \omega$, here $\omega = dp_i \wedge dq_i$ is a standard symplectic form on T^*M induced by the Liouville form $p_i dq_i$, we call Σ the contact manifold of induced type in T^*M with the induced contact form $\lambda = i_V\omega$.

Let (Σ, λ) be a contact manifold of induced type or Weinstein's type in T^*M with contact form λ and X its Reeb vector field, then X integrates to a Reeb flow η_s for $s \in R^1$.

By using the transversal vector field V, one can identify the neighbourhood U of Σ foliated by flow f_t of V and Σ , i.e., $U = \bigcup_t f_t(\Sigma)$ with the neighbourhood of $\{0\} \times \Sigma$ in the symplectization $R \times \Sigma$ by the exact symplectic transformation(see[16]).

Consider the form $d(e^a\lambda)$ at the point (a,x) on the manifold $(R \times \Sigma)$, then one can check that $d(e^a\lambda)$ is a symplectic form on $R \times \Sigma$. Moreover One can check that

$$i_X(e^a\lambda) = e^a \tag{3.3}$$

$$i_X(d(e^a\lambda)) = -de^a \tag{3.4}$$

So, the symplectization of Reeb vector field X is the Hamilton vector field of e^a with respect to the symplectic form $d(e^a\lambda)$. Therefore the Reeb flow lifts to the Hamilton flow h_s on $R \times \Sigma(\text{see}[3, 6])$.

Let \mathcal{L} be a closed Legendre submanifold in (Σ, λ) , i.e., there exists a smooth embedding $Q: \mathcal{L} \to \Sigma$ such that $Q^* \lambda|_{\mathcal{L}} = 0$. Let

$$(V', \omega') = (T^*M, dp_i \wedge dq_i)$$

and

$$W' = \mathcal{L} \times R, \quad W'_s = \mathcal{L} \times \{s\}$$
 (3.5)

define

$$G': W' \to V'$$

 $G'(w') = G'(l, s) = (0, \eta_s(Q(l)))$ (3.6)

Lemma 3.2 There does not exist any Reeb chord connecting Legendre submanifold \mathcal{L} in (Σ, λ) if and only if $G'(W'(s)) \cap G'(W'(s'))$ is empty for $s \neq s'$.

Proof. Obvious.

Lemma 3.3 If there does not exist any Reeb chord for $(X_{\lambda}, \mathcal{L})$ in (Σ, λ) then there exists a smooth embedding $G': W' \to V'$ with $G'(l, s) = (0, \eta_s(Q(l)))$ such that

$$G'_K: \mathcal{L} \times (-K, K) \to V'$$
 (3.7)

is a regular open Lagrangian embedding for any finite positive K.

Proof. One check

$$G'^*(d(e^a\lambda)) = \eta(\cdot, \cdot)^* d\lambda = (\eta_s^* d\lambda + i_X d\lambda \wedge ds) = 0$$
(3.8)

This implies that G' is a Lagrangian embedding, this proves Lemma 3.3.

Note that $d\lambda = dp_i \wedge dq_i$ on Σ by the definition of induced contact type and by assumption $[p_i dq_i - \lambda] = 0 \in H^1(\Sigma)$, we know that

$$p_i dq_i = \lambda + d\beta(\sigma) \text{ on } \Sigma$$
 (3.9)

Then by the proof of Lemma 3.3, one computes

$$G'^*(p_i dq_i) = G'^*(\lambda) + G'^* d\beta = \eta(\cdot, \cdot)^* \lambda + \eta(\cdot, \cdot)^* d\beta$$
$$= \eta_*^* \lambda + i_X \lambda ds + d\eta^* \beta = ds + d\beta$$

here we also use β denote the $\eta^*\beta$.

All above construction is contained in [16]. Now we introduce the upshot construction in [17]:

$$F'_0: \mathcal{L} \times R \times R \to (R \times \Sigma)$$

$$F'_0(((l, s, a) = (a, \eta_s(l)))$$
(3.10)

Now we embed a elliptic curve E long along s-axis and thin along a-axis such that $E \subset [-K,K] \times [0,\varepsilon]$. We parametrize the E by $t' \in S^1$.

Lemma 3.4 If there does not exist any Reeb chord in (Σ, λ) , then

$$F_0: \mathcal{L} \times S^1 \to (R \times \Sigma)$$

$$F_0(l, t') = (a(t'), \eta_{s(t')}(l))$$
(3.11)

is a compact Lagrangian submanifold. Moreover

$$l(R \times \Sigma, F_0(\mathcal{L} \times S^1), d(e^a \lambda))$$

$$= \inf \{ \int_D f^* de^a \lambda > 0 | f : (D, \partial D) \to (R \times \Sigma, F_0(\mathcal{L} \times S^1)) \}$$

$$= area(E)$$
(3.12)

Proof. We check that

$$F_0^*(e^a\lambda) = e^{a(t')}ds(t')$$
 (3.13)

So, F_0 is a Lagrangian embedding.

If the circle C homotopic to $C_1 \subset \mathcal{L} \times s_0$ then we compute

$$\int_{C} F_0^*(e^a \lambda) = \int_{C_1} F_0^*(e^a \lambda) = 0.$$
 (3.14)

since $\lambda | C_1 = 0$ due to $C_1 \subset \mathcal{L}$ and \mathcal{L} is Legendre submanifold. If the circle C homotopic to $C_1 \subset l_0 \times S^1$ then we compute

$$\int_C F_0^*(e^a \lambda) = \int_{C_1} F_0^*(e^{a(t')} ds(t')) = n(area(E)).$$
 (3.15)

This proves the Lemma.

Now we modify the above construction as follows:

$$F': \mathcal{L} \times R \times R \to ([0, \varepsilon] \times \Sigma) \subset T^*M$$

$$F'(l, s, a) = (a, \eta_s(l))$$
(3.16)

Now we embed a elliptic curve E long along s-axis and thin along b-axis such that $E \subset [-s_1, s_2] \times [0, \varepsilon]$. We parametrize the E by t'.

Lemma 3.5 If there does not exist any Reeb chord in (Σ, λ) , then

$$F: \mathcal{L} \times S^1 \to ([0, \varepsilon] \times \Sigma) \subset T^*M$$

$$F(l, t') = (a(t'), \eta_{s(t')}(l))$$
(3.17)

is a compact Lagrangian submanifold. Moreover

$$l(V', F(\mathcal{L} \times S^1), d(p_i dq_i)) = area(E)$$
(3.18)

Proof. We check that

$$F^*(p_i dq_i) = F^*(e^a \lambda + d\beta). \tag{3.19}$$

This proves the Lemma.

Now we construct an isotopy of Lagrangian embeddings as follows:

$$F': \mathcal{L} \times S^{1} \times [0, 1] \to V'$$

$$F'(l, t', t) = h_{t}^{T}(a(t'), \eta_{s(t')}(l))$$

$$F'_{t}(l, t') = F'(l, t', t). \tag{3.20}$$

Lemma 3.6 If there does not exist any Reeb chord for X_{λ} in (Σ, λ) then F' is an weakly exact isotopy of Lagrangian embeddings. Moreover for the choice of $T = T_{\Sigma}$ satisfying $[0, \varepsilon] \times \Sigma \cap h_1^T([0, \varepsilon] \times \Sigma) = \emptyset$, then $F'_0(\mathcal{L} \times S^1) \cap F'_1(\mathcal{L} \times S^1) = \emptyset$.

Proof. By Lemma 3.1-3.5 and below.

Let $(V', \omega') = (T^*M, dp_i \wedge dq_i)$, $W' = F(\mathcal{L} \times S^1)$, and $(V, \omega) = (V' \times C, \omega' \oplus \omega_0)$. As in [10], we use figure eight trick invented by Gromov to construct a Lagrangian submanifold in V through the Lagrange isotopy F' in V'. Fix a positive $\delta < 1$ and take a C^{∞} -map $\rho : S^1 \to [0, 1]$, where the circle S^1 is parametrized by $\Theta \in [-1, 1]$, such that the δ -neighborhood I_0 of $0 \in S^1$ goes to $0 \in [0, 1]$ and δ -neighbourhood I_1 of $\pm 1 \in S^1$ goes $1 \in [0, 1]$. Let $h_{\rho}^T(t, w') = h_{\rho(t)}^T(w')$ and

$$\tilde{l} = h_{\rho}^{T*}(p_{i}dq_{i}) = p_{i}dq_{i} - \rho(\Theta)Tdg$$

$$= e^{a(t')}ds(t') + d\beta) - \rho Tdg = e^{a(t')}ds(t') + d\beta + d\rho Tg) + Tgd\rho$$

$$= e^{a(t')}ds(t') + d\beta + d\rho Tg) - Tg\rho'(\Theta)d\Theta$$

$$= e^{a(t')}ds(t') + d\beta + d\rho Tg) - \Phi d\Theta \tag{3.21}$$

be the pull-back of the form $\tilde{l}' = e^{a(t')}ds(t') + d\beta + d\rho Tg) - \psi(s,t)dt$ to $W' \times S^1$ under the map $(w',\Theta) \to (w',\rho(\Theta))$ and assume without loss of generality Φ vanishes on $W' \times (I_0 \cup I_1)$. Since $[\tilde{l}']|W' \times \{t\} = [e^{a(t')}ds(t')]$ is independent of t, so F' is weakly exact. It is crucial here $|-\psi(s,t)| \leq M_0$ and M_0 is independent of area(E).

Next, consider a map α of the annulus $S^1 \times [\Phi_-, \Phi_+]$ into R^2 , where Φ_- and Φ_+ are the lower and the upper bound of the fuction Φ correspondingly, such that

- (i) The pull-back under α of the form $dx \wedge dy$ on R^2 equals $-d\Phi \wedge d\Theta$.
- (ii) The map α is bijective on $I \times [\Phi_-, \Phi_+]$ where $I \subset S^1$ is some closed subset, such that $I \cup I_0 \cup I_1 = S^1$; furthermore, the origin $0 \in R^2$ is a unique double point of the map α on $S^1 \times 0$, that is

$$0 = \alpha(0,0) = \alpha(\pm 1,0),$$

and α is injective on $S^1 = S^1 \times 0$ minus $\{0, \pm 1\}$.

(iii) The curve $S_0^1 = \alpha(S^1 \times 0) \subset R^2$ "bounds" zero area in R^2 , that is $\int_{S_0^1} x dy = 0$, for the 1-form x dy on R^2 .

Proposition 3.1 Let V', W' and F' as above. Then there exists an exact Lagrangian embedding $F: W' \times S^1 \to V' \times R^2$ given by $F(w', \Theta) = (F'(w', \rho(\Theta)), \alpha(\Theta, \Phi))$. Denote $F(W' \times S^1)$ by $W. W \subset T^*M \times B_{r_0}(0)$ with $4\pi r_0^2 = 8M_0$.

Proof. Similar to $[10, 2.3B_3]$.

3.2 Formulation of Hilbert manifolds

Let (Σ, λ) be a closed (2n-1)- dimensional manifold with a contact form λ of induced type in T^*M , it is well-known that T^*M is a Stein manifold, so it is

exausted by a proper pluri-subharmonic function. In fact since M is an open manifold one can take a proper Morse function g on M and let $f = \frac{|p|^2}{2} + \pi^* g$. Then f is pluri-subharmonic function on T^*M for some complex structure J' on T^*M tamed by $dp_i \wedge dq_i(\text{see}[5])$. Since Σ is compact and $W' = G'(\mathcal{L} \times R)$ is contained in Σ , by our construction we have W' is contained in a compact set f_c for c large enough.

Let $V' = T^*M$ and we choose an almost complex structure J' on T^*M tamed by $\omega' = dp_i \wedge dq_i$ and the metric $g' = \omega'(\cdot, J' \cdot)(\text{see}[10])$. By above discussion we know that all mechanism such as W' or Σ contained in $f_c = \{v' \in T^*M | f(v') \leq c\}$ for c large enough, i.e., contained in a compact set V'_c in T^*M . Then we expanding near $\partial f^{-1}(c)$ to get a complete exact symplectic manifold with a complete Riemann metric with injective radius $r_0 > 0(\text{see}[16])$.

In the following we denote by $(V, \omega) = (V' \times R^2, d(p_i dq_i) \oplus dx \wedge dy))$ with the metric $g = g' \oplus g_0$ induced by $\omega(\cdot, J \cdot)(J = J' \oplus i$ and $W \subset V$ a Lagrangian submanifold which was constructed in section 3.1, moreover we can slightly perturb the $J' \oplus i$ near p such that $J \oplus i$ is integrable near p.

Let

$$\mathcal{D}^k(V,W,p)=\{u\in H^k(D,V)|u(x)\in W\ a.e\ for\ x\in\partial D\ and\ u(1)=p\}$$
 for $k\geq 100.$

Lemma 3.7 Let W be a Lagrangian submanifold in V. Then,

$$\mathcal{D}^k(V, W, p) = \{ u \in H^k(D, V) | u(x) \in W \text{ a.e for } x \in \partial D \text{ and } u(1) = p \}$$

is a pseudo-Hilbert manifold with the tangent bundle

$$T\mathcal{D}^{k}(V, W, p) = \bigcup_{u \in \mathcal{D}^{k}(V, W, p)} \Lambda^{k-1}(u^{*}TV, u|_{\partial D}^{*}TW, p)$$
(3.22)

here

$$\Lambda^{k-1}(u^*TV,u|_{\partial D}^*TW,p) =$$

$$\{H^{k-1} - sections \ of \ (u^*(TV),(u|_{\partial D})^*TL) \ which \ vanishes \ at \ 1\}$$

Proof: See [4, 14].

Now we consider a section from $\mathcal{D}^k(V, W, p)$ to $T\mathcal{D}^k(V, W, p)$ follows as in [4, 10], i.e., let $\bar{\partial}: \mathcal{D}^k(V, W, p) \to T\mathcal{D}^k(V, W, p)$ be the Cauchy-Riemmann section

$$\bar{\partial}u = \frac{\partial u}{\partial s} + J\frac{\partial u}{\partial t} \tag{3.23}$$

for $u \in \mathcal{D}^k(V, W, p)$.

Theorem 3.1 The Cauchy-Riemann section $\bar{\partial}$ defined in (3.23) is a Fredholm section of Index zero.

Proof. According to the definition of the Fredholm section, we need to prove that $u \in \mathcal{D}^k(V, W, p)$, the linearization $D\bar{\partial}(u)$ of $\bar{\partial}$ at u is a linear Fredholm operator. Note that

$$D\bar{\partial}(u) = D\bar{\partial}_{[u]} \tag{3.24}$$

where

$$(D\bar{\partial}_{[u]})v = \frac{\partial v}{\partial s} + J\frac{\partial v}{\partial t} + A(u)v$$
(3.25)

with

$$v|_{\partial D} \in (u|_{\partial D})^*TW$$

here A(u) is $2n \times 2n$ matrix induced by the torsion of almost complex structure, see [4, 10] for the computation.

Observe that the linearization $D\bar{\partial}(u)$ of $\bar{\partial}$ at u is equivalent to the following Lagrangian boundary value problem

$$\frac{\partial v}{\partial s} + J \frac{\partial v}{\partial t} + A(u)v = f, \ v \in \Lambda^k(u^*TV)$$

$$v(t) \in T_{u(t)}W, \ t \in \partial D$$
(3.26)

One can check that (3.26) defines a linear Fredholm operator. In fact, by Proposition 2.2 and Lemma 2.1, since the operator A(u) is a compact, we know that the operator $\bar{\partial}$ is a nonlinear Fredholm operator of the index zero.

Definition 3.1 Let X be a Banach manifold and $P: Y \to X$ the Banach vector bundle. A Fredholm section $F: X \to Y$ is proper if $F^{-1}(0)$ is a compact set and is called generic if F intersects the zero section transversally, see [4, 7, 10].

Definition 3.2 $deg(F, y) = \sharp \{F^{-1}(0)\} mod 2$ is called the Fredholm degree of a Fredholm section (see[4, 7, 10]).

Theorem 3.2 The Fredholm section $F = \bar{\partial} : \mathcal{D}^k(V, W, p) \to T(\mathcal{D}^k(V, W, p))$ constructed in (3.23) is proper near $F^{-1}(0)$ and

$$deg(F,0) = 1$$

Proof: We assume that $u: D \mapsto V$ be a J-holomorphic disk with boundary $u(\partial D) \subset W$ and by the assumption that u is homotopic to the constant map $u_0(D) = p$. Since almost complex structure J tamed by the symplectic form ω , by stokes formula, we conclude $u: D \to V$ is a constant map. Because

u(1) = p, We know that $F^{-1}(0) = p$ which implies the properness. Next we show that the linearizatioon DF(p) of F at p is an isomorphism from $T_p\mathcal{D}(V,W,p)$ to E. This is equivalent to solve the equations

$$\frac{\partial v}{\partial s} + J \frac{\partial v}{\partial t} + Av = f$$

$$v|_{\partial D} \subset T_p W$$
(3.27)

here J = J(p) = i and A(=0) a constant zero matrix. By Lemma 2.1, we know that DF(p) is an isomorphism. Therefore deg(F,0) = 1.

4 Non-properness of a Fredholm section

In this section we shall construct a non-proper Fredholm section $F_1: \mathcal{D} \to E$ by perturbing the Cauchy-Riemann section as in [4, 10].

4.1 Anti-holomorphic section

Let $(V', \omega') = (T^*M, \omega_M)$ and $(V, \omega) = (V' \times C, \omega' \oplus \omega_0)$, and W as in section and $J = J' \oplus i$, $g = g' \oplus g_0$, g_0 the standard metric on C.

Now let $c \in C$ be a non-zero vector. We consider the equations

$$v = (v', f) : D \to V' \times C$$

$$\bar{\partial}_{J'}v' = 0, \bar{\partial}f = c$$

$$v|_{\partial D} : \partial D \to W$$
(4.1)

here v homotopic to constant map $\{p\}$ relative to W. Note that $W \subset V' \times B_{r_0}(0)$.

Lemma 4.1 Let v be the solutions of (4.1), then one has the following estimates

$$E(v) = \int_{D} \left(g'(\frac{\partial v'}{\partial x}, J'\frac{\partial v'}{\partial x}) + g'(\frac{\partial v'}{\partial y}, J'\frac{\partial v'}{\partial y})\right) + g_{0}(\frac{\partial f}{\partial x}, i\frac{\partial f}{\partial x}) + g_{0}(\frac{\partial f}{\partial y}, i\frac{\partial f}{\partial y})\right) d\sigma \leq 4\pi r_{0}^{2}.$$

$$(4.2)$$

Proof: Since v(z) = (v'(z), f(z)) satisfy (4.1) and $v(z) = (v'(z), f(z)) \in V' \times C$ is homotopic to constant map $v_0 : D \to \{p\} \subset W$ in (V, W), by the Stokes formula

$$\int_{D} v^{*}(\omega' \oplus \omega_{0}) = 0 \tag{4.3}$$

Note that the metric g is adapted to the symplectic form ω and J, i.e.,

$$g = \omega(\cdot, J\cdot) \tag{4.4}$$

By the simple algebraic computation, we have

$$\int_{D} v^* \omega = \frac{1}{4} \int_{D^2} (|\partial v|^2 - |\bar{\partial} v|^2) = 0$$
 (4.5)

and

$$|\nabla v|^2 = \frac{1}{2}(|\partial v|^2 + |\bar{\partial}v|^2) \tag{4.6}$$

Then

$$E(v) = \int_{D} |\nabla v|^{2}$$

$$= \int_{D} \{\frac{1}{2}(|\partial v|^{2} + |\bar{\partial}v|^{2})\}d\sigma$$

$$= \pi |c|_{g_{0}}^{2}$$

$$(4.7)$$

By the equations (4.1), one get

$$\bar{\partial}f = c \text{ on } D \tag{4.8}$$

We have

$$f(z) = \frac{1}{2}c\bar{z} + h(z) \tag{4.9}$$

here h(z) is a holomorphic function on D. Note that f(z) is smooth up to the boundary ∂D , then, by Cauchy integral formula

$$\int_{\partial D} f(z)dz = \frac{1}{2}c \int_{\partial D} \bar{z}dz + \int_{\partial D} h(z)dz$$
$$= \pi ic$$
(4.10)

So, we have

$$|c| = \frac{1}{\pi} \left| \int_{\partial D^2} f(z) dz \right| \tag{4.11}$$

Therefore,

$$E(v) \leq \pi |c|^{2} \leq \frac{1}{\pi} |\int_{\partial D} f(z) dz|^{2}$$

$$\leq \frac{1}{\pi} |\int_{\partial D} |f(z)| |dz||^{2}$$

$$\leq 4\pi |diam(pr_{2}(W))^{2}$$

$$\leq 4\pi r_{0}^{2}. \tag{4.12}$$

This finishes the proof of Lemma.

Proposition 4.1 For $|c| \geq 3r_0$, then the equations (4.1) has no solutions.

Proof. By (4.11), we have

$$|c| \leq \frac{1}{\pi} \int_{\partial D} |f(z)| |dz|$$

$$\leq \frac{1}{\pi} \int_{\partial D} diam(pr_2(W)) ||dz|$$

$$\leq 2r_0 \tag{4.13}$$

It follows that $c = 3r_0$ can not be obtained by any solutions.

4.2 Modification of section c

Note that the section c is not a section of the Hilbert bundle in section 3 since c is not tangent to the Lagrangian submanifold W, we must modify it as follows:

Let c as in section 4.1, we define

$$c_{\chi,\delta}(z,v) = \begin{cases} c & \text{if } |z| \le 1 - 2\delta, \\ 0 & \text{otherwise} \end{cases}$$
 (4.14)

Then by using the cut off function $\varphi_h(z)$ and its convolution with section $c_{\chi,\delta}$, we obtain a smooth section c_{δ} satisfying

$$c_{\delta}(z,v) = \begin{cases} c & \text{if } |z| \le 1 - 3\delta, \\ 0 & \text{if } |z| \ge 1 - \delta. \end{cases}$$

$$(4.15)$$

for h small enough, for the convolution theory see [12].

Now let $c \in C$ be a non-zero vector and c_{δ} the induced anti-holomorphic section. We consider the equations

$$v = (v', f) : D \to V' \times C$$

$$\bar{\partial}_{J'}v' = 0, \bar{\partial}f = c_{\delta}$$

$$v|_{\partial D} : \partial D \to W$$
(4.16)

Note that $W \subset V \times B_{r_0}(0)$. Then by repeating the same argument as section 4.1., we obtain

Lemma 4.2 Let v be the solutions of (4.16) and δ small enough, then one has the following estimates

$$E(v) \le 4\pi r_0^2. (4.17)$$

and

Proposition 4.2 For $|c| \geq 3r_0$, then the equations (4.16) has no solutions.

4.3 Modification of $J \oplus i$

Let (Σ, λ) be a closed contact manifold with a contact form λ of induced type in T^*M . Let J_M be an almost complex structure on T^*M and $J_1 = J_M \oplus i$ the almost complex structure on $T^*M \times R^2$ tamed by $\omega' \oplus \omega_0$. Let J_2 be any almost complex structure on $T^*M \times R^2$.

Now we consider the almost conplex structure on the symplectic fibration $D \times V \to D$ which will be discussed in detail in section 5.1., see also [10].

$$J_{\chi,\delta}(z,v) = \begin{cases} i \oplus J_M \oplus i & \text{if } |z| \le 1 - 2\delta, \\ i \oplus J_2 & \text{otherwise} \end{cases}$$
 (4.18)

Then by using the cut off function $\varphi_h(z)$ and its convolution with section $J_{\chi,\delta}$, we obtain a smooth section J_{δ} satisfying

$$J_{\delta}(z,v) = \begin{cases} i \oplus J_M \oplus i & \text{if } |z| \le 1 - 3\delta, \\ i \oplus J_2 & \text{if } |z| \ge 1 - \delta. \end{cases}$$
 (4.19)

as in section 4.2.

Then as in section 4.2, one can also reformulation of the equations (4.16) and get similar estimates of Cauchy-Riemann equations, we leave it as exercises to reader.

Theorem 4.1 The Fredholm sections $F_1 = \bar{\partial} + c_{\delta} : \mathcal{D}^k(V, W, p) \to T(\mathcal{D}^k(V, W, p))$ is not proper for |c| large enough.

Proof. See [4, 10].

5 J-holomorphic section

Recall that $W \subset T^*M \times B_{r_0}(0)$ as in section 3. The Riemann metric g on $M \times R^2$ induces a metric g|W.

Now let $c \in C$ be a non-zero vector and c_{δ} the induced anti-holomorphic section. We consider the nonlinear inhomogeneous equations (4.16) and transform it into \bar{J} -holomorphic map by considering its graph as in [4, 10].

Denote by $Y^{(1)} \to D \times V$ the bundle of homomorphisms $T_s(D) \to T_v(V)$. If D and V are given the disk and the almost Kähler manifold, then we distinguish the subbundle $X^{(1)} \subset Y^{(1)}$ which consists of complex linear homomorphisms and we denote $\bar{X}^{(1)} \to D \times V$ the quotient bundle $Y^{(1)}/X^{(1)}$. Now, we assign to each C^1 -map $v: D \to V$ the section $\bar{\partial}v$ of the bundle

 $\bar{X}^{(1)}$ over the graph $\Gamma_v \subset D \times V$ by composing the differential of v with the quotient homomorphism $Y^{(1)} \to \bar{X}^{(1)}$. If $c_\delta : D \times V \to \bar{X}$ is a H^k - section we write $\bar{\partial}v = c_\delta$ for the equation $\bar{\partial}v = c_\delta|\Gamma_v$.

Lemma 5.1 (Gromov[10]) There exists a unique almost complex structure J_g on $D \times V$ (which also depends on the given structures in D and in V), such that the (germs of) J_{δ} -holomorphic sections $v:D\to D\times V$ are exactly and only the solutions of the equations $\bar{\partial}v=c_{\delta}$. Furthermore, the fibres $z\times V\subset D\times V$ are J_{δ} -holomorphic (i.e. the subbundles $T(z\times V)\subset T(D\times V)$ are J_{δ} -complex) and the structure $J_{\delta}|z\times V$ equals the original structure on $V=z\times V$. Moreover J_{δ} is tamed by $k\omega_0\oplus \omega$ for k large enough which is independent of δ .

6 Proof of Theorem 1.1

Theorem 6.1 There exists a non-constant J-holomorphic map $u:(D,\partial D) \to (T^*M \times C, W)$ with $E(u) \leq 4\pi r_0^2$.

Proof. By Gromov's C^0 -converngence theorem and the results in section 4 shows the solutions of equations (4.16) must denegerate to a cusp curves, i.e., we obtain a Sacks-Uhlenbeck's bubble, i.e., J-holomorphic sphere or disk with boundary in W, the exactness of the symplectic form on $T^*M \times R^2$ rules out the possibility of J-holomorphic sphere. For the more detail, see the proof of Theorem 2.3.B in [10].

Proof of Theorem 1.1. By Theorem 6.1, we know that

$$l(V', F(\mathcal{L} \times S^1), d(p_i dq_i)) = area(E) \le 4\pi r_0^2$$
(6.1)

But if K large enough, $area(E) > 8\pi r_0^2$. This implies the assumption that \mathcal{L} has no self-intersection point under Reeb flow does not hold.

References

- [1] Abbas, C., Finite energy surfaces and the chord problem, Duke Math. Journ., Vol96, No.2(1999), 241-316.
- [2] Arnold, V. I., First steps in symplectic topology, Russian Math. Surveys 41(1986),1-21.
- [3] Arnold, V.& Givental, A., Symplectic Geometry, in: Dynamical Systems IV, edited by V. I. Arnold and S. P. Novikov, Springer-Verlag, 1985.

- [4] Audin, M& Lafontaine, J., eds.: Holomorphic Curves in Symplectic Geometry. Progr. Math. 117, (1994) Birkhaüser, Boston.
- [5] Eliashberg, Y.& Gromov, M., Convex symplectic manifolds, Pro. of Sym. in Pure Math., vol. 52(1991), Part2, 135-162.
- [6] Eliashberg, Y.& Gromov, M., Lagrangian Intersection Theory: Finite-Dimensional Approach, Amer. Math. Soc. Transl. 186(1998): 27-118.
- [7] Floer, A., Hofer, H.& Viterbo, C., The Weinstein conjecture in $P \times C^l$, Math.Z. 203(1990)469-482.
- [8] Givental, A. B., Nonlinear generalization of the Maslov index, Adv. in Sov. Math., V.1, AMS, Providence, RI, 1990.
- [9] Gray, J.W., Some global properties of contact structures. Ann. of Math., 2(69): 421-450, 1959.
- [10] Gromov, M., Pseudoholomorphic Curves in Symplectic manifolds. Inv. Math. 82(1985), 307-347.
- [11] Hofer, H., Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjection in dimension three. Inventions Math., 114(1993), 515-563.
- [12] Hörmander, L., The Analysis of Linear Partial Differential Operators I, Springer-Verlag, 1983.
- [13] Jost, J., Two-Dimensional Geometric Variational Problems, A Wiley-Interscience Publication.
- [14] Klingenberg, K., Lectures on closed Geodesics, Grundlehren der Math. Wissenschaften, vol 230, Spinger-Verlag, 1978.
- [15] Lalonde, F & Sikorav, J.C., Sous-Varitès Lagrangiennes et lagrangiennes exactes des fibrès cotangents, Comment. Math. Helvetici 66(1991) 18-33.
- [16] Ma, R., Legendrian submanifolds and A Proof on Chord Conjecture, Boundary Value Problems, Integral Equations and Related Problems, edited by J K Lu & G C Wen, World Scientific, 135-142,2000.
- [17] Mohnke, K.: Holomorphic Disks and the Chord Conjecture, Annals of Math., (2001), 154:219-222.
- [18] Sacks, J. and Uhlenbeck, K., The existence of minimal 2-spheres. Ann. Math., 113:1-24, 1983.

- [19] Smale, S., An infinite dimensional version of Sard's theorem, Amer. J. Math. 87: 861-866, 1965.
- [20] Thurston, W., The theory of foliations in codimension greater than one, Comm. Math. Helv. 214-231, 49(1974).
- [21] Wendland, W., Elliptic systems in the plane, Monographs and studies in Mathematics 3, Pitman, London-San Francisco, 1979.