C*-ALGEBRAS ASSOCIATED WITH COMPLEX DYNAMICAL SYSTEMS

TSUYOSHI KAJIWARA AND YASUO WATATANI

ABSTRACT. Iteration of a rational function R gives a complex dynamical system on the Riemann sphere. We introduce a C^* -algebra \mathcal{O}_R associated with R as a Cuntz-Pimsner algebra of a Hilbert bimodule over the algebra $A = C(J_R)$ of continuous functions on the Julia set J_R of R. The algebra \mathcal{O}_R is a certain analog of the crossed product by a boundary action. We show that if the degree of Ris at least two, then C^* -algebra \mathcal{O}_R is simple and purely infinite. For example if $R(z) = z^2 - 2$, then the Julia set $J_R = [-2, 2]$ and the restriction $R: J_R \to J_R$ is topologically conjugate to the tent map on [0, 1]. The algebra \mathcal{O}_{z^2-2} is isomorphic to the Cuntz algebra \mathcal{O}_{∞} . We also show that the Lyubich measure associated with R gives a unique KMS state on the C^* -algebra \mathcal{O}_R for the gauge action at inverse temperature log(deg R) if the Julia set contains no critical points.

1. INTRODUCTION

For a branched covering $\pi: M \to M$, Deaconu and Muhly [5] introduced a C^{*}-algebra $C^*(M,\pi)$ as the C^{*}-algebra of the r-discrete groupoid constructed by Renault [25]. In particular they consider rational functions on the Riemann sphere \mathbb{C} and compute the K-groups of the C^* -algebra. See also the previous work [4] by Deaconu on C^* -algebras associated with continuous graphs and Anantharaman-Delaroche's work [1] on purely infinite C^* -algebras for expansive dynamical systems. Although Deaconu and Muhly's work itself is interesting, we introduce a slightly different C^* -algebras $\mathcal{O}_R(\hat{\mathbb{C}}), \mathcal{O}_R = \mathcal{O}_R(J_R)$ and $\mathcal{O}_R(F_R)$ associated with a rational function R on the Riemann sphere, the Julia set J_R and the Fatou set F_R of R in this note. The C^{*}-algebra $\mathcal{O}_R(\mathbb{C})$ is defined as a Cuntz-Pimsner algebra of the Hilbert bimodule $Y = C(\operatorname{graph} R)$ over $B = C(\widehat{\mathbb{C}})$. The C^{*}-algebra \mathcal{O}_R is defined as a Cuntz-Pimsner algebra of the Hilbert bimodule $X = C(\operatorname{graph} R|_{J_R})$ over $A = C(J_R)$. And the C*-algebra $\mathcal{O}_R(F_R)$ is similarly defined. The difference between Deaconu and Muhly's construction and ours is the following: They exclude branched points to construct their groupoids. We include branched points to construct our bimodules. An advantage of theirs is that their C^* -algebra is constructed from both a groupoid and a bimodule. An advantage of ours is that our C^* -algebra \mathcal{O}_R is always simple and purely infinite if the degree of R is at least two.

For example, if $R(z) = z^2 - 2$, then the Julia set $J_R = [-2, 2]$ and the restriction $R|_{J_R} : J_R \to J_R$ is topologically conjugate to the tent map $h : [0,1] \to [0,1]$. Then their C^* -algebra $C^*([0,1],h)$ is not simple and $K_0(C^*([0,1],h)) = \mathbb{Z} \oplus \mathbb{Z}$ and $K_1(C^*([0,1],h)) = \{0\}$. Our C^* algebra \mathcal{O}_{z^2-2} is simple and purely infinite and $K_0(\mathcal{O}_{z^2-2}) = \mathbb{Z}$ and $K_1(\mathcal{O}_{z^2-2}) = \{0\}$. In fact the algebra \mathcal{O}_{z^2-2} is isomorphic to the Cuntz algebra \mathcal{O}_{∞} .

Even if R is a quadratic polynomial $R(z) = z^2 - c$, the structure of the C^* -algebra \mathcal{O}_R is closely related to the property of R as the complex dynamical system. If c is not in the Mandelbrot set \mathcal{M} , then C^* -algebra \mathcal{O}_R is isomorphic to the Cuntz algebra \mathcal{O}_2 . If c is in the interior of the main cardioid, then C^* -algebra \mathcal{O}_R is not isomorphic to \mathcal{O}_2 . In fact we have $K_0(\mathcal{O}_R) = \mathbb{Z}$ and $K_1(\mathcal{O}_R) = \mathbb{Z}$.

We compare our construction with other important constructions by C. Delaroche [1] and M. Laca - J. Spielberg [19]. They showed that a certain boundary action of a Kleinian group on the limit set yields a simple nuclear purely infinite C^* -algebra as groupoid C^* -algebra or crossed product. Recall that Sullivan's dictionary says that there is a strong analogy between the limit set Λ_{Γ} of a Kleinian group Γ and the Julia set J_R of a rational function R. The algebra \mathcal{O}_R is generated by $C(J_R)$ and $\{S_f; f \in C(\operatorname{graph} R|_{J_R})\}$. We regard the algebra \mathcal{O}_R as a certain analog of the crossed product $C(\Lambda_{\Gamma}) \rtimes \Gamma$ of $C(\Lambda_{\Gamma})$ by a boundary action of a Klein group Γ . In fact the crossed product is generated by $C(\Lambda_{\Gamma})$ and $\{\lambda_g : g \in \Gamma\}$. Moreover commutation relations $\alpha(a)S_f = S_f a$ for $a \in C(J_R)$ and $\alpha_g(a)\lambda_g = \lambda_g a$ for $a \in C(\Lambda_{\Gamma})$ are similar, where $\alpha(a)(x) = a(R(x))$ and $\alpha_g(a)(x) = a(g^{-1}x)$.

Applying a result by Fowler, Muhly and Raeburn [11], the quotient algebra $\mathcal{O}_R(\hat{\mathbb{C}})/\mathcal{I}(I)$ by the ideal $\mathcal{I}(I)$ corresponding to the Fatou set is canonically isomorphic to $\mathcal{O}_R = \mathcal{O}_R(J_R)$.

Several people [6], [21], [26], [12] considered conditions for simplicity of Cuntz-Pimsner algebras. We directly show that C^* -algebra \mathcal{O}_R is simple purely infinite through analyzing branched coverings. A criterion by Schweizer [26] can be applied to show the only simplicity of C^* -algebra \mathcal{O}_R . C^* -algebra \mathcal{O}_R is separable and nuclear, and belongs to the UCT class. Thus if deg $R \geq 2$, then the isomorphisms class of \mathcal{O}_R is completely determined by the K-theory together with the class of the unit by the classification theorem by Kirchberg-Phillips [17], [22]. If there exist no critical points in J_R , then simplicity of \mathcal{O}_R is also given by the simplicity of a certain crossed product by an endomorhism with a transfer operator in Exel-Vershik [9], [8]. Katsura [16] studies his new construction which contains \mathcal{O}_R if there exist no critical points in J_R . A difficulty of our analysis stems from the fact that the Julia set J_R contains the critical points (i.e. the branched points.) Many examples of Cuntz-Pimsner algebras arise from Hilbert bimodules which are finitely generated and projective as right module and the image of the left action is contained in the compacts. Another extremal case is studied by A. Kumjian in [18], where the image of the left action has trivial intersection with the compacts. Our case is a third one and the intersection of the image of the left action and the compacts is exactly represented by the branched points, i.e., critical points of R. Hence our study is focused on the behaviour of the branched points under iteration.

We also showed that the Lyubich measure associated with R gives a unique KMS state on the C^* -algebra \mathcal{O}_R for the gauge action at inverse temperature log deg R if the Julia set contains no critical points. But the problem of KMS state is subtle and difficult, if the Julia set contains critical points. We will discuss the case elsewhere.

We should note that B. Brenken also announced a study of Cuntz-Pimsner algebras associated with branched coverings in a satellite conference at Chenge, China in 2002, when we announced a content of the paper in the same conference.

2. RATIONAL FUNCTIONS AND HILBERT BIMODULES

We recall some facts on iteration of rational functions. Let R be a rational function of the form $R(z) = \frac{P(z)}{Q(z)}$ with relative prime polynomials P and Q. The degree of R is denoted by $d = \deg R :=$ max{deg P, deg Q}. If deg R = 1 and the Julia set is not empty, then it is, in fact, one point. Therefore the classical Toeplitz algebra appears and the C^* -algebra \mathcal{O}_R becomes $C(\mathbb{T})$. Hence we need to assume that deg $R \geq 2$ to consider non-trivial ones. We regard a rational function R as a d-fold branched covering map $R : \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ on the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. The sequence $(R^n)_n$ of iterations of R gives a complex analytic dynamical system on $\hat{\mathbb{C}}$. The Fatou set F_R of R is the maximal open subset of $\hat{\mathbb{C}}$ on which $(R^n)_n$ is equicontinuous (or a normal family), and the Julia set J_R of R is the complement of the Fatou set in $\hat{\mathbb{C}}$.

We always assume that a rational function R is not a constant function. Recall that a *critical point* of R is a point z_0 at which R is not locally one to one. It is a zero of R' or a pole of R of order two or higher. The image $w_0 = R(z_0)$ is called a *critical value* of R. Using appropriate local charts, if $R(z) = w_0 + c(z - z_0)^n + (\text{higher terms})$ with $n \ge 1$ and $c \ne 0$ on some neighborhood of z_0 , then the integer $n = e(z_0) = e_R(z_0)$ is called the *branch index* of R at z_0 . Thus $e(z_0) \ge 2$ if z_0 is a critical point and $e(z_0) = 1$ otherwise. Therefore R is an $e(z_0) : 1$ map in a punctured neighborhood of z_0 . By the Riemann-Hurwitz formula, there exist 2d-2 critical points counted with multiplicity, that is,

$$\sum_{z \in \hat{\mathbb{C}}} e(z) - 1 = 2 \deg R - 2$$

Furthermore for each $w \in \hat{\mathbb{C}}$, we have

z

$$\sum_{\in R^{-1}(w)} e(z) = \deg R$$

Let C be the set of critical points of R and R(C) be the set of the critical values of R. We put $B = R^{-1}(R(C))$. Then the restriction $R : \hat{\mathbb{C}} \setminus B \to \hat{\mathbb{C}} \setminus R(C)$ is a d : 1 regular covering, where $d = \deg R$. This means that any point $y \in \hat{\mathbb{C}} \setminus R(C)$ has an open neighborhood V such that $R^{-1}(V)$ has d connected components U_1, \ldots, U_d and the restriction $R|_{U_k} : U_k \to V$ is a homeomorphism for $k = 1, \ldots, d$. Thus R has d analytic local cross sections $S_k = (R|_{U_k})^{-1}$.

We recall Cuntz-Pimsner algebras [23]. Let A be a C^* -algebra and X be a Hilbert right A-module. We denote by L(X) be the algebra of the adjointable bounded operators on X. For ξ , $\eta \in X$, the "rank one" operator $\theta_{\xi,\eta}$ is defined by $\theta_{\xi,\eta}(\zeta) = \xi(\eta|\zeta)$ for $\zeta \in X$. The closure of the linear span of rank one operators is denoted by K(X). We call that X is a Hilbert bimodule over A if X is a Hilbert right A-module with a homomorphism $\phi : A \to L(X)$. We assume that X is full and ϕ is injective.

Let $F(X) = \bigoplus_{n=0}^{\infty} X^{\otimes n}$ be the Fock module of X with a convention $X^{\otimes 0} = A$. For $\xi \in X$, the creation operator $T_{\xi} \in L(F(X))$ is defined by

$$T_{\xi}(a) = \xi a$$
 and $T_{\xi}(\xi_1 \otimes \cdots \otimes \xi_n) = \xi \otimes \xi_1 \otimes \cdots \otimes \xi_n$.

We define $i_{F(X)} : A \to L(F(X))$ by

$$i_{F(X)}(a)(b) = ab$$
 and $i_{F(X)}(a)(\xi_1 \otimes \cdots \otimes \xi_n) = \phi(a)\xi_1 \otimes \cdots \otimes \xi_n$

for $a, b \in A$. The Cuntz-Toeplitz algebra \mathcal{T}_X is the C^* -algebra on F(X) generated by $i_{F(X)}(a)$ with $a \in A$ and T_{ξ} with $\xi \in X$. Let $j_K : K(X) \to \mathcal{T}_X$ be the homomorphism defined by $j_K(\theta_{\xi,\eta}) = T_{\xi}T_{\eta}^*$. We consider the ideal $I_X := \phi^{-1}(K(X))$ of A. Let \mathcal{J}_X be the ideal of \mathcal{T}_X generated by $\{i_{F(X)}(a) - (j_K \circ \phi)(a); a \in I_X\}$. Then the Cuntz-Pimsner algebra \mathcal{O}_X is the the quotient $\mathcal{T}_X/\mathcal{J}_X$. Let $\pi : \mathcal{T}_X \to \mathcal{O}_X$ be the quotient map. Put $S_{\xi} = \pi(T_{\xi})$ and $i(a) = \pi(i_{F(X)}(a))$. Let $i_K : K(X) \to \mathcal{O}_X$ be the homomorphism defined by $i_K(\theta_{\xi,\eta}) = S_{\xi}S_{\eta}^*$. Then $\pi((j_K \circ \phi)(a)) = (i_K \circ \phi)(a)$ for $a \in I_X$. We note that the Cuntz-Pimsner algebra \mathcal{O}_X is the universal C^* -algebra generated by i(a) with $a \in A$ and S_{ξ} with $\xi \in X$ satisfying that $i(a)S_{\xi} = S_{\phi(a)\xi}, S_{\xi}i(a) = S_{\xi a}, S_{\xi}^*S_{\eta} = i((\xi|\eta)_A)$ for $a \in A, \xi, \eta \in X$ and $i(a) = (i_K \circ \phi)(a)$ for $a \in I_X$. We usually identify i(a) with a in A. We denote by \mathcal{O}_X^{alg} the *-algebra generated algebraically by A and S_{ξ} with $\xi \in X$. There exists an

action $\gamma : \mathbb{R} \to Aut \ \mathcal{O}_X$ with $\gamma_t(S_{\xi}) = e^{it}S_{\xi}$, which is called the gauge action. Since we assume that $\phi : A \to L(X)$ is isometric, there is an embedding $\phi_n : L(X^{\otimes n}) \to L(X^{\otimes n+1})$ with $\phi_n(T) = T \otimes id_X$ for $T \in L(X^{\otimes n})$ with the convention $\phi_0 = \phi : A \to L(X)$. We denote by \mathcal{F}_X the C^* -algebra generated by all $K(X^{\otimes n})$, $n \ge 0$ in the inductive limit algebra $\varinjlim L(X^{\otimes n})$. Let \mathcal{F}_n be the C^* -subalgebra of \mathcal{F}_X generated by $K(X^{\otimes k})$, $\overrightarrow{k} = 0, 1, \ldots, n$, with the convention $\mathcal{F}_0 = A = K(X^{\otimes 0})$. Then $\mathcal{F}_X = \varinjlim \mathcal{F}_n$.

Let graph $R = \{(x, y) \in \hat{\mathbb{C}}^2; y = R(x)\}$ be the graph of a rational function R. Consider a C^* -algebra $B = C(\hat{\mathbb{C}})$. Let $Y = C(\operatorname{graph} R)$. Then Y is a B-B bimodule by

$$(a \cdot f \cdot b)(x, y) = a(x)f(x, y)b(y)$$

for $a, b \in B$ and $f \in Y$. We introduce a *B*-valued inner product $(|)_B$ on *Y* by

$$(f|g)_B(y) = \sum_{x \in R^{-1}(y)} e(x)\overline{f(x,y)}g(x,y)$$

for $f, g \in Y$ and $y \in \hat{\mathbb{C}}$. We need branch index e(x) in the formula of the inner product above. Put $||f||_2 = ||(f|f)_B||_{\infty}$.

Lemma 2.1. The above *B*-valued inner product is well defined, that is, $\hat{\mathbb{C}} \ni y \mapsto (f|g)_B(y) \in \mathbb{C}$ is continuous.

Proof. Let $d = \deg R$. If y_0 is not a critical value of R, then there are d distinct continuous cross sections, say S_1, \ldots, S_d , of R defined on some open neighbourhood V of y_0 such that V contains no critical values of R. Then for $y \in V$,

$$(f|g)_B(y) = \sum_{i=1}^d \overline{f(S_i(y), y)}g(S_i(y), y).$$

Thus, $(f|g)_B(y)$ is continuous at y_0 . Next consider the case that y_0 is a critical value of R. Let x_1, \ldots, x_r be the distinct points of $R^{-1}(y_0)$ Then there exist open neighbourhoods V of y_0 and U_j of x_j such that $R : U_j \to V$ is a $e(x_j) : 1$ branched covering and is expressed as $f(z) = z^{e(x_j)}$ by local charts for $j = 1, \ldots, r$. Consider $S_k^{(j)}(z) =$ $r^{1/e(x_j)} \exp(i(\frac{\theta}{e(x_j)} + \frac{2k\pi}{e(x_j)}))$ for $z = r \exp(i\theta)$ and $S_k^{(j)}(0) = 0$ under the local coordinate. Thus there exist cross sections, say $S_k^{(j)}$ of R $k = 1, \ldots, e(x_j)$ for each $j = 1, \ldots, r$ defined on U of y_0 such any $S_k^{(j)}$ is continuous at least at y_0 but not necessarily continuous on V. Then for $y \in V$,

$$\lim_{y \to y_0} (f|g)_B(y) = \lim_{y \to y_0} \sum_{j=1}^r \sum_{k=1}^{e(x_j)} \overline{f(S_k^{(j)}(y), y)} g(S_k^{(j)}(y), y)$$
$$= \sum_{j=1}^r \sum_{k=1}^{e(x_j)} \overline{f(S_k^{(j)}(y_0), y_0)} g(S_k^{(j)}(y_0), y_0)$$
$$= \sum_{j=1}^r e(x_j) \overline{f(x_j, y_0)} g(x_j, y_0) = (f|g)_B(y_0)$$

Thus $(f|g)_B(y)$ is continuous at y_0 .

The left multiplication of B on Y gives the left action $\phi : B \to L(Y)$ such that $(\phi(b)f)(x,y) = b(x)f(x,y)$ for $b \in B$ and $f \in Y$.

Proposition 2.2. If R is a rational function, then $Y = C(\operatorname{graph} R)$ is a full Hilbert bimodule over $B = C(\hat{\mathbb{C}})$ without completion. The left action $\phi : B \to L(Y)$ is unital and faithful.

Proof. Let $d = \deg R$. For any $f \in Y = C(\operatorname{graph} R)$, we have

$$||f||_{\infty} \le ||f||_{2} = (\sup_{y} \sum_{x \in R^{-1}(y)} e(x)|f(x,y)|^{2})^{1/2} \le \sqrt{d} ||f||_{\infty}$$

Therefore two norms $\| \|_2$ and $\| \|_{\infty}$ are equivalent. Since $C(\operatorname{graph} R)$ is complete with respect to $\| \|_{\infty}$, it is also complete with respect to $\| \|_2$.

Since $(1|1)_B(y) = \sum_{x \in R^{-1}(y)} e(x)1 = \deg R$, $(Y|Y)_B$ contains the identity I_B of B. Therefore Y is full. If $b \in B$ is not zero, then there exists $x_0 \in \hat{\mathbb{C}}$ with $b(x_0) \neq 0$. Choose $f \in Y$ with $f(x_0, R(x_0)) \neq 0$. Then $\phi(b)f \neq 0$. Thus ϕ is faithful.

Since the Julia set J_R is completely invariant, i.e., $R(J_R) = J_R = R^{-1}(J_R)$, we can consider the restriction $R|_{J_R} : J_R \to J_R$, which will be often denoted by the same letter R. Let graph $R|_{J_R} = \{(x, y) \in J_R \times J_R ; y = R(x)\}$ be the graph of the restriction map $R|_{J_R}$. Let $A = C(J_R)$ and $X = C(\operatorname{graph} R|_{J_R})$. Through a restriction of the above action, X is an A-A bimodule.

Corollary 2.3. Let R be a rational function with $J_R \neq \phi$, for example, deg $R \geq 2$. Then $X = C(\operatorname{graph} R|_{J_R})$ is a full Hilbert bimodule over $A = C(J_R)$ without completion, where A-valued inner product $(|)_A$ on X is given by

$$(f|g)_A(y) = \sum_{\substack{x \in R^{-1}(y) \\ 6}} e(x)\overline{f(x,y)}g(x,y)$$

for $f, g \in X$ and $y \in J_R$. The left action $\phi : A \to L(X)$ is unital and faithful.

Proof. It is an immediate consequence of the fact that $R(J_R) = J_R = R^{-1}(J_R)$.

Definition. We introduce the C^* -algebras $\mathcal{O}_R(\hat{\mathbb{C}})$, $\mathcal{O}_R(J_R)$ and $\mathcal{O}_R(F_R)$ associated with a rational function R. The C^* -algebra $\mathcal{O}_R(\hat{\mathbb{C}})$ is defined as a Cuntz-Pimsner algebra of the Hilbert bimodule $Y = C(\operatorname{graph} R)$ over $B = C(\hat{\mathbb{C}})$. When the Julia set J_R is not empty, for example deg $R \geq 2$, we introduce the C^* -algebra $\mathcal{O}_R(J_R)$ as a Cuntz-Pimsner algebra of the Hilbert bimodule $X = C(\operatorname{graph} R|_{J_R})$ over $A = C(J_R)$. When the Fatou set F_R is not empty, the C^* -algebra $\mathcal{O}_R(F_R)$ is defined similarly. Sullivan's dictionary says that there is a strong analogy between the limit set of a Kleinian group and the Julia set of a rational function. Therefore we simply denote by \mathcal{O}_R the C^* -algebra $\mathcal{O}_R(J_R)$ to emphasize the analogy.

Proposition 2.4. Let R be a rational function and $Y = C(\operatorname{graph} R)$ be a Hilbert bimodule over $B = C(\hat{\mathbb{C}})$. Then there exists an isomorphism

$$\varphi: Y^{\otimes n} \to C(\operatorname{graph} R^n)$$

as a Hilbert bimodule over B such that

$$(\varphi(f_1 \otimes \cdots \otimes f_n))(x, R^n(x))$$

= $f_1(x, R(x))f_2(R(x), R^2(x)) \dots f_n(R^{n-1}(x), R^n(x))$

for $f_1, \ldots, f_n \in Y$ and $x \in \hat{\mathbb{C}}$. Moreover when the Julia set J_R is not empty, let $A = C(J_R)$ and $X = C(\operatorname{graph} R|_{J_R})$. Then we have a similar isomorphism

$$\varphi: X^{\otimes n} \to C(\operatorname{graph} R^n|_{J_R})$$

as a Hilbert bimodule over A. Here we use the same symbol φ to save the notation.

Proof. It is easy to see that φ is well-defined and a bimodule homomorphism. We show that φ preserves inner product. The point is to use the following chain rule for branch index:

$$e_{R^n}(x) = e_R(x)e_R(R(x))\dots e_R(R^{n-1}(x))$$

7

Consider the case when n = 2 for simplicity of the notation.

$$\begin{aligned} &(f_1 \otimes f_2 | g_1 \otimes g_2)_B(x_2) = (f_2 | (f_1 | g_1)_B g_2)_B(x_2) \\ &= \sum_{x_1 \in R^{-1}(x_2)} e_R(x_1) \overline{f_2(x_1, x_2)} (f_1 | g_1)_B(x_1) g_2(x_1, x_2) \\ &= \sum_{x_1 \in R^{-1}(x_2)} e_R(x_1) \overline{f_2(x_1, x_2)} (\sum_{x_0 \in R^{-1}(x_1)} e_R(x_0) \overline{f_1(x_0, x_1)} g_1(x_0, x_1) g_2(x_1, x_2)) \\ &= \sum_{x_0 \in R^{-2}(x_2)} e_{R^2}(x_0) \overline{f_1(x_0, R(x_0))} f_2(R(x_0), x_2) g_1(x_0, R(x_0)) g_2(R(x_0), x_2) \\ &= \sum_{x_0 \in R^{-2}(x_2)} e_{R^2}(x_0) \overline{(\varphi(f_1 \otimes f_2))(x_0, x_2)} (\varphi(g_1 \otimes g_2))(x_0, x_2) \\ &= (\varphi(f_1 \otimes f_2) | \varphi(g_1 \otimes g_2))(x_2) \end{aligned}$$

Since φ preserves inner product, φ is one to one. The non-trivial one is to show that φ is onto. Since the image of φ is a *-subalgebra of $C(\hat{\mathbb{C}})$ and separates the two points, the image of φ is dense in $C(\hat{\mathbb{C}})$ with respect to $\| \|_{\infty}$ by the Stone-Wierstrass Theorem. Since two norms $\| \|_2$ and $\| \|_{\infty}$ are equivalent and φ is isometric with respect to $\| \|_2$, φ is onto.

We recall that the Julia set $J|_R$ of R is completely invariant under R and the Julia set of R coincides with the Julia set of R^n . Therefore the proof is valid for A and X also.

The set C of critical points of R is described by the ideal $I_Y = \phi^{-1}(K(Y))$ of B.

Proposition 2.5. Let R be a rational function and $Y = C(\operatorname{graph} R)$ be a Hilbert bimodule over $B = C(\widehat{\mathbb{C}})$. Similarly let $A = C(J_R)$ and $X = C(\operatorname{graph} R|_{J_R})$. Consider the ideal $I_Y = \phi^{-1}(K(Y))$ of B and the ideal $I_X = \phi^{-1}(K(X))$ of A. Then $I_Y = \{b \in B; b \text{ vanishes on } C\}$. and $I_X = \{a \in A; a \text{ vanishes on } C \cap J_R\}$.

Proof. Let $d = \deg R$, then there exist 2d - 2 critical points counted with multiplicity. Therefore C is a finite non-empty set. Firstly, let us take $b \in B$ with a compact support $S = \operatorname{supp}(b)$ in $\widehat{\mathbb{C}} \setminus C$. For any $x \in S$, since x is not a critical point, there exists an open neighbourhood U_x of x such that $U_x \cap C = \phi$ and the restriction $R|_{U_x} : U_x \to R(U_x)$ is a homeomorphism. Since S is compact, there exists a finite subset $\{x_1, \ldots, x_m\}$ such that $S \subset \bigcup_{i=1}^m U_{x_i} \subset \widehat{\mathbb{C}} \setminus C$. Let $(f_i)_i$ be a finite family in $C(\widehat{\mathbb{C}})$ such that $0 \leq f_i \leq 1$, $\operatorname{supp}(f_i) \subset U_{x_i}$ for $i = 1, \ldots, m$ and $\sum_{i=1}^m f_i(x) = 1$ for $x \in S$. Define $\xi_i, \eta_i \in C(\operatorname{graph} R)$ by $\xi_i(x, R(x)) =$ $b(x)\sqrt{f_i(x)}$ and $\eta_i(x, R(x)) = \sqrt{f_i(x)}$. Consider $T := \sum_{i=1}^k \theta_{\xi_i, \eta_i} \in K(Y)$. We shall show that $T = \phi(b)$. For any $\zeta \in C(\operatorname{graph} R)$, we have $(\phi(b)\zeta)(x,y) = b(x)\zeta(x,y)$ and

$$(T\zeta)(x,y) = \sum_{i} \xi_{i}(x,y) \sum_{x' \in R^{-1}(y)} e(x') \overline{\eta_{i}(x',y)} \zeta(x',y)$$
$$= \sum_{i} b(x) \sqrt{f_{i}(x)} \sum_{x' \in R^{-1}(y)} e(x') \sqrt{f_{i}(x')} \zeta(x',y).$$

In the case when b(x) = 0, we have

$$(T\zeta)(x,y) = 0 = (\phi(b)\zeta)(x,y).$$

In the case when $b(x) \neq 0$, we have $x \in \operatorname{supp}(b) = S \subset \bigcup_{i=1}^{m} U_{x_i}$. Hence e(x) = 1 and $x \in U_{x_i}$ for some *i*. Put y = R(x). Then for any $x' \in R^{-1}(y)$ with $x' \neq x$, $f_i(x') = 0$, because $x' \in U_{x_i}^c$. Therefore we have

$$(T\zeta)(x,y) = \sum_{\{i;x \in U_{x_i}\}} b(x)f_i(x)\zeta(x,y) = b(x)\zeta(x,y) = (\phi(b)\zeta)(x,y).$$

Thus $\phi(b) = T \in K(Y)$. Now for a general $b \in B$ which vanishes on the critical points C of R, there exists a sequence $(b_n)_n$ in B with compact supports $\operatorname{supp}(b_n) \subset \widehat{\mathbb{C}} \setminus C$ such that $\|b - b_n\|_{\infty} \to 0$. Hence $\phi(b) \in K(Y)$, i.e., $b \in I_Y$.

Conversely let $b \in B$ and $b(c) \neq 0$ for some critical point c. We may assume that b(c) = 1. Put $m = e(c) \geq 2$. We need to show that $\phi(b) \notin K(Y)$. On the contrary suppose that $\phi(b) \in K(Y)$. Then for $\varepsilon = \frac{1}{5\sqrt{d}}$, there exists a finite subset $\{\xi_i, \eta_i \in X; i = 1, \ldots, N\}$ such that $\|\phi(b) - \sum_{i=1}^N \theta_{\xi_i,\eta_i}\| < \varepsilon$. Choose a sequence $(x_n)_n$ in $\mathbb{C} \setminus C$ such that $x_n \to c$. Since x_n is not a critical point, there exists an open neighbourhood U_n of x_n such that $U_x \cap C = \phi$ and the restriction $R|_{U_n} :$ $U_n \to R(U_n)$ is a homeomorphism. There exists $\zeta_n \in Y$ with $\sup \zeta_n \subset$ $\{(x, R(x)); x \in U_n\}, \zeta_n(x_n, R(x_n)) = 1$ and $0 \leq \zeta_n \leq 1$. Then $\|\zeta_n\|_2 \leq \sqrt{d}$. For any $x' \in R^{-1}(R(x_n))$ with $x' \neq x_n, \zeta_n(x', R(x_n)) = 0$, Hence

$$\begin{aligned} |b(x_n) - \sum_{i=1}^{N} \xi_i(x_n, R(x_n)) \overline{\eta_i(x_n, R(x_n))}| \\ &= |b(x_n) - \sum_{i=1}^{N} \xi_i(x_n, R(x_n)) \sum_{x' \in R^{-1}(R(x_n))} e(x') \overline{\eta_i(x', R(x_n))} (\zeta_n)(x', R(x_n)) \\ &= |((\phi(b) - \sum_{i=1}^{N} \theta_{\xi_i, \eta_i}) \zeta_n)(x_n, R(x_n))| \\ &\leq \|(\phi(b) - \sum_{i=1}^{N} \theta_{\xi_i, \eta_i}) \zeta_n\|_2 \leq \|(\phi(b) - \sum_{i=1}^{N} \theta_{\xi_i, \eta_i}\| \|\zeta_n\|_2 \leq \varepsilon \sqrt{d}. \end{aligned}$$

Taking $n \to \infty$, we have

$$|b(c) - \sum_{i=1}^{N} \xi_i(c, R(c)) \overline{\eta_i(c, R(c))}| \le \varepsilon \sqrt{d}$$

On the other hand, consider $\zeta \in Y$ satisfying $\zeta(c, R(c)) = 1, 0 \le \zeta \le 1$ and $\zeta(x', R(x')) = 0$ for $x' \in R^{-1}(R(c))$ with $x' \ne c$ Then

$$\begin{aligned} |b(c) - \sum_{i=1}^{N} \xi_{i}(c, R(c))e(c)\overline{\eta_{i}(c, R(c))}| \\ &= |b(c) - \sum_{i=1}^{N} \xi_{i}(c, R(c)) \sum_{x' \in R^{-1}(R(c))} e(x')\overline{\eta_{i}(x', R(x'))}\zeta(x', R(x'))| \\ &\leq \|(\phi(b) - \sum_{i=1}^{N} \theta_{\xi_{i}, \eta_{i}})\zeta\|_{2} \leq \varepsilon \sqrt{d}. \end{aligned}$$

Since $e(c) \ge 2$ and b(c) = 1, we have

$$\frac{1}{2} \le |b(c) - \frac{1}{e(c)}b(c)| \le \varepsilon\sqrt{d} + \frac{1}{e(c)}\varepsilon\sqrt{d} \le 2\varepsilon\sqrt{d} = \frac{2}{5}$$

This is a contradiction. Therefore $\phi(b) \notin K(Y)$

Corollary 2.6. $^{\#}C = \dim(B/I_Y) \text{ and } ^{\#}(C \cap J_R) = \dim(A/I_X).$

Corollary 2.7. The Julia set contains no critical points if and only if $\phi(A)$ is contained in K(X) if and only if X is finitely generated projective right A module.

3. SIMPLICITY AND PURE INFINITENESS

Let R be a rational function with the Julia set $J_R \neq \phi$. Let $A = C(J_R)$ and $X = C(\operatorname{graph} R|_{J_R})$. Define an endomorphism $\alpha : A \to A$ by

$$(\alpha(a))(x) = a(R(x))$$

for $a \in A$, $x \in J_R$. We also define a unital completely positive map $E_R : A \to A$ by

$$(E_R(a))(y) := \frac{1}{\deg R} \sum_{x \in R^{-1}(y)} e(x)a(x)$$

for $a \in A$, $y \in J_R$. In fact, for a constant function $\xi_0 \in X$ with

$$\xi_0(x,y) := \frac{1}{\sqrt{\deg R}}$$

we have

$$E_R(a) = (\xi_0 | \phi(a) \xi_0)_A$$
 and $E_R(I) = (\xi_0 | \xi_0)_A = I.$

We introduce an operator $D := S_{\xi_0} \in \mathcal{O}_R$.

Lemma 3.1. In the above situation, for $a, b \in A$, we have the following:

(1) $E_R(\alpha(a)b) = aE(b)$ and in particular $E_R(\alpha(a)) = a$ (2) $D^*D = I$. (3) $\alpha(a)D = Da$. (4) $D^*aD = E_R(a)$. (5) $\phi(A)\xi_0 = X$.

Proof. (1): By the definition of E_R , we have

$$E_R(\alpha(a)b)(y) = \frac{1}{\deg R} \sum_{x \in R^{-1}(y)} e(x)a(R(x))b(x) = a(y)E(b)(y).$$

(2): $D^*D = S^*_{\xi_0}S_{\xi_0} = (\xi_0|\xi_0)_A = I.$ (3): We have $\alpha(a)D = S_{\phi(\alpha(a))\xi_0}$ and $Da = S_{\xi_0a}$. Since $(\phi(\alpha(a))\xi_0)(x, R(x)) = (\alpha(a))(x)\xi_0(x, R(x)) = a(R(x)) = (\xi_0a)(x, R(x)),$ we have $\alpha(a)D = Da.$ (4): $D^*aD = S^*_{\xi_0}aS_{\xi_0} = (\xi_0|\phi(a)\xi_0)_A = E_R(a).$ (5)For any $f \in X$, define $\tilde{f} \in A$ by $\tilde{f}(x) = \sqrt{\deg R}f(x, R(x)).$ Then $\phi(\tilde{f})\xi_0 = f.$ Thus $\phi(A)\xi_0 = X.$

R. Exel introduced an interesting construction of a crossed product by an endomorhism with a transfer operator in [8].

Proposition 3.2. C^* -algebra \mathcal{O}_R is isomorphic to a Exel's crossed product by an endomorhism with a transfer operator E_R .

Proof. By the universality, the Toeplitz algebras are isomorphic. Since $\phi(A)S_{\xi_0} = \{S_x; x \in X\}$ can be identified with X by (5), (a, k) is a redundancy if and only if $k = \phi(a)$. Therefore C*-algebra \mathcal{O}_R is isomorphic to his algebra.

Remark. If there exist no critical points in J_R , then the simplicity of \mathcal{O}_R is a consequence of the simplicity of his algebra, which is proved in Exel-Vershik [9]. But we need a further argument because of the existence of critical points in J_R .

Lemma 3.3. In the same situation, for $a \in A$ and $f_1, \ldots, f_n \in X$, we have the following:

$$\alpha^n(a)S_{f_1}\dots S_{f_n}=S_{f_1}\dots S_{f_n}a.$$

Proof. It is enough to show that $\alpha(a)S_f = S_f a$ for $f \in X$. We have $\alpha(a)S_f = S_{\phi(\alpha(a))f}$ and $S_f a = S_{fa}$. Since

$$(\phi(\alpha(a))f)(x, R(x)) = (\alpha(a))(x)f(x, R(x)) = a(R(x))f(x, R(x)) = (fa)(x, R(x)),$$

we have $\alpha(a)S_f = S_f a$.

Lemma 3.4. Let R be a rational function with deg $R \ge 2$. For any non-zero positive element $a \in A$ and for any $\varepsilon > 0$ there exist $n \in \mathbb{N}$ and $f \in X^{\otimes n}$ with $(f|f)_A = I$ such that

$$\|a\| - \varepsilon \le S_f^* a S_f \le \|a\|$$

Proof. Let x_0 be a point in J_R with $|a(x_0)| = ||a||$. For any $\varepsilon > 0$ there exist an open neighbourhood U of x_0 in J_R such that for any $x \in U$ we have $||a|| - \varepsilon \leq a(x) \leq ||a||$. Choose another open neighbourhood V of x_0 in J_R and a compact subset $K \subset J_R$ satisfying $V \subset K \subset U$. Since deg $R \geq 2$, there exists $n \in \mathbb{N}$ such that $R^n(V) = J_R$ by Beardon [2] Theorem 4.2.5. We identify $X^{\otimes n}$ with $C(\operatorname{graph} R^n|_{J_R})$ as in Lemma 3.3. Define closed subsets F_1 and F_2 of $J_R \times J_R$ by

$$F_{1} = \{(x, y) \in J_{R} \times J_{R} | y = R^{n}(x), x \in K\}$$

$$F_{2} = \{(x, y) \in J_{R} \times J_{R} | y = R^{n}(x), x \in U^{c}\}$$

Since $F_1 \cap F_2 = \phi$, there exists $g \in C(\operatorname{graph} R^n|_{J_R})$ such that $0 \leq g(x) \leq 1$ and

$$g(x,y) = \begin{cases} 1, & (x,y) \in F_1 \\ 0, & (x,y) \in F_2 \end{cases}$$

Since $R^n(V) = J_R$, for any $y \in J_R$ there exists $x_1 \in V$ such that $R^n(x_1) = y$, so that $(x_1, y) \in F_1$. Therefore

$$(g|g)_A(y) = \sum_{x \in (R^n)^{-1}(y)} e_{R^n}(x)|g(x,y)|^2$$

$$\ge e_{R^n}(x_1)|g(x_1,y)|^2 \ge 1$$

Let $b := (g|g)_A$. Then $b(y) = (g|g)_A(y) \ge 1$. Thus $b \in A$ is positive and invertible. We put $f := gb^{-1/2} \in X^{\otimes n}$. Then

$$(f|f)_A = (gb^{-1/2}|gb^{-1/2})_A = b^{-1/2}(g|g)_A b^{-1/2} = I.$$

For any $y \in J_R$ and $x \in (\mathbb{R}^n)^{-1}(y)$, if $x \in U$, then $||a|| - \varepsilon \leq a(x)$, and if $x \in U^c$, then $f(x, y) = g(x, y)b^{-1/2}(y) = 0$. Therefore

$$\begin{aligned} \|a\| - \varepsilon &= (\|a\| - \varepsilon)(f|f)_A(y) \\ &= (\|a\| - \varepsilon) \sum_{x \in (R^n)^{-1}(y)} e_{R^n}(x)|f(x,y)|^2 \\ &\leq \sum_{x \in (R^n)^{-1}(y)} e_{R^n}(x)a(x)|f(x,y)|^2 \\ &= (f|af)_A(y) = S_f^* a S_f(y). \end{aligned}$$

We also have that

$$S_f^* a S_f = (f | a f)_A \le ||a|| (f | f)_A = ||a||.$$

Lemma 3.5. Let R be a rational function with deg $R \geq 2$. For any non-zero positive element $a \in A$ and for any $\varepsilon > 0$ with $0 < \varepsilon < ||a||$, there exist $n \in \mathbb{N}$ and $u \in X^{\otimes n}$ such that

$$||u||_2 \le (||a|| - \varepsilon)^{-1/2}$$
 and $S_u^* a S_u = I$

Proof. For any $a \in A$ and $\varepsilon > 0$ as above, we choose $f \in (X)^{\otimes n}$ as in Lemma 3.4. Put $c = S_f^* a S_f$. Since $0 < ||a|| - \varepsilon \le c \le ||a||$, c is positive and invertible. Let $u := fc^{-1/2}$. Then

$$S_u^* a S_u = (u|au)_A = (fc^{-1/2}|afc^{-1/2})_A = c^{-1/2}(f|af)_A c^{-1/2} = I.$$

Since $||a|| - \varepsilon \le c$, we have $c^{-1/2} \le (||a|| - \varepsilon)^{-1/2}$. Hence $||u||_2 = ||fc^{-1/2}||_2 \le ||c^{-1/2}||_2 \le (||a|| - \varepsilon)^{-1/2}$

$$|u||_{2} = ||fc^{-1/2}||_{2} \le ||c^{-1/2}||_{2} \le (||a|| - \varepsilon)^{-1/2}.$$

Lemma 3.6. Let R be a rational function with deg R > 2. For any $m \in \mathbb{N}$, any $n \in \mathbb{N}$, any $T \in L(X^{\otimes n})$ and any $\varepsilon > 0$, there exists a positive element $a \in A$ such that

$$\|\phi(\alpha^n(a))T\|^2 \ge \|T\|^2 - \varepsilon$$

and $a\alpha^k(a) = 0$ for $k = 1, \dots, m$.

Proof. For any $m \in \mathbb{N}$, let $\mathcal{P}_m = \{x \in J_R \mid R^k(x) = x, \text{ for some } k = x\}$ $1, \ldots, m$. Since R is a rational function, \mathcal{P}_m is a finite set.

For any $n \in \mathbb{N}$, any $T \in L(X^{\otimes n})$ and any $\varepsilon > 0$, there exists $f \in X^{\otimes n}$ such that $||f||_2 = 1$ and $||T||^2 \ge ||Tf||_2^2 > ||T||^2 - \varepsilon$. We still identify $X^{\otimes n}$ with $C(\operatorname{graph} R^n|_{J_R})$. Then there exists $y_0 \in J_R$ such that

$$||Tf||_2^2 = \sum_{x \in (R^n)^{-1}(y_0)} e_{R^n}(x) |(Tf)(x, y_0)|^2 > ||T||^2 - \varepsilon.$$

Since $y \mapsto (Tf|Tf)_A(y)$ is continuous and

$$||Tf||_2^2 = \sup_{y \in J_R} \sum_{x \in (R^n)^{-1}(y)} e_{R^n}(x) |(Tf)(x,y)|^2,$$

there exists an open neighbourhood U_{y_0} of y_0 such that for any $y \in U_{y_0}$

$$\sum_{x \in (R^n)^{-1}(y)} e_{R^n}(x) |(Tf)(x,y)|^2 > ||T||^2 - \varepsilon.$$

Since R is a rational function with deg $R \ge 2$, the Julia set J_R is a perfect set and uncountable by Beardon [2] Theorem 4.2.4. Therefore $U_{y_0} \setminus \mathcal{P}_m \neq \phi$. Choose $y_1 \in U_{y_0} \setminus \mathcal{P}_m$. Since $R^k(y_1) \neq y_1$ $(k = 1, \dots, m)$, $y_1 \notin (R^k)^{-1}(y_1)$. For $k = 1, \ldots, m$, choose an open neighbourhood W_k of y_1 such that $W_k \subset U_{y_0}$ and $(R^k)^{-1}(W_k) \cap W_k = \phi$. Put $W = \phi$ $\bigcap_{k=1}^{m} W_k$. Choose $a \in A = C(J_R)$ such that $a(y_1) = 1, 0 \le a \le 1$ and supp $a \subset W$. Since $(R^k)^{-1}(W) \cap W = \phi$,

$$\operatorname{supp}(\alpha^k(a)) \cap \operatorname{supp}(a) = \phi \qquad (k = 1, \dots, m).$$

It implies that $\alpha^k(a)a = 0$ (k = 1, ..., m).

Since $y_1 \in U_{y_0}$,

$$\begin{aligned} |\phi(\alpha^{n}(a))Tf||_{2}^{2} &= \sup_{y \in J_{R}} \sum_{x \in (R^{n})^{-1}(y)} e_{R^{n}}(x) |(a(R^{n}(x))(Tf)(x,y)|^{2} \\ &\geq \sum_{x \in (R^{n})^{-1}(y_{1})} e_{R^{n}}(x) |a(y_{1})(Tf)(x,y_{1})|^{2} \\ &= \sum_{x \in (R^{n})^{-1}(y_{1})} e_{R^{n}}(x) |(Tf)(x,y_{1})|^{2} \\ &> ||T||^{2} - \varepsilon. \end{aligned}$$

Therefore we have $\|\phi(\alpha^n(a))T\|^2 \ge \|T\|^2 - \varepsilon$.

Let \mathcal{F}_n be the C^* -subalgebra of \mathcal{F}_X generated by $K(X^{\otimes k})$, $k = 0, 1, \ldots, n$ and B_n be the C^* -subalgebra of \mathcal{O}_X generated by

$$\bigcup_{k=1}^{n} \{ S_{x_1} \dots S_{x_k} S_{y_k}^* \dots S_{y_1}^* : x_1, \dots x_k, y_1, \dots y_k \in X \} \cup A.$$

In the following Lemma 3.7 we shall use an isomorphism $\varphi : \mathcal{F}_n \to B_n$ as in Pimsner [23] and Fowler-Muhly-Raeburn [11] such that

 $\varphi(\theta_{x_1\otimes\cdots\otimes x_k,y_1\otimes\cdots\otimes y_k})=S_{x_1}\ldots S_{x_k}S_{y_k}^*\ldots S_{y_1}^*.$

To simplify notation, we put $S_x = S_{x_1} \dots S_{x_k}$ for $x = x_1 \otimes \dots \otimes x_k \in X^{\otimes k}$

Lemma 3.7. Let R be a rational function with deg $R \ge 2$. Let $b = c^*c$ for some $c \in \mathcal{O}_X^{alg}$. We decompose $b = \sum_j b_j$ with $\gamma_t(b_j) = e^{ijt}b_j$. For any $\varepsilon > 0$ there exists $P \in A$ with $0 \le P \le I$ satisfying the following:

(1) $Pb_jP = 0$ $(j \neq 0)$ (2) $||Pb_0P|| \ge ||b_0|| - \varepsilon$

Proof. For $x \in X^{\otimes n}$, we define length(x) = n with the convention length(a) = 0 for $a \in A$. We write c as a finite sum $c = a + \sum_i S_{x_i} S_{y_i}^*$. Put $m = 2 \max\{ \text{length}(x_i), \text{length}(y_i); i \}$.

For j > 0, each b_j is a finite sum of terms in the form such that

$$S_x S_y^* \qquad x \in X^{\otimes (k+j)}, \qquad y \in X^{\otimes k} \qquad 0 \le k+j \le m$$

In the case when j < 0, b_j is a finite sum of terms in the form such that

$$S_x S_y^* \qquad x \in X^{\otimes k}, \qquad y \in X^{\otimes (k+|j|)} \qquad 0 \le k+|j| \le m$$

We shall identify b_0 with an element in $A_{m/2} \subset A_m \subset L(X^{\otimes m})$. Apply Lemma 3.6 for m = n and $T = (b_0)^{1/2}$. Then there exists a positive element $a \in A$ such $\|\phi(\alpha^m(a))T\|^2 \ge \|T\|^2 - \varepsilon$ and $a\alpha^j(a) = 0$ for $j = 1, \dots, m$. Define a positive operator $P = \alpha^m(a) \in A$. Then

$$||Pb_0P|| = ||Pb_0^{1/2}||^2 \ge ||b_0^{1/2}||^2 - \varepsilon = ||b_0|| - \varepsilon$$

For j > 0, we have

$$PS_x S_y^* P = \alpha^m(a) S_x S_y^* \alpha^m(a) = S_x \alpha^{m-(k+j)}(a) \alpha^{m-k}(a) S_y^*$$
$$= S_x \alpha^{m-(k+j)}(a\alpha^j(a)) S_y^* = 0$$

For j < 0, we also have that $PS_xS_y^*P = 0$. Hence $Pb_jP = 0$ for $j \neq 0$.

Theorem 3.8. Let R be a rational function with deg $R \ge 2$. Then the C^* -algebra \mathcal{O}_R associated with R is simple and purely infinite.

Proof. Let $w \in \mathcal{O}_R$ be any non-zero positive element. We shall show that there exist $z_1, z_2 \in \mathcal{O}_R$ such that $z_1^*wz_2 = I$. We may assume that ||w|| = 1. Let $E : \mathcal{O}_R \to \mathcal{O}_R^{\gamma}$ be the canonical conditional expectation onto the fixed point algebra by the gauge action γ . Since E is faithful, $E(w) \neq 0$. Choose ε such that

$$0 < \varepsilon < \frac{\|E(w)\|}{4}$$
 and $\varepsilon \|E(w) - 3\varepsilon\|^{-1} \le 1$.

There exists an element $c \in \mathcal{O}_X^{alg}$ such that $||w - c^*c|| < \varepsilon$ and $||c|| \leq 1$. Let $b = c^*c$. Then b is decomposed as a finite sum $b = \sum_j b_j$ with $\gamma_t(b_j) = e^{ijt}b_j$. Since $||b|| \leq 1$, $||b_0|| = ||E(b)|| \leq 1$. By Lemma 3.7, there exists $P \in A$ with $0 \leq P \leq I$ satisfying $Pb_jP = 0$ $(j \neq 0)$ and $||Pb_0P|| \geq ||b_0|| - \varepsilon$. Then we have

$$\|Pb_0P\| \ge \|b_0\| - \varepsilon = \|E(b)\| - \varepsilon$$
$$\ge \|E(w)\| - \|E(w) - E(b)\| - \varepsilon \ge \|E(w)\| - 2\varepsilon$$

For $T := Pb_0P \in L(X^{\otimes m})$, there exists $f \in X^{\otimes m}$ with ||f|| = 1 such that

$$||T^{1/2}f||_2^2 = ||(f|Tf)_A|| \ge ||T|| - \varepsilon$$

Hence we have $||T^{1/2}f||_2^2 \ge ||E(w)|| - 3\varepsilon$. Define $a = S_f^*TS_f = (f|Tf)_A \in A$. Then $||a|| \ge ||E(w)|| - 3\varepsilon > \varepsilon$. By Lemma 3.5, there exists $n \in \mathbb{N}$ and $u \in X^{\otimes n}$ such that

$$|u||_2 \le (||a|| - \varepsilon)^{-1/2}$$
 and $S_u^* a S_u = I$

Then $||u|| \leq (||E(w)|| - 3\varepsilon)^{-1/2}$. Moreover we have

$$\begin{split} \|S_{f}^{*}PwPS_{f} - a\| &= \|S_{f}^{*}PwPS_{f} - S_{f}^{*}TS_{f}\| \\ &= \|S_{f}^{*}PwPS_{f} - S_{f}^{*}Pb_{0}PS_{f}\| \\ &= \|S_{f}^{*}PwPS_{f} - S_{f}^{*}PbPS_{f}\| \\ &\leq \|S_{f}\|^{2}\|P\|^{2}\|w - b\| < \varepsilon \end{split}$$

Therefore

$$\begin{split} \|S_u^* S_f^* P w P S_f S_u - I\| &= \|S_u^* S_f^* P w P S_f S_u - S_u^* a S_u\| \\ &\leq \|u\|^2 \|S_f P w P S_f - a\| \\ &< \|u\|^2 \varepsilon \leq \varepsilon \|E(w) - 3\varepsilon\|^{-1} \leq 1. \end{split}$$

Hence $S_u^* S_f^* P w P S_f S_u$ is invertible. Thus there exists $v \in \mathcal{O}_X$ with $S_u^* S_f^* P w P S_f S_u v = I$. Put $z_1 = S_u^* S_f^* P$ and $z_2 = P S_f S_u v$. Then $z_1 w z_2 = I$.

Remark. J. Schweizer showed a nice criterion of the simplicity of Cuntz-Pimsner algebras in [26]: If a Hilbert bimodule X is minimal and non-periodic, then \mathcal{O}_X is simple. Any X-invariant ideal J of A corresponds to a closed subset K of J_R with $R^{-1}(K) \subset K$. If deg $R \geq 2$, then for any $z \in J_R$ the backward orbit of z is dense in J_R by [2] Theorem 4.2.7. Therefore any X-invariant ideal J of A is A or 0, that is, X is minimal. Since A is commutative and $L(X_A)$ is non-commutative, X is non-periodic. Thus Schweizer's theorem also implies that \mathcal{O}_R is simple. Our theorem gives simplicity and pure infiniteness with a direct proof.

Proposition 3.9. Let R be a rational function with deg $R \ge 2$. Then the C^* -algebra \mathcal{O}_R associated with R is separable and nuclear, and satisfies the Universal Coefficient Theorem.

Proof. Since \mathcal{J}_X and \mathcal{T}_X are KK-equivalent to abelian C^* -algebras I_X and A, the quotient $\mathcal{O}_X \cong \mathcal{T}_X / \mathcal{J}_X$ satisfies the UCT. \mathcal{O}_X is also shown to be nuclear as in an argument of [7].

Remark. If deg $R \geq 2$, then the isomorphisms class of \mathcal{O}_R is completely determined by the K-theory together with the class of the unit by the classification theorem by Kirchberg-Phillips [17], [22].

4. EXAMPLES

If rational functions R_1 and R_2 are topologically conjugate, then their C^* -algebras \mathcal{O}_{R_1} and \mathcal{O}_{R_2} are isomorphic. Therefore the K-groups $K_i(\mathcal{O}_{R_1})$ and $K_i(\mathcal{O}_{R_2})$ are isomorphic. Similarly $K_i(\mathcal{O}_{R_1}(\hat{\mathbb{C}}))$ and $K_i(\mathcal{O}_{R_2}(\hat{\mathbb{C}}))$ are isomorphic. Moreover if rational functions R_1 and R_2 are topologically conjugate, then the gauge actions are also conjugate. Therefore the K-groups of the fixed point algebras are also topologically conjugate invariant. We investigate what kind of information of complex dynamical systems is described by the K-theory.

We calculate the K-groups by the following six-term exact sequence due to Pimsner [23].

$$\begin{array}{cccc} K_0(I_X) & \xrightarrow{id-[X]} & K_0(A) & \xrightarrow{i_*} & K_0(\mathcal{O}_R) \\ & & & & & & \\ \delta_1 \uparrow & & & & & \\ K_1(\mathcal{O}_R) & \xleftarrow{i_*} & K_1(A) & \xleftarrow{id-[X]} & K_1(I_X) \end{array}$$

Example 4.1. Let $P(z) = z^n$ for $n \ge 2$. Then the Julia set J_P is the unit circle S^1 . The map $\phi : A \to L(X)$ can be identified with the *n*-times around embedding. Hence we have $K_0(\mathcal{O}_P) = \mathbb{Z} \oplus \mathbb{Z}/(n-1)\mathbb{Z}$ and $K_1(\mathcal{O}_P) = \mathbb{Z}$. The fixed point algebra \mathcal{O}_P^{γ} by the gauge action γ is a Bunce-Deddence algebra of type n^{∞} .

Example 4.2. Let $P(z) = z^2 - 2$. Then the Julia set J_P is the interval [-2, 2] and it contains a critical point 0. Since $I_X = \{a \in C([-2, 2]); a(0) = 0\}$, $K_0(I_X) = K_1(I_X) = 0$. Applying the above six term exact sequence with $K_0(A) = \mathbb{Z}$ and $K_1(A) = 0$, we have $K_0(\mathcal{O}_P) = \mathbb{Z}$ and $K_1(\mathcal{O}_P) = 0$. Since the identity I of \mathcal{O}_P represents the generator of $K_0(\mathcal{O}_P) = \mathbb{Z}$, the algebra \mathcal{O}_P is isomorphic to the Cuntz algebra \mathcal{O}_{∞} . (J_P, P) is topologically conjugate to a tent map ([0, 1], h) defined by

$$h(x) = \begin{cases} 2x, & 0 \le x \le \frac{1}{2}, \\ -2x+2, & \frac{1}{2} \le x \le 1. \end{cases}$$

Then $\frac{1}{2}$ is the only branched point of h. Deaconu and Muhly [5] associate a C^* -algebra $C^*([0,1],h)$ to the branched covering map h: $[0,1] \rightarrow [0,1]$. They exclude the branched point to construct their groupoid and the groupoid C^* -algebra $C^*([0,1],h)$ is not simple. Moreover $K_0(C^*([0,1],h)) = \mathbb{Z}^2$ and $K_1(C^*([0,1],h)) = 0$. Our C^* -algebra \mathcal{O}_P is simple and purely infinite and is not isomorphic to their C^* algebra $C^*([0,1],h)$. The K-groups are different.

Example 4.3. (quadratic polynomial) Let $P_c(z) = z^2 + c$. If c is not in the Mandelbrot set $\mathcal{M} := \{c \in \mathbb{C}; P_c^n(0) \text{ is bounded }\}$, then \mathcal{O}_{P_c} is isomorphic to the Cuntz algebra \mathcal{O}_2 . In fact (J_{P_c}, P_c) is topologically conjugate to the full two shift.

If c is in the interior of the main cardioid $C = \{\frac{z}{2} - \frac{z^2}{4} \in \mathbb{C}; |z| < 1\}$, then the Julia set is homeomorphic to the unit circle S^1 (see, for example, [10], page 211-212), and P_c is topologically conjugate to $h(z) = z^2$ on S^1 . Hence We have $K_0(\mathcal{O}_P) = \mathbb{Z}$ and $K_1(\mathcal{O}_P) = \mathbb{Z}$.

Example 4.4. Let R be a rational function of degree $d \ge 2$. Let z_0 be a (super)attracting fixed point of R. If all of the critical points of

R lie in the immediate attracting basin of z_0 , then \mathcal{O}_R is isomorphic to the Cuntz algebra \mathcal{O}_d . In fact (J_R, R) is topologically conjugate to the full d-shift by [2] Theorem 9.8.1. See section 4 in [24]. For example, If $R(z) = \frac{2z^2-1}{z}$, then $\mathcal{O}_R \cong \mathcal{O}_2$.

Example 4.5. Tchebychev polynomials T_n are defined by $\cos nz = T_n(\cos z)$. For example, $T_1(z) = z$, $T_2(z) = 2z^2 - 1$. Then the Julia set J_{T_n} is the interval [-1, 1] and J_{T_n} contains n - 1 critical points for $n \ge 2$. Since $K_0(A) = \mathbb{Z}$, $K_1(A) = 0$, $K_0(I_X) = 0$ and $K_1(I_X) = \mathbb{Z}^{n-2}$, we have $K_0(\mathcal{O}_{T_n}) = \mathbb{Z}^{n-1}$ and $K_1(\mathcal{O}_{T_n}) = 0$. Recall that, if the Julia set of a polynomial P of degree $n \ge 2$ is the interval [-1, 1], then $P = T_n$ or $P = -T_n$ ([2], page 11).

Example 4.6. We consider a rational function R by Lattes such that the Julia set is the entire Riemann sphere. Let $R(z) = \frac{(z^2+1)^2}{4z(z^2-1)}$. Then $J_R = \hat{\mathbb{C}}$ contains six critical points and $K_0(I_X) = \mathbb{Z}$ and $K_1(I_X) = \mathbb{Z}^5$. Therefore we have the exact sequence:

Deaconu and Muhly [5] already obtained the similar diagram for their C^* -algebra $C^*(\hat{\mathbb{C}}, R)$:

Since they exclude branched points, the rank of K_0 -groups of them are different with ours and their C^* -algebra is also different with ours.

Example 4.7. Ushiki [27] discovered a rational function whose Julia set is homeomorphic to the Sierpinski gasket. See also [15]. For example, let $R(z) = \frac{z^3 - \frac{16}{27}}{z}$. Then J_R is homeomorphic to the Sierpinski gasket K and J_R contains three critical points. Recall that the usual Sierpinski gasket K is constructed by three contractions $\gamma_1, \gamma_2, \gamma_3$ on the regular triangle T in \mathbb{R}^2 with three vertices $P = (1/2, \sqrt{3}/2)$, Q = (0,0) and R = (1,0) such that $\gamma_1(x,y) = (\frac{x}{2} + \frac{1}{4}, \frac{y}{2} + \frac{\sqrt{3}}{4})$, $\gamma_2(x,y) = (\frac{x}{2}, \frac{y}{2}), \gamma_3(x,y) = (\frac{x}{2} + \frac{1}{2}, \frac{y}{2})$. Then a self-similar set $K \subset T$ satisfying $K = \gamma_1(K) \cup \gamma_2(K) \cup \gamma_3(K)$ is called a Sierpinski gasket. But these three contractions are not inverse branches of a map, because $\gamma_1(Q) = \gamma_2(P)$. Therefore we need to modify the construction of contractions. Put $\tilde{\gamma}_1 = \gamma_1, \tilde{\gamma}_2 = \alpha_{-\frac{2\pi}{3}} \circ \gamma_2$, and $\tilde{\gamma}_3 = \alpha_{\frac{2\pi}{3}} \circ \gamma_3$, where α_{θ} is $\frac{18}{2}$.

a rotation by the angle θ . Then $\tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3$ are inverse branches of a map $h: K \to K$, which is conjugate to $R: J_R \to J_R$. Two correspondences $C = \bigcup_i \{(\gamma_i(z), z); z \in K\}$ and $\tilde{C} = \bigcup_i \{(\tilde{\gamma}_i(z), z); z \in K\}$ generate Hilbert bimodule Z and \tilde{Z} over C(K). Then C^* -algebra $\mathcal{O}_R \cong \mathcal{O}_{\tilde{Z}}$ and $K_0(\mathcal{O}_R)$ contains a torsion free element. But C^* -algebra \mathcal{O}_Z is isomorphic to the Cuntz algebra \mathcal{O}_3 . Therefore C^* -algebra \mathcal{O}_Z and \mathcal{O}_R are not isomorphic. See [14].

5. Fatou set and the corresponding ideal

Recall that the Fatou set F_R of a rational functon R is the maximal open subset of the Riemann sphere $\hat{\mathbb{C}}$ on which $(R^n)_n$ is equicontinuous, and the Julia set J_R of R is the complement of the Fatou set in $\hat{\mathbb{C}}$. We consider the corresponding decomposition for the C^* -algebra $\mathcal{O}_R(\hat{\mathbb{C}})$, which was first pointed out by Deaconu and Muhly [5] in the case of their construction. Let $B = C(\hat{\mathbb{C}})$ and consider the ideal $I = \{b \in B; b | J_R = 0\}$ of B, so $I \cong C_0(F_R)$ and $B/I \cong C(J_R)$. We consider a submodule and quotient module of a Hilbert bimodule $Y = C(\operatorname{graph} R)$ over B. The right Hilbert I-module $YI := \{fb \in Y; f \in Y, b \in I\}$ is also described as $Y_I := \{f \in Y; (f|g)_B \in I \text{ for all } g \in Y\}$. Since $(f|f)_B \in I$ means that

$$(f|f)_B(y) = \sum_{x \in R^{-1}(y)} e(x)|f(x,y)|^2 = 0$$

for all $y \in J_R$, we have

$$YI = \{ f \in Y; f(x, y) = 0 \text{ for all } (x, y) \in \operatorname{graph} R|_{J_R} \}.$$

because J_R is complete invariant.

Invariant ideals for bimodules are introduced in [12] by Pinzari and ours and developed by Fowler, Muhly and Raeburn [11] in general case. In our situation I is a Y-invariant ideal of B, i.e., $\phi(I)Y \subset YI$. In fact, the condition is equivalent to that $(f|\phi(a)g)_B \subset I$ for any $a \in I$ and $f, g \in Y$, and it is easily checked as

$$(f|\phi(a)g)_B(y) = \sum_{x \in R^{-1}(y)} e(x)\overline{f(x,y)}a(x)g(x,y) = 0$$

for $y \in J_R$, because $x \in R^{-1}(J_R) = J_R$ and so a(x) = 0 for $a \in I$.

Therefore Y/YI is naturally a Hilbert bimodule over $B/I \cong C(J_R)$. We can identify Y/YI with a bimodule $X = C(\operatorname{graph} R|_{J_R})$ over $A = C(J_R)$.

Theorem 5.1. Let R be a rational function, $B = C(\hat{\mathbb{C}})$, $A = C(J_R)$ and $I = \{b \in B; b | J_R = 0\}$. Then the ideal $\mathcal{I}(I)$ generated by Iin $\mathcal{O}_R(\hat{\mathbb{C}})$ is Morita equivalent to $\mathcal{O}_R(F_R)$ and the quotient algebra $\mathcal{O}_R(\hat{\mathbb{C}})/\mathcal{I}(I)$ is canonically isomorphic to $\mathcal{O}_R = \mathcal{O}_R(J_R)$. Proof. We apply a result by Fowler, Muhly and Raeburn [11] Corollary 3.3. But we have to be careful, because $\phi(B)$ is not included in K(Y). The only thing we know is that the quotient algebra $\mathcal{O}_R(\hat{\mathbb{C}})/\mathcal{I}(I)$ is canonically isomorphic to the relative Cuntz-Pimsner algebra $\mathcal{O}(q^I(I_Y), Y/YI)$. Here $q^I : B \to B/I \cong A$ is the canonical quotient map. By Proposition 2.5, $I_Y = \{b \in B; b \text{ vanishes on } C\}$. and $I_X = \{a \in A; a \text{ vanishes on } C \cap J_R\}$. Hence we can identify $q^I(I_Y)$ with I_X . Since we also identify a bimodule Y/YI over B/I with a bimodule $X = C(\operatorname{graph} R|_{J_R})$ over $A = C(J_R)$, we have that $\mathcal{O}(q^I(I_Y), Y/YI)$ is isomorphic to $\mathcal{O}_R(J_R)$. \Box

6. Lyubich measure and KMS state

Lyubich [20] constructed an invariant measure μ for a rational function R, called a Lyubich measure, whose support is the Julia set. It derives a lower bound for the Hausdorff dimension of the Julia set. Lyubich showed that μ is a unique measure of maximal entropy and the entropy value $h_{\mu}(R)$ is equal to the topological entropy h(R) of R, which is log(deg R). We show that the Lyubich measure gives a unique KMS state on the C^* -algebra \mathcal{O}_R for the gauge action at inverse temperature log(deg R) if the Julia set contains no critical points.

Let $d = \deg R$ and δ_x be the Dirac measure for x on the Riemann sphere. For any $y \in \hat{\mathbb{C}}$ and each $n \in \mathbb{N}$, we define a probability measure μ_n^y by

$$\mu_n^y = \sum_{\{x \in \widehat{\mathbb{C}}; R^n(x) = y\}} d^{-n} e(x) \delta_x \ .$$

Lyubich showed that the sequence $(\mu_n^y)_n$ converges weakly to a measure μ . The Lyubich measure μ is independent of the choice of y, the support of μ is the Julia set J_R and $\mu(E) = \mu(R^{-1}(E))$ for any Borel set E. Hence $\int a(R(x))d\mu(x) = \int a(x)d\mu(x)$.

Theorem 6.1. Let R be a rational function with deg $R \ge 2$. Consider the C^* -algebra \mathcal{O}_R associated with R and the gauge action $\gamma : \mathbb{R} \to$ Aut \mathcal{O}_R such that $\gamma_t(S_f) = e^{it}S_f$ for $f \in X$. Suppose that the Julia set contains no critical points, (in particular, suppose that R is hyperbolic). Then (\mathcal{O}_R, γ) has a KMS state at the inverse temperature β if and only if $\beta = \log(\deg R)$ and the corresponding KMS state φ is also unique. Moreover the restriction of the state φ to $A = C(J_R)$ is given by the Lyubich measure μ such that $\varphi(a) = \int ad\mu$ for $a \in A$.

Proof. Since the Julia set contains no critical points, A-module X has a finite right basis $\{u_1, \ldots, u_n\}$. Let φ be a state on \mathcal{O}_R and φ_0 the restriction of φ to A. As in [24] section 3, φ is a KMS state at the inverse temperature β if and only if φ_0 satisfies that

(*)
$$\varphi_0(\sum_i (u_i | a u_i)_A) = e^\beta \varphi_0(a)$$
 for all $a \in A$.
20

Moreover any state φ_0 on A satisfying (*) extends uniquely to a KMS state φ at the inverse temperature β . In fact, define a state φ_n on the C^* -algebra B_n generated by

$$\{S_{x_1} \dots S_{x_n} S_{y_n}^* \dots S_{y_1}^*; x_1, \dots x_n, y_1, \dots y_n \in X\},\$$

by the induction:

$$\varphi_{n+1}(T) = e^{-\beta}\varphi_n(\sum_i S_{u_i}^* T S_{u_i})$$

for $T \in B_n$. The sequence $(\varphi_n)_n$ gives a state φ_∞ on the fixed point algebra \mathcal{O}_R^{γ} . Let $E : \mathcal{O}_R \to \mathcal{O}_R^{\gamma}$ be the canonical conditional expectation. Then the KMS state φ is defined by $\varphi = \varphi_\infty \circ E$.

Define a complete positive map $h: A \to A$ by $h(a) = \sum_{i} (u_i | a u_i)_A$ for $a \in A$. Then we have

$$h(a)(y) = \sum_{x \in R^{-1}(y)} a(x)$$

for $y \in J_R$. In fact, since $\{u_1, \ldots, u_n\}$ is a finite basis for X, $\sum_i u_i(u_i|f)_A = f$ for any $f \in X$. Therefore for y = R(x),

$$\sum_{i} u_i(x,y) \sum_{x' \in R^{-1}(y)} \overline{u_i(x',y)} f(x',y) = f(x,y).$$

This implies that $\sum_{i} u_i(x, y) \overline{u_i(x', y)} = \delta_{x,x'}$. Hence we have

$$\sum_{i} u_i(x,y) \overline{u_i(x,y)} = 1$$

Therefore

$$h(a)(y) = \sum_{i} (u_i | au_i)_A(y)$$

= $\sum_{i} \sum_{x \in R^{-1}(y)} \overline{u_i(x, y)} a(x) u_i(x, y) = \sum_{x \in R^{-1}(y)} a(x).$

Hence the condition (*) is written as

$$\varphi_0(h(a)) = e^\beta \varphi_0(a) \text{ for all } a \in A.$$

Putting a = 1, we have that deg $R = e^{\beta}$, that is, β need to be $\log(\deg R)$. For any $a \in A$, $(e^{-\beta}h)^n(a)$ converges uniformly to a constant $\int ad\mu$ by [20]. Thus $\int e^{-\beta}h(a)d\mu = \int ad\mu$. Hence Lyubich measure μ gives a KMS state at the inverse temperature log deg R. Take another KMS state τ . Then $\tau((e^{-\beta}h)^n(a) = \tau(a)$ converges to $\int ad\mu$. Thus $\tau(a) = \int ad\mu$. It shows that KMS state is unique.

References

- C. Anatharaman-Delaroche, Purely infinite C*-algebras arising from dynamical systems, Bull. Soc. Math. France 125(1997), 199-225
- [2] A. F. Beardon, Iteration of rational functions GTM 132, 1991, Springer New York
- [3] J. Cuntz, Simple C^{*}-algebras generated by isometries, Comm. Math. Phys. 57 (1977), 173-185.
- [4] V. Deaconu, generalized solenoids and C*-algebras, Pacific J. Math. 90 (1999), 247-260.
- [5] V. Deaconu and M. Muhly, C^{*}-algebras associated with branched coverings, Proc. AMS. 129 (2001), 1077-1086
- [6] Doplicher, C. Pinzari and R. Zuccante, The C*-algebra of a Hilbert bimodule, Boll. Unione Mat. Ital. 1 (1998), 263-281.
- [7] K. Dykema and D. Shilyakhtenko, Exactness of Cuntz-Pimsner C*-algebras, Proc. Edinburg Math. Soc. 44 (2001), 425-444.
- [8] R. Exel, A new look at the crossed-product of a C*-algebra by an endomorphism, to appear in Ergod. Th. & Dynam. Sys.
- [9] R. Exel and A. Vershik, C^* -algebras of Irreducible dynamical systems preprint
- [10] K. J. Falconer, Fractal Geometry, Wiley, 1997.
- [11] N. J. Fowler, P.S. Muhly and I. Raeburn, Representations of Cuntz-Pimsner Algebras, Indiana Univ. Math. J. 52 (2003), 569-605.
- [12] T. Kajiwara, C. Pinzari and Y. Watatani, Ideal structure and simplicity of the C^{*}-algebras generated by Hilbert bimodules J. Funct. Anal. 159 (1998), 295-322
- [13] T. Kajiwara, C. Pinzari and Y. Watatani, Jones index theory for Hilbert C^{*}-bimodules and its equivalence with conjugation theory, [arXiv: math. OA/0203185].
- [14] T. Kajiwara and Y. Watatani, C^* -algebras generated by self-similar sets, in preparation.
- [15] A. Kameyama, Julia sets of postcritically finite rational maps and topological self-similar sets, Nonlinearity 13, (2000), 165-188.
- [16] T. Katsura, A class of C^{*}-algebras generalizing both graph algebras and homeomorphism C^{*}-algebras, I, fundamental results, [arXiv: math. OA/0207252]
- [17] E. Kirchberg, The classification of purely infinite C^* -algebras using Kasparov's theory, preprint.
- [18] A. Kumjian, On certain Cuntz-Pimsner algebras, [arXive:math.OA/0108194].
- [19] M. Laca and J. Spielberg, Purely infinite C*-algebras from boundary actions of discrete groups, J. reine angew. Math. 480 (1996), 125-139.
- [20] M. Yu. Lyubich, Entropy properties of rational endomorphisms of the Riemann sphere, Ergodic Th. & Dynam. Sys. 3 (1983), 351-385.
- [21] P. S. Muhly and B. Solel, On the simplicity of some Cuntz-Pimsner algebras, Math. Scand. 83 (1998) 53-73.
- [22] N. C. Phillips, A classification theorem for nuclear purely infinite simple C^{*}algebras, Documenta Math. (2000), 5, 49-114.
- [23] M. Pimsner, A class of C^{*}-algebras generating both Cuntz-Krieger algebras and crossed product by Z, Free probability theory, AMS, (1997), 189-212.
- [24] C. Pinzari, Y. Watatani and K. Yonetani, KMS states, entropy and the variational principle in full C^{*}-dynamical systems, Comm. Math. Phys. 213 (2000), 331-379.
- [25] J. Renault, A Groupoid Approach to C*-algebras, Lecture Note in Mathematics, vol 793, Springer, 1980.

- [26] J. Schweizer, Dilations of C*-correspondences and the simplicity of Cuntz-Pimsner algebras, J. Funct Anal. 180 (2001), 404-425
- [27] S. Ushiki, Julia sets with polynomial symmetries, Proc, Int. Conf. on Dynamical systems and Related Topics, World Scientific, 1991.

(Tsuyoshi Kajiwara) Department of Environmental and Mathematical Sciences, Okayama University, Tsushima, 700-8530, Japan

(Yasuo Watatani) Department of Mathematical Sciences, Kyushu University, Hakozaki, Fukuoka, 812-8581, Japan