# Quantization of semi-classical twists and noncommutative geometry

Samsonov M.E.\*

Theoretical department St. Petersburg University Institute of Physics

#### Abstract

A problem of defining the quantum analogues for semi-classical twists in  $U(\mathfrak{g})[[t]]$  is considered. First, we study specialization at q=1 of singular coboundary twists defined in  $U_q(\mathfrak{g})[[t]]$  for  $\mathfrak{g}$  being a nonexceptional Lie algebra, then we consider specialization of noncoboundary twists when  $\mathfrak{g}=\mathfrak{sl}_3$  and obtain q-deformation of the semi-classical twist introduced by Connes and Moscovici in noncommutative geometry.

Keywords: Noncommutative geometry, Hopf algebras

### 1 Introduction

Hopf algebras play an increasingly important role in noncomutative geometry and Quantum Field Theory. One of the sources for producing new types of Hopf algebras is twisting, a deformation of the coalgebraic structure of a given Hopf algebra  $(H, \mu, \eta, \Delta, \epsilon, S)$  preserving the algebraic structure  $(H, \mu, \eta)$ . Such deformations are generated by the twisting elements (twists)  $\mathcal{F} \in (H \otimes H)$  satisfying the conditions

$$\mathcal{F}^{12}(\Delta \otimes \mathrm{id})(\mathcal{F}) = \mathcal{F}^{23}(\mathrm{id} \otimes \Delta)(\mathcal{F})$$

$$(\varepsilon \otimes \mathrm{id})(\mathcal{F}) = (\mathrm{id} \otimes \varepsilon)(\mathcal{F}) = 1$$
(1)

that guarantee  $H^{\mathcal{F}} \equiv (H, \mu, \eta, \operatorname{Ad}\mathcal{F} \circ \Delta, \epsilon, S)$  is a new Hopf algebra. In fact, when H is not finite dimensional,  $\mathcal{F}$  is usually defined in some completion of the tensor product and H is understood to be a topological Hopf algebra.

In this article we consider two types of twists: the semi-classical ones if  $H = U(\mathfrak{g})[[t]]$  and the quantum ones if  $H = U_q(\mathfrak{g})[[t]]$ . Some of the semi-classical deformations such as those defined by the Jordanian twists [5, 10] appear as the limiting cases of the quantum ones (in the sense that specialization

 $<sup>^*</sup>$ E-mail address: samsonov@pink.phys.spbu.ru

at q=1 is extended to work for topological Hopf algebras). It was a motivation for us to study the quantum twists as many computational problems involved into a direct check of (1) drastically resolve when one works with  $U_q(\mathfrak{g})[[t]]$  instead of  $U(\mathfrak{g})[[t]]$  and thus the quantum twists is a source for many universal deformation formulas in the sense of [5].

The work is organized as follows. After preliminary section intended to fix notations, we show that if  $\mathfrak{g}$  is a nonexceptional simple Lie algebra, then a quantum analogue of the Jordanian twist can be taken to be a coboundary twist in  $U_q(\mathfrak{g})[[t]]$ :

$$\mathcal{J}(e_{\lambda}) := (W \otimes W)\Delta(W^{-1}), \text{ where } W = \exp_{q_{\lambda}}(\frac{t}{1 - q_{\lambda}} e_{\lambda}); \qquad q_{\lambda} := q^{(\lambda, \lambda)}$$

with  $e_{\lambda}$  being a quantum highest root generator in some quantum Cartan-Weyl basis. We prove that  $\mathcal{J}(e_{\lambda})$  is nonsingular and specializes to a nontrivial twisting of  $U(\mathfrak{g})[[t]]$ . As an application of the Jordanian twists [5, 10] to noncommutative geometry [1], we prove that there is a homomorphism of the Connes-Moscovici Hopf algebra  $\iota: \mathcal{H}_1 \to U^{\mathcal{F}}(\mathfrak{sl}_3)[[t]]$ , with  $\mathcal{F}$  being a Jordanian twist, where  $\mathcal{H}_1$  has the following structure

$$[Y, X] = X, [Y, \delta_n] = n\delta_n, [X, \delta_n] = \delta_{n+1}, [\delta_k, \delta_l] = 0, k, l \ge 1$$
  

$$\Delta(Y) = Y \otimes 1 + 1 \otimes Y, \Delta(\delta_1) = \delta_1 \otimes 1 + 1 \otimes \delta_1$$
  

$$\Delta(X) = X \otimes 1 + 1 \otimes X + \delta_1 \otimes Y.$$
(2)

Through factoring  $\mathcal{H}'_1 := \mathcal{H}_1/<\delta_2-\frac{1}{2} \ \delta_1^2>$ , one obtains in fact an embedding

$$\iota: \mathcal{H}'_1 \hookrightarrow U^{\mathcal{F}}(\mathfrak{sl}_3)[[t]]$$

and the twist found in [1]:

$$F = \sum_{n>0} t^n \sum_{k=0}^n \frac{S(X)^k}{k!} (2Y + k)_{n-k} \otimes \frac{X^{n-k}}{(n-k)!} (2Y + n - k)_k$$
 (3)

where  $(\alpha)_k = \alpha(\alpha+1)\cdots(\alpha+k-1)$  and  $S(X) = -X + \delta_1 Y$ , can be obtained as a pullback  $F = \iota_*\Phi$  of a semi-classical twist  $\Phi$  in  $U^{\mathcal{F}}(\mathfrak{sl}_3)[[t]]$ . In section ?? we show that  $\iota$  can be "quantized", thus leading to a quantum analogue of  $\mathcal{H}'_1$ :

$$kxk^{-1} = q^2 x, \qquad kzk^{-1} = q^2 z, \qquad q^2xz - zx = -tz^2$$
 
$$\Delta(k) = k \otimes k, \qquad \Delta(z) = z \otimes k + 1 \otimes z$$
 
$$\Delta(x) = x \otimes k^{-1} + 1 \otimes x + t z \otimes \frac{(k - k^{-1})}{1 - q^2}.$$

### Acknowledgment

I would like to thank V. Lyakhovsky, A. Stolin and V. Tolstoy for valuable comments on the subject

### 2 Preliminarities

Let  $\mathfrak{g}$  be a simple Lie algebra with the set of simple roots  $\pi = \{\alpha_1, \ldots, \alpha_N\}$  and the Cartan matrix  $(A)_{ij} = a_{ij} = 2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i)$ . By definition, a Hopf algebra  $U_q(\mathfrak{g})$  is generated by  $\{e_i, f_i, k_i^{\pm 1}\}_{1 \leq i \leq N}$  over  $\mathbb{C}(q)$  which are subject to the following relations

$$k_i e_j k_i^{-1} = q^{(\alpha_i, \alpha_j)} e_j, \quad k_i f_j k_i^{-1} = q^{-(\alpha_i, \alpha_j)} f_j$$
 (4)

$$e_i f_j - f_j e_i = \delta_{ij} \frac{k_i - k_i^{-1}}{q - q^{-1}}$$
 (5)

$$\sum_{n=0}^{1-a_{ij}} (-1)^n \begin{bmatrix} 1-a_{ij} \\ n \end{bmatrix}_{q_i} (e_i)^n e_j(e_i)^{1-a_{ij}-n} = 0 \text{ for } i \neq j$$
 (6)

$$\sum_{n=0}^{1-a_{ij}} (-1)^n \begin{bmatrix} 1-a_{ij} \\ n \end{bmatrix}_{q_i} (f_i)^n f_j(f_i)^{1-a_{ij}-n} = 0 \text{ for } i \neq j$$
 (7)

where  $q_i = q^{\frac{(\alpha_i, \alpha_i)}{2}}$  and

$$\begin{bmatrix} m \\ n \end{bmatrix}_q \equiv \frac{[m]_q!}{[n]_q![m-n]_q!}$$

$$[k]_q! \equiv [1]_q[2]_q \dots [k]_q, \quad [l]_q \equiv (q^l - q^{-l})/(q - q^{-1})$$

The Hopf algebra structure is defined uniquely by fixing the values of the coproduct on the Chevalley generators

$$\Delta(k_i) = k_i \otimes k_i \tag{8}$$

$$\Delta(e_i) = k_i^{-1} \otimes e_i + e_i \otimes 1, \quad \Delta(f_i) = f_i \otimes k_i + 1 \otimes f_i$$
(9)

$$S(k_i) = k_i^{-1}, \quad S(e_i) = -k_i e_i, \quad S(f_i) = -f_i k_i^{-1}$$
 (10)

$$\varepsilon(k_i) = 1, \qquad \varepsilon(e_i) = 0, \qquad \varepsilon(f_i) = 0.$$
 (11)

Letting  $q_h = e^h$  and  $K_i := q_h^{h_i}$  in (4)-(11), we come to definition of  $U_h(\mathfrak{g})[[h]]$ , the topological Hopf algebra over  $\mathbb{C}[[h]]$ .

One introduces a linear ordering on the set of positive roots  $\Delta_+$  by fixing the reduced decomposition of the longest element in the Weyl group  $w_0 = s_{i_1} s_{i_2} \cdots s_{i_M}$ . Then the linear ordering read from the left to the right is the following

$$\Delta_{+} = \{\alpha_{i_1}, s_{i_1}\alpha_{i_2}, \dots, s_{i_1}s_{i_2} \cdots s_{i_{M-1}}\alpha_{i_M}\}. \tag{12}$$

An ordering (12) is normal, namely for each  $\alpha, \beta \in \Delta_+$  such that  $\alpha + \beta \in \Delta_+$  and  $\alpha \prec \beta$ , we have  $\alpha \prec \alpha + \beta \prec \beta$ . There is one-to-one correspondence between the reduced decompositions of the longest element in the Weyl group and the normal orderings given by (12). Following [8], one defines the generators corresponding to the composite roots. For a chosen normal ordering on  $\Delta_+$  let  $\alpha, \beta, \gamma \in \Delta_+$ 

be pairwise noncollinear roots, such that  $\gamma = \alpha + \beta$ . Let  $\alpha$  and  $\beta$  are taken so that there are no other roots  $\alpha'$  and  $\beta'$  with the property  $\gamma = \alpha' + \beta'$ . Then if  $e_{\pm\alpha}$  and  $e_{\pm\beta}$  have already been constructed, we set

$$e_{\gamma} = e_{\alpha}e_{\beta} - q^{-(\alpha,\beta)}e_{\beta}e_{\alpha}, \quad e_{-\gamma} = e_{-\beta}e_{-\alpha} - q^{-(\beta,\alpha)}e_{-\alpha}e_{-\beta}.$$

For any root  $\gamma \in \Delta_+$  define

$$\check{R}_{\gamma} := \exp_{q_{\gamma}}(-(q - q^{-1})a_{\gamma}^{-1} e_{\gamma}k_{\gamma} \otimes k_{\gamma}^{-1}f_{\gamma}),$$

where  $q_{\gamma} = q^{(\gamma,\gamma)}$  and

$$\exp_q(x) := \sum_{n \ge 0} \frac{x^n}{(n)_q!}, \quad (n)_q! \equiv (1)_q(2)_q \dots (n)_q, \quad (k)_q \equiv (1 - q^k)/(1 - q).$$

with factors  $a_{\gamma}$  coming from the relations

$$[e_{\gamma}, e_{-\gamma}] = a_{\gamma} \frac{k_{\gamma} - k_{\gamma}^{-1}}{q - q^{-1}}.$$

The elements  $\check{R}_{\gamma}$  are understood to be taken in some completion of  $U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$  (the Taylor extension or the h-adic [8]). Now the coproducts of composite root generators can be expressed in terms of the adjoint action of the following factors

$$\check{R}_{\prec\beta} := \prod_{\gamma \prec \beta} \check{R}_{\gamma},$$

where the product over all the positive roots such that  $\prec \beta$  is taken in accordance with the chosen normal ordering. Namely, we have the following

**Theorem 1** ([8, 9]). Consider the canonical isomorphism  $\mathfrak{h} \simeq \mathfrak{h}^*$  defined by the bilinear form  $(\ ,\ )$  on  $\mathfrak{h}$ . Let  $h_{\beta} \in \mathfrak{h}$  be the image of a root  $\beta \in \mathfrak{h}^*$  with respect to this isomorphism. Then the following identity holds:

$$\Delta(e_{\beta}) = \check{R}_{\prec\beta}(k_{\beta}^{-1} \otimes e_{\beta} + e_{\beta} \otimes 1)\check{R}_{\prec\beta}^{-1}.$$
 (13)

*Proof.* Note that (13) is equivalent to Proposition 8.3 from [8] if one applies  $(S \otimes S)$  to both parts of

$$\Delta^{op}(e_{\alpha}) = \left(\prod_{\gamma < \alpha} \check{R}_{\gamma}\right) (1 \otimes e_{\alpha} + e_{\alpha} \otimes \overline{k}_{\alpha}) \left(\prod_{\gamma < \alpha} \check{R}_{\gamma}\right)^{-1}$$

(notations are of [8]) and check that our construction of the modified basis differs by the change  $q \leftrightarrow q^{-1}$ .

We conclude this section by adding remarks on specialization [3]. Let  $\mathcal{A} = \mathbb{C}[q,q^{-1}]_{(q-1)}$  be a ring of rational functions regular at q=1 and let  $\hat{U}_q(\mathfrak{g})$  be an  $\mathcal{A}$  subalgebra in  $U_q(\mathfrak{g})$  generated by  $\left\{e_i,f_i,k_i^{-1},\frac{k_i-1}{q-1}\right\}$  then

$$\hat{U}_q(\mathfrak{g}) \otimes_{\mathcal{A}} \mathbb{C} \approx U(\mathfrak{g})$$

where  $\mathbb{C}$  is regarded as an  $\mathcal{A}$  module (q acts as 1). By construction of the quantum Cartan-Weyl basis we have the property

$$\Delta(e_{\beta}) \in \hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})$$

but, in fact, one can deduce from (13) more restrictive property

$$\Delta(e_{\beta}) - k_{\beta}^{-1} \otimes e_{\beta} - e_{\beta} \otimes 1 \in (q - q^{-1}) \ \hat{U}_{q}^{+}(\mathfrak{g}) \otimes \hat{U}_{q}^{+}(\mathfrak{g})$$

$$\tag{14}$$

where  $\hat{U}_q^+(\mathfrak{g})$  is generated by  $\left\{e_i, k_i^{-1}, \frac{k_i - 1}{q - 1}\right\}$ . In what follows we are usually working rather with completions  $\hat{U}_q(\mathfrak{g})[[t]]$  and  $\hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})[[t]]$  in which the twists are to be defined. Let us formulate the following simple result which is of value for further application

**Proposition 1.** Let  $\mathcal{F} \in \hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})[[t]]$  be a twist in  $U_q(\mathfrak{g})[[t]]$ , then its specialization  $\overline{\mathcal{F}}$ , obtained by order-wise specialization of its coefficients from  $\hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})$  at q = 1, is still a twist in  $U(\mathfrak{g})[[t]]$ .

*Proof.* Indeed, representing  $\mathcal{F}$  as a series

$$\mathcal{F} = 1 \otimes 1 + \mathcal{F}_1 t + \mathcal{F}_2 t^2 + \cdots$$

we see that (1) is a equivalent to an infinite set of identities and each of them after specializing q = 1 remain valid. Thus

$$\overline{\mathcal{F}} = 1 \otimes 1 + \overline{\mathcal{F}}_1 t + \overline{\mathcal{F}}_2 t^2 + \cdots$$

П

is a twist in  $U(\mathfrak{g})[[t]]$ .

## 3 Quantum Jordanian twists

We restrict ourselves to consideration of nonexceptional Lie algebra  $\mathfrak{g}$  and define the quantum Jordanian twists as those specializing to semi-classical ones which define quantization of skew-symmetric extended Jordanian r-matrices:

$$r_{\lambda} = H_{\lambda} \wedge E_{\lambda} + 2 \sum_{\gamma_{1} \prec \gamma_{2}, \gamma_{1} + \gamma_{2} = \lambda} E_{\gamma_{1}} \wedge E_{\gamma_{2}}$$

by the rule

$$\mathcal{R} = \mathcal{F}_{\lambda 21} \mathcal{F}_{\lambda}^{-1} = 1 \otimes 1 + t \ r_{\lambda} \bmod t^2,$$

where we have denoted by  $H_{\lambda}, E_{\gamma}$  the elements of the classical Cartan-Weyl basis.

Let us fix some normal ordering on  $\Delta_+$  and define a generator  $e_{\lambda} \in U_q(\mathfrak{g})$  corresponding to the highest root  $\lambda$  according to the recipe from the previous section. Then nonexceptional root systems are remarkable by the following property:

Proposition 2. Let g be a non exceptional Lie algebra, then there is such a normal ordering " $\prec$ " on  $\Delta_+$  so that

$$[e_{\gamma}, e_{\lambda}]_{q^{-(\gamma,\lambda)}} = 0 \text{ for any } \gamma \prec \lambda, \text{ and } e_{\gamma}, e_{\lambda} \in U_q(\mathfrak{g}).$$

*Proof.* The proof is based on the expansion [8, 9]:

$$e_{\gamma}e_{\lambda} - q^{-(\gamma,\lambda)}e_{\lambda}e_{\gamma} = \sum_{\gamma \prec \gamma_1 \prec \cdots \prec \gamma_j \prec \lambda} c_{l,\gamma,\lambda}e_{\gamma_1}^{l_1} \cdots e_{\gamma_j}^{l_j}$$
 (15)

where  $c_{l,\gamma,\lambda} \in \mathbb{C}[q,q^{-1}]$ . Non zero terms in the sum are subject to condition

$$\gamma + \lambda = l_1 \gamma_1 + l_2 \gamma_2 + \dots + l_j \gamma_j. \tag{16}$$

In  $A_N$  we choose a normal ordering as

roots without 
$$\alpha_1$$
  
 $\alpha_1 \succ \alpha_1 + \alpha_2 \succ \cdots \succ \alpha_1 + \alpha_2 + \cdots + \alpha_N \succ \alpha_2 \succ \cdots \succ \alpha_{N-1} + \alpha_N \succ \alpha_N$ 

and  $\lambda = \alpha_1 + \cdots + \alpha_N$ . If  $\lambda \succ \gamma \succ \alpha_N$  we can satisfy (15) only with zero coefficients.

In  $B_N$  we have

and 
$$\lambda = \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{N-1} + 2\beta$$
.

roots without  $\alpha_1$ 

$$\alpha_1 \succ \alpha_1 + \alpha_2 \succ \dots \succ \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{N-1} + 2\beta \succ \alpha_2 \succ \alpha_2 + \alpha_3 \succ \dots \succ \beta$$
and  $\lambda = \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{N-1} + 2\beta$ .

In  $C_N$  we fix the following ordering

roots without 
$$\beta$$

roots without 
$$\beta$$

$$\overbrace{\alpha_1 \prec \alpha_1 + \alpha_2 \prec \cdots \prec \alpha_1 + \cdots + \alpha_{N-1}}^{\text{roots without } \beta} \prec 2(\alpha_1 + \cdots + \alpha_{N-1}) + \beta \prec \alpha_1 + \cdots + \alpha_{N-1} + \beta \prec \cdots \prec \alpha_1 + 2(\alpha_2 + \cdots + \alpha_{N-1}) + \beta \prec \cdots \prec \alpha_2 \prec \cdots \prec \beta,$$

and  $\lambda = 2(\alpha_1 + \cdots + \alpha_{N-1}) + \beta$ . This ordering eliminates all non zero terms on the r.h.s of (15).

In  $D_N$  we have quite a similar situation

$$\alpha_1 \succ \alpha_1 + \alpha_2 \succ \cdots \succ \alpha_1 + \cdots + \alpha_{N-1} \succ \alpha_1 + \cdots + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + 2\alpha_{N-2} + \alpha_{N-1} + \beta \succ \cdots \succ \alpha_1 + 2\alpha_2 + \cdots + 2\alpha_{N-2} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \beta \succ \alpha_1 + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_{N-1} + \alpha_2 + \cdots + \alpha_{N-1} + \alpha_2 + \alpha_2$$

$$\underbrace{\alpha_2 \succ \alpha_2 + \alpha_3 \succ \cdots \succ \beta}_{\text{roots without } \alpha_1},$$

and 
$$\lambda = \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{N-2} + \alpha_{N-1} + \beta$$
.

As a direct consequence of Proposition 2, we obtain

**Proposition 3.** A q-commutation holds

$$(e_{\lambda} \otimes 1)(\Delta(e_{\lambda}) - e_{\lambda} \otimes 1) = q_{\lambda}(\Delta(e_{\lambda}) - e_{\lambda} \otimes 1)(e_{\lambda} \otimes 1)$$
 where  $q_{\lambda} = q^{(\lambda,\lambda)}$ .

*Proof.* The statement follows from Proposition 2 and (13) if one notices that

$$\check{R}_{\prec\lambda}(e_{\lambda}\otimes 1)=(e_{\lambda}\otimes 1)\check{R}_{\prec\lambda}.$$

Let us define a coboundary twist

$$\mathcal{J}(e_{\lambda}) = (W \otimes W)\Delta(W^{-1}), \text{ where } W = \exp_{q_{\lambda}}(\frac{t}{1-q_{\lambda}} e_{\lambda})$$

Then the following is true

**Proposition 4.**  $\mathcal{J}(e_{\lambda})$  is nonsingular and defines a nontrivial twisting of  $U(\mathfrak{g})[[t]]$  in the limit  $q \to 1$ .

Proof. By Proposition 3 we have

$$\mathcal{J}(e_{\lambda}) = \exp_{q_{\lambda}}\left(\frac{t}{1 - q_{\lambda}} \ 1 \otimes e_{\lambda}\right) \exp_{q_{\lambda}^{-1}}\left(-\frac{t}{1 - q_{\lambda}} \left(\Delta(e_{\lambda}) - e_{\lambda} \otimes 1\right)\right).$$

The latter representation is nonsingular and from  $\hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})[[t]]$ , which is obvious if one uses the Campbell-Hausdorff formula after applying the dilogarithmic representation of q-exponent [4]:

$$\exp_{q_{\lambda}}\left(\frac{t}{1-q_{\lambda}} \ x\right) = \exp\left(\sum_{n\geq 1} \frac{t^n}{n(1-q_{\lambda}^n)} \ x^n\right)$$
(17)

along with the properties (14) and

$$[e_{\lambda}, \hat{U}_q^+(\mathfrak{g})] \in (q-1) \; \hat{U}_q^+(\mathfrak{g}).$$

Note, that to be self-consistent one can directly verify that (17) satisfies (20), considering (20) as a functional equation for the function  $\text{Li}_2(t \cdot x, q)$ :  $\text{Li}_2(t \cdot x, q) = \ln(\exp_q(\frac{t}{1-q} x))$ . Finally,  $\overline{\mathcal{J}}$  is a twist by Proposition 1.

**Example 1.** Consider  $\mathfrak{g} = \mathfrak{sl}_{N+1}$ . We have the following formula for the coproduct associated with the chosen normal ordering

roots without 
$$\alpha_1$$

$$\alpha_1 \succ \alpha_1 + \alpha_2 \succ \cdots \succ \alpha_1 + \alpha_2 + \cdots + \alpha_N \succ \alpha_2 \succ \cdots \succ \alpha_{N-1} + \alpha_N \succ \alpha_N$$

given by

$$\Delta(e_{\epsilon_1-\epsilon_{N+1}}) = k_{\epsilon_1-\epsilon_{N+1}}^{-1} \otimes e_{\epsilon_1-\epsilon_{N+1}} + e_{\epsilon_1-\epsilon_{N+1}} \otimes 1 + (1-q^2) \sum_{i=1}^{N-1} e_{\epsilon_1-\epsilon_{i+1}} k_{\epsilon_{i+1}-\epsilon_{N+1}}^{-1} \otimes e_{\epsilon_{i+1}-\epsilon_{N+1}} \otimes 1 + (1-q^2) \sum_{i=1}^{N-1} e_{\epsilon_1-\epsilon_{N+1}} k_{\epsilon_1+1}^{-1} \otimes 1 + (1-q^2) \sum_{i=1}^{N-1} e_{\epsilon_1-\epsilon_{N+1}} k_$$

where  $\alpha_i = \epsilon_i - \epsilon_{i+1}$ .

To calculate specialization  $\overline{\mathcal{J}(e_{\epsilon_1-\epsilon_{N+1}})}$ , we represent  $\mathcal{J}(e_{\epsilon_1-\epsilon_{N+1}})$  in the following form

$$\mathcal{J}(e_{\epsilon_1 - \epsilon_{N+1}}) = \exp_{q^{-2}}(-t \sum_{i=1}^{N-1} e_{\epsilon_1 - \epsilon_{i+1}} k_{\epsilon_{i+1} - \epsilon_{N+1}}^{-1} \otimes e_{\epsilon_{i+1} - \epsilon_{N+1}} C_{1,N+1}) \mathcal{J}_1$$

where

$$C_{1,N+1} = \exp_{q^2}\left(-\frac{q\ t}{1-q^2}\ e_{\epsilon_1-\epsilon_{N+1}}\right) \cdot \exp_{q^{-2}}\left(\frac{t}{1-q^2}\ e_{\epsilon_1-\epsilon_{N+1}}\right)$$
(18)

and

$$\mathcal{J}_1 = \exp_{q^2}\left(\frac{t}{1 - q^2} \ 1 \otimes e_{\epsilon_1 - \epsilon_{N+1}}\right) \exp_{q^{-2}}\left(-\frac{t}{1 - q^2} \ k_{\epsilon_1 - \epsilon_{N+1}}^{-1} \otimes e_{\epsilon_1 - \epsilon_{N+1}}\right). \tag{19}$$

Calculation of (18) - (19) is based on the Heine's formula from [6]:

$$1 + \sum_{n>1} t^n \frac{(\alpha)_q \cdots (\alpha + n - 1)_q}{(n)_q!} x^n = \exp_q(\frac{t}{1-q} x) \exp_{q^{-1}}(-\frac{q^{\alpha}t}{1-q} x).$$
 (20)

Note that (20) can be recast so that to hold in  $\hat{U}_q(\mathfrak{g}) \otimes \hat{U}_q(\mathfrak{g})[[t]]$ :

$$1 \otimes 1 + \sum_{n \ge 1} \frac{t^n}{(n)_{q^2}!} \left( \frac{k_{\epsilon_1 - \epsilon_{N+1}}^{-1} - 1}{q^2 - 1} \right) \cdots \left( \frac{k_{\epsilon_1 - \epsilon_{N+1}}^{-1} q^{2(n-1)} - 1}{q^2 - 1} \right) \otimes e_{\epsilon_1 - \epsilon_{N+1}}^n = \mathcal{J}_1$$
(21)

as one checks  $\frac{k_{\epsilon_1-\epsilon_{N+1}}^{-1}q^{2(n-1)}-1}{q^2-1} \in \hat{U}_q(\mathfrak{g})$ . Applying the specialization map

$$a \mapsto \overline{a} := a \otimes_{\mathcal{A}} 1$$

to each of the tensor factors in  $\mathcal{J}(e_{\epsilon_1-\epsilon_{N+1}})$  we come to a formula of [5, 10]:

$$\overline{\mathcal{J}(e_{\epsilon_1-\epsilon_{N+1}})} = \exp(-t\sum_{i=1}^{N-1} E_{1,i+1} \otimes E_{i+1,N+1} e^{-\frac{1}{2}\sigma_{1,N+1}}) \cdot \left(1 \otimes 1 + \sum_{n\geq 1} (-1)^n t^n \frac{H_{1,N+1}(H_{1,N+1}-1)\cdots(H_{1,N+1}-n+1)}{2^n n!} \otimes E_{1,N+1}\right)$$

where

$$\sigma_{1,N+1} = \ln(1 - t \ E_{1,N+1}) = -\sum_{n>1} \frac{t^n}{n} \ E_{1,N+1}$$

and

$$E_{i,j} = \overline{e_{\epsilon_i - \epsilon_j}}$$
  $H_{1,N} = \overline{\frac{k_{\epsilon_1 - \epsilon_{N+1}} - 1}{q - 1}}$ 

# 4 The Cremmer-Gervais twist and its specialization at $q \rightarrow 1$

In this section we consider nontrivial quantum twists in  $U_q(\mathfrak{sl}_3)[[t]]$  and their semi-classical limits  $q \to 1$ . As is known from the classification of Belavin-Drinfeld triples, there are two possible Belavin-Drinfeld triples for  $\mathfrak{sl}_3$ . The first one, the empty triple, is accounted for the Drinfeld-Jimbo deformation itself, while the second is associated with another deformation which can be called the Cremmer-Gervais quantization [2] and there is a solution to (1) defining the twisting element providing a possibility to deform  $U_h(\mathfrak{sl}_3)[[h]]$  further. To be self-consistent we first recall the construction of this twist from [11] and then study different possibilities to define specialization  $q \to 1$ , unveiling a surprising connection with the Connes-Moscovici algebra  $\mathcal{H}_1$ .

### Proposition 5 ([11]). An element

$$\mathcal{J}_{CG} = \Phi_{CG} \cdot \mathcal{K} = \exp_{q_h^{-2}}(\xi \ e_{32} \otimes e_{12}) \ q_h^{h_{w_2} \otimes h_{w_1}}, \quad \xi \in h \cdot \mathbb{C}[[h]],$$

where

$$h_{w_1} = \frac{2}{3}e_{11} - \frac{1}{3}(e_{22} + e_{33}), \qquad h_{w_2} = \frac{1}{3}(e_{11} + e_{22}) - \frac{2}{3}e_{33}$$
 (22)

with  $w_{1,2}$  being the fundamental weights, is a twist.

*Proof.* It is clear that  $\mathcal{K} = q_h^{h_{w_2} \otimes h_{w_1}}$  defines an abelian twist of  $U_{q_h}(\mathfrak{sl}_3)[[h]]$ . It leads to the following new Hopf algebra  $U_{q_h}^{\mathcal{K}}(\mathfrak{sl}_3)[[h]]$  with the same algebra structure as for  $U_{q_h}(\mathfrak{sl}_3)[[h]]$  and the new deformed coproducts:

$$\Delta_{\mathcal{K}}(e_{12}) = q_h^{-2\ h_{1,-1}} \otimes e_{12} + e_{12} \otimes 1, \qquad \Delta_{\mathcal{K}}(e_{23}) = q_h^{h_{1,-2}} \otimes e_{23} + e_{23} \otimes q_h^{h_{1,0}}$$

$$\Delta_{\mathcal{K}}(e_{21}) = e_{21} \otimes q_h^{h_{2,-1}} + q^{-h_{0,1}} \otimes e_{21}, \qquad \Delta_{\mathcal{K}}(e_{32}) = e_{32} \otimes q_h^{-2h_{1,-1}} + 1 \otimes e_{32}$$

where  $h_{m,n} := m \ h_{w_1} + n \ h_{w_2}$ . We are done if we prove that  $\Phi_{CG}$  is a twist for  $U_{q_h}(\mathfrak{sl}_3)[[h]]$ . Indeed, explicitly (1) reads as the following

$$\exp_{q_h^{-2}}(\xi \ e_{32} \otimes e_{12} \otimes 1) \cdot \exp_{q_h^{-2}}(\xi \ (e_{32} \otimes q_h^{-2h_{1,-1}} + 1 \otimes e_{32}) \otimes e_{12}) =$$

$$\exp_{q_h^{-2}}(\xi \ 1 \otimes e_{32} \otimes e_{12}) \cdot \exp_{q_h^{-2}}(\xi \ e_{32} \otimes (q_h^{-2h_{1,-1}} \otimes e_{12} + e_{12} \otimes 1))$$

and by the characteristic property of q-exponent

$$\exp_a(x+y) = \exp_a(y) \exp_a(x); \qquad xy = q \ yx$$

$$\Box$$
 (1) holds.

Consider now the problem of defining specialization of  $\mathcal{J}_{CG}$ . To do so, we introduce from the beginning a  $\mathbb{C}(q)$  analogue of  $U_{q_h}^{\mathcal{K}}(\mathfrak{sl}_3)[[h]]$ , which we denote by  $U_q'(\mathfrak{sl}_3)$ . As an algebra  $U_q'(\mathfrak{sl}_3)$  is an extension of  $U_q(\mathfrak{sl}_3)$  obtained by

attaching elements  $L_i$  (the maximal lattice), so that  $K_j = \prod_{i=1}^3 L_i^{a_{ij}}$ . On the other hand, as a coalgebra  $U_q'(\mathfrak{sl}_3)$  has a new coproduct fixed uniquely by its values on the Chevalley generators

$$\Delta(L_i) = L_i \otimes L_i$$

$$\Delta(e_1) = L_1^{-2} L_2^2 \otimes e_1 + e_1 \otimes 1,$$
  $\Delta(e_2) = L_1 L_2^{-2} \otimes e_2 + e_2 \otimes L_1$ 

$$\Delta(f_1) = f_1 \otimes L_1^2 L_2^{-1} + L_2^{-1} \otimes f_1, \qquad \Delta(f_2) = f_2 \otimes L_1^{-2} L_2^2 + 1 \otimes f_2,$$

where  $L_i$  are invertible, pairwise commuting and satisfying

$$L_i e_j L_i^{-1} = q^{\delta_{i,j}} e_j, \qquad L_i f_j L_i^{-1} = q^{-\delta_{i,j}} f_j.$$

In the sense of [?] define the regular form  $\hat{U}'_q(\mathfrak{sl}_3)$  as a  $\mathcal{A} := \mathbb{C}[q, q^{-1}]_{(q-1)}$  algebra such that there is an isomorphism

$$\hat{U}'_q(\mathfrak{sl}_3) \otimes_{\mathcal{A}} \mathbb{C}(q) \approx U'_q(\mathfrak{sl}_3).$$

 $\hat{U}_q'(\mathfrak{sl}_3)$  is generated over  $\mathcal{A}$  by the following set of generators

$$\left\{L_i^{-1}, \frac{L_i - 1}{q - 1}, e_i, f_i\right\}$$

If we denote by the "barred" generators the images of generators under specialization map

$$a \mapsto \overline{a} := a \hat{\otimes}_A 1$$
,

then the generators of  $\hat{U}'_q(\mathfrak{sl}_3)$  specialize to the classical Chevalley generators of  $U(\mathfrak{sl}_3)$  as the following:

$$\overline{e}_1 = E_{12}, \qquad \overline{e}_2 = E_{23}, \qquad \frac{\overline{L_1 - 1}}{q - 1} = h_{w_1}$$

$$\overline{f}_1 = E_{21}, \qquad \overline{f}_2 = E_{32}, \qquad \frac{\overline{L_2 - 1}}{q - 1} = h_{w_2}$$
(23)

(see (22)). Let us additionally make completion  $U'_q(\mathfrak{sl}_3)[[t]]$  then we can formulate a q-analogue of Proposition 5:

Proposition 6. An element

$$\hat{\Phi}_{CG} = \exp_{q^{-2}}(\zeta \ f_2 \otimes e_1), \quad \zeta \in t \cdot \mathcal{A}[[t]]$$

is a twist in  $\hat{U}'_q(\mathfrak{sl}_3)[[t]]$ .

On the other hand apart from  $\hat{\Phi}_{CG}$  we can construct the twists in  $U_q'(\mathfrak{sl}_3)[[t]]$  which restrict to  $\hat{U}_q'(\mathfrak{sl}_3)[[t]]$  only after a suitable change of basis which is implemented by some coboundary twist, namely we can prove

Proposition 7. An element

$$\mathcal{F}_{CG} = (V \otimes V) \exp_{q^{-2}} \left( \frac{q^{-2} \cdot \zeta \cdot t}{1 - q^2} \ f_2 \otimes e_1 \right) \Delta(V^{-1}), \quad \zeta \in t \cdot \mathcal{A}[[t]],$$

where

$$V = \exp_{q^{-2}}\left(-\frac{\zeta}{1-q^2} e_1 L_1^2 L_2^{-2}\right) \cdot \exp_{q^2}\left(\frac{t}{1-q^2} f_2\right),$$

restricts to a twist of  $\hat{U}'_q(\mathfrak{sl}_3)[[t]]$ .

*Proof.* By the form of the coproducts

$$\Delta(e_1L_1^2L_2^{-2}) = e_1L_1^2L_2^{-2} \otimes L_1^2L_2^{-2} + 1 \otimes e_1L_1^2L_2^{-2}, \qquad \Delta(f_2) = f_2 \otimes L_1^{-2}L_2^2 + 1 \otimes f_2$$

we have explicitly

$$\mathcal{F}_{CG} = \left( V \otimes \exp_{q^{-2}} \left( -\frac{\zeta}{1 - q^2} \ e_1 L_1^2 L_2^{-2} \right) \right)$$

$$\exp_{q^{-2}} \left( \frac{q^{-2} \cdot \zeta \cdot t}{1 - q^2} \ f_2 \otimes e_1 \right) \exp_{q^{-2}} \left( -\frac{t}{1 - q^2} \ f_2 \otimes L_1^{-2} L_2^2 \right) \Delta \left( \exp_{q^2} \left( \frac{\zeta}{1 - q^2} \ e_1 L_1^2 L_2^{-2} \right) \right).$$

Using the five terms relation, [4]:

If  $[u, [u, v]]_{q^2} = [v, [u, v]]_{q^{-2}} = 0$  then

$$e_{q^2}(u) \cdot e_{q^2}(v) = e_{q^2}(v) \cdot e_{q^2}(\frac{1}{1 - q^2}[u, v]) \cdot e_{q^2}(u)$$
 (24)

we can simplify

$$\begin{split} \exp_{q^{-2}}(-\frac{t}{1-q^2} \ f_2 \otimes L_1^{-2}L_2^2) \exp_{q^{-2}}(-\frac{\zeta}{1-q^2} \ 1 \otimes e_1L_1^2L_2^{-2}) = \\ \exp_{q^{-2}}(-\frac{\zeta}{1-q^2} \ 1 \otimes e_1L_1^2L_2^{-2}) \exp_{q^{-2}}(\frac{q^{-2} \cdot \zeta \cdot t}{1-q^2} \ f_2 \otimes e_1) \exp_{q^{-2}}(-\frac{t}{1-q^2} \ f_2 \otimes L_1^{-2}L_2^2) \end{split}$$

and thus  $\mathcal{F}_{CG}$  transforms to the following form

$$\mathcal{F}_{CG} = \exp_{q^{-2}}\left(-\frac{t}{1-q^2}\left(f_2 \otimes L_1^{-2} L_2^2 + e_1 L_1^2 L_2^{-2} \otimes 1\right)\right) \exp_{q^2}\left(\frac{t}{1-q^2}\left(f_2 \otimes 1 + L_1^2 L_2^{-2} e_1 \otimes L_1^2 L_2^{-2}\right)\right)$$

and finally by the Heine's formula we have

$$\mathcal{F}_{CG} = 1 \otimes 1 + \sum_{n \geq 1} \frac{1}{(n)_{q^2}!} \left( 1 \otimes \frac{L_1^{-2}L_2^2 - 1}{q^2 - 1} \cdots \frac{L_1^{-2}L_2^2q^{2(n-1)} - 1}{q^2 - 1} \right) (t \cdot f_2 \otimes 1 + \zeta \cdot L_1^2 L_2^{-2} e_1 \otimes L_1^2 L_2^{-2})^n$$

П

the latter expression restricts to  $\hat{U}'_q(\mathfrak{sl}_3)[[t]]$  as we have

$$\frac{L_1^{-2}L_2^2q^{2(n-1)}-1}{q^2-1}\in \hat{U}_q'(\mathfrak{sl}_3)$$

# 5 Semi-classical twists and noncommutative geometry

**Proposition 8.** There is a semi-classical twist  $\mathcal{F}$  and a homomorphism

$$\iota:\mathcal{H}_1\to U^{\mathcal{F}}(\mathfrak{sl}_3)[[t]]$$

*Proof.* We solve more general problem of obtaining quantization  $\mathcal{H}'_{1,q}$  in the sense that there is an embedding

$$\iota_q:\mathcal{H}'_{1,q}\hookrightarrow U'_q(\mathfrak{sl}_3)[[t]]$$

where  $\mathcal{H}'_{1,q}$  denotes an appropriately defined q-deformation of  $\mathcal{H}'_1$ . Consider the following coboundary twist in  $U'_q(\mathfrak{sl}_3)[[t]]$ :

$$\mathcal{J} := (W \otimes W)\Delta(W^{-1}), \qquad W = \exp_{q^{-2}}(-\frac{t}{1-q^2} e_{1+2}L_1)$$

where

$$e_{1+2}L_1 := (e_1e_2 - q e_2e_1)L_1$$

has the following coproduct

$$\Delta(e_{1+2}L_1) = e_{1+2}L_1 \otimes L_1^2 + 1 \otimes e_{1+2}L_1 + (1-q^2) \ e_1L_1^2L_2^{-2} \otimes e_2L_1.$$

By the properties

$$(1 \otimes e_{1+2}L_1)(e_{1+2}L_1 \otimes L_1^2 + (1-q^2) e_1L_1^2L_2^{-2} \otimes e_2L_1) =$$

$$q^{-2}(e_{1+2}L_1 \otimes L_1^2 + (1-q^2)e_1L_1^2L_2^{-2} \otimes e_2L_1)(1 \otimes e_{1+2}L_1),$$

$$(e_1L_1^2L_2^{-2} \otimes e_2L_1)(e_{1+2}L_1 \otimes L_1^2) = q^{-2}(e_{1+2}L_1 \otimes L_1^2)(e_1L_1^2L_2^{-2} \otimes e_2L_1)$$

 $\mathcal J$  is nonsingular, the reasoning is same as in Proposition 4, and can be represented in the following form

$$\mathcal{J} = \overbrace{\operatorname{Ad} \exp_{q^{-2}}(-\frac{t}{1-q^2} \ e_{1+2}L_1 \otimes 1)(\exp_{q^2}(t \ e_1L_1^2L_2^{-2} \otimes e_2L_1))}^{\mathcal{J}_1} \cdot \underbrace{\exp_{q^{-2}}(-\frac{t}{1-q^2} \ e_{1+2}L_1 \otimes 1)\exp_{q^2}(\frac{t}{1-q^2} \ e_{1+2}L_1 \otimes L_1^2)}_{\mathcal{J}_2}.$$

Let us check that for both factors we have

$$\mathcal{J}_{1,2} \in \hat{U}'_q(\mathfrak{sl}_3) \otimes \hat{U}'_q(\mathfrak{sl}_3)[[t]].$$

Indeed, it follows from explicit form of the factors  $\mathcal{J}_{1,2}$ :

$$\mathcal{J}_1 = \exp_{q^2} \left( t \ e_1 L_1^2 L_2^{-2} \ \frac{1}{1 - t \ e_{1+2} L_1} \otimes e_2 L_1 \right),$$

$$\mathcal{J}_{2} = \exp_{q^{-2}} \left( \frac{q^{-2} t}{1 - q^{-2}} e_{1+2} L_{1} \otimes 1 \right) \left( \exp_{q^{-2}} \left( \frac{q^{-2} t}{1 - q^{-2}} e_{1+2} L_{1} \otimes L_{1}^{2} \right) \right)^{-1} =$$

$$= 1 \otimes 1 + \sum_{n \geq 1} t^{n} \frac{(-1)^{n}}{(n)_{q^{-2}}!} (e_{1+2} L_{1})^{n} \otimes \frac{(L_{1}^{2} - 1)}{q^{2} - 1} \cdot \frac{(L_{1}^{2} q^{-2} - 1)}{q^{2} - 1} \cdots \frac{(L_{1}^{2} q^{-2(n-1)} - 1)}{q^{2} - 1}.$$

Calculating specialization at q = 1 we come to

$$\overline{\mathcal{J}} = \exp(t \ E_{12} \frac{1}{1 - t \ E_{13}} \otimes E_{23}) \left( 1 \otimes 1 + \sum_{n \geq 1} (-1)^n t^n E_{13}^n \otimes \frac{h_{w_1}(h_{w_1} - 1) \cdots (h_{w_1} - n + 1)}{n!} \right)$$

 $\overline{\mathcal{J}}$  defines a noncoboundary deformation of  $U(\mathfrak{sl}_3)[[t]]$  as it follows from

$$\overline{\mathcal{J}}_{21} \neq \overline{\mathcal{J}}.$$

On the other hand its quantum counter part  $\mathcal{J}$  is coboundary in  $U_q'(\mathfrak{sl}_3)[[t]]$  and amounts to switching to another basis of  $U_q'(\mathfrak{sl}_3)[[t]]$ . In particular, the subset of generators

$$\{L_1L_2^{-1}, e_1, f_2\}$$

is changing in the following way

$$L_1^2 L_2^{-2} \mapsto W(L_1^2 L_2^{-2}) W^{-1} = L_1^2 L_2^{-2}, \qquad e_1 \mapsto W(e_1) W^{-1} = e_1 \frac{1}{1 - t \ e_{1+2} L_1}$$

$$f_2 \mapsto W(f_2)W^{-1} = f_2 - \frac{t}{1 - q^2} q^{-1}L_1^2L_2^{-2}e_1 \frac{1}{1 - t e_{1+2}L_1}$$

and we can form a Hopf subalgebra  $\mathcal{D}_q \subset U_q'^{\mathcal{J}}(\mathfrak{sl}_3)[[t]]$  generated by the set of generators

$$\left\{k := L_1^2 L_2^{-2}, x := f_2, z := q^{-1} L_1^2 L_2^{-2} e_1 \frac{1}{1 - t e_{1+2} L_1} \right\}.$$

with the following structure:

$$kxk^{-1} = q^2 x$$
,  $kzk^{-1} = q^2 z$ ,  $q^2xz - zx = -tz^2$ 

$$\Delta(k) = k \otimes k, \qquad \Delta(z) = z \otimes k + 1 \otimes z$$

$$\Delta(x) = x \otimes k^{-1} + 1 \otimes x + t \ z \otimes \frac{(k - k^{-1})}{1 - q^2}.$$

The structure of its specialization  $\mathcal{D}_1 \approx \mathcal{D}_q \hat{\otimes}_{\mathcal{A}} 1$  is the following:

$$[\overline{y}, \overline{x}] = \overline{x}, \qquad [\overline{y}, \overline{z}] = \overline{z}, \qquad \overline{xz} - \overline{zx} = -t\overline{z}^2$$

$$\Delta(\overline{y}) = \overline{y} \otimes 1 + 1 \otimes \overline{y}, \qquad \Delta(\overline{z}) = \overline{z} \otimes 1 + 1 \otimes \overline{z}$$

$$\Delta(\overline{x}) = \overline{x} \otimes 1 + 1 \otimes \overline{x} - t \ \overline{z} \otimes \overline{y}$$

where

$$\overline{y} := \overline{\frac{k-1}{q-1}}.$$

Finally we obtain the stated map

$$\iota: \mathcal{H}'_1 \to \mathcal{D}_1 \subset U^{\overline{\mathcal{J}}}(\mathfrak{sl}_3)[[t]]$$

by fixing its values on the generators

$$\iota(X) = -\frac{1}{2}\overline{x}, \qquad \iota(Y) = \overline{y}, \qquad \iota(Z) = t\overline{z}.$$

where the generators X, Y, Z fulfills the relations

$$[Y, X] = X,$$
  $[Y, Z] = Z,$   $[X, Z] = \frac{1}{2}Z^2$ 

and the coproducts are obtained from (2) by substitution Z for  $\delta_1$ . Next,  $\iota$  is an isomorphism as  $\iota$  maps the basis of  $\mathcal{H}'_1: \left\{X^kY^lZ^m\right\}_{k,l,m\geq 0}$  onto the basis

of 
$$\mathcal{D}_1: \left\{\overline{x}^k \overline{y}^l \overline{z}^m\right\}_{k,l,m>0}$$

Now we can obtain

#### Proposition 9. An element

$$F_q = (WV \otimes WV) \exp_{q^{-2}}(\frac{q^{-3} \cdot t^3}{(1-q^2)^2} f_2 \otimes e_1) \Delta(V^{-1})(W^{-1} \otimes W^{-1}),$$

where

$$V = \exp_{q^{-2}}\left(-\frac{q^{-1}t}{(1-q^2)^2} e_1 L_1^2 L_2^{-2}\right) \cdot \exp_{q^2}\left(\frac{t}{1-q^2} f_2\right),$$

restricts to a twist in  $\hat{U}_q^{\prime\mathcal{J}}(\mathfrak{sl}_3)[[t]]$ .

*Proof.* It is convenient to introduce notation

$$\tilde{\mathcal{F}}_{CG} = (V \otimes V) \exp_{q^{-2}}(\frac{q^{-3} \cdot t^3}{(1 - q^2)^2} f_2 \otimes e_1) \Delta(V^{-1}).$$

Then by the reasoning of Proposition 6 we can prove that

$$\tilde{\mathcal{F}}_{CG} = 1 \otimes 1 + \sum_{n \geq 1} \frac{1}{(n)_{q^2}!} \left( 1 \otimes \frac{L_1^{-2} L_2^2 - 1}{q^2 - 1} \cdots \frac{L_1^{-2} L_2^2 q^{2(n-1)} - 1}{q^2 - 1} \right) (t \cdot f_2 \otimes 1 + \frac{q^{-1} t^2}{1 - q^2} \cdot L_1^2 L_2^{-2} e_1 \otimes L_1^2 L_2^{-2})^n$$

Next, the following element

$$\tilde{\mathcal{F}}_{CG}^W := (W \otimes W) \tilde{\mathcal{F}}_{CG} \Delta(W^{-1})$$

is a twist in  $U'_q(\mathfrak{sl}_3)[[t]]$  and respectively

$$F_q = (W \otimes W)\tilde{\mathcal{F}}_{CG}(W^{-1} \otimes W^{-1})$$

defines a twist in  $U_q^{\prime\mathcal{J}}(\mathfrak{sl}_3)[[t]]$ . Explicitly

$$F_{q} = 1 \otimes 1 + \sum_{n \geq 1} \frac{1}{(n)_{q^{2}}!} \left( 1 \otimes \frac{L_{1}^{-2}L_{2}^{2} - 1}{q^{2} - 1} \cdots \frac{L_{1}^{-2}L_{2}^{2}q^{2(n-1)} - 1}{q^{2} - 1} \right) \cdot \left( t \cdot f_{2} \otimes 1 - t^{2} \cdot q^{-1} \ e_{1} \frac{1}{1 - t \ e_{1+2}L_{1}} \otimes \frac{L_{1}^{2}L_{2}^{-2} - 1}{q^{2} - 1} \right)^{n}$$

again similarly to Proposition 6 we check that  $F_q$  resticts to a twist in  $\hat{U}'_q^{\mathcal{J}}(\mathfrak{sl}_3)[[t]]$ . If we specialize q=1 and apply Proposition 8 we obtain

$$F_1 = 1 \otimes 1 + \sum_{n \geq 1} \frac{t^n}{n!} (1 \otimes Y(Y-1) \cdots (Y-n+1)) (2X \otimes 1 + Z \otimes Y)^n.$$

Note that a formula for  $F_1$  was obtained in [7] by a direct check in the study of the twists for  $\mathfrak{sl}_2$  Yangian. To make correspondence with (3) we additionally twist  $F_1$  by a coboundary:

$$F_1' = (\exp(t \ XY) \otimes \exp(t \ XY)) F_1 \Delta(\exp(-t \ XY))$$

expanding in t we have

$$F_1' = 1 \otimes 1 + t \cdot (X \otimes Y - Y \otimes X + ZY \otimes Y) + \cdots$$

Thus  $F_1'$  is equivalent to (3).

### References

- [1] A. Connes and H. Moscovici, Rankin-Cohen Brackets and the Hopf Algebra of Transverse Geometry, MMJ: Vol. 4 (2004),N1, math.QA/0304316.
- [2] E. Cremmer and J. -L. Gervais, The quantum group structure associated with non-linearly extended Virasoro algebras, Comm. Math. Phys. 134 (1990), 619-632.
- [3] C. DeConcini and V.G. Kac, Representations of quantum groups at roots of 1, In "Operator Algebras, Unitary Representations, Enveloping Algebras and Invariant Theory", Progress in Mathematics, Vol. 92, pp 471-506, Birkhauser, Boston, 1990.
- [4] L.D. Faddeev and R.M. Kashaev, Quantum dilogarithm, Modern Phys. Lett. A 9 (1994), 427-434, hep-th/9310070.
- [5] A. Giaquinto and J. Zhang, Bialgebra actions, twists, and universal deformation formulas, *Journal of Pure and Applied Algebra* 128(4) (1998), 133-151, hep-th/9411140.
- [6] V. G. Kac and P. Cheung, Quantum calculus, Springer, Berlin, (2002).

- [7] S.M. Khoroshkin, A.A. Stolin and V.N. Tolstoy, q-Power function over q-commuting variables and deformed XXX XXZ chains, *Physics of Atomic Nuclei* **64**(12) (2001), math.QA/0012207.
- [8] S.M. Khoroshkin and V.N. Tolstoy, Universal *R*-matrix for quantized (super)algebras, *Commun. Math. Phys.* **141**(3) (1991), 599-617.
- [9] L. Korogodski and Y. Soibelman, Algebras of functions on quantum groups, Providence, R.I.: AMS, 1998.
- [10] P.P. Kulish, V.D. Lyakhovsky and A.I. Mudrov, Extended jordanian twist for Lie algebras, *Journ. Math. Phys.* 40 (1999), 4569-4586, math.QA/9806014.
- [11] P.P. Kulish and A.I. Mudrov, Universal R-matrix for esoteric quantum group, Lett. Math. Phys. 47 (1999), 139-148, math.QA/9804006.
- [12] V.D. Lyakhovsky and M.E. Samsonov, Elementary parabolic twist, *Journal of Algebra and Its Applications*, **1**(4) (2002), math.QA/0107034.