## THE SMOOTHNESS OF RIEMANNIAN SUBMERSIONS WITH NONNEGATIVE SECTIONAL CURVATURE

JIANGUO CAO AND MEI-CHI SHAW

In this article, we study the smoothness of Riemannian submersions for open manifolds with non-negative sectional curvature. Suppose that  $M^n$  is a  $C^{\infty}$ -smooth, complete and non-compact Riemannian manifold with nonnegative sectional curvature. Cheeger-Gromoll [ChG] established a fundamental theory for such a manifold. Among other things, they showed that  $M^n$  admits a totally convex exhaustion  $\{\Omega_u\}_{u\geq 0}$  of  $M^n$ , where  $\Omega_0 = S$  is a totally geodesic and compact submanifold without boundary. Furthermore,  $M^n$  is diffeomorphic to the normal vector bundle of the soul S.

Sharafutdinov found that there exists a distance non-increasing retraction  $\Psi$ :  $M^n \to \mathcal{S}$  from the open manifold  $M^n$  of non-negative sectional curvature to its soul, (cf. [Sh], [Y2]). Perelman [Per] further showed that such a map  $\Psi$  is indeed a  $C^1$ -smooth Riemannian submersion. Furthermore,  $\Psi[Exp_q(t\vec{v})] = q$  for any  $q \in \mathcal{S}$ and  $\vec{v} \perp T_q(\mathcal{S})$ . Therefore, the fiber  $F_q = \Psi^{-1}(q)$  is a k-dimensional submanifold, which is  $C^\infty$ -smooth almost everywhere, where  $k = \dim(M^n) - \dim(\mathcal{S}) > 0$ .

Guijarro [Gu] proved that the fiber  $F_q$  is indeed a  $C^2$ -smooth submanifold for each  $q \in S$ . In this paper, we prove that the fibres are  $C^{\infty}$ -smooth.

**Theorem 1.** Let  $M^n$  be a complete, non-compact and  $C^{\infty}$ -smooth Riemannian manifold with nonnegative sectional curvature. Suppose S is a soul of  $M^n$ . Then any distance non-increasing retraction  $\Psi: M^n \to S$  must give rise to a  $C^{\infty}$ -smooth Riemannian submersion.

Consequently, if  $\mathbb{R}^k = \mathcal{N}_q(\mathcal{S}, M^n)$  is the normal space of the soul  $\mathcal{S}$  in  $M^n$  at q, then the fiber  $F_q = \Psi^{-1}(q) = Exp_q(\mathbb{R}^k)$  is a k-dimensional  $C^{\infty}$ -smooth submanifold of  $M^n$ , for any  $q \in S$ .

Professor Wilking kindly informed us that he has recently obtained a similar result (cf. [Wi]). His method is completely independent of ours. Our proof of

Both authors are supported by NSF Grants. Research at MSRI is supported in part by NSF grant DMS-9810361. The first author is grateful to MSRI and Max-Planck Institute for Mathematics at Leipzig for their hospitality.

Theorem 1 uses a flat strip theorem associated with Cheeger-Gromoll exhaustion (cf. Theorem 4 below), an uniform estimate for cut-radii of convex subsets in [ChG] and a smooth extension theorem for ruled surfaces.

For each compact convex subset  $\Omega \subset M^n$ , we let  $U_{\epsilon}(\Omega) = \{x \in M^n | d(x, \Omega) < \epsilon\}$ . Its cut-radius is given by  $\delta_{\Omega} = \sup\{\epsilon | \text{ there is a unique nearest point projection } \mathcal{P}_{\Omega} : U_{\epsilon}(\Omega) \to \Omega\}.$ 

For each  $x \in M^n$ , we let  $\operatorname{Inj}_{M^n}(x)$  be the injectivity radius of  $M^n$  at x. Similarly, let  $\operatorname{Inj}_{M^n}(A) = \sup\{\operatorname{Inj}_{M^n}(x) | x \in A\}.$ 

A subset  $\Omega$  of a complete Riemannian manifold  $M^n$  is said to be *totally convex* if for any pair of points  $\{p,q\} \subset \Omega$  and for any geodesic segment  $\sigma$  joining p and q, the geodesic segment  $\sigma$  is contained in  $\Omega$ . There is a totally convex exhaustion  $\{\Omega_u\}_{u\geq 0}$  of  $M^n$  given in [ChG]. By comparing the inner angles of geodesic triangles, we have the following semi-global estimate for cut-radius.

**Lemma 2.** (Lemma 2.4 of [ChG], [CaS]) Let  $A \subset \Omega_T$  be a connected, convex and compact subset in a Riemannian manifold  $M^n$  with nonnegative curvature, let  $K_0 = \max\{K(x)|x \in \Omega_{T+1}\}$ ,  $Inj_{M^n}(\Omega_T)$  be the upper bound of sectional curvature on  $\Omega_{T+1}$  and S be as above. Suppose that  $\dim(\Omega_T) = n$ . Then the subset A has cut-radius bounded below by

$$\delta_A \ge \delta_0(T) = \frac{1}{4} \min\{Inj_{M^n}(\Omega_T), \frac{\pi}{\sqrt{K_0}}, 1\},\$$

where  $\delta_0(T)$  is independent of choices of A with  $A \subset \Omega_T$ .

Let us briefly recall the Cheeger-Gromoll convex exhaustion. According to [ChG], there is a partition  $a_0 = 0 < a_1 < ... a_m < a_{m+1} = \infty$  of  $[0, \infty)$  and an exhaustion  $\{\Omega_u\}_{u>0}$  of  $M^n$  such that the following holds:

(1)  $M^n = \bigcup_{u \ge 0} \Omega_u$ . If  $u > a_m$  then  $dim[\Omega_u] = n$ . If  $u \le a_m$ , then  $\dim[\Omega_u] < n$ .

(2)  $\Omega_0 = S$  is the soul of  $M^n$ , which is a totally geodesic  $C^{\infty}$ -smooth compact submanifold without boundary.

(3) If u > 0,  $\Omega_u$  is a totally convex, compact subset of  $M^n$  and hence  $\Omega_u$  is a compact submanifold with a  $C^{\infty}$ -smooth relative interior. Furthermore, dim $(\Omega_u) = k_u > 0$ and  $\Omega_u$  has a non-empty  $(k_u - 1)$ -dimensional relative boundary  $\partial \Omega_u$ ;

(4) For any  $u_0 \in [a_j, a_{j+1}]$  and  $0 \leq t \leq u_0 - a_j$ , the family  $\{\Omega_{u_0-t}\}_{t \in [0, u_0-a_j]}$  is given by the inward equidistant evolution:

$$\Omega_{u_0-t} = \{ x \in \Omega_{u_0} | d(x, \partial \Omega_{u_0}) \ge t \}.$$

$$(2.1)$$

(5) If  $u > a_m$  then  $u - a_m = \max\{d(x, \partial \Omega_u) | x \in \Omega_u\}$ . If  $0 \le j \le m - 1$  then  $a_{j+1} - a_j = \max\{d(x, \partial \Omega_{a_{j+1}}) | x \in \Omega_{a_{j+1}}\}$  and hence  $\dim[\Omega_{a_j}] < \dim[\Omega_{a_{j+1}}]$  for  $j \ge 0$ .

Assume that  $k = \dim[M^n] - \dim[\mathcal{S}] = \dim(F_q)$  for all  $q \in \mathcal{S}$ . Since  $M^n = \bigcup_{T \ge 0} \Omega_T$ , it is sufficient to verify that the subset  $[U_{\delta_0(T)}(\Omega_T) \cap F_q]$  has a k-dimensional  $C^{\infty}$ -smooth interior, where  $\delta_0(T)$  is given by Lemma 2 and  $T > a_m$ .

For this purpose, we need to study the geometry of the equidistant hypersurfaces from  $\partial \Omega_u$ . Federer [Fe] has studied the smoothness of the outward equidistant hypersurfaces  $\partial [U_{\epsilon}(\Omega)]$  for  $0 < \epsilon < \delta_{\Omega}$ . Following his approach, we consider the outward normal cone of  $\Omega$  as follows:

$$\mathcal{N}^+(\Omega, M^m) = \{ (p, \vec{v}) | p \in \Omega, d(Exp_p(t\vec{v}), \Omega) = t | \vec{v} |, \text{ for } 0 \le t | \vec{v} | < \delta_\Omega \}.$$

If  $\{\Omega_u\}$  is the Cheeger-Gromoll convex exhaustion as above and u > 0, then the relative boundary  $\partial \Omega_u$  is not necessarily smooth. We are going to study the corresponding decomposition of  $\mathcal{N}^+(\Omega, M^m)$ :

$$\mathcal{N}_p^+(\Omega_u, M^n) \subset \left[\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon})) \oplus \mathcal{N}_p^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)\right],$$
(2.2)

where  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  is defined by

$$\mathcal{N}^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon})) = \{(p, \vec{v}) | p \in \Omega_u, d(Exp_p(t\vec{v}), \Omega_u) = t | \vec{v} |, \\ \text{for } 0 \le t | \vec{v} | < \delta_{\Omega_u}, Exp_p(t\vec{v}) \in \operatorname{int}(\Omega_{u+\epsilon}) \}.$$

Our next step is to choose  $\epsilon$  sufficiently small so that (1) there is a nearest point projection  $\mathcal{P}$  :  $\operatorname{int}(\Omega_{u+\epsilon}) \to \Omega_u$ ; and (2)  $\Omega_u = \{x \in \Omega_{u+\epsilon} | d(x, \partial \Omega_{u+\epsilon}) \geq \epsilon\}$  holds. We first find j so that  $a_j \leq u < a_{j+1}$  for some  $0 \leq j \leq m$ . Let  $T = u + a_m + 1$  and  $\delta_0(T)$  be given by Lemma 2. It follows from a result of Yim that there is a constant  $C_T$  such that, for  $0 \leq u_1 < u_2 \leq T$ , we have

$$\max\{d(x,\Omega_{u_1})|x\in\Omega_{u_2}\} \le C_T(u_2-u_1),\tag{2.3}$$

see [Y2, Theorem A.5(3)]. In what follows, we always choose

$$0 < \epsilon = \epsilon_u < \min\{[a_{j+1} - u], \frac{\delta_0(T)}{2C_T}\},$$
(2.4)

where  $u \in [a_j, a_{j+1})$ ,  $T = u + a_m + 1$  and  $\delta_0(T)$  is given by Lemma 2.

With such a choice of  $\epsilon = \epsilon_u$  by (2.4), the geometry of  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  is determined by its minimal normal vectors which we now describe.

**Definition 3.** (Minimal normal vector) Let  $\Omega_u$ ,  $\Omega_{u+\epsilon}$  and  $\mathcal{N}^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  be as above. Let  $\sigma_{(p,\vec{v})} : [0,\epsilon] \to M^n$  be a geodesic given by  $\sigma_{(p,\vec{v})}(t) = \operatorname{Exp}_p(t\frac{\vec{v}}{|\vec{v}|})$ , where  $\vec{v} \neq 0$ . If  $\sigma_{(p,\vec{v})}$  is a length-minimizing geodesic from  $p \in \Omega_u$  to  $\partial \Omega_{u+\epsilon}$ , then  $\vec{v}$  is called a minimal normal vector in  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$ . It is known that any other normal vector  $\vec{w} \in \mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  can be expressed as a linear combination of *minimal normal vectors* at p. Moreover, the convex hull of minimal normal vectors at p is equal to  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$ , (cf. [Y1, Proposition 1.7]).

For each  $p \in M^n$ , we let  $\mathcal{V}_p = T_p(F_{\Psi(p)})$  and  $\mathcal{H}_p = [\mathcal{V}_p]^{\perp}$ . A geodesic  $\alpha : [a, b] \to M^n$  is said to be horizontal, if  $\alpha'(t) \perp F_{\Psi(\alpha(t))}$  for all  $t \in [a, b]$ . We need the following flat strip theorem for the proof of Theorem 1.

**Theorem 4.** Let  $\{\Omega_u\}$  be the Cheeger-Gromoll totally convex exhaustion of  $M^n$ as above. Suppose that  $\Psi: M^n \to S$  be a distance non-increasing retraction and  $F_q = \Psi^{-1}(q)$  be a fibre for some  $q \in S$ . Then for  $p \in F_q \cap \Omega_u$  and any  $(p, \vec{v}) \in \mathcal{N}^+(\Omega_u, M^n)$ , we have

$$\Psi[Exp_p([\mathbb{R}\{\vec{v}\}])] = \Psi(p) = q. \tag{4.1}$$

Moreover, if dim(S)  $\geq 1$  and if  $\vec{w} \in \mathcal{H}_p$  has  $|\vec{w}| = 1 = |\vec{v}|$ , then the surface  $\Sigma^2_{\vec{v},\vec{w}} = Exp_p[\mathbb{R}\{\vec{v}\} \oplus \mathbb{R}\{\vec{w}\}]$  is totally geodesic immersed flat plane in  $M^n$ .

A result similar to Theorem 4 was proved in [CaS] via a totally different method.

Proof of Theorem 4. Theorem 4 was proved by Perelman [Per] for the case of  $\Omega_0 = \mathcal{S}$ . Applying Perelman's argument for the case of  $p \notin \mathcal{S}$ , Guijarro [Gu1] found the following sufficient condition for (4.1).

(4.2)  $\vec{v} \in \mathcal{V}_p$  stays vertical under parallel transport along any horizontal broken geodesic.

Guijarro showed that (4.1) follows from (4.2). Moreover, if (4.2) holds and if  $\vec{w} \in \mathcal{H}_p$  has  $|\vec{w}| = 1 = |\vec{v}|$ , then the surface  $\Sigma^2_{\vec{v},\vec{w}} = Exp_p[\mathbb{R}\{\vec{v}\} \oplus \mathbb{R}\{\vec{w}\}]$  is totally geodesic immersed flat plane in  $M^n$ , (cf. Theorem 3.1 of [Gu1]).

In order to see that  $\mathcal{N}^+(\Omega_u, M^n) \subset \mathcal{V}_p$  holds, we recall that any horizontal geodesic  $\alpha$  is contained a tubular neighborhood of the soul  $\mathcal{S}$ , by Perelman's theorem [Per]. Hence,  $\alpha$  is contained in a compact totally geodesic subset  $\Omega_T$  for a sufficiently large T. It follows from Theorem 5.1 of [ChG] that  $\alpha \subset \partial \Omega_u$  for some u, (cf. [Gu2]).

(4.3) Any horizontal geodesic  $\alpha$  with  $\alpha(0) \in \partial \Omega_u$  must be entirely contained in  $\partial \Omega_u$ . Consequently,  $\mathcal{H}_p \subset T_p^-(\partial \Omega_u)$ , where  $T_p^-(\partial \Omega_u)$  is the tangent cone of  $\partial \Omega_u$  at p.

Recall that by (2.2) we have

$$\mathcal{N}_p^+(\Omega_u, M^n) \subset \left[\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon})) \oplus \mathcal{N}_p^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)\right].$$

For  $\vec{v}$  in either  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  or  $\mathcal{N}_p^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)$ , we will show that such a  $\vec{v}$  satisfies (4.2).

It follows from Theorem 1.10 of [ChG] (or Corollary 1.4 of [Y1]) that any minimal normal vector  $\vec{v}$  of  $\mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  stays minimal under parallel transport along any geodesic in  $\partial\Omega_u$ . Since the convex hull of minimal normal vectors is equal to the outward normal cone (cf. [Y1, Proposition 1.7]), the bundle  $\mathcal{N}^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$ is invariant under parallel transport along any geodesic in  $\partial\Omega_u$ . This together with (4.3) implies that if  $\vec{v} \in \mathcal{N}_p^+(\Omega_u, \operatorname{int}(\Omega_{u+\epsilon}))$  then  $\vec{v}$  satisfies (4.2).

For  $\vec{v} \in \mathcal{N}_p^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)$ , the assertion (4.2) follows from Corollary 3.2 of [Gu1]. In fact, since  $\operatorname{int}(\Omega_{u+\epsilon})$  is totally convex and totally geodesic, both  $T(\operatorname{int}(\Omega_{u+\epsilon}))$ and  $N^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)$  are invariant under parallel transport along any geodesic in  $\operatorname{int}(\Omega_{u+\epsilon})$ . This together with (4.3) implies that (4.2) holds for any  $\vec{v} \in \mathcal{N}_p^+(\operatorname{int}(\Omega_{u+\epsilon}), M^n)$ .

Therefore, (4.2) holds for any  $\vec{v} \in \mathcal{N}_p^+(\Omega_u, M^n)$ . This completes the proof of Theorem 4.  $\Box$ 

In order to see that Theorem 4 implies Theorem 1, we need to establish a bootstrap argument for the smoothness of ruled surfaces. A  $C^1$ -smooth one-parameter family of a straight lines in  $\mathbb{R}^3$  gives rise to a ruled surface. Suppose that  $\{\beta(s), \vec{v}(s)\}$ are  $C^1$ -smooth vector valued functions with  $[\beta'(s) + t\vec{v}'(s)] \wedge \vec{v}(s) \neq 0$  for all  $(s,t) \in (a,b) \times (c,d)$ . Then we have a corresponding  $C^1$ -smooth immersed ruled surface.

$$F: (a,b) \times (c,d) \to \mathbb{R}^3$$
$$(s,t) \to \beta(s) + t\vec{v}(s)$$

Our bootstrap argument is motivated by the following observation.

**Lemma 5.** (The smooth extension for ruled surfaces in  $\mathbb{R}^3$ ) Let  $F((a, b) \times (c, d)) = \Sigma^2$  be an embedded ruled surface in  $\mathbb{R}^3$  and let  $F: (a, b) \times (c, d) \to \mathbb{R}^3$  be a  $C^{1,1}$ -smooth embedding map be as above. Suppose that a subset  $\hat{\Sigma}^2_{\epsilon} = F((a, b) \times (\epsilon_1, \epsilon_2))$  is a  $C^{\infty}$ -smooth embedded surface of  $\mathbb{R}^3$ , where  $(\epsilon_1, \epsilon_2) \subset (c, d)$ . Then the whole ruled surface  $\Sigma^2$  is a  $C^{\infty}$ -smooth surface of  $\mathbb{R}^3$ .

Proof. By our assumption, F is an embedding map, and hence the surface  $\hat{\Sigma}_{\epsilon}^2 = F((a,b) \times (\epsilon_1, \epsilon_2))$  is foliated by straight lines. Because the surface  $\hat{\Sigma}_{\epsilon}^2$  and each orbit (each straight line) are  $C^{\infty}$ , the quotient space  $Q = [\hat{\Sigma}^2/\sim]$  is a  $C^{\infty}$ -smooth 1-dimensional space as well, where  $\sim$  is the equivalent relation induced by the orbits (the ruling straight lines). Thus, we have a fibration  $(\epsilon_1, \epsilon_2) \longrightarrow \hat{\Sigma}_{\epsilon}^2 \longrightarrow Q$ . We may assume that Q = (0, 1). Let  $\pi : \hat{\Sigma}_{\epsilon}^2 \to Q$  be the quotient map. Because the fibration is topologically trivial, we can find two disjoint  $C^{\infty}$ -smooth cross-sections

$$h_i: \quad Q \to \hat{\Sigma}_{\epsilon}^2$$
$$u \to h_i(u)$$

for i = 0, 1, where  $\pi(h_i(u)) = 0$ . (Since the fibre is 1-dimensional line, we may assume that the graph of the cross-section  $h_1$  lies above that of  $h_0$ ). Because

 $h_0(Q)$  and  $h_1(Q)$  are disjoint, we obtain a new  $C^{\infty}$ -smooth parametrization of the ruled surface  $G: \quad Q \times \mathbb{R} \to \mathbb{R}^3$ 

: 
$$Q \times \mathbb{R} \to \mathbb{R}^{3}$$
  
 $(u, \lambda) \to h_0(u) + \lambda \frac{[h_1(u) - h_0(u)]}{\|h_1(u) - h_0(u)\|}$ 

Clearly, G is a  $C^{\infty}$ -smooth map with  $\Sigma^2 \subset G(Q \times \mathbb{R})$ . Because F is an embedding map, on the subset  $G^{-1}(\Sigma^2)$ , one can check that G remains to be injective and with non-vanishing Jacobi  $G_u \wedge G_\lambda \neq 0$ . Hence,  $G|_{G^{-1}(\Sigma^2)}$  is an embedding as well. Thus,  $\Sigma^2$  is a  $C^{\infty}$ -smooth embedded surface.  $\Box$ 

The proof of Lemma 5 can be applied to the proof of Theorem 1 as follows. Let  $\Omega_u$  be a totally convex subset as above. By Federer's Theorem, the hypersurface  $\partial[U_{\delta}(\Omega_u)]$  is  $C^{1,1}$ -smooth if  $\delta$  is less than the cut-radius of  $\Omega_u$ . Assume that T > u and  $d = \delta_T - \delta > 0$ . Let  $\vec{v}(x)$  be the outward unit normal vector of  $\partial[U_{\delta}(\Omega)]$  at x. There is an embedding:

$$F: \quad \partial[U_{\delta}(\Omega)] \times (c, d) \to M^{n}$$
  
(x, t)  $\to \operatorname{Exp}_{x}[t\vec{v}(x)]$  (6.1)

where  $c = -\delta$ .

**Proposition 6.** (The smooth extension for the ruled sub-manifold) For each  $y \in \partial[U_{\delta}(\Omega)]$ , we let  $B_r^{k-1}(y) \subset \partial[U_{\delta}(\Omega)]$  be a small k-dimensional ball around y which is  $C^1$ -diffeomorphic to  $B^{k-1}(0) \subset \mathbb{R}^k$  and let  $\Omega_u, \delta, \delta_T, c, d$  and F be as above. Suppose that F is an  $C^{1,1}$ -smooth embedding and that  $\hat{\Sigma}_{\epsilon}^k = F(B_r^{k-1}(y) \times (\epsilon_1, \epsilon_2))$  is a  $C^{\infty}$ -smooth embedded k-submanifold of  $M^n$ , where  $(\epsilon_1, \epsilon_2) \subset (c, d)$ . Then the whole ruled submanifold  $\Sigma^k = F(B_r^{k-1}(y) \times (c, d))$  is a  $C^{\infty}$ -smooth submanifold of  $M^n$ .

Proof. The proof of Proposition 6 is the same as above with minor modifications. By our assumption,  $\hat{\Sigma}^k_{\epsilon}$  is foliated by  $C^{\infty}$ -smooth open geodesic segments. The quotient space  $Q = [\hat{\Sigma}^k_{\epsilon}/\sim]$  is a  $C^{\infty}$ -smooth (k-1)-dimensional open manifold. Because the fibration  $(\epsilon_1, \epsilon_2) \longrightarrow \hat{\Sigma}^k_{\epsilon} \longrightarrow Q$  is trivial, we can choose two *disjoint* cross sections  $h_0: Q \to \hat{\Sigma}^k_{\epsilon}$  for i = 0, 1. If  $\pi: \hat{\Sigma}^k_{\epsilon} \to Q$  is the quotient map, then  $\pi \circ h_i(u) = u$  for all  $u \in Q$ . Since the two cross-sections are disjoint, we may assume that  $r(h_1(u)) > r(h_0(u))$  for all  $u \in Q$ , where  $r(y) = d(y, \partial[U_{\delta}(\Omega)])$ . For each  $u \in Q$ , we consider the unit vector

$$\vec{\eta}(u) = \frac{\operatorname{Exp}_{h_0(u)}^{-1}[h_1(u)]}{\|\operatorname{Exp}_{h_0(u)}^{-1}[h_1(u)]\|}$$

at the point  $h_0(u)$ . Similarly, we consider a new  $C^{\infty}$ -smooth parametrization

(

$$G: \quad Q \times \mathbb{R} \to M^n$$
$$(u, \lambda) \to \operatorname{Exp}_{h_0(u)}[\lambda \vec{\eta}(u)].$$

Clearly, we have  $\Sigma^k = F(B_r^{k-1}(y) \times (c,d)) \subset G(Q \times \mathbb{R})$ . This completes the proof.  $\Box$ 

With Lemma 2, Theorem 4 and Proposition 6, we are ready to prove Theorem 1.

Proof of Theorem 1. Let  $\{\Omega_u\}$  be a Cheeger-Gromoll convex exhaustion described as above. It is sufficient to verify that the subset  $[U_{\delta_0(T)}(\Omega_T) \cap F_q]$  has a k-dimensional  $C^{\infty}$ -smooth interior for any given  $T > a_m$  and  $q \in \mathcal{S}$ , where  $\delta_0(T)$  is given by Lemma 2.

Fix  $T > a_m$  with dim $[\Omega_T] = n$ . Let  $\delta_0(T)$  be given by Lemma 2 and  $C_T$  be given by (2.3). Choose a partition  $0 = u_0 < u_1 < ... < u_N = T$  of [0, T] such that  $u_j - u_{j-1} < \frac{2C_T}{\delta_0(T)}$  for j = 1, ..., N, where  $N = N_T$  is a number depending on T.

We will prove the following assertion by induction on j = 0, 1, ..., N.

**Assertion j.** The sub-level set  $[U_{\delta_0(T)}(\Omega_{u_j}) \cap F_q]$  has the k-dimensional  $C^{\infty}$ -smooth interior, where  $q \in S$  and  $k = \dim[F_q]$ .

It follows from Perelman's theorem or Theorem 4 that  $\operatorname{Exp}_q[\mathcal{N}_q^+(\mathcal{S}, M^n)] \subset F_q$ . Since the soul  $\mathcal{S}$  has the cut radius  $\geq \delta_0(T)$  and  $\mathcal{S}$  is  $C^{\infty}$ -smooth, Assertion 0 holds. Let  $\epsilon_1 = \frac{\delta_0(T)}{16}$  and  $\epsilon_2 = \frac{\delta_0(T)}{8}$ . We consider

$$A(\Omega_{u_i}, r_1, r_2) = \{ z \in F_q | \quad 0 < r_1 < d(z, \Omega_{u_i}) < r_2 \}$$

It is clear that  $A(\Omega_{u_1}, \epsilon_1, \epsilon_2) \subset U_{\delta_0(T)}(\mathcal{S})$ . It follows from Assertion 0 that the subset  $\hat{\Sigma}^k_{\epsilon} = A(\Omega_{u_1}, \epsilon_1, \epsilon_2) \subset F_q \cap U_{\delta_0(T)}(\mathcal{S})$  is  $C^{\infty}$ -smooth k-dimensional open sub-manifold. By Theorem 4, we let  $\Sigma^k_1 = A(\Omega_{u_1}, \frac{\delta_0(T)}{16}, \delta_0(T))$  be the ruled k-dimensional submanifold. It follows from Proposition 6 (the smooth extension theorem for the ruled submanifold) that  $\Sigma^k_1$  is a  $C^{\infty}$ -smooth k-dimensional sub-manifold of  $M^n$ . Observe that the subset  $[U_{\delta_0(T)}(\Omega_{u_j}) \cap F_q]$  is contained in the union  $\{[U_{\delta_0(T)}(\mathcal{S}) \cap F_q] \cup \Sigma^k_1\}$ . Since  $\Sigma^k_1$  is a  $C^{\infty}$ -smooth, Assertion 1 follows from Assertion 0.

Similarly, using Theorem 4 and Proposition 6 we can verify that Assertion (j-1) is true then Assertion j holds as well for  $j \geq 2$ . In fact, by induction we see that  $A(\Omega_{u_j}, \epsilon_1, \epsilon_2) \subset [U_{\delta_0(T)}(\Omega_{u_{j-1}}) \cap F_q]$  is  $C^{\infty}$ -smooth. It follows from Theorem 4 and Proposition 6 that the ruled submanifold  $\Sigma_j^k = A(\Omega_{u_j}, \frac{\delta_0(T)}{16}, \delta_0(T))$  must be of  $C^{\infty}$ -smooth as well. Since  $[U_{\delta_0(T)}(\Omega_{u_j}) \cap F_q] \subset [U_{\delta_0(T)}(\Omega_{u_{j-1}}) \cap F_q] \cup \Sigma_j^k$ , Assertion j follows. Theorem 1 follows from Assertion  $N_T$  for any arbitrarily large T.  $\Box$ 

The first author is very grateful to Professor B. Wilking for pointing out a mistake in an earlier version of the manuscript [CaS].

## References

- [CaS] Cao, J. and Shaw, M., A new proof of the Cheeger-Gromoll soul conjecture and the Takeuchi theorem, Preprint.
- [ChG] Cheeger, J and Gromoll, D., On the structure of complete manifolds of nonnegative curvature,, Ann. of Math. **96** (1972), 413–443..
- [Fe] Federer, H, Curvature measures, Trans. Amer. Math. Soc. 93 (1959), 418-491.
- [Gu1] Guijarro, L., On the metric structure of open manifolds with nonnegative curvature, Pacific J. Math. **196** (2000), 429-444.
- [Gu2] \_\_\_\_\_, Nonnegative curvature and normal holonomy in open manifolds, In "Geometry and topology: Aarhus, (1998)" Amer. Math. Soc., Providence, RI, Contemp. Math. **258** (2000), 197–202,.
- [Hi] Hirsch, M., *Differential topology*, Corrected reprint of the 1976 original. Graduate Texts in Mathematics, 33. Springer-Verlag, New York, 1994.
- [Per] Perelman, G., Proof of the soul conjecture of Cheeger and Gromoll., J. Differential Geom. 40 (1994), 209–212.
- [Sh] Sharafutdinov, V. A., The Pogorelov-Klingenberg theorem for manifolds that are homeomorphic to R<sup>n</sup>., (Russian) Sibirsk. Mat. Ž. 18 (1977), 915–925; English translation: Siberian Math. J. 18 (1977), no. 4, 649–657 (1978).
- [Wi] Wilking, B., A duality theorem for singular Riemannian foliations in nonnegative sectional curvature, in preparation.
- [Y1] Yim, J.-W., Distance nonincreasing retraction on a complete open manifold of nonnegative sectional curvature., Ann. Global Anal. Geom. 6 (1988,), 191–206.
- [Y2] \_\_\_\_\_, Spaces of souls in a complete open manifold of nonnegative curvature, J. Diff. Geom. **32** (1990,), 429-455.

JIANGUO CAO, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556 USA.

 $E\text{-}mail \ address: \texttt{cao.7}@\texttt{nd.edu}$ 

MEI-CHI SHAW, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556 USA

*E-mail address*: mei-chi.shaw.1@nd.edu