

# Flat manifolds, harmonic spinors, and eta invariants

M. Sadowski, A. Szczepański \*

December 3, 2018

## Abstract

The aim of this paper is to calculate the eta invariants and the dimensions of the spaces of harmonic spinors for two infinite families of closed flat manifolds. The first one  $\mathcal{F}_{CHD}$  consists of some flat manifolds  $M$  with cyclic holonomy groups. The second one  $\mathcal{F}_{HW}$  is the family of generalized oriented Hantzsche-Wendt manifolds. If  $M \in \mathcal{F}_{HW}$ , and  $M$  admits a spin structure, then  $\eta(M) = 0$  and  $\mathfrak{h}(M) = 0$ . If  $M \in \mathcal{F}_{CHD}$ , then we give explicit formulas for  $\eta(M)$  and  $\mathfrak{h}(M)$ . They are expressed in terms of solutions of appropriate congruences in  $\{-1, 1\}^{\lfloor \frac{n-1}{2} \rfloor}$ . The methods used here can be generalized to various other cases.

Key words and phrases: *Spin structure, harmonic spinor, eta invariant, flat manifold.*

2000 Mathematics Subject Classification: 58J28, 53C27, 20H25

## 1 Introduction

In this paper we consider Dirac operators on two infinite families of closed flat manifolds. The first one  $\mathcal{F}_{CHD}$  consists of flat manifolds  $M$  with cyclic holonomy groups of odd order equal to the dimension of  $M$ . The second one

---

\*Supported by University of Gdańsk grant number BW - 5100-5-0319-3

$\mathcal{F}_{HW}$  consists of oriented generalized Hantzsche-Wendt manifolds. We describe the eta invariants of the Dirac operators arising from different spin structures and we give necessary and sufficient conditions of the existence of nontrivial harmonic spinors. The results allow to calculate the eta invariants and the dimension of the space of harmonic spinors using a computer. The family  $\mathcal{F}_{CHD}$  is particularly simple and the investigation of different properties of multidimensional flat manifolds should start with the investigation of them in this particular case. The methods developed here are of independent interest. They can be generalized to many other cases.

To formulate the main results we need some definitions. Let  $\mathfrak{h}(V)$  be the dimension of the vector space of harmonic spinors on a spin-manifold  $V$ . Let  $n = 2k + 1$  be an odd number, and let  $a_1, \dots, a_n$  be a basis of  $\mathbb{R}^n$ . Consider the linear map  $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $A(a_j) = a_{j+1}$  for  $j < n - 1$ ,  $A(a_{n-1}) = -a_1 - \dots - a_{n-1}$ , and  $A(a_n) = a_n$ . Let  $a = \frac{1}{n}a_n$  and let  $g(x) = A(x) + a$ . An  $n$ -dimensional flat manifold  $M \in \mathcal{F}_{CHD}$  can be written as  $\mathbb{R}^n/\Gamma$ , where  $\Gamma = \langle a_1, \dots, a_{n-1}, g \rangle$ . The linear part  $A$  of  $g$  has two lifts  $\alpha_+, \alpha_- \in Spin(n)$  such that  $\alpha_+^n = id$  and  $\alpha_-^n = -id$  (see Section 2). This defines two spin structures on  $M$ . Now we state the main results of the paper.

**Theorem 1.** *Let  $k$  be a positive integer and let  $n = 2k + 1$ . If  $M \in \mathcal{F}_{CHD}$ , then*

- a)  $\mathfrak{h}(M, \alpha_+) > 0$  if and only if  $n \geq 5$ .
- b)  $\mathfrak{h}(M, \alpha_-) = 0$ .

**Theorem 2.** *If  $k$  is odd,  $n = 2k + 1$ , and  $M^n \in \mathcal{F}_{HW}$ , then*

- a)  $M^n$  has exactly  $2^{n-1}$  spin structures,
- b)  $\mathfrak{h}(M^n) = 0$  and  $\eta_{M^n}(0) = 0$ .

It is easy to see that generalized Hantzsche-Wendt manifolds with  $k$  even do not admit spin structures (see Remark 4). To formulate the result describing  $\eta_M^n(0)$  for  $M^n \in \mathcal{F}_{CHD}$  we need some combinatorial invariants. It is known that  $\eta_{M^n}(0) = 0$  if  $k$  is even (cf. [1, p. 61]) so we consider the case when  $k$  is odd. Let

$$c(k) = \begin{cases} 0 & \text{if } \frac{k(k+1)}{2} \text{ is even} \\ \frac{1}{2} & \text{if } \frac{k(k+1)}{2} \text{ is odd} \end{cases} .$$

For every  $\epsilon = (\epsilon_1, \dots, \epsilon_k) \in \{-1, 1\}^k$  consider

$$\mu_\epsilon = \sum_{j=1}^k \epsilon_j j \quad \text{and} \quad \nu(\epsilon) = \epsilon_1 \cdots \epsilon_k.$$

Let  $\mathcal{D}_+ = \{\epsilon \in \{-1, 1\}^k : \nu(\epsilon) = \epsilon_1 \epsilon_2 \cdots \epsilon_k = 1\}$ , let  $r \in \{0, \dots, n-1\}$ , let

$$A_r = 2\#\{\epsilon \in \mathcal{D}_+ : \frac{\mu_\epsilon}{2} + c(k)n \equiv r \pmod{(n)}\}$$

in the case of  $\alpha_+$ , and let

$$A_r = 2\#\{\epsilon \in \mathcal{D}_+ : \frac{\mu_\epsilon}{2} + c(k)n + k \equiv r \pmod{(n)}\}$$

in the case of  $\alpha_-$ . The numbers  $A_r$  are well defined (cf. Remark 1).

**Theorem 3.** *Let  $k$  be an odd positive integer and let  $n = 2k + 1$ . If  $M^n \in \mathcal{F}_{CHD}$ , then*

$$(1) \quad \eta_{M^n, \alpha_+}(0) = \sum_{r=1}^{n-1} A_r \left(1 - \frac{2r}{n}\right),$$

$$(2) \quad \eta_{M^n, \alpha_-}(0) = \sum_{r=0}^{n-1} A_r \left(1 - \frac{2r+1}{n}\right).$$

If  $M^n \in \mathcal{F}_{CHD}$ , then we have a similar description of  $\mathfrak{h}(M^n, \alpha_+)$  (Proposition 2). We give some examples illustrating Theorem 3. The simplest one is Example 1 reproving a result of [12].

The spectra of the Dirac operators on flat tori were described in [6]. The spectra of the Dirac operators on closed 3-dimensional flat manifolds and their eta invariants were calculated in [12]. If the holonomy group of a closed flat spin 3-manifold  $V$  is nontrivial, then  $\mathfrak{h}(V) = 0$  and  $\eta_V(0) \neq 0$  only when the holonomy group of  $V$  is isomorphic to  $\mathbb{Z}_3$  ([12]). Related results about the eta invariants for the signature operators on closed 3-dimensional flat manifolds can be found in [10, p. 176]. For more information we refer to [2] and [7].

Throughout this paper the following notation will be used. If  $G$  is a group and  $g_1, \dots, g_l \in G$ , then  $\langle g_1, \dots, g_l \rangle$  is the subgroup of  $G$  generated by  $g_1, \dots, g_l$ .

The symbol  $X^G$  stands for the set of the fixed points of an action of  $G$  on  $X$ . For every  $g \in G$ ,  $X^g = \{x \in X : gx = x\}$ . By  $\Gamma$  (or  $\Gamma_n$ ) we denote the deck group of a closed flat manifold  $M$ , by  $h$  the holonomy homomorphism of  $M$ , and by  $\widehat{h}$  its lift to  $Spin(n)$ . The standard epimorphism from  $Spin(n)$  to  $SO(n)$  will be denoted by  $\lambda$  (cf. Section 2). The letter  $\Gamma_0$  stands for the maximal abelian subgroup of  $\Gamma$  consisting of all translations belonging to  $\Gamma$ , (cf. [4]). By  $a_1, \dots, a_n$  we usually denote a basis of  $\Gamma_0$ . The subspace of a vector space spanned by vectors  $v_1, \dots, v_l$  will be denoted by  $\text{Span}[v_1, \dots, v_l]$ . The symbols  $\alpha_+$ ,  $\alpha_-$ ,  $\mathfrak{h}(M)$ ,  $c(k)$ ,  $\mu_\epsilon$ ,  $\nu(\epsilon)$ , and  $A_r$  were defined above. The cyclic group  $\langle A \rangle$  will be denoted by  $G$ .

We would like to thank Andrzej Weber for helpful conversations.

## 2 Spin structures on $\mathcal{F}_{CHD}$ -manifolds and Dirac operators

Let  $k \in \mathbb{N} \cup \{0\}$  and let  $n = 2k + 1$ .

**Lemma 1.** *Let  $A$ ,  $a_1, \dots, a_n$  and  $\Gamma$  be as in the introduction, and let  $\langle \cdot, \cdot \rangle_0^*$  be an  $A$ -invariant scalar product in  $\mathbb{R}^n$ . Then*

- a)  $M = \mathbb{R}^n/\Gamma$  is a closed, orientable, flat manifold,
- b) the eigenvalues of  $A$  are equal to  $e^{\frac{2\pi i j}{n}}$ ,  $j = 1, \dots, n$ ,
- c)  $\text{Span}[a_1, \dots, a_{n-1}] \perp a_n$ ,
- d) there is an  $A$ -invariant scalar product  $\langle \cdot, \cdot \rangle$ , similar to  $\langle \cdot, \cdot \rangle_0^*$ , such that  $\langle a_n, a_n \rangle = 1$ .

**Proof .** a) For the sake of completeness we give a simple direct argument (cf. [4]). Assume that  $\gamma x = x$  for  $\gamma \in \Gamma$ . Then  $\gamma^n x = x$ . Since  $\Gamma_0$  acts freely,  $\gamma^n = 1$ . Hence  $\gamma = 1$ . The action of  $\Gamma$  is properly discontinuous, because the action of  $\Gamma_0$  is. The manifold  $\mathbb{R}^n/\Gamma$  is closed and orientable, because  $\mathbb{R}^n/\Gamma_0$  is closed and  $A \in SO(n)$ .

b) For every  $j = 2, \dots, n - 1$ , consider the  $(j \times j)$ -matrix

$$M_j = \begin{bmatrix} 0 & 0 & \dots & 0 & -1 \\ 1 & 0 & \dots & 0 & -1 \\ 0 & 1 & \dots & 0 & -1 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & -1 \end{bmatrix}.$$

Let  $M_j(z) = M_j - zI$ . Then

$$\det(A - zI) = (1 - z) \det M_{n-1}(z).$$

Applying the Laplace expansion with respect to the first row we have

$$\det M_j(z) = -z \det M_{j-1}(z) + (-1)^j.$$

Using this it is easy to check that  $\det M_j(z) = (-1)^j \sum_{l=0}^j z^l$ . Hence

$$\det(A - zI) = (1 - z) \det M_{n-1}(z) = -z^n + 1$$

and the eigenvectors of  $A$  are equal to  $e^{\frac{2\pi i j}{n}}$ ,  $j = 1, \dots, n$ .

c) Let  $W = \text{Span}[a_1, \dots, a_{n-1}]$  and let  $w \in W$ . It is clear that  $W$  is an  $(n-1)$ -dimensional  $A$ -invariant subspace of  $\mathbb{R}^n$  and  $\sum_{j=0}^{n-1} A^j w \in W^A = \{0\}$ . Since  $\left\langle \sum_{j=0}^{n-1} A^j w, a_n \right\rangle = \left\langle w, \sum_{j=0}^{n-1} A^{-j} a_n \right\rangle = n \langle w, a_n \rangle$  we have  $\langle w, a_n \rangle = 0$ .

d) is obvious.  $\square$

Let  $\langle, \rangle^*$  be a scalar product in  $\mathbb{R}^n$  and let  $e_1, \dots, e_n$  be an orthonormal basis in  $\mathbb{R}^n, \langle, \rangle^*$  such that:

- (i)  $e_1, \dots, e_{n-1} \in \text{Span}[a_1, \dots, a_{n-1}]$  and  $e_n = a_n$ ,
- (ii) for every for  $j \leq n-1$  :  $A(e_{2j-1}) = \cos(2\pi j/n)e_{2j-1} + \sin(2\pi j/n)e_{2j}$ , and  $A(e_{2j}) = -\sin(2\pi j/n)e_{2j-1} + \cos(2\pi j/n)e_{2j}$ .

We say that the *Riemannian metric* on  $M$  determined by  $\langle, \rangle^*$  is *special*.

**Lemma 2.** a) Let  $\langle, \rangle_0$  be a flat Riemannian metric on  $M \in \mathcal{F}_{CHD}$ . Then  $\langle, \rangle_0$  is similar to a special flat metric  $\langle, \rangle$ .

b) For each spin structure  $\sigma$  on  $M$  we have  $\eta(M, \langle, \rangle_0, \sigma) = \eta(M, \langle, \rangle, \sigma)$  and  $\mathfrak{h}(M, \langle, \rangle_0, \sigma) = \mathfrak{h}(M, \langle, \rangle, \sigma)$ .

**Proof.** The part a) is an immediate consequence of Lemma 1. The part b) follows from [1, p. 61] and [9, Proposition 1.3].  $\square$

Throughout the rest of the paper we shall always assume that a Riemannian metric on  $M \in \mathcal{F}_{CHD}$  is special and  $e_1, \dots, e_n$  is a basis satisfying (i) and (ii).

Let  $\text{Cliff}(n)$  be the Clifford algebra in  $\mathbb{R}^n$  and let  $\text{Cliff}_{\mathbb{C}}(n)$  be its complexification. The group  $\text{Spin}(n)$  is the set of products  $x_1 \cdots x_{2r}$ , where  $r \in \mathbb{N}$ ,

and where  $x_1, \dots, x_{2r}$  are the elements of the unit sphere in  $\mathbb{R}^n$ . The standard covering map  $\lambda : Spin(n) \rightarrow SO(n)$  carries  $y \in Spin(n)$  onto  $\mathbb{R}^n \ni x \rightarrow yxy^*$ , where  $(e_{j_1} \cdots e_{j_s})^* = e_{j_s} \cdots e_{j_1}$ . A spin structure on an orientable flat manifold  $V = \mathbb{R}^n/\Gamma_V$  is determined by the lift  $\widehat{h} : \Gamma_V \rightarrow Spin(n)$  of the holonomy homomorphism  $h : \Gamma_V \rightarrow SO(n)$ . Recall that  $h$  carries  $\gamma \in \Gamma_V$  onto its linear part  $h(\gamma)$ . For  $M \in \mathcal{F}_{CHD}$  we have  $h(\Gamma) = \langle A \rangle \cong \mathbb{Z}_n$  and any lift  $\widehat{A}$  of  $A$  to  $Spin(n)$  defines the lift  $\widehat{h}$  of  $h$ , given by the formulas  $\widehat{h}(a_j) = 1$  for  $j \leq n-1$ ,  $\widehat{h}(g) = \widehat{A}$ . In order to construct  $\widehat{A}$  consider  $\beta = \frac{\pi}{n}$ ,

$$r_j = \cos(j\beta) + e_{2j-1}e_{2j} \sin(j\beta),$$

and  $\alpha = \prod_{j=1}^k r_j$ . Clearly  $r_i r_j = r_j r_i$  for  $i, j \in \{1, \dots, k\}$ . A direct calculation yields

$$\lambda(r_j)(e_l) = \begin{cases} \cos(2j\beta)e_{2j-1} + \sin(2j\beta)e_{2j} & \text{for } l = 2j-1 \\ -\sin(2j\beta)e_{2j-1} + \cos(2j\beta)e_{2j} & \text{for } l = 2j \\ e_l & \text{for } l \notin \{2j-1, 2j\} \end{cases}.$$

Using this it is easy to check the following.

**Lemma 3.**  $\alpha^n = (-1)^{\frac{k(k+1)}{2}}$  and  $\lambda(\alpha) = A$ .

Now we can define

$$\alpha_+ = (-1)^{\frac{k(k+1)}{2}} \alpha, \quad \alpha_- = -(-1)^{\frac{k(k+1)}{2}} \alpha.$$

Since  $n$  is odd,

$$\alpha_+^n = 1 \quad \text{and} \quad \alpha_-^n = -1.$$

To see that  $\alpha_+$  and  $\alpha_-$  determine all spin structures on  $M$  we need the following lemma, known in the case when  $p$  is a prime number ([4, p. 153]).

**Lemma 4.** a)  $H_1(M, \mathbb{Z}) \cong \mathbb{Z} \oplus S$ , where  $S$  is a finite group of odd order.  
b)  $H^1(M, \mathbb{Z}_2) \cong \mathbb{Z}_2$ .

**Proof.** a) The group  $\Gamma_0 = \langle a_1, \dots, a_n \rangle$  is the maximal abelian subgroup of  $\Gamma$  and the following sequence

$$0 \rightarrow \Gamma_0 \rightarrow \Gamma \rightarrow \langle A \rangle \rightarrow 1$$

is exact (cf. [4, ch. 1, Proposition 4.1], [19, Theorem 3.2.9]). It is well known (cf. [8, Corollary 1.3]) that  $\text{rank } H_1(\Gamma, \mathbb{Z}) = \text{rank } \Gamma_0^A = 1$  so that  $H_1(M, \mathbb{Z}) \cong \mathbb{Z} \oplus S$ , where  $S$  is a finite group. According to [3, ch.3, § 9], there are homomorphisms  $\text{res} : H_*(M, \mathbb{Z}) \cong H_*(\Gamma, \mathbb{Z}) \rightarrow H_*(\Gamma_0, \mathbb{Z})$  and  $\text{cor} : H_*(\Gamma_0, \mathbb{Z}) \rightarrow H_*(M, \mathbb{Z})$  such that  $\text{cor} \circ \text{res}$  is the multiplication by  $n$ . Since the group  $H_*(\Gamma_0, \mathbb{Z}) \cong H_*(T^n, \mathbb{Z})$  is torsion free we have  $nS = 0$ . In particular, the order of  $S$  is odd.

**b)** By a),  $H^1(M, \mathbb{Z}_2) \cong \text{Hom}(H_1(M, \mathbb{Z}), \mathbb{Z}_2) \cong \text{Hom}(\mathbb{Z}, \mathbb{Z}_2) \cong \mathbb{Z}_2$ . This finishes the proof of Lemma 4.  $\square$

Since  $\alpha_+$ ,  $\alpha_-$  are different lifts of the holonomy homomorphism  $h$  to  $\text{Spin}(n)$ , the spin structures determined by them are different. It is known that spin structures on  $M$  correspond to the elements of  $H^1(M, \mathbb{Z}_2)$  ([7, p. 40]). Hence we have.

**Corollary 1.**  *$M$  has exactly two spin structures determined by  $\alpha_+$  and  $\alpha_-$ .*

By [7, Section 1.3], the irreducible complex  $\text{Cliff}_{\mathbb{C}}(n)$ -module  $\Sigma_{2k}$  can be described as follows. Let

$$g_1 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \quad g_2 = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, \quad T = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}.$$

Let  $\Sigma_{2k} = \underbrace{\mathbb{C}^2 \otimes \dots \otimes \mathbb{C}^2}_{k \text{ times}}$  and let  $\alpha(j) = \begin{cases} 1 & \text{if } j \text{ is odd} \\ 2 & \text{if } j \text{ is even} \end{cases}$ . Take an element  $u = u_1 \otimes \dots \otimes u_k$  of  $\Sigma_{2k}$  and the orthonormal basis  $e_1, \dots, e_n$  considered above. Then

$$e_j u = (I \otimes \dots \otimes I \otimes g_{\alpha(j)} \otimes \underbrace{T \otimes \dots \otimes T}_{[\frac{j-1}{2}] \text{ times}})(u),$$

for  $j \leq n-1$ , and

$$e_n u = i(T \otimes \dots \otimes T)u.$$

A spin structure on  $M$  determines a complex spinor bundle  $P\Sigma_{2k}$  with fiber  $\Sigma_{2k}$ . This bundle is the orbit space of  $\mathbf{R}^n \times \Sigma_{2k}$  by the action of  $\Gamma$  given by

$$\gamma(x, v) = (\gamma x, \widehat{h}(\gamma)v),$$

where  $x \in \mathbf{R}^n$  and  $v \in \Sigma_{2k}$ . Clearly

$$\widehat{h}(a_j) = 1 \text{ for } j \leq n-1$$

and

$$\widehat{h}(g) = \pm\alpha.$$

Since  $\text{Span}[e_1, \dots, e_{n-1}] = \text{Span}[a_1, \dots, a_{n-1}]$  and  $a_n = e_n$  we have

$$\widehat{h}(e_j) = 1 \text{ for } j \leq n-1.$$

Consider the covering  $T^n = \mathbb{R}^n/\Gamma_0 \rightarrow M$ , where  $\Gamma_0 = \langle a_1, \dots, a_n \rangle$ . As  $n$  is odd we have

$$\widehat{h}(a_n) = \pm\alpha^n.$$

The lift  $P_T\Sigma_{2k}$  of  $P\Sigma_{2k}$  to  $T^n$  is the orbit space  $(\mathbb{R}^n \times \Sigma_{2k})/\Gamma_0$ , where the action of  $\Gamma_0$  on  $\mathbb{R}^n \times \Sigma_{2k}$  is given by the formula

$$\gamma(x, v) = (\gamma(x), v)$$

in the case of  $\alpha_+$  and

$$a_j(x, v) = (a_j(x), v) \text{ for } j \leq n-1, \quad a_n(x, v) = (a_n(x), -v),$$

in the case of  $\alpha_-$ .

To deal with the spectrum of the Dirac operator  $D$  it is convenient to describe it in terms of the spectrum of  $D^2$ . We state without proofs some related results of [12] that will be used later. Identify the parallel section  $\mathbb{R}^n \ni x \rightarrow (x, v) \in \mathbb{R}^n \times \Sigma_{2k}$  with  $v$ . Every section of the trivial bundle  $\mathbb{R}^n \times \Sigma_{2k}$  (covering our bundle  $P\Sigma_{2k}$ ) can be written as  $fv$  for some  $f \in C^\infty(\mathbb{R}^n)$  and for some parallel section  $v$ . Take the coordinate system  $x_1, \dots, x_n$  determined by  $e_1, \dots, e_n$ . Since  $v$  is parallel,

$$D(fv) = \sum_j e_j \nabla_{e_j}(fv) = \sum_j e_j \left( \frac{\partial}{\partial x_j}(f)v + f \nabla_{e_j} v \right) = \sum_j e_j \frac{\partial f}{\partial x_j} v.$$

Let  $\Gamma_0^* = \{b \in \mathbb{R}^n : \langle b, \gamma \rangle \in \mathbb{Z} \text{ for } \gamma \in \Gamma_0\}$ . Let  $\mathcal{B}$  be  $\Gamma_0^*$  in the case of  $\alpha_+$  and  $\Gamma_0^* + \frac{1}{2}e_n$  in the case of  $\alpha_-$ . The action of  $g$  on the set of sections of  $\mathbb{R}^n \times \Sigma_{2k}$ , induced by the action of  $g$  on  $\mathbb{R}^n$ , is given by the formula

$$g(s)(x) = \widehat{h}(A)s(g^{-1}x).$$

**Lemma 5** (cf. [6], [12, section 4]) *Let  $f \in C^\infty(\mathbb{R}^n)$ , let  $b \in \mathbb{R}^n$ , and let  $f_b(x) = e^{2\pi i \langle b, x \rangle}$ . Then*

- a)  $D^2(fv) = -\sum_{j=1}^n \frac{\partial^2 f}{\partial x_j^2} v$ ,
- b)  $D^2(f_b v) = 4\pi^2 \|b\|^2 f_b v$ ,
- c)  $f_b v$  determines a section over  $T^n$  if and only if  $b \in \mathcal{B}$ ,
- d) the sections  $f_b v, b \in \mathcal{B}, v \in \Sigma_{2k}$ , correspond to eigenvectors of  $D$  on  $T^n$ ,
- e) the elements of  $\{f_b v : v \in \Sigma_{2k}, b \in \mathcal{B}\}^g$  correspond to eigenvectors of  $D$  on  $M$ .

As in [12] we decompose  $D$  onto a symmetric part  $D_S$  and possibly asymmetric part  $D_{Pas}$ . To achieve this goal consider  $b \in \mathcal{B}$  and  $E_b(D^2) = \{f_b v : v \in \Sigma_{2k}\}$ . For  $b \in \mathcal{B} - \{0\}$  take the decomposition  $E_b(D^2) = E_{b+} \oplus E_{b-}$ , where

$$E_{b\pm} = \{s \in E_b(D^2) : Ds = \pm 2\pi \|b\| s\}$$

([12, p. 9]). Then  $\dim E_{b+} = \dim E_{b-} = 2^{k-1}$  ([12, p. 9]). It is easy to see that

$$E_{b\pm} = \{f_b v : v \in \Sigma_{\pm}\},$$

where  $\Sigma_{\pm} = \text{Span}[v_{\epsilon}]$ . Since  $g^{-1}x = A^{-1}(x) - a$  and  $A \in SO(n)$  we have

$$(f_b \circ g^{-1})(x) = e^{-2\pi i \langle A(b), a \rangle} f_{A(b)}(x)$$

so that  $AE_b \subseteq E_{A(b)}$ . Let  $\mathcal{B}_{Sym} = \{b \in \mathcal{B} : \#G(b) = \#G\}$ , let  $\mathcal{B}_{Pas} = \{b \in \mathcal{B} : \#G(b) < \#G\}$ , and let

$$D_S = D|_{[\oplus_{b \in \mathcal{B}_{Sym}} E_b(D^2)]^g}, \quad D_{Pas} = D|_{[\oplus_{b \in \mathcal{B}_{Pas}} E_b(D^2)]^g}.$$

Clearly  $\mathcal{B}$  is the disjoint union of  $\mathcal{B}_{Sym}$  and  $\mathcal{B}_{Pas}$  and the Dirac operator  $D$  on  $M$  can be identified with  $D_S \oplus D_{Pas}$ . In the proof of Theorem 3 we use Theorem 4.2 from [12]. In a particular case, for  $M \in \mathcal{F}_{CHD}$  it can be stated as follows.

**Lemma 6** ([12, Theorem 4.2]) **a)** *Let  $b \in \mathcal{B}_{Sym}$  and let  $V_b^{\pm} = \bigoplus_{h \in G} E_{h(b\pm)}(D^2)$ .*

*Then  $\dim(V_b^{\pm})^g = \dim E_b^{\pm} = 2^{k-1}$ .*

**b)** *The spectrum of  $D_S$  is symmetric.*

The statement b) means that if  $\lambda$  is an eigenvalue of  $D_S$ , then  $-\lambda$  is an eigenvalue of  $D_S$  of the same multiplicity.

### 3 Eta invariants of $\mathcal{F}_{CHD}$ -manifolds

The aim of this section is to prove Theorem 3. Recall that the  $\eta$ -invariant of the Dirac operator on a closed spin manifold  $M$  is defined as follows. As  $D$  is elliptic formally self adjoint, it has discrete real spectrum and the series  $\sum_{\lambda \neq 0} \text{sgn } \lambda |\lambda|^{-z}$  converges for  $z \in \mathbb{C}$  with  $\text{Re}(z)$  sufficiently large ([1, Theorem 3.10]). Here summation is taken over all nonzero eigenvalues  $\lambda$  of  $D$ , each eigenvalue being repeated according to its multiplicity. The function  $z \rightarrow \sum_{\lambda \neq 0} \text{sgn } \lambda |\lambda|^{-z}$  can be extended to a meromorphic function  $\eta_M$  in the whole complex plane such that 0 is a regular point of  $\eta_M$  ([1, Theorem 3.10]). The *eta-invariant* of  $M$  is  $\eta_M(0)$ .

Define an endomorphism  $\rho_1$  of  $\mathbb{C}^2$  by the formula

$$\rho_1(u) = \cos \beta u + \sin \beta g_1 g_2 u.$$

The matrix of  $\rho_1$  is equal to

$$\cos \beta I + \sin \beta \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix}$$

so that the matrix of  $\rho_1^j$  is equal to

$$\cos(\beta j) I + \sin(\beta j) \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

The following lemma is crucial.

**Lemma 7.** *Let  $w_{+1} = (1, -i)$ , let  $w_{-1} = (1, i)$ , let  $\epsilon = (\epsilon_1, \dots, \epsilon_k) \in \{-1, 1\}^k$ , and let  $v_\epsilon = w_{\epsilon_1} \otimes \dots \otimes w_{\epsilon_k}$ . Let  $\beta = \frac{\pi}{n}$  and let  $\mu_\epsilon$  be as in Theorem 3. Take  $u = u_1 \otimes \dots \otimes u_k \in \Sigma_{2k}$ . Then*

- a)  $\alpha u = \rho_1 u_1 \otimes \dots \otimes \rho_1^k u_k$ ,
- b)  $\alpha e_n u = e_n \alpha u$ ,
- c)  $\rho_1(w_{\pm 1}) = e^{\pm i\beta} w_{\pm 1}$ ,
- d)  $\alpha v_\epsilon = e^{i\beta \mu_\epsilon} v_\epsilon$  and  $\{v_\epsilon : \epsilon \in \{-1, 1\}^k\}$  is a basis of  $\Sigma_{2k}$ ,
- e)  $e_n v_\epsilon = -i\nu(\epsilon)v_\epsilon$ .

**Proof.** a) Since  $T^2 = id$ ,

$$e_{2j-1} e_{2j} (u_1 \otimes \dots \otimes u_j \otimes \dots \otimes u_k) = u_1 \otimes \dots \otimes g_1 g_2 (u_j) \otimes \dots \otimes u_k$$

and consequently

$$r_j(u_1 \otimes \dots \otimes u_j \otimes \dots \otimes u_k) = u_1 \otimes \dots \otimes \rho_1^j(u_j) \otimes \dots \otimes u_k.$$

Hence

$$\alpha(u_1 \otimes \dots \otimes u_k) = (r_1 \cdots r_k)(u_1 \otimes \dots \otimes u_k) = \rho_1(u_1) \otimes \dots \otimes \rho_1^k(u_k).$$

b) For  $j \leq k$  we have  $e_{2j-1}e_{2j}e_n = e_n e_{2j-1}e_{2j}$  so that  $r_j e_n = e_n r_j$ .

c) is obvious.

d) By c),  $r_1(w_{\epsilon_j}) = e^{i\beta\epsilon_j} w_{\epsilon_j}$ . Hence

$$\alpha(v_\epsilon) = r_1(w_{\epsilon_1}) \otimes \dots \otimes r_1^k(w_{\epsilon_k}) = e^{i\beta\mu_\epsilon} v_\epsilon.$$

Since  $\#\{v_\epsilon : \epsilon \in \{-1, 1\}^k\} = 2^k = \dim \Sigma_{2k}$  and the vectors  $v_\epsilon$  are linearly independent, they form a basis of  $\Sigma_{2k}$ .

e) We have  $T(w_1) = -w_1$  and  $T(w_{-1}) = w_{-1}$ . It follows that

$$e_n v_\epsilon = iTw_{\epsilon_1} \otimes \dots \otimes Tw_{\epsilon_k} = i(-1)^{\#\{j \in \{1, \dots, k\} : \epsilon_j = 1\}} v_\epsilon = -i\nu(\epsilon)v_\epsilon.$$

This finishes the proof of Lemma 7.  $\square$

**Lemma 8.** *Let  $\mathcal{E}(\lambda, D_{Pas})$  be the eigenspace of  $\lambda$  for  $D_{Pas}$  on  $M$ . Then:*

a)  $\lambda = 2\pi l$  for some  $l \in \mathbb{Z}$ ,

b)  $D(f_{le_n} v_\epsilon) = \nu(\epsilon)2\pi l f_{le_n} v_\epsilon$ ,

c)  $\mathcal{E}(2\pi l, D_{Pas}) = \text{Span}[f_{\nu(\epsilon)le_n} v_\epsilon : \epsilon \in \{-1, 1\}^k]^g$ .

**Proof.** a) By Lemma 5 e), the eigenvectors of  $\lambda$ , can be written as  $f_b v$  for some  $b \in \mathcal{B}_{Pas}$  and  $v \in \Sigma_{2k}$ . The condition  $\#G(b) < \#G$  means that  $A^s(b) = b$  for  $s < n$  and this implies that  $b \in \text{Span}[e_n]$  so that  $b = l_0 e_n$  for some  $l_0 \in \mathbb{Z}$ . According to Lemma 5 b),  $\lambda = \pm 2\pi l_0$ .

b) By Lemma 7 e) we have

$$D(f_{le_n} v_\epsilon) = -\frac{\partial}{\partial x_n}(e^{2\pi i \langle le_n, x \rangle}) i\nu(\epsilon)v_\epsilon = \nu(\epsilon)2\pi l v_\epsilon.$$

c) If  $f_b v \in \mathcal{E}(2\pi l, D_{Pas})$ , then  $v$  is an eigenvector of  $e_n$  so that  $v = v_\epsilon$  for some  $\epsilon \in \{-1, 1\}^k$ . By a) and b),  $f_b v = f_{\nu(\epsilon)le_n} v_\epsilon$ . Clearly  $f_b v$  is  $g$ -invariant.  $\square$

Now we are able to describe the spectrum of  $D_{Pas}$ .

**Proposition 1.** *Let  $n, k, M, \mu_\epsilon, \nu(\epsilon)$ , and  $c(k)$  be as in Theorem 3.*

**a)** *If the spin structure is given by  $\alpha_+$ , then*

- (i)  $gf_{le_n}v_\epsilon = f_{le_n}v_\epsilon \Leftrightarrow l \equiv \frac{\mu_\epsilon}{2} + c(k)n \pmod{n}$ ,
- (ii)  $\mathcal{E}(2\pi l, D_{Pas}) = \text{Span}[f_{\nu(\epsilon)le_n}v_\epsilon : \frac{\mu_\epsilon}{2} + c(k)n \equiv \nu(\epsilon)l \pmod{n}]$ ,

**b)** *If the spin structure is given by  $\alpha_-$ , then*

- (i)  $gf_{(l+\frac{1}{2})e_n}v_\epsilon = f_{(l+\frac{1}{2})e_n}v_\epsilon \Leftrightarrow l \equiv \frac{\mu_\epsilon}{2} + c(k)n + \frac{n-1}{2} \pmod{n}$ ,
- (ii)  $\mathcal{E}(2\pi l, D_{Pas}) = \text{Span}[f_{\nu(\epsilon)(l+\frac{1}{2})e_n}v_\epsilon : \frac{\mu_\epsilon}{2} + c(k)n + \frac{n-1}{2} \equiv \nu(\epsilon)l \pmod{n}]$ .

**Remark 1.** Since  $\epsilon_j - 1$  are even, the difference  $\mu_\epsilon - k(k+1)/2 = \sum_{j=1}^k \epsilon_j j - \sum_{j=1}^k j$  is divisible by 2. Using this and the definition of  $c(k)$  it is easy to see that  $\mu_\epsilon/2 + c(k)n$  is an integer.

**Proof of Proposition 1. a)** Recall that  $a = \frac{1}{n}e_n$ ,  $g(x) = A(x) + a$ , and  $G = \langle A \rangle$ . From the definitions of  $\alpha_+$  and  $c(k)$  it follows that  $\alpha_+ = (-1)^{2c(k)}\alpha$ . Since  $A(a) = a$  and  $AA^* = id$  we have

$$\begin{aligned} \langle le_n, g^{-1}x \rangle &= \langle le_n, A^{-1}x - a \rangle = \langle A(le_n), x - a \rangle \\ &= \langle le_n, x \rangle - \langle le_n, \frac{1}{n}e_n \rangle = \langle le_n, x \rangle - \frac{1}{n}. \end{aligned}$$

Hence

$$gf_{le_n}(x) = (f_{le_n} \circ g^{-1})(x) = e^{2\pi i \langle le_n, g^{-1}x \rangle} = e^{-\frac{2\pi i l}{n}} f_{le_n}(x)$$

and consequently

$$\begin{aligned} g(f_{le_n}(x)v_\epsilon) &= f_{le_n}(g^{-1}x)\alpha_+v_\epsilon = e^{-\frac{2\pi i l}{n}} f_{le_n}(x)(-1)^{2c(k)}\alpha v_\epsilon \\ &= e^{-\frac{2\pi i}{n}(l - \frac{\mu_\epsilon}{2} - c(k)n)} f_{le_n}(x)v_\epsilon. \end{aligned}$$

The condition  $gf_{le_n}v_\epsilon = f_{le_n}v_\epsilon$  is satisfied if and only if  $l \in n\mathbb{Z} + \frac{\mu_\epsilon}{2} + c(k)n$ . The section  $f_{\nu(\epsilon)le_n}v_\epsilon$  is  $g$ -equivariant if and only if

$$\nu(\epsilon)l \equiv \frac{\mu_\epsilon}{2} + c(k)n \pmod{n}.$$

**b)** In the case of  $\alpha_-$  we have  $\mathcal{B} = \Gamma_0^* + \frac{1}{2}e_n$  so that the eigenvectors of  $D_{Pas}$  on  $T^n$  can be written as  $f_{(l+\frac{1}{2})e_n}v$  for  $l \in \mathbb{Z}, v \in \Sigma_{2k}$ . We have

$$\langle (l + \frac{1}{2})e_n, g^{-1}x \rangle = \langle (l + \frac{1}{2})e_n, x \rangle - \frac{2l+1}{2n}.$$

Reasoning as in a) we get

$$gf_{(l+\frac{1}{2})e_n}v_\epsilon = e^{-\frac{2\pi i}{n}(l-c(k)n-\frac{n-1}{2}-\frac{\mu_\epsilon}{2})}f_{(l+\frac{1}{2})e_n}v_\epsilon.$$

Hence  $f_{(l+\frac{1}{2})e_n}v_\epsilon$  is  $g$ -equivariant if and only if  $l \in n\mathbb{Z} + c(k)n + \frac{n-1}{2} + \frac{\mu_\epsilon}{2}$ . The rest of the argument is the same as in a). This finishes the proof of Proposition 1.  $\square$

**Lemma 9.** *Let  $M$  be an  $n$ -dimensional flat manifold belonging to  $\mathcal{F}_{CHD}$  with  $k = \lfloor \frac{n-1}{2} \rfloor$  odd and with a fixed spin structure. Assume that  $f_b v_\epsilon \in \mathcal{E}(\lambda, D_{Pas})$ . Then*

- a)  $f_{-b}v_{-\epsilon} \in \mathcal{E}(\lambda, D_{Pas})$ ,
- b)  $\dim \mathcal{E}(2\pi(mn+r), D_{Pas}) = A_r$ ,

**Proof.** a) If the spin structure on  $M$  is  $\alpha_+$ , then  $b = le_n$  for some  $l \in \mathbb{Z}$ , and, from the equivariance of  $f_b v_\epsilon$ , it follows that

$$l \equiv \frac{\mu_\epsilon}{2} + c(k)n \pmod{n}.$$

Hence

$$-l \equiv \frac{\mu_{-\epsilon}}{2} + c(k)n \pmod{n}$$

and  $f_{-b}v_{-\epsilon}$  is  $g$ -equivariant. By the assumption that  $k$  is odd,  $\nu(-\epsilon) = -\nu(\epsilon)$ . According to Lemma 8,  $f_{-b}v_{-\epsilon} \in \mathcal{E}(\lambda, D_{Pas})$ .

If the spin structure is  $\alpha_-$ , then we use a similar argument. We write  $b$  as  $(l + \frac{1}{2})e_n$  and we use the congruence

$$l \equiv \frac{\mu_\epsilon}{2} + c(k)n + k \pmod{n}.$$

Since  $\mu_{-\epsilon} = -\mu_\epsilon$ ,  $c(k)n \equiv -c(k)n \pmod{n}$ , and  $-k - 1 \equiv k \pmod{n}$  we have

$$-l - 1 \equiv \frac{\mu_{-\epsilon}}{2} + c(k)n + k \pmod{n}$$

and consequently  $f_{-b}v_{-\epsilon}$  is  $g$ -equivariant.

b) If the spin structure is  $\alpha_+$ , then using Proposition 1 and a), we get

$$\begin{aligned} \dim \mathcal{E}(2\pi(mn+r), D_{Pas}) &= \dim \mathcal{E}(2\pi(r), D_{Pas}) \\ &= 2\#\{\epsilon \in \mathcal{D}_+ : \frac{\mu_\epsilon}{2} + c(k)n \equiv r\} = A_r. \end{aligned}$$

A similar argument can be applied to  $\alpha_-$ .  $\square$

In order to prove Theorem 3 we need Lemma 5.5 from [12]. It can be stated as follows.

**Lemma 10.** *Let  $d \in (0, 1)$ , and let  $\sigma, C \in (0, \infty)$ . Consider the function*

$$\{z \in \mathbb{C} : \operatorname{Re}(z) > x_0\} \ni z \rightarrow \eta_{d,C,\sigma}(z) = \sum_{m=-\infty}^{\infty} \frac{C \operatorname{sgn}(\sigma(m+d))}{|\sigma(m+d)|^z} \in \mathbb{R}.$$

*Then  $\eta_{d,C,\sigma}(z)$  has a meromorphic extension to  $\mathbb{C}$ , 0 is a regular point of  $\eta_{d,C,\sigma}(z)$ , and*

$$\eta_{d,C,\sigma}(0) = C(1 - 2d).$$

**Proof of Theorem 3.** a) Let  $\mathcal{S}_{Pas}$  be the set of eigenvalues of  $D_{Pas}$ , let  $r \in \{0, \dots, n-1\}$ ,  $m \in \mathbb{Z}$ , and let  $\mathcal{S}_r = \{2\pi(mn+r) : m \in \mathbb{Z}\}$ . It is clear that  $\mathcal{S}_r$  are disjoint and  $\mathcal{S}_{Pas} \subseteq \cup_{r=0}^{n-1} \mathcal{S}_r$ . By Lemma 6,  $D_S$  has symmetric spectrum so that for  $\operatorname{Re}(z)$  sufficiently large

$$\eta_M(z) = \sum_{\lambda \in \mathcal{S}_{Pas}} \frac{\operatorname{sgn}(\lambda)}{|\lambda|^z} \dim \mathcal{E}(\lambda, D_{Pas}).$$

By Lemma 9 b),  $\dim \mathcal{E}(\lambda, D_{Pas}) = A_r$  for  $\lambda \in \mathcal{S}_r$ . If  $A_0 \neq 0$ , then the eigenvalue  $\lambda \in \mathcal{S}_0$  occur together with  $-\lambda$  with the same multiplicity  $A_0$  so that  $\sum_{\lambda \in \mathcal{S}_0 - \{0\}} \frac{A_0 \operatorname{sgn}(\lambda)}{|\lambda|^z} = 0$  and, for  $\operatorname{Re}(z)$  sufficiently big,

$$\eta_M(z) = \sum_{r=1}^{n-1} \sum_{m=-\infty}^{\infty} \frac{A_r \operatorname{sgn}(2\pi(mn+r))}{|2\pi(mn+r)|^z} = \sum_{r=1}^{n-1} \sum_{m=-\infty}^{\infty} \frac{A_r \operatorname{sgn}(2\pi n(m + \frac{r}{n}))}{|2\pi n(m + \frac{r}{n})|^z}.$$

By Lemma 10,

$$\eta_M(0) = \sum_{r=1}^{n-1} A_r \left(1 - \frac{2r}{n}\right).$$

b) We use similar arguments as those given in the proof of a). Now the component  $\mathcal{S}_0$  is not symmetric so that we do not remove  $r=0$  from the formula describing  $\eta_M(z)$ . The equality

$$2\pi \left(mn + r + \frac{1}{2}\right) = 2\pi n \left(m + \frac{2r+1}{2n}\right)$$

and Lemma 10 implies that

$$\eta_M(0) = \sum_{r=0}^{n-1} A_r \left(1 - \frac{2r+1}{n}\right).$$

This finishes the proof of Theorem 3.  $\square$

**Example 1.** The eta invariants of  $(M^3, -\alpha)$  and  $(M^3, \alpha)$  were calculated in [12]. We explain how to obtain this result using Theorem 3. Consider the first case. We have  $\epsilon = 1$  or  $\epsilon = -1$ . Only  $\epsilon = 1$  satisfies  $\nu(\epsilon) = 1$ . Clearly  $c(1) = \frac{1}{2}$  and  $\mu_1 = 1$ . The equation

$$r \equiv \frac{\mu_1}{2} + 3c(1) \pmod{3}$$

has only one solution  $r = 2$  belonging to  $\{1, 2\}$ . Since

$$\#\left\{\epsilon \in \mathcal{D}_+ : \frac{\mu_\epsilon}{2} + 3c(1) \equiv 2 \pmod{3}\right\} = 1$$

we have  $A_2 = 2$ . By Theorem 3,

$$\eta_M(0) = 2 \left(1 - \frac{4}{3}\right) = -\frac{2}{3}.$$

In the case of the second structure from the equation

$$r \equiv \frac{1}{2} + \frac{3}{2} + 1 \pmod{3}$$

it follows that  $r = 0$ . Since  $A_0 = 2$  we have

$$\eta_M(0) = 2 \left(1 - \frac{2 \cdot 0 + 1}{3}\right) = \frac{4}{3}.$$

**Remark 2.** Example 1 shows that the difference between the eta invariants arising from different spin structures on  $M^3 \in \mathcal{F}_{CHD}$  is an integer. This is a particular case of a general result of M. Dahl ([5, Theorem 3.1]).

**Example 2.** We calculate the eta invariant of the 7-dimensional spin manifold  $M, \alpha_+$  such that  $M \in \mathcal{F}_{CHD}$ . Since  $k(k+1)/2 = 6$  is even our equation is

$$r \equiv \frac{\mu_\epsilon}{2} \pmod{7}.$$

The values of  $\frac{\mu_\epsilon}{2}$  and  $r$  for  $\epsilon \in \mathcal{D}_+$  are given in the following table.

$\epsilon$	$\frac{\mu_\epsilon}{2}$	$r$
(1, 1, 1)	3	3
(1, -1, -1)	-2	5
(-1, 1, -1)	-1	6
(-1, -1, 1)	0	0

It follows that  $A_j = 2$ , for  $j = 0, 3, 5, 6$  and  $A_j = 0$  for other values of  $j$ . By Theorem 3,

$$\eta_M(0) = 2 \left[ \left(1 - \frac{6}{7}\right) + \left(1 - \frac{10}{7}\right) + \left(1 - \frac{12}{7}\right) \right] = -2.$$

## 4 Harmonic spinors on $\mathcal{F}_{CHD}$ -manifolds

A *harmonic spinor* on a closed spin manifold  $M$  is an element of the kernel of the Dirac operator on  $M$ .

**Proof of Theorem 1.** a) We have

$$gv_\epsilon = (-1)^{\frac{k(k+1)}{2}} \alpha v_\epsilon = (-1)^{\frac{k(k+1)}{2}} e^{\frac{2\pi i}{2n} \mu_\epsilon} v_\epsilon.$$

First consider the case when  $k(k+1)/2$  is even. Then  $gv_\epsilon = v_\epsilon$  if and only if

$$\mu_\epsilon \equiv 0 \pmod{2n}$$

and  $k = 4k_0 + 3$  or  $k = 4k_0$ . Let  $\delta_4$  denote the sequence  $1, -1, -1, 1$ . If  $k = 4k_0 + 3$  and

$$\epsilon = (-1, -1, 1, \underbrace{\delta_4, \dots, \delta_4}_{k_0 \text{ times}}),$$

then  $\epsilon$  belongs to  $\{-1, 1\}^k$  and  $\mu_\epsilon = 0$ . If  $k = 4k_0$  and

$$\epsilon = (\underbrace{\delta_4, \dots, \delta_4}_{k_0 \text{ times}}),$$

then  $\epsilon$  belongs to  $\{-1, 1\}^k$  and  $\mu_\epsilon = 0$ . In particular,  $\mathfrak{h}(M) > 0$ .

Now assume that  $k > 2$  and  $k(k+1)/2$  is odd. Then  $gv_\epsilon = v_\epsilon$  if and only if

$$\mu_\epsilon \equiv n \pmod{2n}$$

and  $k = 4k_0 + 1$  or  $k = 4k_0 + 2$ . If  $k = 4k_0 + 1$  consider

$$\epsilon = (1, -1, 1, \underbrace{\delta_4, \dots, \delta_4}_{k_0-1 \text{ times}}, 1, 1)$$

and if  $k = 4k_0 + 2$  consider

$$\epsilon = (-1, 1, -1, 1, \underbrace{\delta_4, \dots, \delta_4}_{k_0-1 \text{ times}}, 1, 1).$$

In both cases  $\epsilon \in \{-1, 1\}^k$  and  $\mu_\epsilon = n$ . It is easily seen that the equation  $\mu_\epsilon \equiv 0 \pmod{(2n)}$  have no solutions for  $k = 1$  or  $2$ .

b) Since  $g^n = -id$ , the equation  $gv = v$  have only one solution  $v = 0$ .  $\square$

It is easy to see that the equality  $\alpha_+v_\epsilon = v_\epsilon$  implies  $\alpha_+v_{-\epsilon} = v_{-\epsilon}$ . Using this and the arguments given in the proof of Theorem 1 we have.

**Proposition 2.** *If  $M \in \mathcal{F}_{CHD}$  and  $\dim M = 2k + 1$ , then*

$$\mathfrak{h}(M, \alpha_+) = 2\#\{\epsilon \in \mathcal{D}_+ : \frac{\mu_\epsilon}{2} + c(k)n \equiv 0 \pmod{(2k + 1)}\}.$$

*In particular,  $\mathfrak{h}(M, \alpha_+)$  is even.*

**Remark 3.** Let  $X$  be a manifold such that  $\partial X = M$  and let  $D_X$  be the operator on  $X$ , defined in [1], with the boundary condition introduced in [1]. By [1, Theorem 4.2],  $\text{index} D_X = \frac{\mathfrak{h}(M) + \eta_M(0)}{2}$ .

## 5 Harmonic spinors and eta invariants for Hantzsche-Wendt manifolds

We start with the definitions. Let  $n = 2k + 1$  be an odd number and let  $a_1, \dots, a_n$  be a basis in  $\mathbb{R}^n$ . Let  $A_i, i = 1, \dots, n - 1$ , be the linear maps such that

$$A_i(a_j) = \begin{cases} -a_j & \text{for } i \neq j \\ a_j & \text{for } i = j \end{cases}.$$

For  $i = 1, \dots, n - 1$  the formula

$$\gamma_i(x) = A_i(x) + \frac{1}{2}(a_i + a_{i+1})$$

defines an isometry of  $\mathbb{R}^n$ . It was shown in [15] that the group  $\Gamma_n = \langle \gamma_1, \dots, \gamma_n \rangle$  acts freely and properly discontinuously on  $\mathbb{R}^n$ , the orbit space  $M^n = \mathbb{R}^n / \Gamma_n$  is a closed orientable manifold, and the holonomy group of  $M^n$  is isomorphic to  $\mathbb{Z}_2^{n-1}$ . We say that  $M^n$  is the *generalized orientable Hantzsche-Wendt manifold*. More general classes of Hantzsche-Wendt manifolds were studied in [16]. These manifolds have several interesting properties (see e.g. [11], [14], [15]). One of them is that they are rational homology spheres. In order to prove Theorem 2 we describe flat Riemannian metrics on  $\mathbb{R}^n / \Gamma_n$ .

**Lemma 11.** a) A scalar product  $\langle, \rangle$  in  $\mathbb{R}^n$  is  $\Gamma_n$ -invariant if and only if  $\langle a_i, a_j \rangle = 0$  for  $i \neq j$ .  
b) Let  $e_i = \|a_i\|^{-1}a_i$  and let  $\gamma'_i(x) = A_i(x) + \frac{1}{2}(e_i + e_{i+1})$ . Let  $\Gamma_e = \langle \gamma'_1, \dots, \gamma'_{n-1} \rangle$  and let  $h : \Gamma \rightarrow SO(n)$ ,  $h_e : \Gamma \rightarrow SO(n)$  be the holonomy homomorphisms of  $\mathbb{R}^n/\Gamma_n$  and  $\mathbb{R}^n/\Gamma_e$ . Then there is an isomorphism  $\Phi : \Gamma \rightarrow \Gamma_e$  such that  $\lambda = \lambda_e \circ \Phi$ .

**Proof.** a)  $\Leftarrow$  Let  $\langle, \rangle$  be a  $\Gamma_n$ -invariant scalar product in  $\mathbb{R}^n$ . If  $i$  belongs to  $\{1, \dots, n-1\}$  and  $j \neq i$ , then

$$\langle a_i, -a_j \rangle = \langle A_i(a_i), A_i(a_j) \rangle = \langle a_i, a_j \rangle$$

so that  $\langle a_i, a_j \rangle = 0$ .

$\Leftarrow$  Assume that  $\langle a_i, a_j \rangle = 0$  for  $i \neq j$ . Let  $x = \sum_{k=1}^n x_k a_k$ . Then

$$\langle A_i(x), A_i(x) \rangle = \sum_{k,l=1}^n x_k x_l \langle A_i(a_k), A_i(a_l) \rangle = \sum_{k=1}^n x_k^2 \langle a_k, a_k \rangle = \langle x, x \rangle.$$

b) The isometries  $\gamma_i$  and  $\gamma'_i$  have the same linear part  $A_i$ . Using this it is easy to see that the formula  $\Phi(\gamma_i) = \gamma'_i$  defines an isomorphism such that  $\lambda = \lambda_e \circ \Phi$ . This completes the proof of Lemma 11.  $\square$

**Lemma 12.**  $H^1(\Gamma_n, \mathbb{Z}_2) \cong \mathbb{Z}_2^{n-1}$ .

**Proof.** For  $n \geq 5$  we have. As  $A_j$  and  $A_l$  commute,  $[\gamma_j, \gamma_l]$  is a translation so that  $[\Gamma_n, \Gamma_n] \subseteq \Gamma_0$  and consequently there is an epimorphism of  $\Gamma_n$  onto  $\Gamma_n/\Gamma_0 \cong \mathbb{Z}_2^{n-1}$ . Using the definition of  $\Gamma_n$  one can check that  $\Gamma_0 \subset [\Gamma_n, \Gamma_n]$ . Hence  $H^1(\Gamma_n, \mathbb{Z}) \cong H^1(\Gamma_n, \mathbb{Z}_2) \cong (\mathbb{Z}_2)^{n-1}$ . By [19, ch. 3, Corollary 3.5.10],  $H_1(\Gamma_3, \mathbb{Z}) \cong \mathbb{Z}_4 \oplus \mathbb{Z}_4$ . As above we obtain our result.  $\square$

**Proof of Theorem 2.** The spin structures on  $M^n$  correspond to homomorphisms  $\widehat{h} : \Gamma_n = \pi_1(M^n) \rightarrow Spin(n)$  such that  $h = \lambda \circ \widehat{h}$ . Let  $e_i = \|a_i\|^{-1}a_i$  be the orthonormal basis from Lemma 11. As in [13] and [16] consider the extra special 2-group  $\mathcal{G}_n \subset Spin(n)$ , generated by  $-1$  and the products of even number of copies of  $e_1, \dots, e_n$ . We show that there is an epimorphism  $\widehat{h}_e : \Gamma_e \rightarrow \mathcal{G}_n$  such that  $\widehat{h}_e(\gamma'_i) \in \lambda^{-1}(A_i)$ ,  $i = 1, \dots, n-1$ . For every  $j = 1, \dots, n-1$ , take  $E_j = e_1 \cdots e_{j-1} e_{j+1} \cdots e_n$ . A direct calculation yields

$$E_j E_l = -E_l E_j \text{ for } l \neq j \quad \text{and} \quad \lambda(E_j) = A_j.$$

Since  $k$  is odd, all  $E_j^2$  are equal to  $-1$ . The group  $\mathcal{G}_n$  is generated by  $E_1, \dots, E_{n-1}$ . By [16, Theorem 2], there is an epimorphism  $f : \Gamma_n \rightarrow \mathcal{G}_n$  whose kernel is equal to

$$\langle 2e_1, e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n \rangle.$$

It follows that  $\widehat{h}_e(\gamma_j) = \delta_j E_j$  where  $\delta_j \in \{-1, 1\}$ . Since  $\lambda(\widehat{h}_e)(\gamma'_i) = A_i = h(\gamma_i)$  we have  $\lambda \circ \widehat{h}_e = h$  so that  $\widehat{h}_e$ , define  $2^{n-1}$  different spin structures on  $M^n$ . By Lemma 12, there are no other all spin structures on  $M^n = \mathbb{R}^n/\Gamma_n$ .

**b)** The maximal abelian subgroup  $\Gamma_0$  of  $\Gamma_n$  is equal to  $\langle a_1, \dots, a_n \rangle$  and  $h(\Gamma) = \langle A_1, \dots, A_{n-1} \rangle$ . As in Section 2 consider  $\mathcal{B} = \{b \in \mathbf{R}^n : \langle b, \gamma \rangle \in \mathbf{Z} \text{ for } \gamma \in \Gamma_0\}$ . For every  $b \in \mathcal{B}$  and every nontrivial element  $h$  of the holonomy group  $\langle A_1, \dots, A_{n-1} \rangle$  we have  $hb \neq b$ . By [12, Theorem 4.2],  $\eta_{M^n}(0) = 0$ .

If  $v$  is a harmonic spinor on  $M^n$ , with the spin structure  $h_\delta$ , then  $gv = v$  for every  $g \in \widehat{h}_\delta(\Gamma_n) = \mathcal{G}_n$ . Since  $-1 \in \mathcal{G}_n$  we have  $-v = v$  and thus  $v = 0$ . This finishes the proof of Theorem 2.  $\square$

**Remark 4.** If  $k$  is even and  $M^n \in \mathcal{F}_{HW}$ , then  $M^n$  does not admit any spin structure. To see this assume the contrary that the holonomy homomorphism  $h : \Gamma_n \rightarrow SO(n)$  has a lift  $\widehat{h} : \Gamma_n \rightarrow Spin(n)$ . Clearly  $\#\text{im } \widehat{h} = 2\#\text{im } h$ . Let  $E_j$  be as in the proof of Theorem 2. Since  $\lambda(E_j) = A_j$  we have  $\widehat{h}(\gamma_j) = \pm E_j$ . As  $k$  is even,  $E_j^2 = 1$  so that  $\widehat{h}(a_j) = \widehat{h}(\gamma_j^2) = 1$  and  $\Gamma_0 \subseteq \ker \widehat{h}$ . In particular,  $\#\text{im } \widehat{h} \leq \#(\Gamma_n/\Gamma_0) = \#\text{im } h$ . This is impossible, because the kernel of the epimorphism  $\lambda|_{\langle E_1, \dots, E_{n-1} \rangle}$  is nontrivial. For other examples of closed flat manifold without any spin structure we refer to [17] and [18]. It is easy to see that any flat manifold, whose holonomy group has odd order, admits a spin structure (cf. [18, Corollary 1.3] and [7, page 40]).

## References

- [1] ATIYAH, M.F.; PATODI, V.K.; SINGER I.M.; Spectral asymmetry and Riemannian geometry I. *Math. Proc. Cambridge Philos. Soc.* **77** (1975), 43-69;
- [2] BÄR CH.; Dependence of the Dirac spectrum on the spin structure. *Sémin. Congr.* **4** (2000), 17-33;

- [3] BROWN K.S.: *Cohomology of groups*. Springer, Berlin 1982.
- [4] CHARLAP L.S.: *Bieberbach Groups and Flat Manifolds*. Springer-Verlag, 1986.
- [5] DAHL M.; Dependence on the spin structure of the eta and Rokhlin invariants. *Topology Appl.* **118** (2002), 345-355;
- [6] FRIEDRICH T.; Zur Abhängigkeit des Dirac-Operators von der Spin-Structur. *Colloq. Math.* **47** (1984), 57-62.
- [7] FRIEDRICH T.: *Dirac Operators in Riemannian Geometry*. AMS, Graduate Studies in Math., Vol. 25 Providence, Rhode Island, 2000.
- [8] HILLER H., SAH C.H.; Holonomy of flat manifolds with  $b_1 = 0$ . *Q. J. Math.* **37** (1986), 177-187;
- [9] HITCHIN N.; Harmonic spinors. *Adv. Math.* **14** (1974), 1-55;
- [10] LONG D.D.; REID A.W.; On the geometric boundaries of hyperbolic 4-manifolds. *Geom. Topol.* **4** (2000), 171-178;
- [11] MIATELLO R.J.; ROSSETTI J.P.; Hantzsche-Wendt manifolds. *J.Reine Angew. Math.* **515** (1999), 1-23;
- [12] PFAFFLE F.; The Dirac spectrum of Bieberbach manifolds. *J. Geom. Phys.* **35** (4) 2000, 367-385;
- [13] QUILLEN D.; The mod 2 cohomology rings of extra-special 2-groups and the spinor groups. *Math. Ann.* **194** (1971), 197-212;
- [14] ROSSETTI J.P.; SZCZEPAŃSKI A. Generalized Hantzsche-Wendt flat manifolds. preprint 2003;
- [15] SZCZEPAŃSKI A.; Aspherical manifolds with the  $\mathbb{Q}$ -homology of a sphere. *Mathematika* **30** (1983), 291-294;
- [16] SZCZEPAŃSKI A.; The euclidean representations of the Fibonacci groups. *Q. J. Math.* **52** (2001), 385-389;
- [17] SUNG MO IM, HEUNG KI KIM; Compact flat manifolds with non-vanishing Stiefel-Whitney classes. *Topology Appl.* **96** (1999), 267-276;

- [18] VASQUEZ A.T.; Flat Riemannian manifolds. *J. Differential Geom.* **4** (1970), 367-382;
- [19] WOLF J.A.: *Spaces of constant curvature*. McGraw-Hill, 1967.

Institute of Mathematics  
University of Gdańsk  
ul. Wita Stwosza 57  
80 - 952 Gdańsk  
Poland  
E - mail : msa @ delta.math.univ.gda.pl  
E - mail : matas @ paula.univ.gda.pl