

# On approximation of topological algebraic systems by finite ones

L.Yu. Glebsky, E.I. Gordon,\* C.W.Henson†

## Abstract

We introduce and discuss a definition of approximation of a topological algebraic system  $A$  by finite algebraic systems of some class  $\mathcal{K}$ . For the case of a discrete algebraic system this definition is equivalent to the well-known definition of a local embedding of an algebraic system  $A$  in a class  $\mathcal{K}$  of algebraic systems. According to this definition  $A$  is locally embedded in  $K$  iff it is a subsystem of an ultraproduct of some systems in  $\mathcal{K}$ . We obtain a similar characterization of approximation of a locally compact system  $A$  by systems in  $\mathcal{K}$ .

We introduce the bounded formulas of the signature of  $A$  and their approximations similar to those introduced by C.W.Henson [8] for Banach spaces. We prove that a positive bounded formula  $\varphi$  holds in  $A$  if all precise enough approximations of  $\varphi$  hold in all precise enough approximations of  $A$ .

We prove that a locally compact field cannot be approximated by finite associative rings (not necessary commutative). Finite approximations of the field  $\mathbb{R}$  can be concedered as computer systems for reals. Thus, it is impossible to construct a computer arithmetic for reals that is an associative ring.

## 1 Introduction

The numerical systems that are implemented in computers for the simulation of the field  $\mathbb{R}$  are finite algebraic systems with two binary operations  $\oplus$  and  $\odot$ . The support of any such system  $R$  (we denote this support by  $R$  also) is a subset of  $\mathbb{R}$  such that  $a \in R$  iff  $-a \in R$ , where  $a \in \mathbb{R}$ . The operations  $\oplus$  and  $\odot$  are defined as follows. Let  $M$  be a maximal element of  $R$ . If  $x, y \in R$  and  $x + y$  ( $x \cdot y$ )  $\in [-M, M]$  then  $x \oplus y$  ( $x \odot y$ ) is the nearest to  $x + y$  ( $x \cdot y$ ) element of  $R$ . Here  $+$  and  $\cdot$  are the addition and the multiplication in  $\mathbb{R}$ . If  $x + y$  ( $x \cdot y$ )  $\notin [-M, M]$  then  $x \oplus y$  ( $x \odot y$ ) are defined more or less arbitrary. If such overfilling of the memory would happen during a computation we obtain a wrong result, so it is necessary to take care for the overfilling of memory not to occur.

Usually  $R$  is an  $\varepsilon$ -grid on some interval  $[-N, N] \subseteq [-M, M]$  for a small enough  $\varepsilon \geq \delta$  – the minimal positive element of  $R$ . So for any  $x, y \in R$  if  $x + y$  ( $x \cdot y$ )  $\in [-N, N]$  then  $x \oplus y$  ( $x \odot y$ ) approximates  $x + y$  ( $x \cdot y$ ) with the accuracy  $\varepsilon$ .

The concrete systems  $R$  that are implemented in working computers are based on the representation of reals in the form with floating point. These systems are discussed in the monograph [12], where it is shown that they are neither associative, nor distributive. These properties hold only approximately with the accuracy that depends on  $M$  and  $\varepsilon$ . A natural question arising here is the following one. How to construct the approximate versions of more complicated theorems about reals? More precisely, given a proposition  $\varphi$  concerning reals is it possible to construct for any  $M$  and  $\varepsilon$  a proposition  $\varphi_{M,\varepsilon}$  such that  $\varphi$  holds for reals if and only if the proposition  $\varphi_{M,\varepsilon}$  holds for all systems  $R$  that approximate the field  $\mathbb{R}$  on the interval  $[-M, M]$  with the accuracy  $\varepsilon$  for all big enough  $M$  and small enough  $\varepsilon$ ? This question may be important for understanding the following problem. Suppose that we use some convergent numerical method for computation of a real function, or a functional, or an operator. The theorem about convergence of this method is a theorem about the field  $\mathbb{R}$  but we use a finite system  $R$  implemented in our computer, which approximates  $\mathbb{R}$ . Can we be sure that the obtained result of our computation is approximately right if we can use big enough numbers and high enough accuracy in our computer? The fact that this problem is natural can be demonstrated by the following example that was discussed in some first texts on FORTRAN.

\*The work was supported by NSF Grant DMS-9970009

†The work was supported by NSF Grant DMS-9970009

Though the Taylor's series for  $\sin x$  converges for all  $x$  the approximate computation of  $\sin x$  based on its Taylor's expansion gives a wrong answer for a big enough  $x$  even if we do not need a very high accuracy  $\delta$ . The reason is that we have to proceed with our computation until we obtain the term of our series that is less than  $\delta$ . If  $a_k$  is the  $k$ -th term of the series, then  $a_{k+1} = -a_k \cdot \mu_k$ , where  $\mu_k = \frac{x^2}{2k(2k+1)}$ . For big enough  $x$  the term  $\mu_k$  becomes less than  $\varepsilon$  - the minimum positive number in our finite system  $R$  - earlier than  $a_{k+1}$  becomes less than  $\delta$ . When  $\mu_k < \varepsilon$  it is considered as zero by a computer. Hence we obtain zero for all  $a_l$  with  $l > k$  and it makes no sense to proceed our computation any more.

Another problem concerning the computer systems approximating  $\mathbb{R}$  is the following one. What properties of reals can hold for approximating finite systems. More precisely, let  $\varphi$  be any theorem about the field  $\mathbb{R}$ . Is it true that for any big enough  $M$  and small enough  $\varepsilon$  there exists a finite system  $R$  that approximates  $\mathbb{R}$  on the interval  $[-M, M]$  with the accuracy  $\varepsilon$  and such that  $\varphi$  holds for  $R$ ? For example, we mentioned above that the operations  $\oplus$  and  $\odot$  in numerical systems based on the floating point representation are neither associative nor distributive. It is easy to construct approximating systems for  $\mathbb{R}$  that are abelian groups for  $\oplus$  (see Example 2 of Section 2). Is it possible to construct finite rings that approximate  $\mathbb{R}$ ?

These problems are discussed in the present paper in a more general setting. We consider a locally compact algebraic system  $\mathcal{A} = \langle A, \sigma \rangle$  of the finite signature  $\sigma$  that contains only functional symbols (a universal algebra) and give a definition of approximation of this system by a finite system  $\mathcal{A}_f$  on a compact set  $C \subset A$  with the accuracy  $U$ , where  $U$  is an element of the canonical uniformity on  $C$  ( $\langle C, U \rangle$ -approximation). For example if the topology on  $A$  is defined by a metric  $\rho$  then  $U = \{ \langle x, y \rangle \in C^2 \mid \rho(x, y) < \varepsilon \}$  for some  $\varepsilon > 0$ . The universal algebra  $\mathcal{A}$  is said to be approximable by finite algebras of a class  $\mathcal{K}$  ( $\mathcal{K}$ -algebras) if for any  $C$  and  $U$  there exists a  $\langle C, U \rangle$ -approximation  $\mathcal{A}_f \in \mathcal{K}$ . The definition of approximation of a locally compact group by finite groups, which is discussed in [6] is a particular case of this definition. It is known [6] that all locally compact commutative groups are approximable by finite groups but this is wrong for noncommutative groups [7]. There exist noncommutative groups that are approximable neither by finite groups, nor by finite semigroups, nor by finite quasigroups [1], [7], [5]. Basing on these results we show here that locally compact fields are not approximable by finite associative rings (Theorem 1). In [7], it was proved under some additional assumptions that the field  $\mathbb{R}$  is not approximable by finite fields. From the point of view of the above discussion of computer numerical systems this result means that it is impossible to implement a numerical system for simulating the reals in a computer that is a finite associative ring.

In [2] (see also [4]) finite approximations of locally compact abelian groups are used for a construction of finite dimensional approximations of pseudodifferential operators. Under this approach the operators are approximated together with the group structures connected with them. This allows to construct approximations, which have some nice properties, e.g. uniform convergence and spectrum convergence. Usually algebraic and geometric structures connected with operators can be considered as finite dimensional manifolds (e.g. the symmetry groups of operators are often Lie groups). Thus the approximations of these structures are based on approximations of the field  $\mathbb{R}$  together with some other continuous functions on  $\mathbb{R}$ . The approximations of the other locally compact fields can be used in  $p$ -adic analysis, adelic analysis, etc. This is another reason for investigation of finite approximations of topological algebraic systems.

We investigate the problem of existence of approximate versions of theorems that was discussed in the beginning of this introduction. We consider here only theorems formulated in  $L_\sigma$  - the first order language of the signature  $\sigma$ . The obtained results are similar to the well-known results about positive bounded formulas of the theory of Banach spaces [8], [9], [10].

Recall that a formula is *positive* if it can be built up from atomic formulas using only conjunction, disjunction and universal and existential quantifiers. We consider *positive bounded* formulas, i.e. positive formulas in the prenex form such that all involved existential quantifiers are bounded by compact subsets of  $A$  and all involved universal quantifiers are bounded by relatively compact open subsets of  $A$ . These formulas are not of the language  $L_\sigma$  but they have the obvious interpretation in  $\mathcal{A}$ . We say that a formula  $\varphi'$  is an *approximation* of a positive bounded formula  $\varphi$  if it is obtained

- 1) by an extending of a compact set, which bounds a variable under each existential quantifier,
- 2) by a shrinking of an open set, which bounds a variable under each universal quantifier,
- 3) by replacing the equality in each atomic subformula by an approximate equality, i.e. by the statement that the ordered pair of terms of this equality is contained in a given element of the uniformity on  $A$ .

We prove (Theorem 6) that  $\mathcal{A} \models \varphi$  iff for any approximation  $\varphi'$  for all big enough  $C$  and small enough

$U$  holds  $\mathcal{A}_{(C,U)} \models \varphi'$ . Here  $\mathcal{A}_{(C,U)}$  is a  $\langle C, U \rangle$ -approximation of  $\mathcal{A}$  and  $\varphi$  is a positive bounded sentence. Hence, positive bounded theorems about  $A$  have approximate versions in the sense discussed above.

The proof of this statement uses the nonstandard analysis.

If the topology on  $A$  is discrete then it is well-known [14] that  $\mathcal{A}$  is approximable by algebras of a class  $\mathcal{K}$  if it is isomorphic to a subalgebra of an ultraproduct of  $\mathcal{K}$ -algebras. Using the language of nonstandard analysis the last statement can be reformulated in the following way.

A discrete universal algebra  $\mathcal{A}$  is approximable by  $\mathcal{K}$ -algebras if it is isomorphic to a subalgebra (maybe external) of an internal  $^*\mathcal{K}$ -algebra.

We prove that if a non-discrete locally compact topological algebra  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -algebras then it is isomorphic to the nonstandard hull of an appropriate hyperfinite  $^*\mathcal{K}$ -algebra  $\mathcal{A}_h$ . We say that  $\mathcal{A}_h$  is a hyperfinite approximation of  $\mathcal{A}$ .

If  $\varphi$  is a sentence in  $L_\sigma$  (not necessary positive bounded) then denote by  $\varphi_\approx$  the formula that is obtained from  $\varphi$  by restricting all quantifiers in  $\varphi$  by the set of nearstandard elements and by replacing all equalities in  $\varphi$  by the relation  $\approx$  - "to be infinitesimally close". Then  $\mathcal{A} \models \varphi$  iff for any hyperfinite approximation  $\mathcal{A}_h$  of  $\mathcal{A}$  holds  $\mathcal{A}_h \models \varphi_\approx$ . For the case of positive bounded formulas the translation of the last statement into the standard language with the help of the Nelson's algorithm [15] gives Theorem 6.

## 2 Approximation of locally compact fields

Let  $\mathcal{A} = \langle A, \sigma \rangle$  be an algebraic system of a finite signature  $\sigma$  that contains only functional symbols. Recall that such algebraic systems are said to be the universal algebras. In what follows we call them simply algebras. We assume also that  $A$  is endowed with a locally compact topology and the functional symbols of  $\sigma$  are interpreted by continuous functions, which we denote by the same letters as the corresponding functional symbols in  $\sigma$ .

Let  $C \subset A$  be a compact set,  $\mathcal{U}$  - a finite covering of  $C$  by relatively compact open sets (r.c.o. covering),  $\mathcal{A}_f = \langle A_f, \sigma \rangle$  - a finite algebra of the signature  $\sigma$  and  $j : A_f \rightarrow A$  - a mapping. The interpretation of a functional symbol  $g \in \sigma$  in  $\mathcal{A}_f$  is denoted by  $g_f$ . For  $a_1, \dots, a_n \in A_f$  we denote by  $j(\langle a_1, \dots, a_n \rangle)$  the  $n$ -tuple  $\langle j(a_1), \dots, j(a_n) \rangle$ . We say that  $a, b \in C$  are  $\mathcal{U}$ -close if  $\exists U \in \mathcal{U} (a \in U \wedge b \in U)$ .

**Definition 1** 1. We say that a set  $M \subset A$  is a  $(C, \mathcal{U})$ -grid if for any  $c \in C$  there exists an  $m \in M$  such that  $c$  and  $m$  are  $\mathcal{U}$ -close.

2. We say that  $j$  is a  $(C, \mathcal{U})$ -homomorphism if for any  $n$ -ary functional symbol  $g \in \sigma$  and for any  $\bar{a} \in A_f^n$  such that  $j(\bar{a}) \in C^n$  and  $g(j(\bar{a})) \in C$ , the elements  $g(j(\bar{a}))$  and  $j(g_f(\bar{a}))$  are  $\mathcal{U}$ -close.
3. We say that the pair  $\langle \mathcal{A}_f, j \rangle$  is a  $(C, \mathcal{U})$ -approximation of  $\mathcal{A}$  if  $j$  is a  $(C, \mathcal{U})$ -homomorphism and  $j(A_f)$  is a  $(C, \mathcal{U})$ -grid.
4. Let  $\mathcal{K}$  be a class of finite algebras of the signature  $\sigma$ . We say that the locally compact algebra  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -algebras if for any compact set  $C \subset A$  and for any finite r.c.o. covering  $\mathcal{U}$  of  $C$  there exists a  $(C, \mathcal{U})$ -approximation  $\langle \mathcal{A}_f, j \rangle$  of  $\mathcal{A}$  such that  $\mathcal{A}_f \in \mathcal{K}$ .

Usually we deal with the case when the set  $A$  is equipped with a uniformity  $\mathcal{W}$  such that there exists an element  $W \in \mathcal{W}$  that satisfies the following condition: for any  $x \in A$  the set  $W(x) = \{y \in A \mid \langle x, y \rangle \in W\}$  is relatively compact. In this case the uniformity  $\mathcal{W}$  determines a locally compact topology on  $A$  and we say that the uniform space  $A$  is uniformly locally compact. For example, all locally compact groups satisfy this condition. When we consider the uniformly compact algebras of the signature  $\sigma$  we assume that the interpretations of functional symbols are some continuous, but not necessary uniformly continuous, functions. For example  $\mathbb{R}$  is a uniformly locally compact space but the multiplication in  $\mathbb{R}$  is not uniformly continuous. It follows from the general theory of the uniform spaces (see, for example, [3]) that the restriction of a continuous function by a compact subset  $C$  is uniformly compact on  $C$ . For the case of uniformly locally compact algebras our definition can be simplified.

We assume now that  $\mathcal{A}$  is a uniformly locally compact algebra of the signature  $\sigma$  and  $W$  is an element of uniformity such that  $\forall x \in A \overline{W(x)}$  is compact (here and below the closure of a set  $E$  is denoted by  $\overline{E}$ ). Without loss of generality we can assume that  $W$  is symmetric i.e.  $\langle x, y \rangle \in W$  iff  $\langle y, x \rangle \in W$ . The items  $C$ ,  $\mathcal{A}_f$  and  $j$  satisfy the same assumptions as above. We say that  $a, b \in C$  are  $W$ -close if  $\langle a, b \rangle \in W$ .

**Definition 2** 1. We say that a set  $M \subset A$  is a  $(C, W)$ -grid if for any  $c \in C$  there exists an  $m \in M$  such that  $c$  and  $m$  are  $W$ -close.

2. We say that  $j$  is a  $(C, M)$ -homomorphism if for any  $n$ -ary functional symbol  $g \in \sigma$  and for any  $\bar{a} \in A_f^n$  such that  $j(\bar{a}) \in C^n$  and  $g(j(\bar{a})) \in C$ , the elements  $g(j(\bar{a}))$  and  $j(g_f(\bar{a}))$  are  $W$ -close.

3. We say that the pair  $\langle \mathcal{A}_f, j \rangle$  is a  $(C, W)$ -approximation of  $\mathcal{A}$  if  $j$  is a  $(C, W)$ -homomorphism and  $j(\mathcal{A}_f)$  is a  $(C, W)$ -grid. If  $\mathcal{A}_f \subset \mathcal{A}$  and  $j$  is the identical inclusion we say that  $\mathcal{A}_f$  is a  $(C, W)$ -approximation of  $\mathcal{A}$ .

4. Let  $\mathcal{K}$  be a class of finite algebras of the signature  $\sigma$ . We say that a locally compact algebra  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -algebras if for any compact set  $C \subset A$  and for any  $W \in \mathcal{W}$  such that  $\forall x \in A \overline{W(x)}$  is compact, there exists a  $(C, W)$ -approximation  $\langle \mathcal{A}_f, j \rangle$  of  $\mathcal{A}$  such that  $\mathcal{A}_f \in \mathcal{K}$ .

**Remark 1** If  $A$  is a compact set then  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -algebras if for any  $W \in \mathcal{W}$  there exists a finite  $\mathcal{K}$ -algebra, which is an  $(A, W)$ -approximation of  $\mathcal{A}$ .

Let us consider some examples of approximations of the field  $\mathbb{R}$ . We use the signature  $\sigma = \langle +, \cdot \rangle$ .

Since any compact  $C \subset \mathbb{R}$  is contained in the interval  $[-a, a]$  for an appropriate  $a$  and the sets  $W_\varepsilon = \{ \langle x, y \rangle \mid |x - y| < \varepsilon \}$ ,  $\varepsilon > 0$  form a base of the uniformity on  $\mathbb{R}$ , it is enough to consider only the  $([-a, a], W_\varepsilon)$ -approximations of  $\mathbb{R}$ . We call these approximations the  $(a, \varepsilon)$ -approximations.

**Example 1** Recall that the normal (computer) form of a real  $\alpha$  is its representation:

$$\alpha = \pm 10^p \cdot 0.a_1a_2\dots, \quad (1)$$

where  $p \in \mathbb{Z}$ , and  $a_1a_2\dots$  is a finite or infinite sequence of decimal digits  $0 \leq a_n \leq 9$ , and  $a_1 \neq 0$ . The integer  $p$  is called the exponent of  $\alpha$ , the sequence  $a_1a_2\dots$  - its mantissa.

Fix two natural numbers  $P > Q$ . Consider the finite set  $A_{PQ}$  of reals in the form (1) such that the exponent  $p$  of  $\alpha$  satisfies the inequality  $|p| \leq P$  and its mantissa contains no more than  $Q$  decimal digits. Define the following two binary operations  $\oplus$  and  $\odot$  on  $A_{PQ}$ . For any two elements  $\alpha, \beta \in A_{PQ}$  let  $\alpha \times \beta$  be either  $\alpha \times \beta$  or  $\alpha + \beta$ . Consider the normal form of  $\alpha \times \beta$ :

$$\alpha \times \beta = \pm 10^r \cdot 0.c_1c_2\dots \quad (2)$$

Notice that the mantissa of  $c_1c_2\dots$  may contain more than  $Q$  digits. Put

$$\alpha \otimes \beta = \begin{cases} \pm 10^r \cdot 0.c_1c_2\dots c_Q, & \text{if } |r| \leq P \\ \pm 10^P \cdot 0.\underbrace{99\dots 9}_Q, & \text{if } r > P \\ 0, & \text{if } r < -P \end{cases}$$

If the mantissa  $c_1c_2\dots$  contains less than  $Q$  digits we complete it to a  $Q$ -digits mantissa by zeros.

We denote by  $\mathcal{A}_{PQ}$  the algebra  $\langle A_{PQ}, \sigma \rangle$ , such that the interpretations of the functional symbols  $+$  and  $\cdot$  are the functions  $\oplus$  and  $\odot$  respectively.

Let us show that for any positive  $a$  and  $\varepsilon$  there exist natural numbers  $P$  and  $Q$  such that the algebra  $\mathcal{A}_{PQ}$  is an  $(a, \varepsilon)$ -approximation of  $\mathbb{R}$ .

Fix an arbitrary  $a > 0$  and a real  $\varepsilon$  such that  $0 < \varepsilon < 1$  and put  $r = [\log_{10}(a + \varepsilon)] + 1$ ,  $Q = r + [\log_{10} \frac{1}{\varepsilon}] + 1$ . Here  $[x]$  is the integer part of a real  $x$ . Then  $a < 10^r - 10^{r-Q}$  and  $10^{r-Q} < \varepsilon$ . Fix an arbitrary natural number  $P > Q$  and consider the universal algebra  $\mathcal{A}_{PQ}$ . Let  $B = \{ \pm k \cdot 10^{r-Q} \mid k = 0, 1, \dots, 10^Q - 1 \}$ . Obviously  $B \subset A_{PQ}$  (the exponent of an arbitrary element of  $B$  does not exceed  $r < P$  and its mantissa does not contain more than  $Q$  digits) and  $B$  is a  $10^{r-Q}$ -grid of the interval  $[-(10^r - 10^{r-Q}), 10^r - 10^{r-Q}]$  and thus it is an  $\varepsilon$ -grid of the interval  $[-a, a]$ . If for  $\alpha, \beta \in B \cap [-a, a]$ , holds  $\alpha \times \beta \in [-a, a]$  then  $|\alpha \times \beta| < 10^r$ . Let the representation of  $\alpha \times \beta$  in the form with the fixed point be  $\alpha \times \beta = m_1\dots m_s.m_{s+1}m_{s+2}\dots$ . Then  $\alpha \otimes \beta = m_1\dots m_s.m_{s+1}m_{s+2}\dots m_{Q-s}$ . By the previous inequality  $Q - s \geq Q - r$ . Thus,  $|\alpha \times \beta - \alpha \otimes \beta| \leq 10^{r-Q} < \varepsilon$ .

The described systems  $\mathcal{A}_{PQ}$  are implemented in working computers. What properties of addition and multiplication of reals hold for  $\oplus$  and  $\odot$ ?

It is easy to see that the operations  $\oplus$  and  $\odot$  are commutative,  $\xi \oplus (-\xi) = 0$  and  $\xi \oplus 0 = \xi$  for any  $\xi \in A_{PQ}$

Let  $\alpha = \beta = 0.\underbrace{60\dots 06}_Q$ ,  $\gamma = 0.\underbrace{60\dots 5}_Q$ . Then  $\alpha \oplus \beta = \alpha \oplus \gamma$ , so the cancellation law fails for  $\oplus$ . Thus, the law of associativity for  $\oplus$  fails also.

The following examples show that the laws of associativity for  $\odot$  and distributivity fail.

$$\left( 0.\underbrace{99\dots 9}_Q \odot 2 \right) \odot 0.5 = 0.\underbrace{99\dots 5}_Q,$$

while

$$\begin{aligned} 0.\underbrace{99\dots 9}_Q \odot (2 \odot 0.5) &= 0.\underbrace{99\dots 9}_Q. \\ 2 \odot 0.\underbrace{99\dots 9}_Q \oplus 1 \odot 0.\underbrace{99\dots 9}_Q &= \underbrace{2.99\dots 8}_Q, \end{aligned}$$

while

$$(2 \oplus 1) \odot 0.\underbrace{99\dots 9}_Q = \underbrace{2.99\dots 7}_Q.$$

We use here that  $(2 \odot 0.5) = 1$  and  $(2 \oplus 1) = 3$ .

**Example 2.** Fix a natural number  $M$  and a positive  $\Delta$ . Put  $A'_{M\Delta} = \{k\Delta \mid k = -M \dots M\}$ . Let  $N = 2M + 1$ . For any  $n \in \mathbb{Z}$  denote by  $n(\pmod N)$  the element of the set  $\{-M, \dots, M\}$  which is congruent  $n$  modulo  $N$ . The operations  $\oplus$  and  $\odot$  on  $A'_{M,\Delta}$  are defined as follows.

$$k\Delta \oplus m\Delta = (k + m)(\pmod N)\Delta \quad (3)$$

$$k\Delta \odot m\Delta = [km\Delta](\pmod N)\Delta. \quad (4)$$

Denote by  $\mathcal{A}'_{M,\Delta}$  the algebra of the signature  $\sigma$  with the underlying set  $A'_{M,\Delta}$  and the interpretation of the functional symbols defined by formulas (3) and (4).

Let us show that for any positive  $a$  and real  $\varepsilon$  such that  $0 < \varepsilon < 1$  there exists an  $(a, \varepsilon)$ -approximation of  $\mathbb{R}$  of the form  $\mathcal{A}'_{M,\Delta}$ . Fix a  $\Delta \leq \varepsilon$ , a  $b \geq a + \Delta$  and put  $M = \lfloor \frac{b}{\Delta} \rfloor$ . Obviously  $A'_{M\Delta}$  is a  $(b, \Delta)$ -grid and thus an  $(a, \varepsilon)$ -grid. If  $k\Delta + m\Delta \in [-a, a]$ , then  $|(k + m)\Delta| \leq a < b$ . Hence  $|(k + m)| \leq \lfloor \frac{b}{\Delta} \rfloor = M$  and  $(k + m)(\pmod N) = k + m$ . It follows from formula (3) that in this case  $k\Delta \oplus m\Delta = k\Delta + m\Delta$ .

If  $k\Delta, m\Delta, km\Delta^2 \in [-a, a]$  then  $|km\Delta| \leq \frac{b}{\Delta} - 1$  and thus  $[km\Delta] \in [-M, M]$ , i.e.  $[km\Delta](\pmod N) = [km\Delta]$ . Hence

$$|km\Delta^2 - k\Delta \odot m\Delta| = (km\Delta - [km\Delta])\Delta \leq \Delta.$$

This proves that  $\mathcal{A}'_{M,\Delta}$  is an  $(a, \varepsilon)$ -approximation of the field  $\mathbb{R}$ .

It is obvious that  $\mathcal{A}'_{M,\Delta}$  is an abelian group with respect to  $\oplus$  (see (3)). Let us show that for any big enough  $M$  and small enough  $\Delta$  the multiplication  $\odot$  satisfies neither the law of associativity, nor the law of distributivity. Fix any  $0 < k < \frac{M}{2}$  such that  $\{k\Delta\} \geq 0.5$ , where  $\{k\Delta\} = k\Delta - [k\Delta]$  is the fractional part of  $k\Delta$ . Such  $k$  exists for any small enough  $\Delta$  since  $\{k\Delta \mid k = -M, \dots, M\}$  is a  $\Delta$ -grid on  $[-a, a]$ . In this case

$$[k\Delta + k\Delta] = [2k\Delta] = 2[k\Delta] + 1. \quad (5)$$

Since  $0 < k < \frac{M}{2}$  we have  $k\Delta \oplus k\Delta = k\Delta + k\Delta = 2k\Delta$ . By formulas (4) and (5) holds

$$(k\Delta \oplus k\Delta) \odot \Delta = (2k\Delta) \odot \Delta = [2k\Delta]\Delta = (2[k\Delta] + 1)\Delta.$$

On the other hand

$$k\Delta \odot \Delta + k\Delta \odot \Delta = [k\Delta]\Delta + [k\Delta]\Delta = 2[k\Delta]\Delta.$$

So the law of distributivity fails in  $\mathcal{A}'_{M,\Delta}$ .

The law of associativity for  $\odot$  also fails in  $\mathcal{A}'_{M,\Delta}$  since, for example,

$$\left( \left[ \frac{1}{2\Delta} \right] \Delta \odot \left[ \frac{2}{\Delta} \right] \Delta \right) \odot M\Delta \neq \left[ \frac{1}{2\Delta} \right] \Delta \odot \left( \left[ \frac{2}{\Delta} \right] \Delta \odot M\Delta \right). \quad (6)$$

Indeed, it is easy to see that for a small enough  $\Delta$  the left-hand side of the inequality (6) is an element of the interval  $[\frac{M}{2}\Delta, M\Delta]$ , while the right-hand side is negative.

**Example 3.** Consider an example of approximation of the field  $\mathbb{Q}_p$  of  $p$ -adic numbers. Any non-archimedean locally compact field can be approximated in a similar way. Recall that any  $p$ -adic number  $\alpha$  can be uniquely represented in the form

$$\alpha = \sum_{\nu=n}^{\infty} a_{\nu} p^{\nu}, \quad (7)$$

where  $n, a_{\nu} \in \mathbb{Z}$ ,  $0 \leq a_{\nu} < p$ ,  $a_n \neq 0$ . The  $p$ -adic norm of  $\alpha$  is given by the formula

$$|\alpha|_p = p^{-n}, \quad (8)$$

The set  $\mathbb{Z}_p = \{\alpha \mid |\alpha|_p \leq 1\}$  is a compact subring of  $\mathbb{Q}_p$  which is called the ring of integer  $p$ -adic numbers. From the multiplicativity of  $p$ -adic norm, we get that for any  $n \in \mathbb{Z}$  the subgroup  $p^n \mathbb{Z}_p = \{\alpha \mid |\alpha|_p \leq p^{-n}\}$ . It is obvious that this subgroup is compact. Thus the sequence  $\{p^n \mathbb{Z}_p \mid n \in \mathbb{Z}\}$  is a decreasing sequence of compact sets that covers  $\mathbb{Q}_p$ . Hence it is enough to consider only the  $(p^n \mathbb{Z}_p, p^{-m})$ -approximations of  $\mathbb{Q}_p$  for all  $n \in \mathbb{Z}$ ,  $n < 0$ ,  $m \in \mathbb{N}$ , which we call simply the  $(m, n)$ -approximations.

For any  $n > 0$  the subgroup  $p^n \mathbb{Z}_p$  is an ideal in  $\mathbb{Z}_p$  and the quotient ring  $K_n$  is equal to  $\mathbb{Z}/p^n \mathbb{Z}$ . Represent the elements of this ring by the least positive residuals modulo  $p^n$ . Then  $K_n = \{0, 1, \dots, p^n - 1\}$ . Represent each  $k \in K$  in the  $p$ -based number system:

$$k = \sum_{\nu=0}^{n-1} a_{\nu} p^{\nu}. \quad (9)$$

Comparing (7) and (9), we see that the inclusion  $K_n \subset \mathbb{Z}_p$  holds for the sets  $K_n$  and  $\mathbb{Z}_p$ . However the ring operations in these sets are distinct. Indeed, the addition  $+$  and multiplication  $\cdot$  of natural numbers in  $\mathbb{Z}_p$  are the same as in  $\mathbb{N}$  while the addition  $\oplus$  and multiplication  $\odot$  in  $K_n$  are the addition and multiplication modulo  $p^n$ . For  $\alpha = \sum_{\nu=0}^{\infty} a_{\nu} p^{\nu} \in \mathbb{Z}_p$  denote by  $\alpha_n$  the number  $\sum_{\nu=0}^{n-1} a_{\nu} p^{\nu} \in K_n$ . Then  $|\alpha - \alpha_n|_p \leq p^{-n}$ . Hence,  $K_n$  is a  $p^{-n}$ -grid in  $\mathbb{Z}_p$ . It is easy to see that

$$|\alpha + \beta - (\alpha_n \oplus \beta_n)|_p, |\alpha \cdot \beta - (\alpha_n \odot \beta_n)|_p \leq p^{-n}.$$

Thus, the identical embedding of  $K_n$  into  $\mathbb{Z}_p$  is a  $p^{-n}$ -homomorphism. Hence, the ring  $K_n$  is a  $(\mathbb{Z}_p, p^{-n})$ -approximation of  $\mathbb{Z}_p$  (see Remark 1). We see that the compact ring  $\mathbb{Z}_p$  is approximable by finite commutative associative rings.

To construct an  $(-n, m)$ -approximation ( $n, m > 0$ ) of  $\mathbb{Q}_p$  consider the set  $H_{n,m} \subset p^{-n} \mathbb{Z}_p$  of all numbers of the form  $\sum_{\nu=-n}^{m-1} a_{\nu} p^{\nu}$ . It is obvious that this set is a  $p^{-m}$ -grid in  $p^{-n} \mathbb{Z}_p$ . Define operations  $\hat{\oplus}$  and  $\hat{\odot}$  such that the identical embedding of  $H_{n,m}$  into  $p^{-n} \mathbb{Z}_p$  is a  $(p^{-n} \mathbb{Z}_p, p^{-m})$ -homomorphism.

Notice that  $\alpha \in H_{n,m}$  iff  $p^n \alpha \in K_{n+m}$ . For any  $\alpha, \beta \in H_{n,m}$  put

$$\alpha \hat{\oplus} \beta = p^{-n} (p^n \alpha \oplus p^n \beta), \quad (10)$$

where  $\oplus$  is the addition in  $K_{n+m}$ .

The definition of  $\hat{\odot}$  is more complicated. Let

$$p^n \alpha \cdot p^n \beta = c_0 + c_1 p + \dots + c_{n-1} p^{n-1} + c_n p^n \dots + c_{2n+2m-2} p^{2n+2m-2}. \quad (11)$$

Put

$$\alpha \hat{\odot} \beta = c_n p^{-n} + \dots + c_{2n+m-1} p^{m-1}. \quad (12)$$

Since  $\alpha, \beta \in p^{-n}\mathbb{Z}_p$  we have  $\alpha + \beta \in p^{-n}\mathbb{Z}_p$ . Thus  $p^n\alpha + p^n\beta \in \mathbb{Z}_p$  and

$$|p^n\alpha + p^n\beta - (p^n\alpha \oplus p^n\beta)|_p \leq p^{-(n+m)}$$

Dividing the left hand side of this inequality by  $p^n$ , using the properties of  $p$ -adic norm and formula (10), we obtain that

$$|\alpha + \beta - (\alpha \hat{\oplus} \beta)|_p \leq p^{-m}.$$

If  $\alpha \cdot \beta \in p^{-n}\mathbb{Z}_p$ , then  $p^n\alpha \cdot p^n\beta \in p^n\mathbb{Z}_p$ . Hence in formula (11)  $c_0 = \dots = \gamma_{n-1} = 0$ . Now, by formula (12),

$$|\alpha \cdot \beta - (\alpha \hat{\odot} \beta)|_p \leq p^{-m}.$$

This proves that the identical embedding of  $H_{n,m}$  in  $\mathbb{Q}_p$  is a  $(p^{-n}\mathbb{Z}_p, p^{-m})$ -homomorphism.

By formula (10), it is obvious that  $\langle H_{n,m}, \hat{\oplus} \rangle$  is an abelian group isomorphic to the additive group of  $K_{n+m}$ .

It is easy to see that for any integer  $c$  such that  $0 \leq c < p$  holds  $\frac{c}{p^n} \hat{\odot} \frac{1}{p} = 0$ . Thus  $\frac{1}{p^n} \hat{\odot} \frac{1}{p} \hat{\oplus} \frac{p-1}{p^n} \hat{\odot} \frac{1}{p} = 0$ , while  $\left(\frac{1}{p^n} \hat{\oplus} \frac{p-1}{p^n}\right) \hat{\odot} \frac{1}{p} = \frac{1}{p^n}$ . This shows that the distributivity law fails for  $\hat{\oplus}$  and  $\hat{\odot}$ .

Since  $0 \hat{\odot} p = 0$  and  $\frac{1}{p} \hat{\odot} p = 1$ , we have  $\left(\frac{1}{p^n} \hat{\odot} \frac{1}{p}\right) \hat{\odot} p = 0$ , while  $\frac{1}{p^n} \hat{\odot} \left(\frac{1}{p} \hat{\odot} p\right) = \frac{1}{p^n}$ . This shows that the associativity law fails for  $\hat{\odot}$ .

In all these examples the finite algebras that approximate the locally compact fields fail to be rings. Indeed, the following theorem holds.

**Theorem 1** *A locally compact field can't be approximated by finite associative rings.*

The proof of this theorem is based on the following two theorems proved in [5].

**Theorem 2** *If a locally compact group is approximable by finite semigroups then it is approximable by finite groups.*

**Theorem 3** *If a locally compact group is approximable by finite quasigroups (in particular, by finite groups) then it is unimodular (i.e. the left and right Haar measures coincide).*

It is well known [13] that if a unimodular group  $G$  acts continuously on a unimodular locally compact group  $H$  by automorphisms and this action does not preserve the Haar measure on  $H$  then the semidirect product of  $G$  and  $H$  is non-unimodular.

**Proof of Theorem 1.** Let  $K$  be a locally compact field,  $K_+$  the additive group of  $K$ , and  $K^*$  the multiplicative group of  $K$ . The multiplication in  $K$  is a continuous action of  $K^*$  on  $K_+$ . It is obvious that this action is not measure preserving. Thus, the semidirect product of these groups is non-unimodular. This semidirect product is isomorphic to the following matrix group:

$$G = \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : a \in K^*, b \in K \right\}.$$

Let us assume that  $K$  is approximable by finite associative rings and prove under this assumption that  $G$  is approximable by finite semigroups. Then by Theorem 2,  $G$  is approximable by finite groups. Thus, by Theorem 3  $G$  is unimodular. This contradiction prove Theorem 1.

The group  $G$  is homeomorphic to  $K^* \times K$  as a topological space. Put

$$W_\varepsilon = \{ \langle (a, b), (a', b') \rangle : a, a' \in K^*, b, b' \in K, \max\{|a - a'|_K, |b - b'|_K\} < \varepsilon \},$$

where  $|\cdot|_K$  is the norm in  $K$

We have to show that for any compact sets  $A \subset K^*$  and  $B \subset K$  there exists a  $(A \times B, W_\varepsilon)$ -approximation  $\langle H, j \rangle$  such that  $H$  is a semigroup.

Let  $D = A \cup B \cup A \cdot B$ . Then  $D$  is a compact subset of  $K$  (here  $A \cdot B = \{a \cdot b \mid a \in A, b \in B\}$ ). For any positive  $\delta$  denote by  $U_\delta$  the set  $\{ \langle a, b \rangle \in K \mid |a - b|_K < \delta \}$  and put  $C = \overline{U_{\varepsilon/2}(D)}$ . Since  $D$  is a compact set and any open ball in  $K$  is relatively compact, we have that  $C$  is a compact set also.

According to our assumption, there exists a finite associative ring  $\langle F, \oplus, \odot \rangle$  and a map  $i : F \rightarrow K$  such that the pair  $\langle F, i \rangle$  is a  $(C, U_{\varepsilon/2})$ -approximation of the field  $K$ . Our group  $G$  is equal to  $K^* \times K$  as a set. The multiplication in  $G$  is given by the formula

$$\langle a, b \rangle \cdot \langle c, d \rangle = \langle ac, ad + b \rangle. \quad (13)$$

Consider the finite  $S$  such that  $S = F \times F$  as a set and the multiplication in  $S$  is given by the formula

$$\langle s, p \rangle \widehat{\odot} \langle q, r \rangle = \langle s \odot q, s \odot r \oplus p \rangle. \quad (14)$$

Since  $F$  is an associative ring, it is easy to see that  $S$  is a semigroup. The map  $i : F \rightarrow K$  is defined by the formula

$$i(\langle a, b \rangle) = \langle j(a), j(b) \rangle.$$

Since  $j(F)$  is an  $\frac{\varepsilon}{2}$ -grid for  $C$  and thus, for  $A$  and  $B$ , it is obvious that  $i(S)$  is an  $\frac{\varepsilon}{2}$ -grid for  $A \times B$ .

Let  $i(\langle s, p \rangle), i(\langle q, r \rangle), i(\langle s, p \rangle \widehat{\odot} \langle q, r \rangle) \in A \times B$ . Then, by the definition of  $i$  and formula (14), we have  $j(s), j(q), j(s \odot q) \in A$ . Hence,

$$|j(s) \cdot j(q) - j(s \odot q)|_K < \frac{\varepsilon}{2} \quad (15)$$

Since  $j(p), j(r) \in B$ , we have  $j(s) \cdot j(r) \in A \cdot B \subset C$ . Thus,  $|j(s) \cdot j(r) - j(s \odot r)|_K < \frac{\varepsilon}{2}$  and  $j(s \odot r) \in C$ . Hence,

$$|j(s) \cdot j(r) + j(p) - (j(s \odot r) + j(p))|_K < \frac{\varepsilon}{2}$$

and

$$|j(s \odot r) + j(p) - j((s \odot r) \oplus p)|_K < \frac{\varepsilon}{2}.$$

Thus

$$|j(s) \cdot j(r) + j(p) - j((s \odot r) \oplus p)|_K < \varepsilon. \quad (16)$$

By the definition of  $i$  and formulas (13)-(16), we obtain that  $i$  is an  $(A \times B, W_\varepsilon)$ -homomorphism.  $\square$ .

### 3 Characterization of approximability in terms of ultraproducts. The language of nonstandard analysis.

If a universal algebra  $\mathcal{A}$  is discrete the definition 1 (4) of approximation of  $\mathcal{A}$  by finite  $\mathcal{K}$ -algebras can be reformulated in the following way.

**Proposition 1** *A discrete algebra  $\mathcal{A}$  of a finite signature  $\sigma$  is approximable by finite  $\mathcal{K}$ -algebras if for any finite subset  $C \subset A$  there exists a finite algebra  $\mathcal{A}_C \in \mathcal{K}$  and a map  $j : A_C \rightarrow A$  such that*

1.  $C \subset j(A_C)$ .
2. For any  $n$ -ary functional symbol  $f \in \sigma$  and for any  $\bar{a} \in A_C^n$  such that  $j(\bar{a}) \in C$  and  $f(j(\bar{a})) \in C$  holds

$$j(f_{\mathcal{A}_C}(\bar{a})) = f(j(\bar{a})),$$

where  $f_{\mathcal{A}_C}$  is the interpretation of  $f$  in  $\mathcal{A}_C$ .  $\square$

It is easy to see that this definition is equivalent to the well known in Model Theory (cf., for example, [14]) definition of a local embedding of  $\mathcal{A}$  in  $\mathcal{K}$ . The following proposition is contained in [14].

**Proposition 2** *A discrete algebra  $\mathcal{A}$  is approximable by  $\mathcal{K}$ -algebras iff  $\mathcal{A}$  is isomorphic to a subalgebra of an ultraproduct of  $\mathcal{K}$ -algebras.*

The characterisation of approximability of an arbitrary locally compact algebra by  $\mathcal{K}$ -algebras in terms of ultraproducts is more complicated. In order to simplify our considerations we use the language of nonstandard analysis.

Let  $I$  be an arbitrary infinite set (a set of indexes) and  $\mathcal{G}$  a free ultrafilter on  $I$ . We assume them fixed in this section. Recall that in nonstandard analysis the ultrapower  $A^{\mathcal{G}}$  is called the nonstandard extension of  $A$  and is denoted by  ${}^*A$ . For any  $c \in A$  the element of  $A^{\mathcal{G}}$ , which is of the form  $\langle c_i \mid i \in I \rangle^{\mathcal{G}}$ , where  $\forall i \in I \ c_i = c$ , is identified with  $c$ . Thus  $A$  is considered as a subset of  ${}^*A$ . In this context the elements of  $A$  are said to be standard elements of  ${}^*A$ .

Let  $A$  be a topological space. Consider an arbitrary family  $\langle c_i \rangle$  of elements of  $A$  indexed by elements  $i \in I$ . We say that  $\lim_{\mathcal{G}} c_i = c$  if for any open  $U \ni c$  holds  $\{i \in I \mid c_i \in U\} \in \mathcal{G}$ . It is well-known (see, for example, [3]) that if  $\{c_i \mid i \in M\} \subset D$ , where  $D$  is a compact set then the  $\lim_{\mathcal{F}} c_i$  exists.

It is obvious that if  $\langle c_i \rangle \equiv_{\mathcal{G}} \langle d_i \rangle$ , i.e.  $\{i \in I \mid c_i = d_i\} \in \mathcal{G}$ , then  $\lim_{\mathcal{G}} c_i = \lim_{\mathcal{F}} d_i$  and both limits either exist or do not exist simultaneously. Thus, the notion of limit is defined for elements of  ${}^*A$ . In the language of nonstandard analysis the  $\lim_{\mathcal{G}} \gamma$ , where  $\gamma \in {}^*A$ , is said to be the standard part of  $\gamma$  or the shadow of  $\gamma$  and is denoted by  ${}^{\circ}\gamma$ . If  ${}^{\circ}\gamma$  exists then  $\gamma$  is called a nearstandard element of  ${}^*A$ . The set of all nearstandard elements is denoted by  $\text{ns}({}^*A)$ . By the discussion in previous paragraph, it is easy to see that  $\bigcup \{ {}^*C \mid C \subseteq A, C \text{ is compact} \} \subseteq \text{ns}({}^*A)$ . It is easy to see that if  $A$  is a locally compact space then the following equality holds:

$$\bigcup \{ {}^*C \mid C \subseteq A, C \text{ is compact} \} = \text{ns}({}^*A). \quad (17)$$

The nonstandard extension  ${}^*\mathcal{A}$  of a locally compact algebra  $\mathcal{A} = \langle A, \sigma \rangle$  of a finite signature  $\sigma$  is the algebra  $\langle {}^*A, \sigma \rangle$ , where any functional symbol  $f \in \sigma$  is interpreted by the operation  ${}^*f$  – the ultrapower of the  $f$  on  $A$ . If  $f \in \sigma$  is an  $n$ -ary functional symbol and  $\bar{\gamma} \in \text{ns}({}^*A^n)$  then, since  $f$  is continuous,  ${}^*f(\bar{\gamma}) \in \text{ns}({}^*A)$  and

$${}^{\circ}({}^*f(\bar{\gamma})) = f({}^{\circ}\bar{\gamma}). \quad (18)$$

Thus, the following proposition holds

**Proposition 3** *The set  $\text{ns}({}^*A)$  is closed in  ${}^*A$  under operations of the signature  $\sigma$ , i.e.  $\langle \text{ns}({}^*A), \sigma \rangle$  is a subalgebra of  ${}^*\mathcal{A}$ .*

We denote this subalgebra by  $\text{ns}({}^*\mathcal{A})$ .

Consider the following binary relation  $\approx$  on  $\text{ns}({}^*A)$ . For any  $\alpha, \beta \in \text{ns}({}^*A)$  put  $\alpha \approx \beta$  iff  ${}^{\circ}\alpha = {}^{\circ}\beta$ . By equality (18), the relation  $\approx$  is a congruence relation on the algebra  $\text{ns}({}^*\mathcal{A})$ .

**Proposition 4** *The quotient algebra  $\text{ns}({}^*\mathcal{A})/\approx$  is isomorphic to  $\mathcal{A}$*

**Proof.** Consider the map  $\varphi : \text{ns}({}^*A) \rightarrow A$  such that  $\varphi(\alpha) = {}^{\circ}\alpha$ . Obviously  $\varphi$  is a surjective homomorphism of the algebra  $\text{ns}({}^*\mathcal{A})$  onto the algebra  $\mathcal{A}$  and  $\varphi(\alpha) = \varphi(\beta)$  iff  ${}^{\circ}\alpha = {}^{\circ}\beta$   $\square$

In nonstandard analysis the ultraproducts of sets (functions, relations) are said to be internal sets (functions, relations). All theorems of classical mathematics hold for internal objects. This follows from the Los's theorem about ultraproduct and is called the Transfer Principle in the context of nonstandard analysis. For example, any bounded from above internal subset  $S \subset {}^*\mathbb{R}$  has the lowest upper bound. Indeed, let  $S = \prod_{i \in I} S_i/\mathcal{G}$ . Since  $S$  is bounded we can assume without loss of generality that all sets  $S_i$  are bounded from above. Then it is easy to see that  $\prod_{i \in I} \sup S_i/\mathcal{G}$  is the  $\sup S$  in  ${}^*\mathbb{R}$ . It is easy to see that the sets  $\mu(0) = \{\alpha \in {}^*\mathbb{R} \mid \alpha \approx 0\}$  and  $\text{ns}({}^*\mathbb{R})$  are not internal. Indeed, it is obvious that  $\text{ns}({}^*\mathbb{R})$  is bounded from above by any infinite  $\Omega$ , but it has no supremum, since it is easy to see that the  $\sup \text{ns}({}^*\mathbb{R})$  can not be either finite or infinite. A similar consideration holds for  $\mu(0)$ . The non-internal sets are called external.

The ultraproducts of finite sets are called the hyperfinite sets. The internal cardinality of a hyperfinite set  $\prod_{i \in I} B_i/\mathcal{G}$  is an element  $\prod_{i \in I} |B_m| \in {}^*\mathbb{N}$  (standard, if  $\exists N \in \mathbb{N}(\{i \in I \mid |B_i| < N\} \in \mathcal{F})$ ). In what follows we use the term "cardinality" only for internal cardinalities. We use the term "external" for classical cardinalities.

It is easy to prove that if the internal cardinality of an internal set is a standard natural number  $n$  then its external cardinality is also equal to  $n$ .

An ultraproduct  $\mathcal{B} = \prod_{i \in I} \mathcal{A}_i / \mathcal{G}$  of algebras  $\mathcal{A}_i$  of a finite signature  $\sigma$  is said to be a hyperfinite algebra of the signature  $\sigma$ . If  $\mathcal{K}$  is a class of algebras of the signature  $\sigma$  and  $\{i \in I \mid \mathcal{A}_i \in \mathcal{K}\} \in \mathcal{G}$  then, by definition,  $\mathcal{B} \in {}^*\mathcal{K}$ . We say in this case that  $\mathcal{B}$  is a  $\mathcal{K}$ -algebra.

In what follows we need the saturation properties of ultraproducts. We use the standard notation  $\lambda^+$  for the cardinal next to a cardinal  $\lambda$ . Recall (see, for example, [11]), that an ultraproduct is said to be  $\lambda^+$ -saturated if any family of cardinality  $\lambda$  of internal subsets that satisfies the finite interesection condition has the non-empty intersection. The saturation properties of ultraproducts depend only on ultrafilters and thus we can speak about the saturation properties of ultrafilters. For example, any ultraproduct over a free ultrafilter over a countable set is  $\omega^+$ -saturated i.e. any free ultrafilter over  $\omega$  is  $\omega^+$ -saturated. For any cardinal  $\lambda$  there exists a  $\lambda^+$ -saturated ultrafilter [11].

**Definition 3** Let  $\mathcal{A}_h = \langle A_h, \sigma \rangle$  is a hyperfinite algebra of a finite signature  $\sigma$ , and  $j : A_h \rightarrow {}^*A$  an internal mapping. We say that a pair  $\langle \mathcal{A}_h, j \rangle$ , is a hyperfinite approximation of a locally compact algebra  $\mathcal{A}$  of the signature  $\sigma$  if the following two conditions hold:

1.  $\forall a \in A \exists \alpha \in A_h (j(\alpha) \approx a)$ ;
2. for any  $n$ -ary functional symbol  $\varphi \in \sigma$  holds  $\forall \alpha \in (j^{-1}(ns({}^*A))^n j(\varphi_h(\alpha)) \approx {}^*\varphi(j(\alpha))$ .

Here and in the sequel  $\varphi_h$  is the interpretation of the functional symbol  $\varphi$  in  $\mathcal{A}_h$ .

**Theorem 4** A locally compact algebra  $\mathcal{A} = \langle A, \sigma \rangle$  is approximable by finite algebras of a class  $\mathcal{K}$  iff there exists a hyperfinite approximation  $\langle \mathcal{A}_h, j \rangle$  of  $\mathcal{A}$  such that  $\mathcal{A}_h = \langle A_h, \sigma \rangle \in {}^*\mathcal{K}$ .

**Proof.**  $\Leftarrow$  Let  ${}^*A = \mathcal{A}^{\mathcal{G}}$ , where  $\mathcal{G}$  is a free ultrafilter over a set  $I$ ,  $\{\mathcal{A}_i \mid i \in I\}$  - a family of finite  $\mathcal{K}$ -algebras,  $\mathcal{A}_h = \prod_{i \in I} \mathcal{A}_i / \mathcal{G}$ ,  $j = \prod_{i \in I} j_i / \mathcal{G}$ , where  $j_i : \mathcal{A}_i \rightarrow A$  for any  $i \in I$ . We assume that  $\mathcal{A}_h$  and  $j$  satisfy the conditions of Definition 3.

Consider a compact subset  $C \subset A$  and a finite covering  $\mathcal{U}$  of  $C$  by relatively compact open sets. For any  $U \in \mathcal{U}$  fix an element  $a_U \in C \cap U$ . By our assumption, there exists an element  $\alpha^U \in \mathcal{A}_h$  such that  $\varphi(j(\alpha^U)) = a_U$ , thus,  $\{i \mid j_i(\alpha_i^U) \in U\} \in \mathcal{G}$ . Hence, for any  $U \in \mathcal{U}$  the set  $I_U = \{i \mid j_i^{-1}A_i \cap U \neq \emptyset\} \in \mathcal{G}$  and, since  $\mathcal{U}$  is finite,  $I_{\mathcal{U}} = \bigcap_{U \in \mathcal{U}} I_U \in \mathcal{G}$ . Obviously, for any  $i \in I_{\mathcal{U}}$  the set  $j_i(A_i)$  is a  $\langle C, \mathcal{U} \rangle$ -grid. We assume, without loss of generality, that  $I_{\mathcal{U}} = I$ .

To prove that  $j$  is a  $\langle C, \mathcal{U} \rangle$ -homomorphism consider for simplicity a unary functional symbol  $\varphi \in \sigma$ . If the set  $\{i \in I \mid \exists a \in A_i (j_i(a), \varphi(j_i(a)) \in C)\} \notin \mathcal{G}$  then its complement is in  $\mathcal{G}$ . For the elements  $i$  of this complement the map  $j_i$  satisfies the conditions of  $\langle C, \mathcal{U} \rangle$ -homomorphism for  $\varphi$  automatically. So, we can assume, without loss of generality, that  $\forall i \in I \exists a \in A_i (j_i(a), \varphi(j_i(a)) \in C)$ . We have to show that for any  $\varphi \in \sigma$  the set

$$I_{\varphi} = \{i \in I \mid \forall a \in A_i (j_i(a) \in C \wedge \varphi(j_i(a)) \in C \longrightarrow j_i(\varphi(a)) \text{ and } \varphi(j_i(a)) \text{ are } \mathcal{U} - \text{close})\} \in \mathcal{G}. \quad (19)$$

Indeed, the condition (19) implies that  $\bigcup_{\varphi \in \sigma} I_{\varphi} \neq \emptyset$ . Obviously, for any  $i \in \bigcup_{\varphi \in \sigma} I_{\varphi} \neq \emptyset$  pair  $\langle \mathcal{A}_i, j_i \rangle$  is  $\langle C, \mathcal{U} \rangle$ -approximation of  $\mathcal{A}$ . Let  $\alpha = \langle \alpha_i \rangle^{\mathcal{G}} \in \mathcal{A}_h$ . Put  $I_{\alpha}^1 = \{i \in I \mid j_i(a_i) \in C \wedge \varphi(j_i(\alpha_i)) \in C\}$ ,  $I_{\alpha}^2 = \{i \in I \mid j_i(\varphi_i(\alpha_i)) \text{ and } \varphi(j_i(\alpha_i)) \text{ are } \mathcal{U} - \text{close}\}$ . By the Los's theorem, the condition (19) follows from the condition

$$\forall \alpha \in \mathcal{A}_h (I_{\alpha}^1 \in \mathcal{G} \longrightarrow I_{\alpha}^2 \in \mathcal{G}). \quad (20)$$

To prove (20) notice that if  $I_{\alpha}^1 \in \mathcal{G}$  then  $\{i \in I \mid j_i(\alpha_i) \in C\} \in \mathcal{G}$ . Hence the  $\varphi(j(\alpha))$  exists. Now, by definition 3, holds  $\varphi(j(\varphi_h(a))) = \varphi(j(a)) = d$ . Thus there exists  $u \in U$  such that  $\{i \mid \varphi(j_i(\alpha_i)) \in U\} \in \mathcal{G}$  and  $\{i \mid j_i(\varphi_i(\alpha_i)) \in U\} \in \mathcal{G}$ . Thus,  $\{i \mid j_i(\varphi_i(\alpha_i)) \text{ and } \varphi(j_i(\alpha_i)) \text{ are } \mathcal{U} - \text{close}\} \in \mathcal{G}$ . This proves (20).

$\implies$  consider the set  $M$  of all pairs  $\langle C, \mathcal{U} \rangle$ , where  $C \subset A$  is a compact set and  $\mathcal{U}$  is a finite r.c.o. covering of  $C$ . Consider the relation  $\leq$  on  $M$  defined by the following formula

$$\langle C_1, \mathcal{U}_1 \rangle \leq \langle C_2, \mathcal{U}_2 \rangle \iff (C_1 \supset C_2) \wedge \forall U \in \mathcal{U}_1 (U \cap C_2 \neq \emptyset \longrightarrow \exists V \in \mathcal{U}_2 (U \subset V))$$

**Lemma 1** 1. If  $\mathcal{A}'$  is a  $\langle C, \mathcal{U} \rangle$ -approximation of  $\mathcal{A}$  and  $\langle C, \mathcal{U} \rangle \leq \langle C_1, \mathcal{U}_1 \rangle$  then  $\mathcal{A}'$  is  $\langle C_1, \mathcal{U}_1 \rangle$ -approximation of  $\mathcal{A}$ ;

2. the relation  $\leq$  is a pre-ordering;

3. for any  $\langle C_1, \mathcal{U}_1 \rangle, \langle C_2, \mathcal{U}_2 \rangle \in M$  there exists a  $\langle C, \mathcal{U} \rangle \in M$  such that  $\langle C, \mathcal{U} \rangle \leq \langle C_1, \mathcal{U}_1 \rangle$  and  $\langle C, \mathcal{U} \rangle \leq \langle C_2, \mathcal{U}_2 \rangle$ .

**Proof.** The first two properties follow immediately from the definition. To prove the property 3 it is enough to put  $C = C_1 \cup C_2$  and  $\mathcal{U} = \{U \cap V \mid U \in \mathcal{U}_1, V \in \mathcal{U}_2\} \cup \{U \setminus C_2 \mid U \in \mathcal{U}_1\} \cup \{V \setminus C_1 \mid V \in \mathcal{U}_2\}$ .  $\square$

Recall that a pre-ordered set that satisfies the condition 3 of Lemma 1 is called a *directed set*.

Let now  $\lambda$  be the weight of the topological space  $A$ , i.e. the least cardinality of a base of topology of  $A$  is equal to  $\lambda$ . Then it is easy to see that the pre-ordered set  $M$  has a cofinal subset  $M_\lambda$ . Obviously,  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -algebras iff for any  $m = \langle C, \mathcal{U} \rangle \in M_\lambda$  there exists a finite  $\mathcal{K}$ -algebra  $\mathcal{A}_m$  that is an  $m$ -approximation of  $\mathcal{A}$ .

Fix a  $\lambda^+$ -saturated ultrafilter  $\mathcal{G}$ . In the rest of this proof we consider only ultraproducts over  $\mathcal{G}$ .

Let  $\mathcal{X}_m$  be the set of all finite  $m$ -approximations  $\langle \mathcal{A}_m, j_m \rangle$  of  $\mathcal{A}$  such that  $\mathcal{A}_m \in \mathcal{K}$ . Since  $\mathcal{A}$  is approximable by finite  $\mathcal{K}$ -systems,  $\mathcal{X}_m \neq \emptyset$ . By Lemma 1 (1), if  $m \leq m_1, \dots, m \leq m_k$  then  $\mathcal{X}_m \subset \mathcal{X}_{m_1} \cap \dots \cap \mathcal{X}_{m_k}$ . Thus, the family  $\{\mathcal{X}_m \mid m \in M_\omega\}$  satisfies the finite intersection condition and, by  $\lambda^+$ -saturation,  $\bigcap_{m \in M_\omega} \mathcal{X}_m \neq \emptyset$ . Let  $\langle \mathcal{A}_h, j \rangle \in \bigcap_{m \in M_\omega} \mathcal{X}_m$ . Then by the Los theorem (transfer principle),  $\mathcal{A}_h$  is hyperfinite and  $\mathcal{A}_h \in \ast\mathcal{K}$ . Let us show that  $\langle \mathcal{A}_h, j \rangle$  is a hyperfinite approximation of  $\mathcal{A}$ .

Let  $a \in A$ . Fix a base  $\mathcal{V}$  of cardinality  $\lambda$  of locally compact neighborhoods of  $a$  and consider the family of internal sets  $\mathcal{V}' = \{j(A_h) \cap \ast U \mid U \in \mathcal{V}\}$ . This family has the cardinality  $\lambda$ . For any  $U \in \mathcal{V}$  put  $m_0 = \langle \{a\}, \{U\} \rangle$ . Since  $M_\lambda$  is cofinal in  $M$ , there exists an  $m \in M_\lambda$  such that  $m \leq m_0$ . By the transfer principle  $\langle \mathcal{A}_h, j \rangle$  is an  $\ast m$ -approximation of  $\ast A$  and, thus, an  $\ast m_0$ -approximation of  $\ast A$ . By Definition 1  $j(A_h) \cap \ast U \neq \emptyset$ . Now it is easy to see that the family  $\mathcal{V}'$  satisfies the finite intersection condition. Hence this family has the non-empty intersection. This means that there exists an  $\alpha \in \mathcal{A}_h$  such that  $\forall U \in \mathcal{V} (j(\alpha) \in \ast U)$ . Thus,  $\circ j(\alpha) = a$ .

Again for simplicity, consider only a unary functional symbol  $\varphi \in \sigma$ . Let  $\alpha \in \mathcal{A}_h$  and  $\circ j(\alpha) = a$  exists. Since  $\varphi$  is a continuous function on  $A$ , there exists the  $\circ \ast \varphi(j(\alpha)) = \varphi(\circ j(\alpha)) = b \in A$ . Consider at first the case when  $a \neq b$ . Fix any  $U \ni a$  and  $V \ni b$  such that  $U \cap V = \emptyset$ . Let  $m_0 = \langle \{a, b\}, \{U, V\} \rangle$ . Then  $\langle \mathcal{A}_h, j \rangle$  is an  $\ast m_0$ -approximation of  $\ast A$  and thus,  $j(\varphi_h(\alpha)), \ast \varphi(j(\alpha)) \in \ast V$ . Since the family of  $V \ni b$  such that  $a \notin \bar{V}$  form a base of neighborhoods of  $b$ , we have  $j(\varphi_h(\alpha)) \approx \ast \varphi(j(\alpha))$ . If  $a = b$  it is enough to put  $m_0 = \langle \{a\}, \{U\} \rangle$  for each  $U \ni a$   $\square$

The following proposition is obvious (compare with Propositions 3 and 4).

**Proposition 5** Let  $\langle \mathcal{A}_h, j \rangle$  be a hyperfinite approximation of  $\mathcal{A}$ . Put  $(\mathcal{A}_h)_b = j^{-1}(ns \ast \mathcal{A})$  and  $\alpha \sim \beta$  iff  $j(\alpha) \approx j(\beta)$  for  $\alpha, \beta \in (\mathcal{A}_h)_b$ . Then  $(\mathcal{A}_h)_b$  is a subalgebra of  $\mathcal{A}_h$ , the relation  $\sim$  restricted on  $(\mathcal{A}_h)_b$  is a congruence relation and  $(\mathcal{A}_h)_b / \sim$  is isomorphic to  $\mathcal{A}$ . The isomorphism is induced by the surjective homomorphism  $\circ j : (\mathcal{A}_h)_b \rightarrow \mathcal{A}$   $\square$

Let  $\psi(x_1, \dots, x_n)$  be a formula of  $L_\sigma$  - the first order language in the signature  $\sigma$ . Denote by  $\psi_\sim$  the formula that is obtained from  $\psi$  by replacement of any atomic subformula  $t_1 = t_2$  of  $\psi$ , where  $t_1$  and  $t_2$  are terms of the signature  $\sigma$ , by  $t_1 \sim t_2$ . Denote by  $\psi_\sim^b$  the formula that is obtained from  $\psi_\sim$  by replacement of any quantifier  $Qx$  in  $\psi$  by  $Q^b x$ . We interpret  $\psi_\sim$  and  $\psi_\sim^b$  in  $\mathcal{A}_h$ . The interpretation of  $t_1 \sim t_2$  is obvious. A quantifier  $Q^b$  is interpreted by a bounded quantifier  $Qx \in (\mathcal{A}_h)_b$ . The following proposition is an easy corollary of Proposition 5.

**Proposition 6** If  $\langle \mathcal{A}_h, j \rangle$  is a hyperfinite approximation of  $\mathcal{A}$  then for any formula  $\psi(x_1, \dots, x_n) \in L_\sigma$  and any  $\alpha_1, \dots, \alpha_n \in (\mathcal{A}_h)_b$  holds

$$\mathcal{A}_h \models \psi_\sim^b(\alpha_1, \dots, \alpha_n) \iff \mathcal{A} \models \psi(\circ j(\alpha_1), \dots, \circ j(\alpha_n)).$$

**Remark 2** From the point of view of computer numerical systems discussed in Introduction Proposition 2 has the following interpretation. Usinig the language of nonstandard analysis we can consider an idealized

computer that has the hyperfinite memory. Then the numerical system for simulating the field of reals implemented in this computer is a hyperfinite algebra  $\mathbb{R}_h$  in the signature  $\sigma = \langle +, \cdot \rangle$  that is a hyperfinite approximation of  $\mathbb{R}$ .

The elements of  $(A_h)_b$  can be considered as the elements that are not too close to the boundary of the memory. It is very natural that the property "to be not too close to the boundary" is an external property: if a natural number  $n$  is "not too close to the boundary" then obviously the same is true for  $n + 1$ . Thus the induction principle fails for this property. The result of Proposition 6 shows that the first order properties of  $\mathbb{R}$  hold approximately for the computer implementation of  $\mathbb{R}$  if we deal with the elements that are not too close to the boundary of the memory. This fact seems to be very clear for those, who use computers for numerical computation. The language of nonstandard analysis makes it possible to formulate it as a rigorous mathematical theorem. In the language of classical mathematics we can consider only the approximate properties of reals that hold eventually when the memory of computers increases to infinity and the accuracy becomes more and more precise. We will show in the next section that only some restricted results can be obtained on this way.

Recall that an external subset of a  $\lambda^+$ -saturated ultrapower is called a  $\sigma$ -set (a  $\pi$ -set) if it is the union (the intersection) of a family of cardinality  $\lambda$  of internal sets<sup>1</sup>. By Proposition 5  $\mathcal{A}$  is isomorphic to the quotient algebra  $(\mathcal{A}_h)_b / \sim$ .

If the weight of the topological space  $A$  is  $\lambda$  then it is easy to see that there exists a family  $\mathcal{C}_\lambda$  of compact subsets of  $A$  such that the cardinality of  $\mathcal{C}_\gamma$  is  $\lambda$  and  $A = \bigcup \{C \mid C \in \mathcal{C}_\lambda\}$ .

It follows from (17) that in this case  $\text{ns}(*\mathcal{A}) = \bigcup \{*C \mid C \in \mathcal{C}_\lambda\}$ . Then

$$A_h = \bigcup \{j^{-1}(*C) \mid C \in \mathcal{C}_\lambda\}.$$

Thus  $(\mathcal{A}_h)_b$  is a  $\sigma$ -subalgebra of  $\mathcal{A}_h$ .

Let us assume now that the topology on  $A$  is induced by some uniformity  $\mathcal{W}$ . It is easy to see that the minimal cardinality of a base of this uniformity does not exceed the weight  $\lambda$  of topology on  $A$ . Then it is easy to see that for any  $\gamma, \delta \in \text{ns}(*A)$

$$\gamma \approx \delta \iff \forall W \in \mathcal{W} (\langle \gamma, \delta \rangle \in *W). \quad (21)$$

The condition (21) allows to extend the equivalence relation  $\approx$  defined on  $\text{ns}(*A)$  on the whole  $*A$ . Thus, the relation  $\sim$  can be extended on the whole  $\mathcal{A}_h$ : for any  $\alpha, \beta \in A_h$

$$\alpha \sim \beta \iff \forall W \in \mathcal{W} (\langle j(\alpha), j(\beta) \rangle \in *W).$$

Let  $\mathcal{W}_\lambda$  be a base of uniformity on  $A$  of the minimal cardinality. Obviously, the family  $\mathcal{W}$  can be replaced by  $\mathcal{W}_\lambda$  in (21). Thus, the relation  $\sim$  is defined by the formula:

$$\sim = \bigcap_{W \in \mathcal{W}_\lambda} (j \times j)^{-1}(W). \quad (22)$$

Hence  $\sim$  is a  $\pi$ -relation. The relation  $\sim$  has also the following important property:

$$\forall \alpha \in (A_h)_b \{ \beta \in A_h \mid \alpha \sim \beta \} \subset (A_h)_b. \quad (23)$$

In what follows the set  $\{ \beta \in A_h \mid \alpha \sim \beta \}$  will be denoted by  $\alpha^\sim$ .

The following theorem shows how a locally compact universal algebra  $\mathcal{A}$  of a finite signature can be constructed from an arbitrary  $\lambda^+$ -saturated ultrapower of finite universal algebras in  $\sigma$ .

Let  $\mathcal{A}_h$  be a  $\lambda^+$ -saturated ultraproduct of finite algebras of a finite signature  $\nu$ . Let  $(A_h)_b$  be a  $\sigma$ -subset of  $A_h$  closed under the  $\nu$ -operations in  $\mathcal{A}_h$ , i.e.  $(A_h)_b$  is a  $\sigma$ -subalgebra of  $\mathcal{A}_h$ . Let  $\sim$  be a  $\pi$ -equivalence relation on  $A_h$  such that its restriction on  $(A_h)_b$  is a congruence relation that satisfies (23).

Let  $F \subset A_h$ . Put  $i(F) = \{ \alpha \in F \mid \alpha^\sim \subset F \}$ . Denote by  $\mathcal{I}$  the family of all internal subsets of  $(A_h)_b$ . Consider the quotient system  $\hat{\mathcal{A}} = (A_h)_b / \sim$  and let  $\# : (A_h)_b \rightarrow \hat{\mathcal{A}}$  be a canonical map. Define the topology  $\mathcal{T}$  on  $\hat{\mathcal{A}}$  such that for any  $\alpha \in (A_h)_b$  the family  $\{ i(F)^\# \mid \alpha \in F \in \mathcal{I} \}$  is a base of neighborhoods of the point  $\alpha^\#$ .

<sup>1</sup>The notation  $\sigma$  is canonical and is in no sense connected with the notation  $\sigma$  that we use for the signature and that also is canonical.

**Theorem 5** 1. The weight of topology  $\mathcal{T}$  does not exceed  $\lambda$ .

2. The algebra  $\hat{\mathcal{A}}$  endowed with the topology  $\mathcal{T}$  is a topological algebra.

3. The topological algebra  $\hat{\mathcal{A}}$  is locally compact iff the following condition holds:

$$\forall F \in \mathcal{IV} \text{ internal } G \supset \sim \exists K \subset F (|K| \in \mathbb{N} \wedge F \subset \bigcup_{\alpha \in K} \{\beta \mid \langle \alpha, \beta \rangle \in G\}).$$

4. If  $\langle \mathcal{A}_h, j \rangle$  is a hyperfinite approximation of a locally compact uniform algebra  $\mathcal{A}$  and  $(\mathcal{A}_h)_b$  and  $\sim$  are defined by formulas (20) and (22) respectively then  $\mathcal{A}$  is topologically isomorphic to  $\hat{\mathcal{A}}$

A proof of this theorem for the case of locally compact groups is contained in [6]. It can be transferred without any changes on the general case.

We say that a locally compact algebra  $\mathcal{A}$  is *abstractly approximable* by finite  $\mathcal{K}$ -algebras if  $\mathcal{A}$  is topologically isomorphic to the algebra  $\hat{\mathcal{A}}$  constructed in Theorem 5, where  $\mathcal{A}_h$  is an ultraproduct of finite  $\mathcal{K}$ -algebras. Theorem 5 (4) shows that if  $\mathcal{A}$  is approximable by  $\mathcal{K}$ -algebras then it is abstractly approximable by  $\mathcal{K}$ -algebras. Is the inverse statement true? This question is open.

It is easy to see that Theorems 1 - 3 hold for abstract approximation. The Corollary 6 of Theorem 4 can be reformulated for abstract approximations also.

Let us say that that a triple  $\langle \mathcal{A}_h, (\mathcal{A}_h)_b, \sim \rangle$  is an abstract approximation of the algebra  $\mathcal{A}$  if this triple and  $\mathcal{A}$  satisfy Theorem 5.

**Proposition 7** If  $\langle \mathcal{A}_h, (\mathcal{A}_h)_b, \sim \rangle$  is an abstract approximation of  $\mathcal{A}$  then for any formula  $\psi(x_1, \dots, x_n) \in L_\sigma$  and any  $\alpha_1, \dots, \alpha_n \in (\mathcal{A}_h)_b$  holds

$$\mathcal{A}_h \models \psi_{\sim}^b(\alpha_1, \dots, \alpha_n) \iff \mathcal{A} \models \psi(\alpha_1^\#, \dots, \alpha_n^\#).$$

## 4 Positive bounded formulas in the theories of a locally compact algebra and their finite approximations

In this section we investigate the approximate versions of first order theorems about a locally compact algebra  $\mathcal{A}$  that hold in its finite approximations. Proposition 6 shows that if we use the language of nonstandard analysis then it is possible to construct an approximate version  $\varphi_{\sim}$  of any first order sentence  $\varphi$  such that  $\varphi \in \text{Th}(\mathcal{A})$  iff  $\varphi_{\sim}$  holds for any nonstandard approximation  $\langle \mathcal{A}_h, j \rangle$  of  $\mathcal{A}$ . If we use only the language of standard mathematics then the situation is more complicated: the similar result can be obtained only for some special class of formulas – the so called positive bounded formulas. We obtain this result by using Nelson’s algorithm of translation of nonstandard sentences into standard ones introduced in [15]. This algorithm is based on Nelson’s Idealization Principle [15] that does not hold for our models, but for the formulas, for which we apply this algorithm, it can be derived from  $\lambda^+$ -saturation.

In this section we assume that a locally compact topology on our algebra  $\mathcal{A}$  is induced by a uniformity  $\mathcal{W}$  on  $\mathcal{A}$  and use Definition 2 of approximation by finite algebras. Obviously all results of section 3 hold for this Definition. We have only to take for  $M$  the set of all pairs  $\langle C, W \rangle$ , where  $C$  is a compact subset of  $\mathcal{A}$  and  $W \in \mathcal{W}$ . The partial ordering  $\leq$  on  $M$  is defined as follows:  $\langle C, W \rangle \leq \langle C', W' \rangle$  if  $C \supseteq C'$  and  $W' \subseteq W$ . It is easy to see that if the weight of induced topology on  $\mathcal{A}$  is  $\lambda$  then there exists a cofinal subset  $M_\lambda \subset M$  of cardinality  $\lambda$ .

We assume that the uniform space  $\mathcal{A}$  is *uniformly locally compact*, i.e. there exist a base of uniformity  $\mathcal{W}_\lambda$  of cardinality  $\lambda$  such that for any  $W \in \mathcal{W}_\lambda$  and for any  $x \in \mathcal{A}$  the open set  $W(x)$  is relatively compact.

We say that a formula  $\varphi \in L_\sigma$  is *positive* if it can be built up from atomic formulas using only conjunction, disjunction and quantifiers. In what follows we work only with positive formulas in the prenex form

$$Q_1 y_1 \dots Q_m y_m \psi(x_1, \dots, x_n, y_1, \dots, y_m), \tag{24}$$

where  $Q_i$  are quantifiers and  $\psi$  is a disjunction of conjunctions of atomic formulas.

An arbitrary (not necessary positive) formula  $\varphi$  is equivalent to a formula in the form (24), where  $\psi$  is a disjunction of conjunctions of atomic formulas and their negations. Let  $\alpha_1, \dots, \alpha_k$  be the list of all atomic formulas and their negations involved in  $\psi$ . For any  $1 \leq i \leq k$  fix a  $W_i \in \mathcal{W}$  and denote by  $\alpha_i[W_i]$  the formula  $\langle t_1, t_2 \rangle \in W_i$  if  $\alpha_i$  is  $t_1 = t_2$  and  $\langle t_1, t_2 \rangle \notin W_i$  if  $\alpha_i$  is  $\neg(t_1 = t_2)$ . Here  $t_1$  and  $t_2$  are terms of the signature  $\sigma$ .

The interpretation of  $\langle t_1, t_2 \rangle \in W$  in  $\mathcal{A}$  is obvious. If  $\langle \mathcal{A}_f, j_f \rangle$  is a  $\langle C', W' \rangle$ -approximation for some  $\langle C', W' \rangle \in M$  then  $\mathcal{A}_f \models \langle t_1, t_2 \rangle \in W$  iff  $\langle j_f(t_1), j_f(t_2) \rangle \in W$ . Denote by  $\varphi[W_1, \dots, W_k]$  the formula that is obtained from  $\varphi$  by replacement of all  $\alpha_i$  by  $\alpha_i[W_i]$  respectively. The formula  $\varphi[W_1, \dots, W_k]$  is called an approximation of  $\varphi$ . Obviously, if  $\varphi$  is positive, then for any  $W_1, \dots, W_k \in \mathcal{W}$  holds  $\varphi \implies \varphi[W_1, \dots, W_k]$  (for both interpretations). This is wrong for non-positive formulas. Similarly if  $W'_i \subseteq W_i$ ,  $i = 1, \dots, k$  then  $\varphi[W'_1, \dots, W'_k] \implies \varphi[W_1, \dots, W_k]$  for positive  $\varphi$  and this is wrong for non-positive  $\varphi$ . For positive formulas we say in this case that  $\varphi[W'_1, \dots, W'_k]$  is finer approximation than  $\varphi[W_1, \dots, W_k]$ . Obviously, for any approximation  $\varphi[W_1, \dots, W_k]$  of a positive formula  $\varphi$  there exists a finer approximation  $\varphi[W'_1, \dots, W'_k]$  such that  $W'_1 = \dots = W'_k = W$  (it is enough to put  $W = W_1 \cap \dots \cap W_k$ ). In this case we write  $\varphi[W]$  instead of  $\varphi[W, \dots, W]$ . In what follows we deal only with approximations of the form  $\varphi[W]$  of a positive formula  $\varphi$ .

If  $B \subseteq A$  and  $Q$  is either  $\forall$  or  $\exists$  then  $Q_B x \dots$  is interpreted in  $\mathcal{A}$  by  $Qx(x \in B \dots)$  and in a finite  $\langle C, W \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathcal{A}$  - by  $Qx(x \in j_f^{-1}(B) \dots)$ .

The quantifiers of the form  $Q_B$  are called bounded quantifiers. If all quantifiers in a formula  $\varphi$  are bounded then we say that  $\varphi$  is bounded.

Let  $c = \langle C_1, \dots, C_n \rangle$  be an  $n$ -tuple of the subsets of  $A$  and  $\varphi$  is a formula in the form (24). Then  $\varphi[c]$  is the formula

$$Q_1 c_1 y_1 \dots Q_m c_n y_m \psi.$$

In what follows we consider only positive bounded formulas  $\varphi[c]$  that satisfy the following condition: for any  $i \leq m$  such that  $Q_i = \forall$  ( $Q_i = \exists$ ) the set  $C_i$  is a relatively compact open (compact) set.

In this case we say that an  $m$ -tuple  $c$  of subsets of  $A$  is  $\varphi$ -regular.

For two  $\varphi$ -regular  $m$ -tuples  $c$  and  $c'$  we say that  $c \ll c'$  if for any  $i \leq m$  the following property holds:

if  $Q_i = \forall$  then  $\overline{C'_i} \subseteq C_i$  and if  $Q_i = \exists$  then  $C_i \subseteq \text{int}(C'_i)$ . Here  $\overline{B}$  is the closure of  $B$  and  $\text{int}(B)$  - the interior of  $B$ .

If  $c' \ll c$  and  $W \in \mathcal{W}$  then the formula  $\varphi[c'][W]$  is called a strong approximation of  $\varphi[c]$ . The following lemma is obvious.

**Lemma 2** *Let  $\varphi(x_1, \dots, x_n)$  be a positive formula of  $L_\sigma$  in the form (24),  $c_2 \ll c_1$ , -  $\varphi$ -regular  $m$ -tuples of subsets of  $A$ ,  $W_2 \subseteq W_1$  - two elements of the uniformity  $\mathcal{W}$ ,  $\langle \mathcal{A}_f, j_f \rangle$  - a  $\langle C, W \rangle$ -approximation of  $\mathcal{A}$ . Then*

1.  $\forall a_1, \dots, a_n \in A_f (\mathcal{A}_f \models \varphi[c_2][W_2](a_1, \dots, a_n) \implies \mathcal{A}_f \models \varphi[c_1][W_2](a_1, \dots, a_n));$
2.  $\forall a_1, \dots, a_n \in A_f (\mathcal{A}_f \models \varphi[c_1][W_2](a_1, \dots, a_n) \implies \mathcal{A}_f \models \varphi[c_1][W_1](a_1, \dots, a_n));$
3.  $\forall a_1, \dots, a_n \in A (\mathcal{A} \models \varphi[c_2](a_1, \dots, a_n) \implies \mathcal{A} \models \varphi[c_1](a_1, \dots, a_n))$

The following theorem is the main result of this section.

**Theorem 6** *Let  $\varphi[c](x_1, \dots, x_n)$  be a positive bounded formula and  $a_1, \dots, a_n \in A$ . Then  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$  iff for any strong approximation  $\varphi[c'][W']$  of  $\varphi[c]$  there exists a pair  $\langle C_0, W_0 \rangle \in M$  such that the following conditions hold:*

- 1)  $\bigcup_{i=1}^n W_0(a_i) \subseteq C_0;$
- 2) for any  $\langle C, W \rangle \leq \langle C_0, W_0 \rangle$ , for any  $\langle C, W \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathcal{A}$ , and for any  $b_1, \dots, b_n \in A_f$  such that  $\langle a_i, j(b_i) \rangle \in W_0$ ,  $i = 1, \dots, n$ , holds  $\mathcal{A}' \models \varphi[c'][W'](b_1, \dots, b_n)$ .

If for some property P there exists a  $\langle C_0, W_0 \rangle \in M$  such that P holds for all  $\langle C, W \rangle$ -approximations of  $\mathcal{A}$  such that  $\langle C, W \rangle \leq \langle C_0, W_0 \rangle$ , then we say that P holds for all precise enough approximations of  $\mathcal{A}$ .

**Corollary 1** *A positive bounded sentence  $\varphi[c]$  holds in  $\mathcal{A}$  iff any its strong approximation  $\varphi[c'][W]$  holds in all precise enough approximations of  $\mathcal{A}$*

From the point of view of numerical systems implemented in computers this corollary means that approximate versions of positive bounded theorems about the reals hold for numerical computer systems that simulate the field of reals in powerful enough computers.

**Example 1** Fix any positive  $d > 1$ . Then the following positive bounded formula holds for the field  $\mathbb{R}$ :

$$\forall_{(d^{-1} < |x| < d)} x \exists_{(d^{-1} \leq |y| \leq d)} y (xy = 1).$$

It is easy to see that for any strong approximation of this formula there exists a finer strong approximation of the following form:

$$\forall_{(c^{-1} < |x| < c)} x \exists_{(b^{-1} \leq |y| \leq b)} y (|xy - 1| < \delta), \quad (25)$$

where  $1 < c < d < b$  and  $\delta > 0$ .

We have to show that there exist  $a_0, \varepsilon_0$  such that for any  $a > a_0$ ,  $\varepsilon < \varepsilon_0$  formula (25) holds for any finite  $\langle a, \varepsilon \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathbb{R}$  (the definition of  $\langle a, \varepsilon \rangle$ -approximation see before the example 1 of section 2). We assume for simplicity, without loss of generality, that the constant symbol 1 is included in the signature  $\sigma$  and that  $j_f(1) = 1$ . Fix any  $x$  such that  $c^{-1} < |x| < c$  and let  $y = x^{-1}$ ,  $b^{-1} < |y| < b$ . Take  $\xi, \eta \in A_f$  such that  $|x - j_f(\xi)| < \varepsilon$  and  $|y - j_f(\eta)| < \varepsilon$ . The  $a$  and  $\varepsilon$  have to satisfy the following conditions:  $j_f(\xi), j_f(\eta), j_f(\xi) \cdot j_f(\eta) \in [-a, a]$ ,  $|j_f(\xi \odot \eta) - 1| < \delta$ , where  $\odot$  is the multiplication in  $\mathcal{A}_f$ . To satisfy the first two conditions we have to take  $a$  such that  $b + \varepsilon < a$ . Since  $xy = 1$  we have  $|1 - j_f(\xi) \cdot j_f(\eta)| < (2b + \varepsilon)\varepsilon$  and thus  $(2b + \varepsilon)\varepsilon + 1 < a$ . By the definition of  $\langle a, \varepsilon \rangle$ -approximation we have under this conditions  $|j_f(\xi) \cdot j_f(\eta) - j_f(\xi \odot \eta)| < \varepsilon$ , thus  $|1 - j_f(\xi \odot \eta)| < (2b + 1 + \varepsilon)\varepsilon$  and thus  $(2b + 1 + \varepsilon)\varepsilon < \delta$ . So

$$\varepsilon_0 = \sqrt{\left(\frac{2b+1}{2}\right)^2 + \delta} - \frac{2b+1}{2},$$

and

$$a_0 = \max\{b + \varepsilon_0, (2b + \varepsilon_0)\varepsilon_0 + 1\}$$

**Example 2** The relation  $\leq$  is defined in  $\mathbb{R}$  by positive formula

$$x \leq y \iff \exists z (x + z^2 = y) = \varphi(x, y)$$

Consider a bounded version of this formula  $\varphi[b](x, y) = \exists_{|z| \leq b} (x + z^2 = y)$ , which defines the relation  $x \leq y \leq x + b^2$ . A strong approximation of this formula is of the form  $\varphi[c][\alpha](x, y) = \exists_{|z| \leq c} (|x + z^2 - y| < \alpha)$  for some  $\alpha > 0$  and  $0 < b < c$ . Let  $x_0, y_0 \in [-d, d]$ . Put  $a_0 = (c + \alpha)^2 + d + \alpha + 1$  and  $\varepsilon_0 = \max\{c - b, \frac{\alpha}{5 + 2a_0}\}$ . Then it is easy to see that for any  $a > a_0$ ,  $\varepsilon < \varepsilon_0$  for any  $\langle a, \varepsilon \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathbb{R}$  and for any  $\xi, \eta \in A_f$  such that  $|x_0 - j_f(\xi)| < \varepsilon_0$ ,  $|y_0 - j_f(\eta)| < \varepsilon_0$  holds  $\mathcal{A}_f \models \varphi[c][\alpha](j_f(\xi), j_f(\eta))$ . If  $x_0 > y_0$  and  $\alpha < \frac{1}{2}(x_0 - y_0)$  then, obviously, there exists  $\varepsilon_0$  such that  $(x_0 - \varepsilon_0) > y_0 + \alpha$ . Then for any  $\langle a, \varepsilon \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathbb{R}$  for any  $\xi, \eta \in A_f$  such that  $|x_0 - j_f(\xi)| < \varepsilon_0$ ,  $|y_0 - j_f(\eta)| < \varepsilon_0$  the formula  $\varphi[c][\alpha](j_f(\xi), j_f(\eta))$  fails in  $\mathcal{A}_f$ . A similar consideration holds for  $y_0 > x_0 + b^2$ .

**Example 3** The relation  $<$  also can be defined by positive formula. Indeed:

$$x < y \iff \exists z ((y - x)z^2 = 1) = \varphi(x, y)$$

A bounded version of this formula  $\varphi[b](x, y) = \exists_{|z| \leq b} ((y - x)z^2 = 1)$  defines the relation  $y > x + \frac{1}{b^2}$ . A strong approximation of this formula is of the form  $\varphi[c][\alpha](x, y) = \exists_{|z| \leq c} (|(y - x)z^2 - 1| < \alpha)$  for some  $\alpha > 0$  and  $0 < b < c$ . It is easy to see that for  $\alpha < 1$ , for small enough  $\varepsilon$ , big enough  $a$  and for any  $\langle a, \varepsilon \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathbb{R}$  if  $j_f(\xi), j_f(\eta) \in [-a, a]$  then  $\mathcal{A}_f \models \varphi[c][\alpha](\xi, \eta) \iff j_f(\eta) > j_f(\xi) + \frac{1 - \alpha}{c^2}$ .

**Remark 3** Since in  $Th(\mathbb{R})$  any formula is equivalent to a quantifier free formula the considered examples show that any formula of the theory of rings is equivalent in  $Th(\mathbb{R})$  to a positive formula and thus has its approximate version.

If the topological space  $A$  is totally disconnected, i.e. the clopen sets form a base of its topology then by Lemma 2 (2) it is enough to consider only positive bounded formulas  $\varphi[c]$  with an  $n$ -tuple  $c$  that consists of clopen sets. In this case we say that  $c$  is clopen. Since for a clopen set  $V$  holds  $\overline{V} \subseteq V$  and  $V \subseteq \text{int}(V)$  then for a clopen  $n$ -tuple  $c$  holds  $c \ll c$  and thus if  $W \in \mathcal{W}$  then  $\varphi[c][W]$  is a strong approximation of  $\varphi[c]$ . So the formulation of Theorem 6 can be simplified for this case.

**Corollary 2** Let  $\mathcal{A}$  be a totally disconnected algebra,  $\varphi[c](x_1, \dots, x_n)$  be a positive bounded formula (24) with a clopen  $m$ -tuple  $c, a_1, \dots, a_n \in A$ . Then  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$  iff for any  $W' \in \mathcal{W}$  of there exists a pair  $\langle C_0, W_0 \rangle \in M$  such that the following conditions hold:

$$1) \bigcup_{i=1}^n W_0(a_i) \subseteq C_0;$$

2) for any  $\langle C, W \rangle \leq \langle C_0, W_0 \rangle$ , for any  $(C, W)$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$  of  $\mathcal{A}$ , and for any  $b_1, \dots, b_n \in A_f$  such that  $\langle a_i, j(b_i) \rangle \in W_0$ ,  $i = 1, \dots, n$ , holds  $\mathcal{A}' \models \varphi[c][W'](b_1, \dots, b_n)$ .

To prove Theorem 6 we consider at first its nonstandard version.

Let  $\langle \mathcal{A}_h, j \rangle$  be a hyperfinite approximation of  $\mathcal{A}$  in the sense of Theorem 4. Then a strong approximation  $\varphi[c][W]$  of a positive formula  $\varphi$  in the form (24) has an obvious interpretation in  $\mathcal{A}_h$ : a quantifier  $Q_C x \dots$  is interpreted by  $Q_x(j(x) \in {}^*C \dots)$  and a formula  $\langle t_1, t_2 \rangle \in W$  - by  $\langle j(t_1), j(t_2) \rangle \in {}^*W$ . Obviously, the statements (1) and (2) of Lemma 2 hold for the hyperfinite approximations of  $\mathcal{A}$ . We need also the formulas of the form  $\varphi[c]_{\sim}$  (The definition of  $\psi_{\sim}$  for any formula  $\psi \in L_{\sigma}$  see on page 11).

**Lemma 3** For any  $\beta_1, \dots, \beta_n \in (A_h)_b$  holds  $\mathcal{A}_h \models \varphi[c]_{\sim}(\beta_1, \dots, \beta_n) \iff \forall W \in \mathcal{W} \mathcal{A}_h \models \varphi[c][W](\beta_1, \dots, \beta_n)$ .

**Proof** Obviously,  $\mathcal{A}_h \models \varphi[c]_{\sim}(\beta_1, \dots, \beta_n) \implies \forall W \in \mathcal{W} \mathcal{A}_h \models \varphi[c][W](\beta_1, \dots, \beta_n)$ . So we have to prove only the inverse implication. Consider at first the case of quantifier free formula, i.e. the case when  $\varphi = \psi$  in the form (24). We have  $\psi = P_1 \vee \dots \vee P_r$ , where each  $P_i$  is a conjunction of atomic formulas. Assume that  $\forall W \in \mathcal{W}_{\lambda}$  holds  $\psi[W]$ . If  $\psi_{\sim}$  is false then for each  $i \leq r$  there exists  $W_i \in \mathcal{W}_{\lambda}$  such that  $P_i[W_i]$  is false. Take  $W \in \mathcal{W}_{\lambda}$  such that  $W \subseteq \bigcap_{i=1}^r W_i$ . Then by Lemma 2 (2) for any  $i \leq r$  the formula  $P_i[W]$  is false and, thus, the formula  $\psi[W]$  is false.

We have to prove now that

$$\forall W \in \mathcal{W}_{\lambda} Q_{1C_1} y_1 \dots Q_{mC_m} \psi[W] \implies Q_{1C_1} y_1 \dots Q_{mC_m} \forall W \in \mathcal{W}_{\lambda} \psi[W]$$

To prove this implication, it is enough to prove that for any positive bounded formula  $\theta(x)$  and a compact set  $C$  holds

$$\forall W \in \mathcal{W}_{\lambda} \exists C x \theta[W](x) \implies \exists C x \forall W \in \mathcal{W}_{\lambda} \theta[W](x). \quad (26)$$

Assume that the left hand side of this implication holds. Put  $B(W) = \{x \mid j(x) \in {}^*C, \theta[W](x)\}$ . Then  $B[W] \neq \emptyset$  and since for any  $W_1, \dots, W_s \in \mathcal{W}_{\lambda}$  there exists  $W \in \mathcal{W}_{\lambda}$  such that  $W \subseteq \bigcap_{i=1}^s W_i$ , using Lemma 2 (2), we obtain that the family  $\{B(W) \mid W \in \mathcal{W}_{\lambda}\}$  has the finite intersection property. Thus, by saturation, we obtain that the right hand side of the implication (26) holds  $\square$

**Lemma 4** Let  $\varphi(x_1, \dots, x_n)$  be a positive formula in  $L_{\sigma}$  of the form (24),  $c = \langle C_1, \dots, C_m \rangle$  -  $\varphi$ -regular  $m$ -tuple of subsets of  $A$ ,  $a_1, \dots, a_n \in A$ ,  $\langle \mathcal{A}_h, j \rangle$  - a hyperfinite approximation of  $\mathcal{A}$ . Then  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$  iff for any  $\varphi$ -regular  $c' = \langle C'_1, \dots, C'_m \rangle$  such that  $c' \gg c$  and for any  $\alpha_1, \dots, \alpha_n \in A_h$  such that  $j(\alpha_i) \approx a_i$ ,  $i = 1, \dots, n$  holds  $\mathcal{A}_h \models \varphi[c']_{\sim}(\alpha_1, \dots, \alpha_n)$ .

**Proof** We prove this lemma by induction over  $m$ . For  $m = 0$  it follows from Proposition 6. Assume that it is proved for  $m - 1$ . Denote by  $\theta(y_1, x_1, \dots, x_n)$  the formula  $Q_2 y_2 \dots Q_m y_m \psi(x_1, \dots, x_n, y_1, \dots, y_m)$ , by  $c_{-}$  - the  $(m - 1)$ -tuple  $\langle C_2, \dots, C_m \rangle$ , by  $c'_{-}$  - the  $(m - 1)$ -tuple  $\langle C'_2, \dots, C'_m \rangle$  so that  $\varphi = Q_1 y_1 \theta$ ,  $\varphi[c] = Q_{1C_1} \theta$ ,  $\varphi[c'] = Q_{1C'_1} \theta$ . Consider two cases.

a).  $Q_1 = \exists$ . In this case  $C_1, C'_1$  are compact sets and  $C_1 \subseteq \text{int} C'_1$ .

$\implies$  Let  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$ . Then there exists  $b \in C_1$  such that  $\mathcal{A} \models \theta(b, a_1, \dots, a_n)$ . Let  $\beta \in A_h$  be such that  $j(\beta) \approx b$ . Then, by the induction assumption,  $\mathcal{A}_h \models \theta[c'_{-}]_{\sim}(\beta, \alpha_1, \dots, \alpha_n)$ . Since  $b \in C_1 \subseteq \text{int}(C'_1)$ ,  $j(\beta) \approx b$  and  $\text{int}(C'_1)$  is an open set, we have  $j(\beta) \in \text{int}({}^*C'_1)$ . This proves that  $\mathcal{A}_h \models \varphi[c']_{\sim}(\alpha_1, \dots, \alpha_n)$ .

$\impliedby$ . Obviously  $C = \bigcap \{ \overline{W(C)} \mid W \in \mathcal{W}_{\lambda} \}$ . Fix any  $V, W \in \mathcal{W}_{\lambda}$ . By the induction assumption for any  $\theta$ -regular  $(m - 1)$ -tuple  $c'_{-}$  holds  $\mathcal{A}_h \models Q_{1\overline{W(C)}} y_1 \theta[c'_{-}]_{\sim} \alpha_1, \dots, \alpha_n$ . Thus, by Lemma 3, we obtain that  $B(W, V, c'_{-}) = \{ \beta \in j^{-1}(\overline{{}^*W(C)}) \mid \mathcal{A}_h \models \theta[c'_{-}][V](\beta, \alpha_1, \dots, \alpha_n) \} \neq \emptyset$ . Let  $\Xi$  be the set of all  $\theta$ -regular

$(m-1)$ -tuples  $c'_-$ . Then it is easy to see that there exists a cofinal subset  $\Xi_\lambda$  (i.e.  $\forall d \in \Xi \exists d_1 \in \Xi_\lambda (d_1 \ll d)$ ) of cardinality  $\lambda$ . By Lemma 2 (3), if for some  $b, \alpha_1, \dots, \alpha_n \in (A_h)_b$  for all  $d \in \Xi_\lambda$  holds  $\mathcal{A}_h \models \theta[d][V]$  then the same holds for all  $d \in \Xi$ . It is easy to see also that similar to  $\mathcal{W}_\lambda$  the family  $\Xi_\lambda$  has the following property: for any  $c^{(1)}, \dots, c^{(s)} \in \Xi_\lambda$  there exists a  $c'_- \in \Xi_\lambda$  such that  $c'_- \ll c^{(1)}, \dots, c'_- \ll c^{(s)}$ . All this shows that the family  $\{B(W, V, c'_-) \mid V, W \in \mathcal{W}_\lambda, c'_- \in \Xi_\lambda\}$  has the finite intersection property and thus, by saturation, the nonempty intersection. By our construction and Lemma 3, any element  $b$  in this intersection has the following properties:  $j(b) \in {}^*C$  and  $\mathcal{A}_h \models \theta[c'_-]_{\sim}(\beta, \alpha_1, \dots, \alpha_n)$  for any  $c'_- \in \Xi$ . By the induction assumption this implies that  $\mathcal{A} \models \theta[c_-](\circ j(\beta), a_1, \dots, a_n)$ . Since  $C$  is a compact set, we obtain that  $\circ j(\beta) \in C$ . This proves a).

b)  $Q_1 = \forall$ . In this case  $C_1$  and  $C'_1$  are relatively compact open sets and  $\overline{C'_1} \subseteq C_1$

$\implies$  Let  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$  and  $\alpha_1, \dots, \alpha_n \in (A_h)_b$  be such that  $j(\alpha_i) \approx a_i$ . Take any  $\beta \in A_h$  such that  $j(\beta) \in {}^*C'_1$ . Then  $\circ j(\beta) \in C$  and, thus,  $\mathcal{A} \models \theta[c_-](j(\beta), a_1, \dots, a_n)$ . By the induction assumption,  $\mathcal{A}_h \models \theta[c'_-]_{\sim}(\beta, \alpha_1, \dots, \alpha_n)$ . This completes the prove.

$\impliedby$ . Let  $b \in C_1$ . Then, obviously, there exists an open  $D$  such that  $\overline{D} \subseteq C_1$  and  $b \in D$ . Since for any  $c'_- \gg c_-$  and for any  $\alpha_1, \dots, \alpha_n \in (A_h)_b$  such that  $j(\alpha_i) \approx a_i$  holds  $\mathcal{A}_h \models \forall_{D y_1} \theta[c'_-]_{\sim}(y_1, \alpha_1, \dots, \alpha_n)$ , we obtain that for any  $\beta \in A_h$  such that  $j(\beta) \approx b$  holds  $\mathcal{A}_h \models \theta[c'_-]_{\sim}(\beta, \alpha_1, \dots, \alpha_n)$ . Thus, by the induction assumption  $\mathcal{A} \models \theta[c](b, a_1, \dots, a_n)$   $\square$ .

We say that a pair  $\langle C, W \rangle \in {}^*M_\lambda$  is infinitesimal if for any  $\langle D, V \rangle \in M_\lambda$  holds  $\langle C, W \rangle \leq \langle {}^*D, {}^*V \rangle$ . Obviously, if  $\langle C, W \rangle \in {}^*M_\lambda$  is infinitesimal then  $C \supset \text{ns}({}^*A)$ . The following lemma is an immediate corollary of  $\lambda$ -saturation.

**Lemma 5** *Let  $N \subseteq {}^*M_\lambda$  be an internal subset. Then*

1. *If  $\forall \langle D, V \rangle \in M_\lambda \langle {}^*D, {}^*V \rangle \in N$  then there exists an infinitesimal element  $\langle C, W \rangle \in N$ ;*
2. *if  $N$  contains all infinitesimal elements of  ${}^*M_\lambda$  then there exists a  $m = \langle D, V \rangle \in M_\lambda$  such that  ${}^*M_m \cap {}^*M_\lambda \subseteq N$*

**Lemma 6** *A pair  $\langle \mathcal{A}_h, j \rangle$  is a hyperfinite approximation of the algebra  $\mathcal{A}$  iff  $\langle \mathcal{A}_h, j \rangle$  is a  $\langle C, W \rangle$ -approximation of  ${}^*A$  for some infinitesimal  $\langle C, W \rangle \in {}^*M_\lambda$*

**Proof**  $\impliedby$  Let  $a \in A$ . Since  $j(A_h)$  is a  $\langle C, W \rangle$ -grid, there exists  $b \in A_h$  such that  $\langle a, j(b) \rangle \in W$ . Thus for any  $V \in \mathcal{W}$  holds  $\langle a, j(b) \rangle \in {}^*V$ , i.e.  $j(b) \approx a$ . Let  $\varphi \in \sigma$  be a unary (for simplicity) functional symbol and  $\varphi_h$  its interpretation in  $\mathcal{A}_h$ ,  $a \in (A_h)_b$ , i.e.  $j(h) \in \text{ns}({}^*A) \subset C$ . Since  $\varphi$  is a continuous function in  $\mathcal{A}$ , we have  ${}^*\varphi(j(a)) \in \text{ns}({}^*A)$ . Hence  $\varphi(j(a)) \in C$ . By Definition 2 and the transfer principle,  $\langle j(\varphi_h(a)), {}^*\varphi(j(a)) \rangle \in W$ . Thus  $j(\varphi_h(a)) \approx {}^*\varphi(j(a))$ . So  $\langle \mathcal{A}_h, j \rangle$  is a hyperfinite approximation of  $\mathcal{A}$ .

$\implies$  Let  $\langle \mathcal{A}_h, j \rangle$  be a hyperfinite approximation of  $\mathcal{A}$ . Then it is easy to see that for any  $\langle D, V \rangle \in M_\lambda$  the pair  $\langle \mathcal{A}_h, j \rangle$  is a  $\langle {}^*D, {}^*V \rangle$ -approximation of  ${}^*A$ . By Lemma 5 (1), there exists an infinitesimal  $\langle C, W \rangle$  such that  $\langle \mathcal{A}_h, j \rangle$  is a  $\langle C, W \rangle$ -approximation of  ${}^*A$   $\square$ .

**Proof of Theorem 6**  $\implies$  Let  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$ . Fix a strong approximation  $\varphi[c'][W'](x_1, \dots, x_n)$  of  $\varphi[c](x_1, \dots, x_n)$ . Consider the internal set  $N$  of all pairs  $\langle C_0, W_0 \rangle \in {}^*M_\lambda$  such that for all  $\langle C, W \rangle \leq \langle C_0, W_0 \rangle$  for any  $\langle C, W \rangle$ -approximation  $\langle \mathcal{A}_h, j \rangle$  of  ${}^*A$  and for any  $\alpha_1, \dots, \alpha_n \in A_h$  such that  $\langle a_i, j(\alpha_i) \rangle \in W_0$  holds  $\mathcal{A}_h \models \varphi[c'][W'](\alpha_1, \dots, \alpha_n)$ . Lemmas 3, 4 and 6 imply that  $N$  contains all infinitesimal pairs  $\langle C_0, W_0 \rangle \in {}^*M_\lambda$ . By Lemma 5 (2), there exists  $\langle C_0, W_0 \rangle \in M_\lambda$  such that  ${}^*C_0, {}^*W_0 \in N$ . By the transfer principle, this completes the proof.

$\impliedby$ . Let  $c' \gg c$ ,  $\langle \mathcal{A}_h, j \rangle$  be a hyperfinite approximation of  $\mathcal{A}$  and  $\alpha_1, \dots, \alpha_n \in A_h$  be such that  $j(\alpha_i) \approx a_i$ . Then for any  $\langle C, W \rangle \in M_\lambda$  the pair  $\langle \mathcal{A}_h, j \rangle$  is a  $\langle {}^*C, {}^*W \rangle$ -approximation of  ${}^*A$  and  $\langle j(\alpha_i), a_i \rangle \in {}^*W$ ,  $i = 1, \dots, n$ . Thus, by the conditions of the theorem,  $\mathcal{A}_h \models \varphi[c'][W'](\alpha_1, \dots, \alpha_n)$  for any  $W' \in \mathcal{W}_\lambda$ . By Lemma 3,  $\mathcal{A}_h \models \varphi[c]_{\sim}(\alpha_1, \dots, \alpha_n)$ . Hence, by Lemma 4 holds  $\mathcal{A} \models \varphi[c](a_1, \dots, a_n)$   $\square$

The following corollary of Theorem 6 shows that approximations of continuous functions on the closed intervals holds for all precise enough approximations of the field  $\mathbb{R}$  (cf. the example concerning  $\sin x$  which was discussed in the Introduction).

**Corollary 3** *Let  $\mathcal{A} = \langle \mathbb{R}, \sigma \rangle$  be such that  $\sigma$  contains the symbols  $+$  and  $\cdot$  and a unary functional symbol  $g$ . Suppose that the continuous function  $g$  is approximable on an interval  $[-d, d]$  by a polynomial  $b_n x^n + \dots + b_1 x + b_0$  with an accuracy  $\delta$ . Then for any  $0 < d' < d$  and for any  $\delta' > \delta$  there exist  $a_0, \varepsilon_0 > 0$*

such that for any  $a > a_0$ ,  $\varepsilon < \varepsilon_0$  for any  $\langle a, \varepsilon \rangle$ -approximation  $\langle \mathcal{A}_f, j_f \rangle$ , for any  $\beta_0, \dots, \beta_n \in A_f$  such that  $|j(\beta_i) - b_i| < \varepsilon_0$ ,  $i = 0, \dots, n$ , and for any  $\xi \in A_f$  such that  $j(\xi) \in [-d', d']$  holds

$$|j_f(g_f(\xi)) - j_f(\beta_n \xi^n + \dots + \beta_1 \xi + \beta_0)| < \delta',$$

where  $g_f$  is the interpretation of the symbol  $g$  in  $\mathcal{A}_f$ .

## References

- [1] Alekseev M.A., Glebskii L.Yu., Gordon E.I. On approximations of groups, group actions and Hopf algebras. *Representation Theory, Dynamical Systems, Combinatorial and Algebraic Methods. III*, A.M.Vershik editor, Russian Academy of Science. St.Petersburg Branch of V.A.Steklov's Mathematical Institute. Zapiski nauchnih seminarov POMI 256 (1999), 224-262. (in Russian; Engl. Transl. in *Journal of Mathematical Sciences*, 107, No.5 (2001), pp.4305-4332)
- [2] Albeverio S., Gordon E., Khrennikov A. (2001) Finite dimensional approximations of operators in the spaces of functions on locally compact abelian groups. *Acta Applicandae Mathematicae*, 64 (2000), no. 1, 33-73
- [3] N. Bourbaki (1966), *General topology*, Part 1, Hermann, Paris and Addison-Wesley, Reading, Mass.
- [4] Digernes T., Husstad E., Varadarajan V. (1999) Finite Approximations of Weyl Systems. *Math. Scand.* Vol. 84, pp. 261-283.
- [5] Glebsky L.Yu, Gordon E.I. (2002) On approximation of topological groups by finite algebraic systems. *Submitted to the Illinois Journal of Mathematics*. arXiv:math.GR/0201101
- [6] Gordon E. (1997) *Nonstandard Methods in Commutative Harmonic Analysis* AMS, Providence, Rhode Island.
- [7] E.I. Gordon, O.A. Rezvova. On hyperfinite approximations of the field  $\mathbb{R}$ . *Reuniting the Antipodes - Constructive and Nonstandard Views of the Continuum, Proceedings of the Symposium in San Servolo/Venice, Italy, May 17 20, 2000*. B.Ulrich, H.Ossvald and P. Schuster, editors. Synthèse Library, volume 306 By Kluwer Academic Publishers, Dordrecht, etc., 2001.
- [8] Henson C.W. Nonstandard Hulls of Banach Spaces (1976) *Israel Journal of Mathematics* Vol. 25, pp. 108-1440
- [9] Heinrich S., Henson C.W. (1986) Banach space model theory II. Isomorphic equivalence. *Math.Nachr.* Vol. 125, pp. 301 - 317.
- [10] Henson C.W., Moore L.C. (1983) Nonstandard Analysis and the theory of Banach spaces. *Nonstandard analysis - recent developments. Ed. A.Hurd*. Lecture Notes in Math. **983**. Springer: Berlin - Heidelberg - New-York.
- [11] Chang C.C., Keisler H.J. (1990) *Model theory*. Third edition. Studies in Logic and the Foundations of Mathematics, 73. North-Holland Publishing Co., Amsterdam.
- [12] Knut D. (1982) *Art of Computer Programming*. V.II. Addison Wesley Pub. Co
- [13] Loomis L. (1953) *An introduction to abstract harmonic analysis*. D. Van Nostrand Company, Inc., Toronto-New York-London.
- [14] Mal'tsev A.I. (1970) *Algebraic systems*. Moscow. (in Russian)
- [15] Nelson E. Internal Set Theory. A New Approach to Nonstandard Analysis, *Bull. Amer. Math. Soc.*, vol. 83 (1977), pp. 1165 - 1198.

Instituto de Investigacion en Comunicacion Optica de Universidad Autonoma de San Luis Potosi,  
Mexico

Eastern Illinois University, USA  
University of Illinois at Urbana-Champaign, USA

IICO-UASLP  
AvKarakorum 1470  
Lomas 4ta Session  
SanLuis Potosi SLP 7820  
Mexico  
Phone: 52-444-825-0892 (ext. 120)  
e-mail:glebsky@cactus.iico.uaslp.mx

Department of Mathematics and Computer Science  
Eastern Illinois University  
600 Lincoln Avenue  
Charleston, IL 61920-3099  
USA  
Phone: 1-217-581-6282  
e-mail: cfyig@eiu.edu

Department of Mathematics  
University of Illinois at Urbana-Champaign  
1409 West Green Street  
Urbana, IL 61801  
USA  
Phone: 1-217-333-2768  
e-mail: henson@math.uiuc.edu

1991 *Mathematics Subject Classification*. Primary 26E35, 03H05; Secondary 28E05, 42A38