

Order units in a C^* -algebra

ANIL K KARN

Department of Mathematics, Deen Dayal Upadhyaya College, New Delhi 110 015,
India
E-mail: anilkarn@vsnl.net

MS received 28 September 2001

Abstract. Order unit property of a positive element in a C^* -algebra is defined. It is proved that precisely projections satisfy this order theoretic property. This way, unital hereditary C^* -subalgebras of a C^* -algebra are characterized.

Keywords. Matrix order unit property; matrix order unit ideal.

1. Introduction

Matrix ordered spaces were introduced and studied by Choi and Effros [1]. Matricially normed spaces were introduced and studied by Ruan [8]. Matrix order unit spaces, studied by Choi and Effros, are matrix ordered spaces with a matrix norm induced by an order unit. Their non-unital counterpart were studied by us (see [2–5]) and Schreiner [9]. We call such spaces L^∞ -matricially Riesz normed spaces (Schreiner call them *matrix regular operator spaces*.) We know that C^* -algebras are L^∞ -matricially Riesz normed spaces.

In this paper we consider the following problem: Which element of an L^∞ -matricially Riesz normed space looks like order unit and also determines the matrix norm as the order unit norm in its possible range? (We know that if two elements determine the (matrix) norm as order unit norms, then the two elements must be identical.) In this paper, we have been able to characterize these elements in C^* -algebras. These are precisely the projections. This is an order theoretic characterization of projections in a C^* -algebra and seems to be an important tool in a possible order theoretic characterization of non-commutative C^* -algebras. (Note that the set of all projections in $B(H)$ have a nice lattice structure.) As a consequence of the above characterization, we have been able to characterize unital hereditary C^* -subalgebras of a C^* -algebra.

Now we recollect some definitions for the sake of completeness.

DEFINITION 1.1.

An L^∞ -matricially Riesz normed space (L^∞ - mRn space) V consists of a matrix order structure $\{M_n(V)^+\}$ and a matrix norm structure $\{\|\cdot\|_n\}$ which are related in the following manner:

(I) For every $n \in \mathbb{N}$ and $v \in M(V)^+$

$$\|v\|_n = \left\{ \max\{\|u_1\|_n, \|u_2\|_n\} : \begin{pmatrix} u_1 & v \\ v^* & u_2 \end{pmatrix} \in M_{2n}(V)^+ \right\}$$

- (II) $\|v\|_n \leq \|\alpha\| \|v\|_n \|\beta\|$
 (III) $M_n(V)^+$ is $\|\cdot\|_n$ -closed or all n .

This space is denoted by $(V, \{\|\cdot\|_n\}, \{M_n(V)^+\})$.

Next, we recall the notion of an order unit. Let $(V, \{M_n(V)^+\})$ be a matrix ordered space. Then $e \in V^+$ is called an *order unit* for V , if for each $v \in V$ there is $k > 0$ such that $\begin{pmatrix} ke & v \\ v^* & ke \end{pmatrix} \in M_2(V)^+$. In this case, $e^n = e \oplus \cdots \oplus e \in M_n(V)^+$ for all n . If, in addition, V^+ is proper and if $M_n(V)^+$ is Archimedean for all n , then we can define for each n , the order unit norm $\|\cdot\|$ on $M_n(V)$ given by

$$\|v_n\| = \inf \left\{ \lambda > 0 : \begin{bmatrix} \lambda e^n & v \\ v^* & \lambda e^n \end{bmatrix} \in M_{2n}(V)^+ \right\}.$$

In this case $(V, \{\|\cdot\|_n\}, \{M_n(V)^+\})$ becomes an L^∞ - mRn space which is called a matrix order unit space and is denoted by (V, e) .

Now, let W be a self-adjoint subspace of a matrix ordered space $(V, \{M_n(V)^+\})$. For each $n \in N$, if we put

$$M_n(W)^+ = M_n(V)^+ \cap M_n(W)_{sa}$$

then $(W, \{M_n(W)^+\})$ becomes a matrix ordered space and is called a *matrix ordered subspace* of $(V, \{M_n(V)^+\})$.

A matrix ordered subspace W of V is called a *matrix order ideal*, if for any $v \in V^+$, $v \in W^+$ whenever $v \leq w$ for some $w \in W^+$. (In this case this property is carried over to $M_n(W)^+$ for all n .)

Finally, let $(V, \{\|\cdot\|_n\}, \{M_n(V)^+\})$ be an L^∞ -matricially Riesz normed space and suppose that W is a matrix order ideal in V with an order unit e . If (W, e) is a matrix order unit space such that e determines the matrix norm of V on W as the matrix order unit norm, then (W, e) is called a matrix order unit ideal of V .

2. Order unit and matrix order ideal

Let $(V, \{M_n(V)^+\})$ be a matrix ordered space and let $a \in V, a \neq 0$. Put

$$V_a = \left\{ v \in V : \begin{pmatrix} ka & v \\ v^* & ka \end{pmatrix} \in M_2(V)^+, \text{ for some } K > 0 \right\}.$$

Then V_a becomes a matrix order ideal in V and a is an order unit for V_a . Conversely, if W is a matrix order ideal of V and $a \in W^+$ is an order unit for W , then $W = V_a$. In particular, if e is an order unit for V , then V have no matrix order ideals containing e , other than itself.

Now, let $(V, \{\|\cdot\|_n\}, \{M_n(V)^+\})$ be an L^∞ - mRn space and suppose that $a \in V^+$ with $\|a\|_1 = 1$. We say that a has *order unit property* in V , if for each $v \in V_a$, we have

$$\begin{pmatrix} \|v\|_1 a & v \\ v^* & \|v\|_1 a \end{pmatrix} \in M_2(V)^+.$$

If a^n have order unit property for all n , we say that a have *matrix order unit property* in V . The following result is a simple consequence of this definition.

PROPOSITION 2.1.

Let $(V, \{\|\cdot\|_n\}, \{M_n(V)^+\})$ be an L^∞ - mRn space and suppose that $a \in V^+$ with $\|a\|_1 = 1$. Then ‘ a ’ have matrix order unit property in V if and only if (V_a, a) is a matrix order unit ideal in V . In particular, if (W, e) is a matrix order unit ideal in V , then ‘ e ’ have matrix order unit property in V and $W = V_e$.

Remark. It is possible to get an order unit $a \in V^+$ with $\|a\|_1 = 1$ which does not have order unit property in V . To see this, consider the matrix order unit space (M_2, I_2) . Then $a = \begin{pmatrix} 1 & 0 \\ 0 & 1/2 \end{pmatrix} \in M_2^+$ and $\|a\|_1 = 1$. Moreover, a determines the operator norm in the subspace defined as $A = \{\alpha a : \alpha \in \mathbf{C}\}$. Now, it is easy to see that a is an order unit for M_2 . Finally, note that the operator norm of I_2 is 1 and its order unit norm determined by a is 2. Thus a does not have order unit property in M_2 .

3. Order units in a \mathbf{C}^* -algebra

Let H be a complex Hilbert space and let $p \in B(H)$ be an (orthogonal) projection. Define

$$\begin{aligned} B_p(H) &= \{x \in B(H) : xp = px = x\} \\ &= \{pxp : x \in B(H)\} \end{aligned}$$

and

$$B^p(H) = \{x \in B(H) : x(H) \subset p(H) \text{ and } x^*(H) \subset p(H)\}.$$

Lemma 3.1. $B(H)_p = B_p(H) = B^p(H) \cong B(pH)$.

Proof. Evidently, $B_p(H) = B^p(H) \cong B(pH)$.

Let $x \in B(H)_p$. Then $\begin{pmatrix} \alpha p & x \\ x^* & \alpha p \end{pmatrix} \geq 0$ for some $\alpha > 0$. Thus, for each pair $\xi, \eta \in H$, we have

$$\left\langle \begin{pmatrix} \alpha p & x \\ x^* & \alpha p \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} \right\rangle \geq 0$$

or equivalently

$$\alpha \langle p\xi, \xi \rangle + \langle x\eta, \xi \rangle + \langle x^*\xi, \eta \rangle + \alpha \langle \eta, \eta \rangle \geq 0.$$

In particular, for $\xi \in p(H)^\perp$ and $\eta \in H$, we have

$$\langle x\eta, \xi \rangle + \langle x^*\xi, \eta \rangle + \alpha \langle \eta, \eta \rangle \geq 0.$$

Fixing η and replacing ξ by $i^k \lambda \xi$ ($k = 0, 1, 2, 3$ and $\lambda > 0$), we conclude that $\langle x\eta, \xi \rangle \geq 0$ for all $\eta \in H$ and $\xi \in p(H)^\perp$. Thus $x(H) \subset p(H)$. Next, interchanging roles of ξ and η , we conclude that $x^*(H) \subset p(H)$. Thus $B(H)_p \subset B^p(H)$.

Conversely, if $x \in B_p(H)$, then as $\begin{pmatrix} \|x\|I & x \\ x^* & \|x\|I \end{pmatrix} \geq 0$, we have

$$0 \leq \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} \begin{pmatrix} \|x\|I & x \\ x^* & \|x\|I \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}$$

$$= \begin{pmatrix} \|x\| p & p x p \\ p x^* p & \|x\| p \end{pmatrix} = \begin{pmatrix} \|x\| p & x \\ x^* & \|x\| p \end{pmatrix}$$

for $p x p = x$ and $x^* = (p x p)^* = p x^* p$. Hence follows the result. \square

COROLLARY 3.2.

Let A be a C^* -algebra and suppose that $p \in A^+$ is a projection. Then $A_p = p A p = \{p x p : x \in A\}$. In particular, every projection in A has the matrix order unit property.

Theorem 3.3. Let A be a C^* -algebra, $p \in A^+$ with $\|p\| = 1$. If p has the order unit property in A , then p is a projection.

Proof. First we show that A_p is a C^* -subalgebra of A . Let $x \in A_p$. Then $\begin{pmatrix} k p & x \\ x^* & k p \end{pmatrix} \in M_2(A)^+$, for some $k > 0$. Identifying A in a suitable $B(H)$, we conclude that $x^* x \leq k^2 p$ ([6], Exercise 3.2(ii)) so that $x^* x \in A_p$. It follows, from polarization identity, that A_p is a $*$ -algebra. To see that A_p is norm closed, let $\{x_n\}$ be any sequence in A_p converging to $x \in A$. Let $\|x_n\| \leq k$ for all n and some $k > 0$. Since p has order unit property in A , we have $\begin{pmatrix} k p & x_n \\ x_n^* & k p \end{pmatrix} \in M_2(A)^+$ for all n . It follows, from norm-closedness of $M_2(A)^+$, that $\begin{pmatrix} k p & x \\ x^* & k p \end{pmatrix} \in M_2(A)^+$. Thus A_p is a C^* -subalgebra of A .

Now identify A_p in some $B(K)$ such that A_p acts non-degenerately on K . Let $\{u_\lambda\}$ be an approximate identity of A_p . Then $u_\lambda \uparrow I$ where I is the identity operator on K (see e.g. [7], 2.2.4). Now for each λ , $\|u_\lambda\| \leq 1$ and $u_\lambda \geq 0$, so that $u_\lambda \leq p$. Thus $I \leq p$. Again, as $\|p\| = 1$ and $p \geq 0$, we have $p \leq I$. Thus $p = I$. In particular, A_p is a unital C^* -algebra. Hence p is a projection in A . \square

Now summing up Corollary 3.2 and Theorem 3.3, we obtain the following characterization.

Theorem 3.4. Let A be a C^* -algebra and suppose that $p \in A^+$ with $\|p\| = 1$. Then p has the order unit property in A if and only if p is a projection in A . In this case, $A_p = p A p$.

Remark. (1) Matrix order unit ideals in a C^* -algebra are precisely unital hereditary C^* -subalgebras. (2) Matrix order unit ideals in a von-Neumann algebra are precisely hereditary von-Neumann subalgebras. (3) Matrix order unit ideals in $B(H)$ are precisely of the form $B(K)$ where K is a closed subspace of H .

References

- [1] Choi M D and Effros E G, Injectivity and operator spaces, *J. Funct. Anal.* **24** (1977) 156–209
- [2] Karn A K and Vasudevan R, Matrix norms in matrix ordered spaces, *Glasnik Matematički* **32(52)** (1997) 87–97
- [3] Karn A K and Vasudevan R, Approximate matrix order unit spaces, *Yokohama Math. J.* **44** (1997) 73–91
- [4] Karn A K and Vasudevan R, Matrix duality for matrix ordered spaces, *Yokohama Math. J.* **45** (1998) 1–18

- [5] Karn A K and Vasudevan R, Characterizations of matricially Riesz normed spaces, *Yokohama Math. J.* **47** (2000) 143–153
- [6] Paulsen V I, Completely bounded maps and dialations, *Pitman Research Notes in Mathematics, Longman Scientific and Technical*, London, 1986
- [7] Pedersen G K, C^* -algebras and their group automorphisms (Academic Press) (1978)
- [8] Ruan Z J, Subspaces of C^* -algebras, *J. Funct. Anal.* **76** (1988) 217–230
- [9] Schreiner W J, Matrix regular operator spaces, *J. Funct. Anal.* **152** (1998) 136–175