

# Configurations in abelian categories. I.

## Basic properties

Dominic Joyce  
Lincoln College, Oxford

### Abstract

This is the first in a series of papers on *configurations* in an abelian category  $\mathcal{A}$ . Given a *finite partially ordered set (poset)*  $(I, \preceq)$ , an  $(I, \preceq)$ -*configuration*  $(\sigma, \iota, \pi)$  is a finite collection of objects  $\sigma(J)$  and morphisms  $\iota(J, K)$  or  $\pi(J, K) : \sigma(J) \rightarrow \sigma(K)$  in  $\mathcal{A}$  satisfying some axioms, where  $J, K$  are subsets of  $I$ . Configurations are a tool for describing *how an object  $X$  in  $\mathcal{A}$  decomposes into subobjects*, and are especially useful for studying *stability conditions* on  $\mathcal{A}$ .

This paper is primarily elementary algebra and category theory. We define and motivate the idea of configurations, and explain some natural operations upon them — subconfigurations, quotient configurations, refinements and improvements. The next paper studies *moduli spaces of  $(I, \preceq)$ -configurations* in  $\mathcal{A}$ , using the theory of *Artin stacks*. We prove well-behaved moduli stacks exist when  $\mathcal{A}$  is an abelian or exact category of *coherent sheaves* or *vector bundles* on a *projective  $\mathbb{K}$ -scheme  $P$* , and if  $\mathcal{A}$  is an abelian category of *representations of a quiver  $Q$* .

Subsequent papers will apply these results to define *systems of invariants* of a projective  $\mathbb{K}$ -scheme  $P$  that ‘count’ (semi)stable coherent sheaves and satisfy interesting identities, and to construct and study *infinite-dimensional algebras* associated to a quiver  $Q$ .

## 1 Introduction

This is the first of a series of papers [3, 4, 5, 6] developing the concept of *configuration* in an abelian category. Given an abelian category  $\mathcal{A}$  and a finite partially ordered set (poset)  $(I, \preceq)$ , we define an  $(I, \preceq)$ -*configuration*  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  to be a collection of objects  $\sigma(J)$  and morphisms  $\iota(J, K)$  or  $\pi(J, K) : \sigma(J) \rightarrow \sigma(K)$  in  $\mathcal{A}$  satisfying certain axioms, where  $J, K$  are subsets of  $I$ .

Configurations are a tool for describing *how an object  $X$  in  $\mathcal{A}$  decomposes into subobjects*. They are especially useful for studying *stability conditions* on  $\mathcal{A}$ . Given a *stability condition*  $(\tau, T, \leq)$  on  $\mathcal{A}$  (such as a *slope function*), objects  $X$  in  $\mathcal{A}$  are called  $\tau$ -*stable*,  $\tau$ -*semistable* or  $\tau$ -*unstable* according to whether subobjects  $S \subset X$  with  $S \neq 0, X$  have  $\tau([S]) < \tau([X])$ ,  $\tau([S]) \leq \tau([X])$ , or  $\tau([S]) > \tau([X])$ .

For some large classes of interesting abelian categories  $\mathcal{A}$ , such as the abelian category  $\text{mod-}\mathbb{K}Q$  of representations of a *quiver  $Q$*  over  $\mathbb{K}$ , or  $\text{coh}(P)$  of *coherent*

sheaves on a projective  $\mathbb{K}$ -scheme  $P$ , for  $\mathbb{K}$  an algebraically closed field, one can define moduli spaces  $\text{Obj}_{\text{ss}}^\alpha(\tau)$  of  $\tau$ -semistable objects in a fixed class  $\alpha \in K(\mathcal{A})$ , which in good cases are quasiprojective  $\mathbb{K}$ -schemes.

We shall regard  $\text{Obj}_{\text{ss}}^\alpha(\tau)$  as the subset of  $\tau$ -semistable geometric points in the algebraic stack  $\mathfrak{Obj}_{\mathcal{A}}^\alpha$  of all objects in class  $\alpha$ . One goal of [3, 4, 5] is to understand the relationship between  $\text{Obj}_{\text{ss}}^\alpha(\tau)$  and  $\text{Obj}_{\text{ss}}^\alpha(\tilde{\tau})$  for two different stability conditions  $(\tau, T, \leq)$ ,  $(\tilde{\tau}, \tilde{T}, \leq)$ . Our key idea is that this is best understood by introducing moduli stacks  $\mathfrak{M}(I, \preceq, \kappa)_{\mathcal{A}}$  of  $(I, \preceq)$ -configurations in  $\mathcal{A}$ .

Write  $\mathcal{M}_{\text{ss}}(I, \preceq, \kappa, \tau)_{\mathcal{A}}$  for the subset of points  $[(\sigma, \iota, \pi)]$  in  $\mathfrak{M}(I, \preceq, \kappa)_{\mathcal{A}}$  with  $\sigma(\{i\})$   $\tau$ -semistable for all  $i \in I$ . When  $I = \{i\}$  is one point  $\mathcal{M}_{\text{ss}}(I, \preceq, \kappa, \tau)_{\mathcal{A}}$  coincides with  $\text{Obj}_{\text{ss}}^{\kappa(i)}(\tau)$ . We shall express  $\mathcal{M}_{\text{ss}}(I, \preceq, \kappa, \tilde{\tau})_{\mathcal{A}}$  in terms of projections of  $\mathcal{M}_{\text{ss}}(K, \trianglelefteq, \mu, \tau)_{\mathcal{A}}$  for finite posets  $(K, \trianglelefteq)$  more complicated than  $(I, \preceq)$ .

The next paper [3] deals with the technicalities of constructing Artin stacks of configurations  $\mathfrak{M}(I, \preceq, \kappa)_{\mathcal{A}}$ . Under some assumptions on  $\mathcal{A}$ ,  $(\tau, T, \leq)$ , in [4, 5] we define systems of invariants of  $\mathcal{A}$ ,  $(\tau, T, \leq)$  by taking weighted Euler characteristics of  $\mathcal{M}_{\text{ss}}(I, \preceq, \kappa, \tau)_{\mathcal{A}}$ , and determine identities the invariants satisfy, and their transformation laws as  $(\tau, T, \leq)$  changes.

The fifth paper [6] will give applications of a different kind. Ringel–Hall algebras are associative algebras of constructible functions on moduli spaces of quiver representations. The multiplication is defined by a kind of convolution product. One can also use perverse sheaves, or homology, in place of constructible functions. Authors such as Ringel, Green, Lusztig and Nakajima have used Ringel–Hall algebras to give geometric interpretations of interesting infinite-dimensional Lie algebras, such as subalgebras of universal enveloping algebras  $U(\mathfrak{g})$  of Kac–Moody Lie algebras  $\mathfrak{g}$ .

We shall reinterpret the Ringel–Hall algebra construction in terms of natural operations on constructible functions on the moduli stacks of configurations  $\mathfrak{M}(I, \preceq, \kappa)$ , using the theory of constructible functions on stacks developed by the author in [7]. The power of the configurations framework (once we have done a lot of work to set it up) then suggests many generalizations, which may lead to progress in understanding infinite-dimensional algebras.

Also, we will be able to import ideas from Ringel–Hall algebras to configurations of vector bundles or coherent sheaves on a projective  $\mathbb{K}$ -scheme  $P$ . We can thus associate infinite-dimensional algebras to  $P$ , which may have a String-Theoretic interpretation as algebras of BPS states.

This is the start of a broader programme. In later papers [8] the author intends to extend the notion of configurations, and the corresponding moduli stacks and invariants, to triangulated categories. Then by applying them to derived categories of coherent sheaves on Calabi–Yau manifolds, we shall formulate some results and conjectures on Homological Mirror Symmetry, branes in String Theory, and  $\Pi$ -stability.

This paper introduces configurations in an abelian category  $\mathcal{A}$ , and studies their basic properties. We begin in §2 with an introduction to abelian categories, focussing on the notion of subobject  $S \subset X$  of an object  $X \in \mathcal{A}$ , and the Jordan–Hölder Theorem for  $\mathcal{A}$  of finite length, which splits each  $X \in \mathcal{A}$  into simple

factors  $S_1, \dots, S_n$ .

Section 3 refines the Jordan–Hölder Theorem in the case when the simple factors  $S_1, \dots, S_n$  of  $X \in \mathcal{A}$  are nonisomorphic. We find that the set of all *subobjects* of  $X$  may be classified using a *partial order*  $\preceq$  on  $I = \{1, \dots, n\}$ , the indexing set for the simple factors of  $X$ . We also classify *quotient objects*, *factors* and *composition series* for  $X$  using  $(I, \preceq)$ .

Motivated by this, §4 defines the notion of  $(I, \preceq)$ -*configuration*  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ , and proves that it captures the properties of the set of all subobjects of  $X \in \mathcal{A}$  when  $X$  has nonisomorphic simple factors  $\{S^i : i \in I\}$ .

Section 5 defines some elementary operations on configurations. Given an  $(I, \preceq)$ -configuration we can make *sub-* and *quotient*  $(K, \trianglelefteq)$ -configurations, where  $(K, \trianglelefteq)$  comes from  $(I, \preceq)$  with  $K \subseteq I$  or using a surjective  $\phi : I \rightarrow K$ . We also construct new configurations by *substituting* one configuration into another.

Let  $\trianglelefteq, \preceq$  be partial orders on a finite  $I$ , with  $\trianglelefteq$  stronger than  $\preceq$ . Then each  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  has a quotient  $(I, \trianglelefteq)$ -configuration  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . We call  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -*improvement* of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . We call  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  *best* if it has no strict improvements. Section 6 shows that improvements can be divided into a sequence of *steps*, classifies *one step improvements*, and gives a criterion for *best configurations* in terms of whether short exact sequences split.

Finally, §7 discusses five short topics: the classes of the objects  $\sigma(J)$  in the *Grothendieck group*  $K_0(\mathcal{A})$ ; configurations in *exact categories*; interpretations of  $(I, \preceq)$ -configurations in terms of *flasque sheaves* on  $I$ , and as *functors*  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  for a category  $\mathcal{C}_{(I, \preceq)}$  constructed from  $(I, \preceq)$ ; and we show that  $(I, \preceq)$ -configurations in  $\mathcal{A}$  form an *exact category*.

*Acknowledgements.* I would like to thank Tom Bridgeland for some inspiring conversations, and helpful comments on early versions of this paper. I would also like to thank Richard Thomas, Paul Seidel, Frances Kirwan and Alastair King for useful conversations. I was supported by an EPSRC Advanced Research Fellowship whilst writing this paper.

## 2 Introduction to abelian categories

We now review material on abelian categories we shall need later. Some useful references are Popescu [9] and Gelfand and Manin [2, §II.5–§II.6].

### 2.1 Abelian categories

Here is the definition of abelian category, taken from [2, §II.5].

**Definition 2.1.** Let  $\mathcal{A}$  be a category. As a shorthand, write  $X \in \mathcal{A}$  or  $X \in \text{Obj}(\mathcal{A})$  when  $X$  is an object of  $\mathcal{A}$ , and  $f \in \text{Mor}(\mathcal{A})$  when  $f$  is a morphism of  $\mathcal{A}$ . When  $X, Y \in \mathcal{A}$  write  $\text{Hom}(X, Y)$  for the set of morphisms  $f : X \rightarrow Y$  in  $\mathcal{A}$ . Write  $\text{id}_X \in \text{Hom}(X, X)$  for the identity map  $\text{id}_X : X \rightarrow X$ .

We call  $\mathcal{A}$  an *additive category* if it has the properties:

- (i)  $\text{Hom}(X, Y)$  is an *abelian group* for all  $X, Y \in \mathcal{A}$ , and composition of morphisms is *biadditive*.
- (ii) There exists a *zero object*  $0 \in \mathcal{A}$  such that  $\text{Hom}(0, 0) = 0$ .
- (iii) For any  $X, Y \in \mathcal{A}$  there exists  $Z \in \mathcal{A}$  and morphisms  $\iota_X : X \rightarrow Z$ ,  $\iota_Y : Y \rightarrow Z$ ,  $\pi_X : Z \rightarrow X$ ,  $\pi_Y : Z \rightarrow Y$  with  $\pi_X \circ \iota_X = \text{id}_X$ ,  $\pi_Y \circ \iota_Y = \text{id}_Y$ ,  $\iota_X \circ \pi_X + \iota_Y \circ \pi_Y = \text{id}_Z$  and  $\pi_X \circ \iota_Y = \pi_Y \circ \iota_X = 0$ . We write  $Z = X \oplus Y$ , the *direct sum* of  $X$  and  $Y$ . Any two such direct sums are canonically isomorphic.

Let  $\mathcal{A}$  be an additive category, and  $f : X \rightarrow Y$  a morphism in  $\mathcal{A}$ . We call  $k : K \rightarrow X$  a *kernel* of  $f$  if  $f \circ k = 0$  and for any  $k' : K' \rightarrow X$  with  $f \circ k' = 0$  there exists a unique  $h : K' \rightarrow K$  with  $k' = k \circ h$ . Similarly we call  $c : Y \rightarrow C$  a *cokernel* of  $f$  if  $c \circ f = 0$  and for any  $c' : Y \rightarrow C'$  with  $c' \circ f = 0$  there exists a unique  $h : C \rightarrow C'$  with  $c' = h \circ c$ .

If a kernel or cokernel exists it is unique up to canonical isomorphism. Often we refer to  $K, C$  as the kernel or cokernel of  $f$ , taking  $k, c$  to be implicitly given. Define a morphism  $f : X \rightarrow Y$  to be *injective* if it has kernel  $0$ , and *surjective* if it has cokernel  $0$ .

We call  $\mathcal{A}$  an *abelian category* if it satisfies (i)–(iii) above and:

- (iv) For any morphism  $f : X \rightarrow Y$  there is a sequence  $K \xrightarrow{k} X \xrightarrow{i} I \xrightarrow{j} Y \xrightarrow{c} C$  in  $\mathcal{A}$  such that  $j \circ i = f$ , and  $K$  is the kernel of  $f$ , and  $C$  the cokernel of  $f$ , and  $I$  is both the cokernel of  $k$  and the kernel of  $c$ .

An abelian category  $\mathcal{A}$  is called  $\mathbb{K}$ -*linear* over a field  $\mathbb{K}$  if  $\text{Hom}(X, Y)$  is a *vector space* over  $\mathbb{K}$  for all  $X, Y \in \mathcal{A}$ , and composition maps are *bilinear*.

We will often use the following properties of abelian categories:

- If  $i \circ f = i \circ g$  and  $i$  is *injective*, then  $f = g$  ( $i$  is *left cancellable*).
- If  $f \circ \pi = g \circ \pi$  and  $\pi$  is *surjective*, then  $f = g$  ( $\pi$  is *right cancellable*).
- If  $f : X \rightarrow Y$  is *injective* and *surjective*, then it is an *isomorphism*.

In an abelian category we can define *exact sequences* [2, §II.6].

**Definition 2.2.** Let  $\mathcal{A}$  be an abelian category, and  $X \xrightarrow{f} Y \xrightarrow{g} Z$  a sequence in  $\mathcal{A}$  with  $g \circ f = 0$ . Let  $k : K \rightarrow Y$  be the kernel of  $g$  and  $c : Y \rightarrow C$  the cokernel of  $f$ . Then there exist unique morphisms  $a : X \rightarrow K$  and  $b : C \rightarrow Z$  with  $f = k \circ a$  and  $g = b \circ c$ . We say that  $X \xrightarrow{f} Y \xrightarrow{g} Z$  is *exact at  $Y$*  if  $a$  is surjective, or equivalently if  $b$  is injective.

A complex in  $\mathcal{A}$  is called *exact* if it is exact at every term. A short exact sequence  $0 \rightarrow X \xrightarrow{f} Y \xrightarrow{g} Z \rightarrow 0$  in  $\mathcal{A}$  is called *split* if there exists an isomorphism  $h : X \oplus Z \rightarrow Y$  such that the following diagram commutes:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & X & \xrightarrow{\iota_X} & X \oplus Z & \xrightarrow{\pi_Z} & Z & \longrightarrow & 0 \\
 & & \text{id}_X \downarrow & & h \downarrow & & \text{id}_Z \downarrow & & \\
 0 & \longrightarrow & X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \longrightarrow & 0.
 \end{array} \tag{1}$$

The *Grothendieck group*  $K_0(\mathcal{A})$  is the abelian group generated by  $\text{Obj}(\mathcal{A})$ , with a relation  $[Y] = [X] + [Z]$  for each short exact sequence  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ .

## 2.2 Subobjects, quotient objects and factors

*Subobjects* of objects in  $\mathcal{A}$  are analogous to subgroups of an abelian group.

**Definition 2.3.** Let  $\mathcal{A}$  be an abelian category. Two injective morphisms  $i : S \rightarrow X$ ,  $i' : S' \rightarrow X$  in  $\mathcal{A}$  are *equivalent* if there exists an isomorphism  $h : S \rightarrow S'$  with  $i = i' \circ h$ . Then  $h$  is unique. A *subobject* of  $X \in \mathcal{A}$  is an equivalence class of injective morphisms  $i : S \rightarrow X$ . Usually we refer to  $S$  as the subobject, taking both  $i$  and the equivalence class to be implicitly given, and write  $S \subset X$  to mean  $S$  is a subobject of  $X$ . We write  $0, X$  for the subobjects of  $X$  which are equivalence classes of  $0 \rightarrow X$  and  $\text{id}_X : X \rightarrow X$ .

Similarly, surjective morphisms  $\pi : X \rightarrow Q$ ,  $\pi' : X \rightarrow Q'$  in  $\mathcal{A}$  are *equivalent* if there is an isomorphism  $h : Q \rightarrow Q'$  with  $\pi' = h \circ \pi$ . A *quotient object* of  $X \in \mathcal{A}$  is an equivalence class of surjective  $\pi : X \rightarrow Q$ . Usually we just refer to  $Q$  as the quotient object.

If  $S, T \subset X$  are represented by  $i : S \rightarrow X$  and  $j : T \rightarrow X$ , we write  $S \subset T \subset X$  if there exists  $a : S \rightarrow T$  with  $i = j \circ a$ . Then  $a$  is injective, and so fits into an exact sequence  $0 \rightarrow S \xrightarrow{a} T \xrightarrow{b} F \rightarrow 0$  for  $b, F$  determined up to canonical isomorphism. We write  $F = T/S$ , and call  $F$  a *factor* of  $X \in \mathcal{A}$ . When  $T = X$  and  $j = \text{id}_X$ ,  $X/S$  is a *quotient object*.

We define operations  $\cap, +$  on subobjects, following Popescu [9, §2.6]. The notation comes from the intersection and sum of *subgroups of abelian groups*.

**Definition 2.4.** Let  $\mathcal{A}$  be an abelian category, let  $X \in \mathcal{A}$ , and suppose injective maps  $i : S \rightarrow X$ ,  $j : T \rightarrow X$  define subobjects  $S, T$  of  $X$ . Apply part (iv) of Definition 2.1 to  $f = i \circ \pi_S + j \circ \pi_T : S \oplus T \rightarrow X$ . This yields  $U, V \in \mathcal{A}$  and morphisms  $k : U \rightarrow S \oplus T$ ,  $l : S \oplus T \rightarrow V$  and  $e : V \rightarrow X$  such that  $i \circ \pi_S + j \circ \pi_T = e \circ l$ , and  $k$  is the kernel of  $i \circ \pi_S + j \circ \pi_T$ , and  $l$  is the cokernel of  $k$ , and  $e$  is the *image* (the kernel of the cokernel) of  $i \circ \pi_S + j \circ \pi_T$ .

Define  $a : U \rightarrow S$  by  $a = k \circ \pi_S$ , and  $b : U \rightarrow T$  by  $b = -k \circ \pi_T$  and  $c : S \rightarrow V$  by  $c = f \circ \iota_S$ , and  $d : T \rightarrow V$  by  $d = f \circ \iota_T$ . Then  $k = \iota_S \circ a - \iota_T \circ b$ ,  $l = c \circ \pi_S + d \circ \pi_T$ ,  $i = e \circ c$  and  $j = e \circ d$ . Now  $0 \rightarrow U \xrightarrow{k} S \oplus T \xrightarrow{l} V \rightarrow 0$  is exact. So  $i \circ a = j \circ b$ , and

$$0 \longrightarrow U \xrightarrow{\iota_S \circ a - \iota_T \circ b} S \oplus T \xrightarrow{c \circ \pi_S + d \circ \pi_T} V \longrightarrow 0 \quad \text{is exact.} \quad (2)$$

As  $i, a$  are injective  $i \circ a = j \circ b : U \rightarrow X$  is injective, and defines a *subobject* of  $X$ . Define  $S \cap T$  to be this subobject. Similarly,  $e : V \rightarrow X$  is injective, and so defines a subobject of  $X$ . Define  $S + T$  to be this subobject. As  $U, V, a, \dots, e$  are unique up to canonical isomorphism,  $S \cap T$  and  $S + T$  depend only on the subobjects  $S, T$  of  $X$ . The morphisms  $a, b, c, d$  give inclusions of subobjects

$$S \cap T \subset S \subset S + T \subset X \quad \text{and} \quad S \cap T \subset T \subset S + T \subset X. \quad (3)$$

By Popescu [9, Prop. 2.6.4, p. 39] there are canonical isomorphisms

$$S/(S \cap T) \cong (S + T)/T \quad \text{and} \quad T/(S \cap T) \cong (S + T)/S. \quad (4)$$

These operations  $\cap, +$  are *commutative* and *associative*. We can therefore form multiple sums and intersections. We shall write  $\sum_{j \in J} T_j$  for the multiple sum  $+$  of a finite set of subobjects  $T_j \subset X$ , in the obvious way.

### 2.3 The Jordan–Hölder Theorem

The *Jordan–Hölder Theorem* in group theory decomposes a (finite) group into simple factors, using chains of normal subgroups. We shall explain the analogue of this in an abelian category.

**Definition 2.5.** Let  $\mathcal{A}$  be an abelian category. We call  $\mathcal{A}$  *artinian* if for all  $X \in \mathcal{A}$ , all descending chains of subobjects  $\cdots \subset A_2 \subset A_1 \subset X$  stabilize, that is,  $A_{n+1} = A_n$  for all  $n \gg 0$ . We call  $\mathcal{A}$  *noetherian* if for all  $X \in \mathcal{A}$ , all ascending chains of subobjects  $A_1 \subset A_2 \subset \cdots \subset X$  stabilize, that is,  $A_n = A_{n+1}$  for all  $n \gg 0$ . We call  $\mathcal{A}$  of *finite length* if it is both artinian and noetherian.

A nonzero object  $X$  in an abelian category  $\mathcal{A}$  is called *simple* if it has no nontrivial proper subobjects. That is,  $X \not\cong 0$ , and if  $i : S \rightarrow X$  is injective then either  $S \cong 0$  or  $i$  is an isomorphism.

Let  $X \in \mathcal{A}$  and consider *filtrations* of subobjects

$$0 = A_0 \subset A_1 \subset \cdots \subset A_n = X \quad (5)$$

in  $\mathcal{A}$ . Such a filtration is called *without repetitions* if none of the inclusions  $i_k : A_k \rightarrow A_{k+1}$  is an isomorphism. A *refinement* of (5) is any filtration obtained by inserting further terms. We allow (5) as a refinement of itself, i.e. by inserting no further terms. We call (5) a *composition series* for  $X$  if the factors  $S_k = A_k/A_{k-1}$  are simple objects in  $\mathcal{A}$ .

Here is the Jordan–Hölder Theorem in an abelian category, [11, Th. 2.1].

**Theorem 2.6.** *Let  $\mathcal{A}$  be an abelian category of finite length. Then every filtration  $0 = A_0 \subset A_1 \subset \cdots \subset A_n = X$  without repetitions can be refined to a composition series for  $X$ . Suppose  $0 = A_0 \subset A_1 \subset \cdots \subset A_m = X$  and  $0 = B_0 \subset B_1 \subset \cdots \subset B_n = X$  are two composition series for  $X \in \mathcal{A}$ , with simple factors  $S_k = A_k/A_{k-1}$  and  $T_k = B_k/B_{k-1}$ . Then  $m = n$ , and for some permutation  $\sigma$  of  $1, \dots, n$  we have  $S_k \cong T_{\sigma(k)}$  for  $k = 1, \dots, n$ .*

### 2.4 Exact categories

Finally we define a class of subcategories of abelian categories called *exact categories*, that we will need in §7.2. They were introduced by Quillen [10, §2], and are discussed by Gelfand and Manin [2, Ex. IV.3.3, p. 275].

**Definition 2.7.** Let  $\mathcal{A}$  be an abelian category, and  $\mathcal{B}$  be a *full additive subcategory* of  $\mathcal{A}$  in the sense of Definition 2.1, which is *closed under extensions*. That is, if  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$  is a short exact sequence in  $\mathcal{A}$  with  $X, Z \in \mathcal{B}$  then  $Y \in \mathcal{B}$ . Let  $\mathcal{E}$  be the class of exact sequences  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$  in  $\mathcal{A}$  with  $X, Y, Z \in \mathcal{B}$ . Then the pair  $(\mathcal{B}, \mathcal{E})$  is called an *exact category*.

Quillen [10, §2] gives necessary and sufficient conditions on  $\mathcal{B}, \mathcal{E}$  for  $\mathcal{B}$  to be embedded in an abelian category  $\mathcal{A}$  in this way, and we take this to be the *definition* of an exact category. An *exact functor*  $F : (\mathcal{B}, \mathcal{E}) \rightarrow (\mathcal{B}', \mathcal{E}')$  of abelian categories is a functor  $F : \mathcal{B} \rightarrow \mathcal{B}'$  taking exact sequences  $\mathcal{E}$  in  $\mathcal{B}$  to exact sequences  $\mathcal{E}'$  in  $\mathcal{B}'$ . Note that  $F$  need *not* come from an exact functor  $\mathcal{A} \rightarrow \mathcal{A}'$  of the enveloping abelian categories.

### 3 Refining the Jordan–Hölder Theorem

We shall study the following situation.

**Definition 3.1.** Let  $\mathcal{A}$  be an abelian category of finite length, and  $X \in \mathcal{A}$ . Then  $X$  admits a *composition series*  $0 = A_0 \subset A_1 \subset \cdots \subset A_n = X$  by Theorem 2.6, and the *simple factors*  $S_k = A_k/A_{k-1}$  for  $k = 1, \dots, n$  of  $X$  are independent of choices, up to isomorphism and permutation of  $1, \dots, n$ . Suppose  $S_k \not\cong S_l$  for  $1 \leq k < l \leq n$ . Then we say that  $X$  has *nonisomorphic simple factors*.

Let  $X$  have nonisomorphic simple factors, and let  $I$  be an *indexing set* for  $\{S_1, \dots, S_n\}$ , so that  $|I| = n$ , and write  $\{S_1, \dots, S_n\} = \{S^i : i \in I\}$ . Then Theorem 2.6 implies that for every composition series  $0 = B_0 \subset B_1 \subset \cdots \subset B_n = X$  for  $X$  with simple factors  $T_k = B_k/B_{k-1}$ , there exists a unique bijection  $\phi : I \rightarrow \{1, \dots, n\}$  such that  $S^i \cong T_{\phi(i)}$  for all  $i \in I$ .

Define a *partial order*  $\preceq$  on  $I$  by  $i \preceq j$  for  $i, j \in I$  if and only if  $\phi(i) \leq \phi(j)$  for all bijections  $\phi : I \rightarrow \{1, \dots, n\}$  constructed from a composition series  $0 = B_0 \subset B_1 \subset \cdots \subset B_n = X$  for  $X$  as above. Then  $(I, \preceq)$  is a *partially ordered set*, or *poset* for short.

The point of this definition is to treat all the Jordan–Hölder composition series  $0 = B_0 \subset B_1 \subset \cdots \subset B_n = X$  for  $X$  on an *equal footing*. Now writing the simple factors of  $X$  as  $S_k$  for  $k = 1, \dots, n$  gives them a preferred order  $S_1, S_2, \dots, S_n$ , and so favours one composition series over the rest. So instead we write the simple factors as  $S^i$  for  $i \in I$ , some arbitrary *indexing set*.

To make our notation easier to follow we shall generally use superscripts  $S^i$  when  $i \in I$  and subscripts  $T_k$  when  $k = 1, \dots, n$ , and write elements of  $I$  as  $i, j$ , and elements of  $\{1, \dots, n\}$  as  $k, l$ . Here is some more notation.

**Definition 3.2.** Let  $(I, \preceq)$  be a finite poset. Define  $J \subseteq I$  to be

- (i) an *s-set* if  $i \in I, j \in J$  and  $i \preceq j$  implies  $i \in J$ ,
- (ii) a *q-set* if  $i \in I, j \in J$  and  $j \preceq i$  implies  $i \in J$ , and
- (iii) an *f-set* if  $i \in I$  and  $h, j \in J$  and  $h \preceq i \preceq j$  implies  $i \in J$ .

The motivation for this is that below s-sets will correspond to *subobjects*  $S \subset X$ , q-sets to *quotient objects*  $X/S$ , and f-sets to *factors*  $T/S$  for  $S \subset T \subset X$ .

Here are some properties of s-sets, q-sets and f-sets.

**Proposition 3.3.** *Let  $(I, \preceq)$  be a finite poset. Then*

- (a)  $I$  and  $\emptyset$  are s-sets. If  $J, K$  are s-sets then  $J \cap K$  and  $J \cup K$  are s-sets.
- (b)  $J$  is an s-set if and only if  $I \setminus J$  is a q-set.
- (c) If  $J$  is an s-set and  $K$  a q-set, then  $J \cap K$  is an f-set. Every f-set is of this form.
- (d) If  $J \subset K$  are s-sets then  $K \setminus J$  is an f-set. Every f-set is of this form.
- (e) If  $J, K$  are f-sets then  $J \cap K$  is an f-set, but  $J \cup K$  may not be an f-set.

The proof is elementary, and left as an exercise. For the last part of (c), if  $F \subseteq I$  is an f-set, define  $J = \{i \in I : i \preceq j \text{ for some } j \in F\}$  and  $K = \{i \in I : j \preceq i \text{ for some } j \in F\}$ . It easily follows that  $J$  is an s-set,  $K$  a q-set, and  $F = J \cap K$ . Similarly, for the last part of (d), if  $F \subseteq I$  is an f-set, define  $K = \{i \in I : i \preceq j \text{ for some } j \in F\}$  and  $J = K \setminus F$ . It easily follows that  $J \subset K$  are s-sets with  $F = K \setminus J$ .

Note that (a) and (b) imply that the collections of s-sets and q-sets are both *topologies* on  $I$ , but (e) shows that the f-sets may not be a topology on  $I$ . Also  $\preceq$  can be reconstructed from the set of s-sets on  $I$ , as  $i \preceq j$  if and only if  $i \in J$  for every s-set  $J \subset I$  with  $j \in J$ . In the following series of results we establish 1-1 correspondences between *subobjects*, *quotient objects* and *factors* of  $X$ , and *s-sets*, *q-sets* and *f-sets* of  $I$ .

**Lemma 3.4.** *In the situation of Definition 3.1, suppose  $S \subset X$  is a subobject. Then there exists a unique s-set  $J \subseteq I$  such that the simple factors in any composition series for  $S$  are isomorphic to  $S^i$  for  $i \in J$ .*

*Proof.* Let  $0 = B_0 \subset \cdots \subset B_l = S$  be a composition series for  $S$ , with simple factors  $T_k = B_k/B_{k-1}$  for  $k = 1, \dots, l$ . Then  $0 = B_0 \subset \cdots \subset B_l \subset X$  is a filtration of  $X$  without repetitions, and can be refined to a composition series by Theorem 2.6. As  $T_k$  is simple, no extra terms are inserted between  $B_{k-1}$  and  $B_k$ . Thus  $X$  has a composition series  $0 = B_0 \subset \cdots \subset B_l \subset \cdots \subset B_n = X$ , with simple factors  $T_k = B_k/B_{k-1}$  for  $k = 1, \dots, n$ .

By Definition 3.1 there is a unique bijection  $\phi : I \rightarrow \{1, \dots, n\}$  such that  $S^i \cong T_{\phi(i)}$  for all  $i \in I$ . Define  $J = \phi^{-1}(\{1, \dots, l\})$ . Then  $J \subseteq I$ , and the simple factors  $T_k$  of the composition series  $0 = B_0 \subset \cdots \subset B_l = S$  are isomorphic to  $S^i$  for  $i \in J$ . Theorem 2.6 then implies that the simple factors in *any* composition series for  $S$  are isomorphic to  $S^i$  for  $i \in J$ .

Uniqueness of  $J$  is now clear, as a different  $J$  would give different simple factors for  $S$ . Suppose  $j \in J$  and  $i \in I \setminus J$ . Then  $1 \leq \phi(j) \leq l$  and  $l+1 \leq \phi(i) \leq n$ , so  $\phi(j) < \phi(i)$ , which implies that  $i \not\preceq j$  by Definition 3.1. Hence if  $j \in J$  and  $i \in I$  with  $i \preceq j$  then  $i \in J$ , and  $J$  is an s-set.  $\square$

**Lemma 3.5.** *Suppose  $S, T \subset X$  correspond to s-sets  $J, K \subseteq I$ , as in Lemma 3.4. Then  $S \cap T$  corresponds to  $J \cap K$ , and  $S + T$  corresponds to  $J \cup K$ .*

*Proof.* Let  $S \cap T$  correspond to the s-set  $L \subseteq I$ , and  $S + T$  to the s-set  $M$ . We must show  $L = J \cap K$  and  $M = J \cup K$ . By Theorem 2.6 we may refine the filtration  $0 \subset S \cap T \subset S$  to a composition series for  $S$  containing one for  $S \cap T$ . Thus the simple factors of  $S$  contain those of  $S \cap T$ , and  $L \subseteq J$ . Similarly  $L \subseteq K$ , so  $L \subseteq J \cap K$ , and  $J \cup K \subseteq M$  as  $S, T \subseteq S \cap T$ .

Now the simple factors of  $S/(S \cap T)$  are  $S^i$  for  $i \in J \setminus L$ , and the simple factors of  $(S + T)/T$  are  $S^i$  for  $i \in M \setminus K$ . As  $S/(S \cap T) \cong (S + T)/T$  by (4) we see that  $J \setminus L = M \setminus K$ . Together with  $L \subseteq J \cap K$  and  $J \cup K \subseteq M$  this implies that  $L = J \cap K$  and  $M = J \cup K$ .  $\square$

**Lemma 3.6.** *Suppose  $S, T \subset X$  correspond to s-sets  $J, K \subseteq I$ . Then  $J \subseteq K$  if and only if  $S \subset T \subset X$ , and  $J = K$  if and only if  $S = T$ .*

*Proof.* If  $S \subset T \subset X$  we can refine  $0 \subset S \subset T \subset X$  to a composition series  $0 = B_0 \subset B_1 \subset \dots \subset B_n = X$  with  $S = B_k$  and  $T = B_l$  for  $0 \leq k \leq l \leq n$ . Let  $T_m = B_m/B_{m-1}$ . Then the simple factors of  $S$  are  $T_1, \dots, T_k$  and of  $T$  are  $T_1, \dots, T_l$ . Hence  $J \subseteq K$ , as  $k \leq l$ . This proves the first ‘if’.

Now suppose  $S, T \subset X$  and  $J \subseteq K$ . Then  $J \cap K = J$ , so  $S \cap T$  corresponds to the s-set  $J$ . But  $S \cap T \subset S$ , so  $S/(S \cap T)$  has no simple factors, and  $S = S \cap T$ . Thus  $S \subset T$ , proving the first ‘only if’. The second part is immediate.  $\square$

**Lemma 3.7.** *Let  $j \in I$  and define  $J^j = \{i \in I : i \preceq j\}$ . Then  $J^j$  is an s-set, and there exists a subobject  $D^j \subset X$  corresponding to  $J^j$ .*

*Proof.* Clearly  $J^j$  is an s-set. By Definition 3.1 each composition series  $0 = B_0 \subset B_1 \subset \dots \subset B_n = X$  for  $X$  gives a bijection  $\phi : I \rightarrow \{1, \dots, n\}$ . Let  $\phi_1, \dots, \phi_r$  be the distinct bijections  $\phi : I \rightarrow \{1, \dots, n\}$  realized by composition series for  $X$ . For each  $k = 1, \dots, r$  choose a composition series  $0 = B_0 \subset B_1 \subset \dots \subset B_n = X$  with bijection  $\phi_k$ , and define  $C_k$  to be the subobject  $B_{\phi_k(j)} \subset X$ .

This defines subobjects  $C_1, \dots, C_r \subset X$ , where  $C_k$  corresponds to the s-set  $\phi_k^{-1}(\{1, \dots, \phi_k(j)\}) \subseteq I$ . Define  $D^j = C_1 \cap C_2 \cap \dots \cap C_r$ . Then  $S \subset X$ , and Lemma 3.5 shows that  $D^j$  corresponds to the s-set

$$\begin{aligned} \bigcap_{k=1}^r \phi_k^{-1}(\{1, \dots, \phi_k(j)\}) &= \bigcap_{k=1}^r \{i \in I : \phi_k(i) \leq \phi_k(j)\} = \\ \{i \in I : \phi_k(i) \leq \phi_k(j) \text{ for all } k = 1, \dots, r\} &= \{i \in I : i \preceq j\} = J^j, \end{aligned}$$

by definition of  $\preceq$ .  $\square$

We can now *classify subobjects* of  $X$  in terms of s-sets.

**Proposition 3.8.** *In the situation of Definitions 3.1 and 3.2, for each s-set  $J \subseteq I$  there exists a unique subobject  $S \subset X$  such that the simple factors in any composition series for  $S$  are isomorphic to  $S^i$  for  $i \in J$ . This defines a 1-1 correspondence between subobjects  $S \subset X$  and s-sets  $J \subseteq I$ .*

*Proof.* For each  $j \in J$  define  $J^j$  and  $D^j$  as in Lemma 3.7. Then  $j \in J^j \subseteq J$ , so  $J = \bigcup_{j \in J} J^j$ . Set  $S = \sum_{j \in J} D^j$ . Then  $S \subset X$  corresponds to the s-set  $\bigcup_{j \in J} J^j = J$  by Lemma 3.5. Uniqueness follows from Lemma 3.6.  $\square$

In the same way we classify *quotient objects* of  $X$  in terms of q-sets.

**Proposition 3.9.** *In the situation of Definitions 3.1 and 3.2, for each q-set  $K \subseteq I$  there exists a unique quotient object  $Q = X/S$  of  $X$  such that the simple factors in any composition series for  $Q$  are isomorphic to  $S^i$  for  $i \in K$ . This defines a 1-1 correspondence between quotient objects and q-sets.*

*Proof.* Let  $S \subset X$  correspond to an s-set  $J \subseteq I$ , and let  $Q = X/S$  be the quotient object. If the simple factors of  $S$  are  $S^i$  for  $i \in J$ , then the simple factors of  $Q$  are  $S^i$  for  $i \in K = I \setminus J$ . But  $K = I \setminus J$  is a q-set if and only if  $J \subseteq I$  is an s-set. The result then follows from Proposition 3.8.  $\square$

We can also classify *composition series* for  $X$ .

**Proposition 3.10.** *In the situation of Definition 3.1, for each bijection  $\phi : I \rightarrow \{1, \dots, n\}$  there exists a unique composition series  $0 = B_0 \subset \dots \subset B_n = X$  with  $S^i \cong B_{\phi(i)}/B_{\phi(i)-1}$  for all  $i \in I$  if and only if  $i \preceq j$  implies  $\phi(i) \leq \phi(j)$ .*

*Proof.* The ‘only if’ part follows from Definition 3.1. For the ‘if’ part, let  $\phi : I \rightarrow \{1, \dots, n\}$  be a bijection for which  $i \preceq j$  implies that  $\phi(i) \leq \phi(j)$ . Then  $\phi^{-1}(\{1, \dots, k\})$  is an s-set for each  $k = 0, 1, \dots, n$ . Let  $B_k \subset X$  be the unique subobject corresponding to  $\phi^{-1}(\{1, \dots, k\})$ , which exists by Proposition 3.8. It easily follows that  $0 = B_0 \subset B_1 \subset \dots \subset B_n = X$  is the unique composition series with  $B_k/B_{k-1} \cong S^{\phi^{-1}(k)}$  for  $k = 1, \dots, n$ , and the result follows.  $\square$

This implies that composition series for  $X$  up to isomorphism are in 1-1 correspondence with *total orders* on  $I$  compatible with the partial order  $\preceq$ . In Definition 3.1 we defined the partial order  $\preceq$  on  $I$  to be the intersection of all the total orders on  $I$  coming from composition series for  $X$ . We now see that *every* total order on  $I$  compatible with  $\preceq$  comes from a composition series.

Recall that a *factor* of  $X$  is a quotient  $T/S$  for  $S \subset T \subset X$ . We classify the factors of  $X$  in terms of f-sets.

**Proposition 3.11.** *In the situation of Definitions 3.1 and 3.2, for each f-set  $L \subseteq I$  there exists  $U \in \mathcal{A}$ , unique up to canonical isomorphism, such that*

- (a) *the simple factors in any composition series for  $U$  are isomorphic to  $S^i$  for  $i \in L$ , and*
- (b) *if  $J \subset K$  are s-sets in  $I$  with  $K \setminus J = L$ , and  $S \subset T$  are the corresponding subobjects of  $X$ , then there is a canonical isomorphism  $U \cong T/S$ .*

*Proof.* Let  $L \subseteq I$  be an f-set, and define  $K' = \{i \in I : i \preceq j \text{ for some } j \in L\}$  and  $J' = K' \setminus L$ . Then  $J', K'$  are s-sets with  $J' \subseteq K'$  and  $L = K' \setminus J'$ , as in part (d) of Proposition 3.3. Let  $S', T' \subset X$  be the unique subobjects corresponding

to the s-sets  $J', K'$ , which exist by Proposition 3.8. Then  $S' \subset T'$  by Lemma 3.6. Define  $U$  to be the quotient  $T'/S'$ , considered as an object of  $\mathcal{A}$ . This is well-defined up to canonical isomorphism in  $\mathcal{A}$ .

By Theorem 2.6 we may refine  $0 \subset S' \subset T' \subset X$  to a composition series  $0 = B_0 \subset B_1 \subset \dots \subset B_n = X$  with  $S' = B_k$  and  $T' = B_l$  for some  $0 \leq k \leq l \leq n$ . Let  $T_m = B_m/B_{m-1}$ . Then the simple factors of  $S'$  are  $T_1, \dots, T_k$  and of  $T'$  are  $T_1, \dots, T_l$ . Also  $0 = (B_k/S') \subset \dots \subset (B_l/S') = U$  is a composition series for  $U$ . Thus the simple factors of  $U$  are  $T_{k+1}, \dots, T_l$ , that is, the simple factors in  $T'$  not appearing in  $S'$ . Hence the simple factors of  $U$  are  $S^i$  for  $i \in K' \setminus J' = L$ , and part (a) follows from Theorem 2.6.

Now let  $J \subset K$  be s-sets in  $I$  with  $K \setminus J = L$ , and  $S \subset T$  be the corresponding subobjects of  $X$ . Then it is easy to see that  $J' \subseteq J$  and  $K' \subseteq K$ , and  $J \cap K' = J'$ , and  $J \cup K' = K$ . As  $S \cap T'$  corresponds to the s-set  $J \cap K'$  by Lemma 3.5,  $J \cap K' = J'$  gives  $S \cap T' = S'$  by Lemma 3.6. Since  $S + T'$  corresponds to the s-set  $J \cup K'$  by Lemma 3.5,  $J \cup K' = K$  gives  $S + T' = T$  by Lemma 3.6.

But (4) gives a canonical isomorphism  $T'/(S \cap T') \cong (S + T')/S$ , so that  $U = T'/S' \cong T/S$ , proving part (b). Note also that any  $U$  satisfying (a) and (b) must be canonically isomorphic to  $T'/S'$  by part (b), so  $U$  is unique up to canonical isomorphism.  $\square$

## 4 Posets $(I, \preceq)$ and $(I, \preceq)$ -configurations in $\mathcal{A}$

Although a *subobject* of  $X$  is an equivalence class of injective  $i : S \rightarrow X$ , in §3 we for simplicity suppressed the morphisms  $i$ , and just wrote  $S \subset X$ . We shall now change our point of view, and investigate the natural morphisms between the factors  $T/S$  of  $X$ . Therefore we adopt some new notation, which stresses morphisms between objects. The following definition encodes the properties we expect of the factors of  $X$ , and their natural morphisms.

**Definition 4.1.** Let  $(I, \preceq)$  be a finite poset, and use the notation of Definition 3.2. Define  $\mathcal{F}_{(I, \preceq)}$  to be the set of f-sets of  $I$ . Define  $\mathcal{G}_{(I, \preceq)}$  to be the subset of  $(J, K) \in \mathcal{F}_{(I, \preceq)} \times \mathcal{F}_{(I, \preceq)}$  such that  $J \subseteq K$ , and if  $j \in J$  and  $k \in K$  with  $k \preceq j$ , then  $k \in J$ . Define  $\mathcal{H}_{(I, \preceq)}$  to be the subset of  $(J, K) \in \mathcal{F}_{(I, \preceq)} \times \mathcal{F}_{(I, \preceq)}$  such that  $K \subseteq J$ , and if  $j \in J$  and  $k \in K$  with  $k \preceq j$ , then  $j \in K$ . It is easy to show that  $\mathcal{G}_{(I, \preceq)}$  and  $\mathcal{H}_{(I, \preceq)}$  have the following properties:

- (a)  $(J, K)$  lies in  $\mathcal{G}_{(I, \preceq)}$  if and only if  $(K, K \setminus J)$  lies in  $\mathcal{H}_{(I, \preceq)}$ .
- (b) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{G}_{(I, \preceq)}$  then  $(J, L) \in \mathcal{G}_{(I, \preceq)}$ .
- (c) If  $(J, K) \in \mathcal{H}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then  $(J, L) \in \mathcal{H}_{(I, \preceq)}$ .
- (d) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$ ,  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then  $(J, J \cap L) \in \mathcal{H}_{(I, \preceq)}$ ,  $(J \cap L, L) \in \mathcal{G}_{(I, \preceq)}$ .

Define an  $(I, \preceq)$ -*configuration*  $(\sigma, \iota, \pi)$  in an abelian category  $\mathcal{A}$  to be maps  $\sigma : \mathcal{F}_{(I, \preceq)} \rightarrow \text{Obj}(\mathcal{A})$ ,  $\iota : \mathcal{G}_{(I, \preceq)} \rightarrow \text{Mor}(\mathcal{A})$ , and  $\pi : \mathcal{H}_{(I, \preceq)} \rightarrow \text{Mor}(\mathcal{A})$ , where

- (i)  $\sigma(J)$  is an object in  $\mathcal{A}$  for  $J \in \mathcal{F}_{(I, \preceq)}$ , with  $\sigma(\emptyset) = 0$ .

- (ii)  $\iota(J, K) : \sigma(J) \rightarrow \sigma(K)$  is *injective* for  $(J, K) \in \mathcal{G}_{(I, \preceq)}$ , and  $\iota(J, J) = \text{id}_{\sigma(J)}$ .
- (iii)  $\pi(J, K) : \sigma(J) \rightarrow \sigma(K)$  is *surjective* for  $(J, K) \in \mathcal{H}_{(I, \preceq)}$ , and  $\pi(J, J) = \text{id}_{\sigma(J)}$ .

These should satisfy the conditions:

- (A) Let  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and set  $L = K \setminus J$ . Then the following is exact in  $\mathcal{A}$ :

$$0 \longrightarrow \sigma(J) \xrightarrow{\iota(J, K)} \sigma(K) \xrightarrow{\pi(K, L)} \sigma(L) \longrightarrow 0. \quad (6)$$

- (B) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{G}_{(I, \preceq)}$  then  $\iota(J, L) = \iota(K, L) \circ \iota(J, K)$ .
- (C) If  $(J, K) \in \mathcal{H}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then  $\pi(J, L) = \pi(K, L) \circ \pi(J, K)$ .
- (D) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then

$$\pi(K, L) \circ \iota(J, K) = \iota(J \cap L, L) \circ \pi(J, J \cap L). \quad (7)$$

Note that (A)–(D) make sense because of properties (a)–(d), respectively.

A *morphism*  $\alpha : (\sigma, \iota, \pi) \rightarrow (\sigma', \iota', \pi')$  of  $(I, \preceq)$ -configurations in  $\mathcal{A}$  is a collection of morphisms  $\alpha(J) : \sigma(J) \rightarrow \sigma'(J)$  for each  $J \in \mathcal{F}_{(I, \preceq)}$  satisfying

$$\begin{aligned} \alpha(K) \circ \iota(J, K) &= \iota'(J, K) \circ \alpha(J) & \text{for all } (J, K) \in \mathcal{G}_{(I, \preceq)}, \text{ and} \\ \alpha(K) \circ \pi(J, K) &= \pi'(J, K) \circ \alpha(J) & \text{for all } (J, K) \in \mathcal{H}_{(I, \preceq)}. \end{aligned} \quad (8)$$

It is an *isomorphism* if  $\alpha(J)$  is an isomorphism for all  $J \in \mathcal{F}_{(I, \preceq)}$ . Morphisms compose in the obvious way.

We now show that Definition 4.1 captures the properties of the families of subobjects  $S^J \subset X$  considered in §3.

**Theorem 4.2.** *Let  $(I, \preceq)$  be a finite poset,  $\mathcal{A}$  an abelian category, and  $X \in \mathcal{A}$ . Suppose that for each s-set  $J \subseteq I$  we are given a subobject  $S^J \subset X$ , such that*

$$S^\emptyset = 0, \quad S^I = X, \quad S^A \cap S^B = S^{A \cap B} \quad \text{and} \quad S^A + S^B = S^{A \cup B} \quad (9)$$

for all s-sets  $A, B \subseteq I$ . Then there exists an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  with  $\sigma(I) = X$  such that  $\iota(J, I) : \sigma(J) \rightarrow X$  represents  $S^J \subset X$  for each s-set  $J \subseteq I$ . This  $(\sigma, \iota, \pi)$  is unique up to canonical isomorphism in  $\mathcal{A}$ .

*Proof.* Throughout (i)–(iii) and (A)–(D) will refer to Definition 4.1. We divide the proof into the following seven steps:

- Step 1.** Define  $\sigma$  and  $\iota$  on s-sets, and prove (B) for s-sets.
- Step 2.** For  $J, K$  s-sets with  $J \cap K = \emptyset$ , show  $\sigma(J \cup K) \cong \sigma(J) \oplus \sigma(K)$ .
- Step 3.** Define  $\sigma$  on f-sets and  $\pi(J, L)$  for s-sets  $J$ .
- Step 4.** Complete the definitions of  $\iota, \pi$ , and prove (A).
- Step 5.** Prove partial versions of (C), (D), mixing s-sets and f-sets.
- Step 6.** Prove (B), (C), and  $\iota(J, J) = \pi(J, J) = \text{id}_{\sigma(J)}$  in (ii) and (iii).

**Step 7.** Prove (D).

**Step 1.** For each s-set  $J \subseteq I$ , choose  $\sigma(J) \in \mathcal{A}$  and an injective morphism  $\iota(J, I) : \sigma(J) \rightarrow X$  representing  $S^J \subset X$ . Then  $\sigma(J)$  and  $\iota(J, I)$  are unique up to canonical isomorphism. In particular, choose  $\sigma(\emptyset) = 0$  as in (i),  $\sigma(I) = X$ , and  $\iota(I, I) = \text{id}_X$ . Suppose  $J \subseteq K$  are s-sets. Then (9) implies that  $S^J \subset S^K \subset X$ . Hence there exists a unique, injective  $\iota(J, K) : \sigma(J) \rightarrow \sigma(K)$  such that

$$\iota(J, I) = \iota(K, I) \circ \iota(J, K) \quad \text{for } J \subseteq K \text{ s-sets, as in (B)}. \quad (10)$$

By uniqueness the two definitions of  $\iota(J, I)$  coincide, and  $\iota(J, J) = \text{id}_{\sigma(J)}$ .

Suppose  $J \subseteq K \subseteq L$  are s-sets. Applying (10) to  $(K, L), (J, K), (J, L)$  gives

$$\iota(L, I) \circ \iota(K, L) \circ \iota(J, K) = \iota(K, I) \circ \iota(J, K) = \iota(J, I) = \iota(L, I) \circ \iota(J, L).$$

Since  $\iota(L, I)$  is injective we can cancel it from both sides, so that

$$\iota(J, L) = \iota(K, L) \circ \iota(J, K), \quad \text{for } J \subseteq K \subseteq L \text{ s-sets, as in (B)}. \quad (11)$$

**Step 2.** Let  $J, K$  be s-sets with  $J \cap K = \emptyset$ . We shall show that

$$\iota(J, J \cup K) \circ \pi_{\sigma(J)} + \iota(K, J \cup K) \circ \pi_{\sigma(K)} : \sigma(J) \oplus \sigma(K) \rightarrow \sigma(J \cup K) \quad (12)$$

is an *isomorphism*. Apply Definition 2.4 with  $\iota(J, I) : \sigma(J) \rightarrow X$  in place of  $i : S \rightarrow X$ , and  $\iota(K, I) : \sigma(K) \rightarrow X$  in place of  $j : T \rightarrow X$ . By (9) we may take  $U = \sigma(I \cap J) = \sigma(\emptyset) = 0$ ,  $V = \sigma(J \cup K)$  and  $e = \iota(J \cup K, I)$ . The definition gives  $c : \sigma(J) \rightarrow \sigma(J \cup K)$  with  $\iota(J, I) = \iota(J \cup K, I) \circ c$ , so  $c = \iota(J, J \cup K)$  by (10) and injectivity of  $\iota(J \cup K, I)$ . Similarly  $d = \iota(K, J \cup K)$ . Thus (12) is the second map in (2). As  $U = 0$ , exactness implies (12) is an isomorphism.

**Step 3.** Let  $L \subseteq I$  be an f-set which is not an s-set or a q-set, and define  $J' = \{i \in I \setminus L : l \not\leq i \text{ for all } l \in L\}$  and  $K' = J' \cup L$ . Then  $J' \subset K'$  are s-sets with  $L = K' \setminus J'$ . Choose  $\sigma(L) \in \mathcal{A}$  and a surjective  $\pi(K', L) : \sigma(K') \rightarrow \sigma(L)$  which is a *cokernel* for  $\iota(J', K') : \sigma(J') \rightarrow \sigma(K')$ . Then  $\sigma(L), \pi(K', L)$  are unique up to canonical isomorphism.

If  $L$  is an s-set then  $J', L$  are s-sets with  $J' \cap L = \emptyset$ , and Step 2 shows that  $\sigma(K') \cong \sigma(J') \oplus \sigma(L)$ , and we take  $\pi(K', L)$  to be the natural projection with  $\pi(K', L) \circ \iota(L, K') = \text{id}_{\sigma(L)}$ .

Now let  $J \subset K$  be s-sets in  $I$  with  $K \setminus J = L$ . Then  $J \subseteq J'$  and  $K \subseteq K'$ , as  $J', K'$  are defined to be as large as possible, and  $J' \cap K = J, J' \cup K' = K'$ . Let  $c : \sigma(K) \rightarrow C$  be a cokernel for  $\iota(J, K)$ , and consider the commutative diagram with rows short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \sigma(J) & \xrightarrow{\iota(J, K)} & \sigma(K) & \xrightarrow{c} & C \longrightarrow 0 \\ & & \downarrow \iota(J, J') & & \downarrow \iota(K, K') & & \downarrow h \\ 0 & \longrightarrow & \sigma(J') & \xrightarrow{\iota(J', K')} & \sigma(K') & \xrightarrow{\pi(K', L)} & \sigma(L) \longrightarrow 0, \end{array} \quad (13)$$

where  $h$  is not yet constructed. The first square of (13) commutes by (11), so

$$0 = \pi(K', L) \circ \iota(J', K') \circ \iota(J, J') = \pi(K', L) \circ \iota(K, K') \circ \iota(J, K).$$

But  $c$  is the cokernel of  $\iota(J, K)$ , so there exists a unique  $h : C \rightarrow \sigma(L)$  such that  $h \circ c = \pi(K', L) \circ \iota(K, K')$ , that is, the second square in (13) commutes.

As  $S^J = S^{J'} \cap S^K$  and  $S^{K'} = S^{J'} + S^K$  by (9), equation (2) implies that

$$0 \longrightarrow \sigma(J) \xrightarrow{\begin{array}{c} \iota_{\sigma(J')} \circ \iota(J, J') - \\ \iota_{\sigma(K)} \circ \iota(J, K) \end{array}} \sigma(J') \oplus \sigma(K) \xrightarrow{\begin{array}{c} \iota(J', K') \circ \pi_{\sigma(J')} + \\ \iota(K, K') \circ \pi_{\sigma(K)} \end{array}} \sigma(K') \longrightarrow 0$$

is exact. As the composition of the first map with  $c \circ \pi_{\sigma(K)}$  is zero we see that

$$c \circ \pi_{\sigma(K)} = l \circ (\iota(J', K') \circ \pi_{\sigma(J')} + \iota(K, K') \circ \pi_{\sigma(K)})$$

for some unique  $l : \sigma(K') \rightarrow C$ , by definition of cokernel. Composing with  $\iota_{\sigma(J')}$  gives  $l \circ \iota(J', K') = 0$ , so  $l = m \circ \pi(K', L)$  for some unique  $m : \sigma(L) \rightarrow C$ , by exactness of the bottom line of (13).

Then  $m \circ h \circ c = m \circ \pi(K', L) \circ \iota(K, K') = l \circ \iota(K, K') = c = \text{id}_C \circ c$ , so as  $c$  is surjective we have  $m \circ h = \text{id}_C$ . Also  $\pi(K', L) = h \circ l$ , since

$$\begin{aligned} h \circ l \circ (\iota(J', K') \circ \pi_{\sigma(J')} + \iota(K, K') \circ \pi_{\sigma(K)}) &= h \circ c \circ \pi_{\sigma(K)} = \\ \pi(K', L) \circ \iota(K, K') \circ \pi_{\sigma(K)} &= \pi(K', L) \circ (\iota(J', K') \circ \pi_{\sigma(J')} + \iota(K, K') \circ \pi_{\sigma(K)}), \end{aligned}$$

and  $\iota(J', K') \circ \pi_{\sigma(J')} + \iota(K, K') \circ \pi_{\sigma(K)}$  is surjective. Hence  $h \circ m \circ \pi(K', L) = h \circ l = \pi(K', L) = \text{id}_{\sigma(L)} \circ \pi(K', L)$ , and  $h \circ m = \text{id}_{\sigma(L)}$  as  $\pi(K', L)$  is surjective. Thus  $m = h^{-1}$ , and  $h$  is an *isomorphism*.

Define  $\pi(K, L) = \pi(K', L) \circ \iota(K, K')$ , so that  $\pi(K, L) = h \circ c$  as (13) is commutative. As  $h$  is an isomorphism and  $c$  is a cokernel for  $\iota(J, K)$ , we see that  $\pi(K, L) : \sigma(K) \rightarrow \sigma(L)$  is a *cokernel* for  $\iota(J, K) : \sigma(J) \rightarrow \sigma(K)$ . Hence  $\pi(K, L)$  is *surjective*, and (6) is exact when  $J \subseteq K$  are s-sets.

Suppose now that  $J, K$  are s-sets and  $L$  is an f-set with  $(J, K) \in \mathcal{G}_{(I, \succeq)}$  and  $(J, L), (K, L) \in \mathcal{H}_{(I, \succeq)}$ . Define  $K'$  using  $L$  as above. Then  $J \subseteq K \subseteq K'$  and

$$\pi(J, L) = \pi(K', L) \circ \iota(J, K') = \pi(K', L) \circ \iota(K, K') \circ \iota(J, K) = \pi(K, L) \circ \iota(J, K),$$

by (11) and the definitions of  $\pi(J, L), \pi(K, L)$ . Hence

$$\pi(J, L) = \pi(K, L) \circ \iota(K, J) \quad \text{when } J, K \text{ are s-sets.} \quad (14)$$

**Step 4.** Let  $(J, K) \in \mathcal{G}_{(I, \succeq)}$ , and define  $A' = \{i \in I \setminus K : k \not\preceq i \text{ for all } k \in K\}$ ,  $B' = A' \cup J$  and  $C' = A' \cup K$ . Then  $A' \subseteq B' \subseteq C'$  are s-sets with  $J = B' \setminus A'$ ,  $K = C' \setminus A'$ , and  $K \setminus J = C' \setminus B'$ , and they are the *largest* s-sets with this

property. Consider the commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \sigma(A') & \xrightarrow{\iota(A', B')} & \sigma(B') & \xrightarrow{\pi(B', J)} & \sigma(J) \longrightarrow 0 \\
& & \text{id}_{\sigma(A')} \parallel & & \downarrow \iota(B', C') & & \downarrow \iota(J, K) \\
0 & \longrightarrow & \sigma(A') & \xrightarrow{\iota(A', C')} & \sigma(C') & \xrightarrow{\pi(C', K)} & \sigma(K) \longrightarrow 0 \\
& & & & \downarrow \pi(C', K \setminus J) & & \downarrow \pi(K, K \setminus J) \\
& & & & \sigma(K \setminus J) & \xrightarrow{\text{id}_{\sigma(K \setminus J)}} & \sigma(K \setminus J) \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0.
\end{array} \tag{15}$$

Here solid arrows ‘ $\rightarrow$ ’ have already been defined, and dashed arrows ‘ $\dashrightarrow$ ’ remain to be constructed. The left hand square commutes by (11).

Now  $\pi(C', K \setminus J) \circ \iota(A', C') = \pi(C', K \setminus J) \circ \iota(B', C') \circ \iota(A', B') = 0$  as the middle column is exact. Since  $\pi(C', K)$  is the cokernel of  $\iota(A', C')$ , there exists a unique  $\pi(K, K \setminus J) : \sigma(K) \rightarrow \sigma(K \setminus J)$  with  $\pi(K, K \setminus J) \circ \pi(C', K) = \pi(C', K \setminus J)$ . As  $\pi(C', K \setminus J)$  is surjective,  $\pi(K, K \setminus J)$  is *surjective*, as in (iii). Thus in (15) the lower dashed arrow exists, and the lower square commutes.

Suppose  $f : D \rightarrow \sigma(B')$  is a morphism with  $\pi(C', K) \circ \iota(B', C') \circ f = 0$ . Then there exists a unique  $h : D \rightarrow \sigma(A')$  with  $\iota(A', C') \circ h = \iota(B', C') \circ f$ , as  $\iota(A', C')$  is the kernel of  $\pi(C', K)$ . But then  $\iota(B', C') \circ \iota(A', B') \circ h = \iota(B', C') \circ f$ , so  $\iota(A', B') \circ h = f$  as  $\iota(B', C')$  is injective. This implies that  $\iota(A', B')$  is the *kernel* of  $\pi(C', K) \circ \iota(B', C') : \sigma(B') \rightarrow \sigma(K)$ .

Suppose  $f : \sigma(K) \rightarrow D$  is a morphism with  $f \circ \pi(C', K) \circ \iota(B', C') = 0$ . Then there exists a unique  $h : \sigma(K \setminus J) \rightarrow D$  with  $h \circ \pi(C', K \setminus J) = f \circ \pi(C', K)$ , as  $\pi(C', K \setminus J)$  is the cokernel of  $\iota(B', C')$ . But then  $h \circ \pi(K, K \setminus J) \circ \pi(C', K) = f \circ \pi(C', K)$ , so  $h \circ \pi(K, K \setminus J) = f$  as  $\pi(C', K)$  is surjective. This implies that  $\pi(K, K \setminus J)$  is the *cokernel* of  $\pi(C', K) \circ \iota(B', C') : \sigma(B') \rightarrow \sigma(K)$ .

Now apply part (iv) of Definition 2.1 to the morphism  $\pi(C', K) \circ \iota(B', C') : \sigma(B') \rightarrow \sigma(K)$ . As this morphism has kernel  $\iota(A', B')$  and cokernel  $\pi(K, K \setminus J)$ , and  $\pi(B', J)$  is the cokernel of  $\iota(A', B')$ , this gives existence of a unique  $\iota(J, K) : \sigma(J) \rightarrow \sigma(K)$  with  $\iota(J, K) \circ \pi(B', J) = \pi(C', K) \circ \iota(B', C')$ , such that  $\iota(J, K)$  is the kernel of  $\pi(K, K \setminus J)$ . This implies that  $\iota(J, K)$  is *injective*, as in (ii), and in (15) the upper dashed arrow exists, and the upper right square commutes, and the right hand column is exact, which proves (A).

We should also check that if  $J, K$  are s-sets, the definition above gives the same answer for  $\iota(J, K)$  as Step 1, and for  $\pi(K, K \setminus J)$  as Step 3. If  $J, K$  are s-sets then  $A', J, K$  are s-sets with  $A' \cap J = A' \cap K = \emptyset$ , so Step 2 gives  $\sigma(B') \cong \sigma(A') \oplus \sigma(J)$  and  $\sigma(C') \cong \sigma(A') \oplus \sigma(K)$ . Substituting these into (15), we find the definitions are consistent.

**Step 5.** Let  $C$  be an s-set and  $D, E$  f-sets with  $(C, D), (C, E), (D, E) \in \mathcal{H}_{(i, \preceq)}$ , so that  $C \supseteq D \supseteq E$ . Apply Step 4 with  $J = D \setminus E$  and  $K = D$ . This gives  $C'$  which is the largest s-set with  $(C', D) \in \mathcal{H}_{(i, \preceq)}$ , so  $C \subseteq C'$ . Therefore

$$\pi(C, E) = \pi(C', E) \circ \iota(C, C') = \pi(D, E) \circ \pi(C', D) \circ \iota(C', C) = \pi(D, E) \circ \pi(C, D),$$

using (14) for the first and third steps, and commutativity of the bottom square in (15) for the second. Hence

$$\pi(C, E) = \pi(D, E) \circ \pi(C, D) \quad \text{for } C \text{ an s-set, as in (C)}. \quad (16)$$

Suppose  $J, K$  are s-sets and  $L$  an f-set with  $(J, K) \in \mathcal{G}_{(i, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(i, \preceq)}$ . Then  $J \cap L$  is an s-set with  $(J, J \cap L) \in \mathcal{H}_{(i, \preceq)}$  and  $(J \cap L, L) \in \mathcal{G}_{(i, \preceq)}$ . As in Step 4 with  $J, K$  replaced by  $J \cap L, L$ , define  $A' = \{i \in I \setminus L : l \not\preceq i \text{ for all } l \in L\}$ ,  $B' = A' \cup (J \cap L)$  and  $C' = A' \cup L$ . Then  $J \subseteq B'$  and  $K \subseteq C'$ . Therefore

$$\begin{aligned} \iota(J \cap L, L) \circ \pi(J, J \cap L) &= \iota(J \cap L, L) \circ \pi(B', J \cap L) \circ \iota(J, B') = \\ \pi(C', L) \circ \iota(B', C') \circ \iota(J, B') &= \pi(C', L) \circ \iota(B, C') = \\ \pi(C', L) \circ \iota(K, C') \circ \iota(J, K) &= \pi(K, L) \circ \iota(J, K), \end{aligned}$$

using (14) at the first and fifth steps, commutativity of the upper right square in (15) at the second, and (11) at the third and fourth. This proves

$$\pi(K, L) \circ \iota(J, K) = \iota(J \cap L, L) \circ \pi(J, J \cap L) \quad \text{for } J, K \text{ s-sets, as in (D)}. \quad (17)$$

**Step 6.** Suppose  $(J, K), (K, L) \in \mathcal{G}_{(i, \preceq)}$ , and define  $D = \{i \in I : i \preceq l \text{ for some } l \in L\}$ ,  $A = D \setminus L$ ,  $B = A \cup J$ , and  $C = A \cup K$ . Then  $A \subseteq B \subseteq C \subseteq D$  are s-sets, with  $J = B \setminus A$ ,  $K = C \setminus A$ , and  $L = D \setminus A$ . Therefore

$$\begin{aligned} \iota(K, L) \circ \iota(J, K) \circ \pi(B, J) &= \iota(K, L) \circ \pi(C, K) \circ \iota(B, C) = \\ \pi(D, L) \circ \iota(C, D) \circ \iota(B, C) &= \pi(D, L) \circ \iota(B, D) = \iota(J, L) \circ \pi(B, J), \end{aligned}$$

using (17) at the first, second and fourth steps with  $J = B \cap K$ ,  $K = C \cap L$  and  $J = B \cap L$  respectively, and (11) at the third. As  $\pi(B, J)$  is surjective this implies that  $\iota(K, L) \circ \iota(J, K) = \iota(J, L)$ , proving (B).

Similarly, suppose  $(J, K), (K, L) \in \mathcal{H}_{(i, \preceq)}$ , and define  $D = \{i \in I : i \preceq j \text{ for some } j \in J\}$ ,  $A = D \setminus J$ ,  $B = D \setminus K$ , and  $C = D \setminus L$ . Then  $A \subseteq B \subseteq C \subseteq D$  are s-sets, with  $J = D \setminus A$ ,  $K = D \setminus B$ , and  $L = D \setminus C$ . Therefore

$$\pi(J, L) \circ \pi(D, J) = \pi(D, L) = \pi(K, L) \circ \pi(D, K) = \pi(K, L) \circ \pi(J, K) \circ \pi(D, J),$$

using (16) three times. As  $\pi(D, J)$  is surjective, this proves (C). Applying (B), (C) with  $J = K = L$  gives  $\iota(J, J) = \pi(J, J) = \text{id}_{\sigma(J)}$ , as in (ii) and (iii).

**Step 7.** Suppose  $(J, K) \in \mathcal{G}_{(i, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(i, \preceq)}$ , and define  $C = \{i \in I : i \preceq k \text{ for some } k \in K\}$ ,  $A = C \setminus K$  and  $B = A \cup J$ . Then  $A \subseteq B \subseteq C$  are s-sets, with  $J = B \setminus A$  and  $K = C \setminus A$ . Therefore

$$\begin{aligned} \pi(K, L) \circ \iota(J, K) \circ \pi(B, J) &= \pi(K, L) \circ \pi(C, K) \circ \iota(B, C) = \pi(C, L) \circ \iota(B, C) = \\ \iota(J \cap L, L) \circ \pi(B, J \cap L) &= \iota(J \cap L, L) \circ \pi(J, J \cap L) \circ \pi(B, J), \end{aligned}$$

using (17) at the first and third steps with  $J = B \cap K$ ,  $B \cap L = J \cap L$  respectively, and (C) at the second and fourth. As  $\pi(B, J)$  is surjective this proves (D).

Hence  $(\sigma, \iota, \pi)$  is an  $(I, \preceq)$ -configuration, in the sense of Definition 4.1. It remains only to show that  $(\sigma, \iota, \pi)$  is *unique up to canonical isomorphism* in  $\mathcal{A}$ . At each stage in the construction the objects and morphisms were determined either uniquely up to canonical isomorphism, or uniquely. Thus, if  $(\sigma, \iota, \pi)$ ,  $(\sigma', \iota', \pi')$  both satisfy the conditions of the theorem, one can go through the steps above and construct a canonical isomorphism between them.  $\square$

Applying the theorem to the situation of §3 gives:

**Corollary 4.3.** *Let  $\mathcal{A}, X, I$  and  $\preceq$  be as in Definition 3.1. Then there exists an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  with  $\sigma(I) = X$  and  $\sigma(\{i\}) \cong S^i$  for  $i \in I$ , such that  $\iota(J, I) : \sigma(J) \rightarrow X$  represents the subobject of  $X$  corresponding to  $J$  under the 1-1 correspondence of Proposition 3.8 for each  $s$ -set  $J \subseteq I$ , and  $\sigma(L)$  is canonically isomorphic to  $U$  in Proposition 3.11 for each  $f$ -set  $L \subseteq I$ . This  $(\sigma, \iota, \pi)$  is unique up to canonical isomorphism in  $\mathcal{A}$ .*

We can also apply Theorem 4.2 to *filtrations*  $0 = A_0 \subset A_1 \subset \dots \subset A_n = X$ , as in Definition 2.5. Set  $I = \{1, \dots, n\}$ , with the usual total order  $\leq$ . Then the  $s$ -sets of  $(I, \leq)$  are  $\{1, 2, \dots, j\}$  for  $j = 0, 1, \dots, n$ . Define  $S^{\{1, \dots, j\}} = A_j \subset X$  for  $j = 0, \dots, n$ . These  $S^{\{1, \dots, j\}}$  satisfy (9), and Theorem 4.2 gives:

**Corollary 4.4.** *Let  $0 = A_0 \subset A_1 \subset \dots \subset A_n = X$  be a filtration in an abelian category  $\mathcal{A}$ . Set  $I = \{1, \dots, n\}$ , with the usual total order  $\leq$ . Then there is an  $(I, \leq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ , unique up to canonical isomorphism, such that  $\iota(\{1, \dots, j\}, I) : \sigma(\{1, \dots, j\}) \rightarrow X$  represents  $A_j \subset X$  for  $j = 0, \dots, n$ .*

This shows that we can regard  $(I, \preceq)$ -configurations as *generalized filtrations*. Here is the converse to Theorem 4.2.

**Theorem 4.5.** *Let  $(I, \preceq)$  be a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Define  $X = \sigma(I)$ , and let  $S^J \subset X$  be represented by  $\iota(J, I) : \sigma(J) \rightarrow X$  for each  $s$ -set  $J \subseteq I$ . Then the  $S^J$  satisfy (9).*

*Proof.* The first two equations of (9) are obvious. So suppose  $A, B \subseteq I$  are  $s$ -sets. Definition 2.4 with  $S = \sigma(A)$ ,  $T = \sigma(B)$ ,  $i = \iota(A, I)$  and  $j = \iota(B, I)$  gives  $U \in \mathcal{A}$  and  $a : U \rightarrow \sigma(A)$ ,  $b : U \rightarrow \sigma(B)$  with  $i \circ a = j \circ b$ , such that  $i \circ a : U \rightarrow X$  represents  $S^A \cap S^B$ . As  $i \circ \iota(A \cap B, A) = \iota(A \cap B, I) = j \circ \iota(A \cap B, B)$ , by exactness in (2) there is a unique  $h : \sigma(A \cap B) \rightarrow U$  with

$$(\iota_{\sigma(A)} \circ a - \iota_{\sigma(B)} \circ b) \circ h = \iota_{\sigma(A)} \circ \iota(A \cap B, A) - \iota_{\sigma(B)} \circ \iota(A \cap B, B).$$

Composing  $\pi_{\sigma(A)}, \pi_{\sigma(B)}$  gives  $\iota(A \cap B, A) = a \circ h$  and  $\iota(A \cap B, B) = b \circ h$ . Now

$$\iota(A \cup B, I) \circ \iota(A, A \cup B) \circ a = i \circ a = j \circ b = \iota(A \cup B, I) \circ \iota(B, A \cup B) \circ b,$$

by Definition 4.1(B). Thus  $\iota(A, A \cup B) \circ a = \iota(B, A \cup B) \circ b$ , as  $\iota(A \cup B, I)$  is injective. Hence

$$\begin{aligned} \pi(A, A \setminus B) \circ a &= \iota(A \setminus B, A \setminus B) \circ \pi(A, A \setminus B) \circ a = \\ \pi(A \cup B, A \setminus B) \circ \iota(A, A \cup B) \circ a &= \pi(A \cup B, A \setminus B) \circ \iota(B, A \cup B) \circ b = 0, \end{aligned}$$

using Definition 4.1(D) at the second step and exactness in (A) at the fourth.

But  $\iota(A \cap B, A)$  is the kernel of  $\pi(A, A \setminus B)$ , so there is a unique  $h' : U \rightarrow \sigma(A \cap B)$  with  $a = \iota(A \cap B, A) \circ h'$ . As  $\iota(A \cap B, A) = a \circ h$  and  $a, \iota(A \cap B, A)$  are injective we see that  $h, h'$  are *inverse*, so  $h$  is *invertible*. This implies that

$$\iota(A \cap B, I) = \iota(A, I) \circ \iota(A \cap B, A) = \iota(A, I) \circ a \circ h : \sigma(A \cap B) \rightarrow X$$

represents  $S^A \cap S^B$ , so that  $S^{A \cap B} = S^A \cap S^B$ . We prove  $S^{A \cup B} = S^A + S^B$  in a similar way.  $\square$

## 5 New $(I, \preceq)$ -configurations from old

Let  $(I, \preceq)$  be a finite poset, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ . Then we can derive  $(K, \trianglelefteq)$ -configurations  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  in  $\mathcal{A}$  from  $(\sigma, \iota, \pi)$  for other, simpler finite posets  $(K, \trianglelefteq)$ , by forgetting some of the objects and morphisms in  $(\sigma, \iota, \pi)$ . The next two definitions give two ways to do this.

**Definition 5.1.** Let  $(I, \preceq)$  be a finite poset, and use the notation of §3 and §4. Suppose  $J$  is an f-set in  $I$ , so that  $J \in \mathcal{F}_{(I, \preceq)}$ . Then  $(J, \preceq)$  is also a finite poset, and  $K \subseteq J$  is an f-set in  $(J, \preceq)$  if and only if it is an f-set in  $(I, \preceq)$ . Hence  $\mathcal{F}_{(J, \preceq)} \subseteq \mathcal{F}_{(I, \preceq)}$ . One can also show that  $\mathcal{G}_{(J, \preceq)} = \mathcal{G}_{(I, \preceq)} \cap (\mathcal{F}_{(J, \preceq)} \times \mathcal{F}_{(J, \preceq)})$  and  $\mathcal{H}_{(J, \preceq)} = \mathcal{H}_{(I, \preceq)} \cap (\mathcal{F}_{(J, \preceq)} \times \mathcal{F}_{(J, \preceq)})$ , so that  $\mathcal{G}_{(J, \preceq)} \subseteq \mathcal{G}_{(I, \preceq)}$  and  $\mathcal{H}_{(J, \preceq)} \subseteq \mathcal{H}_{(I, \preceq)}$ .

Let  $(\sigma, \iota, \pi)$  be an  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ , and define  $\sigma' : \mathcal{F}_{(J, \preceq)} \rightarrow \text{Obj}(\mathcal{A})$ ,  $\iota' : \mathcal{G}_{(J, \preceq)} \rightarrow \text{Mor}(\mathcal{A})$  and  $\pi' : \mathcal{H}_{(J, \preceq)} \rightarrow \text{Mor}(\mathcal{A})$  by  $\sigma' = \sigma|_{\mathcal{F}_{(J, \preceq)}}$ ,  $\iota' = \iota|_{\mathcal{G}_{(J, \preceq)}}$  and  $\pi' = \pi|_{\mathcal{H}_{(J, \preceq)}}$ . Then (A)–(D) of Definition 4.1 for  $(\sigma, \iota, \pi)$  imply (A)–(D) for  $(\sigma', \iota', \pi')$ , so  $(\sigma', \iota', \pi')$  is a  $(J, \preceq)$ -*configuration* in  $\mathcal{A}$ . We call  $(\sigma', \iota', \pi')$  a *subconfiguration* of  $(\sigma, \iota, \pi)$ .

**Definition 5.2.** Let  $(I, \preceq)$  and  $(K, \trianglelefteq)$  be finite posets, and  $\phi : I \rightarrow K$  a *surjective* map with  $\phi(i) \trianglelefteq \phi(j)$  when  $i, j \in I$  with  $i \preceq j$ . Use the notation of §3 and §4. If  $J \subseteq K$  is an f-set in  $K$  then  $\phi^{-1}(J) \subseteq I$  is an f-set in  $I$ . Hence  $\phi^*(\mathcal{F}_{(K, \trianglelefteq)}) \subseteq \mathcal{F}_{(I, \preceq)}$ , where  $\phi^*$  pulls back subsets of  $K$  to subsets of  $I$  in the obvious way. Similarly, if  $(A, B) \in \mathcal{G}_{(K, \trianglelefteq)}$  then  $(\phi^{-1}(A), \phi^{-1}(B)) \in \mathcal{G}_{(I, \preceq)}$ , and if  $(A, B) \in \mathcal{H}_{(K, \trianglelefteq)}$  then  $(\phi^{-1}(A), \phi^{-1}(B)) \in \mathcal{H}_{(I, \preceq)}$ .

Let  $(\sigma, \iota, \pi)$  be an  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ , and define  $\tilde{\sigma} : \mathcal{F}_{(K, \trianglelefteq)} \rightarrow \text{Obj}(\mathcal{A})$ ,  $\tilde{\iota} : \mathcal{G}_{(K, \trianglelefteq)} \rightarrow \text{Mor}(\mathcal{A})$  and  $\tilde{\pi} : \mathcal{H}_{(K, \trianglelefteq)} \rightarrow \text{Mor}(\mathcal{A})$  by  $\tilde{\sigma}(A) = \sigma(\phi^{-1}(A))$ ,  $\tilde{\iota}(A, B) = \iota(\phi^{-1}(A), \phi^{-1}(B))$ , and  $\tilde{\pi}(A, B) = \pi(\phi^{-1}(A), \phi^{-1}(B))$ . Then  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  is a  $(K, \trianglelefteq)$ -*configuration* in  $\mathcal{A}$ . We call  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  a *quotient configuration* of  $(\sigma, \iota, \pi)$ . We also call  $(\sigma, \iota, \pi)$  a *refinement* of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ , generalizing the notion of refinement of filtrations in §2.3.

Compositions of these constructions all behave in the obvious ways. Next we explain a method to *glue two configurations*  $(\sigma', \iota', \pi')$ ,  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  *together*, to get  $(\sigma, \iota, \pi)$  containing  $(\sigma', \iota', \pi')$  as a *subconfiguration*, and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  as a *quotient configuration*. Consider the following situation.

**Definition 5.3.** Let  $(J, \lesssim)$  and  $(K, \trianglelefteq)$  be finite posets and  $L \subset K$  an f-set, with  $J \cap (K \setminus L) = \emptyset$ . Suppose  $\psi : J \rightarrow L$  is a surjective map with  $\psi(i) \trianglelefteq \psi(j)$

when  $i, j \in J$  with  $i \lesssim j$ . Set  $I = J \cup (K \setminus L)$ , and define a binary relation  $\preceq$  on  $I$  by

$$i \preceq j \text{ for } i, j \in I \text{ if } \begin{cases} i \lesssim j, & i, j \in J, \\ i \trianglelefteq j, & i, j \in K \setminus L, \\ \psi(i) \trianglelefteq j, & i \in J, \quad j \in K \setminus L, \\ i \trianglelefteq \psi(j), & i \in K \setminus L, \quad j \in J. \end{cases} \quad (18)$$

It can be shown that  $\preceq$  is a *partial order* on  $I$ , and  $J \subseteq I$  is an *f-set* in  $(I, \preceq)$ . The restriction of  $\preceq$  to  $J$  is  $\lesssim$ . Define  $\phi : I \rightarrow K$  by  $\phi(i) = \psi(i)$  if  $i \in J$  and  $\phi(i) = i$  if  $i \in K \setminus L$ . Then  $\phi$  is *surjective*, with  $\phi(i) \trianglelefteq \phi(j)$  when  $i, j \in I$  with  $i \preceq j$ , as in Definition 5.2.

An  $(I, \preceq)$ -configuration gives the same  $(L, \trianglelefteq)$ -configuration in two ways.

**Lemma 5.4.** *In the situation of Definition 5.3, suppose  $\mathcal{A}$  is an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Let  $(\sigma', \iota', \pi')$  be its  $(J, \lesssim)$ -subconfiguration, and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  its quotient  $(K, \trianglelefteq)$ -configuration from  $\phi$ . Let  $(\hat{\sigma}, \hat{\iota}, \hat{\pi})$  be the quotient  $(L, \trianglelefteq)$ -configuration from  $(\sigma', \iota', \pi')$  and  $\psi$ , and  $(\check{\sigma}, \check{\iota}, \check{\pi})$  the  $(L, \trianglelefteq)$ -subconfiguration from  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . Then  $(\hat{\sigma}, \hat{\iota}, \hat{\pi}) = (\check{\sigma}, \check{\iota}, \check{\pi})$ .*

Our third construction is a kind of converse to Lemma 5.4.

**Theorem 5.5.** *In the situation of Definition 5.3, let  $\mathcal{A}$  be an abelian category,  $(\sigma', \iota', \pi')$  a  $(J, \lesssim)$ -configuration in  $\mathcal{A}$ , and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  a  $(K, \trianglelefteq)$ -configuration in  $\mathcal{A}$ . Define  $(\hat{\sigma}, \hat{\iota}, \hat{\pi})$  to be the quotient  $(L, \trianglelefteq)$ -configuration from  $(\sigma', \iota', \pi')$  and  $\psi$ , and  $(\check{\sigma}, \check{\iota}, \check{\pi})$  to be the  $(L, \trianglelefteq)$ -subconfiguration from  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ .*

*Suppose  $(\hat{\sigma}, \hat{\iota}, \hat{\pi}) = (\check{\sigma}, \check{\iota}, \check{\pi})$ . Then there exists an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ , unique up to canonical isomorphism, such that  $(\sigma', \iota', \pi')$  is its  $(J, \lesssim)$ -subconfiguration, and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  its quotient  $(K, \trianglelefteq)$ -configuration from  $\phi$ .*

*Proof.* We divide the proof into the following five steps:

**Step 1.** Characterize  $(I, \preceq)$  s-sets.

**Step 2.** Define  $\sigma(B)$  for all  $(I, \preceq)$  s-sets, and some morphisms  $\iota(B, C)$ .

**Step 3.** Define  $\iota(B, B')$  for all  $(I, \preceq)$  s-sets  $B \subseteq B' \subseteq I$ , and prove  $\iota = \iota \circ \iota$ .

**Step 4.** Let  $S^B$  be the subobject represented by  $\iota(B, I) : \sigma(B) \rightarrow \sigma(I) = X$  for all  $(I, \preceq)$  s-sets  $B$ . Show that the  $S^B$  satisfy (9).

**Step 5.** Apply Theorem 4.2 to construct  $(\sigma, \iota, \pi)$ , and complete the proof.

**Step 1.** The proof of the next lemma is elementary, and left as an exercise.

**Lemma 5.6.** *In the situation above, let  $B \subseteq I$  be an s-set in  $(I, \preceq)$ . Define*

$$P = \{k \in K : \text{if } i \in I \text{ and } \phi(i) \trianglelefteq k, \text{ then } i \in B\}, \quad \text{and} \quad (19)$$

$$R = \{k \in K : k \trianglelefteq \phi(i) \text{ for some } i \in B\}. \quad (20)$$

*Define  $A = \phi^{-1}(P)$  and  $C = \phi^{-1}(R)$ . Then  $P \subseteq R$  are  $(K, \trianglelefteq)$  s-sets, and  $A \subseteq B \subseteq C$  are  $(I, \preceq)$  s-sets, with  $P \setminus L = R \setminus L = B \setminus J$ . Define  $D = A \cap J$ ,  $E = B \cap J$  and  $F = C \cap J$ . Then  $D \subseteq E \subseteq F$  are  $(J, \lesssim)$  s-sets, with*

$$(P, R) \in \mathcal{G}_{(K, \trianglelefteq)} \quad \text{and} \quad (D, E), (D, F), (E, F) \in \mathcal{G}_{(J, \lesssim)}. \quad (21)$$

Define  $U = P \cap L$  and  $W = R \cap L$ . Then  $U \subseteq W$  are  $(L, \trianglelefteq)$  s-sets with  $\phi^{-1}(U) = \psi^{-1}(U) = D$  and  $\phi^{-1}(W) = \psi^{-1}(W) = F$ . Hence

$$\begin{aligned} \sigma'(D) = \hat{\sigma}(U) = \check{\sigma}(U) = \tilde{\sigma}(U), \text{ and similarly } \sigma'(F \setminus D) = \bar{\sigma}(W \setminus U), \\ \sigma'(F) = \tilde{\sigma}(W), \quad \iota'(D, F) = \tilde{\iota}(U, W), \quad \pi'(F, F \setminus D) = \tilde{\pi}(W, W \setminus U). \end{aligned} \quad (22)$$

Here  $P, R$  are the largest, smallest  $(K, \trianglelefteq)$  s-sets with  $\phi^{-1}(P) \subseteq B \subseteq \phi^{-1}(R)$ .

**Step 2.** Let  $B$  be an  $(I, \preceq)$  s-set, and use the notation of Lemma 5.6. As  $R \setminus P = W \setminus U$  we have  $\bar{\sigma}(R \setminus P) = \tilde{\sigma}(W \setminus U) = \sigma'(F \setminus D)$  by (22). Consider

$$\pi'(F \setminus D, F \setminus E) \circ \tilde{\pi}(R, R \setminus P) : \tilde{\sigma}(R) \rightarrow \sigma'(F \setminus E). \quad (23)$$

Choose  $\sigma(B) \in \mathcal{A}$  and  $\iota(B, C) : \sigma(B) \rightarrow \sigma(C) = \tilde{\sigma}(R)$  to be a *kernel* for (23). If  $B = \phi^{-1}(Q)$  for some  $(K, \trianglelefteq)$  s-set  $Q$  then  $B = C$  and (23) is zero, and we choose  $\sigma(B) = \tilde{\sigma}(R)$  and  $\iota(B, C) = \text{id}_{\sigma(B)}$ . Define  $\iota(B, I) = \tilde{\iota}(R, K) \circ \iota(B, C)$ .

**Step 3.** Let  $B \subseteq B'$  be  $(I, \preceq)$  s-sets. Use the notation of Lemma 5.6 for  $B$ , and  $P', R', A', C', D', E', F'$  for  $B'$  in the obvious way. Then  $P \subseteq P', R \subseteq R'$ , and so on. We have

$$\begin{aligned} & \pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P') \circ \tilde{\iota}(R, R') \circ \iota(B, C) = \\ & \pi'(F' \setminus D', F' \setminus E') \circ \tilde{\iota}(R \setminus P', R' \setminus P') \circ \tilde{\pi}(R, R \setminus P') \circ \iota(B, C) = \\ & \pi'(F' \setminus D', F' \setminus E') \circ \iota'(F \setminus D', F' \setminus D') \circ \tilde{\pi}(R \setminus P, R \setminus P') \circ \tilde{\pi}(R, R \setminus P) \circ \iota(B, C) = \\ & \iota'(F \setminus E', F' \setminus E') \circ \pi'(F \setminus D', F \setminus E') \circ \pi'(F \setminus D, F \setminus D') \circ \tilde{\pi}(R, R \setminus P) \circ \iota(B, C) = \\ & \iota'(F \setminus E', F' \setminus E') \circ \pi'(F \setminus E, F \setminus E') \circ \pi'(F \setminus D, F \setminus E) \circ \tilde{\pi}(R, R \setminus P) \circ \iota(B, C) = 0, \end{aligned}$$

using Definition 4.1(C), (D), and the definition of  $\iota(B, C)$ .

Thus, as  $\iota(B', C')$  is the kernel of  $\pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P')$ , there exists a unique  $\iota(B, B')$  with  $\tilde{\iota}(R, R') \circ \iota(B, C) = \iota(B', C') \circ \iota(B, B')$ . Hence

$$\begin{aligned} \iota(B', I) \circ \iota(B, B') &= \tilde{\iota}(R', K) \circ \iota(B', C') \circ \iota(B, B') = \\ & \tilde{\iota}(R', K) \circ \tilde{\iota}(R, R') \circ \iota(B, C) = \tilde{\iota}(R, K) \circ \iota(B, C) = \iota(B, I). \end{aligned}$$

The proof of (11) from (10) then gives

$$\iota(B, B'') = \iota(B', B'') \circ \iota(B, B') \quad \text{when } B \subseteq B' \subseteq B'' \text{ are } (I, \preceq) \text{ s-sets.} \quad (24)$$

**Step 4.** Set  $X = \sigma(I) = \tilde{\sigma}(K)$ , and for each  $(I, \preceq)$  s-set  $B$  let  $S^B \subset X$  be subobject represented by  $\iota(B, I) : \sigma(B) \rightarrow \sigma(I) = X$ . We must prove that these  $S^B$  satisfy (9). The first two equations of (9) are immediate. Let  $B', B''$  be  $(I, \preceq)$  s-sets, and  $B = B' \cap B''$ . We shall show that  $S^B = S^{B'} \cap S^{B''}$ .

Use the notation of Lemma 5.6 for  $B$ , and  $P', R', \dots$  for  $B'$  and  $P'', R'', \dots$  for  $B''$  in the obvious way. Apply Definition 2.4 with  $i = \iota(B', I)$  and  $j = \iota(B'', I)$ , giving  $U, V \in \mathcal{A}$  and morphisms  $a, b, c, d, e$  with  $i \circ a = j \circ b$ ,  $i = e \circ c$  and  $j = e \circ d$ , such that  $i \circ a : U \rightarrow X$  represents  $S^{B'} \cap S^{B''}$ .

Set  $\hat{C} = C' \cap C''$ ,  $\hat{R} = R' \cap R''$  and  $\hat{F} = F' \cap F''$ , so that  $C \subseteq \hat{C}$ ,  $R \subseteq \hat{R}$ ,  $F \subseteq \hat{F}$ ,  $\hat{C} = \phi^{-1}(\hat{R})$ , and  $\hat{F} = \hat{C} \cap J$ . Define  $\hat{a} : U \rightarrow \sigma(C')$  and  $\hat{b} : U \rightarrow \sigma(C'')$  by  $\hat{a} = \iota(B', C') \circ a$  and  $\hat{b} = \iota(B'', C'') \circ b$ . Then

$$\begin{aligned} \tilde{\iota}(R', K) \circ \hat{a} &= \tilde{\iota}(R', K) \circ \iota(B', C') \circ a = \iota(B', I) \circ a = i \circ a = \\ j \circ b &= \iota(B'', I) \circ b = \tilde{\iota}(R'', K) \circ \iota(B'', C'') \circ b = \tilde{\iota}(R'', K) \circ \hat{b}. \end{aligned}$$

As  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  is a configuration we see that  $S^{\hat{C}} = S^{C'} \cap S^{C''}$  by Theorem 4.5. Thus there is a unique  $\hat{h} : U \rightarrow \tilde{\sigma}(\hat{R})$  with  $\hat{a} = \tilde{\iota}(\hat{R}, R') \circ \hat{h}$  and  $\hat{b} = \tilde{\iota}(\hat{R}, R'') \circ \hat{h}$ .

As  $\iota(B', C')$  is the kernel of  $\pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P')$ , we have

$$\begin{aligned} 0 &= \pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P') \circ \iota(B', C') \circ a = \\ &= \pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P') \circ \hat{a} = \\ &= \pi'(F' \setminus D', F' \setminus E') \circ \tilde{\pi}(R', R' \setminus P') \circ \tilde{\iota}(\hat{R}, R') \circ \hat{h} = \dots = \\ &= \iota'(\hat{F} \setminus E', F' \setminus E') \circ \pi'(\hat{F} \setminus E, \hat{F} \setminus E') \circ \pi'(F \setminus D, \hat{F} \setminus E) \circ \tilde{\pi}(\hat{R}, \hat{R} \setminus P) \circ \hat{h}, \end{aligned}$$

by Definition 4.1(C), (D). Since  $\iota'(\hat{F} \setminus E', F' \setminus E')$  is injective this gives

$$\begin{aligned} \pi'(\hat{F} \setminus E, \hat{F} \setminus E') \circ \pi'(\hat{F} \setminus D, \hat{F} \setminus E) \circ \tilde{\pi}(\hat{R}, \hat{R} \setminus P) \circ \hat{h} &= 0, \\ \text{and } \pi'(\hat{F} \setminus E, \hat{F} \setminus E'') \circ \pi'(\hat{F} \setminus D, \hat{F} \setminus E) \circ \tilde{\pi}(\hat{R}, \hat{R} \setminus P) \circ \hat{h} &= 0, \end{aligned} \tag{25}$$

proving the second equation in the same way using  $B'', C'', \dots$

As  $(\sigma', \iota', \pi')$  is a configuration and  $E = E' \cap E''$ , one can show that

$$\iota_{\sigma'(\hat{F} \setminus E')} \circ \pi'(\hat{F} \setminus E, \hat{F} \setminus E') + \iota_{\sigma'(\hat{F} \setminus E'')} \circ \pi'(\hat{F} \setminus E, \hat{F} \setminus E'')$$

is an *injective* morphism  $\sigma'(\hat{F} \setminus E) \rightarrow \sigma'(\hat{F} \setminus E') \oplus \sigma'(\hat{F} \setminus E'')$ . Therefore

$$\pi'(\hat{F} \setminus D, \hat{F} \setminus E) \circ \tilde{\pi}(\hat{R}, \hat{R} \setminus P) \circ \hat{h} = 0, \tag{26}$$

by (25). Composing (26) with  $\pi'(\hat{F} \setminus E, \hat{F} \setminus F)$  and using Definition 4.1(C) shows that  $\tilde{\pi}(\hat{R}, \hat{R} \setminus R) \circ \hat{h} = 0$ . But  $\tilde{\iota}(R, \hat{R})$  is the kernel of  $\tilde{\pi}(\hat{R}, \hat{R} \setminus R)$ , so  $\hat{h} = \tilde{\iota}(R, \hat{R}) \circ \tilde{h}$  for some unique  $\tilde{h} : U \rightarrow \tilde{\sigma}(R) = \sigma(C)$ .

Substituting  $\hat{h} = \tilde{\iota}(R, \hat{R}) \circ \tilde{h}$  into (26) and using Definition 4.1(D) gives

$$\iota'(F \setminus E, \hat{F} \setminus E) \circ \pi'(F \setminus D, F \setminus E) \circ \tilde{\pi}(R, R \setminus P) \circ \tilde{h} = 0.$$

Hence  $\pi'(F \setminus D, F \setminus E) \circ \tilde{\pi}(R, R \setminus P) \circ \tilde{h} = 0$ , as  $\iota'(F \setminus E, \hat{F} \setminus E)$  is injective. Thus, as  $\iota(B, C)$  is the kernel of (23), there is a unique  $h : U \rightarrow \sigma(B)$  with  $\tilde{h} = \iota(B, C) \circ h$ . Then

$$\begin{aligned} \iota(B', C') \circ a &= \hat{a} = \tilde{\iota}(\hat{R}, R') \circ \hat{h} = \tilde{\iota}(\hat{R}, R') \circ \tilde{\iota}(R, \hat{R}) \circ \tilde{h} = \\ \tilde{\iota}(R, R') \circ \iota(B, C) \circ h &= \iota(B, C') \circ h = \iota(B', C') \circ \iota(B, B') \circ h \end{aligned}$$

by (24), so  $a = \iota(B, B') \circ h$  as  $\iota(B', C')$  is injective, and similarly  $b = \iota(B, B'') \circ h$ .

Recall the definition of  $i, j, U, V, a, \dots, e$  above. By (24) we have

$$\begin{aligned} e \circ c \circ \iota(B, B') &= i \circ \iota(B, B') = \iota(B', I) \circ \iota(B, B') = \iota(B, I) = \\ \iota(B'', I) \circ \iota(B, B'') &= j \circ \iota(B, B'') = e \circ d \circ \iota(B, B''). \end{aligned}$$

Since  $e$  is injective this gives  $c \circ \iota(B, B') = d \circ \iota(B, B'')$ , and hence

$$(c \circ \pi_{\sigma(B')} \oplus d \circ \pi_{\sigma(B'')}) \circ (\iota_{\sigma(B')} \circ \iota(B, B') - \iota_{\sigma(B'')} \circ \iota(B, B'')) = 0,$$

factoring via  $\sigma(B') \oplus \sigma(B'')$ . So by (2) there is a unique  $m : \sigma(B) \rightarrow U$  with

$$\iota_{\sigma(B')} \circ \iota(B, B') - \iota_{\sigma(B'')} \circ \iota(B, B'') = (\iota_{\sigma(B')} \circ a - \iota_{\sigma(B'')} \circ b) \circ m.$$

Composing with  $\pi_{\sigma(B')}$  gives  $\iota(B, B') = a \circ m$ . As  $a = \iota(B, B') \circ h$  and  $a, \iota(B, B')$  are injective, we see that  $m$  and  $h$  are *inverse*, so  $h$  is an *isomorphism*.

Since  $S^{B'} \cap S^{B''}$  is represented by  $\iota(B', I) \circ a = \iota(B', I) \circ \iota(B, B') \circ h = \iota(B, I) \circ h$  and  $S^B$  by  $\iota(B, I)$ , this proves that  $S^B = S^{B'} \cap S^{B''}$  for all  $(I, \preceq)$  s-sets  $B', B''$  and  $B = B' \cap B''$ . A similar proof shows that  $S^B = S^{B'} + S^{B''}$  when  $B = B' \cup B''$ . Hence the  $S^B$  satisfy (9).

**Step 5.** Theorem 4.2 now constructs an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$ , unique up to canonical isomorphism, from the  $S^B$ . It follows from the construction of the  $S^B$  that the  $(J, \lesssim)$ -subconfiguration of  $(\sigma, \iota, \pi)$  is canonically isomorphic to  $(\sigma', \iota', \pi')$ , and the quotient  $(K, \trianglelefteq)$ -configuration from  $\phi$  is canonically isomorphic to  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . It is not difficult to see that we can choose  $(\sigma, \iota, \pi)$  so that these sub- and quotient configurations are equal to  $(\sigma', \iota', \pi')$  and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . This completes the proof of Theorem 5.5.  $\square$

The case when  $L = \{l\}$  is one point will be particularly useful.

**Definition 5.7.** Let  $(J, \lesssim)$  and  $(K, \trianglelefteq)$  be nonempty finite posets with  $J \cap K = \emptyset$ , and  $l \in K$ . Set  $I = J \cup (K \setminus \{l\})$ , and define a partial order  $\preceq$  on  $I$  by

$$i \preceq j \quad \text{for } i, j \in I \text{ if } \begin{cases} i \lesssim j, & i, j \in J, \\ i \trianglelefteq j, & i, j \in K \setminus \{l\}, \\ l \trianglelefteq j, & i \in J, \quad j \in K \setminus \{l\}, \\ i \trianglelefteq l, & i \in K \setminus \{l\}, \quad j \in J, \end{cases} \quad (27)$$

and a surjective map  $\phi : I \rightarrow K$  by  $\phi(i) = l$  if  $i \in J$ , and  $\phi(i) = i$  if  $i \in K \setminus \{l\}$ .

Let  $\mathcal{A}$  be an abelian category,  $(\sigma', \iota', \pi')$  a  $(J, \lesssim)$ -configuration in  $\mathcal{A}$ , and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  a  $(K, \trianglelefteq)$ -configuration in  $\mathcal{A}$  with  $\sigma'(J) = \tilde{\sigma}(\{l\})$ . Then by Theorem 5.5 there exists an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ , unique up to canonical isomorphism, such that  $(\sigma', \iota', \pi')$  is its  $(J, \lesssim)$ -subconfiguration, and  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  its quotient  $(K, \trianglelefteq)$ -configuration from  $\phi$ . We call  $(\sigma, \iota, \pi)$  the *substitution of  $(\sigma', \iota', \pi')$  into  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$* .

## 6 Improvements and best configurations

We now study *quotient configurations* from  $(I, \preceq)$ ,  $(K, \trianglelefteq)$  when  $\phi : I \rightarrow K$  is a *bijection*. So we identify  $I, K$  and regard  $\preceq, \trianglelefteq$  as two partial orders on  $I$ .

**Definition 6.1.** Let  $I$  be a finite set and  $\preceq, \trianglelefteq$  partial orders on  $I$  such that if  $i \preceq j$  then  $i \trianglelefteq j$  for  $i, j \in I$ . Then we say that  $\trianglelefteq$  *dominates*  $\preceq$ , and  $\trianglelefteq$  *strictly dominates*  $\preceq$  if  $\preceq, \trianglelefteq$  are distinct. Let  $s$  be the number of pairs  $(i, j) \in I \times I$  with  $i \trianglelefteq j$  but  $i \not\preceq j$ . Then we say that  $\trianglelefteq$  *dominates*  $\preceq$  *by  $s$  steps*. Clearly,  $\trianglelefteq$  strictly dominates  $\preceq$  if and only if  $s > 0$ . Also

$$\mathcal{F}_{(I, \trianglelefteq)} \subseteq \mathcal{F}_{(I, \preceq)}, \quad \mathcal{G}_{(I, \trianglelefteq)} \subseteq \mathcal{G}_{(I, \preceq)} \quad \text{and} \quad \mathcal{H}_{(I, \trianglelefteq)} \subseteq \mathcal{H}_{(I, \preceq)}. \quad (28)$$

We shall see below that for distinct  $\preceq, \trianglelefteq$  the second two inclusions are strict.

For each  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in an abelian category  $\mathcal{A}$  we have a quotient  $(I, \trianglelefteq)$ -configuration  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ , as in Definition 5.2 with  $\phi = \text{id} : I \rightarrow I$ . We call  $(\sigma, \iota, \pi)$  an *improvement* or an  $(I, \preceq)$ -*improvement* of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ , and a *strict improvement* if  $\preceq, \trianglelefteq$  are distinct. If  $\trianglelefteq$  dominates  $\preceq$  by  $s$  steps we also call  $(\sigma, \iota, \pi)$  an  *$s$  step improvement* of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ .

We call an  $(I, \trianglelefteq)$ -configuration  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  *best* if there exists no strict improvement  $(\sigma, \iota, \pi)$  of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . Note that improvements are a special kind of *refinement*, in the sense of Definition 5.2.

Our first result is simple. An  $(I, \trianglelefteq)$ -configuration  $(\sigma, \iota, \pi)$  cannot have an infinite sequence of strict improvements, as  $I$  has finitely many partial orders. Thus, after finitely many steps we must reach an  $(I, \preceq)$ -configuration with no strict improvements, that is, a *best* configuration. This gives:

**Lemma 6.2.** *Let  $(I, \trianglelefteq)$  be a finite poset, and  $(\sigma, \iota, \pi)$  an  $(I, \trianglelefteq)$ -configuration in an abelian category  $\mathcal{A}$ . Then  $(\sigma, \iota, \pi)$  can be improved to a best  $(I, \preceq)$ -configuration  $(\sigma', \iota', \pi')$ , for some partial order  $\preceq$  on  $I$  dominated by  $\trianglelefteq$ .*

After some preliminary results on partial orders in §6.1, section 6.2 proves a criterion for *best configurations* in terms of *split* short exact sequences.

### 6.1 Partial orders $\trianglelefteq, \preceq$ where $\trianglelefteq$ dominates $\preceq$

We study partial orders  $\trianglelefteq, \preceq$  on  $I$  where  $\trianglelefteq$  strictly dominates  $\preceq$ .

**Lemma 6.3.** *Let  $\trianglelefteq, \preceq$  be partial orders on a finite set  $I$ , where  $\trianglelefteq$  strictly dominates  $\preceq$ . Then there exist  $i, j \in I$  with  $i \trianglelefteq j$  and  $i \not\preceq j$ , such that there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \trianglelefteq k \trianglelefteq j$ . Also*

$$(\{j\}, \{i, j\}) \in \mathcal{G}_{(I, \preceq)} \setminus \mathcal{G}_{(I, \trianglelefteq)} \quad \text{and} \quad (\{i, j\}, \{i\}) \in \mathcal{H}_{(I, \preceq)} \setminus \mathcal{H}_{(I, \trianglelefteq)}. \quad (29)$$

*Proof.* As  $\trianglelefteq$  strictly dominates  $\preceq$  there exist  $i, j \in I$  with  $i \trianglelefteq j$  and  $i \not\preceq j$ . Suppose there exists  $k \in I$  with  $i \neq k \neq j$  and  $i \trianglelefteq k \trianglelefteq j$ . Then as  $i \not\preceq j$  either

(a)  $i \not\leq k$ , or (b)  $k \not\leq j$ . In case (a) we replace  $j$  by  $k$ , and in case (b) we replace  $i$  by  $k$ . Then the new  $i, j$  satisfy the original conditions, but are ‘closer together’ than the old  $i, j$ . As  $I$  is finite, repeating this process finitely many times we reach  $i, j$  for which there exists no such  $k$ . Equation (29) easily follows.  $\square$

This implies that if  $\trianglelefteq$  strictly dominates  $\preceq$  then  $\mathcal{G}_{(I, \trianglelefteq)} \subseteq \mathcal{G}_{(I, \preceq)}$  and  $\mathcal{H}_{(I, \trianglelefteq)} \subseteq \mathcal{H}_{(I, \preceq)}$  in (28) are *strict* inclusions. But  $\mathcal{F}_{(I, \trianglelefteq)} \subseteq \mathcal{F}_{(I, \preceq)}$  need not be strict. For example, if  $I = \{1, 2\}$  with  $1 \trianglelefteq 2$  the only nontrivial relation, then  $\mathcal{F}_{(I, \trianglelefteq)} = \mathcal{F}_{(I, \preceq)}$  are both the set of all subsets of  $I$ .

The following elementary lemma characterizes  $\trianglelefteq, \preceq$  differing by one step.

**Lemma 6.4.** *Let  $(I, \trianglelefteq)$  be a finite poset, and suppose  $i \neq j \in I$  with  $i \trianglelefteq j$  but there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \trianglelefteq k \trianglelefteq j$ . Define  $\preceq$  on  $I$  by  $a \preceq b$  if and only if  $a \trianglelefteq b$  and  $a \neq i, b \neq j$ . Then  $\preceq$  is a partial order and  $\trianglelefteq$  dominates  $\preceq$  by one step. Conversely, if  $\preceq$  is a partial order and  $\trianglelefteq$  dominates  $\preceq$  by one step then  $\preceq$  arises as above for some unique  $i, j \in I$ .*

If  $\trianglelefteq$  dominates  $\preceq$  by  $s$  steps, we can interpolate a chain of  $s + 1$  partial orders differing by one step.

**Proposition 6.5.** *Let  $I$  be a finite set and  $\preceq, \trianglelefteq$  partial orders on  $I$ , where  $\trianglelefteq$  dominates  $\preceq$  by  $s$  steps. Then there exist partial orders  $\trianglelefteq = \lesssim_0, \lesssim_1, \dots, \lesssim_s = \preceq$  on  $I$  such that  $\lesssim_{r-1}$  dominates  $\lesssim_r$  by one step, for  $r = 1, \dots, s$ .*

*Proof.* Define  $\lesssim_0 = \trianglelefteq$ . Suppose by induction that  $\lesssim_0, \dots, \lesssim_m$  have been chosen for  $0 \leq m < s$  such that  $\lesssim_{r-1}$  dominates  $\lesssim_r$  by one step and  $\lesssim_r$  dominates  $\preceq$  by  $s - r$  steps for  $r = 1, \dots, m$ . Then  $\lesssim_m$  strictly dominates  $\preceq$  as  $s - m > 0$ , so by Lemma 6.3 there exist  $i, j \in I$  with  $i \lesssim_m j$  but  $i \not\leq j$ , and such that there exist no  $i \neq k \neq j$  with  $i \lesssim_m k \lesssim_m j$ .

Define  $\lesssim_{m+1}$  by  $a \lesssim_{m+1} b$  if  $a \lesssim_m b$  and  $a \neq i, b \neq j$ . Then  $\lesssim_{m+1}$  is a partial order on  $I$  and  $\lesssim_m$  dominates  $\lesssim_{m+1}$  by one step by Lemma 6.4, and  $\lesssim_{m+1}$  dominates  $\preceq$  by  $s - m - 1$  steps. Therefore by induction  $\lesssim_0, \dots, \lesssim_s$  exist, and as  $\lesssim_s$  dominates  $\preceq$  by 0 steps we have  $\lesssim = \preceq$ .  $\square$

## 6.2 Best $(I, \preceq)$ -configurations and split sequences

We now prove a criterion for *best*  $(I, \trianglelefteq)$ -configurations. First we decompose certain objects  $\sigma(J \cup K)$  as *direct sums*  $\sigma(J) \oplus \sigma(K)$ .

**Proposition 6.6.** *Suppose  $(I, \preceq)$  is a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Let  $J, K \in \mathcal{F}_{(I, \preceq)}$  with  $j \not\leq k$  and  $k \not\leq j$  for all  $j \in J$  and  $k \in K$ . Then  $J \cup K \in \mathcal{F}_{(I, \preceq)}$  is an  $f$ -set and there is a canonical isomorphism  $\sigma(J) \oplus \sigma(K) \cong \sigma(J \cup K)$  identifying  $\iota_{\sigma(J)}, \iota_{\sigma(K)}, \pi_{\sigma(J)}, \pi_{\sigma(K)}$  with  $\iota(J, J \cup K), \iota(K, J \cup K), \pi(J \cup K, J), \pi(J \cup K, K)$  respectively. Hence*

$$\iota(J, J \cup K) \circ \pi(J \cup K, J) + \iota(K, J \cup K) \circ \pi(J \cup K, K) = \text{id}_{\sigma(J \cup K)}. \quad (30)$$

*Proof.* The conditions on  $J, K$  imply that  $J \cap K = \emptyset$  and  $J \cup K \in \mathcal{F}_{(i, \preceq)}$  with  $(J, J \cup K), (K, J \cup K) \in \mathcal{G}_{(i, \preceq)}$  and  $(J \cup K, J), (J \cup K, K) \in \mathcal{H}_{(i, \preceq)}$ . Definition 4.1(A) applied to  $(J, J \cup K)$  shows that  $\pi(J \cup K, K) \circ \iota(J, J \cup K) = 0$ , and similarly  $\pi(J \cup K, J) \circ \iota(K, J \cup K) = 0$ . Parts (ii), (iii) and (D) of Definition 4.1 with  $J, J \cup K, J$  in place of  $J, K, L$  give  $\pi(J \cup K, J) \circ \iota(J, J \cup K) = \iota(J, J) \circ \pi(J, J) = \text{id}_{\sigma(J)}$ , and similarly  $\pi(J \cup K, K) \circ \iota(K, J \cup K) = \text{id}_{\sigma(K)}$ . The proposition then quickly follows from Popescu [9, Cor. 2.7.4, p. 48].  $\square$

Recall from Definition 2.2 that a short exact sequence  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$  in  $\mathcal{A}$  is called *split* if there is a compatible isomorphism  $Y \cong X \oplus Z$ .

**Proposition 6.7.** *Suppose  $(I, \trianglelefteq)$  is a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \trianglelefteq)$ -configuration in  $\mathcal{A}$  which is not best. Then there exist  $i \neq j \in I$  with  $i \trianglelefteq j$  but there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \trianglelefteq k \trianglelefteq j$ , such that the following short exact sequence is split:*

$$0 \longrightarrow \sigma(\{i\}) \xrightarrow{\iota(\{i\}, \{i, j\})} \sigma(\{i, j\}) \xrightarrow{\pi(\{i, j\}, \{j\})} \sigma(\{j\}) \longrightarrow 0. \quad (31)$$

*Proof.* As  $(\sigma, \iota, \pi)$  is not best it has a strict  $(I, \preceq)$ -improvement  $(\sigma', \iota', \pi')$ , for some  $\preceq$  dominated by  $\trianglelefteq$ . Let  $i, j$  be as in Lemma 6.3. Then  $i \neq j$  as  $i \not\trianglelefteq j$ , and there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \preceq k \preceq j$ . As  $i \not\trianglelefteq j, j \not\trianglelefteq i$  Proposition 6.6 shows that  $\sigma'(\{i, j\}) \cong \sigma'(\{i\}) \oplus \sigma'(\{j\})$ . But  $\sigma'(\{i\}) = \sigma(\{i\}), \sigma'(\{i, j\}) = \sigma(\{i, j\}), \sigma'(\{j\}) = \sigma(\{j\})$ , so  $\sigma(\{i, j\}) \cong \sigma(\{i\}) \oplus \sigma(\{j\})$ .

Proposition 6.6 and equalities between  $\iota, \iota'$  and  $\pi, \pi'$  show that the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \sigma(\{i\}) & \xrightarrow{\iota_{\sigma(\{i\})}} & \sigma(\{i\}) \oplus \sigma(\{j\}) & \xrightarrow{\pi_{\sigma(\{j\})}} & \sigma(\{j\}) \longrightarrow 0 \\ & & \downarrow \text{id}_{\sigma(\{i\})} & & \downarrow h & & \downarrow \text{id}_{\sigma(\{j\})} \\ 0 & \longrightarrow & \sigma(\{i\}) & \xrightarrow{\iota(\{i\}, \{i, j\})} & \sigma(\{i, j\}) & \xrightarrow{\pi(\{i, j\}, \{j\})} & \sigma(\{j\}) \longrightarrow 0 \end{array}$$

commutes, where  $h = \iota(\{i\}, \{i, j\}) \circ \pi_{\sigma(\{i\})} + \iota'(\{j\}, \{i, j\}) \circ \pi_{\sigma(\{j\})}$  is an isomorphism. Therefore by (1), the short exact sequence (6) is split.  $\square$

We *classify improvements* for a two point indexing set  $K = \{i, j\}$ .

**Lemma 6.8.** *Define partial orders  $\trianglelefteq, \lesssim$  on  $K = \{i, j\}$  by  $i \trianglelefteq i, i \trianglelefteq j, j \trianglelefteq j, i \lesssim i$  and  $j \lesssim j$ . Let  $(\sigma, \iota, \pi)$  be a  $(K, \trianglelefteq)$ -configuration in an abelian category  $\mathcal{A}$ . Then there exists a  $(K, \lesssim)$ -improvement  $(\sigma', \iota', \pi')$  of  $(\sigma, \iota, \pi)$  if and only if the short exact sequence (31) is split, and then such  $(K, \lesssim)$ -improvements  $(\sigma', \iota', \pi')$  are in 1-1 correspondence with  $\text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$ .*

*Proof.* If there exists a  $(K, \lesssim)$ -improvement of  $(\sigma, \iota, \pi)$  then (31) is split by Proposition 6.7, which proves the ‘only if’ part. For the ‘if’ part, suppose (31) is split. Then we can choose morphisms  $\iota'(\{j\}, K) : \sigma(\{j\}) \rightarrow \sigma(K)$  and  $\pi'(K, \{i\}) : \sigma(K) \rightarrow \sigma(\{i\})$  with

$$\pi'(K, \{i\}) \circ \iota(\{i\}, K) = \text{id}_{\sigma(\{i\})} \quad \text{and} \quad \pi(K, \{j\}) \circ \iota'(\{j\}, K) = \text{id}_{\sigma(\{j\})}. \quad (32)$$

Defining  $\sigma' = \sigma$ ,  $\iota'|_{\mathcal{G}(K, \trianglelefteq)} = \iota$ ,  $\pi'|_{\mathcal{H}(K, \trianglelefteq)} = \pi$  then gives a  $(K, \lesssim)$ -improvement  $(\sigma', \iota', \pi')$  of  $(\sigma, \iota, \pi)$ , proving the ‘if’ part.

Finally, fix  $\iota'_0(\{j\}, K)$ ,  $\pi'_0(K, \{i\})$  satisfying (32). We can easily prove that every  $(K, \lesssim)$ -improvement  $(\sigma', \iota', \pi')$  of  $(\sigma, \iota, \pi)$  is defined uniquely by  $\sigma' = \sigma$ ,  $\iota'|_{\mathcal{G}(K, \trianglelefteq)} = \iota$ ,  $\pi'|_{\mathcal{H}(K, \trianglelefteq)} = \pi$  and

$$\iota'(\{j\}, K) = \iota'_0(\{j\}, K) + \iota(\{i\}, K) \circ f, \quad \pi'(K, \{i\}) = \pi'_0(K, \{i\}) - f \circ \pi(K, \{j\})$$

for some unique  $f \in \text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$ , and every  $f \in \text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$  gives a  $(K, \lesssim)$ -improvement. This establishes a 1-1 correspondence between  $(K, \lesssim)$ -improvements  $(\sigma', \iota', \pi')$  and  $f \in \text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$ .  $\square$

Here is the converse to Proposition 6.7.

**Proposition 6.9.** *Suppose  $(I, \trianglelefteq)$  is a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \trianglelefteq)$ -configuration in  $\mathcal{A}$ . Let  $i \neq j \in I$  with  $i \trianglelefteq j$  but there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \trianglelefteq k \trianglelefteq j$ , such that the short exact sequence (31) is split. Define a partial order  $\preceq$  on  $I$  by  $a \preceq b$  if  $a \trianglelefteq b$  and  $a \neq i, b \neq j$ , so that  $\trianglelefteq$  dominates  $\preceq$  by one step. Then there exists an  $(I, \preceq)$ -improvement  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  of  $(\sigma, \iota, \pi)$ . Such improvements up to canonical isomorphism are in 1-1 correspondence with  $\text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$ .*

*Proof.* Set  $K = \{i, j\}$ , and let  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  be the  $(K, \trianglelefteq)$ -subconfiguration of  $(\sigma, \iota, \pi)$ . As (31) is split, Lemma 6.8 shows that there exists a  $(K, \lesssim)$ -improvement  $(\sigma', \iota', \pi')$  of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . Then  $(\sigma, \iota, \pi)$  and  $(\sigma', \iota', \pi')$  satisfy the conditions of Theorem 5.5 with  $\phi = \text{id}$ ,  $I$  in place of  $K$ , and  $K$  in place of both  $J$  and  $L$ . Therefore Theorem 5.5 gives the  $(I, \preceq)$ -improvement  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  that we want.

For the last part, note that every  $(I, \preceq)$ -improvement  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  of  $(\sigma, \iota, \pi)$  may be constructed this way, taking  $(\sigma', \iota', \pi')$  to be the  $(K, \preceq)$ -subconfiguration of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . Thus, uniqueness up to canonical isomorphism in Theorem 5.5 shows that such improvements  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  up to canonical isomorphism are in 1-1 correspondence with  $(K, \lesssim)$ -improvements  $(\sigma', \iota', \pi')$  of  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$ . But Lemma 6.8 shows that these are in 1-1 correspondence with  $\text{Hom}(\sigma(\{j\}), \sigma(\{i\}))$ .  $\square$

Propositions 6.7 and 6.9 imply a *criterion for best configurations*:

**Theorem 6.10.** *Let  $(\sigma, \iota, \pi)$  be an  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ . Then  $(\sigma, \iota, \pi)$  is best if and only if for all  $i \neq j \in I$  with  $i \preceq j$  but there exists no  $k \in I$  with  $i \neq k \neq j$  and  $i \preceq k \preceq j$ , the short exact sequence (31) is split.*

If this criterion holds, it also holds for any *subconfiguration* of  $(\sigma, \iota, \pi)$ , giving:

**Corollary 6.11.** *Suppose  $(\sigma, \iota, \pi)$  is a best  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ . Then all subconfigurations of  $(\sigma, \iota, \pi)$  are also best.*

## 7 Further topics

We finish with short discussions of five other subjects. Section 7.1 studies the classes of the objects  $\sigma(J)$  in the *Grothendieck group*  $K_0(\mathcal{A})$ , and §7.2 configurations in *exact categories*. Interpretations of  $(I, \preceq)$ -configurations in terms of *flasque sheaves* on  $I$  are given in §7.3, and as *functors*  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  for a category  $\mathcal{C}_{(I, \preceq)}$  constructed from  $(I, \preceq)$  in §7.4–§7.5. Finally, §7.6–§7.7 show that  $(I, \preceq)$ -configurations in an abelian or exact category  $\mathcal{A}$  form an *exact category*  $\mathbf{Conf}(I, \preceq, \mathcal{A})$ .

### 7.1 Configurations and $K_0(\mathcal{A})$

Let  $\mathcal{A}$  be an abelian category and  $K_0(\mathcal{A})$  its *Grothendieck group*, as in Definition 2.2. Let  $(\sigma, \iota, \pi)$  be an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Then each object  $\sigma(J)$  for  $J \in \mathcal{F}_{(I, \preceq)}$  has a class  $[\sigma(J)]$  in  $K_0(\mathcal{A})$ . The following proposition shows how these classes are related.

**Proposition 7.1.** *Suppose  $(I, \preceq)$  is a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Then there exists a unique map  $\kappa : I \rightarrow K_0(\mathcal{A})$  such that  $[\sigma(J)] = \sum_{j \in J} \kappa(j)$  in  $K_0(\mathcal{A})$  for all  $f$ -sets  $J \subseteq I$ .*

*Proof.* Combining Definitions 2.2 and 3.1(A) shows that

$$[\sigma(K)] = [\sigma(J)] + [\sigma(K \setminus J)] \quad \text{for all } (J, K) \in \mathcal{G}_{(I, \preceq)}. \quad (33)$$

Define  $\kappa : I \rightarrow K_0(\mathcal{A})$  by  $\kappa(i) = [\sigma(\{i\})]$ . As  $\{i\} \in \mathcal{F}_{(I, \preceq)}$  for all  $i \in I$  this is well-defined, and also any  $\kappa$  satisfying  $[\sigma(J)] = \sum_{j \in J} \kappa(j)$  for  $J \in \mathcal{F}_{(I, \preceq)}$  has  $\kappa(i) = [\sigma(\{i\})]$ , so  $\kappa$  is *unique*.

Suppose  $K \in \mathcal{F}_{(I, \preceq)}$  with  $|K| \geq 1$ . Let  $j \in K$  be *minimal*. Then  $(\{j\}, K) \in \mathcal{G}_{(I, \preceq)}$ , so (33) gives  $[\sigma(K)] = \kappa(j) + [\sigma(K \setminus \{j\})]$ . We then easily prove  $[\sigma(J)] = \sum_{j \in J} \kappa(j)$  for all  $J \in \mathcal{F}_{(I, \preceq)}$  by induction on  $|J|$ , completing the proof.  $\square$

### 7.2 Configurations in exact categories

Let  $(\mathcal{B}, \mathcal{E})$  be an *exact category*, as in §2.4. Then Definition 4.1 makes sense in  $(\mathcal{B}, \mathcal{E})$ : we take  $\sigma(J) \in \text{Obj}(\mathcal{B})$ ,  $\iota(J, K), \pi(J, K) \in \text{Mor}(\mathcal{B})$ , and use  $\mathcal{E}$  to say what it means for (6) to be exact. Thus we have a concept of a  $(I, \preceq)$ -configuration in an exact category  $(\mathcal{B}, \mathcal{E})$ . If  $\mathcal{B}$  is embedded in an abelian category  $\mathcal{A}$ , then an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $(\mathcal{B}, \mathcal{E})$  is just an  $(I, \preceq)$ -configuration in  $\mathcal{A}$  with  $\sigma(J) \in \text{Obj}(\mathcal{B})$  for all  $J \in \mathcal{F}_{(I, \preceq)}$ .

Here is an interesting example we will study in [3, 4, 5]. Let  $P$  be a projective  $\mathbb{K}$ -scheme over an algebraically closed field  $\mathbb{K}$ . Then the *abelian category*  $\text{coh}(P)$  of coherent sheaves on  $P$  contains as a subcategory the *exact category*  $\text{vect}(P)$  of vector bundles on  $P$  (that is, locally free sheaves on  $P$  of finite rank). Thus we can form  $(I, \preceq)$ -configurations of vector bundles on  $P$ , which will be useful tools for studying *stability* of vector bundles on  $P$ .

All the material of §3–§6 extends simply to exact categories  $(\mathcal{B}, \mathcal{E})$ . In particular, *subconfigurations* and *quotient configurations* in §5 are well-defined in  $(\mathcal{B}, \mathcal{E})$ . Less trivial is the extension of Theorem 5.5 to exact categories:

**Proposition 7.2.** *Theorem 5.5 holds in an exact category  $(\mathcal{B}, \mathcal{E})$ . That is, if  $\mathcal{B}$  is embedded in an abelian category  $\mathcal{A}$ , and  $(\sigma', \iota', \pi')$ ,  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  in Theorem 5.5 are configurations in  $(\mathcal{B}, \mathcal{E})$ , and so in  $\mathcal{A}$ , then the  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  constructed in Theorem 5.5 is a configuration in  $(\mathcal{B}, \mathcal{E})$ .*

*Proof.* Theorem 5.5 applies in the abelian category  $\mathcal{A}$ , and yields an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ . We must show that  $\sigma(J) \in \text{Obj}(\mathcal{B})$  for all  $J \in \mathcal{F}_{(I, \preceq)}$ , so that  $(\sigma, \iota, \pi)$  is a configuration in  $(\mathcal{B}, \mathcal{E})$ . The theorem then holds in  $(\mathcal{B}, \mathcal{E})$ , independently of the choice of enveloping abelian category  $\mathcal{A}$ .

As  $(\sigma', \iota', \pi')$  is the  $(J, \lesssim)$ -subconfiguration of  $(\sigma, \iota, \pi)$  we have  $\sigma(\{i\}) = \sigma'(\{i\}) \in \text{Obj}(\mathcal{B})$  for  $i \in J$ . And as  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  is the quotient  $(K, \preceq)$ -configuration of  $(\sigma, \iota, \pi)$  from  $\phi$  we have  $\sigma(\{i\}) = \tilde{\sigma}(\{i\}) \in \text{Obj}(\mathcal{B})$  for  $i \in I \setminus J = K \setminus L$ . Hence  $\sigma(\{i\}) \in \text{Obj}(\mathcal{B})$  for all  $i \in I$ . Also  $\sigma(\emptyset) = 0 \in \text{Obj}(\mathcal{B})$ . Hence  $\sigma(A) \in \text{Obj}(\mathcal{B})$  for all  $A \in \mathcal{F}_{(I, \preceq)}$  with  $|A| \leq 1$ .

Suppose by induction that  $\sigma(A) \in \text{Obj}(\mathcal{B})$  for all  $A \in \mathcal{F}_{(I, \preceq)}$  with  $|A| \leq k$ , for  $1 \leq k < |I|$ . Let  $B \in \mathcal{F}_{(I, \preceq)}$  with  $|B| = k + 1$ , let  $i$  be  $\preceq$ -maximal in  $B$ , and set  $A = B \setminus \{i\}$ . Then  $(A, B) \in \mathcal{G}_{(I, \preceq)}$ , so (6) gives a *short exact sequence*  $0 \rightarrow \sigma(A) \rightarrow \sigma(B) \rightarrow \sigma(\{i\}) \rightarrow 0$ . Now  $\sigma(A) \in \text{Obj}(\mathcal{B})$  by induction, and  $\sigma(\{i\}) \in \text{Obj}(\mathcal{B})$  from above, and  $\mathcal{B}$  is *closed under extensions* as it is an exact category. Therefore  $\sigma(B) \in \text{Obj}(\mathcal{B})$ . So by induction  $\sigma(A) \in \text{Obj}(\mathcal{B})$  for all  $A \in \mathcal{F}_{(I, \preceq)}$ , and  $(\sigma, \iota, \pi)$  is an  $(I, \preceq)$ -configuration in  $(\mathcal{B}, \mathcal{E})$ .  $\square$

### 7.3 Configurations and flasque sheaves

We briefly describe an alternative point of view on configurations, explained to me by Tom Bridgeland. Let  $(I, \preceq)$  be a finite poset, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in an abelian category  $\mathcal{A}$ . Then as in §3 the q-sets in  $I$  are the open sets of a *topology* on  $I$ .

It can be shown that the data  $\sigma(J)$  and  $\pi(J, K) : \sigma(J) \rightarrow \sigma(K)$  for all q-sets  $I \supseteq J \supseteq K$  comprises a *flasque sheaf*  $\mathcal{S}$  on  $I$  with the q-set topology, with values in  $\mathcal{A}$ . The rest of the data  $(\sigma, \iota, \pi)$  can be reconstructed, up to canonical isomorphism, from the  $\sigma(J)$  and  $\pi(J, K)$  for q-sets  $J, K$  — this is essentially Theorems 4.2 and 4.5 with s-sets replaced by q-sets.

The f-sets  $J \subseteq I$  are the *locally closed sets* in the q-set topology. In fact, a topology on a finite set  $I$  comes from a (unique) partial order  $\preceq$  if and only if every point is locally closed. For f-sets  $J$  we interpret  $\sigma(J)$  as the sections of  $\mathcal{S}$  near  $J$  locally supported on  $J$ .

Alternatively, we can take the s-sets to be the open sets of a topology on  $I$ , which is more compatible with §3 and §4. Then  $(\sigma, \iota, \pi)$  is equivalent to a *flasque cosheaf* with values in  $\mathcal{A}$ , or equivalently, a flasque sheaf with values in the *opposite category*  $\mathcal{A}^\circ$ .

Probably one could use this to reduce parts of our theory to known facts on sheaves, and so shorten the proofs. In particular, Theorems 4.2 and 5.5 look like

instances of general sheaf results. But I have been unable to find appropriate references. Note however that Baclawski [1, §1] studies sheaves on posets with the q-set topology. He calls q-sets *increasing subsets*, or *order filters*.

#### 7.4 Compositions of morphisms $\iota(*, *)$ , $\pi(*, *)$

In §7.5 we shall show that an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  is equivalent to an exact functor  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  for  $\mathcal{C}_{(I, \preceq)}$  a category constructed using  $(I, \preceq)$ . As a preparation for this, we must first study the collection of morphisms  $f : \sigma(A) \rightarrow \sigma(B)$  obtained by compositions of morphisms  $\iota(*, *)$ ,  $\pi(*, *)$ . The results below and in §7.5 are elementary and will not be used in the sequels [3, 4, 5], so to save space we leave all proofs as an exercise for the reader.

**Proposition 7.3.** *Suppose  $(I, \preceq)$  is a finite poset,  $\mathcal{A}$  an abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ , with  $\sigma(K) \neq \sigma(L)$  for  $K \neq L$ . Then every morphism  $f : \sigma(A) \rightarrow \sigma(B)$  which is a composition of finitely many  $\iota(*, *)$ ,  $\pi(*, *)$  may be written  $\iota(J, B) \circ \pi(A, J)$  for some  $J \subseteq A \cap B$  with  $(A, J) \in \mathcal{H}_{(I, \preceq)}$  and  $(J, B) \in \mathcal{G}_{(I, \preceq)}$ .*

Here we assume  $\sigma(K) \neq \sigma(L)$  to avoid compositions like  $\iota(L, M) \circ \iota(J, K)$ .

**Definition 7.4.** Let  $(I, \preceq)$  be a finite poset, and  $A, B \in \mathcal{F}_{(I, \preceq)}$ . Define an equivalence relation  $\sim_{A, B}$  on  $A \cap B$  by  $i \sim_{A, B} j$  for  $i, j \in A \cap B$  if there exist  $i = i_0, i_1, \dots, i_n = j$  with either  $i_{m-1} \preceq i_m$  or  $i_m \preceq i_{m-1}$  for  $m = 1, \dots, n$ .

Define  $\mathcal{P}(A, B)$  to be the set of  $\sim_{A, B}$ -equivalence classes  $J \subseteq A \cap B$  satisfying

- (a) there do not exist  $i \in A \setminus B$  and  $j \in J$  with  $j \preceq i$ , and
- (b) there do not exist  $i \in B \setminus A$  and  $j \in J$  with  $i \preceq j$ .

Then  $\mathcal{P}(A, B)$  is a finite set of disjoint subsets of  $A \cap B$ .

We use  $\mathcal{P}(A, B)$  to classify  $J \in \mathcal{F}_{(I, \preceq)}$  with  $(A, J) \in \mathcal{H}_{(I, \preceq)}$ ,  $(J, B) \in \mathcal{G}_{(I, \preceq)}$ , and the compositions  $\iota(J, B) \circ \pi(A, J)$ .

**Proposition 7.5.** *Let  $(I, \preceq)$  be a finite poset,  $A, B \in \mathcal{F}_{(I, \preceq)}$ , and  $J \subseteq I$ . Then  $(A, J) \in \mathcal{G}_{(I, \preceq)}$  and  $(J, B) \in \mathcal{H}_{(I, \preceq)}$  if and only if  $J = \coprod_{K \in R} K$  for some unique  $R \subseteq \mathcal{P}(A, B)$ . In this case, let  $(\sigma, \iota, \pi)$  be an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Then*

$$\iota(J, B) \circ \pi(A, J) = \sum_{K \in R} \iota(K, B) \circ \pi(A, K).$$

Propositions 7.3 and 7.5 show that all morphisms  $f : \sigma(A) \rightarrow \sigma(B)$  constructed from  $\iota(*, *)$ ,  $\pi(*, *)$  are linear combinations of  $\iota(K, B) \circ \pi(A, K)$  for  $K \in \mathcal{P}(A, B)$ . The next definition shows how to compose such morphisms.

**Definition 7.6.** Let  $(I, \preceq)$  be a finite poset. For  $A, B, C \in \mathcal{F}_{(I, \preceq)}$ , define  $\mathcal{T}(A, B, C)$  to be the set of triples  $(J, K, L)$  with  $J \in \mathcal{P}(A, B)$ ,  $K \in \mathcal{P}(B, C)$ ,  $L \in \mathcal{P}(A, C)$ , and  $L \subseteq J \cap K$ .

Let  $J \in \mathcal{P}(A, B)$  and  $K \in \mathcal{P}(B, C)$ . Then  $(J, B) \in \mathcal{G}_{(I, \preceq)}$  and  $(B, K) \in \mathcal{H}_{(I, \preceq)}$ , so  $(J, J \cap K) \in \mathcal{H}_{(I, \preceq)}$  and  $(J \cap K, K) \in \mathcal{G}_{(I, \preceq)}$  by Definition 4.1(d). Also  $(A, J) \in \mathcal{H}_{(I, \preceq)}$  and  $(K, C) \in \mathcal{G}_{(I, \preceq)}$ , so  $(A, J \cap K) \in \mathcal{H}_{(I, \preceq)}$  and  $(J \cap K, C) \in \mathcal{G}_{(I, \preceq)}$  by Definition 4.1(b),(c). Thus  $J \cap K$  is a disjoint union of  $L \in \mathcal{P}(A, C)$  by Proposition 7.5. Hence by definition of  $\mathcal{T}(A, B, C)$  we have

$$J \cap K = \coprod_{\substack{L \in \mathcal{P}(A, C): \\ (J, K, L) \in \mathcal{T}(A, B, C)}} L, \quad \text{for } J \in \mathcal{P}(A, B) \text{ and } K \in \mathcal{P}(B, C).$$

Using Proposition 7.5 and Definition 4.1(B),(C) then gives

$$(\iota(K, C) \circ \pi(B, K)) \circ (\iota(J, B) \circ \pi(A, J)) = \sum_{\substack{L \in \mathcal{P}(A, C): \\ (J, K, L) \in \mathcal{T}(A, B, C)}} \iota(L, C) \circ \pi(A, L). \quad (34)$$

## 7.5 Configurations as functors

We shall now show that an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  is equivalent to an *exact functor*  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  for a certain category  $\mathcal{C}_{(I, \preceq)}$ .

**Definition 7.7.** Let  $(I, \preceq)$  be a finite poset, and  $\mathbb{K}$  be a field. Define a category  $\mathcal{C}_{(I, \preceq)}$  to have objects  $\text{Obj}(\mathcal{C}_{(I, \preceq)}) = \mathcal{F}_{(I, \preceq)}$ , the set of f-sets of  $(I, \preceq)$ , and morphisms  $\text{Mor}(\mathcal{C}_{(I, \preceq)})$  given by  $\text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B) = \{f : \mathcal{P}(A, B) \rightarrow \mathbb{K}\}$ , the set of functions from  $\mathcal{P}(A, B)$  to  $\mathbb{K}$ , for  $A, B \in \mathcal{F}_{(I, \preceq)}$ . Then  $\text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$  is a *vector space over*  $\mathbb{K}$ .

Let  $A, B, C \in \mathcal{F}_{(I, \preceq)}$  and  $f \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$ ,  $g \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(B, C)$ , so  $f, g$  map  $\mathcal{P}(A, B), \mathcal{P}(B, C) \rightarrow \mathbb{K}$ . Define the *composition*  $g \circ f \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, C)$  by

$$(g \circ f)(L) = \sum_{\substack{J \in \mathcal{P}(A, B), K \in \mathcal{P}(B, C): \\ (J, K, L) \in \mathcal{T}(A, B, C)}} f(J)g(K), \quad \text{for all } L \in \mathcal{P}(A, C). \quad (35)$$

Then composition of morphisms is bilinear over  $\mathbb{K}$ . Define the *identity morphism*  $\text{id}_A$  for  $A \in \mathcal{F}_{(I, \preceq)}$  by  $\text{id}_A(J) = 1$  for all  $J \in \mathcal{P}(A, A)$ . For  $(A, B) \in \mathcal{G}_{(I, \preceq)}$ , define  $\alpha(A, B) \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$  by  $\alpha(A, B)(J) = 1$  for all  $J \in \mathcal{P}(A, B)$ . For  $(A, B) \in \mathcal{H}_{(I, \preceq)}$ , define  $\beta(A, B) \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$  by  $\beta(A, B)(J) = 1$  for all  $J \in \mathcal{P}(A, B)$ .

**Theorem 7.8.** *In the situation of Definition 7.7, the composition of morphisms (35) is associative, and  $f \circ \text{id}_A = \text{id}_B \circ f = f$  for  $f \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$ , so  $\mathcal{C}_{(I, \preceq)}$  is a category. Also  $\emptyset \in \text{Obj}(\mathcal{C}_{(I, \preceq)}) = \mathcal{F}_{(I, \preceq)}$  is a zero object in  $\mathcal{C}_{(I, \preceq)}$ .*

*Fix  $f \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$ , let  $D$  be the union of all  $J \in \mathcal{P}(A, B)$  with  $f(J) \neq 0$ , and set  $C = A \setminus D$  and  $E = B \setminus D$ . Then  $(C, A), (D, B) \in \mathcal{G}_{(I, \preceq)}$  and  $(A, D), (D, E) \in \mathcal{H}_{(I, \preceq)}$ . Consider the sequence*

$$C \xrightarrow{\alpha(C, A)} A \xrightarrow{\beta(A, D)} D \xrightarrow{g} B \xrightarrow{\beta(B, E)} E$$

in  $\mathcal{C}_{(I, \preceq)}$ , where  $\mathcal{P}(D, B)$  is the subset of  $J \in \mathcal{P}(A, B)$  with  $f(J) \neq 0$ , and  $g \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(D, B)$  is the restriction of  $f$  to  $\mathcal{P}(D, B) \subseteq \mathcal{P}(A, B)$ .

Then  $g \circ \beta(A, D) = f$ , and  $\alpha(C)$  is a kernel for  $f$  and  $\beta(A, D)$ , and  $\beta(B, E)$  is a cokernel for  $f$  and  $g$ , and  $\beta(A, D)$  is a cokernel for  $\alpha(C, A)$ , and  $g$  is a kernel for  $\beta(B, E)$ . That is,  $\mathcal{C}_{(I, \preceq)}$  satisfies Definition 2.1(iv). Hence kernels and cokernels exist, and exact sequences make sense, in  $\mathcal{C}_{(I, \preceq)}$ .

The following analogues of Definition 4.1(A)–(D) hold in  $\mathcal{C}_{(I, \preceq)}$ .

(A) Let  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and set  $L = K \setminus J$ . Then the following is exact in  $\mathcal{C}_{(I, \preceq)}$ :

$$0 \longrightarrow J \xrightarrow{\alpha(J, K)} K \xrightarrow{\beta(K, L)} L \longrightarrow 0.$$

(B) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{G}_{(I, \preceq)}$  then  $\alpha(J, L) = \alpha(K, L) \circ \alpha(J, K)$ .

(C) If  $(J, K) \in \mathcal{H}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then  $\beta(J, L) = \beta(K, L) \circ \beta(J, K)$ .

(D) If  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $(K, L) \in \mathcal{H}_{(I, \preceq)}$  then

$$\beta(K, L) \circ \alpha(J, K) = \alpha(J \cap L, L) \circ \beta(J, J \cap L).$$

The theorem shows that  $\mathcal{C}_{(I, \preceq)}$  satisfies parts (i), (ii) and (iv) of Definition 2.1, but it does *not* satisfy part (iii) if  $I \neq \emptyset$ , as  $\mathcal{C}_{(I, \preceq)}$  is not closed under direct sums. Therefore  $\mathcal{C}_{(I, \preceq)}$  is *not* an abelian category. One can define a natural abelian category containing  $\mathcal{C}_{(I, \preceq)}$  by a kind of tensor product of  $\mathcal{C}_{(I, \preceq)}$  with the category of finite-dimensional vector spaces over  $\mathbb{K}$ , but we will not do this.

The assumption in Definition 7.7 that  $\mathbb{K}$  is a *field* is needed to prove the statements on kernels and cokernels in the theorem. In particular, they do not hold for  $\mathbb{K} = \mathbb{Z}$ . Now we define the functor  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  associated to  $(\sigma, \iota, \pi)$ .

**Definition 7.9.** Let  $(I, \preceq)$ ,  $\mathbb{K}$  and  $\mathcal{C}_{(I, \preceq)}$  be as in Definition 7.7. Suppose that  $\mathcal{A}$  is a  $\mathbb{K}$ -linear abelian category, and  $(\sigma, \iota, \pi)$  an  $(I, \preceq)$ -configuration in  $\mathcal{A}$ . Define  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  by  $F(A) = \sigma(A)$  for  $A \in \text{Obj}(\mathcal{C}_{(I, \preceq)}) = \mathcal{F}_{(I, \preceq)}$ , and for  $f \in \text{Hom}_{\mathcal{C}_{(I, \preceq)}}(A, B)$  define  $F(f) \in \text{Hom}_{\mathcal{A}}(\sigma(A), \sigma(B))$  by

$$F(f) = \sum_{J \in \mathcal{P}(A, B)} f(J) \iota(J, B) \circ \pi(A, J). \quad (36)$$

**Theorem 7.10.** In Definition 7.9,  $F$  is a  $\mathbb{K}$ -linear functor which takes exact sequences to exact sequences, and satisfies  $F(\alpha(A, B)) = \iota(A, B)$  for  $(A, B) \in \mathcal{G}_{(I, \preceq)}$  and  $F(\beta(A, B)) = \pi(A, B)$  for  $(A, B) \in \mathcal{H}_{(I, \preceq)}$ .

Conversely, if  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  is a  $\mathbb{K}$ -linear exact functor then setting  $\sigma(A) = F(A)$  for  $A \in \mathcal{F}_{(I, \preceq)}$  and  $\iota(A, B) = F(\alpha(A, B))$  for  $(A, B) \in \mathcal{G}_{(I, \preceq)}$  and  $\pi(A, B) = F(\beta(A, B))$  for  $(A, B) \in \mathcal{H}_{(I, \preceq)}$  defines an  $(I, \preceq)$ -configuration  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$ .

The theorem provides the motivation for Definitions 7.7 and 7.9. In particular, the composition rule (35) for morphisms in  $\mathcal{C}_{(I, \preceq)}$  follows from (34) and (36).

To study  $(I, \preceq)$ -configurations in an abelian category  $\mathcal{A}$  which is not  $\mathbb{K}$ -linear over some field  $\mathbb{K}$ , we can define  $\mathcal{C}_{(I, \preceq)}$  using  $\mathbb{K} = \mathbb{Z}$  in Definition 7.7. Then  $\mathcal{C}_{(I, \preceq)}$  is still a category, and  $F : \mathcal{C}_{(I, \preceq)} \rightarrow \mathcal{A}$  is well-defined in Definition 7.9. However, as kernels and cokernels do not exist for all morphisms in  $\mathcal{C}_{(I, \preceq)}$ , we need to be more careful about what we mean by  $F$  being an *exact* functor.

The constructions of §5 can be explained in this functorial notation. If  $(\sigma, \iota, \pi)$  is an  $(I, \preceq)$ -configuration and  $(\sigma', \iota', \pi')$  its  $(J, \preceq)$ -subconfiguration for  $J \in \mathcal{F}_{(I, \preceq)}$ , and  $F, F'$  are the associated functors, then  $F' = F \circ \mathcal{S}(I, \preceq, J)$  for a natural functor  $\mathcal{S}(I, \preceq, J) : \mathcal{C}_{(J, \preceq)} \rightarrow \mathcal{C}_{(I, \preceq)}$ . Similarly, if  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi})$  is the quotient  $(K, \trianglelefteq)$ -configuration of  $(\sigma, \iota, \pi)$  from  $\phi$  with functor  $\tilde{F}$  then  $\tilde{F} = F \circ \mathcal{Q}(I, \preceq, K, \trianglelefteq, \phi)$  for a natural functor  $\mathcal{Q}(I, \preceq, K, \trianglelefteq, \phi) : \mathcal{C}_{(K, \trianglelefteq)} \rightarrow \mathcal{C}_{(I, \preceq)}$ .

Theorem 5.5 has a more complicated explanation, as follows. The functors

$$\begin{aligned} \mathcal{Q}(I, \preceq, K, \trianglelefteq, \phi) : \mathcal{C}_{(K, \trianglelefteq)} &\rightarrow \mathcal{C}_{(I, \preceq)}, & \mathcal{S}(I, \preceq, J) : \mathcal{C}_{(J, \preceq)} &\rightarrow \mathcal{C}_{(I, \preceq)}, \\ \mathcal{Q}(J, \preceq, L, \trianglelefteq, \psi) : \mathcal{C}_{(L, \trianglelefteq)} &\rightarrow \mathcal{C}_{(J, \preceq)}, & \mathcal{S}(K, \trianglelefteq, L) : \mathcal{C}_{(L, \trianglelefteq)} &\rightarrow \mathcal{C}_{(K, \trianglelefteq)}, \end{aligned}$$

satisfy  $\mathcal{Q}(I, \preceq, K, \trianglelefteq, \phi) \circ \mathcal{S}(K, \trianglelefteq, L) = \mathcal{S}(I, \preceq, J) \circ \mathcal{Q}(J, \preceq, L, \trianglelefteq, \psi)$ , and so  $\mathcal{Q}(I, \preceq, K, \trianglelefteq, \phi), \mathcal{S}(I, \preceq, J)$  induce a functor from the *fibre product category*

$$\mathcal{C}_{(K, \trianglelefteq)} \times_{\mathcal{S}(K, \trianglelefteq, L), \mathcal{C}_{(L, \trianglelefteq)}, \mathcal{Q}(J, \preceq, L, \trianglelefteq, \psi)} \mathcal{C}_{(J, \preceq)} \rightarrow \mathcal{C}_{(I, \preceq)}.$$

This is an *equivalence of categories*, and so a functor from the left hand side to  $\mathcal{A}$  induced by  $(\tilde{\sigma}, \tilde{\iota}, \tilde{\pi}), (\sigma', \iota', \pi')$ , extends to a functor from the right hand side, uniquely up to canonical isomorphism.

## 7.6 The exact category of exact sequences in $\mathcal{A}$

Let  $\mathcal{A}$  be an *abelian category*, which may be  $\mathbb{K}$ -linear over a field  $\mathbb{K}$ . In §7.7 we will show that the category  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  of  $(I, \preceq)$ -configurations in  $\mathcal{A}$  is an *exact category*. As a warm-up exercise, we first show that the category  $\mathbf{Exact}(\mathcal{A})$  of *short exact sequences* in  $\mathcal{A}$  is an exact category. Define the category  $\mathbf{Comp}(\mathcal{A})$  of *complexes*  $X_1 \xrightarrow{\phi_1} X_2 \xrightarrow{\phi_2} X_3$  in  $\mathcal{A}$  as follows.

*Objects* in  $\mathbf{Comp}(\mathcal{A})$  are  $\mathbf{X} = (X_1, X_2, X_3, \phi_1, \phi_2)$  with  $X_1, X_2, X_3 \in \text{Obj}(\mathcal{A})$  and  $\phi_1 : X_1 \rightarrow X_2, \phi_2 : X_2 \rightarrow X_3$  in  $\text{Mor}(\mathcal{A})$  with  $\phi_2 \circ \phi_1 = 0$ . *Morphisms* in  $\mathbf{Comp}(\mathcal{A})$  are  $\mathbf{f} = (f_1, f_2, f_3) : (X_1, X_2, X_3, \phi_1, \phi_2) \rightarrow (Y_1, Y_2, Y_3, \psi_1, \psi_2)$ , where  $f_a : X_a \rightarrow Y_a$  lie in  $\text{Mor}(\mathcal{A})$  for  $a = 1, 2, 3$  with  $\psi_1 \circ f_1 = f_2 \circ \phi_1, \psi_2 \circ f_2 = f_3 \circ \phi_2$ . Morphisms compose by  $(f_1, f_2, f_3) \circ (g_1, g_2, g_3) = (f_1 \circ g_1, f_2 \circ g_2, f_3 \circ g_3)$ .

**Proposition 7.11.** *This  $\mathbf{Comp}(\mathcal{A})$  is an abelian category. If  $\mathcal{A}$  is  $\mathbb{K}$ -linear then  $\mathbf{Comp}(\mathcal{A})$  is  $\mathbb{K}$ -linear.*

*Proof.* We shall verify Definition 2.1(i)–(iv). Let  $\mathbf{X} = (X_1, X_2, X_3, \phi_1, \phi_2)$  and  $\mathbf{Y} = (Y_1, Y_2, Y_3, \psi_1, \psi_2)$  lie in  $\text{Obj}(\mathbf{Comp}(\mathcal{A}))$ . For (i), the group structures on  $\text{Hom}(X_a, Y_a)$  induce one on  $\text{Hom}(\mathbf{X}, \mathbf{Y})$ , so that addition is defined by  $(f_1, f_2, f_3) + (\beta_1, \beta_2, \beta_3) = (f_1 + \beta_1, f_2 + \beta_2, f_3 + \beta_3)$ , and so on. As the conditions  $\psi_1 \circ f_1 = f_2 \circ \phi_1, \psi_2 \circ f_2 = f_3 \circ \phi_2$  are linear in  $(f_1, f_2, f_3)$ ,  $\text{Hom}(\mathbf{X}, \mathbf{Y})$  is closed under addition, etc. Composition of morphisms is also clearly biadditive, proving (i).

If  $\mathcal{A}$  is  $\mathbb{K}$ -linear then  $\text{Hom}(\mathbf{X}, \mathbf{Y})$  is a  $\mathbb{K}$ -vector space and composition is bilinear, which will prove  $\mathbf{Comp}(\mathcal{A})$  is  $\mathbb{K}$ -linear once we have verified (ii)–(iv). For (ii), the zero object is  $(0, 0, 0, 0, 0)$ . For (iii), direct sums are given by

$$\mathbf{X} \oplus \mathbf{Y} = (X_1 \oplus Y_1, X_2 \oplus Y_2, X_3 \oplus Y_3, \iota_{X_2} \circ \phi_1 \circ \pi_{X_1} + \iota_{Y_2} \circ \psi_1 \circ \pi_{Y_1}, \\ \iota_{X_3} \circ \phi_2 \circ \pi_{X_2} + \iota_{Y_3} \circ \psi_2 \circ \pi_{Y_2}),$$

with  $\iota_{\mathbf{X}} : \mathbf{X} \rightarrow \mathbf{X} \oplus \mathbf{Y}$  equal to  $(\iota_{X_1}, \iota_{X_2}, \iota_{X_3})$ , and so on.

For (iv), let  $\mathbf{f} : \mathbf{X} \rightarrow \mathbf{Y}$  be as above. Applying Definition 2.1(iv) to  $f_a : X_a \rightarrow Y_a$  in the abelian category  $\mathcal{A}$  for  $a = 1, 2, 3$  gives a sequence  $K_a \xrightarrow{k_a} X_a \xrightarrow{i_a} I_a \xrightarrow{j_a} Y_a \xrightarrow{c_a} C_a$  in  $\mathcal{A}$  such that  $j_a \circ i_a = f_a$ , and  $K_a$  is the kernel of  $f_a$ , and  $C_a$  the cokernel of  $f_a$ , and  $I_a$  is both the cokernel of  $k_a$  and the kernel of  $c_a$ . We shall show these fit into a commutative diagram

$$\begin{array}{ccccc} K_1 & \overset{\kappa_1}{\dashrightarrow} & K_2 & \overset{\kappa_2}{\dashrightarrow} & K_3 \\ \downarrow k_1 & & \downarrow k_2 & & \downarrow k_3 \\ X_1 & \xrightarrow{\phi_1} & X_2 & \xrightarrow{\phi_2} & X_3 \\ \downarrow f_1 & \searrow i_1 & \downarrow f_2 & \searrow i_2 & \downarrow f_3 \\ Y_1 & \xrightarrow{\psi_1} & Y_2 & \xrightarrow{\psi_2} & Y_3 \\ \downarrow c_1 & & \downarrow c_2 & & \downarrow c_3 \\ C_1 & \overset{\gamma_1}{\dashrightarrow} & C_2 & \overset{\gamma_2}{\dashrightarrow} & C_3 \end{array} \quad (37)$$

where the morphisms ‘ $\dashrightarrow$ ’ have yet to be constructed.

Since  $\psi_1 \circ f_1 = f_2 \circ \phi_1$  we have  $f_2 \circ (\phi_1 \circ k_1) = \psi_1 \circ f_1 \circ k_1 = 0$ , as  $k_1 = \text{Ker } f_1$ . But  $k_2 = \text{Ker } f_2$ , so there is a unique  $\kappa_1 : K_1 \rightarrow K_2$  with  $k_2 \circ \kappa_1 = \phi_1 \circ k_1$ , making the top left square commute. We construct  $\kappa_2$  in the same way. Then

$$k_3 \circ \kappa_2 \circ \kappa_1 = \phi_2 \circ k_2 \circ \kappa_1 = \phi_2 \circ \phi_1 \circ \kappa_1 = 0,$$

as  $\phi_1 \circ \phi_1 = 0$  by definition. But  $k_2$  is *injective* as it is a kernel, so  $\kappa_2 \circ \kappa_1 = 0$ . Hence  $\mathbf{K} = (K_1, K_2, K_3, \kappa_1, \kappa_2)$  lies in  $\text{Obj}(\mathbf{Comp}(\mathcal{A}))$ . By a dual proof, using cokernels instead of kernels, we construct  $\gamma_1, \gamma_2$  so the bottom squares commute and  $\gamma_2 \circ \gamma_1 = 0$ , so that  $\mathbf{C} = (C_1, C_2, C_3, \gamma_1, \gamma_2)$  lies in  $\text{Obj}(\mathbf{Comp}(\mathcal{A}))$ .

Now  $i_2 \circ \phi_1 \circ k_1 = i_2 \circ k_2 \circ \kappa_1 = 0$ , as  $i_2 = \text{Coker } k_2$ . But  $i_1 = \text{Coker } k_1$ , so there is a unique  $\iota_1 : I_1 \rightarrow I_2$  with  $\iota_1 \circ i_1 = i_2 \circ \phi_1$ , making one of the diamonds in (37) commute. Then

$$j_2 \circ \iota_1 \circ i_1 = j_2 \circ i_2 \circ \phi_1 = f_2 \circ \phi_1 = \psi_1 \circ f_1 = \psi_1 \circ j_1 \circ i_1.$$

But  $i_1$  is *surjective* as it is a cokernel, so  $j_2 \circ \iota_1 = \psi_1 \circ j_1$ , and another diamond commutes. In the same way we construct  $\iota_2 : I_2 \rightarrow I_3$  with  $\iota_2 \circ i_2 = i_3 \circ \phi_2$  and  $j_3 \circ \iota_2 = \psi_2 \circ j_2$ , so the whole of (37) commutes. And

$$\iota_2 \circ \iota_1 \circ i_1 = \iota_2 \circ i_2 \circ \phi_1 = i_3 \circ \phi_2 \circ \phi_1 = 0,$$

so  $\iota_2 \circ \iota_1 = 0$  as  $i_1$  is surjective, and  $\mathbf{I} = (I_1, I_2, I_3, \iota_1, \iota_2)$  lies in  $\text{Obj}(\mathbf{Comp}(\mathcal{A}))$ . It is now easy to show that the sequence  $\mathbf{K} \xrightarrow{k} \mathbf{X} \xrightarrow{i} \mathbf{I} \xrightarrow{j} \mathbf{Y} \xrightarrow{c} \mathbf{C}$  in  $\mathbf{Comp}(\mathcal{A})$  fulfils Definition 2.1(iv).  $\square$

Now define the category  $\mathbf{Exact}(\mathcal{A})$  of *exact sequences* in  $\mathcal{A}$  to be the full subcategory of  $\mathbf{Comp}(\mathcal{A})$  with objects  $\mathbf{X} = (X_1, X_2, X_3, \phi_1, \phi_2)$  such that  $0 \rightarrow X_1 \xrightarrow{\phi_1} X_2 \xrightarrow{\phi_2} X_3 \rightarrow 0$  is a *short exact sequence* in  $\mathcal{A}$ . Define short exact sequences in  $\mathbf{Exact}(\mathcal{A})$  to be short exact sequences in  $\mathbf{Comp}(\mathcal{A})$  with objects in  $\mathbf{Exact}(\mathcal{A})$ . Then we have:

**Proposition 7.12.** *This  $\mathbf{Exact}(\mathcal{A})$  is an exact category. If  $\mathcal{A}$  is  $\mathbb{K}$ -linear then  $\mathbf{Exact}(\mathcal{A})$  is  $\mathbb{K}$ -linear.*

*Proof.* Clearly  $\mathbf{Exact}(\mathcal{A})$  is a *full subcategory* of  $\mathbf{Comp}(\mathcal{A})$  which contains zero and is closed under direct sums, so it is an *additive subcategory*. By Definition 2.7, it remains to verify it is *closed under extensions*. Suppose  $0 \rightarrow \mathbf{X} \xrightarrow{f} \mathbf{Y} \xrightarrow{g} \mathbf{Z} \rightarrow 0$  is a short exact sequence in  $\mathbf{Comp}(\mathcal{A})$  with  $\mathbf{X}, \mathbf{Z} \in \mathbf{Exact}(\mathcal{A})$ . We must show that  $\mathbf{Y} \in \mathbf{Exact}(\mathcal{A})$ .

That is, we have a commutative diagram

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & X_1 & \xrightarrow{\phi_1} & X_2 & \xrightarrow{\phi_2} & X_3 \longrightarrow 0 \\
& & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 \\
0 & \longrightarrow & Y_1 & \xrightarrow{\psi_1} & Y_2 & \xrightarrow{\psi_2} & Y_3 \longrightarrow 0 \\
& & \downarrow g_1 & \swarrow i & \downarrow g_2 & \searrow k & \downarrow g_3 \\
& & Z_1 & \xrightarrow{\chi_1} & Z_2 & \xrightarrow{\chi_2} & Z_3 \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array} \tag{38}$$

in  $\mathcal{A}$ , with  $K$  and arrows ‘ $\dashrightarrow$ ’ yet to be constructed, and with the columns and first and last rows exact. We must show the middle row is exact.

Suppose  $\alpha : A \rightarrow Y_1$  lies in  $\text{Mor}(\mathcal{A})$  with  $\psi_1 \circ \alpha = 0$ . Then  $\chi_1 \circ g_1 \circ \alpha = g_2 \circ \psi_1 \circ \alpha = 0$ , so  $g_1 \circ \alpha = 0$  as  $\chi_1$  is *injective*. But  $f_1 = \text{Ker } g_1$ , so there is a unique  $\beta : A \rightarrow X_1$  with  $\alpha = f_1 \circ \beta$ . Thus  $f_2 \circ \phi_1 \circ \beta = \psi_1 \circ f_1 \circ \beta = \psi_1 \circ \alpha = 0$ . Now  $f_2, \phi_1$  are injective, so  $f_2 \circ \phi_1$  is injective, giving  $\beta = 0$ , so that  $\alpha = f_1 \circ \beta = 0$ . Therefore  $\psi_1 \circ \alpha = 0$  implies  $\alpha = 0$ , and  $\psi_1$  is *injective*. A dual proof shows that  $\psi_2$  is *surjective*.

Let  $k : K \rightarrow Y_2$  be a *kernel* for  $\psi_2$ . As  $\psi_2 \circ \psi_1 = 0$  there exists a unique  $i : Y_1 \rightarrow K$  with  $k \circ i = \psi_1$ . As  $\chi_2 \circ g_2 \circ k = g_2 \circ \psi_2 \circ k = 0$  and  $\chi_1 = \text{ker } \chi_2$  there exists a unique  $j : K \rightarrow Z_1$  with  $g_2 \circ k = \chi_1 \circ j$ . Also  $\chi_1 \circ j \circ i = g_2 \circ k \circ i = g_2 \circ \psi_1 = \chi_1 \circ g_1$ , so  $j \circ i = g_1$  as  $\chi_1$  is injective. This shows (38) commutes. We shall show  $i$  is an *isomorphism*.

As  $j \circ i = \psi_1$  and  $\psi_1$  is injective, we see that  $i$  is *injective*. Since  $j \circ i = g_1$  and  $g_1$  is surjective,  $j$  is *surjective*. We claim  $i \circ f_1$  is a *kernel* for  $j$ . Suppose

$\alpha : A \rightarrow K$  with  $j \circ \alpha = 0$ . Then  $g_2 \circ k \circ \alpha = \chi_1 \circ j \circ \alpha = 0$ , so as  $f_2 = \text{Ker } g_2$  there is a unique  $\beta : A \rightarrow X_2$  with  $f_2 \circ \beta = k \circ \alpha$ . Then

$$f_3 \circ \phi_2 \circ \beta = \psi_2 \circ f_2 \circ \beta = \psi_2 \circ k \circ \alpha = 0,$$

as  $k = \text{Ker } \psi_2$ . But  $f_3$  is injective, so  $\phi_2 \circ \beta = 0$ . Hence there exists  $\gamma : A \rightarrow X_1$  with  $\beta = \phi_1 \circ \gamma$ , as  $\phi_1 = \text{Ker } \phi_2$ . Therefore

$$k \circ \alpha = f_2 \circ \beta = f_2 \circ \phi_1 \circ \gamma = \psi_1 \circ f_1 \circ \gamma = k \circ i \circ f_1 \circ \gamma.$$

So  $\alpha = i \circ f_1 \circ \gamma$  since  $k$  is injective, as it is a kernel. Thus every  $\alpha : A \rightarrow K$  with  $j \circ \alpha = 0$  factors via  $i \circ f_1$ , and  $i \circ f_1 = \text{Ker } j$ . As  $j$  is surjective, we have  $j = \text{Coker}(i \circ f_1)$ .

Suppose  $\delta : K \rightarrow D$  with  $\delta \circ i = 0$ . Then  $\delta \circ i \circ f_1 = 0$ , so there exists  $\epsilon : Z_1 \rightarrow D$  with  $\delta = \epsilon \circ j$ , as  $j = \text{Coker}(i \circ f_1)$ . Thus  $\epsilon \circ g_1 = \epsilon \circ j \circ i = \delta \circ i = 0$ , which implies  $\epsilon = 0$  as  $g_1$  is surjective, and so  $\delta = 0$ . Hence  $\delta \circ i = 0$  implies  $\delta = 0$ , and  $i$  is *surjective*, so it is an *isomorphism*. Since  $k$  is a kernel for  $\psi_2$ , this implies  $\psi_1$  is a kernel for  $\psi_2$ . As  $\psi_2$  is surjective, the middle row of (38) is *exact*, and the proof is complete.  $\square$

Note that  $\mathbf{Exact}(\mathcal{A})$  is *never* an abelian category if  $\mathcal{A}$  is nonzero. For if  $X \in \mathcal{A}$  with  $X \not\cong 0$  then  $(0, \text{id}_X, 0) : (0, X, X, 0, \text{id}_X) \rightarrow (X, X, 0, \text{id}_X, 0)$  is a morphism in  $\mathbf{Exact}(\mathcal{A})$ , whose kernel  $(0, 0, X, 0, 0)$  and cokernel  $(X, 0, 0, 0, 0)$  lie in  $\mathbf{Comp}(\mathcal{A})$  but not in  $\mathbf{Exact}(\mathcal{A})$ , so  $\mathbf{Exact}(\mathcal{A})$  is not closed under kernels or cokernels.

## 7.7 The exact category of $(I, \preceq)$ -configurations in $\mathcal{A}$

Let  $\mathcal{A}$  be an abelian category, and  $(I, \preceq)$  a finite poset. We now generalize the proofs of §7.7 to show that the  $(I, \preceq)$ -configurations in  $\mathcal{A}$  form an *exact category*  $\mathbf{Conf}(I, \preceq, \mathcal{A})$ . First we need an abelian category to embed  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  in, analogous to the category  $\mathbf{Comp}(\mathcal{A})$  of *complexes* in  $\mathcal{A}$ .

Define an  $(I, \preceq)$ -*preconfiguration*  $(\sigma, \iota, \pi)$  in  $\mathcal{A}$  as in Definition 4.1, except that in (ii), (iii) we do not require  $\iota(J, K)$  to be *injective* or  $\pi(J, K)$  *surjective*, and instead of supposing (6) *exact* we require only that  $\pi(K, L) \circ \iota(J, K) = 0$ . Define *morphisms* of  $(I, \preceq)$ -preconfigurations as in Definition 4.1. Write  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  for the category of  $(I, \preceq)$ -preconfigurations. Here is the analogue of Proposition 7.11.

**Proposition 7.13.** *This  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  is an abelian category. If  $\mathcal{A}$  is  $\mathbb{K}$ -linear then  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  is  $\mathbb{K}$ -linear.*

*Proof.* We verify Definition 2.1(i)–(iv). Parts (i)–(iii) and  $\mathbb{K}$ -linearity are an easy generalization of Proposition 7.11. For (iv), suppose  $\alpha : (\sigma_2, \iota_2, \pi_2) \rightarrow (\sigma_4, \iota_4, \pi_4)$  lies in  $\text{Mor}(\mathbf{PConf}(I, \preceq, \mathcal{A}))$ . For each  $J \in \mathcal{F}_{(I, \preceq)}$ , apply Definition 2.1(iv) to  $\alpha(J) : \sigma_2(J) \rightarrow \sigma_4(J)$  in  $\mathcal{A}$ . This gives a sequence

$$\sigma_1(J) \xrightarrow{\beta(J)} \sigma_2(J) \xrightarrow{\gamma(J)} \sigma_3(J) \xrightarrow{\delta(J)} \sigma_4(J) \xrightarrow{\epsilon(J)} \sigma_5(J)$$

in  $\mathcal{A}$  with  $\delta(J) \circ \gamma(J) = \alpha(J)$ , such that  $\sigma_1(J)$  is the kernel and  $\sigma_5(J)$  the cokernel of  $\alpha(J)$ , and  $\sigma_3(J)$  is both the cokernel of  $\beta(J)$  and the kernel of  $\epsilon(J)$ .

For  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and  $L = K \setminus J$ , the proof of Proposition 7.11 yields unique morphisms  $\iota_a(J, K), \pi_a(K, L)$  for  $a = 1, 3, 5$  with  $\pi_a(K, L) \circ \iota_a(J, K) = 0$  fitting into a commutative diagram

$$\begin{array}{ccccc}
\sigma_1(J) & \overset{\iota_1(J, K)}{\dashrightarrow} & \sigma_1(K) & \overset{\pi_1(K, L)}{\dashrightarrow} & \sigma_1(L) \\
\downarrow \beta(J) & & \downarrow \beta(K) & & \downarrow \beta(L) \\
\sigma_2(J) & \xrightarrow{\iota_2(J, K)} & \sigma_2(K) & \xrightarrow{\iota_2(K, L)} & \sigma_2(L) \\
\downarrow \alpha(J) & \searrow \gamma(J) & \downarrow \alpha(K) & \searrow \gamma(K) & \downarrow \alpha(L) \\
\sigma_3(J) & \overset{\iota_3(J, K)}{\dashrightarrow} & \sigma_3(K) & \overset{\pi_3(K, L)}{\dashrightarrow} & \sigma_3(L) \\
\downarrow \delta(J) & \swarrow \gamma(J) & \downarrow \delta(K) & \swarrow \gamma(K) & \downarrow \delta(L) \\
\sigma_4(J) & \xrightarrow{\iota_4(J, K)} & \sigma_4(K) & \xrightarrow{\iota_4(K, L)} & \sigma_4(L) \\
\downarrow \epsilon(J) & & \downarrow \epsilon(K) & & \downarrow \epsilon(L) \\
\sigma_5(J) & \overset{\iota_5(J, K)}{\dashrightarrow} & \sigma_5(K) & \overset{\pi_5(K, L)}{\dashrightarrow} & \sigma_5(L)
\end{array} \quad (39)$$

We must show  $(\sigma_a, \iota_a, \pi_a)$  is an  $(I, \preceq)$ -preconfiguration for  $i = 1, 3, 5$ . That is, we must prove that  $\iota_a(J, J) = \pi_a(J, J) = \text{id}_{\sigma_a(J)}$  for  $J \in \mathcal{F}_{(I, \preceq)}$ , and Definition 4.1(B)–(D) hold.

Putting  $J = K$  and  $L = \emptyset$  in (39) and using  $\iota_2(J) = \text{id}_{\sigma_2(J)}$  and  $\iota_4(J) = \text{id}_{\sigma_4(J)}$ , uniqueness in (39) implies that  $\iota_a(J, J) = \text{id}_{\sigma_a(J)}$  for  $a = 1, 3, 5$ . A similar proof gives  $\pi_a(J, J) = \text{id}_{\sigma_a(J)}$ . For (B), if  $(J, K), (K, L) \in \mathcal{G}_{(I, \preceq)}$  then

$$\begin{aligned}
\beta(L) \circ \iota_1(K, L) \circ \iota_1(J, K) &= \iota_2(K, L) \circ \beta(K) \circ \iota_1(J, K) = \\
&= \iota_2(K, L) \circ \iota_2(J, K) \circ \beta(J) = \iota_2(J, L) \circ \beta(J) = \beta(L) \circ \iota_1(J, L),
\end{aligned}$$

as  $(\sigma_2, \iota_2, \pi_2) \in \text{Obj}(\mathbf{PConf}(I, \preceq, \mathcal{A}))$  and the top left square in (39) commutes. But  $\beta(L)$  is *injective*, so  $\iota_1(K, L) \circ \iota_1(J, K) = \iota_1(J, L)$ , giving (B) for  $(\sigma_1, \iota_1, \pi_1)$ .

Similar proofs using injectivity of  $\delta(L)$  and surjectivity of  $\gamma(J), \epsilon(J)$  prove (B)–(D) for  $(\sigma_a, \iota_a, \pi_a)$  for  $a = 1, 3, 5$ . Therefore

$$(\sigma_1, \iota_1, \pi_1) \xrightarrow{\beta} (\sigma_2, \iota_2, \pi_2) \xrightarrow{\gamma} (\sigma_3, \iota_3, \pi_3) \xrightarrow{\delta} (\sigma_4, \iota_4, \pi_4) \xrightarrow{\epsilon} (\sigma_5, \iota_5, \pi_5)$$

is a sequence in  $\mathbf{PConf}(I, \preceq, \mathcal{A})$ , which is easily seen to fulfil Definition 2.1(iv). So  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  is an *abelian category*.  $\square$

Write  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  for the category of  $(I, \preceq)$ -configurations in  $\mathcal{A}$ , a full subcategory of  $\mathbf{PConf}(I, \preceq, \mathcal{A})$ . Following Proposition 7.12 we prove:

**Theorem 7.14.** *The category  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  of  $(I, \preceq)$ -configurations in  $\mathcal{A}$  is an exact category. If  $\mathcal{A}$  is  $\mathbb{K}$ -linear then  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  is  $\mathbb{K}$ -linear.*

*Proof.* Clearly  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  is a *full subcategory* of  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  which contains zero and is closed under direct sums, so it is an *additive subcategory*. We show it is *closed under extensions*. Suppose

$$0 \rightarrow (\sigma_1, \iota_1, \pi_1) \xrightarrow{\alpha} (\sigma_2, \iota_2, \pi_2) \xrightarrow{\beta} (\sigma_3, \iota_3, \pi_3) \rightarrow 0$$

is a short exact sequence in  $\mathbf{PConf}(I, \preceq, \mathcal{A})$  with  $(\sigma_1, \iota_1, \pi_1)$ ,  $(\sigma_3, \iota_3, \pi_3)$  in  $\text{Obj}(\mathbf{Conf}(I, \preceq, \mathcal{A}))$ . We must show that  $(\sigma_2, \iota_2, \pi_2) \in \mathbf{Conf}(I, \preceq, \mathcal{A})$ .

Let  $(J, K) \in \mathcal{G}_{(I, \preceq)}$  and set  $L = K \setminus J$ . Then we have a commutative diagram

$$\begin{array}{ccccccccc}
& & 0 & & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \sigma_1(J) & \xrightarrow{\iota_1(J,K)} & \sigma_1(K) & \xrightarrow{\pi_1(K,L)} & \sigma_1(L) & \longrightarrow & 0 \\
& & \downarrow \alpha(J) & & \downarrow \alpha(K) & & \downarrow \alpha(L) & & \\
0 & \longrightarrow & \sigma_2(J) & \xrightarrow{\iota_2(J,K)} & \sigma_2(K) & \xrightarrow{\pi_2(K,L)} & \sigma_2(L) & \longrightarrow & 0 \\
& & \downarrow \beta(J) & & \downarrow \beta(K) & & \downarrow \beta(L) & & \\
0 & \longrightarrow & \sigma_3(J) & \xrightarrow{\iota_3(J,K)} & \sigma_3(K) & \xrightarrow{\pi_3(K,L)} & \sigma_3(L) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
& & 0 & & 0 & & 0 & & 
\end{array}$$

in  $\mathcal{A}$ , with the columns and first and last rows exact. The proof of Proposition 7.12 now shows that the middle row is exact. As this holds for all  $(J, K) \in \mathcal{G}_{(I, \preceq)}$ ,  $(\sigma_2, \iota_2, \pi_2)$  is an  $(I, \preceq)$ -configuration, and  $\mathbf{Conf}(I, \preceq, \mathcal{A})$  is closed under extensions. This completes the proof.  $\square$

Using the ideas of §7.2, it is easy to generalize this to show that if  $(\mathcal{B}, \mathcal{E})$  is an exact category then the category  $\mathbf{Conf}(I, \preceq, \mathcal{B})$  of  $(I, \preceq)$ -configurations in  $\mathcal{B}$  is also an *exact category*.

Let  $(J, \lesssim)$  be another finite poset. Then we can study  $(J, \lesssim)$ -configurations in the exact category  $\mathbf{Conf}(I, \preceq, \mathcal{A})$ , that is,  $(J, \lesssim)$ -configurations of  $(I, \preceq)$ -configurations in  $\mathcal{A}$ . The category  $\mathbf{Conf}(J, \lesssim, \mathbf{Conf}(I, \preceq, \mathcal{A}))$  of such ‘configurations of configurations’ is an *exact category* from above, so we can iterate this process indefinitely.

Define  $K = I \times J$ , and a partial order  $\trianglelefteq$  on  $K$  by  $(i, j) \trianglelefteq (i', j')$  if  $i \preceq i'$  and  $j \lesssim j'$ . Then it can be shown that  $(J, \lesssim)$ -configurations of  $(I, \preceq)$ -configurations in  $\mathcal{A}$  are essentially the same thing as  $(K, \trianglelefteq)$ -configurations in  $\mathcal{A}$ . To be more precise, there is a natural *forgetful functor*

$$\mathbf{Conf}(K, \trianglelefteq, \mathcal{A}) \rightarrow \mathbf{Conf}(J, \lesssim, \mathbf{Conf}(I, \preceq, \mathcal{A}))$$

which is an *equivalence of categories*, since each  $(K, \trianglelefteq)$ -configuration induces a unique  $(J, \lesssim)$ -configuration of  $(I, \preceq)$ -configurations, and conversely each  $(J, \lesssim)$ -configuration of  $(I, \preceq)$ -configurations comes from a  $(K, \trianglelefteq)$ -configuration which is unique up to canonical isomorphism.

One moral of this is that  $(I, \preceq)$ -configurations are a useful, universal idea, since *operations on configurations* such as those of §5, or these ‘configurations of configurations’, *tend to produce  $(K, \trianglelefteq)$ -configurations for another finite poset  $(K, \trianglelefteq)$* , rather than some more general object. We will see more examples of this in [4, 5], where we show that the transformations of moduli spaces of stable  $(I, \preceq)$ -configurations under change of stability condition can be fully described in terms of moduli spaces of  $(K, \trianglelefteq)$ -configurations, making configurations a good tool for understanding stability.

## References

- [1] K. Baclawski, *Whitney Numbers of Geometric Lattices*, Advances in Math. 16 (1975), 125–138.
- [2] S.I. Gelfand and Y.I. Manin, *Methods of Homological Algebra*, second edition, Springer Monographs in Mathematics, Springer, Berlin, 2003.
- [3] D.D. Joyce, *Configurations in abelian categories. II. Moduli stacks*, math.AG/0312192, version 3, 2004.
- [4] D.D. Joyce, *Configurations in abelian categories. III. Stability conditions and invariants*, math.AG/0410267, 2004.
- [5] D.D. Joyce, *Configurations in abelian categories. IV. Changing stability conditions*, math.AG/0410268, 2004.
- [6] D.D. Joyce, *Configurations in abelian categories. V. Infinite-dimensional algebras*, in preparation, 2004.
- [7] D.D. Joyce, *Constructible functions on schemes and stacks*, math.AG/0403305, 2004.
- [8] D.D. Joyce, in preparation, 2004.
- [9] N. Popescu, *Abelian Categories with Applications to Rings and Modules*, L.M.S. Monographs 3, Academic Press, London, 1973.
- [10] D. Quillen, *Higher algebraic K-theory. I*, pages 85–147 in H. Bass, editor, *Algebraic K-theory. I*, Springer Lecture Notes 341, Springer-Verlag, Berlin, 1973.
- [11] C.S. Seshadri, *Space of unitary vector bundles on a compact Riemann surface*, Ann. Math. 85 (1967), 303–336.