# Free $A_{\infty}$ -categories

Volodymyr Lyubashenko<sup>\*</sup> and Oleksandr Manzyuk<sup>†</sup>

February 8, 2020

#### Abstract

For a differential graded k-quiver  $\Omega$  we define the free  $A_{\infty}$ -category  $\mathcal{F}\Omega$  generated by  $\Omega$ . The main result is that the restriction  $A_{\infty}$ -functor  $A_{\infty}(\mathcal{F}\Omega, \mathcal{A}) \to A_1(\Omega, \mathcal{A})$  is an equivalence, where objects of the last  $A_{\infty}$ -category are morphisms of differential graded k-quivers  $\Omega \to \mathcal{A}$ .

 $A_{\infty}$ -categories defined by Fukaya [Fuk93] and Kontsevich [Kon95] are generalizations of differential graded categories for which the binary composition is associative only up to a homotopy. They also generalize  $A_{\infty}$ -algebras introduced by Stasheff [Sta63, II].  $A_{\infty}$ -functors are the corresponding generalizations of usual functors, see e.g. [Fuk93, Kel01]. Homomorphisms of  $A_{\infty}$ -algebras (e.g. [Kad82]) are particular cases of  $A_{\infty}$ -functors.  $A_{\infty}$ -transformations are certain coderivations. Examples of such structures are encountered in studies of mirror symmetry (e.g. [Kon95, Fuk02]) and in homological algebra.

For an  $A_{\infty}$ -category there is a notion of units up to a homotopy (homotopy identity morphisms) [Lyu03]. Given two  $A_{\infty}$ -categories  $\mathcal{A}$  and  $\mathcal{B}$ , one can construct a third  $A_{\infty}$ -category  $A_{\infty}(\mathcal{A}, \mathcal{B})$ , whose objects are  $A_{\infty}$ -functors  $f : \mathcal{A} \to \mathcal{B}$ , and morphisms are  $A_{\infty}$ -transformations between such functors (Fukaya [Fuk02], Kontsevich and Soibelman [KS02, KS], Lefèvre-Hasegawa [LH03], as well as [Lyu03]). This allows to define a 2-category, whose objects are unital  $A_{\infty}$ -categories, 1-morphisms are unital  $A_{\infty}$ -functors and 2-morphisms are equivalence classes of natural  $A_{\infty}$ -transformations [Lyu03]. We continue to study this 2-category.

The notations and conventions are explained in the first section. We also describe  $A_N$ -categories,  $A_N$ -functors and  $A_N$ -transformations – truncated at  $N < \infty$  versions of  $A_\infty$ -categories. For instance,  $A_1$ -categories and  $A_1$ -functors are differential graded k-quivers and their morphisms. However,  $A_1$ -transformations bring new 2-categorical features to the theory. In particular, for any differential graded k-quiver Q and any  $A_\infty$ -category

<sup>\*</sup>Institute of Mathematics, National Academy of Sciences of Ukraine, 3 Tereshchenkivska st., Kyiv-4, 01601 MSP, Ukraine; lub@imath.kiev.ua

<sup>&</sup>lt;sup>†</sup>Department of Algebra, Faculty of Mechanics and Mathematics, Kyiv Taras Shevchenko University, 64 Volodymyrska st., Kyiv, 01033, Ukraine; manzyuk@univ.kiev.ua

 $\mathcal{A}$  there is an  $A_{\infty}$ -category  $A_1(\mathcal{Q}, \mathcal{A})$ , whose objects are morphisms of differential graded  $\mathbb{k}$ -quivers  $\mathcal{Q} \to \mathcal{A}$ , and morphisms are  $A_1$ -transformations. We recall the terminology related to trees in Section 1.7.

In the second section we define the free  $A_{\infty}$ -category  $\mathcal{FQ}$  generated by a differential graded k-quiver Q. We classify functors from a free  $A_{\infty}$ -category  $\mathcal{FQ}$  to an arbitrary  $A_{\infty}$ -category  $\mathcal{A}$  in Proposition 2.3. In particular, the restriction map gives a bijection between the set of strict  $A_{\infty}$ -functors  $\mathcal{FQ} \to \mathcal{A}$  and the set of morphisms of differential graded k-quivers  $\mathcal{Q} \to (\mathcal{A}, m_1)$  (Corollary 2.4). We classify chain maps into complexes of transformations whose source is a free  $A_{\infty}$ -category in Proposition 2.8. Description of homotopies between such chain maps is given in Corollary 2.10. Assuming in addition that  $\mathcal{A}$  is unital, we obtain our main result: the restriction  $A_{\infty}$ -functor restr :  $A_{\infty}(\mathcal{FQ}, \mathcal{A}) \to$  $A_1(\mathcal{Q}, \mathcal{A})$  is an equivalence (Theorem 2.12).

In the third section we interpret  $A_{\infty}(\mathfrak{FQ}, \_)$  and  $A_1(\mathfrak{Q}, \_)$  as strict  $A_{\infty}^u$ -2-functors  $A_{\infty}^u \to A_{\infty}^u$ . Moreover, we interpret restr :  $A_{\infty}(\mathfrak{FQ}, \_) \to A_1(\mathfrak{Q}, \_)$  as an  $A_{\infty}^u$ -2-equivalence. In this sense the  $A_{\infty}$ -category  $\mathfrak{FQ}$  represents the  $A_{\infty}^u$ -2-functor  $A_1(\mathfrak{Q}, \_)$ . This is the 2-categorical meaning of freeness of  $\mathfrak{FQ}$ .

#### 1. Conventions and preliminaries

We keep the notations and conventions of [Lyu03, LO02], sometimes without explicit mentioning. Some of the conventions are recalled here.

We assume as in [Lyu03, LO02] that most quivers,  $A_{\infty}$ -categories, etc. are small with respect to some universe  $\mathscr{U}$ .

The ground ring  $\mathbb{k} \in \mathscr{U}$  is a unital associative commutative ring. A  $\mathbb{k}$ -module means a  $\mathscr{U}$ -small  $\mathbb{k}$ -module.

We use the right operators: the composition of two maps (or morphisms)  $f: X \to Y$ and  $g: Y \to Z$  is denoted  $fg: X \to Z$ ; a map is written on elements as  $f: x \mapsto xf = (x)f$ . However, these conventions are not used systematically, and f(x) might be used instead.

 $\mathbb{Z}$ -graded k-modules are functions  $X : \mathbb{Z} \ni d \mapsto X^d \in \mathbb{k}$ -mod. A simple computation shows that the product  $X = \prod_{\iota \in I} X_\iota$  in the category of  $\mathbb{Z}$ -graded k-modules of a family  $(X_\iota)_{\iota \in I}$  of objects  $X_\iota : d \mapsto X_\iota^d$  is given by  $X : \mathbb{Z} \ni d \mapsto X^d = \prod_{\iota \in I} X_\iota^d$ . Everywhere in this article the product of graded k-modules means the above product.

If P is a  $\mathbb{Z}$ -graded k-module, then sP = P[1] denotes the same k-module with the grading  $(sP)^d = P^{d+1}$ . The "identity" map  $P \to sP$  of degree -1 is also denoted s. The map s commutes with the components of the differential in an  $A_{\infty}$ -category ( $A_{\infty}$ -algebra) in the following sense:  $s^{\otimes n}b_n = m_n s$ .

Let C = C(k -mod) denote the differential graded category of complexes of k-modules. Actually, it is a symmetric closed monoidal category.

The cone of a chain of a chain map  $\alpha : P \to Q$  of complexes of k-modules is the graded k-module  $\operatorname{Cone}(\alpha) = Q \oplus P[1]$  with the differential  $(q, ps)d = (qd^Q + p\alpha, psd^{P[1]}) = (qd^Q + p\alpha, -pd^Ps).$ 

**1.1.**  $A_N$ -categories. For a positive integer N we define some  $A_N$ -notions similarly to the case  $N = \infty$ . We may say that all data, equations and constructions for  $A_N$ -case are the same as in  $A_\infty$ -case (e.g. [Lyu03]), however, taken only up to level N.

A differential graded  $\Bbbk$ -quiver  $\Omega$  is the following data: a  $\mathscr{U}$ -small set of objects  $Ob \Omega$ ; a chain complex of  $\Bbbk$ -modules  $\Omega(X, Y)$  for each pair of objects X, Y. A morphism of differential graded  $\Bbbk$ -quivers  $f : \Omega \to \mathcal{A}$  is given by a map  $f : Ob \Omega \to Ob \mathcal{A}, X \mapsto Xf$ and by a chain map  $\Omega(X, Y) \to \mathcal{A}(Xf, Yf)$  for each pair of objects X, Y of  $\Omega$ . The category of differential graded  $\Bbbk$ -quivers is denoted  $A_1$ .

The category of  $\mathscr{U}$ -small graded k-linear quivers, whose set of objects is S, admits a symmetric monoidal structure with the tensor product  $\mathcal{A} \times \mathcal{B} \mapsto \mathcal{A} \otimes \mathcal{B}$ ,  $(\mathcal{A} \otimes \mathcal{B})(X, Y) = \bigoplus_{Z \in S} \mathcal{A}(X, Z) \otimes_{\Bbbk} \mathcal{B}(Z, Y)$ . In particular, we have tensor powers  $T^n \mathcal{A} = \mathcal{A}^{\otimes n}$  of a given graded k-quiver  $\mathcal{A}$ , such that  $\operatorname{Ob} T^n \mathcal{A} = \operatorname{Ob} \mathcal{A}$ . Explicitly,

$$T^{n}\mathcal{A}(X,Y) = \bigoplus_{X=X_{0},X_{1},\dots,X_{n}=Y\in Ob\mathcal{A}} \mathcal{A}(X_{0},X_{1}) \otimes_{\Bbbk} \mathcal{A}(X_{1},X_{2}) \otimes_{\Bbbk} \dots \otimes_{\Bbbk} \mathcal{A}(X_{n-1},X_{n}).$$

In particular,  $T^0\mathcal{A}(X,Y) = \mathbb{k}$  if X = Y and vanishes otherwise. The graded  $\mathbb{k}$ -quiver  $T^{\leq N}\mathcal{A} = \bigoplus_{n\geq 0}^{N} T^n\mathcal{A}$  is called the restricted tensor coalgebra of  $\mathcal{A}$ . It is equipped with the cut comultiplication

$$\Delta: T^{\leqslant N} \mathcal{A}(X, Y) \to \bigoplus_{Z \in Ob \mathcal{A}} T^{\leqslant N} \mathcal{A}(X, Z) \bigotimes_{\Bbbk} T^{\leqslant N} \mathcal{A}(Z, Y),$$
$$h_1 \otimes h_2 \otimes \cdots \otimes h_n \mapsto \sum_{k=0}^n h_1 \otimes \cdots \otimes h_k \bigotimes h_{k+1} \otimes \cdots \otimes h_n,$$

and the counit  $\varepsilon = (T^{\leq N} \mathcal{A}(X, Y) \xrightarrow{\mathrm{pr}_0} T^0 \mathcal{A}(X, Y) \to \Bbbk)$ , where the last map is  $\mathrm{id}_{\Bbbk}$  if X = Y, or 0 if  $X \neq Y$  (and  $T^0 \mathcal{A}(X, Y) = 0$ ). We write  $T\mathcal{A}$  instead of  $T^{\leq \infty} \mathcal{A}$ . If  $g: T\mathcal{A} \to T\mathcal{B}$  is a map of  $\Bbbk$ -quivers, then  $g_{ac}$  denotes its matrix coefficient  $T^a \mathcal{A} \subset T^a \mathcal{A} \xrightarrow{\mathrm{pr}_c} T\mathcal{B} \xrightarrow{\mathrm{pr}_c} T^c \mathcal{B}$ . The matrix coefficient  $g_{a1}$  is abbreviated to  $g_a$ .

**1.2 Definition.** An  $A_N$ -category  $\mathcal{A}$  consists of the following data: a graded k-quiver  $\mathcal{A}$ ; a system of k-linear maps of degree 1

$$b_n: s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \dots \otimes s\mathcal{A}(X_{n-1}, X_n) \to s\mathcal{A}(X_0, X_n), \qquad 1 \leqslant n \leqslant N,$$

such that for all  $1 \leq k \leq N$ 

$$\sum_{r+n+t=k} (1^{\otimes r} \otimes b_n \otimes 1^{\otimes t}) b_{r+1+t} = 0 : T^k s \mathcal{A} \to s \mathcal{A}.$$
(1.2.1)

The system  $b_n$  is interpreted as a (1,1)-coderivation  $b: T^{\leq N} s \mathcal{A} \to T^{\leq N} s \mathcal{A}$  of degree 1 determined by

$$b_{kl} = (b\big|_{T^k s \mathcal{A}}) \operatorname{pr}_l : T^k s \mathcal{A} \to T^l s \mathcal{A}, \qquad b_{kl} = \sum_{\substack{r+n+t=k\\r+1+t=l}} 1^{\otimes r} \otimes b_n \otimes 1^{\otimes t}, \qquad k, l \leqslant N,$$

which is a differential.

**1.3 Definition.** A pointed cocategory homomorphism consists of the following data:  $A_N$ -categories  $\mathcal{A}$  and  $\mathcal{B}$ , a map  $f : \operatorname{Ob} \mathcal{A} \to \operatorname{Ob} \mathcal{B}$  and a system of k-linear maps of degree 0

$$f_n: s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \dots \otimes s\mathcal{A}(X_{n-1}, X_n) \to s\mathcal{B}(X_0 f, X_n f), \qquad 1 \leqslant n \leqslant N.$$

The above data are equivalent to a cocategory homomorphism  $f: T^{\leq N} s \mathcal{A} \to T^{\leq N} s \mathcal{B}$  of degree 0 such that

$$f_{01} = (f|_{T^0 s \mathcal{A}}) \operatorname{pr}_1 = 0 : T^0 s \mathcal{A} \to T^1 s \mathcal{B},$$
 (1.3.1)

(this condition was implicitly assumed in [Lyu03, Definition 2.4]). The components of f are

$$f_{kl} = (f|_{T^k s \mathcal{A}}) \operatorname{pr}_l : T^k s \mathcal{A} \to T^l s \mathcal{B}, \qquad f_{kl} = \sum_{i_1 + \dots + i_l = k} f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_l}, \qquad (1.3.2)$$

where  $k, l \leq N$ . Indeed, the claim follows from the following diagram, commutative for all  $l \geq 0$ :

$$T^{\leqslant N} s\mathcal{A} \xrightarrow{f} T^{\leqslant N} s\mathcal{B} \xrightarrow{\operatorname{pr}_{l}} T^{l} s\mathcal{B}$$

$$\Delta^{(l)} \downarrow = \Delta^{(l)} \downarrow = \parallel$$

$$(T^{\leqslant N} s\mathcal{A})^{\otimes l} \xrightarrow{f^{\otimes l}} (T^{\leqslant N} s\mathcal{B})^{\otimes l} \xrightarrow{\operatorname{pr}_{1}^{\otimes l}} (s\mathcal{B})^{\otimes l}$$

where  $\Delta^{(0)} = \varepsilon$ ,  $\Delta^{(1)} = id$ ,  $\Delta^{(2)} = \Delta$  and  $\Delta^{(l)}$  means the cut comultiplication, iterated l-1 times. Notice that condition (1.3.1) can be written as  $f_0 = 0$ .

**1.4 Definition.** An  $A_N$ -functor  $f : \mathcal{A} \to \mathcal{B}$  is a pointed cocategory homomorphism, which commutes with the differential b, that is, for all  $1 \leq k \leq N$ 

$$\sum_{l>0;i_1+\cdots+i_l=k} (f_{i_1}\otimes f_{i_2}\otimes\cdots\otimes f_{i_l})b_l = \sum_{r+n+t=k} (1^{\otimes r}\otimes b_n\otimes 1^{\otimes t})f_{r+1+t}: T^k s\mathcal{A} \to s\mathcal{B}.$$

We are interested mostly in the case N = 1. Clearly,  $A_1$ -categories are differential graded quivers and  $A_1$ -functors are their morphisms. In the case of one object these reduce to chain complexes and chain maps. The following notion seems interesting even in this case.

**1.5 Definition.** An  $A_N$ -transformation  $r : f \to g : \mathcal{A} \to \mathcal{B}$  of degree d consists of the following data:  $A_N$ -categories  $\mathcal{A}$  and  $\mathcal{B}$ ; pointed cocategory homomorphisms  $f, g : T^{\leq N} s \mathcal{A} \to T^{\leq N} s \mathcal{B}$  (or  $A_N$ -functors  $f, g : \mathcal{A} \to \mathcal{B}$ ); a system of k-linear maps of degree d

$$r_n: s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \cdots \otimes s\mathcal{A}(X_{n-1}, X_n) \to s\mathcal{B}(X_0f, X_ng), \qquad 0 \leqslant n \leqslant N.$$

To give a system  $r_n$  is equivalent to specifying an (f,g)-coderivation  $r: T^{\leq N} s \mathcal{A} \to T^{\leq N+1} s \mathcal{B}$  of degree d

$$r_{kl} = (r|_{T^{k}s\mathcal{A}}) \operatorname{pr}_{l} : T^{k}s\mathcal{A} \to T^{l}s\mathcal{B}, \qquad k \leq N, \ l \leq N+1$$
  
$$r_{kl} = \sum_{\substack{q+1+t=l\\i_{1}+\dots+i_{q}+n+j_{1}+\dots+j_{t}=k}} f_{i_{1}} \otimes \dots \otimes f_{i_{q}} \otimes r_{n} \otimes g_{j_{1}} \otimes \dots \otimes g_{j_{t}}, \qquad (1.5.1)$$

that is, a k-quiver morphism r, satisfying  $r\Delta = \Delta(f \otimes r + r \otimes g)$ . This follows from the commutative diagram

$$\begin{array}{cccc} T^{\leqslant N} s\mathcal{A} & & \xrightarrow{r} & T^{\leqslant N+1} s\mathcal{B} & \xrightarrow{\operatorname{pr}_{l}} & T^{l} s\mathcal{B} \\ & & & & & \\ \Delta^{(l)} \downarrow & = & & & \\ (T^{\leqslant N} s\mathcal{A})^{\otimes l} & & & \xrightarrow{\sum_{q+1+t=l} f^{\otimes q} \otimes r \otimes g^{\otimes t}} & (T^{\leqslant N+1} s\mathcal{B})^{\otimes l} & \xrightarrow{\operatorname{pr}_{1}^{\otimes l}} & (s\mathcal{B})^{\otimes l} \end{array}$$

Let  $\mathcal{A}$ ,  $\mathcal{B}$  be  $A_N$ -categories, and let  $f^0, f^1, \ldots, f^n : T^{\leq N} s \mathcal{A} \to T^{\leq N} s \mathcal{B}$  be pointed cocategory homomorphisms. Consider coderivations  $r_1, \ldots, r_n$  as in

$$f^0 \xrightarrow{r^1} f^1 \xrightarrow{r^2} \dots f^{n-1} \xrightarrow{r^n} f^n : T^{\leqslant N} s \mathcal{A} \to T^{\leqslant N} s \mathcal{B}.$$

We construct the following system of k-linear maps  $\theta_{kl} : T^k s \mathcal{A} \to T^l s \mathcal{B}, k \leq N, l \leq N+n$  of degree deg  $r^1 + \cdots + \deg r^n$  from these data:

$$\theta_{kl} = \sum f_{i_1^0}^0 \otimes \cdots \otimes f_{i_{m_0}^0}^0 \otimes r_{j_1}^1 \otimes f_{i_1^1}^1 \otimes \cdots \otimes f_{i_{m_1}^1}^1 \otimes \cdots \otimes r_{j_n}^n \otimes f_{i_1^n}^n \otimes \cdots \otimes f_{i_{m_n}^n}^n, \quad (1.5.2)$$

where summation is taken over all terms with

 $m_0 + m_1 + \dots + m_n + n = l, \quad i_1^0 + \dots + i_{m_0}^0 + j_1 + i_1^1 + \dots + i_{m_1}^1 + \dots + j_n + i_1^n + \dots + i_{m_n}^n = k.$ Equivalently, we write

$$\theta_{kl} = \sum_{\substack{m_0+m_1+\dots+m_n+n=l\\p_0+j_1+p_1+\dots+j_n+p_n=k}} f_{p_0m_0}^0 \otimes r_{j_1}^1 \otimes f_{p_1m_1}^1 \otimes \dots \otimes r_{j_n}^n \otimes f_{p_nm_n}^n$$

The component  $\theta_{kl}$  vanishes unless  $n \leq l \leq k+n$ . If n = 0, then  $\theta_{kl}$  is expansion (1.3.2) of  $f^0$ . If n = 1, then  $\theta_{kl}$  is expansion (1.5.1) of  $r^1$ .

Given an  $A_K$ -category  $\mathcal{A}$  and an  $A_{K+N}$ -category  $\mathcal{B}$ ,  $1 \leq K, N \leq \infty$ , we construct an  $A_N$ -category  $A_K(\mathcal{A}, \mathcal{B})$  out of these. The objects of  $A_K(\mathcal{A}, \mathcal{B})$  are  $A_K$ -functors  $f : \mathcal{A} \to \mathcal{B}$ . Given two such functors  $f, g : \mathcal{A} \to \mathcal{B}$  we define the graded k-module  $A_K(\mathcal{A}, \mathcal{B})(f, g)$  as the space of all  $A_K$ -transformations  $r : f \to g$ , namely,

$$\begin{split} & [A_K(\mathcal{A}, \mathcal{B})(f, g)]^{d+1} \\ & = \{r: f \to g \mid A_K \text{-transformation } r: T^{\leqslant K} s \mathcal{A} \to T^{\leqslant K+1} s \mathcal{B} \text{ has degree } d\}. \end{split}$$

The system of differentials  $B_n$ ,  $n \leq N$ , is defined as follows:

$$B_{1}: A_{K}(\mathcal{A}, \mathcal{B})(f, g) \to A_{K}(\mathcal{A}, \mathcal{B})(f, g), \quad r \mapsto (r)B_{1} = [r, b] = rb - (-)^{r}br,$$

$$[(r)B_{1}]_{k} = \sum_{i_{1}+\dots+i_{q}+n+j_{1}+\dots+j_{t}=k} (f_{i_{1}} \otimes \dots \otimes f_{i_{q}} \otimes r_{n} \otimes g_{j_{1}} \otimes \dots \otimes g_{j_{t}})b_{q+1+t}$$

$$-(-)^{r} \sum_{\alpha+n+\beta=k} (1^{\otimes \alpha} \otimes b_{n} \otimes 1^{\otimes \beta})r_{\alpha+1+\beta}, \quad k \leq K,$$

$$B_{n}: A_{K}(\mathcal{A}, \mathcal{B})(f^{0}, f^{1}) \otimes \dots \otimes A_{K}(\mathcal{A}, \mathcal{B})(f^{n-1}, f^{n}) \to A_{K}(\mathcal{A}, \mathcal{B})(f^{0}, f^{n}),$$

$$r^{1} \otimes \dots \otimes r^{n} \mapsto (r^{1} \otimes \dots \otimes r^{n})B_{n}, \text{ for } 1 < n \leq N,$$

where the last  $A_K$ -transformation is defined by its components:

$$[(r^1 \otimes \cdots \otimes r^n)B_n]_k = \sum_{l=n}^{n+k} (r^1 \otimes \cdots \otimes r^n)\theta_{kl}b_l, \qquad k \leqslant K.$$

The category of graded k-linear quivers admits a symmetric monoidal structure with the tensor product  $\mathcal{A} \times \mathcal{B} \mapsto \mathcal{A} \boxtimes \mathcal{B}$ , where  $\operatorname{Ob} \mathcal{A} \boxtimes \mathcal{B} = \operatorname{Ob} \mathcal{A} \times \operatorname{Ob} \mathcal{B}$  and  $(\mathcal{A} \boxtimes \mathcal{B})((X,U),(Y,V)) = \mathcal{A}(X,Y) \otimes_{\Bbbk} \mathcal{B}(U,V)$ . The same tensor product was denoted  $\otimes$ in [Lyu03], but we will keep notation  $\mathcal{A} \otimes \mathcal{B}$  only for tensor product from Section 1.1, defined when  $\operatorname{Ob} \mathcal{A} = \operatorname{Ob} \mathcal{B}$ . The two tensor products obey

**Distributivity law.** Let  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{C}$ ,  $\mathcal{D}$  be graded k-linear quivers, such that  $Ob \mathcal{A} = Ob \mathcal{B}$  and  $Ob \mathcal{C} = Ob \mathcal{D}$ . Then the middle four interchange map  $1 \otimes c \otimes 1$  is an isomorphism of quivers

$$(\mathcal{A} \otimes \mathcal{B}) \boxtimes (\mathcal{C} \otimes \mathcal{D}) \xrightarrow{\sim} (\mathcal{A} \boxtimes \mathcal{C}) \otimes (\mathcal{B} \boxtimes \mathcal{D}), \tag{1.5.3}$$

identity on objects.

Indeed, the both quivers in (1.5.3) have the same set of objects  $R \times S$ , where  $R = Ob \mathcal{A} = Ob \mathcal{B}$  and  $S = Ob \mathcal{C} = Ob \mathcal{D}$ . Let  $X, Z \in R$  and  $U, W \in S$ . The sets of morphisms from (X, U) to (Z, W) are isomorphic via

$$((\mathcal{A} \otimes \mathcal{B}) \boxtimes (\mathfrak{C} \otimes \mathcal{D}))((X, U), (Z, W)) = (\oplus_{Y \in R} \mathcal{A}(X, Y) \otimes_{\Bbbk} \mathcal{B}(Y, Z)) \otimes_{\Bbbk} (\oplus_{V \in S} \mathfrak{C}(U, V) \otimes_{\Bbbk} \mathcal{D}(V, W)) \downarrow_{l} \oplus_{(Y,V) \in R \times S} \mathcal{A}(X, Y) \otimes_{\Bbbk} \mathcal{B}(Y, Z) \otimes_{\Bbbk} \mathfrak{C}(U, V) \otimes_{\Bbbk} \mathcal{D}(V, W) \downarrow_{1 \otimes c \otimes 1} \oplus_{(Y,V) \in R \times S} \mathcal{A}(X, Y) \otimes_{\Bbbk} \mathfrak{C}(U, V) \otimes_{\Bbbk} \mathcal{B}(Y, Z) \otimes_{\Bbbk} \mathcal{D}(V, W) = ((\mathcal{A} \boxtimes \mathfrak{C}) \otimes (\mathcal{B} \boxtimes \mathcal{D}))((X, U), (Z, W)).$$

The notion of a pointed cocategory homomorphism extends to the case of several arguments, that is, to degree 0 cocategory homomorphisms  $\psi: T^{\leq L^1}s\mathfrak{C}^1\boxtimes\cdots\boxtimes T^{\leq L^q}s\mathfrak{C}^q \to$ 

 $T^{\leq N}s\mathcal{B}$ , where  $N \geq L^1 + \cdots + L^q$ . We always assume that  $\psi_{00\dots 0} : T^0s\mathcal{C}^1 \boxtimes \cdots \boxtimes T^0s\mathcal{C}^q \to s\mathcal{B}$  vanishes. We call  $\psi$  an A-functor if it commutes with the differential, that is,

$$(b \boxtimes 1 \boxtimes \cdots \boxtimes 1 + 1 \boxtimes b \boxtimes \cdots \boxtimes 1 + \cdots + 1 \boxtimes 1 \boxtimes \cdots \boxtimes b)\psi = \psi b.$$

For example, the map  $\alpha : T^{\leq K} s \mathcal{A} \boxtimes T^{\leq N} s A_K(\mathcal{A}, \mathcal{B}) \to T^{\leq K+N} s \mathcal{B}, \ a \boxtimes r^1 \otimes \cdots \otimes r^n \mapsto a.[(r^1 \otimes \cdots \otimes r^n)\theta]$ , is an A-functor.

**1.6 Proposition (cf. Proposition 5.5 of [Lyu03]).** Let  $\mathcal{A}$  be an  $A_K$ -category, let  $\mathcal{C}^t$  be an  $A_{L^t}$ -category for  $1 \leq t \leq q$ , and let  $\mathcal{B}$  be an  $A_N$ -category, where  $N \geq K + L^1 + \cdots + L^q$ . For any A-functor  $\phi : T^{\leq K} s \mathcal{A} \boxtimes T^{\leq L^1} s \mathcal{C}^1 \boxtimes \cdots \boxtimes T^{\leq L^q} s \mathcal{C}^q \to T^{\leq N} s \mathcal{B}$  there is a unique A-functor  $\psi : T^{\leq L^1} s \mathcal{C}^1 \boxtimes \cdots \boxtimes T^{\leq L^q} s \mathcal{C}^q \to T^{\leq N-K} s A_K(\mathcal{A}, \mathcal{B})$ , such that

$$\phi = \left( T^{\leqslant K} s \mathcal{A} \boxtimes T^{\leqslant L^1} s \mathfrak{C}^1 \boxtimes \cdots \boxtimes T^{\leqslant L^q} s \mathfrak{C}^q \xrightarrow{1 \boxtimes \psi} T^{\leqslant K} s \mathcal{A} \boxtimes T^{\leqslant N-K} s A_K(\mathcal{A}, \mathcal{B}) \xrightarrow{\alpha} T^{\leqslant N} s \mathcal{B} \right).$$

Let  $\mathcal{A}$  be an  $A_N$ -category, let  $\mathcal{B}$  be an  $A_{N+K}$ -category, and let  $\mathcal{C}$  be an  $A_{N+K+L}$ -category. The above proposition implies the existence of an A-functor (cf. [Lyu03, Proposition 4.1])

$$M: T^{\leqslant K} sA_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leqslant L} sA_{N+K}(\mathcal{B}, \mathcal{C}) \to T^{\leqslant K+L} sA_N(\mathcal{A}, \mathcal{C}),$$

in particular,  $(1 \boxtimes B + B \boxtimes 1)M = MB$ . It has the components

$$M_{nm} = M \big|_{T^n \boxtimes T^m} \operatorname{pr}_1 : T^n s A_N(\mathcal{A}, \mathcal{B}) \boxtimes T^m s A_{N+K}(\mathcal{B}, \mathcal{C}) \to s A_N(\mathcal{A}, \mathcal{C}),$$

 $n \leq K$ ,  $m \leq L$ . We have  $M_{00} = 0$  and  $M_{nm} = 0$  for m > 1. If m = 0 and n is positive,  $M_{n0}$  is given by the formula:

$$M_{n0}: sA_N(\mathcal{A}, \mathcal{B})(f^0, f^1) \otimes \cdots \otimes sA_N(\mathcal{A}, \mathcal{B})(f^{n-1}, f^n) \boxtimes \Bbbk_{g^0} \to sA_N(\mathcal{A}, \mathcal{C})(f^0g^0, f^ng^0),$$
$$r^1 \otimes \cdots \otimes r^n \boxtimes 1 \mapsto (r^1 \otimes \cdots \otimes r^n \mid g^0)M_{n0},$$

$$[(r^1 \otimes \cdots \otimes r^n \mid g^0) M_{n0}]_k = \sum_{l=n}^{n+k} (r^1 \otimes \cdots \otimes r^n) \theta_{kl} g_l^0, \qquad k \leqslant N,$$

where | separates the arguments in place of  $\boxtimes$ . If m = 1, then  $M_{n1}$  is given by the formula:

$$M_{n1}: sA_N(\mathcal{A}, \mathcal{B})(f^0, f^1) \otimes \cdots \otimes sA_N(\mathcal{A}, \mathcal{B})(f^{n-1}, f^n) \boxtimes sA_{N+K}(\mathcal{B}, \mathcal{C})(g^0, g^1) \to sA_N(\mathcal{A}, \mathcal{C})(f^0g^0, f^ng^1), \qquad r^1 \otimes \cdots \otimes r^n \boxtimes t^1 \mapsto (r^1 \otimes \cdots \otimes r^n \boxtimes t^1)M_{n1},$$

$$[(r^1 \otimes \cdots \otimes r^n \boxtimes t^1)M_{n1}]_k = \sum_{l=n}^{n+k} (r^1 \otimes \cdots \otimes r^n)\theta_{kl}t_l^1, \qquad k \leqslant N.$$

Note that equations

$$[(r^1 \otimes \cdots \otimes r^n)B_n]_k = [(r^1 \otimes \cdots \otimes r^n \boxtimes b)M_{n1}]_k - (-)^{r^1 + \cdots + r^n}[(b \boxtimes r^1 \otimes \cdots \otimes r^n)M_{1n}]_k$$

imply that

$$(r^{1} \otimes \cdots \otimes r^{n})B_{n} = (r^{1} \otimes \cdots \otimes r^{n} \boxtimes b)M_{n1} - (-)^{r^{1} + \cdots + r^{n}}(b \boxtimes r^{1} \otimes \cdots \otimes r^{n})M_{1n},$$
  
$$B = (1 \boxtimes b)M - (b \boxtimes 1)M : \mathrm{id} \to \mathrm{id} : A_{N}(\mathcal{A}, \mathcal{B}) \to A_{N}(\mathcal{A}, \mathcal{B}).$$

Proposition 1.6 implies the existence of a unique  $A_L$ -functor

$$A_N(\mathcal{A},\_): A_{N+K}(\mathcal{B},\mathfrak{C}) \to A_K(A_N(\mathcal{A},\mathfrak{B}),A_N(\mathcal{A},\mathfrak{C})),$$

such that

$$M = \left[ T^{\leqslant K} s A_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leqslant L} s A_{N+K}(\mathcal{B}, \mathcal{C}) \xrightarrow{1\boxtimes A_N(\mathcal{A}, \cdot)} T^{\leqslant K} s A_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leqslant L} s A_K(A_N(\mathcal{A}, \mathcal{B}), A_N(\mathcal{A}, \mathcal{C})) \xrightarrow{\alpha} T^{\leqslant K+L} s A_N(\mathcal{A}, \mathcal{C}) \right].$$

The  $A_L$ -functor  $A_N(\mathcal{A}, \_)$  is strict, cf. [Lyu03, Proposition 6.2].

Let  $\mathcal{A}$  be an  $A_N$ -category, and let  $\mathcal{B}$  be a unital  $A_\infty$ -category with a unit transformation  $\mathbf{i}^{\mathcal{B}}$ . Then  $A_N(\mathcal{A}, \mathcal{B})$  is a unital  $A_\infty$ -category with the unit transformation  $(1 \boxtimes \mathbf{i}^{\mathcal{B}})M$  (cf. [Lyu03, Proposition 7.7]). The unit element for an object  $f \in \operatorname{Ob} A_N(\mathcal{A}, \mathcal{B})$  is  $_{\mathbf{i}}\mathbf{i}_0^{A_N(\mathcal{A},\mathcal{B})} : \mathbb{k} \to (sA_N)^{-1}(\mathcal{A}, \mathcal{B}), 1 \mapsto f\mathbf{i}^{\mathcal{B}}$ .

When  $\mathcal{A}$  is an  $A_K$ -category and N < K, we may forget part of its structure and view  $\mathcal{A}$ as an  $A_N$ -category. If furthermore,  $\mathcal{B}$  is an  $A_{K+L}$ -category, we have the restriction strict  $A_L$ -functor restr<sub>K,N</sub> :  $A_K(\mathcal{A}, \mathcal{B}) \to A_N(\mathcal{A}, \mathcal{B})$ . To prove the results mentioned above, we notice that they are restrictions of their  $A_\infty$ -analogs to finite N. Since the proofs of  $A_\infty$ -results are obtained in [Lyu03] by induction, an inspection shows that the proofs of the above  $A_N$ -statements are obtained as a byproduct.

1.7. Trees. Since the notions related to trees might be interpreted with some variations, we give precise definitions and fix notation. A *tree* is a non-empty connected graph without cycles. A vertex which belongs to only one edge is called *external*, other vertices are *internal*. A *plane tree* is a tree equipped for each internal vertex v with a cyclic ordering of the set  $E_v$  of edges, adjacent to v. Plane trees can be drawn on an oriented plane in a unique way (up to an ambient isotopy) so that the cyclic ordering of each  $E_v$  agrees with the orientation of the plane. An external vertex distinct from the root is called *input vertex*.

A rooted tree is a tree with a distinguished external vertex, called root. The set of vertices V(t) of a rooted tree t has a canonical ordering:  $x \preccurlyeq y$  iff the minimal path connecting the root with y contains x. A linearly ordered tree is a rooted tree t equipped with a linear order  $\leq$  of the set of internal vertices IV(t), such that for all internal vertices x, y the relation  $x \preccurlyeq y$  implies  $x \leqslant y$ . For each vertex  $v \in V(t) - {\text{root}}$  of a rooted tree, the set  $E_v$  has a distinguished element  $e_v$  – the beginning of a minimal path from v to the root. Therefore, for each vertex  $v \in V(t) - {\text{root}}$  of a rooted plane tree, the set  $E_v$  admits a unique linear order <, for which  $e_v$  is minimal and the induced cyclic order is the given one. An internal vertex v has degree d, if  $Card(E_v) = d + 1$ .

For any  $y \in V(t)$  let  $P_y = \{x \in V(t) \mid x \preccurlyeq y\}$ . With each plane rooted tree t is associated a linearly ordered tree  $t_{\leq} = (t, \leq)$  as follows. If  $x, y \in IV(t)$  are such that  $x \preccurlyeq y$  and  $y \preccurlyeq x$ , then  $P_x \cap P_y = P_z$  for a unique  $z \in IV(t)$ , distinct from x and y. Let  $a \in E_z - \{e_z\}$  (resp.  $b \in E_z - \{e_z\}$ ) be the beginning of the minimal path connecting z and x (resp. y). If a < b, we set x < y. Graphically we <-order the internal vertices by height. Thus, an internal vertex x on the left is depicted lower than a  $\preccurlyeq$ -incomparable internal vertex y on the right:



A forest is a sequence of plane rooted trees. Concatenation of forests is denoted  $\sqcup$ . The vertical composition  $F_1 \cdot F_2$  of forests  $F_1$ ,  $F_2$  is well-defined if the sum of lengths of sequences  $F_1$  and  $F_2$  equals the number of external vertices of  $F_2$ . These operations allow to construct any tree from elementary ones

$$1 = |$$
, and  $\mathfrak{t}_k =$  (k input vertices).

Namely, any linearly ordered tree  $(t, \leq)$  has a unique presentation of the form

$$(t,\leqslant) = (1^{\sqcup\alpha_1} \sqcup \mathfrak{t}_{k_1} \sqcup 1^{\sqcup\beta_1}) \cdot (1^{\sqcup\alpha_2} \sqcup \mathfrak{t}_{k_2} \sqcup 1^{\sqcup\beta_2}) \cdot \ldots \cdot \mathfrak{t}_{k_N}, \qquad (1.7.1)$$

where  $N = |t| \stackrel{\text{def}}{=} \operatorname{Card}(IV(t))$  is the number of internal vertices. Here

$$1^{\sqcup \alpha} \sqcup \mathfrak{t}_k \sqcup 1^{\sqcup \beta} = \overbrace{\left[\begin{array}{c} \\ \\ \\ \end{array}\right]}^{\alpha} \cdots \overbrace{\left[\begin{array}{c} \\ \\ \\ \end{array}\right]}^{k} \overbrace{\left[\begin{array}{c} \\ \\ \\ \\ \end{array}\right]}^{m}} \overbrace{\left[\begin{array}{c} \\ \\ \\ \\ \end{array}\right]}^{m} \overbrace{\left[\begin{array}{c} \\ \\ \\ \\ \end{array}\right]}^{m} \overbrace{\left[\begin{array}{c} \\ \\ \\ \\ \end{array}\right]}^{\beta} \overbrace{\left[\begin{array}{c} \\ \\ \\ \\ \\ \end{array}\right]}^{m}$$

In (1.7.1) the highest vertex is indexed by 1, the lowest – by N.

## 2. Properties of free $A_{\infty}$ -categories

**2.1.** Construction of a free  $A_{\infty}$ -category. The category strict  $A_{\infty}$  has  $A_{\infty}$ -categories as objects and strict  $A_{\infty}$ -functors as morphisms. There is a functor  $\mathcal{U}$ : strict  $A_{\infty} \to A_1$ ,  $\mathcal{A} \mapsto (\mathcal{A}, m_1)$  which sends an  $A_{\infty}$ -category to the underlying differential graded k-quiver, forgetting all higher multiplications. Following Kontsevich and Soibelman [KS02] we are going to prove that  $\mathcal{U}$  has a left adjoint functor  $\mathcal{F} : A_1 \to \text{strict} A_{\infty}, \mathcal{Q} \mapsto \mathcal{FQ}$ . The  $A_{\infty}$ -category  $\mathcal{FQ}$  is called free. Below we describe its structure for an arbitrary differential graded k-quiver  $\mathcal{Q}$ . We shall work with its shift  $(s\mathcal{Q}, d)$ . Let us define an  $A_{\infty}$ -category  $\mathcal{FQ}$  via the following data. The class of objects  $Ob \mathcal{FQ}$  is  $Ob \mathcal{Q}$ . The  $\mathbb{Z}$ -graded k-modules of morphisms between  $X, Y \in Ob \mathcal{Q}$  are

$$s\mathcal{FQ}(X,Y) = \bigoplus_{n \ge 1} \bigoplus_{t \in \mathfrak{T}_{\ge 2}^n} s\mathcal{F}_t \mathcal{Q}(X,Y),$$
  
$$s\mathcal{F}_t \mathcal{Q}(X,Y) = \bigoplus_{X_0,\dots,X_n \in Ob \ \mathcal{Q}}^{X_0=X, X_n=Y} s\mathcal{Q}(X_0,X_1) \otimes \dots \otimes s\mathcal{Q}(X_{n-1},X_n) [-|t|],$$

where  $\mathbb{T}_{\geq 2}^n$  is the class of plane rooted trees with n + 1 external vertices, such that  $\operatorname{Card}(E_v) \geq 3$  for all  $v \in IV(t)$ . We use the following convention: if M, N are (differential) graded k-modules, then<sup>1</sup>

$$(M \otimes N)[k] = M \otimes (N[k]),$$
$$(M \otimes N \xrightarrow{s^k} (M \otimes N)[k]) = (M \otimes N \xrightarrow{1 \otimes s^k} M \otimes (N[k])).$$

The quiver  $\mathcal{FQ}$  is equipped with the following operations. For k > 1 the operation  $b_k$  is a direct sum of maps

$$b_k = s^{|t_1|} \otimes \cdots \otimes s^{|t_{k-1}|} \otimes s^{|t_k| - |t|} : s\mathcal{F}_{t_1} \mathcal{Q}(Y_0, Y_1) \otimes \cdots \otimes s\mathcal{F}_{t_k} \mathcal{Q}(Y_{k-1}, Y_k) \to s\mathcal{F}_t \mathcal{Q}(Y_0, Y_k),$$
(2.1.1)

where  $t = (t_1 \sqcup \cdots \sqcup t_k) \cdot \mathbf{t}_k$ . In particular,  $|t| = |t_1| + \cdots + |t_k| + 1$ . The operation  $b_1$  restricted to  $s\mathcal{F}_t \mathcal{Q}$  is

$$b_1 = d \oplus (-1)^{\beta(t')} s^{-1} : s \mathcal{F}_t \mathcal{Q}(X, Y) \to s \mathcal{F}_t \mathcal{Q}(X, Y) \oplus \bigoplus_{t' = t + \text{edge}} s \mathcal{F}_{t'} \mathcal{Q}(X, Y), \qquad (2.1.2)$$

where the sum extends over all trees  $t' \in \mathcal{T}_{\geq 2}^n$  with a distinguished edge e, such that contracting e we get t from t'. The sign is determined by

$$\beta(t') = \beta(t', e) = 1 + h(\text{highest vertex of } e),$$

where an isomorphism of ordered sets

$$h: IV(t'_{<}) \xrightarrow{\sim} [1, |t'|] \cap \mathbb{Z}$$

is simply the height of a vertex in the linearly ordered tree  $t'_{<}$ , canonically associated with t'. In (2.1.2) d means  $d \otimes 1 \otimes \cdots \otimes 1 + \cdots + 1 \otimes \cdots \otimes d \otimes 1 + 1 \otimes \cdots \otimes 1 \otimes d$ , where the last d is  $d_{sQ[-|t|]} = (-)^{|t|} s^{|t|} \cdot d_{sQ} \cdot s^{-|t|}$ , as usual. According to our conventions,  $s^{-1}$  in (2.1.2) means  $1^{\otimes n-1} \otimes s^{-1}$ .

#### **2.2 Proposition.** $\mathfrak{FQ}$ is an $A_{\infty}$ -category.

*Proof.* First we prove that  $b_1^2 = 0$  on  $s \mathcal{F}_t Q$ . Indeed,

$$b_1^2 = d^2 \oplus (-)^{\beta(t',e)} (s^{-1}d + ds^{-1}) \oplus \left[ (-1)^{\beta(t'_1,e_1) + \beta(t'',e_2)} + (-1)^{\beta(t'_2,e_2) + \beta(t'',e_1)} \right] s^{-2} :$$

$$s \mathcal{F}_t \mathcal{Q} \to s \mathcal{F}_t \mathcal{Q} \oplus \bigoplus_{t'=t+e} s \mathcal{F}_{t'} \mathcal{Q} \oplus \bigoplus_{t''=t+e_1+e_2} s \mathcal{F}_{t''} \mathcal{Q},$$

<sup>1</sup>Another gauge choice  $(M \otimes N)[1] = M[1] \otimes N$ ,  $s = s \otimes 1$  seems less convenient.

where t'' contains two distinguished edges  $e_1$ ,  $e_2$ , contraction along which gives t;  $t'_2$  is t'' contracted along  $e_1$ , and  $t'_1$  is t'' contracted along  $e_2$ . We may assume that highest vertex of  $e_1$  is lower than highest vertex of  $e_2$  in  $t''_{<}$ . Then  $\beta(t'_1, e_1) = \beta(t'', e_1)$  and  $\beta(t'', e_2) = \beta(t'_2, e_2) + 1$ , hence,

$$(-1)^{\beta(t'_1,e_1)+\beta(t'',e_2)} + (-1)^{\beta(t'_2,e_2)+\beta(t'',e_1)} = 0.$$

Obviously,  $d^2 = 0$  and  $s^{-1}d + ds^{-1} = 0$ , hence,  $b_1^2 = 0$ .

Let us prove for each n > 1 that

$$b_{n}b_{1} + \sum_{p=1}^{n} (1^{\otimes p-1} \otimes b_{1} \otimes 1^{\otimes n-p})b_{n} + \sum_{\substack{\alpha+k+\beta=n\\\alpha+\beta>0}}^{k>1} (1^{\otimes \alpha} \otimes b_{k} \otimes 1^{\otimes \beta})b_{\alpha+1+\beta} = 0:$$

$$s\mathcal{F}_{t_{1}}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_{n}}\mathcal{Q} \to s\mathcal{F}_{t}\mathcal{Q} \oplus \bigoplus_{p,t'_{p}} s\mathcal{F}_{t'}\mathcal{Q} \oplus \bigoplus_{\substack{\alpha+k+\beta=n\\\alpha+\beta>0}}^{k>1,t''} s\mathcal{F}_{t''}\mathcal{Q}.$$

where

$$t = (t_1 \sqcup \cdots \sqcup t_n) \cdot \mathfrak{t}_n = \underbrace{t_1 \qquad t_2 \qquad t_{n-1} \qquad t_n}_{t_1 \qquad t_2 \qquad t_{n-1} \qquad t_n}, \qquad (2.2.1)$$
$$t' = (t_1 \sqcup \cdots \sqcup t'_p \sqcup \cdots \sqcup t_n) \cdot \mathfrak{t}_n = \underbrace{t_1 \qquad t'_p \qquad t_n}_{t_1 \qquad t_2 \qquad t_n},$$

$$t'' = (t_1 \sqcup \cdots \sqcup t_n) \cdot (1^{\sqcup \alpha} \sqcup \mathfrak{t}_k \sqcup 1^{\sqcup \beta}) \cdot \mathfrak{t}_{\alpha+1+\beta}$$

$$= \underbrace{t_1 \qquad t_\alpha \qquad t_{\alpha+1}}_{=} \underbrace{t_{\alpha+k+1} \qquad t_n}_{=} ,$$

and contraction of  $t'_p$  along distinguished edge  $e_p$  gives  $t_p$ . According to the three types of summands in the target, the required equation follows from anticommutativity of the following three diagrams:

$$\begin{array}{c} s\mathcal{F}_{t_1} \mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n} \mathcal{Q} \xrightarrow{s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n| - |t|}}{b_n} s\mathcal{F}_t \mathcal{Q} \\ \xrightarrow{1^{\otimes n-1} \otimes d + \dots + d \otimes 1^{\otimes n-1}} \int & - & \int d \\ s\mathcal{F}_{t_1} \mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n} \mathcal{Q} \xrightarrow{s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n| - |t|}}{b_n} s\mathcal{F}_t \mathcal{Q} \end{array}$$

that is,

$$(1^{\otimes n-1} \otimes d + \dots + d \otimes 1^{\otimes n-1})(s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}) + (s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|})(1^{\otimes n-1} \otimes d + \dots + d \otimes 1^{\otimes n-1}) = 0;$$

$$s\mathcal{F}_{t_1} \mathbb{Q} \otimes \cdots \otimes s\mathcal{F}_{t_p} \mathbb{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n} \mathbb{Q} \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} s\mathcal{F}_t \mathbb{Q}$$

$$\downarrow^{(-)^{\beta(t'_p)} 1 \otimes p - 1} \otimes s^{-1} \otimes 1^{\otimes n - p} - (-)^{1 + |t_1| + \dots + |t_{p-1}| + \beta(t'_p)_{s^{-1}}} \downarrow$$

$$s\mathcal{F}_{t_1} \mathbb{Q} \otimes \cdots \otimes s\mathcal{F}_{t'_p} \mathbb{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n} \mathbb{Q} \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{p-1}|} \otimes s^{|t_p|+1} \otimes s^{|t_p|+1} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}} s\mathcal{F}_{t'} \mathbb{Q}$$

that is,

$$(-1)^{\beta(t'_{p})}(1^{\otimes p-1}\otimes s^{-1}\otimes 1^{\otimes n-p})(s^{|t_{1}|}\otimes\cdots\otimes s^{|t_{p-1}|}\otimes s^{|t_{p}|+1}\otimes s^{|t_{p+1}|}\otimes\cdots\otimes s^{|t_{n-1}|}\otimes s^{|t_{n}|-|t|-1}) + (s^{|t_{1}|}\otimes\cdots\otimes s^{|t_{n-1}|}\otimes s^{|t_{n-1}|}\otimes s^{|t_{n}|-|t|})(-1)^{1+|t_{1}|+\cdots+|t_{p-1}|+\beta(t'_{p})}(1^{\otimes n-1}\otimes s^{-1}) = 0,$$

in the particular case p = n it holds as well;

$$s\mathcal{F}_{t_1} Q \otimes \cdots \otimes s\mathcal{F}_{t_n} Q \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} s\mathcal{F}_t Q$$

$$1^{\otimes \alpha} \otimes s^{|t_{\alpha+1}|} \otimes \cdots \otimes s^{|t_{\alpha+k-1}|} \otimes s^{-|t_{\alpha+1}|-\cdots -|t_{\alpha+k-1}|-1} \otimes 1^{\otimes \beta} \xrightarrow{-} (-)^{1+|t_1|+\cdots +|t_{\alpha}|} s^{-1} \downarrow$$

$$s\mathcal{F}_{t_1} Q \otimes \cdots \otimes s\mathcal{F}_{t_{\alpha}} Q \otimes s\mathcal{F}_t Q \otimes s\mathcal{F}_{t_{\alpha}+k+1} Q \otimes \cdots \otimes s\mathcal{F}_{t_n} Q \xrightarrow{|t_1| \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}+k+1}| \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}+k+1}| \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}+k+1}| \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}+k+1}| \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha}+k+1}| \otimes$$

where  $\hat{t} = (t_{\alpha+1} \sqcup \cdots \sqcup t_{\alpha+k}) \cdot \mathfrak{t}_k$ , that is,

$$(1^{\otimes \alpha} \otimes s^{|t_{\alpha+1}|} \otimes \cdots \otimes s^{|t_{\alpha+k-1}|} \otimes s^{-|t_{\alpha+1}|-\cdots-|t_{\alpha+k-1}|-1} \otimes 1^{\otimes \beta}) \cdot (s^{|t_1|} \otimes \cdots \otimes s^{|t_\alpha|} \otimes 1^{\otimes k-1} \otimes s^{|t_{\alpha+1}|+\cdots+|t_{\alpha+k}|+1} \otimes s^{|t_{\alpha+k+1}|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}) + (s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|})(-1)^{1+|t_1|+\cdots+|t_{\alpha}|} (1^{\otimes n-1} \otimes s^{-1}) = 0,$$

in the particular case  $\beta = 0$  it holds as well.

Therefore,  $\mathfrak{FQ}$  is an  $A_\infty\text{-}\mathrm{category.}$ 

Let us establish a property of free  $A_{\infty}$ -categories, which explains why they are called free.

**2.3 Proposition**  $(A_{\infty}$ -functors from a free  $A_{\infty}$ -category). Let  $\mathcal{Q}$  be a differential graded quiver, and let  $\mathcal{A}$  be an  $A_{\infty}$ -category. Let  $f_1 : s\mathcal{Q} \to (s\mathcal{A}, b_1)$  be a chain morphism of differential graded quivers with the underlying mapping of objects  $Ob f : Ob \mathcal{Q} \to Ob \mathcal{A}$ . Suppose given  $\Bbbk$ -quiver morphisms  $f_k : T^k s \mathcal{F} \mathcal{Q} \to s\mathcal{A}$  of degree 0 with the same underlying map Ob f for all k > 1. Then there exists a unique extension of  $f_1$  to a quiver morphism  $f_1 : s\mathcal{F}\mathcal{Q} \to s\mathcal{A}$  such that  $(f_1, f_2, \ldots)$  are components of an  $A_{\infty}$ -functor  $f : \mathcal{F}\mathcal{Q} \to \mathcal{A}$ .

*Proof.* For each n > 1 we have to satisfy the equation

$$b_n f_1 = \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) b_l - \sum_{\alpha + k + \beta = n}^{\alpha + \beta > 0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) f_{\alpha + 1 + \beta} : T^n s \mathcal{F} \Omega \to s \mathcal{A}.$$
(2.3.1)

It is used to define recursively  $f_1$  on  $s\mathcal{FQ}$ . Suppose that  $t_1, \ldots, t_n$  are trees, n > 1, and  $f_1 : s\mathcal{F}_{t_i}\mathcal{Q} \to s\mathcal{A}$  is already defined for all  $1 \leq i \leq n$ . Since

$$b_n = s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n| - |t|} : s\mathcal{F}_{t_1} \mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n} \mathcal{Q} \to s\mathcal{F}_t \mathcal{Q}$$

is invertible for  $t = (t_1 \sqcup \cdots \sqcup t_n) \cdot \mathfrak{t}_n$ , formula (2.3.1) determines  $f_1 : s\mathcal{F}_t \mathcal{Q} \to s\mathcal{A}$  uniquely as

$$f_1 = \left(s\mathcal{F}_t Q \xrightarrow{b_n^{-1}} s\mathcal{F}_{t_1} Q \otimes \cdots \otimes s\mathcal{F}_{t_n} Q \xrightarrow{\sum (f_{i_1} \otimes \cdots \otimes f_{i_l})b_l - \sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta})f_{\alpha+1+\beta}} s\mathcal{A}\right).$$

This proves uniqueness of the extension of  $f_1$ .

Let us prove that the cocategory homomorphism f with so defined components  $(f_1, f_2, ...)$ is an  $A_{\infty}$ -functor. Equations (2.3.1) are satisfied by construction of  $f_1$ . So it remains to prove that  $f_1$  is a chain map. Equation  $f_1b_1 = b_1f_1$  holds on  $s\mathcal{F}_{|}\Omega$  by assumption. We are going to prove by induction on |t| that it holds on  $s\mathcal{F}_{t}\Omega$ . Considering  $t = (t_1 \sqcup \cdots \sqcup t_n) \cdot \mathfrak{t}_n$ , n > 1, we assume that  $f_1b_1 = b_1f_1 : s\mathcal{F}_{t'}\Omega \to s\mathcal{A}$  for all trees t' with |t'| < |t|. To prove that  $f_1b_1 = b_1f_1 : s\mathcal{F}_{t}\Omega \to s\mathcal{A}$  it suffices to show that  $b_nf_1b_1 = b_nb_1f_1$  for all n > 1 due to invertibility of  $b_n$ . Using (2.3.1) and the equation  $b^2 \operatorname{pr}_1 = 0$  we find

$$\begin{split} b_n f_1 b_1 - b_n b_1 f_1 &= \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) b_l b_1 - \sum_{\alpha + k + \beta = n}^{\alpha + \beta > 0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) f_{\alpha + 1 + \beta} b_1 \\ &+ \sum_{\alpha + k + \beta = n}^{\alpha + \beta > 0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) b_{\alpha + 1 + \beta} f_1 \\ &= - \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) \sum_{\gamma + p + \delta = l}^{\gamma + \delta > 0} (1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta}) b_{\gamma + 1 + \delta} \\ &+ \sum_{\alpha + k + \beta = n}^{\alpha + \beta > 0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) \left[ \sum_{j_1 + \dots + j_r = \alpha + 1 + \beta}^{r > 1} (f_{j_1} \otimes \dots \otimes f_{j_r}) b_r \right. \\ &- \left. \sum_{\gamma + p + \delta = \alpha + 1 + \beta} (1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta}) f_{\gamma + 1 + \delta} \right] \\ &= \sum_{r > 1} \left[ \sum_{\alpha + k + \beta = n}^{\alpha + \beta > 0} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) \sum_{j_1 + \dots + j_r = \alpha + 1 + \beta} 1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta} \right] b_r \\ &- \sum_{r > 1} \left[ \sum_{\alpha + k + \beta = n} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) \sum_{\gamma + p + \delta = \alpha + 1 + \beta} 1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta} \right] b_r \\ &- \sum_{r > 1} \left[ \sum_{\alpha + k + \beta = n} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) \sum_{\gamma + p + \delta = \alpha + 1 + \beta} 1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta} \right] f_r. \end{split}$$

Let us show that the expressions in square brackets vanish. The first one is the matrix coefficient  $bf - fb : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \to T^r s\mathcal{A}$ . Indeed, for r > 1 the inequality  $r \leq j_1 + \cdots + j_r = \alpha + 1 + \beta$  automatically implies that  $\alpha + \beta > 0$ , so this condition can be omitted. Using the induction hypothesis one can transform the left hand side of equation

$$\sum_{\alpha+k+\beta=n} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) \sum_{\substack{j_1+\dots+j_r=\alpha+1+\beta\\ j_1+\dots+i_l=n}} f_{j_1} \otimes \dots \otimes f_{j_r}$$
$$= \sum_{\substack{i_1+\dots+i_l=n\\ \gamma+1+\delta=r}} (f_{i_1} \otimes \dots \otimes f_{i_l}) \sum_{\substack{\gamma+p+\delta=l\\ \gamma+1+\delta=r}} 1^{\otimes \gamma} \otimes b_p \otimes 1^{\otimes \delta} : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \dots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \to T^r s\mathcal{A}$$

into the right hand side for all  $n, r \ge 1$ .

The second expression

$$\sum_{\substack{\alpha+k+\beta=n\\\alpha+\beta>0}} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \sum_{\substack{\gamma+p+\delta=\alpha+1+\beta\\\gamma+1+\delta=r}} 1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta}$$
(2.3.2)

is the matrix coefficient

 $(b - b \operatorname{pr}_1) b \operatorname{pr}_r : T^n s \mathcal{FQ} \to T^r s \mathcal{FQ}$ 

of the endomorphism  $(b - b \operatorname{pr}_1)b : Ts \mathfrak{FQ} \to Ts \mathfrak{FQ}$ . However,

$$(b - b\operatorname{pr}_1)b\operatorname{pr}_r = b^2\operatorname{pr}_r - b\operatorname{pr}_1 b\operatorname{pr}_r = -b\operatorname{pr}_1 b\operatorname{pr}_1 \operatorname{pr}_r = 0$$

for r > 1, because  $pr_1 b = pr_1 b pr_1$ . Therefore, (2.3.2) vanishes and equation  $b_n f_1 b_1 = b_n b_1 f_1$  is proven.

Let strict  $A_{\infty}(\mathcal{FQ}, \mathcal{A}) \subset A_{\infty}(\mathcal{FQ}, \mathcal{A})$  be a full  $A_{\infty}$ -subcategory, whose objects are strict  $A_{\infty}$ -functors. Recall that  $\operatorname{Ob} A_1(\mathcal{Q}, \mathcal{A})$  is the set of chain morphisms  $\mathcal{Q} \to \mathcal{A}$  of differential graded quivers.

**2.4 Corollary.** A chain morphism  $f : \mathfrak{Q} \to \mathcal{A}$  admits a unique extension to a strict  $A_{\infty}$ -functor  $\hat{f} : \mathfrak{FQ} \to \mathcal{A}$ . The maps  $f \mapsto \hat{f}$  and

restr : Ob strict 
$$A_{\infty}(\mathfrak{FQ}, \mathcal{A}) \to \operatorname{Ob} A_1(\mathfrak{Q}, \mathcal{A}), \qquad g \mapsto (\operatorname{Ob} g, g_1|_{\mathfrak{s}^0})$$

are inverse to each other.

Indeed, strict  $A_{\infty}$ -functors g are distinguished by conditions  $g_k = 0$  for k > 1.

We may view strict  $A_{\infty}$  as a category, whose objects are  $A_{\infty}$ -categories and morphisms are strict  $A_{\infty}$ -functors. We may also view  $A_1$  as a category consisting of differential graded quivers and their morphisms. There is a functor  $\mathcal{U}$ : strict  $A_{\infty} \to A_1$ ,  $\mathcal{A} \mapsto (\mathcal{A}, m_1)$ , which sends an  $A_{\infty}$ -category to the underlying differential graded k-quiver, forgetting all higher multiplications. The restriction map

restr : strict 
$$A_{\infty}(\mathcal{FQ},\mathcal{A}) \to A_1(\mathcal{Q},\mathcal{UA}), \qquad g \mapsto (\operatorname{Ob} g, g_1|_{s\mathcal{Q}})$$
 (2.4.1)

is functorial in  $\mathcal{A}$ .

**2.5 Corollary.** There is a functor  $\mathcal{F}: A_1 \to \operatorname{strict} A_\infty, \Omega \mapsto \mathcal{F}\Omega$ , left adjoint to  $\mathcal{U}$ .

**2.6.** Explicit formula for the constructed strict  $A_{\infty}$ -functor. Let us obtain a more explicit formula for  $\widehat{f}_1|_{s\mathcal{F}_t Q}$ . We define  $\widehat{f}_1$  for  $t = (t_1 \sqcup \cdots \sqcup t_n) \cdot \mathfrak{t}_n$  recursively by a commutative diagram

Notice that the top map is invertible. Here  $n, t_1, \ldots, t_n$  are uniquely determined by decomposition (2.2.1) of t.

Let  $(t, \leq)$  be a linearly ordered tree with the underlying given plane rooted tree t. Decompose  $(t, \leq)$  into a vertical composition of forests as in (1.7.1). Then the following diagram commutes

$$\begin{array}{c} s \mathbb{Q}^{\otimes n} \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} s \mathbb{Q}^{\otimes \alpha_1} \otimes s \mathbb{F}_{\mathbf{t}_{k_1}} \mathbb{Q} \otimes s \mathbb{Q}^{\otimes \beta_1} \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} \cdots \xrightarrow{b_{k_N}} s \mathbb{F}_t \mathbb{Q} \\ f_1^{\otimes n} \downarrow & & \\ s \mathbb{A}^{\otimes n} \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} s \mathbb{A}^{\otimes \alpha_1} \otimes s \mathbb{A} \otimes s \mathbb{A}^{\otimes \beta_1} \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} \cdots \xrightarrow{b_{k_N}} s \mathbb{A} \end{array}$$

The upper row consists of invertible maps. One can prove by induction that the composition of maps in the upper row equals  $\pm s^{-|t|}$ . When  $(t, \leq) = t_{<}$  is the linearly ordered tree, canonically associated with t, then the composition of maps in the upper row equals  $s^{-|t|}$ . This is also proved by induction: if t is presented as  $t = (t_1 \sqcup \cdots \sqcup t_k) \cdot \mathfrak{t}_k$ , then the composition of maps in the upper row is

$$(1^{\otimes k-1} \otimes s^{-|t_k|})(1^{\otimes k-2} \otimes s^{-|t_{k-1}|} \otimes 1) \dots (s^{-|t_1|} \otimes 1^{\otimes k-1})(s^{|t_1|} \otimes \dots \otimes s^{|t_{k-1}|} \otimes s^{|t_k|-|t|}) = 1^{\otimes k-1} \otimes s^{-|t|} = s^{-|t|}$$

Therefore, for an arbitrary tree  $t\in \mathbb{T}^n_{\geqslant 2}$  the map  $\left.\widehat{f_1}\right|_{s\mathcal{F}_t\Omega}$  is

$$\widehat{f}_1 = \left(s\mathcal{F}_t \mathcal{Q} \xrightarrow{s^{|t|}} s\mathcal{Q}^{\otimes n} \xrightarrow{f_1^{\otimes n}} s\mathcal{A}^{\otimes n} \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} s\mathcal{A}^{\otimes \alpha_1 + 1 + \beta_1} \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} \dots \xrightarrow{b_{k_N}} s\mathcal{A}\right),$$

where the factors correspond to decomposition (1.7.1) of  $t_{<}$ .

**2.7.** Transformations between functors from a free  $A_{\infty}$ -category. Let  $\mathcal{Q}$  be a differential graded quiver, and let  $\mathcal{A}$  be an  $A_{\infty}$ -category. Then  $A_1(\mathcal{Q}, \mathcal{A})$  is an  $A_{\infty}$ -category as well. The differential graded quiver  $(sA_1(\mathcal{Q}, \mathcal{A}), B_1)$  is described as follows. Objects are chain quiver maps  $\phi : (s\mathcal{Q}, b_1) \to (s\mathcal{A}, b_1)$ , the graded k-module of morphisms  $\phi \to \psi$  is the product of graded k-modules

$$sA_1(\mathcal{Q},\mathcal{A})(\phi,\psi) = \prod_{X \in Ob \ \mathcal{Q}} s\mathcal{A}(X\phi, X\psi) \times \prod_{X,Y \in Ob \ \mathcal{Q}} \mathsf{C}(s\mathcal{Q}(X,Y), s\mathcal{A}(X\phi, Y\psi)), \quad r = (r_0, r_1)$$

The differential  $B_1$  is given by

$$(rB_1)_0 = r_0 b_1,$$
  

$$(rB_1)_1 = r_1 b_1 + (\phi_1 \otimes r_0) b_2 + (r_0 \otimes \psi_1) b_2 - (-)^r b_1 r_1.$$
(2.7.1)

Restrictions  $\phi, \psi : \mathfrak{Q} \to \mathcal{A}$  of arbitrary  $A_{\infty}$ -functors  $\phi, \psi : \mathfrak{FQ} \to \mathcal{A}$  to  $\mathfrak{Q}$  are  $A_1$ -functors (chain quiver maps).

**2.8 Proposition.** Let  $\phi, \psi : \mathfrak{FQ} \to \mathcal{A}$  be  $A_{\infty}$ -functors. For an arbitrary complex P of  $\Bbbk$ -modules chain maps  $u : P \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  are in bijection with the following data:  $(u', u_k)_{k>1}$ 

- 1. a chain map  $u': P \to sA_1(Q, \mathcal{A})(\phi, \psi)$ ,
- 2. k-linear maps

$$u_k: P \to \prod_{X,Y \in Ob \, \mathcal{Q}} \mathsf{C}\big((s\mathfrak{FQ})^{\otimes k}(X,Y), s\mathcal{A}(X\phi, Y\psi)\big)$$

of degree 0 for all k > 1.

The bijection maps u to  $(u', u_k)_{k>1}$ , where  $u_k = u \cdot \operatorname{pr}_k$  and

$$u' = \left(P \xrightarrow{u} sA_{\infty}(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_{1}(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_{1}(\mathcal{Q}, \mathcal{A})(\phi, \psi)\right). \quad (2.8.1)$$

The inverse bijection can be recovered from the recurrent formula

$$(-)^{p}b_{k}^{\mathfrak{FQ}}(pu_{1}) = -(pd)u_{k} + \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_{q} \otimes \psi_{c\beta})b_{\alpha+1+\beta}^{\mathcal{A}} - (-)^{p}\sum_{\alpha+q+\beta=k}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_{q}^{\mathfrak{FQ}} \otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta}) : (s\mathfrak{FQ})^{\otimes k} \to s\mathcal{A},$$

where k > 1,  $p \in P$ , and  $\phi_{a\alpha}$ ,  $\psi_{c\beta}$  are matrix elements of  $\phi$ ,  $\psi$ .

*Proof.* Since the k-module of  $(\phi, \psi)$ -coderivations  $sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  is a product, k-linear maps  $u : P \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  of degree 0 are in bijection with sequences of k-linear maps  $(u_k)_{k\geq 0}$  of degree 0:

$$\begin{split} u_0 &: P \to \prod_{X \in \operatorname{Ob} \Omega} s\mathcal{A}(X\phi, X\psi), \qquad \qquad p \mapsto pu_0, \\ u_k &: P \to \prod_{X,Y \in \operatorname{Ob} \Omega} \mathsf{C}\big((s\mathcal{FQ})^{\otimes k}(X,Y), s\mathcal{A}(X\phi, Y\psi)\big), \qquad p \mapsto pu_k, \end{split}$$

for  $k \ge 1$ . The complex  $\Phi_0 = (sA_\infty(\mathfrak{FQ}, \mathcal{A})(\phi, \psi), B_1)$  admits a filtration by subcomplexes

$$\Phi_n = 0 \times \cdots \times 0 \times \prod_{k=n}^{\infty} \prod_{X,Y \in Ob \ \Omega} \mathsf{C}\big((s\mathfrak{F}\mathfrak{Q})^{\otimes k}(X,Y), s\mathcal{A}(X\phi,Y\psi)\big).$$

In particular,  $\Phi_2$  is a subcomplex, and

$$\Phi_0/\Phi_2 = \prod_{X \in \operatorname{Ob} \mathcal{Q}} s\mathcal{A}(X\phi, X\psi) \times \prod_{X,Y \in \operatorname{Ob} \mathcal{Q}} \mathsf{C}\big(s\mathcal{FQ}(X,Y), s\mathcal{A}(X\phi, Y\psi)\big)$$

is the quotient complex with differential (2.7.1). Since  $(s\mathcal{FQ}, b_1)$  splits into a direct sum of two subcomplexes  $s\mathcal{Q} \oplus (\bigoplus_{|t|>0} s\mathcal{F}_t \mathcal{Q})$ , the complex  $\Phi_0/\Phi_2$  has a subcomplex

$$\left(0 \times \prod_{X,Y \in Ob \ \Omega} \mathsf{C}\left(\oplus_{|t|>0} s \mathcal{F}_t \Omega(X,Y), s \mathcal{A}(X\phi,Y\psi)\right), [\_, b_1]\right).$$

The corresponding quotient complex is  $sA_1(\mathcal{Q},\mathcal{A})(\phi,\psi)$ . The resulting quotient map restr<sub>1</sub> :  $sA_{\infty}(\mathcal{FQ},\mathcal{A})(\phi,\psi) \rightarrow sA_1(\mathcal{Q},\mathcal{A})(\phi,\psi)$  is the restriction map. Denoting  $u' = u \cdot \text{restr}_1$ , we get the discussed assignment  $u \mapsto (u', u_n)_{n>1}$ . The claim is that if u is a chain map, then the missing part

$$u_1'': P \to \prod_{X,Y \in \operatorname{Ob} \mathcal{Q}} \mathsf{C}\big(\oplus_{|t|>0} s\mathfrak{F}_t \mathcal{Q}(X,Y), s\mathcal{A}(X\phi, Y\psi)\big),$$

of  $u_1 = u'_1 \times u''_1$  is recovered in a unique way.

Let us prove that the map  $u \mapsto (u', u_n)_{n>1}$  is injective. The chain map u satisfies  $pdu = puB_1$  for all  $p \in P$ . That is,  $pdu_k = (puB_1)_k$  for all  $k \ge 0$ . Since  $puB_1 = (pu)b^{\mathcal{A}} - (-)^p b^{\mathcal{FQ}}(pu)$ , these conditions can be rewritten as

$$pdu_k = \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta} - (-)^p \sum_{\alpha+q+\beta=k} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta}) (pu_{\alpha+1+\beta}), \quad (2.8.2)$$

where  $\phi_{a\alpha}: T^a s \mathcal{FQ}(X, Y) \to T^\alpha s \mathcal{A}(X\phi, Y\phi)$  are matrix elements of  $\phi$ , and  $\psi_{c\beta}$  are matrix elements of  $\psi$ . The same formula can be rewritten as

$$(-)^{p}b_{k}^{\mathcal{F}\mathcal{Q}}(pu_{1}) = -(pd)u_{k} + \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_{q} \otimes \psi_{c\beta})b_{\alpha+1+\beta}^{\mathcal{A}}$$
$$- (-)^{p}\sum_{\alpha+q+\beta=k}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_{q}^{\mathcal{F}\mathcal{Q}} \otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta}) : s\mathcal{F}_{t_{1}}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_{k}}\mathcal{Q} \to s\mathcal{A}. \quad (2.8.3)$$

When k > 1, the map  $b_k^{\mathcal{F}\mathcal{Q}} : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_k}\mathcal{Q} \to s\mathcal{F}_t\mathcal{Q}, t = (t_1 \sqcup \cdots \sqcup t_k)\mathfrak{t}_k$  is invertible, thus,  $pu_1 : s\mathcal{F}_t\mathcal{Q} \to s\mathcal{A}$  in the left hand side is determined in a unique way by  $u_0, u_n$  for n > 1 and by  $pu_1 : s\mathcal{F}_{t_i} \mathcal{Q} \to s\mathcal{A}, \ 1 \leq i \leq k$ , occurring in the right hand side. Since the restriction  $u'_1$  of  $u_1$  to  $s\mathcal{F}_{|}\mathcal{Q} = s\mathcal{Q}$  is known by 1), the map  $u''_1$  is recursively recovered from  $(u_0, u'_1, u_n)_{n>1}$ .

Let us prove that the map  $u \mapsto (u', u_n)_{n>1}$  is surjective. Given  $(u_0, u'_1, u_n)_{n>1}$  we define maps  $u''_1$  of degree 0 recursively by (2.8.3). This implies equation (2.8.2) for k > 1. For k = 0 this equation in the form  $pdu_0 = pu_0b_1$  holds due to condition 1). It remains to prove equation (2.8.2) for k = 1:

$$(pd)u_{1} = (pu_{1})b_{1}^{\mathcal{A}} + (\phi_{1} \otimes pu_{0})b_{2}^{\mathcal{A}} + (pu_{0} \otimes \psi_{1})b_{2}^{\mathcal{A}} - (-)^{p}b_{1}(pu_{1}) :$$
  
$$s\mathcal{F}_{t}\mathcal{Q}(X,Y) \to s\mathcal{A}(X\phi,Y\psi) \quad (2.8.4)$$

for all trees  $t \in \mathcal{T}_{\geq 2}$ . For t = | it holds due to assumption 1). Let N > 1 be an integer. Assume that equation (2.8.4) holds for all trees  $t \in \mathcal{T}_{\geq 2}$  with the number of input leaves in(t) < N. Let  $t \in \mathcal{T}_{\geq 2}^N$  be a tree (with in(t) = N). Then  $t = (t_1 \sqcup \cdots \sqcup t_k) \mathfrak{t}_k$  for some k > 1 and some trees  $t_i \in \mathcal{T}_{\geq 2}$ ,  $in(t_i) < N$ . For such t equation (2.8.4) is equivalent to

$$(-)^{p}b_{k}(pd)u_{1} = (-)^{p}b_{k}(pu_{1})b_{1}^{\mathcal{A}} + (-)^{p}b_{k}(\phi_{1} \otimes pu_{0})b_{2}^{\mathcal{A}} + (-)^{p}b_{k}(pu_{0} \otimes \psi_{1})b_{2}^{\mathcal{A}} + \sum_{\gamma+j+\delta=k}^{\gamma+\delta>0} (1^{\otimes\gamma} \otimes b_{j} \otimes 1^{\otimes\delta})b_{\gamma+1+\delta}(pu_{1}) : s\mathcal{F}_{t_{1}}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_{k}}\mathcal{Q} \to s\mathcal{A}.$$

Substituting definition (2.8.3) of  $u_1$  we turn the above equation into an identity

$$-\sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pdu_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}$$
(2.8.5)

$$-(-)^{p}\sum_{\alpha+q+\beta=k}^{\alpha+\beta>0}(1^{\otimes\alpha}\otimes b_{q}\otimes 1^{\otimes\beta})(pdu_{\alpha+1+\beta})$$
(2.8.6)

$$= -(pdu_k)b_1 \tag{2.8.7}$$

$$+\sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta} b_1 - (-)^p \sum_{\alpha+\beta>0}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta}) (pu_{\alpha+1+\beta}) b_1$$
(2.8.8)

$$-(-)^{p}\sum_{\alpha+q+\beta=k}(1^{\otimes\alpha}\otimes b_{q}\otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta})b_{1}$$
(2.8.8)

$$+ (-)^{p} b_{k}(\phi_{1} \otimes p u_{0}) b_{2} + (-)^{p} b_{k}(p u_{0} \otimes \psi_{1}) b_{2}$$

$$(2.8.9)$$

$$+ (-)^{p} \sum_{\gamma+j+\delta=k}^{\gamma+\delta>0} (1^{\otimes\gamma} \otimes b_{j} \otimes 1^{\otimes\delta}) \bigg[ -p du_{\gamma+1+\delta}$$

$$(2.8.10)$$

$$+\sum_{a+q+c=\gamma+1+\delta}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}$$
(2.8.11)

$$-(-)^{p}\sum_{\alpha+q+\beta=\gamma+1+\delta}^{\alpha+\beta>0}(1^{\otimes\alpha}\otimes b_{q}\otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta})\bigg],\qquad(2.8.12)$$

whose validity we are going to prove now. First of all, terms (2.8.6) and (2.8.10) cancel each other. Term (2.8.12) vanishes because for an arbitrary integer g the sum

$$\sum_{\substack{\gamma+j+\delta=k\\\alpha+q+\beta=\gamma+1+\delta}}^{\alpha+1+\beta=g} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})(1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta})$$
(2.8.13)

is the matrix coefficient  $b^2 = 0$ :  $T^k s \mathcal{FQ} \to T^g s \mathcal{FQ}$ , thus, it vanishes. Notice that condition  $\alpha + \beta > 0$  in (2.8.12) automatically implies  $\gamma + \delta > 0$ . Furthermore, term (2.8.7) cancels one of the terms of sum (2.8.5). In the remaining terms of (2.8.5) we may use the induction assumptions and replace  $pdu_q$  with the right hand side of (2.8.2). We also absorb terms (2.8.9) into sum (2.8.11), allowing  $\gamma = \delta = 0$  in it and allowing simultaneously  $\alpha = \beta = 0$  in (2.8.8) to compensate the missing term  $b_k(pu_1)b_1$ :

$$-\sum_{a+q+c=k}^{\alpha+\beta>0}\sum_{\substack{e+j+f=q\\\alpha+\beta>0}}^{\gamma,\delta} \left[\phi_{a\alpha}\otimes(\phi_{e\gamma}\otimes pu_{j}\otimes\psi_{f\delta})b^{\mathcal{A}}_{\gamma+1+\delta}\otimes\psi_{c\beta}\right]b^{\mathcal{A}}_{\alpha+1+\beta}$$
(2.8.14)

$$+ (-)^{p} \sum_{a+q+c=k}^{\alpha+\beta>0} \sum_{\gamma+j+\delta=q} \left[ \phi_{a\alpha} \otimes (1^{\otimes \gamma} \otimes b_{j} \otimes 1^{\otimes \delta})(pu_{\gamma+1+\delta}) \otimes \psi_{c\beta} \right] b_{\alpha+1+\beta}^{\mathcal{A}}$$
(2.8.15)

$$=\sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta} b^{\mathcal{A}}_1$$
(2.8.16)

$$-(-)^{p} \sum_{\alpha+q+\beta=k} (1^{\otimes \alpha} \otimes b_{q} \otimes 1^{\otimes \beta}) (pu_{\alpha+1+\beta}) b_{1}^{\mathcal{A}}$$

$$(2.8.17)$$

$$+ (-)^p \sum_{\gamma+j+\delta=k} \sum_{a+q+c=\gamma+1+\delta}^{\alpha,\beta} (1^{\otimes \gamma} \otimes b_j \otimes 1^{\otimes \delta}) (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}.$$

Recall that  $\phi_{a0}$  vanish for all *a* except a = 0. Therefore, we may absorb term (2.8.16) into sum (2.8.14) and term (2.8.17) into sum (2.8.15), allowing terms with  $\alpha = \beta = 0$  in them. Denote  $r = pu \in sA_{\infty}(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ . The proposition follows immediately form the following

**2.9 Lemma.** For all  $r \in sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  and all  $k \ge 0$  we have

$$-\sum_{a+q+c=k}^{\alpha,\beta}\sum_{e+j+f=q}^{\gamma,\delta} \left[\phi_{a\alpha}\otimes(\phi_{e\gamma}\otimes r_{j}\otimes\psi_{f\delta})b_{\gamma+1+\delta}^{\mathcal{A}}\otimes\psi_{c\beta}\right]b_{\alpha+1+\beta}^{\mathcal{A}}$$
$$+(-)^{r}\sum_{a+q+c=k}^{\alpha,\beta}\sum_{\gamma+j+\delta=q} \left[\phi_{a\alpha}\otimes(1^{\otimes\gamma}\otimes b_{j}\otimes1^{\otimes\delta})r_{\gamma+1+\delta}\otimes\psi_{c\beta}\right]b_{\alpha+1+\beta}^{\mathcal{A}}$$

$$= (-)^r \sum_{\gamma+j+\delta=k} \sum_{a+q+c=\gamma+1+\delta}^{\alpha,\beta} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta}) (\phi_{a\alpha} \otimes r_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}.$$
(2.9.1)

*Proof.* Sum (2.9.1) is split into three sums accordingly to output of  $b_j$  being an input of  $\phi_{a\alpha}$  or  $r_q$  or  $\psi_{c\beta}$ :

$$-\sum_{a+e+j+f+c=k}^{\alpha,\beta,\gamma,\delta} (\phi_{a\alpha} \otimes \phi_{e\gamma} \otimes r_j \otimes \psi_{f\delta} \otimes \psi_{c\beta}) (1^{\otimes \alpha} \otimes b_{\gamma+1+\delta}^{\mathcal{A}} \otimes 1^{\otimes \beta}) b_{\alpha+1+\beta}^{\mathcal{A}}$$
(2.9.2)

$$+ (-)^{r} \sum_{\substack{a+\gamma+j+\delta+c=k\\a,\alpha,\beta}}^{\alpha,\beta} (1^{\otimes a+\gamma} \otimes b_{j} \otimes 1^{\otimes \delta+c}) (\phi_{a\alpha} \otimes r_{\gamma+1+\delta} \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}$$
(2.9.3)

$$= (-)^r \sum_{x+q+c=k}^{a,\alpha,\beta} (b_{xa}\phi_{a\alpha} \otimes r_q \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}$$
(2.9.4)

$$+ (-)^{r} \sum_{a+y+c=k}^{\alpha,q,\beta} (\phi_{a\alpha} \otimes b_{yq} r_{q} \otimes \psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}$$

$$(2.9.5)$$

$$+\sum_{a+q+z=k}^{\alpha,\beta,c} (\phi_{a\alpha} \otimes r_q \otimes b_{zc}\psi_{c\beta}) b^{\mathcal{A}}_{\alpha+1+\beta}.$$
(2.9.6)

Here  $b_{xa}: T^x s \mathcal{FQ} \to T^a s \mathcal{FQ}$  is a matrix element of  $b^{\mathcal{FQ}}$ . Terms (2.9.3) and (2.9.5) cancel each other. We shall use  $A_{\infty}$ -functor identities  $b\phi = \phi b$ ,  $b\psi = \psi b$  for terms (2.9.4) and (2.9.6). Being a cocategory homomorphism,  $\phi$  satisfies the identity

$$\sum_{a+e=h} \phi_{a\alpha} \otimes \phi_{e\gamma} = \left[ \Delta(\phi \otimes \phi) \right]_{h;\alpha,\gamma} = \phi_{h,\alpha+\gamma} \Delta_{\alpha+\gamma;\alpha,\gamma}$$

for all non-negative integers h, where  $\Delta$  is the cut comultiplication. Similarly for  $\psi$ . Using this identity in (2.9.2) we get the equation to verify:

$$-\sum_{x+q+z=k}^{v,w} (\phi_{xv} \otimes r_q \otimes \psi_{zw}) \sum_{\alpha+y+\beta=v+1+w}^{\alpha \leqslant v,\beta \leqslant w} (1^{\otimes \alpha} \otimes b_y^{\mathcal{A}} \otimes 1^{\otimes \beta}) b_{\alpha+1+\beta}^{\mathcal{A}}$$
$$=\sum_{x+q+z=k}^{v,w,\alpha} (\phi_{xv} \otimes r_q \otimes \psi_{zw}) (b_{v\alpha}^{\mathcal{A}} \otimes 1^{\otimes 1+w}) b_{\alpha+1+w}^{\mathcal{A}}$$
$$+\sum_{x+q+z=k}^{v,w,\beta} (\phi_{xv} \otimes r_q \otimes \psi_{zw}) (1^{\otimes v+1} \otimes b_{w\beta}^{\mathcal{A}}) b_{v+1+\beta}^{\mathcal{A}}.$$

It follows from the identity  $b^2 \operatorname{pr}_1 = 0 : T^{v+1+w} s \mathcal{A} \to s \mathcal{A}$  valid for arbitrary non-negative integers v, w, which we may rewrite like this:

$$\sum_{\alpha+y+\beta=v+1+w}^{\alpha\leqslant v,\beta\leqslant w} (1^{\otimes\alpha}\otimes b_y^{\mathcal{A}}\otimes 1^{\otimes\beta})b_{\alpha+1+\beta}^{\mathcal{A}} + \sum_{\alpha} (b_{v\alpha}^{\mathcal{A}}\otimes 1^{\otimes1+w})b_{\alpha+1+w}^{\mathcal{A}} + \sum_{\beta} (1^{\otimes v+1}\otimes b_{w\beta}^{\mathcal{A}})b_{v+1+\beta}^{\mathcal{A}} = 0.$$

So the lemma is proved.

The proposition follows.

Let us consider now the question, when the discussed chain map is null-homotopic.

**2.10 Corollary.** Let  $\phi, \psi : \mathfrak{FQ} \to \mathcal{A}$  be  $A_{\infty}$ -functors. Let P be a complex of k-modules. Let  $u : P \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  be a chain map. The set (possibly empty) of homotopies  $h : P \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$ , deg h = -1, such that  $u = dh + hB_1$  is in bijection with the set of data  $(h', h_k)_{k>1}$ , consisting of

- 1. a homotopy  $h': P \to sA_1(\Omega, \mathcal{A})(\phi, \psi)$ , deg h' = -1, such that  $dh' + h'B_1 = u'$ , where u' is given by (2.8.1);
- 2. k-linear maps

$$h_k: P \to \prod_{X,Y \in Ob \ \Omega} \mathsf{C}((s\mathfrak{FQ})^{\otimes k}(X,Y), s\mathcal{A}(X\phi, Y\psi))$$

of degree -1 for all k > 1.

The bijection maps h to  $(h', h_k)_{k>1}$ , where  $h_k = h \cdot pr_k$  and

$$h' = \left(P \xrightarrow{h} sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_{1}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_{1}(\mathfrak{Q}, \mathcal{A})(\phi, \psi)\right).$$

The inverse bijection can be recovered from the recurrent formula

$$(-)^{p}b_{k}(ph_{1}) = pu_{k} - (pd)h_{k} - \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes ph_{q} \otimes \psi_{c\beta})b_{\alpha+1+\beta} - (-)^{p} \sum_{a+q+c=k}^{a+c>0} (1^{\otimes a} \otimes b_{q} \otimes 1^{\otimes c})(ph_{a+1+c}) : (s\mathcal{FQ})^{\otimes k} \to s\mathcal{A},$$

where k > 1,  $p \in P$ , and  $\phi_{a\alpha}$ ,  $\psi_{c\beta}$  are matrix elements of  $\phi$ ,  $\psi$ .

Proof. We shall apply Proposition 2.8 to the complex Cone(id :  $P \to P$ ) instead of P. The graded k-module Cone(id<sub>P</sub>) =  $P \oplus P[1]$  is equipped with the differential  $(q, ps)d = (qd + p, -pds), p, q \in P$ . The chain maps  $\overline{u}$  : Cone(id<sub>P</sub>)  $\to C$  to an arbitrary complex C are in bijection with pairs  $(u : P \to C, h : P \to C)$ , where u = dh + hd and deg h = -1. The pair  $(u, h) = (in_1 \overline{u}, s in_2 \overline{u})$  is assigned to  $\overline{u}$ , and the map  $\overline{u} : P \oplus P[1] \to C$ ,  $(q, ps) \mapsto qu + ph$  is assigned to a pair (u, h). Indeed,  $\overline{u}$  being chain map is equivalent to

$$(q, ps)d\overline{u} = qdu + pu - pdh = qud + phd = (q, ps)\overline{u}d,$$

that is, to conditions du = ud, u = dh + hd.

Thus, for a fixed chain map  $u: P \to C$  the set of homotopies  $h: P \to C$ , such that u = dh + hd, is in bijection with the set of chain maps  $\overline{u}$ : Cone(id<sub>P</sub>)  $\to C$  such that in<sub>1</sub> $\overline{u} = u: P \to C$ . Applying this statement to  $u: P \to C = sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  we find by Proposition 2.8 that the set of homotopies  $h: P \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(\phi, \psi)$  such that  $u = dh + hB_1$  is in bijection with the set of data  $(\overline{u}', \overline{u}_k)_{k>1}$ , such that

$$\overline{u}': \operatorname{Cone}(\operatorname{id}_P) \to sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi) \quad \text{is a chain map,} \quad \operatorname{in}_1 \overline{u}' = u', \\ \overline{u}_k: \operatorname{Cone}(\operatorname{id}_P) \to \prod_{X, Y \in \operatorname{Ob} \mathcal{Q}} \mathsf{C}\big((s\mathfrak{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi)\big), \quad \deg \overline{u}_k = 0, \quad \operatorname{in}_1 \overline{u}_k = u_k,$$

therefore, in bijection with the set of data  $(h', h_k)_{k>1} = (s \operatorname{in}_2 \overline{u}', s \operatorname{in}_2 \overline{u}_k)_{k>1}$ , as stated in corollary.

**2.11.** Restriction as an  $A_{\infty}$ -functor. Let Q be a ( $\mathscr{U}$ -small) differential graded k-quiver. Denote by  $\mathcal{F}Q$  the free  $A_{\infty}$ -category generated by Q. Let  $\mathcal{A}$  be a ( $\mathscr{U}$ -small) unital  $A_{\infty}$ -category. There is the restriction strict  $A_{\infty}$ -functor

restr : 
$$A_{\infty}(\mathfrak{FQ},\mathcal{A}) \to A_1(\mathbb{Q},\mathcal{A}), \qquad (f:\mathfrak{FQ}\to\mathcal{A})\mapsto (\overline{f}=(f_1|_{\mathbb{Q}}):\mathbb{Q}\to\mathcal{A}).$$

In fact, it is the composition of two strict  $A_{\infty}$ -functors:  $A_{\infty}(\mathcal{FQ},\mathcal{A}) \xrightarrow{\operatorname{restr}_{\infty,1}} A_1(\mathcal{FQ},\mathcal{A}) \to A_1(\mathcal{Q},\mathcal{A})$ , where the second comes from the full embedding  $\mathcal{Q} \hookrightarrow \mathcal{FQ}$ . Its first component is

$$\operatorname{restr}_{1} : sA_{\infty}(\mathcal{FQ},\mathcal{A})(f,g) \to sA_{1}(\mathcal{Q},\mathcal{A})(\overline{f},\overline{g}), \qquad (2.11.1)$$
$$r = (r_{0},r_{1},\ldots,r_{n},\ldots) \mapsto (r_{0},r_{1}|_{\mathcal{Q}}) = \overline{r}.$$

**2.12 Theorem.** The  $A_{\infty}$ -functor restr :  $A_{\infty}(\mathfrak{FQ}, \mathcal{A}) \to A_1(\mathbb{Q}, \mathcal{A})$  is an equivalence.

*Proof.* Let us prove that restriction map (2.11.1) is homotopy invertible. We construct a chain map going in the opposite direction

$$u: sA_1(\mathfrak{Q}, \mathcal{A})(\overline{f}, \overline{g}) \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(f, g)$$

via Proposition 2.8 taking  $P = sA_1(\mathfrak{Q}, \mathcal{A})(\overline{f}, \overline{g})$ . We choose

$$u': sA_1(\mathfrak{Q}, \mathcal{A})(\overline{f}, \overline{g}) \to sA_1(\mathfrak{Q}, \mathcal{A})(\overline{f}, \overline{g})$$

to be the identity map and  $u_k = 0$  for k > 1. Therefore,

$$u \cdot \operatorname{restr}_1 = u' = \operatorname{id}_{sA_1(Q,\mathcal{A})(\overline{f},\overline{g})}.$$

Denote

$$v = \mathrm{id}_{sA_{\infty}(\mathfrak{FQ},\mathcal{A})(f,g)} - \left[ sA_{\infty}(\mathfrak{FQ},\mathcal{A})(f,g) \xrightarrow{\mathrm{restr}_1} sA_1(\mathcal{Q},\mathcal{A})(\overline{f},\overline{g}) \xrightarrow{u} sA_{\infty}(\mathfrak{FQ},\mathcal{A})(f,g) \right].$$

Let us prove that v is null-homotopic via Corollary 2.10, taking  $P = sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(f, g)$ . A homotopy  $h : sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(f, g) \to sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(f, g)$ , deg h = -1, such that  $v = B_1h + hB_1$  is specified by  $h' = 0 : sA_{\infty}(\mathfrak{FQ}, \mathcal{A})(f, g) \to sA_1(\mathfrak{Q}, \mathcal{A})(\overline{f}, \overline{g})$  and  $h_k = 0$  for k > 1. Indeed,

$$v' = v \cdot \operatorname{restr}_1 = \operatorname{restr}_1 - \operatorname{restr}_1 \cdot u \cdot \operatorname{restr}_1 = \operatorname{restr}_1 - \operatorname{restr}_1 = 0,$$

so  $v' = B_1 h' + h' B_1$  and condition 1 of Corollary 2.10 is satisfied<sup>2</sup>. Therefore, u is homotopy inverse to restr<sub>1</sub>.

Let  $\mathbf{i}^{\mathcal{A}}$  be a unit transformation of the unital  $A_{\infty}$ -category  $\mathcal{A}$ . Then  $A_1(\mathcal{Q}, \mathcal{A})$  is a unital  $A_{\infty}$ -category with the unit transformation  $(1 \otimes \mathbf{i}^{\mathcal{A}})M$  (cf. [Lyu03, Proposition 7.7]). The unit element for an object  $\phi \in \text{Ob} A_1(\mathcal{Q}, \mathcal{A})$  is  ${}_{\phi}\mathbf{i}_0^{A_1(\mathcal{Q}, \mathcal{A})} : \mathbb{k} \to sA_1(\mathcal{Q}, \mathcal{A}), 1 \mapsto \phi\mathbf{i}^{\mathcal{A}}$ . The  $A_{\infty}$ -category  $A_{\infty}(\mathcal{FQ}, \mathcal{A})$  is also unital. To establish equivalence of these two  $A_{\infty}$ -categories via restr :  $A_{\infty}(\mathcal{FQ}, \mathcal{A}) \to A_1(\mathcal{Q}, \mathcal{A})$  we verify the conditions of Theorem 8.8 from [Lyu03].

Consider the mapping  $\operatorname{Ob} A_1(\mathfrak{Q}, \mathcal{A}) \to \operatorname{Ob} A_\infty(\mathfrak{FQ}, \mathcal{A}), \phi \mapsto \widehat{\phi}$ , which extends a given chain map to a strict  $A_\infty$ -functor, constructed in Corollary 2.4. Clearly,  $\overline{\widehat{\phi}} = \phi$ . It remains to give two mutually inverse cycles, which we choose as follows:

$${}_{\phi}r_{0}: \mathbb{k} \to sA_{1}(\mathbb{Q}, \mathcal{A})(\phi, \overline{\widehat{\phi}}), \qquad 1 \mapsto \phi \mathbf{i}^{\mathcal{A}},$$

$${}_{\phi}p_{0}: \mathbb{k} \to sA_{1}(\mathbb{Q}, \mathcal{A})(\overline{\widehat{\phi}}, \phi), \qquad 1 \mapsto \phi \mathbf{i}^{\mathcal{A}}.$$

Clearly,  $_{\phi}r_0B_1 = 0$ ,  $_{\phi}p_0B_1 = 0$ ,

$$({}_{\phi}r_0 \otimes {}_{\phi}p_0)B_2 - {}_{\phi}\mathbf{i}_0^{A_1(\Omega,\mathcal{A})} : 1 \mapsto (\phi\mathbf{i}^{\mathcal{A}} \otimes \phi\mathbf{i}^{\mathcal{A}})B_2 - \phi\mathbf{i}^{\mathcal{A}} \in \operatorname{Im} B_1, ({}_{\phi}p_0 \otimes {}_{\phi}r_0)B_2 - {}_{\phi}\mathbf{i}_0^{A_1(\Omega,\mathcal{A})} : 1 \mapsto (\phi\mathbf{i}^{\mathcal{A}} \otimes \phi\mathbf{i}^{\mathcal{A}})B_2 - \phi\mathbf{i}^{\mathcal{A}} \in \operatorname{Im} B_1.$$

Therefore, all assumptions of Theorem 8.8 [Lyu03] are satisfied. Thus, restr :  $A_{\infty}(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$  is an  $A_{\infty}$ -equivalence.

**2.13 Corollary.** Every  $A_{\infty}$ -functor  $f : \mathfrak{FQ} \to \mathcal{A}$  is isomorphic to the strict  $A_{\infty}$ -functor  $\widehat{f} : \mathfrak{FQ} \to \mathcal{A}$ .

Proof. Note that  $\overline{f} = \overline{\overline{f}}$ . The  $A_1$ -transformation  $\overline{f}\mathbf{i}^{\mathcal{A}} : \overline{f} \to \overline{\overline{f}} : \mathcal{Q} \to \mathcal{A}$  with the components  $({}_{Xf}\mathbf{i}_0^{\mathcal{A}}, \overline{f}_1\mathbf{i}_1^{\mathcal{A}})$  is natural. It is mapped by u into a natural  $A_{\infty}$ -transformation  $(\overline{f}\mathbf{i}^{\mathcal{A}})u : f \to \overline{\overline{f}} : \mathcal{F}\mathcal{Q} \to \mathcal{A}$ . Its zero component  ${}_{Xf}\mathbf{i}_0^{\mathcal{A}}$  is invertible, therefore  $(\overline{f}\mathbf{i}^{\mathcal{A}})u$  is invertible by [Lyu03, Proposition 7.15].

<sup>&</sup>lt;sup>2</sup>By the way, the only non-vanishing component of h is  $h_1$ .

## 3. Representable 2-functors $A^u_{\infty} \to A^u_{\infty}$

Recall that unital  $A_{\infty}$ -categories, unital  $A_{\infty}$ -functors and equivalence classes of natural  $A_{\infty}$ -transformations form a 2-category [Lyu03]. In order to distinguish between the  $A_{\infty}$ -category  $A_{\infty}^{u}(\mathcal{C}, \mathcal{D})$  and the ordinary category, whose morphisms are equivalence classes of natural  $A_{\infty}$ -transformations, we denote the latter by

$$\overline{A^u_{\infty}}(\mathcal{C}, \mathcal{D}) = H^0(A^u_{\infty}(\mathcal{C}, \mathcal{D}), m_1).$$

The corresponding notation for the 2-category is  $\overline{A^u_{\infty}}$ . We will see that arbitrary  $A_N$ -categories can be viewed as 2-functors  $\overline{A^u_{\infty}} \to \overline{A^u_{\infty}}$ . Moreover, they come from certain generalizations called  $A^u_{\infty}$ -2-functors. There is a notion of representability of such 2-functors, which explains some constructions of  $A_{\infty}$ -categories. For instance, a differential graded  $\Bbbk$ -quiver  $\Omega$  will be represented by the free  $A_{\infty}$ -category  $\mathcal{F}\Omega$  generated by it.

**3.1 Definition.** A (strict)  $A^u_{\infty}$ -2-functor  $F: A^u_{\infty} \to A^u_{\infty}$  consists of

- 1. a map  $F : \operatorname{Ob} A^u_{\infty} \to \operatorname{Ob} A^u_{\infty};$
- 2. a unital  $A_{\infty}$ -functor  $F = F_{\mathcal{C},\mathcal{D}} : A^{u}_{\infty}(\mathcal{C},\mathcal{D}) \to A^{u}_{\infty}(F\mathcal{C},F\mathcal{D})$  for each pair  $\mathcal{C},\mathcal{D}$  of unital  $A_{\infty}$ -categories; such that
- 3.  $\operatorname{id}_{F\mathcal{C}} = F(\operatorname{id}_{\mathcal{C}})$  for any unital  $A_{\infty}$ -category  $\mathcal{C}$ ;
- 4. the equation

$$TsA_{\infty}^{u}(\mathcal{C}, \mathcal{D}) \boxtimes TsA_{\infty}^{u}(\mathcal{D}, \mathcal{E}) \xrightarrow{M} TsA_{\infty}^{u}(\mathcal{C}, \mathcal{E})$$

$$F_{\mathcal{C}, \mathcal{D}} \boxtimes F_{\mathcal{D}, \mathcal{E}} \downarrow = \downarrow F_{\mathcal{C}, \mathcal{E}}$$

$$TsA_{\infty}^{u}(F\mathcal{C}, F\mathcal{D}) \boxtimes TsA_{\infty}^{u}(F\mathcal{D}, F\mathcal{E}) \xrightarrow{M} TsA_{\infty}^{u}(F\mathcal{C}, F\mathcal{E})$$

$$(3.1.1)$$

holds strictly for each triple  $\mathcal{C}, \mathcal{D}, \mathcal{E}$  of unital  $A_{\infty}$ -categories.

The  $A_{\infty}$ -functor  $F: A^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \to A^{u}_{\infty}(F\mathcal{C}, F\mathcal{D})$  consists of the mapping of objects

$$\operatorname{Ob} F : \operatorname{Ob} A^u_{\infty}(\mathcal{C}, \mathcal{D}) \to \operatorname{Ob} A^u_{\infty}(F\mathcal{C}, F\mathcal{D}), \qquad f \mapsto Ff,$$

and the components  $F_k, k \ge 1$ :

$$F_1: sA^u_{\infty}(\mathcal{C}, \mathcal{D})(f, g) \to sA^u_{\infty}(F\mathcal{C}, F\mathcal{D})(Ff, Fg),$$
  
$$F_2: sA^u_{\infty}(\mathcal{C}, \mathcal{D})(f, g) \otimes sA^u_{\infty}(\mathcal{C}, \mathcal{D})(g, h) \to sA^u_{\infty}(F\mathcal{C}, F\mathcal{D})(Ff, Fh),$$

and so on.

Weak versions of  $A^u_{\infty}$ -2-functors and 2-transformations between them might be considered elsewhere.

**3.2 Definition.** A (strict)  $A^u_{\infty}$ -2-transformation  $\lambda : F \to G : A^u_{\infty} \to A^u_{\infty}$  of strict  $A^u_{\infty}$ -2-functors is

- 1. a family of unital  $A_{\infty}$ -functors  $\lambda_{\mathfrak{C}} : F\mathfrak{C} \to G\mathfrak{C}, \ \mathfrak{C} \in \operatorname{Ob} A^{u}_{\infty}$ ; such that
- 2. the diagram of  $A_{\infty}$ -functors

$$A^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \xrightarrow{F} A^{u}_{\infty}(F\mathcal{C}, F\mathcal{D})$$

$$G \downarrow = \downarrow_{(1\boxtimes\lambda_{\mathcal{D}})M}$$

$$A^{u}_{\infty}(G\mathcal{C}, G\mathcal{D}) \xrightarrow{(\lambda_{\mathcal{C}}\boxtimes 1)M} A^{u}_{\infty}(F\mathcal{C}, G\mathcal{D})$$
(3.2.1)

strictly commutes.

An  $A^u_{\infty}$ -2-transformation  $\lambda = (\lambda_c)$  for which  $\lambda_c$  are  $A_{\infty}$ -equivalences is called a *natural*  $A^u_{\infty}$ -2-equivalence.

Let us show now that the above notions induce ordinary strict 2-functors and strict 2-transformations in 0-th cohomology. Recall that the strict 2-category  $\overline{A^u_{\infty}}$  consists of objects – unital  $A_{\infty}$ -categories, the category  $\overline{A^u_{\infty}}(\mathcal{C}, \mathcal{D})$  for any pair of objects  $\mathcal{C}, \mathcal{D}$ , the identity functor id<sub>c</sub> for any unital  $A_{\infty}$ -category  $\mathcal{C}$ , and the composition functor [Lyu03]

$$\overline{A^{u}_{\infty}}(\mathcal{C},\mathcal{D})(f,g) \times \overline{A^{u}_{\infty}}(\mathcal{D},\mathcal{E})(h,k) \xrightarrow{\bullet^{2}} \overline{A^{u}_{\infty}}(\mathcal{C},\mathcal{E})(fh,gk),$$
$$(rs^{-1},ps^{-1}) \longmapsto (rhs^{-1} \otimes gps^{-1})m_{2}$$

Given a strict  $A^u_{\infty}$ -2-functor F as in Definition 3.1 we construct from it an ordinary strict 2-functor  $\overline{F} = F$ :  $\operatorname{Ob} \overline{A^u_{\infty}} \to \operatorname{Ob} \overline{A^u_{\infty}}, \ \overline{F} = H^0(sF_1s^{-1}) : \overline{A^u_{\infty}}(\mathcal{C}, \mathcal{D}) \to \overline{A^u_{\infty}}(F\mathcal{C}, F\mathcal{D})$  as follows.

Denote

$$M_{10} \odot M_{01} = \left\{ sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \boxtimes sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \xrightarrow{\Delta_{10} \boxtimes \Delta_{01}} \\ \left[ sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \otimes T^{0}sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \right] \boxtimes \left[ T^{0}sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \otimes sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \right] \\ \xrightarrow{\sim} \left[ sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \boxtimes T^{0}sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \right] \otimes \left[ T^{0}sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \boxtimes sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \right] \\ \xrightarrow{M_{10} \otimes M_{01}} sA^{u}_{\infty}(\mathcal{C}, \mathcal{E}) \otimes sA^{u}_{\infty}(\mathcal{C}, \mathcal{E}) \right\}, \quad (3.2.2)$$

where the obvious isomorphisms  $\Delta_{10}$  and  $\Delta_{01}$  are components of the comultiplication  $\Delta$ , the middle isomorphism is that of distributivity law (1.5.3), and the components  $M_{10}$  and  $M_{01}$  of M are the composition maps.

Property (3.1.1) of F implies that

$$(M_{10} \odot M_{01})(F_1 \otimes F_1) = (F_1 \boxtimes F_1)(M_{10} \odot M_{01}).$$
(3.2.3)

Indeed, the following diagram commutes

$$sA^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \boxtimes T^{0}sA^{u}_{\infty}(\mathcal{D}, \mathcal{E}) \xrightarrow{M_{10}} sA^{u}_{\infty}(\mathcal{C}, \mathcal{E})$$

$$\downarrow F_{1} \boxtimes Ob F \downarrow \qquad \qquad \qquad \downarrow F_{1}$$

$$sA^{u}_{\infty}(F\mathcal{C}, F\mathcal{D}) \boxtimes T^{0}sA^{u}_{\infty}(F\mathcal{D}, F\mathcal{E}) \xrightarrow{M_{10}} sA^{u}_{\infty}(F\mathcal{C}, F\mathcal{E})$$

due to (3.1.1).  $\otimes$ -tensoring it with one more similar diagram we get

$$(M_{10} \otimes M_{01})(F_1 \otimes F_1) = [(F_1 \boxtimes \operatorname{Ob} F) \otimes (\operatorname{Ob} F \boxtimes F_1)](M_{10} \otimes M_{01}).$$

The isomorphisms in (3.2.2) commute with F in expected way, so (3.2.3) follows.

We claim that the diagram

$$sA_{\infty}^{u}(\mathcal{C},\mathcal{D}) \boxtimes sA_{\infty}^{u}(\mathcal{D},\mathcal{E}) \xrightarrow{(M_{10} \odot M_{01})B_{2}} sA_{\infty}^{u}(\mathcal{C},\mathcal{E})$$

$$\downarrow F_{1} \boxtimes F_{1} \qquad \qquad \qquad \downarrow F_{1}$$

$$sA_{\infty}^{u}(F\mathcal{C},F\mathcal{D}) \boxtimes sA_{\infty}^{u}(F\mathcal{D},F\mathcal{E}) \xrightarrow{(M_{10} \odot M_{01})B_{2}} sA_{\infty}^{u}(F\mathcal{C},F\mathcal{E})$$

$$(3.2.4)$$

homotopically commutes. Indeed, since

$$(1 \otimes B_1 + B_1 \otimes 1)F_2 + B_2F_1 = (F_1 \otimes F_1)B_2 + F_2B_1,$$

we get

$$(M_{10} \odot M_{01})B_2F_1$$
  
=  $(M_{10} \odot M_{01})(F_1 \otimes F_1)B_2 + (M_{10} \odot M_{01})F_2B_1 - (M_{10} \odot M_{01})(1 \otimes B_1 + B_1 \otimes 1)F_2$   
=  $(F_1 \boxtimes F_1)(M_{10} \odot M_{01})B_2 + (M_{10} \odot M_{01})F_2B_1 - (1 \boxtimes B_1 + B_1 \boxtimes 1)(M_{10} \odot M_{01})F_2.$ 

We have used equations

$$(M_{10} \odot M_{01})(1 \otimes B_1) = (1 \boxtimes B_1)(M_{10} \odot M_{01}), (M_{10} \odot M_{01})(B_1 \otimes 1) = (B_1 \boxtimes 1)(M_{10} \odot M_{01}),$$

which can be proved similarly to (3.2.3) due to M being an  $A_{\infty}$ -functor. Passing to cohomology we get from (3.2.4) a strictly commutative diagram of functors

since  $\cdot^2 = H^0((s \boxtimes s)(M_{10} \odot M_{01})B_2s^{-1})$ . Using the Künneth map we come to strictly commutative diagram of functors

that is, to a usual strict 2-functor  $\overline{F}: \overline{A^u_\infty} \to \overline{A^u_\infty}$ .

Let us show that an  $A^u_{\infty}$ -2-transformation  $\lambda : F \to G : A^u_{\infty} \to A^u_{\infty}$  as in Definition 3.2 induces an ordinary strict 2-transformation  $\overline{\lambda} : \overline{F} \to \overline{G} : \overline{A^u_{\infty}} \to \overline{A^u_{\infty}}$  in cohomology. Indeed, diagram (3.2.1) implies commutativity of diagram

Passing to cohomology we get

$$\frac{\overline{A_{\infty}^{u}}(\mathcal{C},\mathcal{D}) \xrightarrow{H^{0}(sF_{1}s^{-1})} \overline{A_{\infty}^{u}}(F\mathcal{C},F\mathcal{D})}{= \cdot \lambda_{\mathcal{D}} \downarrow = \overline{A_{\infty}^{u}}(F\mathcal{C},\lambda_{\mathcal{D}})} = \frac{\cdot \lambda_{\mathcal{D}} \downarrow = \overline{A_{\infty}^{u}}(F\mathcal{C},\lambda_{\mathcal{D}})}{\overline{A_{\infty}^{u}}(G\mathcal{C},G\mathcal{D}) \xrightarrow{\lambda_{\mathcal{C}} \cdot \cdot} \overline{A_{\infty}^{u}}(F\mathcal{C},G\mathcal{D})}$$

Therefore,  $\overline{\lambda_{\mathfrak{C}}} \in \operatorname{Ob} \overline{A_{\infty}^{u}}(F\mathfrak{C}, G\mathfrak{C})$  form a strict 2-transformation  $\overline{\lambda} : \overline{F} \to \overline{G} : \overline{A_{\infty}^{u}} \to \overline{A_{\infty}^{u}}$ .

**3.3.** Examples of  $A^u_{\infty}$ -2-functors. Let  $\mathcal{A}$  be an  $A_N$ -category,  $1 \leq N \leq \infty$ . It determines an  $A^u_{\infty}$ -2-functor  $F = A_N(\mathcal{A}, \_) : A^u_{\infty} \to A^u_{\infty}$ , given by the following data:

- 1. the map  $F : Ob A^u_{\infty} \to Ob A^u_{\infty}, \mathcal{C} \mapsto A_N(\mathcal{A}, \mathcal{C})$  (the category  $A_N(\mathcal{A}, \mathcal{C})$  is unital by [Lyu03, Proposition 7.7]);
- 2. the unital strict  $A_{\infty}$ -functor  $F = A_N(\mathcal{A}, \_) : A_{\infty}^u(\mathcal{C}, \mathcal{D}) \to A_{\infty}^u(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{D}))$ for each pair  $\mathcal{C}, \mathcal{D}$  of unital  $A_{\infty}$ -categories (cf. [Lyu03, Propositions 6.2, 8.4]).

Clearly,  $\mathrm{id}_{A_N(\mathcal{A},\mathcal{C})} = (1 \boxtimes \mathrm{id}_{\mathcal{C}})M = A_N(\mathcal{A}, \mathrm{id}_{\mathcal{C}})$ . We want to prove now that the equation

$$\begin{bmatrix} TsA_{\infty}^{u}(\mathcal{C},\mathcal{D}) \boxtimes TsA_{\infty}^{u}(\mathcal{D},\mathcal{E}) & \xrightarrow{M} TsA_{\infty}^{u}(\mathcal{C},\mathcal{E}) & \xrightarrow{A_{N}(\mathcal{A},-)} TsA_{\infty}^{u}(A_{N}(\mathcal{A},\mathcal{C}),A_{N}(\mathcal{A},\mathcal{E})) \end{bmatrix} \\ &= \begin{bmatrix} TsA_{\infty}^{u}(\mathcal{C},\mathcal{D}) \boxtimes TsA_{\infty}^{u}(\mathcal{D},\mathcal{E}) & \xrightarrow{A_{N}(\mathcal{A},-)\boxtimes A_{N}(\mathcal{A},-)} \\ TsA_{\infty}^{u}(A_{N}(\mathcal{A},\mathcal{C}),A_{N}(\mathcal{A},\mathcal{D})) \boxtimes TsA_{\infty}^{u}(A_{N}(\mathcal{A},\mathcal{D}),A_{N}(\mathcal{A},\mathcal{E})) \\ & \xrightarrow{M} TsA_{\infty}^{u}(A_{N}(\mathcal{A},\mathcal{C}),A_{N}(\mathcal{A},\mathcal{E})) \end{bmatrix}$$
(3.3.1)

holds strictly for each triple  $\mathcal{C}, \mathcal{D}, \mathcal{E}$  of unital  $A_{\infty}$ -categories. In fact, this F is a restriction of an  $A_{\infty}$ -2-functor  $F : \operatorname{Ob} A_{\infty} \to \operatorname{Ob} A_{\infty}, \mathcal{C} \mapsto A_N(\mathcal{A}, \mathcal{C})$ , which is defined just as in Definition 3.1 without mentioning the unitality. Equation (3.3.1) follows from a similar equation without the unitality index u. To prove it we consider the compositions

$$\begin{split} \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathbb{D}, \mathbb{E}) \xrightarrow{\text{IEM}} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{E}) \\ \xrightarrow{1\boxtimes A_N(\mathcal{A}, \mathbb{C})} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \xrightarrow{\alpha} TsA_N(\mathcal{A}, \mathbb{E}) \end{bmatrix} \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathbb{D}, \mathbb{E}) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{E}) \xrightarrow{M} TsA_N(\mathcal{A}, \mathbb{E}) \end{bmatrix} \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathbb{D}, \mathbb{E}) \\ \xrightarrow{M\boxtimes 1} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{E}) \xrightarrow{M} TsA_N(\mathcal{A}, \mathbb{E}) \end{bmatrix} \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ \xrightarrow{M\boxtimes 1} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ \xrightarrow{M\boxtimes 1} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{D}) \otimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ \xrightarrow{\alpha\boxtimes 1} TsA_N(\mathcal{A}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{A}, \mathbb{D}), A_N(\mathcal{A}, \mathbb{E}) \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ \xrightarrow{1\boxtimes A_N(\mathcal{A}, \mathbb{C})} \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ &= \begin{bmatrix} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathbb{C}, \mathbb{D}) \boxtimes TsA_{\infty}(\mathcal{D}, \mathbb{E}) \\ \xrightarrow{1\boxtimes A_N(\mathcal{A}, \mathbb{C})} \boxtimes TsA_{\infty}(\mathbb{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{D}) \otimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{D}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(A_N(\mathcal{A}, \mathbb{C}), A_N(\mathcal{A}, \mathbb{E})) \\ \xrightarrow{1\boxtimes M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathcal{A}, \mathbb{C}) \\ \xrightarrow{1\le M} TsA_N(\mathcal{A}, \mathbb{C}) \boxtimes TsA_{\infty}(\mathcal{A}, \mathbb{C}), A_N$$

By Proposition 1.6 we deduce equation (3.3.1) (see also [Lyu03, Proposition 5.5]).

Let now  $\mathcal{A}$  be a unital  $A_{\infty}$ -category. It determines an  $A_{\infty}^{u}$ -2-functor  $G = A_{\infty}^{u}(\mathcal{A}, \_)$ :  $A_{\infty}^{u} \to A_{\infty}^{u}$ , given by the following data:

- 1. the map  $G : \operatorname{Ob} A^u_{\infty} \to \operatorname{Ob} A^u_{\infty}, \mathfrak{C} \mapsto A^u_{\infty}(\mathcal{A}, \mathfrak{C})$  (the category  $A^u_{\infty}(\mathcal{A}, \mathfrak{C})$  is unital by [Lyu03, Proposition 7.7]);
- 2. the unital strict  $A_{\infty}$ -functor  $G = A^{u}_{\infty}(\mathcal{A}, \_) : A^{u}_{\infty}(\mathcal{C}, \mathcal{D}) \to A^{u}_{\infty}(A^{u}_{\infty}(\mathcal{A}, \mathcal{C}), A^{u}_{\infty}(\mathcal{A}, \mathcal{D}))$ for each pair  $\mathcal{C}, \mathcal{D}$  of unital  $A_{\infty}$ -categories, determined from

$$M = \begin{bmatrix} TsA_{\infty}^{u}(\mathcal{A}, \mathcal{B}) \boxtimes TsA_{\infty}^{u}(\mathcal{B}, \mathcal{C}) \xrightarrow{1\boxtimes A_{\infty}^{u}(\mathcal{A}, -)} \\ TsA_{\infty}^{u}(\mathcal{A}, \mathcal{B}) \boxtimes TsA_{\infty}^{u}(\mathcal{A}_{\infty}^{u}(\mathcal{A}, \mathcal{B}), A_{\infty}^{u}(\mathcal{A}, \mathcal{C})) \xrightarrow{\alpha} TsA_{\infty}^{u}(\mathcal{A}, \mathcal{C}) \end{bmatrix}$$

(cf. [Lyu03, Propositions 6.2, 8.4]).

Clearly, GC are full  $A_{\infty}$ -subcategories of FC for the  $A_{\infty}^{u}$ -2-functor  $F = A_{\infty}(\mathcal{A}, \_)$ . Furthermore,  $A_{\infty}$ -functors  $G_{\mathcal{C},\mathcal{D}}(f)$  are restrictions of  $A_{\infty}$ -functors  $F_{\mathcal{C},\mathcal{D}}(f)$ , so G is a full  $A_{\infty}^{u}$ -2-subfunctor of F. In particular, G satisfies equation (3.1.1). Another way to prove that G is an  $A_{\infty}^{u}$ -2-functor is to repeat the reasoning concerning F.

**3.4.** Example of an  $A^u_{\infty}$ -2-equivalence. Assume that  $\Omega$  is a differential graded k-quiver. As usual,  $\mathcal{F}\Omega$  denotes the free  $A_{\infty}$ -category generated by it. We claim that restr :

 $A_{\infty}(\mathfrak{FQ}, \_) \to A_1(\mathfrak{Q}, \_) : A_{\infty}^u \to A_{\infty}^u$  is a strict 2-natural  $A_{\infty}$ -equivalence. Indeed, it is given by the family of unital  $A_{\infty}$ -functors  $\operatorname{restr}_{\mathfrak{C}} : A_{\infty}(\mathfrak{FQ}, \mathfrak{C}) \to A_1(\mathfrak{Q}, \mathfrak{C}), \mathfrak{C} \in \operatorname{Ob} A_{\infty}^u$ , which are equivalences by Theorem 2.12. We have to prove that the diagram of  $A_{\infty}$ -functors

$$\begin{array}{cccc}
 & A^{u}_{\infty}(\mathcal{C}, \mathcal{D}) & \xrightarrow{A_{\infty}(\mathcal{FQ}, \_)} & A^{u}_{\infty}(A_{\infty}(\mathcal{FQ}, \mathcal{C}), A_{\infty}(\mathcal{FQ}, \mathcal{D})) \\ & & & \downarrow \\ & A_{1}(\mathcal{Q}, \_) & \downarrow & \downarrow \\ & & \downarrow \\ & A^{u}_{\infty}(A_{1}(\mathcal{Q}, \mathcal{C}), A_{1}(\mathcal{Q}, \mathcal{D})) & \xrightarrow{(\operatorname{restr}_{\mathcal{C}} \boxtimes 1)M} & A^{u}_{\infty}(A_{\infty}(\mathcal{FQ}, \mathcal{C}), A_{1}(\mathcal{Q}, \mathcal{D})) \end{array} \tag{3.4.1}$$

commutes. Notice that all arrows in this diagram are strict  $A_{\infty}$ -functors. Indeed,  $A_{\infty}(\mathcal{FQ}, \_)$ and  $A_1(\mathbb{Q}, \_)$  are strict by [Lyu03, Proposition 6.2]. For an arbitrary  $A_{\infty}$ -functor f the components  $[(f \boxtimes 1)M]_n = (f \boxtimes 1)M_{0n}$  vanish for all n except for n = 1, thus,  $(f \boxtimes 1)M$ is strict. The  $A_{\infty}$ -functor  $g = \operatorname{restr}_{\mathcal{D}}$  is strict, hence, the n-th component

$$[(1 \boxtimes g)M]_n : r^1 \otimes \cdots \otimes r^n \mapsto (r^1 \otimes \cdots \otimes r^n \mid g)M_{n0}$$

of the  $A_{\infty}$ -functor  $(1 \boxtimes g)M$  satisfies the equation

$$[(r^1 \otimes \cdots \otimes r^n \mid g)M_{n0}]_k = (r^1 \otimes \cdots \otimes r^n)\theta_{k1}g_1.$$

If the right hand side does not vanish, then  $n \leq 1 \leq k + n$ , so n = 1 and  $(1 \boxtimes g)M$  is strict.

Given an  $A_\infty$ -transformation  $t: g \to h: \mathbb{C} \to \mathcal{D}$  between unital  $A_\infty$ -functors we find that

$$\begin{aligned} A_{\infty}(\mathfrak{FQ},\_)(t) &= [(1\boxtimes t)M : (1\boxtimes g)M \to (1\boxtimes h)M : A_{\infty}(\mathfrak{FQ},\mathfrak{C}) \to A_{\infty}(\mathfrak{FQ},\mathfrak{D})], \\ A_{1}(\mathfrak{Q},\_)(t) &= [(1\boxtimes t)M : (1\boxtimes g)M \to (1\boxtimes h)M : A_{1}(\mathfrak{Q},\mathfrak{C}) \to A_{1}(\mathfrak{Q},\mathfrak{D})], \\ [(1\boxtimes \operatorname{restr}_{\mathfrak{D}})M]A_{\infty}(\mathfrak{FQ},\_)(t) &= [((1\boxtimes t)M) \cdot \operatorname{restr}_{\mathfrak{D}} : ((1\boxtimes g)M) \cdot \operatorname{restr}_{\mathfrak{D}} \\ &\to ((1\boxtimes h)M) \cdot \operatorname{restr}_{\mathfrak{D}} : A_{\infty}(\mathfrak{FQ},\mathfrak{C}) \to A_{1}(\mathfrak{Q},\mathfrak{D})], \\ [(\operatorname{restr}_{\mathfrak{C}}\boxtimes 1)M]A_{1}(\mathfrak{Q},\_)(t) &= [\operatorname{restr}_{\mathfrak{C}} \cdot ((1\boxtimes t)M) : \operatorname{restr}_{\mathfrak{C}} \cdot ((1\boxtimes g)M) \\ &\to \operatorname{restr}_{\mathfrak{C}} \cdot ((1\boxtimes h)M) : A_{\infty}(\mathfrak{FQ},\mathfrak{C}) \to A_{1}(\mathfrak{Q},\mathfrak{D})]. \end{aligned}$$

We have to verify that the last two  $A_{\infty}$ -transformations are equal. First of all, let us show that mappings of objects in (3.4.1) commute. Given a unital  $A_{\infty}$ -functor  $g : \mathcal{C} \to \mathcal{D}$ , we are going to check that

$$[(1 \boxtimes g)M]_n \cdot \operatorname{restr}_1 = \operatorname{restr}_1^{\otimes n} \cdot [(1 \boxtimes g)M]_n \tag{3.4.2}$$

for any  $n \ge 1$ . Indeed, for any *n*-tuple of composable  $A_{\infty}$ -transformations

$$f^0 \xrightarrow{r^1} f^1 \longrightarrow \dots \xrightarrow{r^n} f^n : \mathfrak{FQ} \to \mathcal{C},$$

we have in both cases

$$\{(r^{1} \otimes \cdots \otimes r^{n})[(1 \boxtimes g)M]_{n}\}_{0} = [(r^{1} \otimes \cdots \otimes r^{n}|g)M_{n0}]_{0} = (r_{0}^{1} \otimes \cdots \otimes r_{0}^{n})g_{n},$$
  

$$\{(r^{1} \otimes \cdots \otimes r^{n})[(1 \boxtimes g)M]_{n}\}_{1} = [(r^{1} \otimes \cdots \otimes r^{n}|g)M_{n0}]_{1}$$
  

$$= \sum_{i=1}^{n} (r_{0}^{1} \otimes \cdots \otimes r_{0}^{i-1} \otimes r_{1}^{i} \otimes r_{0}^{i+1} \otimes \cdots \otimes r_{0}^{n})g_{n}$$
  

$$+ \sum_{i=1}^{n} (r_{0}^{1} \otimes \cdots \otimes r_{0}^{i-1} \otimes f_{1}^{i} \otimes r_{0}^{i+1} \otimes \cdots \otimes r_{0}^{n})g_{n+1}.$$

Note that the right hand sides depend only on 0-th and 1-st components of  $r^i$ ,  $f^i$ . This is precisely what is claimed by equation (3.4.2).

The coincidence of  $A_{\infty}$ -transformations  $((1 \boxtimes t)M) \cdot \operatorname{restr}_{\mathcal{D}} = \operatorname{restr}_{\mathcal{C}} \cdot ((1 \boxtimes t)M)$  follows similarly from the computation:

$$\{(r^{1} \otimes \cdots \otimes r^{n})[(1 \boxtimes t)M]_{n}\}_{0} = [(r^{1} \otimes \cdots \otimes r^{n} \boxtimes t)M_{n1}]_{0} = (r_{0}^{1} \otimes \cdots \otimes r_{0}^{n})t_{n},$$
  
$$\{(r^{1} \otimes \cdots \otimes r^{n})[(1 \boxtimes t)M]_{n}\}_{1} = [(r^{1} \otimes \cdots \otimes r^{n} \boxtimes t)M_{n1}]_{1}$$
  
$$= \sum_{i=1}^{n} (r_{0}^{1} \otimes \cdots \otimes r_{0}^{i-1} \otimes r_{1}^{i} \otimes r_{0}^{i+1} \otimes \cdots \otimes r_{0}^{n})t_{n}$$
  
$$+ \sum_{i=1}^{n} (r_{0}^{1} \otimes \cdots \otimes r_{0}^{i-1} \otimes f_{1}^{i} \otimes r_{0}^{i+1} \otimes \cdots \otimes r_{0}^{n})t_{n+1}.$$

**3.5.** Representability. An  $A^u_{\infty}$ -2-functor  $F : A^u_{\infty} \to A^u_{\infty}$  is called *representable*, if it is naturally  $A^u_{\infty}$ -2-equivalent to the  $A^u_{\infty}$ -2-functor  $A_{\infty}(\mathcal{A}, \_) : A^u_{\infty} \to A^u_{\infty}$  for some  $A_{\infty}$ -category  $\mathcal{A}$ . The above results imply that the  $A^u_{\infty}$ -2-functor  $A_1(\mathcal{Q}, \_)$  corresponding to a differential graded k-quiver  $\mathcal{Q}$  is represented by the free  $A_{\infty}$ -category  $\mathcal{F}\mathcal{Q}$  generated by  $\mathcal{Q}$ .

This definition of representability has a disadvantage: many different  $A_{\infty}$ -categories can represent the same  $A^u_{\infty}$ -2-functor. More attractive notion is the following. An  $A^u_{\infty}$ -2-functor  $F: A^u_{\infty} \to A^u_{\infty}$  is called *unitally representable*, if it is naturally  $A^u_{\infty}$ -2-equivalent to the  $A^u_{\infty}$ -2-functor  $A^u_{\infty}(\mathcal{A}, \_): A^u_{\infty} \to A^u_{\infty}$  for some unital  $A_{\infty}$ -category  $\mathcal{A}$ . Such  $\mathcal{A}$ is unique up to an  $A_{\infty}$ -equivalence. Indeed, composing a natural 2-equivalence  $\overline{\lambda}:$  $\overline{A^u_{\infty}(\mathcal{A},\_)} \to \overline{A^u_{\infty}(\mathcal{B},\_)}: \overline{A^u_{\infty}} \to \overline{A^u_{\infty}}$  with the 0-th cohomology 2-functor  $H^0: \overline{A^u_{\infty}} \to \mathbb{C}at$ , we get a natural 2-equivalence  $H^0\overline{\lambda}: H^0\overline{A^u_{\infty}(\mathcal{A},\_)} \to H^0\overline{A^u_{\infty}(\mathcal{B},\_)}: \overline{A^u_{\infty}} \to \mathbb{C}at$ . However,  $H^0\overline{A^u_{\infty}(\mathcal{A},\_)} = \overline{A^u_{\infty}(\mathcal{A},\_)}$ , so using a 2-category version of Yoneda lemma one can deduce that  $\mathcal{A}$  and  $\mathcal{B}$  are equivalent in  $\overline{A^u_{\infty}}$ . We shall present an example of unital representability in subsequent publication [LM04].

**3.6.** Acknowledgements. We are grateful to all the participants of the  $A_{\infty}$ -category seminar at the Institute of Mathematics, Kyiv, for attention and fruitful discussions, especially to Yu. Bespalov and S. Ovsienko. One of us (V.L.) is grateful to Max-Planck-Institut für Mathematik for warm hospitality and support at the final stage of this research.

### References

- [Fuk93] K. Fukaya, Morse homotopy, A<sub>∞</sub>-category, and Floer homologies, Proc. of GARC Workshop on Geometry and Topology '93 (H. J. Kim, ed.), Lecture Notes, no. 18, Seoul Nat. Univ., Seoul, 1993, pp. 1–102.
- [Fuk02] K. Fukaya, Floer homology and mirror symmetry. II, Minimal surfaces, geometric analysis and symplectic geometry (Baltimore, MD, 1999), Adv. Stud. Pure Math., vol. 34, Math. Soc. Japan, Tokyo, 2002, pp. 31–127.
- [Kad82] T. V. Kadeishvili, The algebraic structure in the homology of an  $A(\infty)$ -algebra, Soobshch. Akad. Nauk Gruzin. SSR **108** (1982), no. 2, 249–252, in Russian.
- [Kel01] B. Keller, Introduction to A-infinity algebras and modules, Homology, Homotopy and Applications 3 (2001), no. 1, 1-35, http://arXiv.org/abs/math.RA/ 9910179, http://www.rmi.acnet.ge/hha/.
- [Kon95] M. Kontsevich, Homological algebra of mirror symmetry, Proc. Internat. Cong. Math., Zürich, Switzerland 1994 (Basel), vol. 1, Birkhäuser Verlag, 1995, 120– 139.
- [KS02] M. Kontsevich and Y. S. Soibelman,  $A_{\infty}$ -categories and non-commutative geometry, in preparation, 2002.
- [KS] M. Kontsevich and Y. S. Soibelman, *Deformation theory*, book in preparation.
- [LH03] K. Lefèvre-Hasegawa, Sur les  $A_{\infty}$ -catégories, Ph.D. thesis, Université Paris 7, U.F.R. de Mathématiques, 2003, http://arXiv.org/abs/math.CT/0310337.
- [LM04] V. V. Lyubashenko and O. Manzyuk, *Quotients of unital*  $A_{\infty}$ -categories, http: //arXiv.org/abs/math.CT/0306018, 2004.
- [LO02] V. V. Lyubashenko and S. A. Ovsienko, A construction of an  $A_{\infty}$ -category, http: //arXiv.org/abs/math.CT/0211037, 2002.
- [Lyu03] V. V. Lyubashenko, Category of  $A_{\infty}$ -categories, Homology, Homotopy and Applications 5 (2003), no. 1, 1–48, http://arXiv.org/abs/math.CT/0210047, http://www.rmi.acnet.ge/hha/.
- [Sta63] J. D. Stasheff, Homotopy associativity of H-spaces, I & II, Trans. Amer. Math. Soc. 108 (1963), 275–292, 293–312.