

Free A_∞ -categories

Volodymyr Lyubashenko* and Oleksandr Manzyuk†

February 8, 2020

Abstract

For a differential graded \mathbb{k} -quiver \mathcal{Q} we define the free A_∞ -category \mathcal{FQ} generated by \mathcal{Q} . The main result is that the restriction A_∞ -functor $A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$ is an equivalence, where objects of the last A_∞ -category are morphisms of differential graded \mathbb{k} -quivers $\mathcal{Q} \rightarrow \mathcal{A}$.

A_∞ -categories defined by Fukaya [Fuk93] and Kontsevich [Kon95] are generalizations of differential graded categories for which the binary composition is associative only up to a homotopy. They also generalize A_∞ -algebras introduced by Stasheff [Sta63, II]. A_∞ -functors are the corresponding generalizations of usual functors, see e.g. [Fuk93, Kel01]. Homomorphisms of A_∞ -algebras (e.g. [Kad82]) are particular cases of A_∞ -functors. A_∞ -transformations are certain coderivations. Examples of such structures are encountered in studies of mirror symmetry (e.g. [Kon95, Fuk02]) and in homological algebra.

For an A_∞ -category there is a notion of units up to a homotopy (homotopy identity morphisms) [Lyu03]. Given two A_∞ -categories \mathcal{A} and \mathcal{B} , one can construct a third A_∞ -category $A_\infty(\mathcal{A}, \mathcal{B})$, whose objects are A_∞ -functors $f : \mathcal{A} \rightarrow \mathcal{B}$, and morphisms are A_∞ -transformations between such functors (Fukaya [Fuk02], Kontsevich and Soibelman [KS02, KS], Lefèvre-Hasegawa [LH03], as well as [Lyu03]). This allows to define a 2-category, whose objects are unital A_∞ -categories, 1-morphisms are unital A_∞ -functors and 2-morphisms are equivalence classes of natural A_∞ -transformations [Lyu03]. We continue to study this 2-category.

The notations and conventions are explained in the first section. We also describe A_N -categories, A_N -functors and A_N -transformations – truncated at $N < \infty$ versions of A_∞ -categories. For instance, A_1 -categories and A_1 -functors are differential graded \mathbb{k} -quivers and their morphisms. However, A_1 -transformations bring new 2-categorical features to the theory. In particular, for any differential graded \mathbb{k} -quiver \mathcal{Q} and any A_∞ -category

*Institute of Mathematics, National Academy of Sciences of Ukraine, 3 Tereshchenkivska st., Kyiv-4, 01601 MSP, Ukraine; lub@imath.kiev.ua

†Fachbereich Mathematik, Postfach 3049, 67653 Kaiserslautern, Germany; manzyuk@mathematik.uni-kl.de

\mathcal{A} there is an A_∞ -category $A_1(\mathcal{Q}, \mathcal{A})$, whose objects are morphisms of differential graded \mathbb{k} -quivers $\mathcal{Q} \rightarrow \mathcal{A}$, and morphisms are A_1 -transformations. We recall the terminology related to trees in Section 1.7.

In the second section we define the free A_∞ -category \mathcal{FQ} generated by a differential graded \mathbb{k} -quiver \mathcal{Q} . We classify functors from a free A_∞ -category \mathcal{FQ} to an arbitrary A_∞ -category \mathcal{A} in Proposition 2.3. In particular, the restriction map gives a bijection between the set of strict A_∞ -functors $\mathcal{FQ} \rightarrow \mathcal{A}$ and the set of morphisms of differential graded \mathbb{k} -quivers $\mathcal{Q} \rightarrow (\mathcal{A}, m_1)$ (Corollary 2.4). We classify chain maps into complexes of transformations whose source is a free A_∞ -category in Proposition 2.8. Description of homotopies between such chain maps is given in Corollary 2.10. Assuming in addition that \mathcal{A} is unital, we obtain our main result: the restriction A_∞ -functor $\text{restr} : A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$ is an equivalence (Theorem 2.12).

In the third section we interpret $A_\infty(\mathcal{FQ}, _)$ and $A_1(\mathcal{Q}, _)$ as strict A_∞^u -2-functors $A_\infty^u \rightarrow A_\infty^u$. Moreover, we interpret $\text{restr} : A_\infty(\mathcal{FQ}, _) \rightarrow A_1(\mathcal{Q}, _)$ as an A_∞^u -2-equivalence. In this sense the A_∞ -category \mathcal{FQ} represents the A_∞^u -2-functor $A_1(\mathcal{Q}, _)$. This is the 2-categorical meaning of freeness of \mathcal{FQ} .

1. Conventions and preliminaries

We keep the notations and conventions of [Lyu03, LO02], sometimes without explicit mentioning. Some of the conventions are recalled here.

We assume as in [Lyu03, LO02] that most quivers, A_∞ -categories, etc. are small with respect to some universe \mathcal{U} .

The ground ring $\mathbb{k} \in \mathcal{U}$ is a unital associative commutative ring. A \mathbb{k} -module means a \mathcal{U} -small \mathbb{k} -module.

We use the right operators: the composition of two maps (or morphisms) $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ is denoted $fg : X \rightarrow Z$; a map is written on elements as $f : x \mapsto xf = (x)f$. However, these conventions are not used systematically, and $f(x)$ might be used instead.

\mathbb{Z} -graded \mathbb{k} -modules are functions $X : \mathbb{Z} \ni d \mapsto X^d \in \mathbb{k}\text{-mod}$. A simple computation shows that the product $X = \prod_{\iota \in I} X_\iota$ in the category of \mathbb{Z} -graded \mathbb{k} -modules of a family $(X_\iota)_{\iota \in I}$ of objects $X_\iota : d \mapsto X_\iota^d$ is given by $X : \mathbb{Z} \ni d \mapsto X^d = \prod_{\iota \in I} X_\iota^d$. Everywhere in this article the product of graded \mathbb{k} -modules means the above product.

If P is a \mathbb{Z} -graded \mathbb{k} -module, then $sP = P[1]$ denotes the same \mathbb{k} -module with the grading $(sP)^d = P^{d+1}$. The “identity” map $P \rightarrow sP$ of degree -1 is also denoted s . The map s commutes with the components of the differential in an A_∞ -category (A_∞ -algebra) in the following sense: $s^{\otimes n} b_n = m_n s$.

Let $\mathbf{C} = \mathbf{C}(\mathbb{k}\text{-mod})$ denote the differential graded category of complexes of \mathbb{k} -modules. Actually, it is a symmetric closed monoidal category.

The cone of a chain of a chain map $\alpha : P \rightarrow Q$ of complexes of \mathbb{k} -modules is the graded \mathbb{k} -module $\text{Cone}(\alpha) = Q \oplus P[1]$ with the differential $(q, ps)d = (qd^Q + p\alpha, psd^{P[1]}) = (qd^Q + p\alpha, -pd^P s)$.

1.1. A_N -categories. For a positive integer N we define some A_N -notions similarly to the case $N = \infty$. We may say that all data, equations and constructions for A_N -case are the same as in A_∞ -case (e.g. [Lyu03]), however, taken only up to level N .

A *differential graded \mathbb{k} -quiver* \mathcal{Q} is the following data: a \mathcal{U} -small set of objects $\text{Ob } \mathcal{Q}$; a chain complex of \mathbb{k} -modules $\mathcal{Q}(X, Y)$ for each pair of objects X, Y . A morphism of differential graded \mathbb{k} -quivers $f : \mathcal{Q} \rightarrow \mathcal{A}$ is given by a map $f : \text{Ob } \mathcal{Q} \rightarrow \text{Ob } \mathcal{A}$, $X \mapsto Xf$ and by a chain map $\mathcal{Q}(X, Y) \rightarrow \mathcal{A}(Xf, Yf)$ for each pair of objects X, Y of \mathcal{Q} . The category of differential graded \mathbb{k} -quivers is denoted A_1 .

The category of \mathcal{U} -small graded \mathbb{k} -linear quivers, whose set of objects is S , admits a monoidal structure with the tensor product $\mathcal{A} \times \mathcal{B} \mapsto \mathcal{A} \otimes \mathcal{B}$, $(\mathcal{A} \otimes \mathcal{B})(X, Y) = \bigoplus_{Z \in S} \mathcal{A}(X, Z) \otimes_{\mathbb{k}} \mathcal{B}(Z, Y)$. In particular, we have tensor powers $T^n \mathcal{A} = \mathcal{A}^{\otimes n}$ of a given graded \mathbb{k} -quiver \mathcal{A} , such that $\text{Ob } T^n \mathcal{A} = \text{Ob } \mathcal{A}$. Explicitly,

$$T^n \mathcal{A}(X, Y) = \bigoplus_{X=X_0, X_1, \dots, X_n=Y \in \text{Ob } \mathcal{A}} \mathcal{A}(X_0, X_1) \otimes_{\mathbb{k}} \mathcal{A}(X_1, X_2) \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} \mathcal{A}(X_{n-1}, X_n).$$

In particular, $T^0 \mathcal{A}(X, Y) = \mathbb{k}$ if $X = Y$ and vanishes otherwise. The graded \mathbb{k} -quiver $T^{\leq N} \mathcal{A} = \bigoplus_{n \geq 0}^N T^n \mathcal{A}$ is called the restricted tensor coalgebra of \mathcal{A} . It is equipped with the cut comultiplication

$$\begin{aligned} \Delta : T^{\leq N} \mathcal{A}(X, Y) &\rightarrow \bigoplus_{Z \in \text{Ob } \mathcal{A}} T^{\leq N} \mathcal{A}(X, Z) \bigotimes_{\mathbb{k}} T^{\leq N} \mathcal{A}(Z, Y), \\ h_1 \otimes h_2 \otimes \cdots \otimes h_n &\mapsto \sum_{k=0}^n h_1 \otimes \cdots \otimes h_k \bigotimes_{\mathbb{k}} h_{k+1} \otimes \cdots \otimes h_n, \end{aligned}$$

and the counit $\varepsilon = (T^{\leq N} \mathcal{A}(X, Y) \xrightarrow{\text{pr}_0} T^0 \mathcal{A}(X, Y) \rightarrow \mathbb{k})$, where the last map is $\text{id}_{\mathbb{k}}$ if $X = Y$, or 0 if $X \neq Y$ (and $T^0 \mathcal{A}(X, Y) = 0$). We write $T\mathcal{A}$ instead of $T^{\leq \infty} \mathcal{A}$. If $g : T\mathcal{A} \rightarrow T\mathcal{B}$ is a map of \mathbb{k} -quivers, then g_{ac} denotes its matrix coefficient $T^a \mathcal{A} \xrightarrow{\text{in}_a} T\mathcal{A} \xrightarrow{g} T\mathcal{B} \xrightarrow{\text{pr}_c} T^c \mathcal{B}$. The matrix coefficient g_{a1} is abbreviated to g_a .

1.2 Definition. An A_N -category \mathcal{A} consists of the following data: a graded \mathbb{k} -quiver \mathcal{A} ; a system of \mathbb{k} -linear maps of degree 1

$$b_n : s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \cdots \otimes s\mathcal{A}(X_{n-1}, X_n) \rightarrow s\mathcal{A}(X_0, X_n), \quad 1 \leq n \leq N,$$

such that for all $1 \leq k \leq N$

$$\sum_{r+n+t=k} (1^{\otimes r} \otimes b_n \otimes 1^{\otimes t}) b_{r+1+t} = 0 : T^k s\mathcal{A} \rightarrow s\mathcal{A}. \quad (1.2.1)$$

The system b_n is interpreted as a (1,1)-coderivation $b : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N} s\mathcal{A}$ of degree 1 determined by

$$b_{kl} = (b|_{T^k s\mathcal{A}}) \text{pr}_l : T^k s\mathcal{A} \rightarrow T^l s\mathcal{A}, \quad b_{kl} = \sum_{\substack{r+n+t=k \\ r+1+t=l}} 1^{\otimes r} \otimes b_n \otimes 1^{\otimes t}, \quad k, l \leq N,$$

which is a differential.

1.3 Definition. A *pointed cocategory homomorphism* consists of the following data: A_N -categories \mathcal{A} and \mathcal{B} , a map $f : \text{Ob } \mathcal{A} \rightarrow \text{Ob } \mathcal{B}$ and a system of \mathbb{k} -linear maps of degree 0

$$f_n : s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \cdots \otimes s\mathcal{A}(X_{n-1}, X_n) \rightarrow s\mathcal{B}(X_0 f, X_n f), \quad 1 \leq n \leq N.$$

The above data are equivalent to a cocategory homomorphism $f : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N} s\mathcal{B}$ of degree 0 such that

$$f_{01} = (f|_{T^0 s\mathcal{A}}) \text{pr}_1 = 0 : T^0 s\mathcal{A} \rightarrow T^1 s\mathcal{B}, \quad (1.3.1)$$

(this condition was implicitly assumed in [Lyu03, Definition 2.4]). The components of f are

$$f_{kl} = (f|_{T^k s\mathcal{A}}) \text{pr}_l : T^k s\mathcal{A} \rightarrow T^l s\mathcal{B}, \quad f_{kl} = \sum_{i_1 + \cdots + i_l = k} f_{i_1} \otimes f_{i_2} \otimes \cdots \otimes f_{i_l}, \quad (1.3.2)$$

where $k, l \leq N$. Indeed, the claim follows from the following diagram, commutative for all $l \geq 0$:

$$\begin{array}{ccccc} T^{\leq N} s\mathcal{A} & \xrightarrow{f} & T^{\leq N} s\mathcal{B} & \xrightarrow{\text{pr}_l} & T^l s\mathcal{B} \\ \Delta^{(l)} \downarrow & = & \Delta^{(l)} \downarrow & = & \parallel \\ (T^{\leq N} s\mathcal{A})^{\otimes l} & \xrightarrow{f^{\otimes l}} & (T^{\leq N} s\mathcal{B})^{\otimes l} & \xrightarrow{\text{pr}_1^{\otimes l}} & (s\mathcal{B})^{\otimes l} \end{array}$$

where $\Delta^{(0)} = \varepsilon$, $\Delta^{(1)} = \text{id}$, $\Delta^{(2)} = \Delta$ and $\Delta^{(l)}$ means the cut comultiplication, iterated $l - 1$ times. Notice that condition (1.3.1) can be written as $f_0 = 0$.

1.4 Definition. An A_N -functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is a pointed cocategory homomorphism, which commutes with the differential b , that is, for all $1 \leq k \leq N$

$$\sum_{l > 0; i_1 + \cdots + i_l = k} (f_{i_1} \otimes f_{i_2} \otimes \cdots \otimes f_{i_l}) b_l = \sum_{r+n+t=k} (1^{\otimes r} \otimes b_n \otimes 1^{\otimes t}) f_{r+1+t} : T^k s\mathcal{A} \rightarrow s\mathcal{B}.$$

We are interested mostly in the case $N = 1$. Clearly, A_1 -categories are differential graded quivers and A_1 -functors are their morphisms. In the case of one object these reduce to chain complexes and chain maps. The following notion seems interesting even in this case.

1.5 Definition. An A_N -transformation $r : f \rightarrow g : \mathcal{A} \rightarrow \mathcal{B}$ of degree d consists of the following data: A_N -categories \mathcal{A} and \mathcal{B} ; pointed cocategory homomorphisms $f, g : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N} s\mathcal{B}$ (or A_N -functors $f, g : \mathcal{A} \rightarrow \mathcal{B}$); a system of \mathbb{k} -linear maps of degree d

$$r_n : s\mathcal{A}(X_0, X_1) \otimes s\mathcal{A}(X_1, X_2) \otimes \cdots \otimes s\mathcal{A}(X_{n-1}, X_n) \rightarrow s\mathcal{B}(X_0 f, X_n g), \quad 0 \leq n \leq N.$$

To give a system r_n is equivalent to specifying an (f, g) -coderivation $r : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N+1} s\mathcal{B}$ of degree d

$$\begin{aligned} r_{kl} &= (r|_{T^k s\mathcal{A}}) \text{pr}_l : T^k s\mathcal{A} \rightarrow T^l s\mathcal{B}, \quad k \leq N, l \leq N+1 \\ r_{kl} &= \sum_{\substack{q+1+t=l \\ i_1+\dots+i_q+n+j_1+\dots+j_t=k}} f_{i_1} \otimes \dots \otimes f_{i_q} \otimes r_n \otimes g_{j_1} \otimes \dots \otimes g_{j_t}, \end{aligned} \quad (1.5.1)$$

that is, a \mathbb{k} -quiver morphism r , satisfying $r\Delta = \Delta(f \otimes r + r \otimes g)$. This follows from the commutative diagram

$$\begin{array}{ccccc} T^{\leq N} s\mathcal{A} & \xrightarrow{r} & T^{\leq N+1} s\mathcal{B} & \xrightarrow{\text{pr}_l} & T^l s\mathcal{B} \\ \Delta^{(l)} \downarrow & & \Delta^{(l)} \downarrow & & \parallel \\ (T^{\leq N} s\mathcal{A})^{\otimes l} & \xrightarrow{\sum_{q+1+t=l} f^{\otimes q} \otimes r \otimes g^{\otimes t}} & (T^{\leq N+1} s\mathcal{B})^{\otimes l} & \xrightarrow{\text{pr}_1^{\otimes l}} & (s\mathcal{B})^{\otimes l} \end{array}$$

Let \mathcal{A}, \mathcal{B} be A_N -categories, and let $f^0, f^1, \dots, f^n : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N} s\mathcal{B}$ be pointed cocategory homomorphisms. Consider coderivations r_1, \dots, r_n as in

$$f^0 \xrightarrow{r^1} f^1 \xrightarrow{r^2} \dots f^{n-1} \xrightarrow{r^n} f^n : T^{\leq N} s\mathcal{A} \rightarrow T^{\leq N} s\mathcal{B}.$$

We construct the following system of \mathbb{k} -linear maps $\theta_{kl} : T^k s\mathcal{A} \rightarrow T^l s\mathcal{B}$, $k \leq N, l \leq N+n$ of degree $\deg r^1 + \dots + \deg r^n$ from these data:

$$\theta_{kl} = \sum f_{i_1^0}^0 \otimes \dots \otimes f_{i_{m_0}^0}^0 \otimes r_{j_1}^1 \otimes f_{i_1^1}^1 \otimes \dots \otimes f_{i_{m_1}^1}^1 \otimes \dots \otimes r_{j_n}^n \otimes f_{i_1^n}^n \otimes \dots \otimes f_{i_{m_n}^n}^n, \quad (1.5.2)$$

where summation is taken over all terms with

$$m_0 + m_1 + \dots + m_n + n = l, \quad i_1^0 + \dots + i_{m_0}^0 + j_1 + i_1^1 + \dots + i_{m_1}^1 + \dots + j_n + i_1^n + \dots + i_{m_n}^n = k.$$

Equivalently, we write

$$\theta_{kl} = \sum_{\substack{m_0+m_1+\dots+m_n+n=l \\ p_0+j_1+p_1+\dots+j_n+p_n=k}} f_{p_0 m_0}^0 \otimes r_{j_1}^1 \otimes f_{p_1 m_1}^1 \otimes \dots \otimes r_{j_n}^n \otimes f_{p_n m_n}^n.$$

The component θ_{kl} vanishes unless $n \leq l \leq k+n$. If $n=0$, then θ_{kl} is expansion (1.3.2) of f^0 . If $n=1$, then θ_{kl} is expansion (1.5.1) of r^1 .

Given an A_K -category \mathcal{A} and an A_{K+N} -category \mathcal{B} , $1 \leq K, N \leq \infty$, we construct an A_N -category $A_K(\mathcal{A}, \mathcal{B})$ out of these. The objects of $A_K(\mathcal{A}, \mathcal{B})$ are A_K -functors $f : \mathcal{A} \rightarrow \mathcal{B}$. Given two such functors $f, g : \mathcal{A} \rightarrow \mathcal{B}$ we define the graded \mathbb{k} -module $A_K(\mathcal{A}, \mathcal{B})(f, g)$ as the space of all A_K -transformations $r : f \rightarrow g$, namely,

$$\begin{aligned} [A_K(\mathcal{A}, \mathcal{B})(f, g)]^{d+1} \\ = \{r : f \rightarrow g \mid A_K\text{-transformation } r : T^{\leq K} s\mathcal{A} \rightarrow T^{\leq K+1} s\mathcal{B} \text{ has degree } d\}. \end{aligned}$$

The system of differentials B_n , $n \leq N$, is defined as follows:

$$\begin{aligned}
B_1 : A_K(\mathcal{A}, \mathcal{B})(f, g) &\rightarrow A_K(\mathcal{A}, \mathcal{B})(f, g), \quad r \mapsto (r)B_1 = [r, b] = rb - (-)^r br, \\
[(r)B_1]_k &= \sum_{i_1 + \dots + i_q + n + j_1 + \dots + j_t = k} (f_{i_1} \otimes \dots \otimes f_{i_q} \otimes r_n \otimes g_{j_1} \otimes \dots \otimes g_{j_t}) b_{q+1+t} \\
&\quad - (-)^r \sum_{\alpha + n + \beta = k} (1^{\otimes \alpha} \otimes b_n \otimes 1^{\otimes \beta}) r_{\alpha+1+\beta}, \quad k \leq K, \\
B_n : A_K(\mathcal{A}, \mathcal{B})(f^0, f^1) \otimes \dots \otimes A_K(\mathcal{A}, \mathcal{B})(f^{n-1}, f^n) &\rightarrow A_K(\mathcal{A}, \mathcal{B})(f^0, f^n), \\
r^1 \otimes \dots \otimes r^n &\mapsto (r^1 \otimes \dots \otimes r^n)B_n, \text{ for } 1 < n \leq N,
\end{aligned}$$

where the last A_K -transformation is defined by its components:

$$[(r^1 \otimes \dots \otimes r^n)B_n]_k = \sum_{l=n}^{n+k} (r^1 \otimes \dots \otimes r^n) \theta_{kl} b_l, \quad k \leq K.$$

The category of graded \mathbb{k} -linear quivers admits a symmetric monoidal structure with the tensor product $\mathcal{A} \times \mathcal{B} \mapsto \mathcal{A} \boxtimes \mathcal{B}$, where $\text{Ob } \mathcal{A} \boxtimes \mathcal{B} = \text{Ob } \mathcal{A} \times \text{Ob } \mathcal{B}$ and $(\mathcal{A} \boxtimes \mathcal{B})((X, U), (Y, V)) = \mathcal{A}(X, Y) \otimes_{\mathbb{k}} \mathcal{B}(U, V)$. The same tensor product was denoted \otimes in [Lyu03], but we will keep notation $\mathcal{A} \otimes \mathcal{B}$ only for tensor product from Section 1.1, defined when $\text{Ob } \mathcal{A} = \text{Ob } \mathcal{B}$. The two tensor products obey

Distributivity law. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ be graded \mathbb{k} -linear quivers, such that $\text{Ob } \mathcal{A} = \text{Ob } \mathcal{B}$ and $\text{Ob } \mathcal{C} = \text{Ob } \mathcal{D}$. Then the middle four interchange map $1 \otimes c \otimes 1$ is an isomorphism of quivers

$$(\mathcal{A} \otimes \mathcal{B}) \boxtimes (\mathcal{C} \otimes \mathcal{D}) \xrightarrow{\sim} (\mathcal{A} \boxtimes \mathcal{C}) \otimes (\mathcal{B} \boxtimes \mathcal{D}), \quad (1.5.3)$$

identity on objects.

Indeed, the both quivers in (1.5.3) have the same set of objects $R \times S$, where $R = \text{Ob } \mathcal{A} = \text{Ob } \mathcal{B}$ and $S = \text{Ob } \mathcal{C} = \text{Ob } \mathcal{D}$. Let $X, Z \in R$ and $U, W \in S$. The sets of morphisms from (X, U) to (Z, W) are isomorphic via

$$\begin{aligned}
&((\mathcal{A} \otimes \mathcal{B}) \boxtimes (\mathcal{C} \otimes \mathcal{D}))((X, U), (Z, W)) = \\
&(\oplus_{Y \in R} \mathcal{A}(X, Y) \otimes_{\mathbb{k}} \mathcal{B}(Y, Z)) \otimes_{\mathbb{k}} (\oplus_{V \in S} \mathcal{C}(U, V) \otimes_{\mathbb{k}} \mathcal{D}(V, W)) \\
&\quad \downarrow \wr \\
&\oplus_{(Y, V) \in R \times S} \mathcal{A}(X, Y) \otimes_{\mathbb{k}} \mathcal{B}(Y, Z) \otimes_{\mathbb{k}} \mathcal{C}(U, V) \otimes_{\mathbb{k}} \mathcal{D}(V, W) \\
&\quad \downarrow 1 \otimes c \otimes 1 \\
&\oplus_{(Y, V) \in R \times S} \mathcal{A}(X, Y) \otimes_{\mathbb{k}} \mathcal{C}(U, V) \otimes_{\mathbb{k}} \mathcal{B}(Y, Z) \otimes_{\mathbb{k}} \mathcal{D}(V, W) \\
&= ((\mathcal{A} \boxtimes \mathcal{C}) \otimes (\mathcal{B} \boxtimes \mathcal{D}))((X, U), (Z, W)).
\end{aligned}$$

The notion of a pointed cocategory homomorphism extends to the case of several arguments, that is, to degree 0 cocategory homomorphisms $\psi : T^{\leq L^1} s \mathcal{C}^1 \boxtimes \dots \boxtimes T^{\leq L^q} s \mathcal{C}^q \rightarrow$

$T^{\leq N} s\mathcal{B}$, where $N \geq L^1 + \dots + L^q$. We always assume that $\psi_{00\dots 0} : T^0 s\mathcal{C}^1 \boxtimes \dots \boxtimes T^0 s\mathcal{C}^q \rightarrow s\mathcal{B}$ vanishes. We call ψ an A -functor if it commutes with the differential, that is,

$$(b \boxtimes 1 \boxtimes \dots \boxtimes 1 + 1 \boxtimes b \boxtimes \dots \boxtimes 1 + \dots + 1 \boxtimes 1 \boxtimes \dots \boxtimes b)\psi = \psi b.$$

For example, the map $\alpha : T^{\leq K} s\mathcal{A} \boxtimes T^{\leq N} sA_K(\mathcal{A}, \mathcal{B}) \rightarrow T^{\leq K+N} s\mathcal{B}$, $a \boxtimes r^1 \otimes \dots \otimes r^n \mapsto a.[(r^1 \otimes \dots \otimes r^n)\theta]$, is an A -functor.

1.6 Proposition (cf. Proposition 5.5 of [Lyu03]). *Let \mathcal{A} be an A_K -category, let \mathcal{C}^t be an A_{L^t} -category for $1 \leq t \leq q$, and let \mathcal{B} be an A_N -category, where $N \geq K + L^1 + \dots + L^q$. For any A -functor $\phi : T^{\leq K} s\mathcal{A} \boxtimes T^{\leq L^1} s\mathcal{C}^1 \boxtimes \dots \boxtimes T^{\leq L^q} s\mathcal{C}^q \rightarrow T^{\leq N} s\mathcal{B}$ there is a unique A -functor $\psi : T^{\leq L^1} s\mathcal{C}^1 \boxtimes \dots \boxtimes T^{\leq L^q} s\mathcal{C}^q \rightarrow T^{\leq N-K} sA_K(\mathcal{A}, \mathcal{B})$, such that*

$$\phi = (T^{\leq K} s\mathcal{A} \boxtimes T^{\leq L^1} s\mathcal{C}^1 \boxtimes \dots \boxtimes T^{\leq L^q} s\mathcal{C}^q \xrightarrow{1 \boxtimes \psi} T^{\leq K} s\mathcal{A} \boxtimes T^{\leq N-K} sA_K(\mathcal{A}, \mathcal{B}) \xrightarrow{\alpha} T^{\leq N} s\mathcal{B}).$$

Let \mathcal{A} be an A_N -category, let \mathcal{B} be an A_{N+K} -category, and let \mathcal{C} be an A_{N+K+L} -category. The above proposition implies the existence of an A -functor (cf. [Lyu03, Proposition 4.1])

$$M : T^{\leq K} sA_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leq L} sA_{N+K}(\mathcal{B}, \mathcal{C}) \rightarrow T^{\leq K+L} sA_N(\mathcal{A}, \mathcal{C}),$$

in particular, $(1 \boxtimes B + B \boxtimes 1)M = MB$. It has the components

$$M_{nm} = M|_{T^n \boxtimes T^m} \text{pr}_1 : T^n sA_N(\mathcal{A}, \mathcal{B}) \boxtimes T^m sA_{N+K}(\mathcal{B}, \mathcal{C}) \rightarrow sA_N(\mathcal{A}, \mathcal{C}),$$

$n \leq K$, $m \leq L$. We have $M_{00} = 0$ and $M_{nm} = 0$ for $m > 1$. If $m = 0$ and n is positive, M_{n0} is given by the formula:

$$M_{n0} : sA_N(\mathcal{A}, \mathcal{B})(f^0, f^1) \otimes \dots \otimes sA_N(\mathcal{A}, \mathcal{B})(f^{n-1}, f^n) \boxtimes \mathbb{k}_{g^0} \rightarrow sA_N(\mathcal{A}, \mathcal{C})(f^0 g^0, f^n g^0),$$

$$r^1 \otimes \dots \otimes r^n \boxtimes 1 \mapsto (r^1 \otimes \dots \otimes r^n | g^0)M_{n0},$$

$$[(r^1 \otimes \dots \otimes r^n | g^0)M_{n0}]_k = \sum_{l=n}^{n+k} (r^1 \otimes \dots \otimes r^n) \theta_{kl} g_l^0, \quad k \leq N,$$

where $|$ separates the arguments in place of \boxtimes . If $m = 1$, then M_{n1} is given by the formula:

$$M_{n1} : sA_N(\mathcal{A}, \mathcal{B})(f^0, f^1) \otimes \dots \otimes sA_N(\mathcal{A}, \mathcal{B})(f^{n-1}, f^n) \boxtimes sA_{N+K}(\mathcal{B}, \mathcal{C})(g^0, g^1)$$

$$\rightarrow sA_N(\mathcal{A}, \mathcal{C})(f^0 g^0, f^n g^1), \quad r^1 \otimes \dots \otimes r^n \boxtimes t^1 \mapsto (r^1 \otimes \dots \otimes r^n \boxtimes t^1)M_{n1},$$

$$[(r^1 \otimes \dots \otimes r^n \boxtimes t^1)M_{n1}]_k = \sum_{l=n}^{n+k} (r^1 \otimes \dots \otimes r^n) \theta_{kl} t_l^1, \quad k \leq N.$$

Note that equations

$$[(r^1 \otimes \dots \otimes r^n)B_n]_k = [(r^1 \otimes \dots \otimes r^n \boxtimes b)M_{n1}]_k - (-)^{r^1 + \dots + r^n} [(b \boxtimes r^1 \otimes \dots \otimes r^n)M_{1n}]_k$$

imply that

$$(r^1 \otimes \cdots \otimes r^n)B_n = (r^1 \otimes \cdots \otimes r^n \boxtimes b)M_{n1} - (-)^{r^1+\cdots+r^n}(b \boxtimes r^1 \otimes \cdots \otimes r^n)M_{1n},$$

$$B = (1 \boxtimes b)M - (b \boxtimes 1)M : \text{id} \rightarrow \text{id} : A_N(\mathcal{A}, \mathcal{B}) \rightarrow A_N(\mathcal{A}, \mathcal{B}).$$

Proposition 1.6 implies the existence of a unique A_L -functor

$$A_N(\mathcal{A}, -) : A_{N+K}(\mathcal{B}, \mathcal{C}) \rightarrow A_K(A_N(\mathcal{A}, \mathcal{B}), A_N(\mathcal{A}, \mathcal{C})),$$

such that

$$M = [T^{\leq K} sA_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leq L} sA_{N+K}(\mathcal{B}, \mathcal{C}) \xrightarrow{1 \boxtimes A_N(\mathcal{A}, -)} T^{\leq K} sA_N(\mathcal{A}, \mathcal{B}) \boxtimes T^{\leq L} sA_K(A_N(\mathcal{A}, \mathcal{B}), A_N(\mathcal{A}, \mathcal{C})) \xrightarrow{\alpha} T^{\leq K+L} sA_N(\mathcal{A}, \mathcal{C})].$$

The A_L -functor $A_N(\mathcal{A}, -)$ is strict, cf. [Lyu03, Proposition 6.2].


Let \mathcal{A} be an A_N -category, and let \mathcal{B} be a unital A_∞ -category with a unit transformation $\mathbf{i}^{\mathcal{B}}$. Then $A_N(\mathcal{A}, \mathcal{B})$ is a unital A_∞ -category with the unit transformation $(1 \boxtimes \mathbf{i}^{\mathcal{B}})M$ (cf. [Lyu03, Proposition 7.7]). The unit element for an object $f \in \text{Ob } A_N(\mathcal{A}, \mathcal{B})$ is $f \mathbf{i}_0^{A_N(\mathcal{A}, \mathcal{B})} : \mathbb{k} \rightarrow (sA_N)^{-1}(\mathcal{A}, \mathcal{B}), 1 \mapsto f \mathbf{i}^{\mathcal{B}}$.

When \mathcal{A} is an A_K -category and $N < K$, we may forget part of its structure and view \mathcal{A} as an A_N -category. If furthermore, \mathcal{B} is an A_{K+L} -category, we have the restriction strict A_L -functor $\text{restr}_{K,N} : A_K(\mathcal{A}, \mathcal{B}) \rightarrow A_N(\mathcal{A}, \mathcal{B})$. To prove the results mentioned above, we notice that they are restrictions of their A_∞ -analogs to finite N . Since the proofs of A_∞ -results are obtained in [Lyu03] by induction, an inspection shows that the proofs of the above A_N -statements are obtained as a byproduct.

1.7. Trees. Since the notions related to trees might be interpreted with some variations, we give precise definitions and fix notation. A *tree* is a non-empty connected graph without cycles. A vertex which belongs to only one edge is called *external*, other vertices are *internal*. A *plane tree* is a tree equipped for each internal vertex v with a cyclic ordering of the set E_v of edges, adjacent to v . Plane trees can be drawn on an oriented plane in a unique way (up to an ambient isotopy) so that the cyclic ordering of each E_v agrees with the orientation of the plane. An external vertex distinct from the root is called *input vertex*.

A *rooted tree* is a tree with a distinguished external vertex, called *root*. The set of vertices $V(t)$ of a rooted tree t has a canonical ordering: $x \preccurlyeq y$ iff the minimal path connecting the root with y contains x . A *linearly ordered tree* is a rooted tree t equipped with a linear order \leq of the set of internal vertices $IV(t)$, such that for all internal vertices x, y the relation $x \preccurlyeq y$ implies $x \leq y$. For each vertex $v \in V(t) - \{\text{root}\}$ of a rooted tree, the set E_v has a distinguished element e_v – the beginning of a minimal path from v to the root. Therefore, for each vertex $v \in V(t) - \{\text{root}\}$ of a rooted plane tree, the set E_v admits a unique linear order $<$, for which e_v is minimal and the induced cyclic order is the given one. An internal vertex v has degree d , if $\text{Card}(E_v) = d + 1$.

For any $y \in V(t)$ let $P_y = \{x \in V(t) \mid x \preccurlyeq y\}$. With each plane rooted tree t is associated a linearly ordered tree $t_{\leq} = (t, \leq)$ as follows. If $x, y \in IV(t)$ are such that $x \not\preccurlyeq y$ and $y \not\preccurlyeq x$, then $P_x \cap P_y = P_z$ for a unique $z \in IV(t)$, distinct from x and y . Let $a \in E_z - \{e_z\}$ (resp. $b \in E_z - \{e_z\}$) be the beginning of the minimal path connecting z and x (resp. y). If $a < b$, we set $x < y$. Graphically we $<$ -order the internal vertices by height. Thus, an internal vertex x on the left is depicted lower than a \preccurlyeq -incomparable internal vertex y on the right:



A *forest* is a sequence of plane rooted trees. Concatenation of forests is denoted \sqcup . The vertical composition $F_1 \cdot F_2$ of forests F_1, F_2 is well-defined if the sum of lengths of sequences F_1 and F_2 equals the number of external vertices of F_2 . These operations allow to construct any tree from elementary ones

$$1 = \begin{array}{c} | \\ | \\ | \end{array}, \quad \text{and} \quad \mathbf{t}_k = \begin{array}{c} \cdots \\ \diagup \quad \diagdown \\ | \end{array} \quad (k \text{ input vertices}).$$

Namely, any linearly ordered tree (t, \leq) has a unique presentation of the form

$$(t, \leq) = (1^{\sqcup \alpha_1} \sqcup \mathbf{t}_{k_1} \sqcup 1^{\sqcup \beta_1}) \cdot (1^{\sqcup \alpha_2} \sqcup \mathbf{t}_{k_2} \sqcup 1^{\sqcup \beta_2}) \cdot \dots \cdot \mathbf{t}_{k_N}, \quad (1.7.1)$$

where $N = |t| \stackrel{\text{def}}{=} \text{Card}(IV(t))$ is the number of internal vertices. Here

$$1^{\sqcup \alpha} \sqcup \mathbf{t}_k \sqcup 1^{\sqcup \beta} = \begin{array}{c} \overbrace{\begin{array}{|c|c|c|} \hline | & \dots & | \\ \hline \end{array}}^{\alpha} \quad \begin{array}{c} \overbrace{\begin{array}{c} \cdots \\ \diagup \quad \diagdown \\ | \end{array}}^k \\ \end{array} \quad \begin{array}{c} \overbrace{\begin{array}{|c|c|c|} \hline | & \dots & | \\ \hline \end{array}}^{\beta} \end{array}.$$

In (1.7.1) the highest vertex is indexed by 1, the lowest – by N .

2. Properties of free A_∞ -categories

2.1. Construction of a free A_∞ -category. The category $\text{strict}A_\infty$ has A_∞ -categories as objects and strict A_∞ -functors as morphisms. There is a functor $\mathcal{U} : \text{strict}A_\infty \rightarrow A_1$, $\mathcal{A} \mapsto (\mathcal{A}, m_1)$ which sends an A_∞ -category to the underlying differential graded \mathbb{k} -quiver, forgetting all higher multiplications. Following Kontsevich and Soibelman [KS02] we are going to prove that \mathcal{U} has a left adjoint functor $\mathcal{F} : A_1 \rightarrow \text{strict}A_\infty$, $\mathcal{Q} \mapsto \mathcal{F}\mathcal{Q}$. The A_∞ -category $\mathcal{F}\mathcal{Q}$ is called free. Below we describe its structure for an arbitrary differential graded \mathbb{k} -quiver \mathcal{Q} . We shall work with its shift $(s\mathcal{Q}, d)$.

Let us define an A_∞ -category \mathcal{FQ} via the following data. The class of objects $\text{Ob } \mathcal{FQ}$ is $\text{Ob } \mathcal{Q}$. The \mathbb{Z} -graded \mathbb{k} -modules of morphisms between $X, Y \in \text{Ob } \mathcal{Q}$ are

$$\begin{aligned} s\mathcal{FQ}(X, Y) &= \bigoplus_{n \geq 1} \bigoplus_{t \in \mathcal{T}_{\geq 2}^n} s\mathcal{F}_t\mathcal{Q}(X, Y), \\ s\mathcal{F}_t\mathcal{Q}(X, Y) &= \bigoplus_{\substack{X_0=X, X_n=Y \\ X_0, \dots, X_n \in \text{Ob } \mathcal{Q}}} s\mathcal{Q}(X_0, X_1) \otimes \cdots \otimes s\mathcal{Q}(X_{n-1}, X_n)[-|t|], \end{aligned}$$

where $\mathcal{T}_{\geq 2}^n$ is the class of plane rooted trees with $n+1$ external vertices, such that $\text{Card}(E_v) \geq 3$ for all $v \in IV(t)$. We use the following convention: if M, N are (differential) graded \mathbb{k} -modules, then¹

$$\begin{aligned} (M \otimes N)[k] &= M \otimes (N[k]), \\ (M \otimes N \xrightarrow{s^k} (M \otimes N)[k]) &= (M \otimes N \xrightarrow{1 \otimes s^k} M \otimes (N[k])). \end{aligned}$$

The quiver \mathcal{FQ} is equipped with the following operations. For $k > 1$ the operation b_k is a direct sum of maps

$$b_k = s^{|t_1|} \otimes \cdots \otimes s^{|t_{k-1}|} \otimes s^{|t_k| - |t|} : s\mathcal{F}_{t_1}\mathcal{Q}(Y_0, Y_1) \otimes \cdots \otimes s\mathcal{F}_{t_k}\mathcal{Q}(Y_{k-1}, Y_k) \rightarrow s\mathcal{F}_t\mathcal{Q}(Y_0, Y_k), \quad (2.1.1)$$

where $t = (t_1 \sqcup \cdots \sqcup t_k) \cdot t_k$. In particular, $|t| = |t_1| + \cdots + |t_k| + 1$. The operation b_1 restricted to $s\mathcal{F}_t\mathcal{Q}$ is

$$b_1 = d \oplus (-1)^{\beta(t')} s^{-1} : s\mathcal{F}_t\mathcal{Q}(X, Y) \rightarrow s\mathcal{F}_t\mathcal{Q}(X, Y) \oplus \bigoplus_{t'=t+\text{edge}} s\mathcal{F}_{t'}\mathcal{Q}(X, Y), \quad (2.1.2)$$

where the sum extends over all trees $t' \in \mathcal{T}_{\geq 2}^n$ with a distinguished edge e , such that contracting e we get t from t' . The sign is determined by

$$\beta(t') = \beta(t', e) = 1 + h(\text{highest vertex of } e),$$

where an isomorphism of ordered sets

$$h : IV(t'_<) \xrightarrow{\sim} [1, |t'|] \cap \mathbb{Z}$$

is simply the height of a vertex in the linearly ordered tree $t'_<$, canonically associated with t' . In (2.1.2) d means $d \otimes 1 \otimes \cdots \otimes 1 + \cdots + 1 \otimes \cdots \otimes d \otimes 1 + 1 \otimes \cdots \otimes 1 \otimes d$, where the last d is $d_{s\mathcal{Q}[-|t|]} = (-1)^{|t|} s^{|t|} \cdot d_{s\mathcal{Q}} \cdot s^{-|t|}$, as usual. According to our conventions, s^{-1} in (2.1.2) means $1^{\otimes n-1} \otimes s^{-1}$.

2.2 Proposition. \mathcal{FQ} is an A_∞ -category.

Proof. First we prove that $b_1^2 = 0$ on $s\mathcal{F}_t\mathcal{Q}$. Indeed,

$$\begin{aligned} b_1^2 &= d^2 \oplus (-1)^{\beta(t', e)} (s^{-1}d + ds^{-1}) \oplus [(-1)^{\beta(t'_1, e_1) + \beta(t'', e_2)} + (-1)^{\beta(t'_2, e_2) + \beta(t'', e_1)}] s^{-2} : \\ &\quad s\mathcal{F}_t\mathcal{Q} \rightarrow s\mathcal{F}_t\mathcal{Q} \oplus \bigoplus_{t'=t+e} s\mathcal{F}_{t'}\mathcal{Q} \oplus \bigoplus_{t''=t+e_1+e_2} s\mathcal{F}_{t''}\mathcal{Q}, \end{aligned}$$

¹Another gauge choice $(M \otimes N)[1] = M[1] \otimes N$, $s = s \otimes 1$ seems less convenient.

where t'' contains two distinguished edges e_1, e_2 , contraction along which gives t ; t'_2 is t'' contracted along e_1 , and t'_1 is t'' contracted along e_2 . We may assume that highest vertex of e_1 is lower than highest vertex of e_2 in t'' . Then $\beta(t'_1, e_1) = \beta(t'', e_1)$ and $\beta(t'', e_2) = \beta(t'_2, e_2) + 1$, hence,

$$(-1)^{\beta(t'_1, e_1) + \beta(t'', e_2)} + (-1)^{\beta(t'_2, e_2) + \beta(t'', e_1)} = 0.$$

Obviously, $d^2 = 0$ and $s^{-1}d + ds^{-1} = 0$, hence, $b_1^2 = 0$.

Let us prove for each $n > 1$ that

$$b_n b_1 + \sum_{p=1}^n (1^{\otimes p-1} \otimes b_1 \otimes 1^{\otimes n-p}) b_n + \sum_{\substack{\alpha+k+\beta=n \\ \alpha+\beta>0}}^{k>1} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) b_{\alpha+1+\beta} = 0 :$$

$$s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \rightarrow s\mathcal{F}_t\mathcal{Q} \oplus \bigoplus_{p, t'_p} s\mathcal{F}_{t'_p}\mathcal{Q} \oplus \bigoplus_{\substack{k>1, t'' \\ \alpha+k+\beta=n \\ \alpha+\beta>0}} s\mathcal{F}_{t''}\mathcal{Q},$$

where

$$\begin{aligned} t &= (t_1 \sqcup \cdots \sqcup t_n) \cdot \mathbf{t}_n = \begin{array}{c} \boxed{t_1} \quad \boxed{t_2} \quad \cdots \quad \boxed{t_{n-1}} \quad \boxed{t_n} \\ \diagdown \quad \diagup \quad \quad \quad \diagdown \quad \diagup \\ \bullet \\ \downarrow \end{array}, \\ t' &= (t_1 \sqcup \cdots \sqcup t'_p \sqcup \cdots \sqcup t_n) \cdot \mathbf{t}_n = \begin{array}{c} \boxed{t_1} \quad \boxed{t'_p} \quad \boxed{t_n} \\ \diagdown \quad \diagup \quad \quad \quad \diagdown \quad \diagup \\ \bullet \\ \downarrow \end{array}, \end{aligned} \quad (2.2.1)$$

$$\begin{aligned} t'' &= (t_1 \sqcup \cdots \sqcup t_n) \cdot (1^{\sqcup \alpha} \sqcup \mathbf{t}_k \sqcup 1^{\sqcup \beta}) \cdot \mathbf{t}_{\alpha+1+\beta} \\ &= \begin{array}{c} \boxed{t_1} \quad \boxed{t_\alpha} \quad \boxed{t_{\alpha+1}} \quad \boxed{t_{\alpha+k}} \quad \boxed{t_{\alpha+k+1}} \quad \boxed{t_n} \\ \diagdown \quad \diagup \quad \quad \quad \diagdown \quad \diagup \\ \bullet \\ \downarrow \end{array}, \end{aligned}$$

and contraction of t'_p along distinguished edge e_p gives t_p . According to the three types of summands in the target, the required equation follows from anticommutativity of the following three diagrams:

$$\begin{array}{ccc} s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow[s_n]{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} & s\mathcal{F}_t\mathcal{Q} \\ \downarrow 1^{\otimes n-1} \otimes d + \cdots + d \otimes 1^{\otimes n-1} & \quad \quad \quad \downarrow d & \\ s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow[s_n]{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} & s\mathcal{F}_t\mathcal{Q} \end{array}$$

that is,

$$\begin{aligned} &(1^{\otimes n-1} \otimes d + \cdots + d \otimes 1^{\otimes n-1})(s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}) \\ &\quad + (s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|})(1^{\otimes n-1} \otimes d + \cdots + d \otimes 1^{\otimes n-1}) = 0; \end{aligned}$$

$$\begin{array}{ccc}
s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_p}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} & s\mathcal{F}_t\mathcal{Q} \\
\downarrow (-)^{\beta(t'_p)} 1^{\otimes p-1} \otimes s^{-1} \otimes 1^{\otimes n-p} & & \downarrow (-)^{1+|t_1|+\cdots+|t_{p-1}|+\beta(t'_p)} s^{-1} \\
s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t'_p}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{p-1}|} \otimes s^{|t_p|+1} \otimes s^{|t_{p+1}|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}} & s\mathcal{F}_{t'}\mathcal{Q}
\end{array}$$

that is,

$$\begin{aligned}
& (-1)^{\beta(t'_p)} (1^{\otimes p-1} \otimes s^{-1} \otimes 1^{\otimes n-p}) (s^{|t_1|} \otimes \cdots \otimes s^{|t_{p-1}|} \otimes s^{|t_p|+1} \otimes s^{|t_{p+1}|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}) \\
& + (s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}) (-1)^{1+|t_1|+\cdots+|t_{p-1}|+\beta(t'_p)} (1^{\otimes n-1} \otimes s^{-1}) = 0,
\end{aligned}$$

in the particular case $p = n$ it holds as well;

$$\begin{array}{ccc}
s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}} & s\mathcal{F}_t\mathcal{Q} \\
\downarrow 1^{\otimes \alpha} \otimes s^{|t_{\alpha+1}|} \otimes \cdots \otimes s^{|t_{\alpha+k-1}|} \otimes s^{-|t_{\alpha+1}|-\cdots-|t_{\alpha+k-1}|-1} \otimes 1^{\otimes \beta} & & \downarrow (-)^{1+|t_1|+\cdots+|t_{\alpha}|} s^{-1} \\
s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_{\alpha}}\mathcal{Q} \otimes s\mathcal{F}_{\hat{t}}\mathcal{Q} \otimes s\mathcal{F}_{t_{\alpha+k+1}}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow{s^{|t_1|} \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes s^{|t_{\alpha+k+1}|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}} & s\mathcal{F}_{t'}\mathcal{Q}
\end{array}$$

where $\hat{t} = (t_{\alpha+1} \sqcup \cdots \sqcup t_{\alpha+k}) \cdot t_k$, that is,

$$\begin{aligned}
& (1^{\otimes \alpha} \otimes s^{|t_{\alpha+1}|} \otimes \cdots \otimes s^{|t_{\alpha+k-1}|} \otimes s^{-|t_{\alpha+1}|-\cdots-|t_{\alpha+k-1}|-1} \otimes 1^{\otimes \beta}). \\
& (s^{|t_1|} \otimes \cdots \otimes s^{|t_{\alpha}|} \otimes 1^{\otimes k-1} \otimes s^{|t_{\alpha+1}|+\cdots+|t_{\alpha+k}|+1} \otimes s^{|t_{\alpha+k+1}|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|-1}) \\
& + (s^{|t_1|} \otimes \cdots \otimes s^{|t_{n-1}|} \otimes s^{|t_n|-|t|}) (-1)^{1+|t_1|+\cdots+|t_{\alpha}|} (1^{\otimes n-1} \otimes s^{-1}) = 0,
\end{aligned}$$

in the particular case $\beta = 0$ it holds as well.

Therefore, \mathcal{FQ} is an A_{∞} -category. \square

Let us establish a property of free A_{∞} -categories, which explains why they are called free.

2.3 Proposition (A_{∞} -functors from a free A_{∞} -category). *Let \mathcal{Q} be a differential graded quiver, and let \mathcal{A} be an A_{∞} -category. Let $f_1 : s\mathcal{Q} \rightarrow (s\mathcal{A}, b_1)$ be a chain morphism of differential graded quivers with the underlying mapping of objects $\text{Ob } f : \text{Ob } \mathcal{Q} \rightarrow \text{Ob } \mathcal{A}$. Suppose given \mathbb{k} -quiver morphisms $f_k : T^k s\mathcal{FQ} \rightarrow s\mathcal{A}$ of degree 0 with the same underlying map $\text{Ob } f$ for all $k > 1$. Then there exists a unique extension of f_1 to a quiver morphism $f_1 : s\mathcal{FQ} \rightarrow s\mathcal{A}$ such that (f_1, f_2, \dots) are components of an A_{∞} -functor $f : \mathcal{FQ} \rightarrow \mathcal{A}$.*

Proof. For each $n > 1$ we have to satisfy the equation

$$b_n f_1 = \sum_{i_1+\cdots+i_l=n} (f_{i_1} \otimes \cdots \otimes f_{i_l}) b_l - \sum_{\alpha+\beta>0}^{\alpha+k+\beta=n} (1^{\otimes \alpha} \otimes b_k \otimes 1^{\otimes \beta}) f_{\alpha+1+\beta} : T^n s\mathcal{FQ} \rightarrow s\mathcal{A}. \quad (2.3.1)$$

It is used to define recursively f_1 on $s\mathcal{F}\mathcal{Q}$. Suppose that t_1, \dots, t_n are trees, $n > 1$, and $f_1 : s\mathcal{F}_{t_i}\mathcal{Q} \rightarrow s\mathcal{A}$ is already defined for all $1 \leq i \leq n$. Since

$$b_n = s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n| - |t|} : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \dots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \rightarrow s\mathcal{F}_t\mathcal{Q}$$

is invertible for $t = (t_1 \sqcup \dots \sqcup t_n) \cdot \mathbf{t}_n$, formula (2.3.1) determines $f_1 : s\mathcal{F}_t\mathcal{Q} \rightarrow s\mathcal{A}$ uniquely as

$$f_1 = (s\mathcal{F}_t\mathcal{Q} \xrightarrow{b_n^{-1}} s\mathcal{F}_{t_1}\mathcal{Q} \otimes \dots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \xrightarrow{\sum (f_{i_1} \otimes \dots \otimes f_{i_l}) b_l - \sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) f_{\alpha+1+\beta}} s\mathcal{A}).$$

This proves uniqueness of the extension of f_1 .

Let us prove that the cocategory homomorphism f with so defined components (f_1, f_2, \dots) is an A_∞ -functor. Equations (2.3.1) are satisfied by construction of f_1 . So it remains to prove that f_1 is a chain map. Equation $f_1 b_1 = b_1 f_1$ holds on $s\mathcal{F}_|\mathcal{Q}$ by assumption. We are going to prove by induction on $|t|$ that it holds on $s\mathcal{F}_t\mathcal{Q}$. Considering $t = (t_1 \sqcup \dots \sqcup t_n) \cdot \mathbf{t}_n$, $n > 1$, we assume that $f_1 b_1 = b_1 f_1 : s\mathcal{F}_{t'}\mathcal{Q} \rightarrow s\mathcal{A}$ for all trees t' with $|t'| < |t|$. To prove that $f_1 b_1 = b_1 f_1 : s\mathcal{F}_t\mathcal{Q} \rightarrow s\mathcal{A}$ it suffices to show that $b_n f_1 b_1 = b_n b_1 f_1$ for all $n > 1$ due to invertibility of b_n . Using (2.3.1) and the equation $b^2 \text{pr}_1 = 0$ we find

$$\begin{aligned} b_n f_1 b_1 - b_n b_1 f_1 &= \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) b_l b_1 - \sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) f_{\alpha+1+\beta} b_1 \\ &\quad + \sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) b_{\alpha+1+\beta} f_1 \\ &= - \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) \sum_{\gamma+p+\delta=l}^{\gamma+\delta>0} (1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta}) b_{\gamma+1+\delta} \\ &\quad + \sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \left[\sum_{j_1 + \dots + j_r = \alpha+1+\beta}^{r>1} (f_{j_1} \otimes \dots \otimes f_{j_r}) b_r \right. \\ &\quad \left. - \sum_{\gamma+p+\delta=\alpha+1+\beta}^{\gamma+\delta>0} (1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta}) f_{\gamma+1+\delta} \right] \\ &= \sum_{r>1} \left[\sum_{\alpha+k+\beta=n}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \sum_{j_1 + \dots + j_r = \alpha+1+\beta} f_{j_1} \otimes \dots \otimes f_{j_r} \right. \\ &\quad \left. - \sum_{i_1 + \dots + i_l = n} (f_{i_1} \otimes \dots \otimes f_{i_l}) \sum_{\substack{\gamma+p+\delta=l \\ \gamma+1+\delta=r}} 1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta} \right] b_r \\ &\quad - \sum_{r>1} \left[\sum_{\substack{\alpha+k+\beta=n \\ \alpha+\beta>0}} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \sum_{\substack{\gamma+p+\delta=\alpha+1+\beta \\ \gamma+1+\delta=r}} 1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta} \right] f_r. \end{aligned}$$

Let us show that the expressions in square brackets vanish. The first one is the matrix coefficient $bf - fb : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \rightarrow T^r s\mathcal{A}$. Indeed, for $r > 1$ the inequality $r \leq j_1 + \cdots + j_r = \alpha + 1 + \beta$ automatically implies that $\alpha + \beta > 0$, so this condition can be omitted. Using the induction hypothesis one can transform the left hand side of equation

$$\begin{aligned} & \sum_{\alpha+k+\beta=n} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \sum_{j_1+\cdots+j_r=\alpha+1+\beta} f_{j_1} \otimes \cdots \otimes f_{j_r} \\ &= \sum_{i_1+\cdots+i_l=n} (f_{i_1} \otimes \cdots \otimes f_{i_l}) \sum_{\substack{\gamma+p+\delta=l \\ \gamma+1+\delta=r}} 1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta} : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_n}\mathcal{Q} \rightarrow T^r s\mathcal{A} \end{aligned}$$

into the right hand side for all $n, r \geq 1$.

The second expression

$$\sum_{\substack{\alpha+k+\beta=n \\ \alpha+\beta>0}} (1^{\otimes\alpha} \otimes b_k \otimes 1^{\otimes\beta}) \sum_{\substack{\gamma+p+\delta=\alpha+1+\beta \\ \gamma+1+\delta=r}} 1^{\otimes\gamma} \otimes b_p \otimes 1^{\otimes\delta} \quad (2.3.2)$$

is the matrix coefficient

$$(b - b \operatorname{pr}_1) b \operatorname{pr}_r : T^n s\mathcal{F}\mathcal{Q} \rightarrow T^r s\mathcal{F}\mathcal{Q}$$

of the endomorphism $(b - b \operatorname{pr}_1) b : T s\mathcal{F}\mathcal{Q} \rightarrow T s\mathcal{F}\mathcal{Q}$. However,

$$(b - b \operatorname{pr}_1) b \operatorname{pr}_r = b^2 \operatorname{pr}_r - b \operatorname{pr}_1 b \operatorname{pr}_r = -b \operatorname{pr}_1 b \operatorname{pr}_1 \operatorname{pr}_r = 0$$

for $r > 1$, because $\operatorname{pr}_1 b = \operatorname{pr}_1 b \operatorname{pr}_1$. Therefore, (2.3.2) vanishes and equation $b_n f_1 b_1 = b_n b_1 f_1$ is proven. \square

Let $\operatorname{strict} A_\infty(\mathcal{F}\mathcal{Q}, \mathcal{A}) \subset A_\infty(\mathcal{F}\mathcal{Q}, \mathcal{A})$ be a full A_∞ -subcategory, whose objects are strict A_∞ -functors. Recall that $\operatorname{Ob} A_1(\mathcal{Q}, \mathcal{A})$ is the set of chain morphisms $\mathcal{Q} \rightarrow \mathcal{A}$ of differential graded quivers.

2.4 Corollary. *A chain morphism $f : \mathcal{Q} \rightarrow \mathcal{A}$ admits a unique extension to a strict A_∞ -functor $\hat{f} : \mathcal{F}\mathcal{Q} \rightarrow \mathcal{A}$. The maps $f \mapsto \hat{f}$ and*

$$\operatorname{restr} : \operatorname{Ob} \operatorname{strict} A_\infty(\mathcal{F}\mathcal{Q}, \mathcal{A}) \rightarrow \operatorname{Ob} A_1(\mathcal{Q}, \mathcal{A}), \quad g \mapsto (\operatorname{Ob} g, g_1|_{s\mathcal{Q}})$$

are inverse to each other.

Indeed, strict A_∞ -functors g are distinguished by conditions $g_k = 0$ for $k > 1$.

We may view $\operatorname{strict} A_\infty$ as a category, whose objects are A_∞ -categories and morphisms are strict A_∞ -functors. We may also view A_1 as a category consisting of differential graded quivers and their morphisms. There is a functor $\mathcal{U} : \operatorname{strict} A_\infty \rightarrow A_1$, $\mathcal{A} \mapsto (\mathcal{A}, m_1)$, which sends an A_∞ -category to the underlying differential graded \mathbb{k} -quiver, forgetting all higher multiplications. The restriction map

$$\operatorname{restr} : \operatorname{strict} A_\infty(\mathcal{F}\mathcal{Q}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{U}\mathcal{A}), \quad g \mapsto (\operatorname{Ob} g, g_1|_{s\mathcal{Q}}) \quad (2.4.1)$$

is functorial in \mathcal{A} .

2.5 Corollary. *There is a functor $\mathcal{F} : A_1 \rightarrow \operatorname{strict} A_\infty$, $\mathcal{Q} \mapsto \mathcal{F}\mathcal{Q}$, left adjoint to \mathcal{U} .*

2.6. Explicit formula for the constructed strict A_∞ -functor. Let us obtain a more explicit formula for $\widehat{f}_1|_{s\mathcal{F}_t\mathcal{Q}}$. We define \widehat{f}_1 for $t = (t_1 \sqcup \dots \sqcup t_n) \cdot \mathbf{t}_n$ recursively by a commutative diagram

$$\begin{array}{ccc} s\mathcal{F}_{t_1}\mathcal{Q} \otimes \dots \otimes s\mathcal{F}_{t_n}\mathcal{Q} & \xrightarrow{s^{|t_1|} \otimes \dots \otimes s^{|t_{n-1}|} \otimes s^{|t_n| - |t|}} & s\mathcal{F}_t\mathcal{Q} \\ \widehat{f}_1^{\otimes n} \downarrow & \stackrel{\text{def}}{=} & \downarrow \widehat{f}_1 \\ s\mathcal{A} \otimes \dots \otimes s\mathcal{A} & \xrightarrow{b_n} & s\mathcal{A} \end{array}$$

Notice that the top map is invertible. Here n, t_1, \dots, t_n are uniquely determined by decomposition (2.2.1) of t .

Let (t, \leq) be a linearly ordered tree with the underlying given plane rooted tree t . Decompose (t, \leq) into a vertical composition of forests as in (1.7.1). Then the following diagram commutes

$$\begin{array}{ccccccc} s\mathcal{Q}^{\otimes n} & \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} & s\mathcal{Q}^{\otimes \alpha_1} \otimes s\mathcal{F}_{t_{k_1}}\mathcal{Q} \otimes s\mathcal{Q}^{\otimes \beta_1} & \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} & \dots & \xrightarrow{b_{k_N}} & s\mathcal{F}_t\mathcal{Q} \\ f_1^{\otimes n} \downarrow & & \widehat{f}_1^{\otimes \alpha_1 + 1 + \beta_1} \downarrow & & \widehat{f}_1^{\otimes \alpha_2 + 1 + \beta_2} \downarrow & \dots & \widehat{f}_1^{\otimes k_N} \downarrow \\ s\mathcal{A}^{\otimes n} & \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} & s\mathcal{A}^{\otimes \alpha_1} \otimes s\mathcal{A} \otimes s\mathcal{A}^{\otimes \beta_1} & \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} & \dots & \xrightarrow{b_{k_N}} & s\mathcal{A} \end{array}$$

The upper row consists of invertible maps. One can prove by induction that the composition of maps in the upper row equals $\pm s^{-|t|}$. When $(t, \leq) = t_<$ is the linearly ordered tree, canonically associated with t , then the composition of maps in the upper row equals $s^{-|t|}$. This is also proved by induction: if t is presented as $t = (t_1 \sqcup \dots \sqcup t_k) \cdot \mathbf{t}_k$, then the composition of maps in the upper row is

$$\begin{aligned} (1^{\otimes k-1} \otimes s^{-|t_k|})(1^{\otimes k-2} \otimes s^{-|t_{k-1}|} \otimes 1) \dots (s^{-|t_1|} \otimes 1^{\otimes k-1})(s^{|t_1|} \otimes \dots \otimes s^{|t_{k-1}|} \otimes s^{|t_k| - |t|}) \\ = 1^{\otimes k-1} \otimes s^{-|t|} = s^{-|t|}. \end{aligned}$$

Therefore, for an arbitrary tree $t \in \mathcal{T}_{\geq 2}^n$ the map $\widehat{f}_1|_{s\mathcal{F}_t\mathcal{Q}}$ is

$$\widehat{f}_1 = (s\mathcal{F}_t\mathcal{Q} \xrightarrow{s^{|t|}} s\mathcal{Q}^{\otimes n} \xrightarrow{f_1^{\otimes n}} s\mathcal{A}^{\otimes n} \xrightarrow{1^{\otimes \alpha_1} \otimes b_{k_1} \otimes 1^{\otimes \beta_1}} s\mathcal{A}^{\otimes \alpha_1 + 1 + \beta_1} \xrightarrow{1^{\otimes \alpha_2} \otimes b_{k_2} \otimes 1^{\otimes \beta_2}} \dots \xrightarrow{b_{k_N}} s\mathcal{A}),$$

where the factors correspond to decomposition (1.7.1) of $t_<$.

2.7. Transformations between functors from a free A_∞ -category. Let \mathcal{Q} be a differential graded quiver, and let \mathcal{A} be an A_∞ -category. Then $A_1(\mathcal{Q}, \mathcal{A})$ is an A_∞ -category as well. The differential graded quiver $(sA_1(\mathcal{Q}, \mathcal{A}), B_1)$ is described as follows. Objects are chain quiver maps $\phi : (s\mathcal{Q}, b_1) \rightarrow (s\mathcal{A}, b_1)$, the graded \mathbb{k} -module of morphisms $\phi \rightarrow \psi$ is the product of graded \mathbb{k} -modules

$$sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi) = \prod_{X \in \text{Ob } \mathcal{Q}} s\mathcal{A}(X\phi, X\psi) \times \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathcal{C}(s\mathcal{Q}(X, Y), s\mathcal{A}(X\phi, Y\psi)), \quad r = (r_0, r_1).$$

The differential B_1 is given by

$$\begin{aligned}(rB_1)_0 &= r_0b_1, \\ (rB_1)_1 &= r_1b_1 + (\phi_1 \otimes r_0)b_2 + (r_0 \otimes \psi_1)b_2 - (-)^r b_1 r_1.\end{aligned}\tag{2.7.1}$$

Restrictions $\phi, \psi : \mathcal{Q} \rightarrow \mathcal{A}$ of arbitrary A_∞ -functors $\phi, \psi : \mathcal{FQ} \rightarrow \mathcal{A}$ to \mathcal{Q} are A_1 -functors (chain quiver maps).

2.8 Proposition. *Let $\phi, \psi : \mathcal{FQ} \rightarrow \mathcal{A}$ be A_∞ -functors. For an arbitrary complex P of \mathbb{k} -modules chain maps $u : P \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ are in bijection with the following data: $(u', u_k)_{k>1}$*

1. a chain map $u' : P \rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)$,
2. \mathbb{k} -linear maps

$$u_k : P \rightarrow \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}((s\mathcal{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi))$$

of degree 0 for all $k > 1$.

The bijection maps u to $(u', u_k)_{k>1}$, where $u_k = u \cdot \text{pr}_k$ and

$$u' = (P \xrightarrow{u} sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_1(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)). \tag{2.8.1}$$

The inverse bijection can be recovered from the recurrent formula

$$\begin{aligned}(-)^p b_k^{\mathcal{FQ}}(pu_1) &= -(pd)u_k + \sum_{\alpha+q+c=k}^{\alpha, \beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}^{\mathcal{A}} \\ &\quad - (-)^p \sum_{\alpha+q+\beta=k}^{\alpha+\beta>0} (1^{\otimes \alpha} \otimes b_q^{\mathcal{FQ}} \otimes 1^{\otimes \beta})(pu_{\alpha+1+\beta}) : (s\mathcal{FQ})^{\otimes k} \rightarrow s\mathcal{A},\end{aligned}$$

where $k > 1$, $p \in P$, and $\phi_{a\alpha}, \psi_{c\beta}$ are matrix elements of ϕ, ψ .

Proof. Since the \mathbb{k} -module of (ϕ, ψ) -coderivations $sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ is a product, \mathbb{k} -linear maps $u : P \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ of degree 0 are in bijection with sequences of \mathbb{k} -linear maps $(u_k)_{k \geq 0}$ of degree 0:

$$\begin{aligned}u_0 : P &\rightarrow \prod_{X \in \text{Ob } \mathcal{Q}} s\mathcal{A}(X\phi, X\psi), & p &\mapsto pu_0, \\ u_k : P &\rightarrow \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}((s\mathcal{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi)), & p &\mapsto pu_k,\end{aligned}$$

for $k \geq 1$. The complex $\Phi_0 = (sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi), B_1)$ admits a filtration by subcomplexes

$$\Phi_n = 0 \times \cdots \times 0 \times \prod_{k=n}^{\infty} \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}((s\mathcal{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi)).$$

In particular, Φ_2 is a subcomplex, and

$$\Phi_0/\Phi_2 = \prod_{X \in \text{Ob } \mathcal{Q}} s\mathcal{A}(X\phi, X\psi) \times \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}(s\mathcal{FQ}(X, Y), s\mathcal{A}(X\phi, Y\psi))$$

is the quotient complex with differential (2.7.1). Since $(s\mathcal{FQ}, b_1)$ splits into a direct sum of two subcomplexes $s\mathcal{Q} \oplus (\oplus_{|t|>0} s\mathcal{F}_t\mathcal{Q})$, the complex Φ_0/Φ_2 has a subcomplex

$$(0 \times \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}(\oplus_{|t|>0} s\mathcal{F}_t\mathcal{Q}(X, Y), s\mathcal{A}(X\phi, Y\psi)), [-, b_1]).$$

The corresponding quotient complex is $sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)$. The resulting quotient map $\text{restr}_1 : sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)$ is the restriction map. Denoting $u' = u \cdot \text{restr}_1$, we get the discussed assignment $u \mapsto (u', u_n)_{n \geq 1}$. The claim is that if u is a chain map, then the missing part

$$u_1'' : P \rightarrow \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}(\oplus_{|t|>0} s\mathcal{F}_t\mathcal{Q}(X, Y), s\mathcal{A}(X\phi, Y\psi)),$$

of $u_1 = u_1' \times u_1''$ is recovered in a unique way.

Let us prove that the map $u \mapsto (u', u_n)_{n \geq 1}$ is injective. The chain map u satisfies $pdu = puB_1$ for all $p \in P$. That is, $pdu_k = (puB_1)_k$ for all $k \geq 0$. Since $puB_1 = (pu)b^A - (-)^p b^{\mathcal{FQ}}(pu)$, these conditions can be rewritten as

$$pdu_k = \sum_{\alpha+q+c=k}^{\alpha, \beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}^A - (-)^p \sum_{\alpha+q+\beta=k} (1^{\otimes \alpha} \otimes b_q \otimes 1^{\otimes \beta})(pu_{\alpha+1+\beta}), \quad (2.8.2)$$

where $\phi_{a\alpha} : T^a s\mathcal{FQ}(X, Y) \rightarrow T^\alpha s\mathcal{A}(X\phi, Y\phi)$ are matrix elements of ϕ , and $\psi_{c\beta}$ are matrix elements of ψ . The same formula can be rewritten as

$$\begin{aligned} (-)^p b_k^{\mathcal{FQ}}(pu_1) &= -(pd)u_k + \sum_{\alpha+q+c=k}^{\alpha, \beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}^A \\ &\quad - (-)^p \sum_{\alpha+q+\beta=k}^{\alpha+\beta>0} (1^{\otimes \alpha} \otimes b_q^{\mathcal{FQ}} \otimes 1^{\otimes \beta})(pu_{\alpha+1+\beta}) : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_k}\mathcal{Q} \rightarrow s\mathcal{A}. \end{aligned} \quad (2.8.3)$$

When $k > 1$, the map $b_k^{\mathcal{FQ}} : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \cdots \otimes s\mathcal{F}_{t_k}\mathcal{Q} \rightarrow s\mathcal{F}_t\mathcal{Q}$, $t = (t_1 \sqcup \cdots \sqcup t_k)\mathbf{t}_k$ is invertible, thus, $pu_1 : s\mathcal{F}_t\mathcal{Q} \rightarrow s\mathcal{A}$ in the left hand side is determined in a unique way by u_0, u_n for

$n > 1$ and by $pu_1 : s\mathcal{F}_{t_i}\mathcal{Q} \rightarrow s\mathcal{A}$, $1 \leq i \leq k$, occurring in the right hand side. Since the restriction u'_1 of u_1 to $s\mathcal{F}_1\mathcal{Q} = s\mathcal{Q}$ is known by 1), the map u''_1 is recursively recovered from $(u_0, u'_1, u_n)_{n>1}$.

Let us prove that the map $u \mapsto (u', u_n)_{n>1}$ is surjective. Given $(u_0, u'_1, u_n)_{n>1}$ we define maps u''_1 of degree 0 recursively by (2.8.3). This implies equation (2.8.2) for $k > 1$. For $k = 0$ this equation in the form $pdu_0 = pu_0b_1$ holds due to condition 1). It remains to prove equation (2.8.2) for $k = 1$:

$$(pd)u_1 = (pu_1)b_1^A + (\phi_1 \otimes pu_0)b_2^A + (pu_0 \otimes \psi_1)b_2^A - (-)^pb_1(pu_1) : s\mathcal{F}_t\mathcal{Q}(X, Y) \rightarrow s\mathcal{A}(X\phi, Y\psi) \quad (2.8.4)$$

for all trees $t \in \mathcal{T}_{\geq 2}$. For $t = |$ it holds due to assumption 1). Let $N > 1$ be an integer. Assume that equation (2.8.4) holds for all trees $t \in \mathcal{T}_{\geq 2}$ with the number of input leaves $\text{in}(t) < N$. Let $t \in \mathcal{T}_{\geq 2}^N$ be a tree (with $\text{in}(t) = N$). Then $t = (t_1 \sqcup \dots \sqcup t_k)t_k$ for some $k > 1$ and some trees $t_i \in \mathcal{T}_{\geq 2}$, $\text{in}(t_i) < N$. For such t equation (2.8.4) is equivalent to

$$\begin{aligned} (-)^pb_k(pd)u_1 &= (-)^pb_k(pu_1)b_1^A + (-)^pb_k(\phi_1 \otimes pu_0)b_2^A + (-)^pb_k(pu_0 \otimes \psi_1)b_2^A \\ &+ \sum_{\substack{\gamma+\delta>0 \\ \gamma+j+\delta=k}} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})b_{\gamma+1+\delta}(pu_1) : s\mathcal{F}_{t_1}\mathcal{Q} \otimes \dots \otimes s\mathcal{F}_{t_k}\mathcal{Q} \rightarrow s\mathcal{A}. \end{aligned}$$

Substituting definition (2.8.3) of u_1 we turn the above equation into an identity

$$- \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pdu_q \otimes \psi_{c\beta})b_{\alpha+1+\beta}^A \quad (2.8.5)$$

$$- (-)^p \sum_{\substack{\alpha+\beta>0 \\ \alpha+q+\beta=k}} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta})(pdu_{\alpha+1+\beta}) \quad (2.8.6)$$

$$= -(pdu_k)b_1 \quad (2.8.7)$$

$$\begin{aligned} &+ \sum_{a+q+c=k}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta})b_{\alpha+1+\beta}b_1 \\ &- (-)^p \sum_{\substack{\alpha+\beta>0 \\ \alpha+q+\beta=k}} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta})b_1 \end{aligned} \quad (2.8.8)$$

$$+ (-)^pb_k(\phi_1 \otimes pu_0)b_2 + (-)^pb_k(pu_0 \otimes \psi_1)b_2 \quad (2.8.9)$$

$$+ (-)^p \sum_{\substack{\gamma+\delta>0 \\ \gamma+j+\delta=k}} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta}) \left[-pdu_{\gamma+1+\delta} \right] \quad (2.8.10)$$

$$+ \sum_{a+q+c=\gamma+1+\delta}^{\alpha,\beta} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta})b_{\alpha+1+\beta} \quad (2.8.11)$$

$$- (-)^p \sum_{\alpha+q+\beta=\gamma+1+\delta}^{\alpha+\beta>0} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta}) \Big], \quad (2.8.12)$$

whose validity we are going to prove now. First of all, terms (2.8.6) and (2.8.10) cancel each other. Term (2.8.12) vanishes because for an arbitrary integer g the sum

$$\sum_{\substack{\alpha+1+\beta=g \\ \gamma+j+\delta=k \\ \alpha+q+\beta=\gamma+1+\delta}} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})(1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta}) \quad (2.8.13)$$

is the matrix coefficient $b^2 = 0 : T^k s\mathcal{FQ} \rightarrow T^g s\mathcal{FQ}$, thus, it vanishes. Notice that condition $\alpha + \beta > 0$ in (2.8.12) automatically implies $\gamma + \delta > 0$. Furthermore, term (2.8.7) cancels one of the terms of sum (2.8.5). In the remaining terms of (2.8.5) we may use the induction assumptions and replace pdu_q with the right hand side of (2.8.2). We also absorb terms (2.8.9) into sum (2.8.11), allowing $\gamma = \delta = 0$ in it and allowing simultaneously $\alpha = \beta = 0$ in (2.8.8) to compensate for the missing term $b_k(pu_1)b_1$:

$$- \sum_{\alpha+\beta>0} \sum_{\substack{\gamma,\delta \\ a+q+c=k \\ e+j+f=q}} [\phi_{a\alpha} \otimes (\phi_{e\gamma} \otimes pu_j \otimes \psi_{f\delta}) b_{\gamma+1+\delta}^A \otimes \psi_{c\beta}] b_{\alpha+1+\beta}^A \quad (2.8.14)$$

$$+ (-)^p \sum_{\alpha+\beta>0} \sum_{\substack{\gamma,\delta \\ a+q+c=k \\ \gamma+j+\delta=q}} [\phi_{a\alpha} \otimes (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})(pu_{\gamma+1+\delta}) \otimes \psi_{c\beta}] b_{\alpha+1+\beta}^A \quad (2.8.15)$$

$$= \sum_{\substack{\alpha,\beta \\ a+q+c=k}} (\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}^A b_1^A \quad (2.8.16)$$

$$- (-)^p \sum_{\alpha+q+\beta=k} (1^{\otimes\alpha} \otimes b_q \otimes 1^{\otimes\beta})(pu_{\alpha+1+\beta}) b_1^A \quad (2.8.17)$$

$$+ (-)^p \sum_{\gamma+j+\delta=k} \sum_{\substack{\alpha,\beta \\ a+q+c=\gamma+1+\delta}} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})(\phi_{a\alpha} \otimes pu_q \otimes \psi_{c\beta}) b_{\alpha+1+\beta}^A.$$

Recall that ϕ_{a0} vanish for all a except $a = 0$. Therefore, we may absorb term (2.8.16) into sum (2.8.14) and term (2.8.17) into sum (2.8.15), allowing terms with $\alpha = \beta = 0$ in them. Denote $r = pu \in sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$. The proposition follows immediately from the following

2.9 Lemma. *For all $r \in sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ and all $k \geq 0$ we have*

$$\begin{aligned} & - \sum_{\alpha,\beta} \sum_{\substack{\gamma,\delta \\ a+q+c=k \\ e+j+f=q}} [\phi_{a\alpha} \otimes (\phi_{e\gamma} \otimes r_j \otimes \psi_{f\delta}) b_{\gamma+1+\delta}^A \otimes \psi_{c\beta}] b_{\alpha+1+\beta}^A \\ & + (-)^r \sum_{\alpha,\beta} \sum_{\substack{\gamma,\delta \\ a+q+c=k \\ \gamma+j+\delta=q}} [\phi_{a\alpha} \otimes (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta}) r_{\gamma+1+\delta} \otimes \psi_{c\beta}] b_{\alpha+1+\beta}^A \end{aligned}$$

$$= (-)^r \sum_{\gamma+j+\delta=k} \sum_{a+q+c=\gamma+1+\delta}^{\alpha,\beta} (1^{\otimes\gamma} \otimes b_j \otimes 1^{\otimes\delta})(\phi_{a\alpha} \otimes r_q \otimes \psi_{c\beta})b_{\alpha+1+\beta}^A. \quad (2.9.1)$$

Proof. Sum (2.9.1) is split into three sums accordingly to output of b_j being an input of $\phi_{a\alpha}$ or r_q or $\psi_{c\beta}$:

$$- \sum_{a+e+j+f+c=k}^{\alpha,\beta,\gamma,\delta} (\phi_{a\alpha} \otimes \phi_{e\gamma} \otimes r_j \otimes \psi_{f\delta} \otimes \psi_{c\beta})(1^{\otimes\alpha} \otimes b_{\gamma+1+\delta}^A \otimes 1^{\otimes\beta})b_{\alpha+1+\beta}^A \quad (2.9.2)$$

$$+ (-)^r \sum_{a+\gamma+j+\delta+c=k}^{\alpha,\beta} (1^{\otimes a+\gamma} \otimes b_j \otimes 1^{\otimes\delta+c})(\phi_{a\alpha} \otimes r_{\gamma+1+\delta} \otimes \psi_{c\beta})b_{\alpha+1+\beta}^A \quad (2.9.3)$$

$$= (-)^r \sum_{x+q+c=k}^{a,\alpha,\beta} (b_{xa}\phi_{a\alpha} \otimes r_q \otimes \psi_{c\beta})b_{\alpha+1+\beta}^A \quad (2.9.4)$$

$$+ (-)^r \sum_{a+y+c=k}^{\alpha,q,\beta} (\phi_{a\alpha} \otimes b_{yq}r_q \otimes \psi_{c\beta})b_{\alpha+1+\beta}^A \quad (2.9.5)$$

$$+ \sum_{a+q+z=k}^{\alpha,\beta,c} (\phi_{a\alpha} \otimes r_q \otimes b_{zc}\psi_{c\beta})b_{\alpha+1+\beta}^A. \quad (2.9.6)$$

Here $b_{xa} : T^x s\mathcal{FQ} \rightarrow T^a s\mathcal{FQ}$ is a matrix element of $b^{\mathcal{FQ}}$. Terms (2.9.3) and (2.9.5) cancel each other. We shall use A_∞ -functor identities $b\phi = \phi b$, $b\psi = \psi b$ for terms (2.9.4) and (2.9.6). Being a cocategory homomorphism, ϕ satisfies the identity

$$\sum_{a+e=h} \phi_{a\alpha} \otimes \phi_{e\gamma} = [\Delta(\phi \otimes \phi)]_{h;\alpha,\gamma} = \phi_{h,\alpha+\gamma} \Delta_{\alpha+\gamma;\alpha,\gamma}$$

for all non-negative integers h , where Δ is the cut comultiplication. Similarly for ψ . Using this identity in (2.9.2) we get the equation to verify:

$$\begin{aligned} & - \sum_{x+q+z=k}^{v,w} (\phi_{xv} \otimes r_q \otimes \psi_{zw}) \sum_{\alpha+y+\beta=v+1+w}^{\alpha \leq v, \beta \leq w} (1^{\otimes\alpha} \otimes b_y^A \otimes 1^{\otimes\beta})b_{\alpha+1+\beta}^A \\ & = \sum_{x+q+z=k}^{v,w,\alpha} (\phi_{xv} \otimes r_q \otimes \psi_{zw})(b_{v\alpha}^A \otimes 1^{\otimes 1+w})b_{\alpha+1+w}^A \\ & + \sum_{x+q+z=k}^{v,w,\beta} (\phi_{xv} \otimes r_q \otimes \psi_{zw})(1^{\otimes v+1} \otimes b_{w\beta}^A)b_{v+1+\beta}^A. \end{aligned}$$

It follows from the identity $b^2 \text{pr}_1 = 0 : T^{v+1+w} s\mathcal{A} \rightarrow s\mathcal{A}$ valid for arbitrary non-negative integers v, w , which we may rewrite like this:

$$\sum_{\alpha+y+\beta=v+1+w}^{\alpha \leq v, \beta \leq w} (1^{\otimes\alpha} \otimes b_y^A \otimes 1^{\otimes\beta})b_{\alpha+1+\beta}^A + \sum_{\alpha} (b_{v\alpha}^A \otimes 1^{\otimes 1+w})b_{\alpha+1+w}^A + \sum_{\beta} (1^{\otimes v+1} \otimes b_{w\beta}^A)b_{v+1+\beta}^A = 0.$$

So the lemma is proved. \square

The proposition follows. \square

Let us consider now the question, when the discussed chain map is null-homotopic.

2.10 Corollary. *Let $\phi, \psi : \mathcal{FQ} \rightarrow \mathcal{A}$ be A_∞ -functors. Let P be a complex of \mathbb{k} -modules. Let $u : P \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ be a chain map. The set (possibly empty) of homotopies $h : P \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$, $\deg h = -1$, such that $u = dh + hB_1$ is in bijection with the set of data $(h', h_k)_{k>1}$, consisting of*

1. a homotopy $h' : P \rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)$, $\deg h' = -1$, such that $dh' + h'B_1 = u'$, where u' is given by (2.8.1);
2. \mathbb{k} -linear maps

$$h_k : P \rightarrow \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathbb{C}((s\mathcal{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi))$$

of degree -1 for all $k > 1$.

The bijection maps h to $(h', h_k)_{k>1}$, where $h_k = h \cdot \text{pr}_k$ and

$$h' = (P \xrightarrow{h} sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_1(\mathcal{FQ}, \mathcal{A})(\phi, \psi) \xrightarrow{\text{restr}} sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi)).$$

The inverse bijection can be recovered from the recurrent formula

$$\begin{aligned} (-)^p b_k(ph_1) &= pu_k - (pd)h_k - \sum_{a+q+c=k}^{\alpha, \beta} (\phi_{a\alpha} \otimes ph_q \otimes \psi_{c\beta})b_{\alpha+1+\beta} \\ &\quad - (-)^p \sum_{a+q+c=k}^{a+c>0} (1^{\otimes a} \otimes b_q \otimes 1^{\otimes c})(ph_{a+1+c}) : (s\mathcal{FQ})^{\otimes k} \rightarrow s\mathcal{A}, \end{aligned}$$

where $k > 1$, $p \in P$, and $\phi_{a\alpha}, \psi_{c\beta}$ are matrix elements of ϕ, ψ .

Proof. We shall apply Proposition 2.8 to the complex $\text{Cone}(\text{id} : P \rightarrow P)$ instead of P . The graded \mathbb{k} -module $\text{Cone}(\text{id}_P) = P \oplus P[1]$ is equipped with the differential $(q, ps)d = (qd + p, -pds)$, $p, q \in P$. The chain maps $\bar{u} : \text{Cone}(\text{id}_P) \rightarrow C$ to an arbitrary complex C are in bijection with pairs $(u : P \rightarrow C, h : P \rightarrow C)$, where $u = dh + hd$ and $\deg h = -1$. The pair $(u, h) = (\text{in}_1 \bar{u}, \text{sin}_2 \bar{u})$ is assigned to \bar{u} , and the map $\bar{u} : P \oplus P[1] \rightarrow C$, $(q, ps) \mapsto qu + ph$ is assigned to a pair (u, h) . Indeed, \bar{u} being chain map is equivalent to

$$(q, ps)d\bar{u} = qdu + pu - pdh = qud + phd = (q, ps)\bar{u}d,$$

that is, to conditions $du = ud$, $u = dh + hd$.

Thus, for a fixed chain map $u : P \rightarrow C$ the set of homotopies $h : P \rightarrow C$, such that $u = dh + hd$, is in bijection with the set of chain maps $\bar{u} : \text{Cone}(\text{id}_P) \rightarrow C$ such that $\text{in}_1 \bar{u} = u : P \rightarrow C$. Applying this statement to $u : P \rightarrow C = sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ we find by Proposition 2.8 that the set of homotopies $h : P \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(\phi, \psi)$ such that $u = dh + hB_1$ is in bijection with the set of data $(\bar{u}', \bar{u}_k)_{k>1}$, such that

$$\begin{aligned} \bar{u}' : \text{Cone}(\text{id}_P) &\rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\phi, \psi) && \text{is a chain map,} && \text{in}_1 \bar{u}' = u', \\ \bar{u}_k : \text{Cone}(\text{id}_P) &\rightarrow \prod_{X, Y \in \text{Ob } \mathcal{Q}} \mathcal{C}((s\mathcal{FQ})^{\otimes k}(X, Y), s\mathcal{A}(X\phi, Y\psi)), && \deg \bar{u}_k = 0, && \text{in}_1 \bar{u}_k = u_k, \end{aligned}$$

therefore, in bijection with the set of data $(h', h_k)_{k>1} = (s \text{in}_2 \bar{u}', s \text{in}_2 \bar{u}_k)_{k>1}$, as stated in corollary. \square

2.11. Restriction as an A_∞ -functor. Let \mathcal{Q} be a (\mathcal{U} -small) differential graded \mathbb{k} -quiver. Denote by \mathcal{FQ} the free A_∞ -category generated by \mathcal{Q} . Let \mathcal{A} be a (\mathcal{U} -small) unital A_∞ -category. There is the restriction strict A_∞ -functor

$$\text{restr} : A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A}), \quad (f : \mathcal{FQ} \rightarrow \mathcal{A}) \mapsto (\bar{f} = (f_1|_{\mathcal{Q}}) : \mathcal{Q} \rightarrow \mathcal{A}).$$

In fact, it is the composition of two strict A_∞ -functors: $A_\infty(\mathcal{FQ}, \mathcal{A}) \xrightarrow{\text{restr}_{\infty, 1}} A_1(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$, where the second comes from the full embedding $\mathcal{Q} \hookrightarrow \mathcal{FQ}$. Its first component is

$$\begin{aligned} \text{restr}_1 : sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g) &\rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g}), \\ r = (r_0, r_1, \dots, r_n, \dots) &\mapsto (r_0, r_1|_{\mathcal{Q}}) = \bar{r}. \end{aligned} \tag{2.11.1}$$

2.12 Theorem. *The A_∞ -functor $\text{restr} : A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$ is an equivalence.*

Proof. Let us prove that restriction map (2.11.1) is homotopy invertible. We construct a chain map going in the opposite direction

$$u : sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g}) \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g)$$

via Proposition 2.8 taking $P = sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g})$. We choose

$$u' : sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g}) \rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g})$$

to be the identity map and $u_k = 0$ for $k > 1$. Therefore,

$$u \cdot \text{restr}_1 = u' = \text{id}_{sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g})}.$$

Denote

$$v = \text{id}_{sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g)} - [sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g) \xrightarrow{\text{restr}_1} sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g}) \xrightarrow{u} sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g)].$$

Let us prove that v is null-homotopic via Corollary 2.10, taking $P = sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g)$. A homotopy $h : sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g) \rightarrow sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g)$, $\deg h = -1$, such that $v = B_1h + hB_1$ is specified by $h' = 0 : sA_\infty(\mathcal{FQ}, \mathcal{A})(f, g) \rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\bar{f}, \bar{g})$ and $h_k = 0$ for $k > 1$. Indeed,

$$v' = v \cdot \text{restr}_1 = \text{restr}_1 - \text{restr}_1 \cdot u \cdot \text{restr}_1 = \text{restr}_1 - \text{restr}_1 = 0,$$

so $v' = B_1h' + h'B_1$ and condition 1 of Corollary 2.10 is satisfied². Therefore, u is homotopy inverse to restr_1 .

Let $\mathbf{i}^{\mathcal{A}}$ be a unit transformation of the unital A_∞ -category \mathcal{A} . Then $A_1(\mathcal{Q}, \mathcal{A})$ is a unital A_∞ -category with the unit transformation $(1 \otimes \mathbf{i}^{\mathcal{A}})M$ (cf. [Lyu03, Proposition 7.7]). The unit element for an object $\phi \in \text{Ob } A_1(\mathcal{Q}, \mathcal{A})$ is ${}_{\phi}\mathbf{i}_0^{A_1(\mathcal{Q}, \mathcal{A})} : \mathbb{k} \rightarrow sA_1(\mathcal{Q}, \mathcal{A})$, $1 \mapsto \phi\mathbf{i}^{\mathcal{A}}$. The A_∞ -category $A_\infty(\mathcal{FQ}, \mathcal{A})$ is also unital. To establish equivalence of these two A_∞ -categories via $\text{restr} : A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$ we verify the conditions of Theorem 8.8 from [Lyu03].

Consider the mapping $\text{Ob } A_1(\mathcal{Q}, \mathcal{A}) \rightarrow \text{Ob } A_\infty(\mathcal{FQ}, \mathcal{A})$, $\phi \mapsto \widehat{\phi}$, which extends a given chain map to a strict A_∞ -functor, constructed in Corollary 2.4. Clearly, $\overline{\widehat{\phi}} = \phi$. It remains to give two mutually inverse cycles, which we choose as follows:

$$\begin{aligned} {}_{\phi}r_0 : \mathbb{k} &\rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\phi, \widehat{\widehat{\phi}}), & 1 &\mapsto \phi\mathbf{i}^{\mathcal{A}}, \\ {}_{\phi}p_0 : \mathbb{k} &\rightarrow sA_1(\mathcal{Q}, \mathcal{A})(\widehat{\widehat{\phi}}, \phi), & 1 &\mapsto \phi\mathbf{i}^{\mathcal{A}}. \end{aligned}$$

Clearly, ${}_{\phi}r_0B_1 = 0$, ${}_{\phi}p_0B_1 = 0$,

$$\begin{aligned} ({}_{\phi}r_0 \otimes {}_{\phi}p_0)B_2 - {}_{\phi}\mathbf{i}_0^{A_1(\mathcal{Q}, \mathcal{A})} : 1 &\mapsto (\phi\mathbf{i}^{\mathcal{A}} \otimes \phi\mathbf{i}^{\mathcal{A}})B_2 - \phi\mathbf{i}^{\mathcal{A}} \in \text{Im } B_1, \\ ({}_{\phi}p_0 \otimes {}_{\phi}r_0)B_2 - {}_{\phi}\mathbf{i}_0^{A_1(\mathcal{Q}, \mathcal{A})} : 1 &\mapsto (\phi\mathbf{i}^{\mathcal{A}} \otimes \phi\mathbf{i}^{\mathcal{A}})B_2 - \phi\mathbf{i}^{\mathcal{A}} \in \text{Im } B_1. \end{aligned}$$

Therefore, all assumptions of Theorem 8.8 [Lyu03] are satisfied. Thus, $\text{restr} : A_\infty(\mathcal{FQ}, \mathcal{A}) \rightarrow A_1(\mathcal{Q}, \mathcal{A})$ is an A_∞ -equivalence. \square

2.13 Corollary. *Every A_∞ -functor $f : \mathcal{FQ} \rightarrow \mathcal{A}$ is isomorphic to the strict A_∞ -functor $\widehat{\widehat{f}} : \mathcal{FQ} \rightarrow \mathcal{A}$.*

Proof. Note that $\bar{f} = \widehat{\widehat{f}}$. The A_1 -transformation $\bar{f}\mathbf{i}^{\mathcal{A}} : \bar{f} \rightarrow \widehat{\widehat{f}} : \mathcal{Q} \rightarrow \mathcal{A}$ with the components $({}_xf\mathbf{i}_0^{\mathcal{A}}, \bar{f}_1\mathbf{i}_1^{\mathcal{A}})$ is natural. It is mapped by u into a natural A_∞ -transformation $(\bar{f}\mathbf{i}^{\mathcal{A}})u : f \rightarrow \widehat{\widehat{f}} : \mathcal{FQ} \rightarrow \mathcal{A}$. Its zero component ${}_xf\mathbf{i}_0^{\mathcal{A}}$ is invertible, therefore $(\bar{f}\mathbf{i}^{\mathcal{A}})u$ is invertible by [Lyu03, Proposition 7.15]. \square

²By the way, the only non-vanishing component of h is h_1 .

3. Representable 2-functors $A_\infty^u \rightarrow A_\infty^u$

Recall that unital A_∞ -categories, unital A_∞ -functors and equivalence classes of natural A_∞ -transformations form a 2-category [Lyu03]. In order to distinguish between the A_∞ -category $A_\infty^u(\mathcal{C}, \mathcal{D})$ and the ordinary category, whose morphisms are equivalence classes of natural A_∞ -transformations, we denote the latter by

$$\overline{A_\infty^u}(\mathcal{C}, \mathcal{D}) = H^0(A_\infty^u(\mathcal{C}, \mathcal{D}), m_1).$$

The corresponding notation for the 2-category is $\overline{A_\infty^u}$. We will see that arbitrary A_N -categories can be viewed as 2-functors $\overline{A_\infty^u} \rightarrow \overline{A_\infty^u}$. Moreover, they come from certain generalizations called A_∞^u -2-functors. There is a notion of representability of such 2-functors, which explains some constructions of A_∞ -categories. For instance, a differential graded \mathbb{k} -quiver \mathcal{Q} will be represented by the free A_∞ -category \mathcal{FQ} generated by it.

3.1 Definition. A (strict) A_∞^u -2-functor $F : A_\infty^u \rightarrow A_\infty^u$ consists of

1. a map $F : \text{Ob } A_\infty^u \rightarrow \text{Ob } A_\infty^u$;
2. a unital A_∞ -functor $F = F_{\mathcal{C}, \mathcal{D}} : A_\infty^u(\mathcal{C}, \mathcal{D}) \rightarrow A_\infty^u(F\mathcal{C}, F\mathcal{D})$ for each pair \mathcal{C}, \mathcal{D} of unital A_∞ -categories;
such that
3. $\text{id}_{F\mathcal{C}} = F(\text{id}_{\mathcal{C}})$ for any unital A_∞ -category \mathcal{C} ;
4. the equation

$$\begin{array}{ccc} TsA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty^u(\mathcal{D}, \mathcal{E}) & \xrightarrow{M} & TsA_\infty^u(\mathcal{C}, \mathcal{E}) \\ \downarrow F_{\mathcal{C}, \mathcal{D}} \boxtimes F_{\mathcal{D}, \mathcal{E}} & = & \downarrow F_{\mathcal{C}, \mathcal{E}} \\ TsA_\infty^u(F\mathcal{C}, F\mathcal{D}) \boxtimes TsA_\infty^u(F\mathcal{D}, F\mathcal{E}) & \xrightarrow{M} & TsA_\infty^u(F\mathcal{C}, F\mathcal{E}) \end{array} \quad (3.1.1)$$

holds strictly for each triple $\mathcal{C}, \mathcal{D}, \mathcal{E}$ of unital A_∞ -categories.

The A_∞ -functor $F : A_\infty^u(\mathcal{C}, \mathcal{D}) \rightarrow A_\infty^u(F\mathcal{C}, F\mathcal{D})$ consists of the mapping of objects

$$\text{Ob } F : \text{Ob } A_\infty^u(\mathcal{C}, \mathcal{D}) \rightarrow \text{Ob } A_\infty^u(F\mathcal{C}, F\mathcal{D}), \quad f \mapsto Ff,$$

and the components $F_k, k \geq 1$:

$$\begin{aligned} F_1 &: sA_\infty^u(\mathcal{C}, \mathcal{D})(f, g) \rightarrow sA_\infty^u(F\mathcal{C}, F\mathcal{D})(Ff, Fg), \\ F_2 &: sA_\infty^u(\mathcal{C}, \mathcal{D})(f, g) \otimes sA_\infty^u(\mathcal{C}, \mathcal{D})(g, h) \rightarrow sA_\infty^u(F\mathcal{C}, F\mathcal{D})(Ff, Fh), \end{aligned}$$

and so on.

Weak versions of A_∞^u -2-functors and 2-transformations between them might be considered elsewhere.

3.2 Definition. A (strict) A_∞^u -2-transformation $\lambda : F \rightarrow G : A_\infty^u \rightarrow A_\infty^u$ of strict A_∞^u -2-functors is

1. a family of unital A_∞ -functors $\lambda_{\mathcal{C}} : F\mathcal{C} \rightarrow G\mathcal{C}$, $\mathcal{C} \in \text{Ob } A_\infty^u$;
such that
2. the diagram of A_∞ -functors

$$\begin{array}{ccc} A_\infty^u(\mathcal{C}, \mathcal{D}) & \xrightarrow{F} & A_\infty^u(F\mathcal{C}, F\mathcal{D}) \\ G \downarrow & = & \downarrow (1 \boxtimes \lambda_{\mathcal{D}})M \\ A_\infty^u(G\mathcal{C}, G\mathcal{D}) & \xrightarrow{(\lambda_{\mathcal{C}} \boxtimes 1)M} & A_\infty^u(F\mathcal{C}, G\mathcal{D}) \end{array} \quad (3.2.1)$$

strictly commutes.

An A_∞^u -2-transformation $\lambda = (\lambda_{\mathcal{C}})$ for which $\lambda_{\mathcal{C}}$ are A_∞ -equivalences is called a *natural A_∞^u -2-equivalence*.

Let us show now that the above notions induce ordinary strict 2-functors and strict 2-transformations in 0-th cohomology. Recall that the strict 2-category $\overline{A_\infty^u}$ consists of objects – unital A_∞ -categories, the category $\overline{A_\infty^u}(\mathcal{C}, \mathcal{D})$ for any pair of objects \mathcal{C}, \mathcal{D} , the identity functor $\text{id}_{\mathcal{C}}$ for any unital A_∞ -category \mathcal{C} , and the composition functor [Lyu03]

$$\begin{aligned} \overline{A_\infty^u}(\mathcal{C}, \mathcal{D})(f, g) \times \overline{A_\infty^u}(\mathcal{D}, \mathcal{E})(h, k) &\xrightarrow{\bullet^2} \overline{A_\infty^u}(\mathcal{C}, \mathcal{E})(fh, gk), \\ (rs^{-1}, ps^{-1}) &\longmapsto (rhs^{-1} \otimes gps^{-1})m_2. \end{aligned}$$

Given a strict A_∞^u -2-functor F as in Definition 3.1 we construct from it an ordinary strict 2-functor $\overline{F} = F : \text{Ob } \overline{A_\infty^u} \rightarrow \text{Ob } \overline{A_\infty^u}$, $\overline{F} = H^0(sF_1s^{-1}) : \overline{A_\infty^u}(\mathcal{C}, \mathcal{D}) \rightarrow \overline{A_\infty^u}(F\mathcal{C}, F\mathcal{D})$ as follows.

Denote

$$\begin{aligned} M_{10} \odot M_{01} &= \{ sA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes sA_\infty^u(\mathcal{D}, \mathcal{E}) \xrightarrow[\sim]{\Delta_{10} \boxtimes \Delta_{01}} \\ &\quad [sA_\infty^u(\mathcal{C}, \mathcal{D}) \otimes T^0 sA_\infty^u(\mathcal{C}, \mathcal{D})] \boxtimes [T^0 sA_\infty^u(\mathcal{D}, \mathcal{E}) \otimes sA_\infty^u(\mathcal{D}, \mathcal{E})] \\ &\xrightarrow{\sim} [sA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes T^0 sA_\infty^u(\mathcal{D}, \mathcal{E})] \otimes [T^0 sA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes sA_\infty^u(\mathcal{D}, \mathcal{E})] \\ &\xrightarrow{M_{10} \otimes M_{01}} sA_\infty^u(\mathcal{C}, \mathcal{E}) \otimes sA_\infty^u(\mathcal{C}, \mathcal{E}) \}, \end{aligned} \quad (3.2.2)$$

where the obvious isomorphisms Δ_{10} and Δ_{01} are components of the comultiplication Δ , the middle isomorphism is that of distributivity law (1.5.3), and the components M_{10} and M_{01} of M are the composition maps.

Property (3.1.1) of F implies that

$$(M_{10} \odot M_{01})(F_1 \otimes F_1) = (F_1 \boxtimes F_1)(M_{10} \odot M_{01}). \quad (3.2.3)$$

Indeed, the following diagram commutes

$$\begin{array}{ccc} sA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes T^0 sA_\infty^u(\mathcal{D}, \mathcal{E}) & \xrightarrow{M_{10}} & sA_\infty^u(\mathcal{C}, \mathcal{E}) \\ F_1 \boxtimes \text{Ob } F \downarrow & & \downarrow F_1 \\ sA_\infty^u(F\mathcal{C}, F\mathcal{D}) \boxtimes T^0 sA_\infty^u(F\mathcal{D}, F\mathcal{E}) & \xrightarrow{M_{10}} & sA_\infty^u(F\mathcal{C}, F\mathcal{E}) \end{array}$$

due to (3.1.1). \otimes -tensoring it with one more similar diagram we get

$$(M_{10} \otimes M_{01})(F_1 \otimes F_1) = [(F_1 \boxtimes \text{Ob } F) \otimes (\text{Ob } F \boxtimes F_1)](M_{10} \otimes M_{01}).$$

The isomorphisms in (3.2.2) commute with F in expected way, so (3.2.3) follows.

We claim that the diagram

$$\begin{array}{ccc} sA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes sA_\infty^u(\mathcal{D}, \mathcal{E}) & \xrightarrow{(M_{10} \odot M_{01})B_2} & sA_\infty^u(\mathcal{C}, \mathcal{E}) \\ F_1 \boxtimes F_1 \downarrow & & \downarrow F_1 \\ sA_\infty^u(F\mathcal{C}, F\mathcal{D}) \boxtimes sA_\infty^u(F\mathcal{D}, F\mathcal{E}) & \xrightarrow{(M_{10} \odot M_{01})B_2} & sA_\infty^u(F\mathcal{C}, F\mathcal{E}) \end{array} \quad (3.2.4)$$

homotopically commutes. Indeed, since

$$(1 \otimes B_1 + B_1 \otimes 1)F_2 + B_2F_1 = (F_1 \otimes F_1)B_2 + F_2B_1,$$

we get

$$\begin{aligned} & (M_{10} \odot M_{01})B_2F_1 \\ &= (M_{10} \odot M_{01})(F_1 \otimes F_1)B_2 + (M_{10} \odot M_{01})F_2B_1 - (M_{10} \odot M_{01})(1 \otimes B_1 + B_1 \otimes 1)F_2 \\ &= (F_1 \boxtimes F_1)(M_{10} \odot M_{01})B_2 + (M_{10} \odot M_{01})F_2B_1 - (1 \boxtimes B_1 + B_1 \boxtimes 1)(M_{10} \odot M_{01})F_2. \end{aligned}$$

We have used equations

$$\begin{aligned} (M_{10} \odot M_{01})(1 \otimes B_1) &= (1 \boxtimes B_1)(M_{10} \odot M_{01}), \\ (M_{10} \odot M_{01})(B_1 \otimes 1) &= (B_1 \boxtimes 1)(M_{10} \odot M_{01}), \end{aligned}$$

which can be proved similarly to (3.2.3) due to M being an A_∞ -functor. Passing to cohomology we get from (3.2.4) a strictly commutative diagram of functors

$$\begin{array}{ccc} H^0(A_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes A_\infty^u(\mathcal{D}, \mathcal{E})) & \xrightarrow{\bullet^2} & H^0(A_\infty^u(\mathcal{C}, \mathcal{E})) \\ H^0(sF_1s^{-1} \boxtimes sF_1s^{-1}) \downarrow & = & \downarrow H^0(sF_1s^{-1}) \\ H^0(A_\infty^u(F\mathcal{C}, F\mathcal{D}) \boxtimes A_\infty^u(F\mathcal{D}, F\mathcal{E})) & \xrightarrow{\bullet^2} & H^0(A_\infty^u(F\mathcal{C}, F\mathcal{E})) \end{array}$$

since $\bullet^2 = H^0((s \boxtimes s)(M_{10} \odot M_{01})B_2s^{-1})$. Using the Künneth map we come to strictly commutative diagram of functors

$$\begin{array}{ccc} \overline{A}_\infty^u(\mathcal{C}, \mathcal{D}) \times \overline{A}_\infty^u(\mathcal{D}, \mathcal{E}) & \xrightarrow{\bullet^2} & \overline{A}_\infty^u(\mathcal{C}, \mathcal{E}) \\ H^0(sF_1s^{-1}) \times H^0(sF_1s^{-1}) \downarrow & = & \downarrow H^0(sF_1s^{-1}) \\ \overline{A}_\infty^u(F\mathcal{C}, F\mathcal{D}) \times \overline{A}_\infty^u(F\mathcal{D}, F\mathcal{E}) & \xrightarrow{\bullet^2} & \overline{A}_\infty^u(F\mathcal{C}, F\mathcal{E}) \end{array}$$

that is, to a usual strict 2-functor $\overline{F} : \overline{A_\infty^u} \rightarrow \overline{A_\infty^u}$.

Let us show that an A_∞^u -2-transformation $\lambda : F \rightarrow G : A_\infty^u \rightarrow A_\infty^u$ as in Definition 3.2 induces an ordinary strict 2-transformation $\overline{\lambda} : \overline{F} \rightarrow \overline{G} : \overline{A_\infty^u} \rightarrow \overline{A_\infty^u}$ in cohomology. Indeed, diagram (3.2.1) implies commutativity of diagram

$$\begin{array}{ccc} sA_\infty^u(\mathcal{C}, \mathcal{D}) & \xrightarrow{F_1} & sA_\infty^u(F\mathcal{C}, F\mathcal{D}) \\ G_1 \downarrow & = & \downarrow (1 \boxtimes \lambda_{\mathcal{D}})M_{10} \\ sA_\infty^u(G\mathcal{C}, G\mathcal{D}) & \xrightarrow{(\lambda_{\mathcal{C}} \boxtimes 1)M} & sA_\infty^u(F\mathcal{C}, G\mathcal{D}) \end{array}$$

$$r : f \rightarrow g : G\mathcal{C} \rightarrow G\mathcal{D} \longmapsto \lambda_{\mathcal{C}}r : \lambda_{\mathcal{C}}f \rightarrow \lambda_{\mathcal{C}}g : F\mathcal{C} \rightarrow G\mathcal{D}.$$

Passing to cohomology we get

$$\begin{array}{ccc} \overline{A_\infty^u}(\mathcal{C}, \mathcal{D}) & \xrightarrow{H^0(sF_1s^{-1})} & \overline{A_\infty^u}(F\mathcal{C}, F\mathcal{D}) \\ H^0(sG_1s^{-1}) \downarrow & = & \downarrow \text{--}\lambda_{\mathcal{D}} = \overline{A_\infty^u}(F\mathcal{C}, \lambda_{\mathcal{D}}) \\ \overline{A_\infty^u}(G\mathcal{C}, G\mathcal{D}) & \xrightarrow[\overline{A_\infty^u}(\lambda_{\mathcal{C}}, G\mathcal{D})]{\lambda_{\mathcal{C}} \text{--}} & \overline{A_\infty^u}(F\mathcal{C}, G\mathcal{D}) \end{array}$$

Therefore, $\overline{\lambda}_{\mathcal{C}} \in \text{Ob } \overline{A_\infty^u}(F\mathcal{C}, G\mathcal{C})$ form a strict 2-transformation $\overline{\lambda} : \overline{F} \rightarrow \overline{G} : \overline{A_\infty^u} \rightarrow \overline{A_\infty^u}$.

3.3. Examples of A_∞^u -2-functors. Let \mathcal{A} be an A_N -category, $1 \leq N \leq \infty$. It determines an A_∞^u -2-functor $F = A_N(\mathcal{A}, -) : A_\infty^u \rightarrow A_\infty^u$, given by the following data:

1. the map $F : \text{Ob } A_\infty^u \rightarrow \text{Ob } A_\infty^u$, $\mathcal{C} \mapsto A_N(\mathcal{A}, \mathcal{C})$ (the category $A_N(\mathcal{A}, \mathcal{C})$ is unital by [Lyu03, Proposition 7.7]);
2. the unital strict A_∞ -functor $F = A_N(\mathcal{A}, -) : A_\infty^u(\mathcal{C}, \mathcal{D}) \rightarrow A_\infty^u(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{D}))$ for each pair \mathcal{C}, \mathcal{D} of unital A_∞ -categories (cf. [Lyu03, Propositions 6.2, 8.4]).

Clearly, $\text{id}_{A_N(\mathcal{A}, \mathcal{C})} = (1 \boxtimes \text{id}_{\mathcal{C}})M = A_N(\mathcal{A}, \text{id}_{\mathcal{C}})$. We want to prove now that the equation

$$\begin{aligned} [TsA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty^u(\mathcal{D}, \mathcal{E}) &\xrightarrow{M} TsA_\infty^u(\mathcal{C}, \mathcal{E}) \xrightarrow{A_N(\mathcal{A}, -)} TsA_\infty^u(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{E}))] \\ &= [TsA_\infty^u(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty^u(\mathcal{D}, \mathcal{E}) \xrightarrow{A_N(\mathcal{A}, -) \boxtimes A_N(\mathcal{A}, -)} \\ &\quad TsA_\infty^u(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{D})) \boxtimes TsA_\infty^u(A_N(\mathcal{A}, \mathcal{D}), A_N(\mathcal{A}, \mathcal{E})) \\ &\quad \xrightarrow{M} TsA_\infty^u(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{E}))] \quad (3.3.1) \end{aligned}$$

holds strictly for each triple $\mathcal{C}, \mathcal{D}, \mathcal{E}$ of unital A_∞ -categories. In fact, this F is a restriction of an A_∞ -2-functor $F : \text{Ob } A_\infty \rightarrow \text{Ob } A_\infty$, $\mathcal{C} \mapsto A_N(\mathcal{A}, \mathcal{C})$, which is defined just as in Definition 3.1 without mentioning the unitality. Equation (3.3.1) follows from a similar

equation without the unitality index u . To prove it we consider the compositions

$$\begin{aligned}
& [TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \xrightarrow{1 \boxtimes M} TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{E}) \\
& \xrightarrow{1 \boxtimes A_N(\mathcal{A}, -)} TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{E})) \xrightarrow{\alpha} TsA_N(\mathcal{A}, \mathcal{E})] \\
& = [TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \\
& \xrightarrow{1 \boxtimes M} TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{E}) \xrightarrow{M} TsA_N(\mathcal{A}, \mathcal{E})] \\
& = [TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \\
& \xrightarrow{M \boxtimes 1} TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{E}) \xrightarrow{M} TsA_N(\mathcal{A}, \mathcal{E})] \\
& = [TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \xrightarrow{1 \boxtimes A_N(\mathcal{A}, -) \boxtimes 1} \\
& TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{D})) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \\
& \xrightarrow{\alpha \boxtimes 1} TsA_N(\mathcal{A}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \\
& \xrightarrow{1 \boxtimes A_N(\mathcal{A}, -)} TsA_N(\mathcal{A}, \mathcal{D}) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{D}), A_N(\mathcal{A}, \mathcal{E})) \xrightarrow{\alpha} TsA_N(\mathcal{A}, \mathcal{E})] \\
& = [TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(\mathcal{C}, \mathcal{D}) \boxtimes TsA_\infty(\mathcal{D}, \mathcal{E}) \xrightarrow{1 \boxtimes A_N(\mathcal{A}, -) \boxtimes A_N(\mathcal{A}, -)} \\
& TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{D})) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{D}), A_N(\mathcal{A}, \mathcal{E})) \\
& \xrightarrow{1 \boxtimes M} TsA_N(\mathcal{A}, \mathcal{C}) \boxtimes TsA_\infty(A_N(\mathcal{A}, \mathcal{C}), A_N(\mathcal{A}, \mathcal{E})) \xrightarrow{\alpha} TsA_N(\mathcal{A}, \mathcal{E})].
\end{aligned}$$

By Proposition 1.6 we deduce equation (3.3.1) (see also [Lyu03, Proposition 5.5]).

Let now \mathcal{A} be a unital A_∞ -category. It determines an A_∞^u -2-functor $G = A_\infty^u(\mathcal{A}, -) : A_\infty^u \rightarrow A_\infty^u$, given by the following data:

1. the map $G : \text{Ob } A_\infty^u \rightarrow \text{Ob } A_\infty^u$, $\mathcal{C} \mapsto A_\infty^u(\mathcal{A}, \mathcal{C})$ (the category $A_\infty^u(\mathcal{A}, \mathcal{C})$ is unital by [Lyu03, Proposition 7.7]);
2. the unital strict A_∞ -functor $G = A_\infty^u(\mathcal{A}, -) : A_\infty^u(\mathcal{C}, \mathcal{D}) \rightarrow A_\infty^u(A_\infty^u(\mathcal{A}, \mathcal{C}), A_\infty^u(\mathcal{A}, \mathcal{D}))$ for each pair \mathcal{C}, \mathcal{D} of unital A_∞ -categories, determined from

$$\begin{aligned}
M &= [TsA_\infty^u(\mathcal{A}, \mathcal{B}) \boxtimes TsA_\infty^u(\mathcal{B}, \mathcal{C}) \xrightarrow{1 \boxtimes A_\infty^u(\mathcal{A}, -)} \\
& TsA_\infty^u(\mathcal{A}, \mathcal{B}) \boxtimes TsA_\infty^u(A_\infty^u(\mathcal{A}, \mathcal{B}), A_\infty^u(\mathcal{A}, \mathcal{C})) \xrightarrow{\alpha} TsA_\infty^u(\mathcal{A}, \mathcal{C})].
\end{aligned}$$

(cf. [Lyu03, Propositions 6.2, 8.4]).

Clearly, $G\mathcal{C}$ are full A_∞ -subcategories of $F\mathcal{C}$ for the A_∞^u -2-functor $F = A_\infty(\mathcal{A}, -)$. Furthermore, A_∞ -functors $G_{\mathcal{C}, \mathcal{D}}(f)$ are restrictions of A_∞ -functors $F_{\mathcal{C}, \mathcal{D}}(f)$, so G is a full A_∞^u -2-subfunctor of F . In particular, G satisfies equation (3.1.1). Another way to prove that G is an A_∞^u -2-functor is to repeat the reasoning concerning F .

3.4. Example of an A_∞^u -2-equivalence. Assume that \mathcal{Q} is a differential graded \mathbb{k} -quiver. As usual, $\mathcal{F}\mathcal{Q}$ denotes the free A_∞ -category generated by it. We claim that $\text{restr} :$

$A_\infty(\mathcal{FQ}, _) \rightarrow A_1(\mathcal{Q}, _) : A_\infty^u \rightarrow A_\infty^u$ is a strict 2-natural A_∞ -equivalence. Indeed, it is given by the family of unital A_∞ -functors $\text{restr}_\mathcal{C} : A_\infty(\mathcal{FQ}, \mathcal{C}) \rightarrow A_1(\mathcal{Q}, \mathcal{C})$, $\mathcal{C} \in \text{Ob } A_\infty^u$, which are equivalences by Theorem 2.12. We have to prove that the diagram of A_∞ -functors

$$\begin{array}{ccc} A_\infty^u(\mathcal{C}, \mathcal{D}) & \xrightarrow{A_\infty(\mathcal{FQ}, _)} & A_\infty^u(A_\infty(\mathcal{FQ}, \mathcal{C}), A_\infty(\mathcal{FQ}, \mathcal{D})) \\ \downarrow A_1(\mathcal{Q}, _) & = & \downarrow (1 \boxtimes \text{restr}_\mathcal{D})M \\ A_\infty^u(A_1(\mathcal{Q}, \mathcal{C}), A_1(\mathcal{Q}, \mathcal{D})) & \xrightarrow{(\text{restr}_\mathcal{C} \boxtimes 1)M} & A_\infty^u(A_\infty(\mathcal{FQ}, \mathcal{C}), A_1(\mathcal{Q}, \mathcal{D})) \end{array} \quad (3.4.1)$$

commutes. Notice that all arrows in this diagram are strict A_∞ -functors. Indeed, $A_\infty(\mathcal{FQ}, _)$ and $A_1(\mathcal{Q}, _)$ are strict by [Lyu03, Proposition 6.2]. For an arbitrary A_∞ -functor f the components $[(f \boxtimes 1)M]_n = (f \boxtimes 1)M_{0n}$ vanish for all n except for $n = 1$, thus, $(f \boxtimes 1)M$ is strict. The A_∞ -functor $g = \text{restr}_\mathcal{D}$ is strict, hence, the n -th component

$$[(1 \boxtimes g)M]_n : r^1 \otimes \cdots \otimes r^n \mapsto (r^1 \otimes \cdots \otimes r^n \mid g)M_{n0}$$

of the A_∞ -functor $(1 \boxtimes g)M$ satisfies the equation

$$[(r^1 \otimes \cdots \otimes r^n \mid g)M_{n0}]_k = (r^1 \otimes \cdots \otimes r^n)\theta_{k1}g_1.$$

If the right hand side does not vanish, then $n \leq 1 \leq k + n$, so $n = 1$ and $(1 \boxtimes g)M$ is strict.

Given an A_∞ -transformation $t : g \rightarrow h : \mathcal{C} \rightarrow \mathcal{D}$ between unital A_∞ -functors we find that

$$\begin{aligned} A_\infty(\mathcal{FQ}, _)(t) &= [(1 \boxtimes t)M : (1 \boxtimes g)M \rightarrow (1 \boxtimes h)M : A_\infty(\mathcal{FQ}, \mathcal{C}) \rightarrow A_\infty(\mathcal{FQ}, \mathcal{D})], \\ A_1(\mathcal{Q}, _)(t) &= [(1 \boxtimes t)M : (1 \boxtimes g)M \rightarrow (1 \boxtimes h)M : A_1(\mathcal{Q}, \mathcal{C}) \rightarrow A_1(\mathcal{Q}, \mathcal{D})], \\ [(1 \boxtimes \text{restr}_\mathcal{D})M]A_\infty(\mathcal{FQ}, _)(t) &= [((1 \boxtimes t)M) \cdot \text{restr}_\mathcal{D} : ((1 \boxtimes g)M) \cdot \text{restr}_\mathcal{D} \\ &\quad \rightarrow ((1 \boxtimes h)M) \cdot \text{restr}_\mathcal{D} : A_\infty(\mathcal{FQ}, \mathcal{C}) \rightarrow A_1(\mathcal{Q}, \mathcal{D})], \\ [(\text{restr}_\mathcal{C} \boxtimes 1)M]A_1(\mathcal{Q}, _)(t) &= [\text{restr}_\mathcal{C} \cdot ((1 \boxtimes t)M) : \text{restr}_\mathcal{C} \cdot ((1 \boxtimes g)M) \\ &\quad \rightarrow \text{restr}_\mathcal{C} \cdot ((1 \boxtimes h)M) : A_\infty(\mathcal{FQ}, \mathcal{C}) \rightarrow A_1(\mathcal{Q}, \mathcal{D})]. \end{aligned}$$

We have to verify that the last two A_∞ -transformations are equal. First of all, let us show that mappings of objects in (3.4.1) commute. Given a unital A_∞ -functor $g : \mathcal{C} \rightarrow \mathcal{D}$, we are going to check that

$$[(1 \boxtimes g)M]_n \cdot \text{restr}_1 = \text{restr}_1^{\otimes n} \cdot [(1 \boxtimes g)M]_n \quad (3.4.2)$$

for any $n \geq 1$. Indeed, for any n -tuple of composable A_∞ -transformations

$$f^0 \xrightarrow{r^1} f^1 \longrightarrow \cdots \xrightarrow{r^n} f^n : \mathcal{FQ} \rightarrow \mathcal{C},$$

we have in both cases

$$\begin{aligned}
\{(r^1 \otimes \cdots \otimes r^n)[(1 \boxtimes g)M]_n\}_0 &= [(r^1 \otimes \cdots \otimes r^n|g)M_{n0}]_0 = (r_0^1 \otimes \cdots \otimes r_0^n)g_n, \\
\{(r^1 \otimes \cdots \otimes r^n)[(1 \boxtimes g)M]_n\}_1 &= [(r^1 \otimes \cdots \otimes r^n|g)M_{n0}]_1 \\
&= \sum_{i=1}^n (r_0^1 \otimes \cdots \otimes r_0^{i-1} \otimes r_1^i \otimes r_0^{i+1} \otimes \cdots \otimes r_0^n)g_n \\
&\quad + \sum_{i=1}^n (r_0^1 \otimes \cdots \otimes r_0^{i-1} \otimes f_1^i \otimes r_0^{i+1} \otimes \cdots \otimes r_0^n)g_{n+1}.
\end{aligned}$$

Note that the right hand sides depend only on 0-th and 1-st components of r^i , f^i . This is precisely what is claimed by equation (3.4.2).

The coincidence of A_∞ -transformations $((1 \boxtimes t)M) \cdot \text{restr}_{\mathcal{D}} = \text{restr}_{\mathcal{C}} \cdot ((1 \boxtimes t)M)$ follows similarly from the computation:

$$\begin{aligned}
\{(r^1 \otimes \cdots \otimes r^n)[(1 \boxtimes t)M]_n\}_0 &= [(r^1 \otimes \cdots \otimes r^n \boxtimes t)M_{n1}]_0 = (r_0^1 \otimes \cdots \otimes r_0^n)t_n, \\
\{(r^1 \otimes \cdots \otimes r^n)[(1 \boxtimes t)M]_n\}_1 &= [(r^1 \otimes \cdots \otimes r^n \boxtimes t)M_{n1}]_1 \\
&= \sum_{i=1}^n (r_0^1 \otimes \cdots \otimes r_0^{i-1} \otimes r_1^i \otimes r_0^{i+1} \otimes \cdots \otimes r_0^n)t_n \\
&\quad + \sum_{i=1}^n (r_0^1 \otimes \cdots \otimes r_0^{i-1} \otimes f_1^i \otimes r_0^{i+1} \otimes \cdots \otimes r_0^n)t_{n+1}.
\end{aligned}$$

3.5. Representability. An A_∞^u -2-functor $F : A_\infty^u \rightarrow A_\infty^u$ is called *representable*, if it is naturally A_∞^u -2-equivalent to the A_∞^u -2-functor $A_\infty(\mathcal{A}, -) : A_\infty^u \rightarrow A_\infty^u$ for some A_∞ -category \mathcal{A} . The above results imply that the A_∞^u -2-functor $A_1(\mathcal{Q}, -)$ corresponding to a differential graded \mathbb{k} -quiver \mathcal{Q} is represented by the free A_∞ -category \mathcal{FQ} generated by \mathcal{Q} .

This definition of representability has a disadvantage: many different A_∞ -categories can represent the same A_∞^u -2-functor. More attractive notion is the following. An A_∞^u -2-functor $F : A_\infty^u \rightarrow A_\infty^u$ is called *unitally representable*, if it is naturally A_∞^u -2-equivalent to the A_∞^u -2-functor $A_\infty^u(\mathcal{A}, -) : A_\infty^u \rightarrow A_\infty^u$ for some unital A_∞ -category \mathcal{A} . Such \mathcal{A} is unique up to an A_∞ -equivalence. Indeed, composing a natural 2-equivalence $\bar{\lambda} : A_\infty^u(\mathcal{A}, -) \rightarrow A_\infty^u(\mathcal{B}, -) : \overline{A_\infty^u} \rightarrow \overline{A_\infty^u}$ with the 0-th cohomology 2-functor $H^0 : \overline{A_\infty^u} \rightarrow \mathcal{Cat}$, we get a natural 2-equivalence $H^0\bar{\lambda} : H^0\overline{A_\infty^u}(\mathcal{A}, -) \rightarrow H^0\overline{A_\infty^u}(\mathcal{B}, -) : \overline{A_\infty^u} \rightarrow \mathcal{Cat}$. However, $H^0\overline{A_\infty^u}(\mathcal{A}, -) = \overline{A_\infty^u}(\mathcal{A}, -)$, so using a 2-category version of Yoneda lemma one can deduce that \mathcal{A} and \mathcal{B} are equivalent in $\overline{A_\infty^u}$. We shall present an example of unital representability in subsequent publication [LM04].

3.6. Acknowledgements. We are grateful to all the participants of the A_∞ -category seminar at the Institute of Mathematics, Kyiv, for attention and fruitful discussions, especially to Yu. Bєspalov and S. Ovsienko. One of us (V.L.) is grateful to Max-Planck-Institut für Mathematik for warm hospitality and support at the final stage of this research.

References

- [Fuk93] K. Fukaya, *Morse homotopy, A_∞ -category, and Floer homologies*, Proc. of GARC Workshop on Geometry and Topology '93 (H. J. Kim, ed.), Lecture Notes, no. 18, Seoul Nat. Univ., Seoul, 1993, pp. 1–102.
- [Fuk02] K. Fukaya, *Floer homology and mirror symmetry. II*, Minimal surfaces, geometric analysis and symplectic geometry (Baltimore, MD, 1999), Adv. Stud. Pure Math., vol. 34, Math. Soc. Japan, Tokyo, 2002, pp. 31–127.
- [Kad82] T. V. Kadeishvili, *The algebraic structure in the homology of an $A(\infty)$ -algebra*, Soobshch. Akad. Nauk Gruzin. SSR **108** (1982), no. 2, 249–252, in Russian.
- [Kel01] B. Keller, *Introduction to A -infinity algebras and modules*, Homology, Homotopy and Applications **3** (2001), no. 1, 1–35, [arXiv:math.RA/9910179](http://arxiv.org/abs/math.RA/9910179), <http://intlpress.com/HHA/v3/n1/a1/>.
- [Kon95] M. Kontsevich, *Homological algebra of mirror symmetry*, Proc. Internat. Cong. Math., Zürich, Switzerland 1994 (Basel), vol. 1, Birkhäuser Verlag, 1995, 120–139.
- [KS02] M. Kontsevich and Y. S. Soibelman, *A_∞ -categories and non-commutative geometry*, in preparation, 2002.
- [KS] M. Kontsevich and Y. S. Soibelman, *Deformation theory*, book in preparation.
- [LH03] K. Lefèvre-Hasegawa, *Sur les A_∞ -catégories*, Ph.D. thesis, Université Paris 7, U.F.R. de Mathématiques, 2003, [arXiv:math.CT/0310337](http://arxiv.org/abs/math.CT/0310337).
- [LM04] V. V. Lyubashenko and O. Manzyuk, *Quotients of unital A_∞ -categories*, 2004, [arXiv:math.CT/0306018](http://arxiv.org/abs/math.CT/0306018).
- [LO02] V. V. Lyubashenko and S. A. Ovsienko, *A construction of quotient A_∞ -categories*, Homology, Homotopy Appl. **8** (2006), no. 2, 157–203, [arXiv:math.CT/0211037](http://arxiv.org/abs/math.CT/0211037) <http://intlpress.com/HHA/v8/n2/a9/>.
- [Lyu03] V. V. Lyubashenko, *Category of A_∞ -categories*, Homology, Homotopy and Applications **5** (2003), no. 1, 1–48, [arXiv:math.CT/0210047](http://arxiv.org/abs/math.CT/0210047), <http://intlpress.com/HHA/v5/n1/a1/>.
- [Sta63] J. D. Stasheff, *Homotopy associativity of H -spaces, I & II*, Trans. Amer. Math. Soc. **108** (1963), 275–292, 293–312.

Contents

| | | |
|----------|--|-----------|
| 1 | Conventions and preliminaries | 2 |
| 1.1 | A_N -categories. | 3 |
| 1.7 | Trees. | 8 |
| 2 | Properties of free A_∞-categories | 9 |
| 2.1 | Construction of a free A_∞ -category. | 9 |
| 2.6 | Explicit formula for the constructed strict A_∞ -functor. | 15 |
| 2.7 | Transformations between functors from a free A_∞ -category. | 15 |
| 2.11 | Restriction as an A_∞ -functor. | 22 |
| 3 | Representable 2-functors $A_\infty^u \rightarrow A_\infty^u$ | 24 |
| 3.3 | Examples of A_∞^u -2-functors. | 27 |
| 3.4 | Example of an A_∞^u -2-equivalence. | 28 |
| 3.5 | Representability. | 30 |
| 3.6 | Acknowledgements. | 30 |