### A Note on the Cohomology Ring of Spherical Varieties

Kiumars Kaveh Department of Mathematics University of British Columbia

December 7, 2018

**Abstract**. Let X be a (smooth) projective spherical G-variety. We give a combinatorial description of the subalgebra of the cohomology ring  $H^*(X, \mathbb{Q})$  generated by  $H^2(X, \mathbb{Q})$ , in terms of the volume of polyhedra. This generalizes the Khovanksii-Pukhlikov description of the cohomology ring of toric varieties. In particular, we obtain a unified description for the cohomology rings of the flag variety G/B and toric varieties.

Key words: Spherical Variety, flag variety, toric variety, cohomology ring, Newton polytope, Gelfand-Cetlin polytope. Subject Classification: 14M17.

#### Contents

1	Intr	oduction	<b>2</b>
<b>2</b>	АТ	Theorem about Commutative Algebras	3
3	Subalgebra of Cohomology Generated by $H^2(X, \mathbb{Q})$		6
	3.1	Poincare Duality for the Subalgebra Generated by $H^2(X, \mathbb{Q})$ .	6
	3.2	Degree of a Spherical Variety	8
	3.3	Gelfand-Cetlin Polytopes, Newton Polytope and Degree of a	
		Spherical Variety	8
	3.4	Additivity of the Moment Polytope	10
	3.5	Main Theorem	12

<b>4</b>	Examples		
	4.1	Cohomology Ring of Toric Varieties	13
	4.2	Cohomology Ring of the Flag Variety $G/B$	14

#### 1 Introduction

The well-known theorem of Bernstein and Kušnirenko (see [BKK]) on the number of solutions of a system of Laurent polynomial equations in terms of volumes of their Newton polyhedra, can be interpreted as giving a formula for the intersection numbers of divisors in a projective toric variety. Khovanskii and Pukhlikov (see [Kh-P, §1.4]) observed that the Bernstein-Kušnirenko theorem, in fact, completely determines the cohomology ring of any (smooth) projective toric variety (see Theorem 4.1). For this, it is important that the cohomology ring of a (smooth) projective toric variety is generated in degree 2.

Bernstein-Kušnirenko theorem has been generalized to reductive groups by B. Kazarnovskii (see [Kaz]) and more generally, to spherical varieties by M. Brion (see [Br2]). Let G be a complex connected reductive group. A normal G-variety X is called *spherical*, if a Borel subgroup of G has a dense open orbit. Let  $X \hookrightarrow \mathbb{P}(V)$  be a spherical G-variety X of dimension n. Here V is a finite dimensional G-module. In [Br2, §4.1], Brion gives a formula for the degree of X in  $\mathbb{P}(V)$  as the integral of an explicitly defined function over the moment polytope  $\mu(X)$  (see Theorem 3.3). If L is the line bundle on X obtained by the restriction of the universal subbundle on  $\mathbb{P}(V)$  to X, the degree of X in  $\mathbb{P}(V)$  is just  $c_1(L)^n$  where  $c_1$  denote the first Chern class. Let  $L_1, \ldots, L_n$  be n very ample G-line bundles on X. Brion's formula readily gives a formula for  $c_1(L_1) \cdots c_1(L_n)$  generalizing Bernstein-Kušnirenko theorem.

When G is a classical group over  $\mathbb{C}$ , A. Okounkov [Ok] associates a polytope  $\Delta(X)$  to a projective spherical G-variety  $X \hookrightarrow \mathbb{P}(V)$ , where V is a finite dimensional G-module. Using the above result of Brion, he shows that  $\deg(X)$  in  $\mathbb{P}(V)$  is equal to  $n! \operatorname{Vol}(\Delta)$ . The polytope  $\Delta(X)$  is the polytope fibred over the moment polytope of X with the Gelfand-Cetlin polytopes as fibers. It follows that the product of Chern classes of n line bundles  $L_1, \ldots, L_n$  over X can be computed as a mixed volume of polyhedra.

It is known that any spherical variety has a paving by affine spaces and hence does not have any odd cohomology. Unlike the toric case, it can be shown that the cohomology ring of a (smooth) projective spherical variety, in general, is not generated in degree 2 (for example the Grassmanian of 2-planes in  $\mathbb{C}^5$ ). In this note, using Brion's (and Okounkov's) formula for the degree, we give a description of the subalgebra of the cohomology ring of a (smooth) projective spherical variety generated by 2-cocycles in terms of the volume of polyhedra. This is the direct generalization of Khovanskii-Pukhlikov description of the cohomology ring of projective toric varieties.

As a minor but interesting observation in the course of the work, we show that the moment polytope is additive. More precisely, let  $\mu(L)$  denote the moment polytope of X in the embedding given by a very ample G-line bundle L. Let  $L_1$  and  $L_2$  be two very ample G-line bundles over X. Then  $\mu(L_1 \otimes L_2) = \mu(L_1) + \mu(L_2)$ , where the summation in the right-hand side is the Minkowski sum of convex polytopes. It then follows that the Okounkov's polytope  $\Delta$ , defined when G is a classical group, is also additive. That is if  $L_1$  and  $L_2$  are two very ample G-line bundles over X and  $\Delta(L_1)$  and  $\Delta(L_2)$  the Okounkov polytopes for the corresponding embeddings of X, then  $\Delta(L_1 \otimes L_2) = \Delta(L_1) + \Delta(L_2)$ .

From the Bruhat decomposition, flag varieties are spherical. It is known that the cohomology ring of the flag variety G/B is generated in degree 2. When G is a classical group, our theorem gives a description of the cohomology ring of the flag variety G/B in terms of the volume of the Gelfand-Cetlin polytopes. It is interesting to note that this way, we obtain similar descriptions for the cohomology rings of toric varieties and the flag variety G/B. It is possible, but non-trivial, to derive the usual description of the cohomology ring of G/B (due to Borel) from our description.

This note is closely related to the work of Guillemin and Sternberg in [GS].

**Acknowledgment:** The author would like to thank A.G. Khovanskii, E. Meinrenken and V. Timorin for helpful discussions.

#### 2 A Theorem about Commutative Algebras

Let A be a finite dimensional graded commutative algebra over a field k of characteristic zero. In this section we prove a theorem which says, if A is generated by  $A_1$ , the subspace of degree 1 elements, and satisfies few good properties, then the algebra A can be described as a quotient of the algebra of constant coefficient differential operators on the vector space  $A_1$ . Later we will apply this theorem to the subalgebra of the cohomology algebra of a (smooth) projective variety generated by 2-cocycles. This theorem is not new and is implicit in the work of Khovanskii-Pukhlikov (see [Kh-P]).

**Theorem 2.1.** Let A be a commutative finite dimensional graded algebra over a field k of characteristic zero. Let  $A = \bigoplus_{i=0}^{n} A_i$ , where  $A_i$  is the *i*-th graded piece of A.

Suppose the following conditions hold:

- (i) A is generated, as an algebra, by  $A_1$ .
- (*ii*)  $A_0 \cong A_n \cong k$ .
- (iii) The bilinear map  $A_i \times A_{n-i} \to A_n \cong k$  given by  $(x, y) \mapsto xy$  is nondegenerate for all i = 0, ..., n.

Let  $\{\mathbf{a_1}, \ldots, \mathbf{a_r}\}$  be a basis for  $A_1$  and  $x_1, \ldots, x_n$  the coordinates with respect to this basis. Let  $P : A_1 \to k$  be the polynomial defined by  $P(x) = x^n \in$  $A_n \cong k, \forall x \in A_1$ . Then A, as a graded algebra, is isomorphic to the algebra  $k[x_1, \ldots, x_r]/I$ , where I the ideal defined by

$$I = \{ f(x_1, \dots, x_r) \mid f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = 0 \},\$$

where  $f(\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_r})$  is the differential operator obtained by replacing  $x_i$  by  $\partial/\partial x_i$  in f.

Proof. It is straight forward to see that I is indeed an ideal of  $k[x_1, \ldots, x_r]$ . Next, consider the surjective homomorphism  $\Phi : k[x_1, \ldots, x_r] \to A$  given by  $f \mapsto f(\mathbf{a_1}, \ldots, \mathbf{a_r})$ , where in the righthand side f is regarded as a polynomial in A. We wish to prove that  $\ker(\Phi) = I$ . Note that from the definition, both I and  $\ker(\Phi)$  are generated by homogeneous elements. From the definition of the polynomial P we have:

$$P(\sum_{i=1}^{r} x_{i} \mathbf{a}_{i}) = (\sum_{i=1}^{r} x_{i} \mathbf{a}_{i})^{n}$$
$$= \sum_{\alpha_{1} + \dots + \alpha_{r} = n} n_{\alpha_{1}, \dots, \alpha_{r}} \mathbf{a}_{1}^{\alpha_{1}} \cdots \mathbf{a}_{r}^{\alpha_{r}} x_{1}^{\alpha_{1}} \cdots x_{r}^{\alpha_{r}},$$

where  $n_{\alpha_1,...,\alpha_r} = \frac{n!}{\alpha_1!\cdots\alpha_r!}$ . Let

$$f(x_1,\ldots,x_n) = \sum_{\beta_1+\cdots+\beta_r=n} c_{\beta_1,\ldots,\beta_r} x_1^{\beta_1} \cdots x_r^{\beta_r}$$

be a homogeneous polynomial of degree n. One can see that

$$f(\frac{\partial}{\partial x_1},\ldots,\frac{\partial}{\partial x_r})\cdot P = n! \sum_{\beta_1+\cdots+\beta_r=n} c_{\beta_1,\ldots,\beta_r} \mathbf{a_1}^{\beta_1}\cdots \mathbf{a_r}^{\beta_r}$$

Thus  $f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = 0$  if and only if  $f(\mathbf{a_1}, \dots, \mathbf{a_r}) = 0$ . Next, let

$$f(x_1,\ldots,x_n) = \sum_{\beta_1+\cdots+\beta_r=m} c_{\beta_1,\ldots,\beta_r} x_1^{\beta_1} \cdots x_r^{\beta_r}$$

be a homogeneous polynomial of degree m < n. Suppose  $f(\mathbf{a_1}, \ldots, \mathbf{a_r}) \neq 0$ . By assumption (ii), there exists a homogeneous polynomial  $g(x_1, \ldots, x_r)$  of degree (n-m) such that  $g(\mathbf{a_1}, \ldots, \mathbf{a_r}) f(\mathbf{a_1}, \ldots, \mathbf{a_r}) \neq 0$ . Since gf is a homogeneous polynomial of degree n, by what we proved,  $(gf)(\partial/\partial x_1, \ldots, \partial/\partial x_r) \cdot P \neq 0$  which implies that  $f(\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_r}) \cdot P \neq 0$ . So we showed that  $f \in I$  implies  $f \in \ker(\Phi)$ . Conversely, suppose  $f(\mathbf{a_1}, \ldots, \mathbf{a_r}) = 0$ , we have

$$f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = \sum_{\alpha_1 + \dots + \alpha_r = n; \beta_1 + \dots + \beta_r = m; \beta_1 \le \alpha_1, \dots, \beta_r \le \alpha_r} n_{\alpha_1, \dots, \alpha_r} \mathbf{a_1}^{\beta_1} \cdots \mathbf{a_r}^{\beta_r}$$
$$\frac{\alpha_1!}{(\alpha_1 - \beta_1)!} \cdots \frac{\alpha_r!}{(\alpha_r - \beta_r)!} c_{\beta_1, \dots, \beta_r} x_1^{\alpha_1 - \beta_1} \cdots x_r^{\alpha_r - \beta_r}$$

We can write the above expression as

$$f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = \left(\sum_{\substack{\beta_1 + \dots + \beta_r = m \\ \gamma_1 + \dots + \gamma_r = n - m; 0 \le \gamma_i \le n - \beta_i}} c_{\beta_1, \dots, \beta_r} \mathbf{a_1}^{\beta_1} \cdots \mathbf{a_r}^{\beta_r}\right) \cdot \left(\sum_{\substack{\gamma_1 + \dots + \gamma_r = n - m; 0 \le \gamma_i \le n - \beta_i \\ \gamma_1 ! \cdots \gamma_r !}} \frac{n!}{\gamma_1! \cdots \gamma_r!} (\mathbf{a_1} x_1)^{\gamma_1} \cdots (\mathbf{a_r} x_r)^{\gamma_r}\right) = 0.$$

which shows that  $f \in I$ . Hence we prove that  $\ker(\Phi) = I$  and thus  $A \cong k[x_1, \ldots, x_r]/I$ .

In Section 3.5, we will apply this theorem to a cohomology subalgebra of a (smooth) projective variety X with no odd cohomology. We take A to be the subalgebra generated by  $H^2(X, \mathbb{Q})$ . For each *i*, the *i*-th graded piece  $A_i$ is  $A \cap H^{2i}(X, \mathbb{Q})$ .

# 3 Subalgebra of Cohomology Generated by $H^2(X, \mathbb{Q})$

Let X be a (smooth) projective spherical G-variety of dimension n, where G is a connected complex reductive group. We recall that X is spherical if it is normal and a Borel subgroup of G has a dense orbit in X. In this section we apply Theorem 2.1 to  $H^*(X, \mathbb{Q})$  to obtain a description of the subalgebra A of  $H^*(X, \mathbb{Q})$  generated by  $H^2(X, \mathbb{Q})$ .

## 3.1 Poincare Duality for the Subalgebra Generated by $H^2(X, \mathbb{Q})$

Let X be a (smooth) projective variety of dimension n over  $\mathbb{C}$  and no odd cohomology. In this section, we show that the subalgebra A of the cohomology generated by  $H^2(X, \mathbb{Q})$  satisfies the Poincare duality. More precisely, let  $A_i \subset H^{2i}(X, \mathbb{Q})$  denote the *i*-th graded piece of A. Then the cup product pairing

$$A_i \times A_{n-i} \to \mathbb{Q}$$

is non-degenerate for  $i = 0, \ldots, n$ .

Suppose X is a subvariety of a (smooth) projective variety Y of dimension m over  $\mathbb{C}$  and no odd cohomology. Let  $i: X \hookrightarrow Y$  denote the inclusion map. The following lemma shows that the image of the cohomology algebra of Y in X satisfies the Poincare duality.

**Lemma 3.1.** The graded subalgebra  $i^*(H^*(Y, \mathbb{Q}))$  of  $H^*(X, \mathbb{Q})$  satisfies the Poincare duality, that is the cup product pairing

$$i^*(H^{2k}(Y,\mathbb{Q})) \times i^*(H^{2n-2k}(Y,\mathbb{Q})) \to \mathbb{Q}$$

is non-degenerate for all  $0 \le k \le n$ .

*Proof.* Let  $[X] \in H^{2m-2n}(Y, \mathbb{Q})$  denote the Poincare dual of X in Y. Let  $\sigma \in H^{2k}(Y, \mathbb{Q}), \ 0 \le k \le n$ , be a 2k-cocycle. Note that  $i^*(\sigma) = 0$  if and only if  $\sigma \cap [X] = 0$ . Suppose  $\sigma$  is such that

$$i^*(\sigma) \cap i^*(\tau) = 0, \quad \forall \tau \in H^{2n-2k}(Y, \mathbb{Q}).$$

Then,  $i^*(\sigma \cap \tau) = 0$  which means that  $(\sigma \cap \tau) \cap [X] = 0$  in  $H^*(Y, \mathbb{Q}), \forall \tau \in H^{2n-2k}(Y, \mathbb{C})$ . The Poincare duality for Y then implies that  $\sigma \cap [X] = 0$ . Hence  $i^*(\sigma) = 0$ , which proves that the above paring is non-degenerate.  $\Box$  It remains to show that X can be embedded in a (smooth) projective variety Y such that the image of the cohomology of Y in X is the cohomology subalgebra generated by  $H^2(X, \mathbb{Q})$ . Choose very ample line bundles  $L_1, \ldots, L_r$  over X such that  $c_1(L_1), \ldots, c_1(L_r)$  is a vector space basis for  $H^2(X, \mathbb{Q})$ , where  $c_1$  denotes the first Chern class. Each  $L_i$  gives rise to an embedding  $\phi_i : X \hookrightarrow \mathbb{P}(V_i)$ , where  $V_i$  is the finite dimensional vector space of sections of  $L_i$ . Let

$$Y = \prod_{i=1}^{r} \mathbb{P}(V_i).$$

Using the  $\phi_i$ , we get the diagonal embedding:

$$\phi: X \hookrightarrow Y.$$

Since Y does not have odd cohomology, by Künneth formula we have:

$$H^{2}(Y,\mathbb{Q}) = \bigoplus_{i=1}^{r} H^{2}(\mathbb{P}(V_{i}),\mathbb{Q}),$$

and hence

$$\phi^*(H^2(Y,\mathbb{Q})) = \operatorname{Span}_{\mathbb{Q}}(c_1(L_1),\ldots,c_1(L_r)) = H^2(X,\mathbb{Q}).$$

Now, since  $H^*(\mathbb{P}(V_i), \mathbb{Q})$  is generated, as a  $\mathbb{Q}$ -algebra, by  $c_1(L_i)$ , the cohomology algebra  $H^*(Y, \mathbb{Q})$  is generated by the images of  $c_1(L_1), \ldots, c_1(L_r)$ in  $H^*(Y, \mathbb{Q})$ . We conclude that

$$\phi^*(H^*(Y,\mathbb{Q})) = \text{subalgebra generated by } \phi^*(H^2(Y,\mathbb{Q})),$$
  
= subalgebra generated by  $H^2(X,\mathbb{Q}),$ 

i.e. the image of the cohomology of Y in X is the same as the subalgebra generated by  $H^2(X, \mathbb{Q})$ .

After applying Lemma 3.1 to  $\phi: X \hookrightarrow Y$ , we obtain:

**Proposition 3.2.** Let X be a (smooth) projective variety of dimension n over  $\mathbb{C}$  and no odd cohomology. The subalgebra A of  $H^*(X, \mathbb{Q})$  generated by  $H^2(X, \mathbb{Q})$  satisfies the Poincare duality, i.e. for any  $0 \le i \le n$ , the cup product pairing

$$A_i \times A_{n-i} \to \mathbb{Q}$$

is non-degenerate, where  $A_i \subset H^{2i}(X, \mathbb{Q})$  denotes the *i*-th graded piece of A.

#### **3.2** Degree of a Spherical Variety

Let X be a (smooth) projective spherical variety and L a very ample G-line bundle on X giving rise to an embedding  $X \hookrightarrow \mathbb{P}(V)$ , where V is the finite dimensional G-module of all the sections of L. In this section we state a nice combinatorial formula due to Brion for the degree of X in  $\mathbb{P}(V)$ . It generalizes the famous Kušnirenko-Bernstein theorem on the degree of toric varieties (see [BKK]). Before stating the result, we need a bit of notation. Fix a Borel subgroup B for G with a dense orbit . Let T be the maximal torus contained in B and t (respectively  $\mathfrak{t}^*$ ) be the Lie algebra of T (respectively dual of the Lie algebra of T). Equip X with the symplectic structure obtained by the restriction of the symplectic structure of  $\mathbb{P}(V)$  given by the Fubini-Study form. Let  $\mu(L)$  denote the moment polytope of X. Recall that the moment polytope is the intersection of the image of the moment map with the positive Weyl chamber for the choice of B. Denote by  $R^+$  the set of positive roots defined by the choice of B. Also, let  $d\gamma$  be the Lebesgue measure on  $\mathfrak{t}^*$ normalized so that the cell of the weight lattice has volume 1.

**Theorem 3.3** ([Br2], §4.1). The degree of X in  $\mathbb{P}(V)$  is equal to

$$n!\int_{\mu(L)}f(\gamma)d\gamma,$$

where

$$f(\gamma) = \prod_{\alpha \in R^+; \langle \gamma, \alpha \rangle \neq 0} \frac{\langle \gamma, \alpha \rangle}{\langle \rho, \alpha \rangle}$$

and  $\rho$  is half of the sum of positive roots.

#### 3.3 Gelfand-Cetlin Polytopes, Newton Polytope and Degree of a Spherical Variety

Let G be a classical group over  $\mathbb{C}$  and X a (smooth) projective spherical Gvariety of dimension n. In this section, following [Ok] we associate a polytope  $\Delta$  to a G-variety  $X \hookrightarrow \mathbb{P}(V)$ , where V is a finite dimensional G-module. In [Ok], as a corollary of a theorem of Brion, it is shown that the polytope  $\Delta(X)$ has the property that the Hilbert polynomial of the variety X coincides with the Ehrhardt polynomial of the polytope  $\Delta(X)$ . In particular, deg(X) in  $\mathbb{P}(V)$  is equal to  $n! \operatorname{Vol}(\Delta(X))$ . We call  $\Delta$  the Newton polytope of  $X \hookrightarrow \mathbb{P}(V)$ . We start by recalling the Gelfand-Cetlin polytopes. To each dominant weight  $\lambda$  of G, there corresponds a Gelfand-Cetlin (or briefly G-C) polytope  $\Delta_{\lambda}$ . The convex polytope  $\Delta_{\lambda}$  has the property that the number of integral points in  $\Delta_{\lambda}$  is equal to the dimension of the irreducible G-module  $V_{\lambda}$  with highest weight  $\lambda$ . The dimension of the Gelfand-Cetlin polytope is equal to the complex dimension of the maximal unipotent subgroup U of G, i.e.  $\frac{1}{2}(\dim(G) - \operatorname{rank}(G))$ .

Following [GC], we recall the construction of G-C polytopes for  $GL(n, \mathbb{C})$ . For the construction of G-C polytopes for the symplectic group and the orthogonal group see [BZ].

**Definition 3.4 (G-C polytope for GL(n, \mathbb{C})).** Let  $\lambda = (\lambda_1 \geq \cdots \geq \lambda_n)$  be a decreasing sequence of integers representing a dominant weight in  $GL(n, \mathbb{C})$ . The G-C polytope  $\Delta_{\lambda}$  is the set of all real numbers  $x_1, x_2, \ldots, x_{n-1}, y_1, \ldots, y_{n-2}, \ldots, z$ , such that the following inequalities hold:



means  $a \ge c \ge b$ .

If the components of the weight  $\lambda$  are real, we still can define  $\Delta_{\lambda}$  by the above inequalities. So we can extend the definition of  $\Delta_{\lambda}$  to all real  $\lambda$ .

**Lemma 3.5.** The assignment  $\lambda \mapsto \Delta_{\lambda}$  is linear, i.e.  $\Delta_{c\lambda} = c\Delta_{\lambda}$  for any positive c and,  $\Delta_{\lambda+\mu} = \Delta_{\lambda} + \Delta_{\mu}$ , where the addition in the righthand side is the Minkowski sum of convex polytopes.

*Proof.* The proof is immediate from the definition in each of the three cases of classical groups.  $\Box$ 

Now, let  $X \subset \mathbb{P}(V)$  be a (smooth) projective spherical *G*-variety and  $\mu(X)$  its moment polytope. As before, let  $\Lambda$  denote the weight lattice and  $\Lambda_{\mathbb{Q}} \subset \mathfrak{t}^*$  the real vector space spanned by  $\Lambda$ .

Definition 3.6 (Newton polytope of a spherical variety). Define the set  $\Delta(X) \subset \Lambda_{\mathbb{Q}} \oplus \mathbb{R}^{\dim U} = \mathbb{R}^{\dim B}$ , by

$$\Delta(X) = \bigcup_{\lambda \in \mu(X)} (\lambda, \Delta_{\lambda}).$$

From Lemma 3.5, it follows that  $\Delta(X)$  is a convex polytope.

**Proposition 3.7 (See [Ok]).** With notation as above,  $\deg(X)$  in  $\mathbb{P}(V) = n! \operatorname{Vol}(\Delta(X))$ .

#### **3.4** Additivity of the Moment Polytope

Let X be a (smooth) projective (spherical?) G-variety. Let L be a very ample G-line bundle over X and  $i: X \hookrightarrow \mathbb{P}(V)$  the corresponding embedding, where V is the finite dimensional G-module of the sections of L. Let  $\mu_L: X \to \mathfrak{g}^*$  (respectively  $\mu(L)$ ) denote the moment map (respectively moment polytope) of X explained in the beginning of 3.2. In this section we wish to show that the moment polytope  $\mu(L)$  depends on L additively. That is, the moment polytope of the tensor product of two line bundles, is the Minkowski sum of the moment polytopes.

Let S be the semi-group of the very ample G-line bundles over X with respect to tensor product. In this section we wish to prove the following:

**Theorem 3.8.** Let X be a projective spherical variety. Then

 $\mu(L_1 \otimes L_2) = \mu(L_1) + \mu(L_2) \quad \forall L_1, L_2 \in S,$ 

where the addition in the left-hand side is the Minkowski sum of the convex polytopes.

*Proof.* Let V be finite dimensional G-module and  $X \hookrightarrow \mathbb{P}(V)$  a G-equivariant embedding of X. Fix a maximal compact subgroup K for G and let  $\mathfrak{k}$  (respectively  $\mathfrak{k}^*$ ) denote the Lie algebra of K (respectively dual of the Lie algebra of K). Choose a K-invariant Hermitian product  $\langle, \rangle$  on V. The moment map  $\mu : \mathbb{P}(V) \to \mathfrak{k}^*$  for the Hamiltonian action of K on  $\mathbb{P}(V)$ , and in particular X, is defined by

$$\mu(x)(\xi) = \frac{\langle \xi \cdot x, x \rangle}{\langle x, x \rangle}, \quad x \in \mathbb{P}(V), \xi \in \mathfrak{k}.$$

Now, let  $X \hookrightarrow \mathbb{P}(V_i)$  be the embeddings corresponding to the  $L_i$ . It can be shown that the line bundle defined on X by restricting the universal subbundle on  $\mathbb{P}(V_1 \otimes V_2)$  is  $L_1 \otimes L_2$ . Fix K-invariant Hermitian products on  $V_1$  and  $V_2$ . They define a Hermitian product on  $V_1 \otimes V_2$  by defining

$$\langle v_1 \otimes v_2, w_1 \otimes w_2 \rangle = \langle v_1, w_1 \rangle \langle v_2, w_2 \rangle, \quad v_i, w_i \in V_i \ (i = 1, 2).$$

Let  $\mu_i$ , i = 1, 2 (respectively  $\mu$ ) denote the moment map for the action of Kon  $\mathbb{P}(V_i)$  (respectively  $\mathbb{P}(V_1 \otimes V_2)$ ). Also, let  $\mu(L_i)$ , i = 1, 2 and  $\mu(L_1 \otimes L_2)$ denote the corresponding moment polytopes. From the above, we have the following for these moment maps:

$$\mu(x_1 \otimes x_2)(\xi) = \frac{\langle \xi \cdot (x_1 \otimes x_2), x_1 \otimes x_2 \rangle}{\langle x_1 \otimes x_2, x_1 \otimes x_2 \rangle}$$
  
= 
$$\frac{\langle (\xi \cdot x_1) \otimes x_2 + x_1 \otimes (\xi \cdot x_2), x_1 \otimes x_2 \rangle}{\langle x_1, x_1 \rangle \langle x_2, x_2 \rangle}$$
  
= 
$$\frac{\langle \xi \cdot x_1, x_1 \rangle}{\langle x_1, x_1 \rangle} + \frac{\langle \xi \cdot x_2, x_2 \rangle}{\langle x_2, x_2 \rangle}$$
  
= 
$$(\mu(x_1) + \mu(x_2))(\xi)$$

That is,  $\mu(L_1 \otimes L_2) = \mu(L_1) + \mu(L_2)$ .

Now we consider the situation in Section 3.3. Let G be one of the classical groups  $GL(n, \mathbb{C})$ ,  $SP(2n, \mathbb{C})$  or  $O(n, \mathbb{C})$ . Let X be a (smooth) projective spherical G-variety. Let L be a very ample G-line bundle on X. Denote by  $\Delta(L)$  the Newton polytope of X and L, defined in Definition 3.6.

**Corollary 3.9.** Let X be a projective spherical G-variety and  $L_i(i = 1, 2)$  two very ample G-line bundles over X. Then

$$\Delta(L_1 \otimes L_2) = \Delta(L_1) + \Delta(L_2),$$

where the addition in the left-hand side is the Minkowski sum of the convex polytopes.

*Proof.* For a dominant weight  $\lambda$ , from Lemma 3.5, the dependence of the G-C polytope  $\Delta_{\lambda}$  on  $\lambda$  is linear. The corollary now follows immediately from Theorem 3.8 and the definition of  $\Delta(L)$ .

#### 3.5 Main Theorem

As before, let X be a (smooth) projective spherical variety of dimension n. We want to combine the above formulae for the degree (Theorem 3.3 and Proposition 3.7) and Theorem 2.1 to give a description of the subalgebra Aof the cohomology generated by the 2-cocycles.

In Theorem 2.1, let  $k = \mathbb{Q}$  and A be the subalgebra of the cohomology generated by 2-cocycles. We take  $A_i$ , the *i*-th graded piece of A be  $A \cap$  $H^{2i}(X,\mathbb{Q})$ . Since X is (smooth) projective and has no odd cohomology, by Theorem 3.2, A satisfies the conditions in the statement of Theorem 2.1. Let  $P(x) = x^n, x \in A_1 = H^2(X,\mathbb{Q})$ , be the homogeneous polynomial of degree n in Theorem 2.1.

It is known, by the theory of Bialynicki-Birula (see [BB]), that X has a paving by the affine spaces and hence  $H^2(X, \mathbb{Z}) \cong \operatorname{Pic}(X)$ . Let S denote the set of all very ample line bundles on X. It is a semi-group in  $\operatorname{Pic}(X)$  and generates  $H^2(X, \mathbb{Q})$  as a vector space. It suffices to know the polynomial P on S. Also, let S' denote the set of all very ample G-line bundles on X, that is, S' is the set of all  $(L, \rho)$  where L is a very ample line bundle on X and  $\rho$  is a linear action of G on L compatible with the action of G on X. It is known that for any line bundle over X, the action of G on X can be lifted to a linear action of G on L (not necessarily in a unique way). Thus, one has a surjective homomorphism  $\pi$  from S' to S. Let  $(L, \rho)$  be a very ample G-line bundle. Let  $X \hookrightarrow \mathbb{P}(V)$  be the corresponding embedding of X, where V is the G-module of all the sections of X with the G-action given by  $\rho$ . From Theorem 3.3, we have

$$P(c_1(L)) = (c_1(L))^n$$
  
= deg(X) in  $\mathbb{P}(V)$   
=  $n! \int_{\mu(L)} f(\gamma) d\gamma$ ,

where  $f : \mathfrak{t}^* \to \mathbb{R}$  is the function defined in Theorem 3.3. When G is a classical group, from Proposition 3.7, this takes the simpler form

$$P(c_1(L)) = n! \operatorname{Vol}(\Delta(L)).$$

Recall (Corollary 3.9) that the map  $L \mapsto \Delta(L)$  from the semigroup of very ample line bundles to the semigroup of convex polytopes is additive. Now the above formula for P means that the homogeneous degree n polynomial P on  $H^2(X, \mathbb{Q})$  is the composition of the linear map  $L \mapsto \Delta(L)$  and the homogeneous degree n polynomial Vol on the set of convex polytopes.

From Theorem 2.1 we have:

**Theorem 3.10.** Let X be a (smooth) projective spherical variety. Let  $x_1, \ldots, x_r$ denote the coordinates for  $H^2(X, \mathbb{Q})$  in some basis. Let P be the polynomial defined above. Then, the subalgebra A of  $H^*(X, \mathbb{Q})$  generated by  $H^2(X, \mathbb{Q})$ is isomorphic to the algebra  $\mathbb{Q}[x_1, \ldots, x_r]/I$  and I is the ideal defined by

$$I = \{ f(x_1, \dots, x_r) \mid f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = 0 \},\$$

where  $f(\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_r})$  is the differential operator obtained by replacing  $x_i$  with  $\partial/\partial x_i$  in f.

#### 4 Examples

#### 4.1 Cohomology Ring of Toric Varieties

Let  $G = T = (\mathbb{C} \setminus \{0\})^n$  be a torus and, X a (smooth) projective T-toric variety. It is well-known that X is determined by a fan  $\Sigma \subset \mathfrak{t}$ , where t is the Lie algebra of T. Every convex lattice polytope  $\Delta \subset \mathfrak{t}^*$  normal to the fan  $\Sigma$ determines a line bundle  $L_{\Delta}$  on X. Here  $\mathfrak{t}^*$  denotes the dual of  $\mathfrak{t}$ . The set of convex lattice polytopes normal to  $\Sigma$  form a semi-group S with respect to the Minkowski sum of convex sets. Let us denote by  $\mathcal{V}$  the vector space generated by S over  $\mathbb{Q}$ . One can identify  $\mathcal{V}$  with  $\mathbb{Q}^r$  where r = number of 1-dimensional cones in  $\Sigma$ . One has a natural isomorphism between  $\operatorname{Pic}(X) \cong S/ \sim$  and  $H^2(X, \mathbb{Q}) \cong \mathcal{V}/ \sim$ , where  $\Delta \sim \Delta'$  if they can be identified by a translation.

From Bernstein-Kušnirenko theorem one has

 $\deg(X)$  embedded in  $\mathbb{P}(V_{\Delta}) = n! Vol(\Delta)$ ,

where  $V_{\Delta}$  is the vector space of all sections of  $L_{\Delta}$  (note that this is a special case of Theorem 3.3). The function volume extends to a homogeneous polynomial P of degree n on  $\mathcal{V}$ , which we denote by P. It is well-known that the cohomology ring of a toric variety is generated in degree 2. From Theorem 3.10, we then obtain the following description of  $H^*(X, \mathbb{Q})$ :

**Theorem 4.1 (Khovaksii-Pukhlikov, [Kh-P] Theorem §1.4).** With notation as above, we have: the cohomology algebra  $H^*(X, \mathbb{Q})$  is isomorphic to the algebra  $\mathbb{Q}[x_1, \ldots, x_r]/I$ , where I the ideal defined by

$$I = \{ f(x_1, \dots, x_r) \mid f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = 0 \}$$

where P(x) is the homogeneous polynomial of degree n on  $\mathcal{V}$  defined by

$$P(\Delta) = \operatorname{Vol}(\Delta),$$

for a polytope  $\Delta \in \mathcal{V}$ .

One can recover the usual description of the cohomology ring of a toric variety by generators and relations, due to Danilov-Jurkiewicz, from the above theorem (see [Timorin]).

#### 4.2 Cohomology Ring of the Flag Variety G/B

Let G be a classical group and B its standard Borel subgroup. In this section, we obtain a description of the cohomology ring of G/B, similar to the description of the cohomology ring of toric varieties, using Theorem 3.10.

Let  $\mathfrak{t}^*$  denote the dual of the Lie algebra of the maximal torus  $T \subset B$ . Let  $\Lambda \subset \mathfrak{t}^*$  be the weight lattice and  $t^*_+$  the positive Weyl chamber for the choice of B. For every weight  $\lambda$ , let  $V_{\lambda}$  be the irreducible representation with highest weight  $\lambda$ .

Let X = G/B and  $n = \dim(X)$ . If  $\lambda$  is in the interior of the Weyl chamber, one can embed the flag variety G/B in  $\mathbb{P}(V_{\lambda})$  via the orbit of a highest weight vector. Let  $L_{\lambda}$  denote the line bundle on G/B obtained by restricting the universal subbundle of the projective space  $\mathbb{P}(V_{\lambda})$  to G/B. It is known that the map  $\lambda \mapsto L_{\lambda}$  extends to an isomorphism of  $\Lambda$  and  $\operatorname{Pic}(G/B)$ , and consequently an isomorphism of  $\mathfrak{t}^*$  and  $H^2(G/B, \mathbb{Q})$ . It is also known that  $H^*(G/B, \mathbb{Q})$  is generated in degree 2. Hence we conclude that  $H^*(G/B, \mathbb{Q})$  is generated by  $c_1(L_{\lambda})$  where  $\lambda$  is in the interior of the Weyl chamber. Recall that  $\operatorname{deg}(G/B) \hookrightarrow \mathbb{P}(V_{\lambda})$  is equal to  $n!\operatorname{Vol}(\Delta_{\lambda})$ , where  $\Delta_{\lambda}$  is the Gelfand-Cetlin polytope of  $\lambda$ . Note that from Lemma 3.5, the map  $\lambda \mapsto \Delta_{\lambda}$  is linear. From Theorem 3.10 we have the following description of the cohomology ring of X: **Theorem 4.2.** Let G be a classical group over  $\mathbb{C}$  and  $r = \operatorname{rank}(G)$ . With notation as above we have: the cohomology algebra  $H^*(G/B, \mathbb{Q})$  is isomorphic to the algebra  $\mathbb{Q}[x_1, \ldots, x_r]/I$ , where I is the ideal defined by

$$I = \{ f(x_1, \dots, x_r) \mid f(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_r}) \cdot P = 0 \},\$$

and P is the homogeneous polynomial of degree n on  $t^*$  defined by

$$P(\lambda) = \operatorname{Vol}(\Delta_{\lambda})$$

for any  $\lambda$  in the interior of the Weyl chamber.

It is interesting to note the similarity between this description of  $H^*(G/B, \mathbb{Q})$ and the description of the cohomology ring of toric varieties (Theorem 4.1).

**Remark:** We would like to point out that it is possible but non-trivial to derive the usual description of the cohomology ring of the flag variety G/B due to Borel, from the above theorem. According to Borel's theorem  $H^*(G/B, \mathbb{Q})$  is isomorphic to  $\mathbb{Q}[x_1, \ldots, x_r]/J$  where J is the ideal generated by the W-invariant polynomials of positive degree and W is the Weyl group. The above theorem in fact asserts that I = J, i.e. a polynomial f is in the ideal generated by the W-invariant polynomials of positive degree if and only if  $f(\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_r}) \cdot P = 0$ . Even in the case of  $G = \operatorname{GL}(n, \mathbb{C})$  it is not trivial.

#### References

- [BKK] Bernstein, D. N.; Kušnirenko, A. G.; Khovanskiĭ, A. G. Newton polyhedra. Usp. Mat. Nauk. 31, No. 3, (1976) 201–202.
- [BB] Bialynicki-Birula, A. Some theorems on actions of algebraic groups Ann. of Math. 98 (1973) 480–497.
- [Br1] Brion, M. Sur l'image de l'application moment. Seminaire d'algebre Paul Dubreil et Marie-Paule Malliavin, Paris 1986 (Lecture notes in Mathematics 1296). Springer, Berlin, 1987. pp. 177-192.
- [Br2] Brion, M. Groupe de Picard et nombres caractristiques des varits sphriques. [Picard group and characteristic numbers of spherical varieties] Duke Math. J. 58 (1989), no. 2, 397–424.

- [BZ] Berenstein, A.D.; Zelevisnky, A. Tensor product multiplicities and convex polytopes in partition space, J. Geom. Phys. 5 (1988), no. 3, 453–472.
- [GC] Gelfand, I.M.; Cetlin, M.L. Finite dimensional representations of the group of unimodular matrices, Doklady Akad. Nauk USSR (N.S.) ,71 , (1950), 825–828.
- [GS] Guillemin, V.; Sternberg, S. The coefficients of the Duistermaat-Heckman polynomial and the cohomology ring of reduced spaces. Geometry, topology, & physics, 202–213, Conf. Proc. Lecture Notes Geom. Topology, IV, Internat. Press, Cambridge, MA, 1995.
- [Kaz] Kazarnovskii, B. Newton polyhedra and the Bezout formula for matrixvalued functions of finite dimensional representations. Functional Analysis and its applications, v. 21, no. 4, 73–74 (1987).
- [Kh-P] Pukhlikov, A. V.; Khovanskiĭ, A. G. The Riemann-Roch theorem for integrals and sums of quasipolynomials on virtual polytopes. Algebra i Analiz 4 (1992), no. 4, 188–216; translation in St. Petersburg Math. J. 4 (1993), no. 4, 789–812
- [Ok] Okounkov, A. A remark on the Hilbert polynomial of a spherical variety, Func. Anal. and Appl., 31 (1997), 82–85.
- [Timorin] Timorin, V. A. An analogue of the Hodge-Riemann relations for simple convex polyhedra. Uspekhi Mat. Nauk 54 (1999), no. 2(326), 113– 162; translation in Russian Math. Surveys 54 (1999), no. 2, 381–426

University of British Columbia, Vancouver, B.C. *Email address:* kaveh@math.ubc.ca