

# Isothermic submanifolds of Euclidean space

Ruy Tojeiro

**Abstract:** We study the problem posed by F. Burstall of developing a theory of isothermic Euclidean submanifolds of dimension greater than or equal to three. As a natural extension of the definition in the surface case, we call a Euclidean submanifold *isothermic* if it is locally the image of a conformal immersion of a Riemannian product of Riemannian manifolds whose second fundamental form is adapted to the product net of the manifold. Our main result is a complete classification of all such conformal immersions of Riemannian products of dimension greater than or equal to three. We derive several consequences of this result. We also study whether the classical characterizations of isothermic surfaces as solutions of Christoffel's problem and as envelopes of nontrivial conformal sphere congruences extend to higher dimensions.

**MSC 2000:** 53 C15, 53 C12, 53 C50, 53 B25.

**Key words:** *isothermic submanifold, conformal sphere congruence, Darboux transform, Christoffel transform, conformal immersion, isothermic net, Riemannian product.*

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries</b>	<b>6</b>
2.1	Nets and net morphisms . . . . .	6
2.2	Twisted and warped products . . . . .	7
2.3	Conformal product nets . . . . .	8
<b>3</b>	<b>Moebius geometry in the light cone</b>	<b>9</b>
3.1	Euclidean space as a hypersurface of the light cone . . . . .	9
3.2	Families of spheres and affine subspaces in Euclidean space . . . . .	11
<b>4</b>	<b>Orthogonal families of spheres in <math>\mathbb{R}^N</math></b>	<b>14</b>
<b>5</b>	<b>Extrinsic spheres of Euclidean submanifolds</b>	<b>18</b>

<b>6</b>	<b>The decomposition theorem of Nölker</b>	<b>20</b>
6.1	Warped product representations of space forms . . . . .	20
6.2	Nölker's and Molzan's theorems . . . . .	22
<b>7</b>	<b>Conformal immersions of Riemannian products</b>	<b>24</b>
<b>8</b>	<b>Christoffel problem and Darboux transformations</b>	<b>33</b>
8.1	Codazzi tensors . . . . .	33
8.2	Christoffel problem. . . . .	35
8.3	Conformal sphere congruences . . . . .	37

# 1 Introduction

The classical theory of isothermic surfaces has received a recent renaissance in interest mostly due to its connection with the modern theory of integrable systems (cf. [BHPP], [Bu], [Ci], [CGS], [BDPT]). Recall that a surface in Euclidean space  $\mathbb{R}^3$  is *isothermic* if, away from umbilic points, its curvature lines form an isothermic net, that is, there exist locally conformal coordinates that diagonalize the second fundamental form. Standard examples include cylinders, cones, surfaces of revolution, quadrics, surfaces of constant mean curvature and, the concept being a conformally invariant one, the images of such surfaces by an inversion in  $\mathbb{R}^3$ .

Isothermic surfaces have a number of characterizations from various points of view. One of them comes with the problem posed by Christoffel [Ch] of determining the surfaces  $f: M^2 \rightarrow \mathbb{R}^3$  that admit locally a *dual* surface  $\mathcal{F}: M^2 \rightarrow \mathbb{R}^3$  with parallel tangent planes to those of  $f$  which induces the same conformal structure but opposite orientation on  $M^2$ . It was shown by Christoffel himself that this property characterizes isothermic surfaces, and dual isothermic surfaces are now said to be *Christoffel transforms* one of each other.

Another characterization involves the notion of *Ribaucour transformation*. Classically, two surfaces in  $\mathbb{R}^3$  are said to be related by a Ribaucour transformation when they are the enveloping surfaces of a two-parameter sphere congruence such that curvature lines correspond. It was proved by Darboux ([Da<sub>1</sub>]) that if two surfaces in  $\mathbb{R}^3$  are related by a *conformal* Ribaucour transformation that reverts orientation then they must be a *Darboux pair* of isothermic surfaces.

Let us conclude our (rather incomplete) list of characterizations of isothermic surfaces by mentioning the interesting fact that they are precisely the surfaces that admit an infinitesimal isometric deformation preserving the principal curvatures.

A nice account of classical aspects of the subject as well as from the perspective of the modern theory of integrable systems and loop groups was given in [Bu]. In the concluding remarks of that paper, a list of open problems is posed that suggests interesting guides for future research. We quote here the first of them:

*Is there any interesting theory of isothermic submanifolds of  $\mathbb{R}^n$  of dimension greater than two? The problem here is to find a suitable definition that is not too restrictive... One way forward might be to study submanifolds admitting a conformal sphere congruence.*

This article grew out as an attempt to give an answer to this question. The preliminary step of finding a suitable extension to higher dimensions of the notion of an isothermic net was considered in [To<sub>1</sub>]. This turned out to fit into the general problem of studying local decomposability properties of *orthogonal nets* on Riemannian manifolds. Recall that an orthogonal  $k$ -net  $\mathcal{E} = (E_i)_{i=1,\dots,k}$  on a Riemannian manifold  $M$  is a splitting  $TM = \oplus_{i=1}^k E_i$  of the tangent bundle of  $M$  into a family of mutually orthogonal integrable subbundles. Looking for an extension of the notion of an isothermic net amounts to finding geometric properties of an orthogonal  $k$ -net  $\mathcal{E} = (E_i)_{i=1,\dots,k}$  implying that each point  $p \in M$  lies in an open neighborhood  $U$  such that there exists a conformal diffeomorphism of a Riemannian product  $\Pi_{i=1}^k M_i$  onto  $U$  that maps the leaves of the product foliation induced by  $M_i$  onto the leaves of  $E_i$ ,  $1 \leq i \leq k$ . This was achieved in Theorem 12 of [To<sub>1</sub>] (see Theorem 7 below), which may be regarded as a conformal version of the local decomposition theorem of de Rham. Following a more suggestive terminology used in [MRS] for other types of orthogonal nets, higher dimensional isothermic nets were named *conformal product nets* in [To<sub>1</sub>], or *CP*-nets for short.

The next step is to look for the appropriate extension of the property that *the curvature lines* form an isothermic net. In fact, our formulation of this condition in the beginning of the paper makes it natural to call a Euclidean submanifold *isothermic*, or more precisely, *k-isothermic*, if it carries a  $k$ -CP-net  $\mathcal{E} = (E_i)_{i=1,\dots,k}$  to which its second fundamental form is adapted, in the sense that each distribution  $E_i$  is invariant by all shape operators of the submanifold.

Summing things up, a submanifold is isothermic if it is locally given by a conformal immersion of a Riemannian product whose second fundamental form is adapted to the product net of the manifold. Thus, in studying isothermic Euclidean submanifolds of dimension  $n \geq 3$  we are faced with the following problem:

*Describe the conformal immersions of a Riemannian product of dimension  $n \geq 3$  into Euclidean space whose second fundamental forms are adapted to the product net of the manifold.*

The isometric version of this problem (in any dimension  $n \geq 2$ ) has led to a useful result due to Moore ([Mo]), who proved that an isometric immersion of a Riemannian product into Euclidean space whose second fundamental form is adapted to the product net of the manifold must be an extrinsic product of isometric immersions of the factors. Notice that the composition of such an extrinsic product of  $k$  isometric immersions with a conformal transformation in Euclidean space provides trivial examples of  $k$ -isothermic submanifolds for any value of  $k$ . These examples may be regarded as the

higher dimensional analogues of the images by an inversion of cylinders in  $\mathbb{R}^3$  over plane curves. Similar examples can be produced by composing an extrinsic product of isometric immersions into a sphere (see Section 6 for the precise definition of this concept) with a stereographic projection of that sphere onto Euclidean space.

Further examples of isothermic submanifolds arise by means of the diffeomorphism  $\Theta: \mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N$ ,  $1 \leq m \leq N-1$ , onto the complement of an  $(m-1)$ -dimensional affine subspace (a point if  $m=1$ ) in  $\mathbb{R}^N$  (see Lemma 23 below). Here  $\mathbb{H}^m(-c) = \{X \in \mathbb{L}^{m+1}: \langle X, X \rangle = -1/c\}$  and  $\mathbb{S}^{N-m}(c) = \{X \in \mathbb{R}^{N-m+1}: \langle X, X \rangle = 1/c\}$ , where  $\langle \cdot, \cdot \rangle$  stands for the Lorentz inner product in Minkowski space  $\mathbb{L}^{m+1}$  in the first case, and the standard Euclidean inner product in the latter. Thus, if  $m \geq 2$  (resp.,  $N-m \geq 2$ ) then  $\mathbb{H}^m(-c)$  (resp.,  $\mathbb{S}^{N-m}(c)$ ) is the standard model of hyperbolic space (resp., the sphere) with constant sectional curvature  $-c$  (resp.,  $c$ ) as an umbilical hypersurface of  $\mathbb{L}^{m+1}$  (resp.,  $\mathbb{R}^{N-m+1}$ ). Given an isometric immersion  $f: M_1 \rightarrow \mathbb{H}^m(-c)$  and an extrinsic product  $g: M_2 \times \dots \times M_k \rightarrow \mathbb{S}^{N-m}(c)$  of isometric immersions, then  $I \circ \Theta \circ (f \times g)$  defines a  $k$ -isothermic submanifold for any inversion  $I$  in  $\mathbb{R}^N$ . In two dimensions, these examples reduce to conformal parameterizations of the images by an inversion of cones and surfaces of revolution.

The abundance of isothermic surfaces in  $\mathbb{R}^3$  gives a hint that their higher dimensional analogues should be equally as plentiful. The hint turns out to be surprisingly misleading: we are able to show that the examples just described actually comprise all isothermic submanifolds of dimension  $n \geq 3$ . We leave the precise statement for Section 7 (see Theorem 25). From the point of view of the theory of integrable systems, the result is rather disappointing: there is no such theory in connection with isothermic submanifolds of dimension  $n \geq 3$ . The discrepancy between the richness of the two-dimensional theory and the shortage of examples in higher dimensions should be compared to that between the classes of conformal maps between open subsets of Euclidean space in two and higher dimensions. In fact, our result contains as a special case Liouville's theorem on the classification of such conformal maps in dimension  $n \geq 3$ .

Having our definition of isothermic submanifolds not led to as rich a theory as in the two-dimensional case, one might think that it should be better to choose a definition based on an extension of one of the properties that characterize isothermic surfaces in Euclidean three-space. For instance, one might look for those submanifolds that admit a nontrivial conformal sphere congruence, as suggested by Burstall, or search for the solutions of some natural extension of Christoffel's problem. We carry this out and give a complete classification of both classes of submanifolds. Unfortunately, they turn out to be proper (and different) subclasses of the class of isothermic submanifolds as previously defined.

In mentioning the third characterization of isothermic surfaces in terms of infinitesimal deformations, our purpose was to indicate that some of such characterizations seem to be exclusively of a two-dimensional nature, and are not likely to admit any extension to higher dimensions.

Before we draw this already overlong introduction to a close, we provide the reader a brief guide to the contents of each section. We also describe some applications of our main result and indicate some of the tools used in its proof.

We start Section 2 by reviewing from [RS] the notions of nets and net morphisms between netted manifolds, a suitable setting for dealing with decomposition problems for manifolds and for maps between them. Then we recall from [MRS] some basic facts on twisted and warped products and state the local decomposition theorem of a Riemannian manifold as a twisted product, which extends the local versions of de Rham and Hiepko's theorems. We close the section by summarizing some results of [To<sub>1</sub>], where the intrinsic notion of an isothermic net was extended for arbitrary Riemannian manifolds, leading to a conformal version of the local de Rham theorem.

In Section 3 we first give a brief description of the model of Euclidean space as an umbilical hypersurface of the light cone in Minkowski space, an extremely convenient framework to deal with Moebius geometric notions. This is applied to study families of spheres and affine subspaces in Euclidean space, as a preparation for the characterization in the following section of certain mutually orthogonal families of spheres and affine subspaces in Euclidean space (Lemma 11), the principal tool in the proof of our main result. In Section 5 we derive two facts (Propositions 15 and 16) on extrinsic spheres of Euclidean submanifolds that also play a key rôle in that proof. As a first application, we provide in Section 6 a new geometric proof (for Euclidean space) of Nölker's decomposition theorem (Theorem 19) of isometric immersions of warped products into space forms, after reviewing from [No] the notions of warped product representations of space forms and warped products of isometric immersions.

Our main results are contained in Section 7. We obtain a local classification of isothermic Euclidean submanifolds of dimension  $n \geq 3$ , or equivalently, we describe all conformal immersions of a Riemannian product of dimension  $n \geq 3$  into Euclidean space whose second fundamental forms are adapted to the product net of the manifold. We follow with several applications. First, we specialize to surfaces in Euclidean space and generalize to arbitrary codimension a classical result of Bonnet on surfaces whose curvature lines have constant geodesic curvature (Corollary 27). Then, we classify all conformal representations of Euclidean space of dimension  $n \geq 3$  as a Riemannian product, that is, we determine all local diffeomorphisms of a Riemannian product of dimension  $n \geq 3$  onto an open subset of Euclidean space (Corollary 28). Besides containing Liouville's theorem as a special case, this also provides a geometric proof of the known classification of conformally flat Riemannian products. It yields as well a classification of all conformal local diffeomorphisms of a warped product of dimension  $n \geq 3$  with two factors onto an open subset of Euclidean space, which in the two dimensional case recovers a classical theorem on cartography due to Lagrange. We conclude the section with a decomposition theorem for isometric immersions of a twisted product of two Riemannian manifolds into Euclidean space (Corollary 33), which contains as a special case the main result of [DFT]. The latter classifies the Euclidean submanifolds

that carry a Dupin principal normal vector field with umbilical conullity (see the end of Section 7 for the meaning of these terms), which include in particular the cyclides of Dupin of arbitrary dimension.

In the last section, after stating some basic facts on Codazzi tensors we formulate an extension of Christoffel problem for Euclidean submanifolds of arbitrary dimension and codimension, and determine all of its solutions. We conclude the paper with a classification of all Euclidean submanifolds that admit nontrivial conformal sphere congruences.

In a forthcoming paper [To<sub>2</sub>] we consider the more general problem of describing all conformal immersions into Euclidean space of a *warped* product of dimension  $n \geq 3$  with arbitrarily many factors.

## 2 Preliminaries

### 2.1 Nets and net morphisms

A suitable setting for treating decomposition results for manifolds and for maps between them was developed in [RS] by introducing the category of *netted manifolds*.

A *net* on a connected  $C^\infty$ -manifold  $M$  is a splitting  $TM = \oplus_{i=1}^k E_i$  into a family of integrable subbundles. If  $M$  is a Riemannian manifold and the subbundles are mutually orthogonal then the net is said to be an *orthogonal net*. The canonical net on a product manifold  $M = \prod_{i=1}^k M_i$  is called the *product net*. A  $C^\infty$ -map  $\psi: M \rightarrow N$  between two *netted manifolds*  $(M, \mathcal{E})$ ,  $(N, \mathcal{F})$ , that is, manifolds  $M, N$  equipped with nets  $\mathcal{E} = (E_i)_{i=1, \dots, k}$  and  $\mathcal{F} = (F_i)_{i=1, \dots, k}$ , respectively, is called a *net morphism* if  $\psi_* E_i(p) \subset F_i(\psi(p))$  for all  $p \in M$ ,  $1 \leq i \leq k$ , or equivalently, if for any  $p \in M$  the restriction  $\psi|_{L_i^\mathcal{E}(p)}$  of  $\psi$  to the leaf  $L_i^\mathcal{E}(p)$  of  $E_i$  through  $p$  is a  $C^\infty$ -map into the leaf  $L_i^\mathcal{F}(\psi(p))$  of  $F_i$  through  $\psi(p)$ . The net morphism  $\psi$  is said to be a *net isomorphism* if in addition it is a diffeomorphism and  $\psi^{-1}$  is also a net morphism. A net  $\mathcal{E}$  on  $M$  is said to be *locally decomposable* if for every point  $p \in M$  there exist a neighborhood  $U$  of  $p$  in  $M$  and a net isomorphism  $\psi$  from  $(U, \mathcal{E}|_U)$  onto a product manifold  $\prod_{i=1}^k M_i$ . The map  $\psi^{-1}: \prod_{i=1}^k M_i \rightarrow U$  is called a *product representation* of  $(U, \mathcal{E}|_U)$ . The following is a basic criterion for local decomposability of a net (cf. Theorem 1 of [RS]).

**Proposition 1.** [RS] *A  $k$ -net  $\mathcal{E} = (E_i)_{i=1, \dots, k}$  on a  $C^\infty$ -manifold  $M$  is locally decomposable if and only if  $E_i^\perp := \oplus_{j \neq i} E_j$  is integrable for every  $1 \leq i \leq k$ .*

The task of verifying whether a  $C^\infty$ -map  $f: M \rightarrow N$  between product manifolds  $M = \prod_{i=1}^k M_i$  and  $N = \prod_{i=1}^k N_i$  decomposes as a product of  $C^\infty$ -maps between the corresponding factors is simplified by the following key observation (cf. [RS], Lemma 1). Hereafter, we denote by  $\pi_i: M \rightarrow M_i$  the canonical projection of  $M$  onto  $M_i$ . Moreover, for a fixed point  $\bar{p} \in M$  we denote by  $\tau_i^{\bar{p}}: M_i \rightarrow M$  the canonical embedding of  $M_i$  into  $M$  given by  $\tau_i^{\bar{p}}(p_i) = (\bar{p}_0, \dots, p_i, \dots, \bar{p}_k)$ .

**Lemma 2 [RS]** *Let  $f: M \rightarrow N$  be a  $C^\infty$ -map between product manifolds  $M = \Pi_{i=1}^k M_i$  and  $N = \Pi_{i=1}^k N_i$ . For a fixed point  $\bar{p} \in M$  define  $f_i = \pi_i \circ f \circ \tau_i^{\bar{p}}: M_i \rightarrow N_i$ . Then  $f = f_1 \times \cdots \times f_k$ , that is,  $f(p_1, \dots, p_k) = (f_1(p_1), \dots, f_k(p_k))$  for every  $p = (p_1, \dots, p_k) \in M$ , if and only if  $f$  is a net morphism with respect to the product nets of  $M$  and  $N$ .*

## 2.2 Twisted and warped products

We now recall from [MRS] some basic definitions and results on warped and twisted products. If  $M = \Pi_{i=0}^k M_i$  is the product of  $C^\infty$  manifolds  $M_0, \dots, M_k$ , then a metric  $\langle \cdot, \cdot \rangle$  on  $M$  is called a *twisted product metric* if there exist Riemannian metrics  $\langle \cdot, \cdot \rangle_i$  on  $M_i$ ,  $0 \leq i \leq k$ , and a  $C^\infty$  twist-function  $\rho = (\rho_0, \dots, \rho_k): M \rightarrow \mathbb{R}_+^{k+1}$  such that

$$\langle \cdot, \cdot \rangle = \sum_{i=0}^k \rho_i^2 \pi_i^* \langle \cdot, \cdot \rangle_i.$$

Then  $(M, \langle \cdot, \cdot \rangle)$  is said to be a *twisted product* and is denoted by  ${}^\rho \Pi_{i=0}^k (M_i, \langle \cdot, \cdot \rangle_i)$ . If there exists  $\bar{p} \in M$  such that  $\rho_i|_{L_i^\xi(\bar{p})} = 1$  for every  $0 \leq i \leq k$ , then the representation of  $M$  as a twisted product is said to be *normalized* with respect to  $\bar{p}$ . When  $\rho_1, \dots, \rho_k$  are independent of  $M_1, \dots, M_k$ , that is, there exist  $\tilde{\rho}_i \in C^\infty(M_0)$  such that  $\rho_i = \tilde{\rho}_i \circ \pi_0$  for all  $1 \leq i \leq k$ , and, in addition,  $\rho_0$  is identically 1, then  $\langle \cdot, \cdot \rangle$  is called a *warped product metric* and  $(M, \langle \cdot, \cdot \rangle) := (M_0, \langle \cdot, \cdot \rangle_0) \times_{\tilde{\rho}} \Pi_{i=1}^k (M_i, \langle \cdot, \cdot \rangle_i)$  a *warped product* with *warping function*  $\tilde{\rho} = (\tilde{\rho}_1, \dots, \tilde{\rho}_k)$ . If  $\rho_i$  is identically 1 for all  $0 \leq i \leq k$ , the metric  $\langle \cdot, \cdot \rangle$  is the usual *Riemannian product metric*, in which case  $(M, \langle \cdot, \cdot \rangle)$  is called a *Riemannian product*.

An orthogonal net  $\mathcal{E} = (E_i)_{i=0, \dots, k}$  on a Riemannian manifold  $M$  is called a *TP-net* if  $E_i$  is umbilical and  $E_i^\perp$  is integrable for every  $0 \leq i \leq k$ . Recall that a subbundle  $E$  of  $TM$  is *umbilical* if there exists a vector field  $\eta$  in  $E^\perp$  such that

$$(\nabla_X Y)_{E^\perp} = \langle X, Y \rangle \eta \text{ for all } X, Y \in \Gamma(E).$$

Here and in the sequel, the space of smooth local sections of a vector bundle  $E$  over  $M$  is denoted by  $\Gamma(E)$ , whereas writing a vector field with a vector subbundle as a subscript indicates taking the section of that vector subbundle obtained by orthogonally projecting the vector field pointwise onto the corresponding fiber of the subbundle. The vector field  $\eta$  is called the *mean curvature normal* of  $E$ . If, in addition,  $(\nabla_X \eta)_{E^\perp} = 0$  for all  $X \in \Gamma(E)$ , then  $E$  is said to be *spherical*. If  $E$  is umbilical and its mean curvature normal vanishes identically, then it is called *totally geodesic* (or *auto-parallel*). An umbilical distribution is automatically integrable, and the leaves are umbilical submanifolds of  $M$ . When  $E$  is totally geodesic or spherical, its leaves are totally geodesic or spherical submanifolds, respectively. By a *spherical submanifold*, or an *extrinsic sphere*, we mean an umbilical submanifold whose mean curvature vector is parallel with respect to the normal connection.

An orthogonal net  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  is called a *WP-net* if  $E_i$  is spherical and  $E_i^\perp$  is totally geodesic for every  $1 \leq i \leq k$ . If  $E_i$  and  $E_i^\perp$  are totally geodesic for every  $1 \leq i \leq k$ , then  $\mathcal{E}$  is called an *RP-net*. In a *WP-net* the subbundle  $E_0$  is automatically totally geodesic and  $E_0^\perp$  is integrable (see Proposition 3 in [MRS]); in particular, every *WP-net* is also a *TP-net*. The terminologies *TP-net* and *WP-net* are justified by the following result (see [MRS], Proposition 4).

**Proposition 3 [MRS]** *On a connected product manifold  $M =: \Pi_{i=0}^k M_i$  the product net  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  is a *TP-net* (resp., *WP-net*) with respect to a Riemannian metric  $\langle \cdot, \cdot \rangle$  on  $M$  if and only if  $\langle \cdot, \cdot \rangle$  is a twisted product (resp., warped product) metric on  $M$ . Moreover, if  $\rho = (\rho_0, \dots, \rho_k): M \rightarrow \mathbb{R}_+^{k+1}$  is the twist function and  $U_i = -\nabla(\log \circ \rho_i)$ ,  $0 \leq i \leq k$ , where the gradient is calculated with respect to  $\langle \cdot, \cdot \rangle$ , then the mean curvature normal of  $E_i$  is  $\eta_i = (U_i)_{E_i^\perp}$  for all  $0 \leq i \leq k$ .*

By means of Propositions 1 and 3 the following extension of de Rham and Hiepko local decomposition theorems was derived in [MRS].

**Theorem 4 [MRS]** *Let  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  be a *TP-net* (resp., *WP-net*) on a Riemannian manifold  $M$ . Then, for every point  $p \in M$  there exists a local product representation  $\psi: \Pi_{i=0}^k M_i \rightarrow U$  of  $\mathcal{E}$  with  $p \in U \subset M$ , which is an isometry with respect to a twisted product (resp., warped product) metric on  $\Pi_{i=0}^k M_i$ .*

## 2.3 Conformal product nets

An orthogonal net  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  on a Riemannian manifold is a *conformal product net*, or a *CP-net* for short, if both  $E_i$  and  $E_i^\perp$  are umbilical for every  $0 \leq i \leq k$  and

$$\langle \nabla_{X_{\perp i}} \eta_i, X_i \rangle = \langle \nabla_{X_i} H_i, X_{\perp i} \rangle \quad \text{for all } X_i \in \Gamma(E_i), \quad X_{\perp i} \in \Gamma(E_i^\perp), \quad 1 \leq i \leq k, \quad (1)$$

where  $H_i$  and  $\eta_i$  are the mean curvature normals of  $E_i$  and  $E_i^\perp$ , respectively. It is easily seen that condition (1) then holds automatically also for  $i = 0$ . Moreover, we have the following other immediate consequences of the definition.

**Proposition 5** (i) *If  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  is an orthogonal net such that  $E_i$  and  $E_i^\perp$  are spherical for every  $0 \leq i \leq k$  then it is a *CP-net*;*

(ii) *If  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  is a *CP-net*, then for each  $0 \leq i \leq k$  one of  $E_i$  or  $E_i^\perp$  being spherical implies the same for the other.*

The terminology is justified by the next result.

**Proposition 6** *On a connected product manifold  $M = \Pi_{i=0}^k M_i$  the product net  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  is a *CP-net* with respect to a Riemannian metric  $\langle \cdot, \cdot \rangle$  on  $M$  if and only if  $\langle \cdot, \cdot \rangle$  is conformal to a Riemannian product metric on  $M$ .*



The following conformal version of the local de Rham Theorem was obtained in [To<sub>1</sub>] by putting together Propositions 1 and 6. It shows that conformal product nets are natural generalizations of isothermic nets on surfaces.

**Theorem 7** *Let  $\mathcal{E} = (E_i)_{i=0,\dots,k}$  be a CP-net on a Riemannian manifold  $M$ . Then for every point  $p \in M$  there exists a local product representation  $\psi: \Pi_{i=0}^k M_i \rightarrow U$  of  $\mathcal{E}$  with  $p \in U \subset M$ , which is a conformal diffeomorphism with respect to a Riemannian product metric on  $\Pi_{i=0}^k M_i$ .*

### 3 Moebius geometry in the light cone

We start this section with a brief description of the model of Euclidean space as an umbilical hypersurface of the light cone of Minkowski space. Then we use the model to study families of affine subspaces and spheres in Euclidean space.

#### 3.1 Euclidean space as a hypersurface of the light cone

Let  $\mathbb{L}^{N+2}$  be the  $(N+2)$ -dimensional Minkowski space endowed with a Lorentz scalar product of signature  $(+, \dots, +, -)$ , and let

$$\mathbb{V}^{N+1} = \{p \in \mathbb{L}^{N+2} : \langle p, p \rangle = 0\}$$

denote the light cone in  $\mathbb{L}^{N+2}$ . For any  $w \in \mathbb{V}^{N+1}$ , the intersection

$$\mathbb{E}^N = \{p \in \mathbb{V}^{N+1} : \langle p, w \rangle = 1\}$$

of the light cone with the affine hyperplane  $\langle p, w \rangle = 1$  is a model of the  $N$ -dimensional Euclidean space. Namely, choose  $p_0 \in \mathbb{E}^N$  and orthogonally decompose

$$\mathbb{L}^{N+2} = \mathbb{R}^N \oplus \text{span}\{p_0, w\}.$$

Then we have an isometry  $\Psi: \mathbb{R}^N \rightarrow \mathbb{E}^N$  given by

$$x \in \mathbb{R}^N \rightarrow x + p_0 - (1/2)\|x\|^2 w.$$

This model provides a convenient setting for dealing with Moebius geometric notions. The main reason is that conformal transformations of  $\mathbb{R}^N$  are linearized as follows. Given a conformal immersion  $G: M^n \rightarrow \mathbb{V}^{N+1}$  with conformal factor  $\varphi \in C^\infty(M)$  and  $\mu \in C^\infty(M^n)$ , the map  $G_\mu: M^n \rightarrow \mathbb{V}^{N+1}$ ,  $G_\mu(p) = \mu(p)G(p)$  for all  $p \in M^n$ , is conformal with conformal factor  $\mu\varphi$ . Thus, any conformal immersion  $G: M^n \rightarrow \mathbb{R}^N$  with conformal factor  $\varphi \in C^\infty(M^n)$  gives rise to an isometric immersion  $(\Psi \circ G)_{\varphi^{-1}}: M^n \rightarrow \mathbb{V}^{N+1}$ .

Conversely, given an isometric immersion  $F: M^n \rightarrow \mathbb{V}^{N+1}$  with  $F(M^n) \subset \mathbb{V}^{N+1} \setminus \mathbb{R}_w$ , where  $\mathbb{R}_w = \{tw : t > 0\}$ , let  $\mathcal{C}(F): M^n \rightarrow \mathbb{R}^N$  be defined by

$$\Psi \circ \mathcal{C}(F) = \Pi \circ F, \quad (2)$$

where  $\Pi: \mathbb{V}^{N+1} \setminus \mathbb{R}_w \rightarrow \mathbb{E}^N$  is the projection onto  $\mathbb{E}^N$  given by  $\Pi(x) = x/\langle x, w \rangle$ . Since  $\Pi$  is conformal with conformal factor  $\varphi_\Pi(x) = \langle x, w \rangle^{-1}$ , it follows that  $\mathcal{C}(F)$  is also conformal with conformal factor  $\varphi_\Pi \circ F = \langle F, w \rangle^{-1}$ . In particular, any  $T \in \mathbb{O}_1(N+2)$  gives rise to a conformal (Moebius) transformation  $\mathcal{C}(T)$  of  $\mathbb{R}^N$  and, conversely, any Moebius transformation of  $\mathbb{R}^N$  is given by means of some  $T \in \mathbb{O}_1(N+2)$  in the way just described.

Another reason is that hyperspheres in  $\mathbb{R}^N$  can be nicely described in  $\mathbb{E}^N$  as follows (cf. [HTD]). Given a hypersphere  $S$  in  $\mathbb{E}^N$  with (constant) mean curvature  $h$  with respect to a unit normal vector field  $n$  along  $S$ , then  $v = n_p + hp \in \mathbb{L}^{N+2}$  is a constant space-like unit vector such that  $\langle v, p \rangle = 0$  for all  $p \in S$ . Thus, the (nonoriented) sphere  $S$  is given in the model by  $S = \mathbb{E}^N \cap \{v\}^\perp$ . Since  $h = \langle v, w \rangle$ , then  $v$  represents a hyperplane in  $\mathbb{E}^N$  if and only if  $\langle v, w \rangle = 0$ .

The intersection angle of two (oriented) hyperspheres has also a simple description in this model: given hyperspheres  $S_i = \mathbb{E}^N \cap \{v_i\}^\perp$  with unit normal vectors  $n_p^i$ ,  $1 \leq i \leq 2$ , at a common point  $p$ , their intersection angle at  $p$  is given by  $\langle n_p^1, n_p^2 \rangle = \langle v_1, v_2 \rangle$ . Thus  $S_1$  and  $S_2$  intersect orthogonally if and only if  $\langle v_1, v_2 \rangle = 0$ .

A hypersphere  $S = \mathbb{E}^N \cap \{v\}^\perp$  has center at  $p_0 \in \mathbb{E}^N$  and mean curvature  $h \neq 0$  if and only if  $v = hp_0 + (2h)^{-1}w$ . In fact, this follows from  $\langle v, w \rangle = h$ ,  $\langle v, v \rangle = 1$  and  $\text{span}\{p_0, w\}^\perp \subset v^\perp$ , the latter being due to the fact that any hyperplane through  $p_0$  is orthogonal to  $S$ .

Since any  $m$ -dimensional sphere  $S$  in  $\mathbb{E}^N$  is the intersection of  $(N-m)$  mutually orthogonal hyperspheres  $S_i = \mathbb{E}^N \cap \{v_i\}^\perp$ ,  $\langle v_i, v_j \rangle = \delta_{ij}$  for  $1 \leq i, j \leq N-m$ , it follows that  $S = \mathbb{E}^N \cap V^\perp$ , where  $V$  is the space-like  $(N-m)$ -dimensional subspace of  $\mathbb{L}^{N+2}$  spanned by  $v_1, \dots, v_{N-m}$ . Therefore, the space of (non-oriented)  $m$ -dimensional spheres in  $\mathbb{E}^N$  is naturally identified with the Grassmann manifold  $G_+(N-m, m+2)$  of space-like  $(N-m)$ -dimensional subspaces of  $\mathbb{L}^{N+2}$ . The  $m$ -dimensional affine subspaces in  $\mathbb{E}^N$  are given by those space-like  $(N-m)$ -dimensional subspaces of  $\mathbb{L}^{N+2}$  orthogonal to  $w$ .

We say that two  $m_i$ -dimensional spheres or affine subspaces  $S_i$  in  $\mathbb{R}^N$ ,  $1 \leq i \leq 2$ , with  $m_1 + m_2 \geq N$ , intersect orthogonally if their normal spaces  $H_1$  and  $H_2$  in  $\mathbb{R}^N$ , respectively, at one (and hence at any) point in  $S_1 \cap S_2$  satisfy  $H_1 \subset H_2^\perp$ . If  $S_i$  is given in the model  $\mathbb{E}^N$  by  $S_i = \mathbb{E}^N \cap V_i^\perp$ ,  $1 \leq i \leq 2$ , then this is equivalent to  $V_1 \subset V_2^\perp$ .

The relative position of spheres and affine subspaces can be easily recovered from the corresponding space like subspaces. For instance, let  $H_i$ ,  $1 \leq i \leq 2$ , be  $m_i$ -dimensional affine subspaces in  $\mathbb{R}^N$  with, say,  $m_1 \geq m_2$ . Let  $H_i$  be given in the model  $\mathbb{E}^N$  by  $H_i = \mathbb{E}^N \cap V_i^\perp$ , where  $V_i$  is a  $(N-m_i)$ -dimensional space-like subspace with  $w \in V_i^\perp$ ,  $1 \leq i \leq 2$ . Then  $H_1$  is parallel to  $H_2$ , meaning that  $H_1$  contains a parallel translate of  $H_2$ , if and only if  $V_1 \subset V_2 \oplus \text{span}\{w\}$ . In fact,  $H_1$  is parallel to  $H_2$  if and only if

every affine subspace orthogonal to  $H_2$  is also orthogonal to  $H_1$ , which is equivalent to  $(V_2 \oplus \text{span}\{w\})^\perp \subset V_1^\perp$ . Arguing in a similar way, it follows that an  $m_1$ -dimensional sphere  $S_1 = \mathbb{E}^N \cap V_1^\perp$  is tangent at  $p \in \mathbb{E}^N$  to an  $m_2$ -dimensional sphere or affine subspace  $S_2 = \mathbb{E}^N \cap V_2^\perp$ ,  $m_1 \geq m_2$ , if and only if  $V_1 \subset V_2 \oplus \text{span}\{p\}$ .

### 3.2 Families of spheres and affine subspaces in Euclidean space

The geometric nature of a family  $\mathcal{F} = (H^\alpha)_{\alpha \in \Lambda}$  of at least two affine subspaces in  $\mathbb{R}^N$  of a fixed dimension  $\ell$  can be captured from the type of a subspace  $V_{\mathcal{F}}$  of  $\mathbb{L}^{N+2}$  naturally associated to it. Namely, for each  $\alpha \in \Lambda$  let  $V^\alpha$  be the space-like subspace of  $\mathbb{L}^{N+2}$  such that  $H^\alpha = \mathbb{E}^N \cap V^{\alpha\perp}$ , and let  $V_{\mathcal{F}}$  be the subspace spanned by all  $V^\alpha$ ,  $\alpha \in \Lambda$ . Since  $w \in V_{\mathcal{F}}^\perp$ , the subspace  $V_{\mathcal{F}}$  is either degenerate or space-like. We say that  $\mathcal{F}$  is a *light-like* (respectively, *space-like*) family if  $V_{\mathcal{F}}$  is degenerate (respectively, space-like). As  $\mathcal{F}$  has at least two elements and each  $V^\alpha$  has dimension  $N - \ell$ , the dimension  $m + 1$  of  $V_{\mathcal{F}}$  is at least  $N - \ell + 1$ .

**Proposition 8** (i) *If  $\mathcal{F}$  is space-like, then  $\cap_{\alpha \in \Lambda} H^\alpha = \mathbb{E}^N \cap V_{\mathcal{F}}^\perp := \mathcal{H}_{\mathcal{F}}$  is either an affine subspace of dimension  $N - m - 1$  if  $m \leq N$  or a single point  $\bar{q} \in \mathbb{R}^N$  otherwise. Conversely, if every  $H^\alpha \in \mathcal{F}$  contains a point or an affine subspace  $\tilde{\mathcal{H}} = \mathbb{E}^N \cap \tilde{V}^\perp$  then  $\mathcal{F}$  is space-like and  $\tilde{\mathcal{H}} \subset \mathcal{H}_{\mathcal{F}}$ .*

(ii) *If  $\mathcal{F}$  is light-like, write  $V_{\mathcal{F}} = U \oplus \text{span}\{w\}$ , where  $U$  is a (non-unique) space-like subspace of dimension  $m$ , and let  $\mathbb{R}_{\mathcal{F}}^{N-m}$  be the  $(N-m)$ -dimensional subspace of  $\mathbb{R}^N$  that is parallel to  $\mathcal{H} = \mathbb{E}^N \cap U^\perp$  (if  $m$  assumes its maximal possible value  $N$  then  $\mathbb{R}_{\mathcal{F}}^{N-m}$  is the trivial subspace and  $\mathcal{H}$  reduces to a single point). Then every  $H^\alpha \in \mathcal{F}$  is parallel to  $\mathbb{R}_{\mathcal{F}}^{N-m}$ , which is the maximal subspace with this property.*

*Proof:* (i) Since  $V^\alpha \subset V_{\mathcal{F}}$  for every  $\alpha \in \Lambda$ , then  $\mathcal{H}_{\mathcal{F}} = \mathbb{E}^N \cap V_{\mathcal{F}}^\perp \subset \cap_{\alpha \in \Lambda} H^\alpha$ . On the other hand, if every  $H^\alpha \in \mathcal{F}$  contains a point or an affine subspace  $\tilde{\mathcal{H}} = \mathbb{E}^N \cap \tilde{V}^\perp$ , where  $\tilde{V}$  is a space-like subspace of  $\mathbb{L}^{N+2}$  with  $w \in \tilde{V}^\perp$ , then  $V^\alpha \subset \tilde{V}$  for every  $\alpha \in \Lambda$ . Thus  $V_{\mathcal{F}} \subset \tilde{V}$ , and hence  $V_{\mathcal{F}}$  is space-like and  $\tilde{\mathcal{H}} \subset \mathcal{H}_{\mathcal{F}}$ .

(ii) Given  $H^\alpha = \mathbb{E}^N \cap V^{\alpha\perp} \in \mathcal{F}$ , we have that  $V^\alpha \subset V_{\mathcal{F}} = U \oplus \text{span}\{w\}$ . Thus  $H^\alpha$  is parallel to  $\mathbb{E}^N \cap U^\perp = \mathcal{H}$ , and hence to  $\mathbb{R}_{\mathcal{F}}^{N-m}$ . Moreover, if  $\mathbb{R}^s = \mathbb{E}^N \cap \tilde{U}^\perp$  is a subspace of  $\mathbb{R}^N$  such that every  $H^\alpha \in \mathcal{F}$  is parallel to  $\mathbb{R}^s$ , then  $V^\alpha \subset \tilde{U} \oplus \text{span}\{w\}$  for every  $\alpha \in \Lambda$ . Therefore  $V_{\mathcal{F}} \subset \tilde{U} \oplus \text{span}\{w\}$ , which implies that  $\mathbb{R}^s \subset \mathbb{R}_{\mathcal{F}}^{N-m}$ . ■

If  $\mathcal{F}$  is a space-like (resp., light-like) family of affine subspaces of dimension  $\ell$  in  $\mathbb{R}^N$ , then we call the affine subspace  $\mathcal{H}_{\mathcal{F}}$  (resp., the subspace  $\mathbb{R}_{\mathcal{F}}^{N-m}$ ) given by Proposition 8 the *axis* of  $\mathcal{F}$ . We obtain from Proposition 8 the following characterization of the spherical hull of an immersion  $f: M^n \rightarrow \mathbb{R}^N$  in terms of the family of affine normal spaces of  $f$ . Recall that the *spherical hull* of an immersion  $f: M^n \rightarrow \mathbb{Q}_c^N$  into a simply connected complete space form of constant sectional curvature  $c$  is the complete spherical submanifold of minimal dimension of  $\mathbb{Q}_c^N$  that contains  $f(M)$ .

**Proposition 9** *Let  $f: M^n \rightarrow \mathbb{R}^N$  be an immersion, let  $\mathcal{F} = (H(p))_{p \in M}$  be the family of affine normal spaces of  $f$ , and let  $m+1 \geq n+1$  be the dimension of  $V_{\mathcal{F}}$ . Then the following hold:*

- (i) *If  $\mathcal{F}$  is light-like and  $\mathbb{R}^N = \mathbb{R}^{N-m} \oplus \mathbb{R}^m$  is the orthogonal decomposition of  $\mathbb{R}^N$  with  $\mathbb{R}^{N-m} = \mathbb{R}_{\mathcal{F}}^{N-m}$ , then the spherical hull of  $f(M^n)$  is the affine subspace  $f(q) + \mathbb{R}^m$  for any  $q \in M^n$ .*
- (ii) *If  $\mathcal{F}$  is space-like, let  $\mathbb{R}^N = \mathbb{R}^{N-m-1} \oplus \mathbb{R}^{m+1}$  be the orthogonal decomposition of  $\mathbb{R}^N$  such that  $\mathbb{R}^{N-m-1}$  is the  $(N-m-1)$ -dimensional subspace of  $\mathbb{R}^N$  that is parallel to  $\mathcal{H}_{\mathcal{F}}$ . Then for any  $q \in M^n$  the spherical hull of  $f(M^n)$  is the  $m$ -dimensional sphere in  $f(q) + \mathbb{R}^{m+1}$  centered at the intersection of  $\mathcal{H}_{\mathcal{F}}$  with  $f(q) + \mathbb{R}^{m+1}$ .*

*Proof:* (i) Since the subspace  $\mathbb{R}^{N-m}$  is contained in the normal space of  $f$  at  $p$  for every  $p \in M^n$  (regarded as a subspace of  $\mathbb{R}^N$ ), it follows that  $f(M^n)$  is contained in  $f(q) + \mathbb{R}^m$  for any  $q \in M^n$ . If  $f(M^n)$  were contained in a lower dimensional affine subspace  $f(q) + \mathbb{R}^\ell \subset f(q) + \mathbb{R}^m$ ,  $\ell < m$ , then  $\mathbb{R}^{N-\ell}$  would be a subspace of  $\mathbb{R}^N$  of dimension  $N-\ell > N-m$  parallel to every  $H(p)$ ,  $p \in M$ , contradicting the maximality of  $\mathbb{R}_{\mathcal{F}}^{N-m}$  with respect to this property. This shows that  $f(q) + \mathbb{R}^m$  is the *affine* hull of  $f(M^n)$  for any  $q \in M^n$ , that is, the affine subspace of minimum dimension that contains  $f(M^n)$ . It remains to be shown that  $f(M^n)$  can not be contained in a sphere. In fact, otherwise every affine normal space  $H(p) \in \mathcal{F}$  would contain the center of that sphere, and hence  $\mathcal{F}$  would be a space-like family by Proposition 8-(i).

(ii) Arguing as in the preceding case, we obtain that  $f(q) + \mathbb{R}^{m+1}$  is the affine hull of  $f(M^n)$  for any  $q \in M^n$ . Thus, it suffices to show that  $f(M^n)$  is contained in the hypersphere centered at the intersection  $\bar{q}$  of  $\mathcal{H}_{\mathcal{F}}$  with  $f(q) + \mathbb{R}^{m+1}$ . This follows from the fact that the position vector  $f(q) - \bar{q}$  belongs to  $H(q)$  for any  $q \in M^n$ , and is therefore a normal vector field along  $f(M^n)$ . ■

Let  $T \in \mathbb{O}_1(N+2)$  be the reflection

$$T(p) = p - 2\langle p, v \rangle v \quad (3)$$

with respect to the hyperplane in  $\mathbb{L}^{N+2}$  orthogonal to the unit space-like vector  $v$ , with  $\langle v, w \rangle \neq 0$ . Then the conformal transformation  $\mathcal{C}(T)$  of  $\mathbb{R}^N$  given by (2) is the inversion with respect to the hypersphere  $S = \mathbb{E}^N \cap \{v\}^\perp$ . This is used next to extend to families of spheres in  $\mathbb{R}^N$  the description of families of affine subspaces in Proposition 8.

Let  $\mathcal{F} = (S^\alpha)_{\alpha \in \Lambda}$  be a family of at least two spheres or affine subspaces of dimension  $\ell$  in  $\mathbb{R}^N$ . As before, for each  $\alpha \in \Lambda$  let  $V^\alpha$  be the space-like subspace of  $\mathbb{L}^{N+2}$  such that  $S^\alpha = \mathbb{E}^N \cap V^{\alpha\perp}$  and let  $V_{\mathcal{F}}$  be the subspace spanned by all  $V^\alpha$ ,  $\alpha \in \Lambda$ . The subspace  $V_{\mathcal{F}}$  can now be either space-like, degenerate or time-like. We say accordingly that  $\mathcal{F}$  is a *space-like*, *light-like* or *time-like family*. Basic examples of time-like families arise as follows. Let  $\mathcal{H} = \mathbb{E}^N \cap V^\perp$  be an affine subspace of  $\mathbb{R}^N$  of dimension  $m$  (a point  $\bar{q}$  if

$m = 0$ ), where  $V$  is a space-like subspace of  $\mathbb{L}^{N+2}$  of dimension  $N - m$  with  $w \in V^\perp$ . If  $m > 0$ , let  $\mathcal{F}$  be the family of all spheres or affine subspaces of a fixed dimension  $\ell$ ,  $\ell + m \geq N$ , that are orthogonal to  $\mathcal{H}$ . If  $m = 0$ , let  $\mathcal{F}$  be the family of hyperspheres centered at  $\bar{q}$ . Then  $V_{\mathcal{F}} = V^\perp$  is a time-like subspace, and hence  $\mathcal{F}$  is a time-like family. We say that it is a *standard time-like family* of spheres or affine subspaces in  $\mathbb{R}^N$  and call  $\mathcal{H} = \mathcal{H}_{\mathcal{F}}$  the *axis* of  $\mathcal{F}$ .

**Proposition 10** *Let  $\mathcal{F} = (S^\alpha)_{\alpha \in \Lambda}$  be a family of at least two spheres or affine subspaces of dimension  $\ell$  in  $\mathbb{R}^N$ , and let  $m + 1 \geq N - \ell + 1$  be the dimension of  $V_{\mathcal{F}}$ .*

- (i) *If  $\mathcal{F}$  is space-like and  $m + 1 \leq N$  then either  $\mathcal{F}$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}} = \mathbb{E}^N \cap V_{\mathcal{F}}^\perp$ , or this holds for the image of  $\mathcal{F}$  under an inversion in  $\mathbb{R}^N$ .*
- (ii) *If  $\mathcal{F}$  is space-like and  $m = N$  then there exist a space-like family of affine subspaces  $\hat{\mathcal{F}}$  in  $\mathbb{R}^{N+1}$  with axis  $\mathcal{H}_{\hat{\mathcal{F}}} = \{0\}$ , a sphere  $\mathbb{S}^N(c) \subset \mathbb{R}^{N+1}$  centered at the origin and a stereographic projection  $\mathcal{P}: \mathbb{S}^N(c) \setminus \{z\} \rightarrow \mathbb{R}^N$ , where  $\mathbb{R}^N$  is identified with the tangent space of  $\mathbb{S}^N(c)$  at  $-z$ , such that each  $S^\alpha \in \mathcal{F}$  is given by  $S^\alpha = \mathcal{P}(\hat{S}^\alpha \cap \mathbb{S}^N(c))$  for some  $\hat{S}^\alpha \in \hat{\mathcal{F}}$ .*
- (iii) *If  $\mathcal{F}$  is light-like then either  $\mathcal{F}$  is a light-like family of affine subspaces as in Proposition 8-(ii) or this holds for the image of  $\mathcal{F}$  under an inversion in  $\mathbb{R}^N$ .*
- (iv) *If  $\mathcal{F}$  is time-like and  $m \leq N$  then either  $\mathcal{F}$  is a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}} = \mathbb{E}^N \cap V_{\mathcal{F}}$ , or this holds for the image of  $\mathcal{F}$  under an inversion in  $\mathbb{R}^N$ .*

*Proof:* (i) If  $w \in V_{\mathcal{F}}^\perp$ , then  $\mathcal{F}$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}} = \mathbb{E}^N \cap V_{\mathcal{F}}^\perp$  by Proposition 8-(i). Otherwise, since  $m + 1 = \dim V_{\mathcal{F}} \leq N$ , there exists  $\zeta \in \mathbb{E}^N \cap V_{\mathcal{F}}^\perp$ . Let  $I$  be the inversion in  $\mathbb{R}^N$  determined by  $T \in \mathbb{O}_1(N+2)$  as in (3) with  $v = \zeta + (1/2)w$ , so that  $w = -2T(\zeta)$ . Then  $\tilde{\mathcal{F}} = I(\mathcal{F}) := \{I(S^\alpha) : S^\alpha \in \mathcal{F}\}$  satisfies  $V_{\tilde{\mathcal{F}}} = T(V_{\mathcal{F}})$ . Since  $w \in T(V_{\mathcal{F}})^\perp$ , it follows that  $\tilde{\mathcal{F}}$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\tilde{\mathcal{F}}} = \mathbb{E}^N \cap V_{\tilde{\mathcal{F}}}^\perp$ .

(ii) Consider the model  $\mathbb{E}^{N+1} \subset \mathbb{L}^{N+3}$  of  $\mathbb{R}^{N+1}$ , and let  $\mathbb{R}^N$  be represented in this model by  $\mathbb{E}^N = \mathbb{E}^{N+1} \cap \{e_0\}^\perp$ , where  $e_0$  is a unit space-like vector orthogonal to  $w$ . Also, let  $\mathbb{L}^{N+2}$  be identified with  $\{e_0\}^\perp$ . For each  $S^\alpha = \mathbb{E}^N \cap V^{\alpha\perp} \in \mathcal{F}$ , define  $\hat{S}^\alpha = \mathbb{E}^{N+1} \cap V^{\alpha\perp}$  and  $\hat{\mathcal{F}} = (\hat{S}^\alpha)_{\alpha \in \Lambda}$ , where  $V^\alpha$  is now regarded as a subspace of  $\mathbb{L}^{N+3}$ . Then  $\hat{S}^\alpha$  is the unique sphere or affine subspace in  $\mathbb{R}^{N+1}$  that intersects  $\mathbb{R}^N$  orthogonally along  $S^\alpha$ . Now let  $\mathbb{L}^2$  be the time-like plane spanned by  $e_0$  and the one-dimensional orthogonal complement  $V^\perp$  of  $V$  in  $\mathbb{L}^{N+2}$ . Let  $\zeta \in \mathbb{E}^{N+1} \cap \mathbb{L}^2$  and extend  $\zeta$  to a pseudo-orthonormal basis  $\{\zeta, \eta\}$  of  $\mathbb{L}^2$ . Define

$$v = \frac{1}{\langle \zeta, e_0 \rangle} \zeta + \frac{1}{2} \langle \zeta, e_0 \rangle w. \quad (4)$$

Then  $\langle v, e_0 \rangle = 1 = \langle v, v \rangle$ , and hence  $p_0 = v - e_0$  belongs to  $\mathbb{E}^{N+1}$  and satisfies  $p_0 \in \mathbb{R}^N \cap S$ , where  $S = \mathbb{E}^{N+1} \cap \{v\}^\perp$ . Therefore  $S$  and  $\mathbb{R}^N$  are tangent at  $p_0$ . Consider the inversion  $I$  in  $\mathbb{R}^{N+1}$  determined by  $T \in \mathcal{O}_1(N+3)$  as in (3), with  $v$  given by (4). Then  $w = -2\langle \zeta, e_0 \rangle^{-2}T(\zeta) \in T(V)^\perp$  and  $\bar{q} = -(\langle \zeta, e_0 \rangle^2/2)T(\eta) \in \mathbb{E}^{N+1} \cap T(V)^\perp$ , and hence  $I$  takes  $\tilde{\mathcal{F}}$  into a space-like family of affine subspaces with axis  $\{\bar{q}\}$ . Define  $\mathbb{S}^N(c) = I(\mathbb{R}^N) = \mathbb{E}^{N+1} \cap \{T(e_0)\}^\perp$ . Since  $e_0 = \alpha\eta + (2\alpha)^{-1}\zeta$ , with  $\alpha = \langle e_0, \zeta \rangle$ , it follows that  $T(e_0) = \beta\bar{q} + (2\beta)^{-1}w$ , with  $\beta = -2\alpha\langle \zeta, e_0 \rangle^{-2}$ , and thus  $\mathbb{S}^N(c)$  has center at  $\bar{q}$ . Now, the restriction of the inversion  $I$  itself to  $\mathbb{S}^N(c) \setminus \{z\}$  can be regarded as a stereographic projection  $\mathcal{P}$  of  $\mathbb{S}^N(c) \setminus \{z\}$  onto  $\mathbb{R}^N$ . Moreover, we have that  $\mathcal{P}(I(\hat{S}^\alpha) \cap \mathbb{S}^N(c)) = \hat{S}^\alpha \cap \mathbb{R}^N = S^\alpha$  for every  $\hat{S}^\alpha \in \hat{\mathcal{F}}$ . Finally, we may assume that  $\bar{q}$  is the origin in  $\mathbb{R}^{N+1}$ .

(iii) If  $w \in V_{\mathcal{F}} \cap V_{\tilde{\mathcal{F}}}^\perp$ , then there exists a (non-unique) space-like subspace  $U$  of dimension  $m$  such that  $V_{\mathcal{F}} = U \oplus \text{span}\{w\}$ , and it follows from Proposition 8-(ii) that  $\mathcal{F}$  is a light-like family of affine subspaces whose axis is the  $(N-m)$ -dimensional subspace  $\mathbb{R}_{\mathcal{F}}^{N-m}$  that is parallel to the affine subspace  $\mathbb{E}^N \cap U^\perp$ . Otherwise, we may assume that there exists  $\zeta \in \mathbb{E}^N$  such that  $\zeta \in V_{\mathcal{F}} \cap V_{\tilde{\mathcal{F}}}^\perp$ . Arguing as in case (i), we find that there exists an inversion  $I$  in  $\mathbb{R}^N$  determined by  $T \in \mathcal{O}_1(N+2)$  as in (3) such that  $\tilde{\mathcal{F}} = I(\mathcal{F})$  is a light-like family of affine subspaces whose axis is the subspace  $\mathbb{R}_{\tilde{\mathcal{F}}}^{N-m}$  that is parallel to the affine subspace  $\mathbb{E}^N \cap T(U)^\perp$ .

(iv) If  $w \in V_{\mathcal{F}}$ , then  $\mathcal{F}$  is a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}} = \mathbb{E}^N \cap V_{\mathcal{F}}$ . Otherwise, arguing again as in case (i) we obtain an inversion  $I$  in  $\mathbb{R}^N$  determined by  $T \in \mathcal{O}_1(N+2)$  as in (3) such that  $\tilde{\mathcal{F}} = I(\mathcal{F})$  is a standard time-like family of spheres with axis  $\tilde{\mathcal{H}}_{\mathcal{F}} = \mathbb{E}^N \cap T(V_{\mathcal{F}})$ . ■

## 4 Orthogonal families of spheres in $\mathbb{R}^N$

Our aim in this section is to give a characterization of certain orthogonal families of spheres and affine subspaces in Euclidean space, which is the principal tool in the proof of Theorem 25.

**Lemma 11** *Let  $\Lambda = \Lambda_0 \times \cdots \times \Lambda_k$ , where each  $\Lambda_i$  is a set with at least two elements, and let  $\mathcal{F}_i = (S_i^\alpha)_{\alpha \in \Lambda_i}$ ,  $0 \leq i \leq k$ , be families of spheres or affine subspaces in  $\mathbb{R}^N$  with the following properties:*

- (i) *Each  $S_i^\alpha \in \mathcal{F}_i$  has codimension  $n_i \geq 1$ , and  $\sum_{i=0}^k n_i = n \leq N$ .*
- (ii) *Two elements  $S_i^\alpha, S_i^\beta \in \mathcal{F}_i$  coincide if and only if  $\pi_i(\alpha) = \pi_i(\beta)$ , where  $\pi_i: \Lambda \rightarrow \Lambda_i$  denotes the projection onto  $\Lambda_i$ .*
- (iii)  *$S_i^\alpha$  is orthogonal to  $S_j^\alpha$  for every  $\alpha \in \Lambda$  and for every  $0 \leq i \neq j \leq k$ .*

*Then one of the following holds:*

- (a) After reordering the families, there exist  $-1 \leq s \leq k$  with  $k-s \leq N-n$ , an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  with  $m_i \geq n_i$  for  $0 \leq i \leq s$ ,  $m_i \geq n_i + 1$  for  $s+1 \leq i \leq k$  and  $m_{k+1} \geq 0$ , and affine  $(N-m_i)$ -dimensional subspaces  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i^\perp}$ ,  $s+1 \leq i \leq k$ , such that, up to an inversion in  $\mathbb{R}^N$ ,  $\mathcal{F}_i$  is a light-like family of affine subspaces with axis  $\mathbb{R}^{m_i^\perp}$  for  $0 \leq i \leq s$  and a space-like family of affine subspaces with axis  $\mathcal{H}_i$  for  $s+1 \leq i \leq k$ .
- (b)  $k \leq N-n$  and there exist an orthogonal decomposition  $\mathbb{R}^{N+1} = \oplus_{i=0}^k \mathbb{R}^{m_i}$  with  $m_i \geq n_i + 1$ , space-like families of affine subspaces  $\hat{\mathcal{F}}_i$  with axis  $\mathcal{H}_{\hat{\mathcal{F}}_i} = \mathbb{R}^{m_i^\perp}$ ,  $0 \leq i \leq k$ , a sphere  $\mathbb{S}^N(c) \subset \mathbb{R}^{N+1}$  centered at the origin and a stereographic projection  $\mathcal{P}: \mathbb{S}^N(c) \setminus \{z\} \rightarrow \mathbb{R}^N$ , where  $\mathbb{R}^N$  is identified with the tangent space of  $\mathbb{S}^N(c)$  at  $-z$ , such that each  $S_i^\alpha \in \mathcal{F}_i$  is given by  $S_i^\alpha = \mathcal{P}(\hat{S}_i^\alpha \cap \mathbb{S}^N(c))$  for some  $\hat{S}_i^\alpha \in \hat{\mathcal{F}}_i$ .
- (c)  $k \leq N-n+1$  and for exactly one  $j \in \{0, \dots, k\}$  there exist an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  with  $m_j \geq n_j - 1$ ,  $m_i \geq n_i + 1$  for  $0 \leq i \neq j \leq k$  and  $m_{k+1} \geq 0$ , affine  $(N-m_i)$ -dimensional subspaces  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i^\perp}$ ,  $0 \leq i \neq j \leq k$ , and an affine  $m_j$ -dimensional affine subspace  $\mathcal{H}_j \subset \cap_{i \neq j} \mathcal{H}_i$  parallel to  $\mathbb{R}^{m_j}$  (which reduces to a point  $\bar{q}$  if  $m_j = 0$ ), such that, up to an inversion in  $\mathbb{R}^N$ ,  $\mathcal{F}_j$  is a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_j} = \mathcal{H}_j$  (which is a family of hyperspheres centered at  $\bar{q}$  if  $m_j = 0$ ) and  $\mathcal{F}_i$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_i} = \mathcal{H}_i$  for  $0 \leq i \neq j \leq k$ .

*Proof:* Write  $S_i^\alpha = \mathbb{E}^N \cap V_i^{\alpha^\perp}$ , where  $V_i^\alpha$  is a space-like subspace of  $\mathbb{L}^{N+2}$  of dimension  $n_i$  and let  $V_i = V_{\mathcal{F}_i}$  be the subspace of  $\mathbb{L}^{N+2}$  spanned by all  $V_i^\alpha$ ,  $\alpha \in \Lambda$ . By (ii) and the assumption that  $\Lambda_i$  has at least two elements, we have that the dimension of  $V_i$  is at least  $n_i + 1$ . On the other hand, it follows from (ii) and (iii) that each  $S_i^\alpha \in \mathcal{F}_i$  is orthogonal to every  $S_j^\beta \in \mathcal{F}_j$ ,  $0 \leq i \neq j \leq k$ . Therefore,  $V_i^\alpha \subset V_j^{\beta^\perp}$  for every  $\alpha, \beta \in \Lambda$ , which implies that  $V_i \subset V_j^\perp$ ,  $0 \leq i \neq j \leq k$ . Then one of the following holds:

- (i)  $V_i$  is either degenerate or space-like for  $0 \leq i \leq k$  and if  $V_i$  is space-like for every  $0 \leq i \leq k$  then  $V = \oplus_{i=0}^k V_i$  has dimension at most  $N$ .
- (ii)  $V$  is a space-like subspace of dimension  $N+1$ .
- (iii) There exists (exactly) one  $j \in \{0, \dots, k\}$  such that  $V_j$  is time-like.

Assume first that (i) holds. After reordering the families, there exist  $-1 \leq s \leq k$  and  $\zeta \in \mathbb{V}^{N+1}$  such that  $V_i$  orthogonally decomposes as  $V_i = U_i \oplus \text{span}\{\zeta\}$  for  $0 \leq i \leq s$  and  $V_i$  is space-like for  $s+1 \leq i \leq k$ . In the latter case set  $U_i = V_i$ . Then  $U_i$  is a space-like subspace for every  $0 \leq i \leq k$ , whose dimension  $m_i$  is at least  $n_i$  for  $0 \leq i \leq s$  and at least  $n_i + 1$  for  $s+1 \leq i \leq k$ . By taking a multiple of  $\zeta$  we may assume either that  $\zeta = w$  or that  $\zeta \in \mathbb{E}^N$ , that is,  $\langle \zeta, w \rangle = 1$ . In the first case, define the affine subspaces  $\mathcal{H}_i$  by

$\mathcal{H}_i = \mathbb{E}^N \cap U_i^\perp$ ,  $0 \leq i \leq k$ , let  $\mathbb{R}^{m_i}$  be the subspace of  $\mathbb{R}^N$  such that  $\mathbb{R}^{m_i \perp}$  is parallel to  $\mathcal{H}_i$ , and let  $\mathbb{R}^{m_{k+1}} = \bigcap_{i=0}^k \mathbb{R}^{m_i \perp}$ . Then  $\mathbb{R}^N = \bigoplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  is an orthogonal decomposition as in the statement and  $\mathcal{F}_i$ ,  $0 \leq i \leq s$  (respectively,  $s+1 \leq i \leq k$ ), is a light-like (respectively, space-like) family of affine subspaces with axis  $\mathbb{R}^{m_i \perp}$  (respectively,  $\mathcal{H}_i$ ).

Now suppose  $\zeta \in \mathbb{E}^N$ . Let  $I$  be the inversion in  $\mathbb{R}^N$  determined by  $T \in \mathcal{O}_1(N+2)$  as in (3) with  $v = \zeta + (1/2)w$ , so that  $w = -2T(\zeta)$ . Arguing as before with  $U_i$  replaced by  $T(U_i)$ ,  $0 \leq i \leq k$ , we obtain an orthogonal decomposition  $\mathbb{R}^N = \bigoplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  as in the statement and affine  $(N-m_i)$ -dimensional subspaces  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i \perp}$ ,  $0 \leq i \leq k$ , such that  $I$  takes  $\mathcal{F}_i$ ,  $0 \leq i \leq s$  (respectively,  $s+1 \leq i \leq k$ ) into a light-like (respectively, space-like) family of affine subspaces with axis  $\mathbb{R}^{m_i \perp}$  (respectively,  $\mathcal{H}_i$ ).

In case (ii), we must have  $n+k+1 = \sum_{i=0}^k (n_i+1) \leq \dim V = N+1$ , and hence  $k \leq N-n$ . We argue as in the proof of Proposition 10-(ii): consider the model  $\mathbb{E}^{N+1} \subset \mathbb{L}^{N+3}$  of  $\mathbb{R}^{N+1}$ , identify  $\mathbb{L}^{N+2}$  with  $e_0^\perp$ , where  $e_0$  is a unit space-like vector orthogonal to  $w$ , and let  $\mathbb{R}^N$  be represented by  $\mathbb{E}^N = \mathbb{E}^{N+1} \cap e_0^\perp$ . For each  $S_i^\alpha = \mathbb{E}^N \cap V_i^{\alpha \perp} \in \mathcal{F}_i$ , let  $\hat{S}_i^\alpha$  be the unique sphere in  $\mathbb{R}^{N+1}$  that intersects  $\mathbb{R}^N$  orthogonally along  $S_i^\alpha$ , that is,  $\hat{S}_i^\alpha = \mathbb{E}^{N+1} \cap V_i^{\alpha \perp}$ , where  $V_i^\alpha$  is now regarded as a subspace of  $\mathbb{L}^{N+3}$ . Then we have an orthogonal decomposition  $\mathbb{L}^{N+3} = \bigoplus_{i=0}^k V_i \oplus \mathbb{L}^2$ , where  $\mathbb{L}^2$  is the time-like plane spanned by  $e_0$  and the one-dimensional orthogonal complement  $V^\perp$  of  $V$  in  $\mathbb{L}^{N+2}$ . Proceeding exactly as in the proof of Proposition 10-(ii), we obtain an inversion  $I$  in  $\mathbb{R}^{N+1}$  determined by  $T \in \mathcal{O}_1(N+3)$ , a point  $\bar{q} \in \mathbb{R}^{N+1}$  (which we may assume to be the origin), and an orthogonal decomposition  $\mathbb{R}^{N+1} = \bigoplus_{i=0}^k \mathbb{R}^{m_i}$ , where  $\mathbb{R}^{m_i}$  is the subspace of  $\mathbb{R}^{N+1}$  defined by  $\bar{q} + \mathbb{R}^{m_i \perp} = \mathbb{E}^{N+1} \cap T(V_i)^\perp$  for  $0 \leq i \leq k$ , such that  $I$  takes every  $\hat{\mathcal{F}}_i$  into a space-like family of affine subspaces with axis  $\mathcal{H}_{\hat{\mathcal{F}}_i} = \bar{q} + \mathbb{R}^{m_i \perp}$ . As before, the restriction of the inversion  $I$  itself to  $\mathbb{S}^N(c) \setminus \{z\}$ , where  $\mathbb{S}^N(c) = I(\mathbb{R}^N)$  is a sphere centered at  $\bar{q}$ , can be regarded as a stereographic projection  $\mathcal{P}$  of  $\mathbb{S}^N(c) \setminus \{z\}$  onto  $\mathbb{R}^N$ , and  $\mathcal{P}(I(\hat{S}_i^\alpha) \cap \mathbb{S}^N(c)) = \hat{S}_i^\alpha \cap \mathbb{R}^N = S_i^\alpha$  for every  $\hat{S}_i^\alpha \in \hat{\mathcal{F}}_i$ .

Finally, if (iii) holds, we have that  $n+k+1 = \sum_{i=0}^k (n_i+1) \leq \dim V \leq N+2$ , thus  $k \leq N-n+1$ . Assume first that  $w \in V_j$  and set  $m_j = \dim V_j - 2 \geq n_j - 1$ . Define the affine subspaces  $\mathcal{H}_i$ ,  $0 \leq i \neq j \leq k$ , by  $\mathcal{H}_i = \mathbb{E}^N \cap V_i^\perp$ , and set  $\mathcal{H}_j = \mathbb{E}^N \cap V_j$ . Then  $\mathcal{H}_j$  is either an affine subspace of dimension  $m_j$  if  $m_j > 0$  or a single point  $\bar{q} \in \mathbb{R}^N$  otherwise, in which case  $n_j$  must be equal to one, and the affine subspace  $\mathcal{H}_i$  has codimension  $m_i \geq n_i + 1$  and contains  $\mathcal{H}_j$  for every  $0 \leq i \neq j \leq k$ . Let  $\mathbb{R}^{m_i}$  be the subspace of  $\mathbb{R}^N$  such that  $\mathbb{R}^{m_i \perp}$  is parallel to  $\mathcal{H}_i$  and define  $\mathbb{R}^{m_j}$  as the subspace parallel to the affine subspace  $\mathcal{H}_j$  if  $m_j > 0$ , or the trivial subspace otherwise. Then we obtain an orthogonal decomposition  $\mathbb{R}^N = \bigoplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  as in the statement such that  $\mathcal{F}_j$  is a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_j} = \mathcal{H}_j$  (which is a family of hyperspheres centered at  $\bar{q}$  if  $m_j = 0$ ) and  $\mathcal{F}_i$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_i} = \mathcal{H}_i$  for  $0 \leq i \neq j \leq k$ .

If  $w \notin V_j$ , choose  $\zeta \in \mathbb{E}^N \cap V_j$  and let  $I$  be the inversion in  $\mathbb{R}^N$  determined by  $T \in \mathcal{O}_1(N+2)$  as in (3) with  $v = \zeta + (1/2)w$ . Then  $w = -2T(\zeta) \in T(V_j)$  and we can argue as before with  $V_j$ ,  $V$  and  $V_i$  replaced by  $T(V_j)$ ,  $T(V)$  and  $T(V_i)$ , respectively. We



obtain affine subspaces  $\mathcal{H}_i$ ,  $0 \leq i \neq j \leq k$ , of codimension  $m_i \geq n_i + 1$  and either an affine subspace  $\mathcal{H}_j$  of dimension  $m_j = \dim V_j - 2 \geq n_j - 1$  if  $m_j > 0$  or a point  $\bar{q}$  if  $m_j = 0$ , and an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  such that  $I$  takes  $\mathcal{F}_j$  into a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_j} = \mathcal{H}_j$  and  $\mathcal{F}_i$  into a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_i} = \mathcal{H}_i$  for  $0 \leq i \neq j \leq k$ . ■

**Remarks 12.** (i) In case (b) we have the following alternative description:

(b') For any  $j \in \{0, \dots, k\}$  there exist an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m'_i}$  with  $m'_j \geq n_j$  and  $m'_i \geq n_i + 1$  for  $0 \leq i \neq j \leq k$ , and affine  $(N - m'_i)$ -dimensional subspaces  $\mathcal{H}'_i$  parallel to  $\mathbb{R}^{m'_i \perp}$ , all of which contain an affine  $m'_j$ -dimensional subspace  $\mathcal{H}'_j$  parallel to  $\mathbb{R}^{m'_j}$ , such that, up to an inversion in  $\mathbb{R}^N$ ,  $\mathcal{F}_j$  is a standard time-like family of spheres or affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_j} = \mathcal{H}'_j$  and  $\mathcal{F}_i$  is a space-like family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_i} = \mathcal{H}'_i$  for  $0 \leq i \neq j \leq k$ . This can be seen by arguing as in the proof of part (c) with  $V_j$  replaced by the time-like subspace  $\tilde{V}_j = V_j \oplus V^\perp$ .

(ii) The inversion  $I$  in parts (a) and (c) of Lemma 11 can be taken (as it is in the proof) with respect to a sphere of *unit* radius.

The next two lemmas will be used only in our geometric proof of Nölker's theorem (in Euclidean space) in Section 6.

**Lemma 13** *Under the assumptions of Lemma 11, assume further that any element  $S_i^\alpha \in \mathcal{F}_i$ ,  $0 \leq i \leq k$ , is an affine subspace of  $\mathbb{R}^N$ . Then, after reordering the families, there exist  $-1 \leq s \leq k$  with  $k - s \leq N - n$ , an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  with  $m_i \geq n_i$  for  $0 \leq j \leq s$ ,  $m_i \geq n_i + 1$  for  $s + 1 \leq i \leq k$  and  $m_{k+1} \geq 0$ , and affine  $(N - m_i)$ -dimensional subspaces  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i \perp}$ ,  $s + 1 \leq i \leq k$ , such that, for  $0 \leq i \leq s$  (respectively,  $s + 1 \leq i \leq k$ ),  $\mathcal{F}_i$  is a light-like (respectively, space-like) family of affine subspaces with axis  $\mathbb{R}^{m_i \perp}$  (respectively,  $\mathcal{H}_i$ ).*

*Proof:* Define  $V_i = V_{\mathcal{F}_i}$ ,  $0 \leq i \leq k$ , and  $V = \oplus_{i=0}^k V_i$  as in the proof of Lemma 11. By the assumption that any element  $S_i^\alpha \in \mathcal{F}_i$ ,  $0 \leq i \leq k$ , is an affine subspace of  $\mathbb{R}^N$ , we have that  $w \in V^\perp$ . After reordering the families, we may assume that there exists  $-1 \leq s \leq k$  such that  $w \in V_i$  for  $0 \leq i \leq s$ . Write  $V_i = U_i \oplus \text{span}\{w\}$  for  $0 \leq i \leq s$  and  $U_i = V_i$  otherwise. Then  $U_i$  is a space-like subspace for every  $0 \leq i \leq k$ , whose dimension  $m_i$  is at least  $n_i$  for  $0 \leq i \leq s$  and at least  $n_i + 1$  for  $s + 1 \leq i \leq k$ . Define the affine subspaces  $\mathcal{H}_i$  by  $\mathcal{H}_i = \mathbb{E}^N \cap U_i^\perp$ ,  $0 \leq i \leq k$ , let  $\mathbb{R}^{m_i}$  be the subspace of  $\mathbb{R}^N$  such that  $\mathbb{R}^{m_i \perp}$  is parallel to  $\mathcal{H}_i$ , and set  $\mathbb{R}^{m_{k+1}} = \cap_{i=1}^k \mathbb{R}^{m_i \perp}$ . Then  $\mathbb{R}^N = \oplus_{i=0}^{k+1} \mathbb{R}^{m_i}$  is an orthogonal decomposition as in the statement and, by Proposition 8, for  $0 \leq i \leq s$  (respectively,  $s + 1 \leq i \leq k$ ),  $\mathcal{F}_i$  is a light-like (respectively, space-like) family of affine subspaces with axis  $\mathbb{R}^{m_i \perp}$  (respectively,  $\mathcal{H}_i$ ). ■

**Lemma 14** *Under the assumptions of Lemma 11, assume further that  $k = 1$  and that any element  $S_1^\alpha \in \mathcal{F}_1$  is an affine subspace of  $\mathbb{R}^N$ . Then one of the following holds:*

- (a) *There exist an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^2 \mathbb{R}^{m_i}$  such that, for  $0 \leq i \leq 1$ ,  $\mathcal{F}_i$  is either a light-like family of affine subspaces with axis  $\mathbb{R}^{m_i^\perp}$ , in which case  $m_i \geq n_i$ , or a space-like family of affine subspaces whose axis is an affine  $(N-m_i)$ -dimensional subspace  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i^\perp}$ , in which case  $m_i \geq n_i + 1$ .*
- (b) *There exist an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^{m_0} \oplus \mathbb{R}^{m_1}$ , with  $m_0 \geq n_0 - 1$ ,  $m_1 \geq n_1 + 1$ , and an  $m_0$ -dimensional affine subspace  $\mathcal{H}$  parallel to  $\mathbb{R}^{m_0}$  such that  $\mathcal{F}_1$  is a space-like family of affine subspaces with axis  $\mathcal{H}$  and  $\mathcal{F}_0$  is contained in a standard time-like family of affine subspaces or spheres with axis  $\mathcal{H}$ .*

*Proof:* Define  $V_0, V_1$  and  $V$  as in the proof of Lemma 11. The assumption that any element  $S_1^\alpha \in \mathcal{F}_1$  is an affine subspace of  $\mathbb{R}^N$  implies that  $w \in V_1^\perp$ . Therefore either  $w \in V_0^\perp$  and both  $V_0$  and  $V_1$  are degenerate or space-like, or  $w \notin V_0^\perp$  and  $V_1$  is necessarily space-like. If the first possibility holds, then for  $0 \leq i \leq 1$  let  $U_i$  be a space-like subspace of dimension  $m_i \geq n_i$  such that  $V_i = U_i \oplus \mathbb{R}w$  if  $V_i$  is degenerate, and set  $U_i = V_i$  if  $V_i$  is space-like. Define  $\mathcal{H}_i = \mathbb{E}^N \cap U_i^\perp$ ,  $0 \leq i \leq 1$ , and let  $\mathbb{R}^{m_i}$  be the subspace such that  $\mathbb{R}^{m_i^\perp}$  is parallel to  $\mathcal{H}_i$  for  $0 \leq i \leq 1$ . Then we obtain an orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^2 \mathbb{R}^{m_i}$  such that  $\mathcal{F}_i$  is either a light-like family of affine subspaces with axis  $\mathbb{R}^{m_i^\perp}$  or a space-like family of affine subspaces with axis  $\mathcal{H}_i$ , according to  $V_i$  being degenerate or space-like, respectively.

Now suppose that  $w \notin V_0^\perp$ , and hence that  $V_1$  is space-like. Define  $\mathcal{H} = \mathbb{E}^N \cap V_1^\perp$ . Then the statement in part (b) holds by defining  $\mathbb{R}^{m_1}$  as the subspace of  $\mathbb{R}^N$  such that  $\mathbb{R}^{m_0} = \mathbb{R}^{m_1^\perp}$  is parallel to  $\mathcal{H}$ . ■

## 5 Extrinsic spheres of Euclidean submanifolds

In this section we derive two results on extrinsic spheres of Euclidean submanifolds that play a key rôle in the proof of Theorem 25 and are also interesting on their own.

**Proposition 15** *Let  $f: M^n \rightarrow \mathbb{R}^N$  and  $g: L^k \rightarrow M^n$  be isometric immersions. Then the following assertions are equivalent:*

- i)  *$g$  is an extrinsic sphere whose mean curvature vector has length  $1/r$ ,  $r > 0$  (respectively,  $g$  is totally geodesic) and*

$$\alpha_f(g_*X, Z) = 0 \text{ for all } X \in \Gamma(TL), \quad Z \in \Gamma(T_g^\perp L). \quad (5)$$

- ii)  *$f(g(L))$  is contained in a sphere with radius  $r$  (respectively, affine subspace) in  $\mathbb{R}^N$  of dimension  $(N - n + k)$  whose normal space along  $f(g(L))$  is  $f_*T_g^\perp L$ .*

*Proof:* Assume that  $g$  is an extrinsic sphere whose mean curvature vector  $H_g$  has length  $1/r$ ,  $r > 0$ . Then

$$\alpha_g(X, Y) = \langle X, Y \rangle H_g \quad (6)$$

and

$$\langle \nabla_X H_g, Z \rangle = 0 \text{ for all } X \in \Gamma(TL), Z \in \Gamma(T_g^\perp L), \quad (7)$$

where  $\nabla$  denotes the connection in  $M^n$ . In particular,

$$\nabla_X H_g = -r^{-2} g_* X \text{ for all } X \in \Gamma(TL). \quad (8)$$

Set  $h = f \circ g + r^2 f_* H_g$ . Using (5) and (8), we obtain

$$h_* X = f_* g_* X + r^2 (-r^{-2} f_* g_* X + \alpha_f(g_* X, H_g)) = 0$$

for all  $X \in \Gamma(TL)$ , and hence  $h = P_0 \in \mathbb{R}^N$  is a constant map. It follows that  $f(g(L))$  is contained in a hypersphere with radius  $r$  centered at  $P_0$ . Moreover, let  $W$  be the subbundle of  $T_g^\perp L$  orthogonal to  $H_g$ . By (6), for any  $X, Y \in \Gamma(TL)$  and  $Z \in \Gamma(W)$  we have

$$\langle \nabla_X Z, g_* Y \rangle = -\langle \nabla_X g_* Y, Z \rangle = \langle \alpha_g(X, Y), Z \rangle = 0,$$

whereas (7) implies that  $\langle \nabla_X Z, H_g \rangle = 0$ , and thus  $\nabla_X Z \in \Gamma(W)$ . Using (5) we obtain

$$\tilde{\nabla}_X f_* Z = f_* \nabla_X Z + \alpha_f(g_* X, Z) = f_* \nabla_X Z \in f_* W.$$

Hence  $f_* W$  is a parallel subbundle in  $\mathbb{R}^N$  orthogonal to  $f(g(L))$ , and *ii*) follows. The proof for the case where  $g$  is totally geodesic is similar and easier.

Conversely, assume that  $f(g(L))$  is contained in a sphere in  $\mathbb{R}^N$  of dimension  $(N-n+k)$  with center at  $P_0$  and radius  $r$  whose normal space along  $f(g(L))$  is  $f_* T_g^\perp L$ . Then, there exists a unit vector field  $\eta \in T_g^\perp L$  such that  $f \circ g + r\eta = P_0$ . Moreover, if  $W$  is the subbundle of  $T_g^\perp L$  orthogonal to  $\eta$ , then  $f_* W$  is parallel in  $\mathbb{R}^N$  along  $f(g(L))$ . Hence

$$f_* g_* X + r \tilde{\nabla}_X \eta = 0 \quad (9)$$

and

$$(\tilde{\nabla}_X f_* Z)_{(f_* W)^\perp} = 0 \text{ for all } X \in \Gamma(TL), Z \in \Gamma(W). \quad (10)$$

Taking the component in  $T_f^\perp M$  of (9) and (10) yields (5). On the other hand, taking the inner product of (9) and (10) with  $f_* g_* Y$ ,  $Y \in TL$ , implies that

$$\alpha_g(X, Y) = (1/r) \langle X, Y \rangle \eta \text{ for all } X, Y \in \Gamma(TL),$$

and thus  $g$  is umbilical with mean curvature vector  $H_g = (1/r)\eta$ . Finally, parallelism of  $H_g$  in the normal connection of  $g$  follows by taking the  $\eta$ -component of (10). ■

**Proposition 16** *Let  $f: M^n \rightarrow \mathbb{R}^N$  and  $g: L^k \rightarrow M^n$ ,  $k \geq 2$ , be isometric immersions. Assume that  $g$  is umbilical and that (5) holds. Then  $g$  is an extrinsic sphere.*

*Proof:* The Codazzi equation for  $g$  is

$$(\nabla_X^\perp \alpha_g)(Y, T) - (\nabla_Y^\perp \alpha_g)(X, T) = (R(g_*X, g_*Y)g_*T)_{T_g^\perp L} \quad (11)$$

for any  $X, Y, T \in \Gamma(TL)$ , where  $R$  stands for the curvature tensor of  $M^n$ . By the Gauss equation for  $f$  and (5),

$$R(g_*X, g_*Y)Z = A_{\alpha_f(g_*Y, Z)}^f g_*X - A_{\alpha_f(g_*X, Z)}^f g_*Y = 0$$

for all  $X, Y \in \Gamma(TL)$ ,  $Z \in \Gamma(T_g^\perp L)$ , hence the right-hand-side of (11) vanishes. It follows that the mean curvature vector  $H_g$  of  $g$  satisfies  $\langle X, T \rangle \nabla_Y^\perp H_g = \langle Y, T \rangle \nabla_X^\perp H_g$  for all  $X, Y, T \in \Gamma(TL)$ . Choosing  $Y = T$  orthogonal to  $X$ , we conclude that  $H_g$  is parallel in the normal connection of  $g$ . ■

**Remark 17.** Propositions 15 and 16 can be easily extended to the case of nonflat ambient space forms.

## 6 The decomposition theorem of Nölker

We start this section by reviewing from [No] the notions of *warped product representations* of space forms and *warped product of isometric immersions*. Then we state Nölker's decomposition theorem for isometric immersions of warped products into space forms and give a short geometric proof of it in Euclidean space.

### 6.1 Warped product representations of space forms

We first define a *WP-net*  $\hat{\mathcal{E}} = (\hat{E}_i)_{i=0, \dots, k}$  on an open dense subset of  $\mathbb{Q}_c^N$ ,  $c \geq 0$ , as follows. Assume first that  $c = 0$ . Let  $\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m_i}$  be an orthogonal decomposition of  $\mathbb{R}^N$  with  $\mathbb{R}^{m_0}$  possibly a trivial subspace and  $\mathbb{R}^{m_i}$  nontrivial for  $1 \leq i \leq k$ . Let  $\mathcal{H}$  be an affine  $m_0$ -dimensional subspace of  $\mathbb{R}^N$  parallel to  $\mathbb{R}^{m_0}$  ( $\mathcal{H}$  reduces to a single point  $\bar{q} \in \mathbb{R}^N$  if  $\mathbb{R}^{m_0}$  is trivial). Fix a number  $s \in \{0, \dots, k\}$ . For  $s+1 \leq i \leq k$ , let  $\mathcal{H}_i$  be an  $(N-m_i)$ -dimensional affine subspace that contains  $\mathcal{H}$  and is parallel to  $\mathbb{R}^{m_i^\perp}$ . For each  $q \in G = \mathbb{R}^N \setminus \bigcup_{i=s+1}^k \mathcal{H}_i$ , define  $\hat{S}_i(q) = q + \mathbb{R}^{m_i}$  for  $1 \leq i \leq s$  and let  $\hat{S}_i(q)$  be the sphere in  $q + \mathbb{R}^{m_i}$  through  $q$  centered at the intersection of  $q + \mathbb{R}^{m_i}$  with  $\mathcal{H}_i$  for  $s+1 \leq i \leq k$ . Set  $\hat{E}_i(q) = T_q \hat{S}_i(q)$ ,  $1 \leq i \leq k$ . Finally, let  $-a_i(q)$  be the mean curvature vector of  $\hat{S}_i(q)$  at  $q$  for  $s+1 \leq i \leq k$  and set  $\hat{E}_0(q) = \mathbb{R}^{m_0} \oplus \text{span}\{a_i(q) : s+1 \leq i \leq k\}$ . If  $s < k$ , we say that  $\hat{\mathcal{E}} = (\hat{E}_i)_{i=0, \dots, k}$  is the *WP-net* on  $G \subset \mathbb{R}^N$  determined by  $(\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m_i}, \mathcal{H}, s)$ . For  $s = k$ , the net  $\hat{\mathcal{E}}$  reduces to the *RP-net* on  $\mathbb{R}^N$  determined by the orthogonal decomposition  $\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m_i}$ .

Assume now that  $c > 0$  and consider the standard model of  $\mathbb{Q}_c^N$  as an umbilical hypersurface of Euclidean space  $\mathbb{R}^{N+1}$ . Let  $\mathbb{R}^{N+1} = \oplus_{i=0}^k \mathbb{R}^{m_i}$  be an orthogonal decomposition of  $\mathbb{R}^{N+1}$  with  $\mathbb{R}^{m_0}$  possibly a trivial subspace and  $\mathbb{R}^{m_i}$  nontrivial for  $1 \leq i \leq k$ .

For each  $q \in G = \mathbb{Q}_c^N \setminus \cup_{i=1}^k \mathbb{R}^{m_i \perp}$  and each  $1 \leq i \leq k$ , let  $\hat{S}_i(q)$  be the sphere in  $q + \mathbb{R}^{m_i}$  through  $q$  centered at the intersection of  $q + \mathbb{R}^{m_i}$  with  $\mathbb{R}^{m_i \perp}$ . Set  $\hat{E}_i(q) = T_q \hat{S}_i(q)$ ,  $1 \leq i \leq k$ . Let  $-a_i(q)$  be the mean curvature vector of  $\hat{S}_i(q)$  at  $q$  in  $\mathbb{R}^{N+1}$  and set  $\hat{E}_0(q) = \mathbb{Q}_c^N \cap (\mathbb{R}^{m_0} \oplus \text{span}\{a_i(q) : 1 \leq i \leq k\})$ . We say that  $\hat{\mathcal{E}}$  is the  $WP$ -net on  $G \subset \mathbb{Q}_c^N$  determined by the orthogonal decomposition  $\mathbb{R}^{N+1} = \oplus_{i=0}^k \mathbb{R}^{m_i}$ .

In either case, fix  $\bar{p} \in G$  and let  $N_i = \hat{S}_i(\bar{p})$ ,  $0 \leq i \leq k$ , be the leaf of  $\hat{E}_i$  through  $\bar{p}$ . Then  $(\bar{p}, N_1, \dots, N_k)$  satisfies the following conditions. If  $V_i = T_{\bar{p}} N_j$ ,  $0 \leq j \leq k$ , and  $z_i$  is the mean curvature vector of  $N_i$  in  $\mathbb{Q}_c^N$  at  $\bar{p}$  for  $1 \leq i \leq k$ , then  $T_{\bar{p}} \mathbb{Q}_c^N = \oplus_{i=0}^k V_i$  is an orthogonal decomposition of  $T_{\bar{p}} \mathbb{Q}_c^N$  into nontrivial subspaces,  $z_1, \dots, z_k \in V_0$  and  $\langle z_i, z_j \rangle = -c$  for  $i \neq j$ . The latter equality in the case  $c > 0$  follows from the fact that the mean curvature vectors  $-a_i = z_i - c\bar{p}$  of  $N_i$  in  $\mathbb{R}^{N+1}$  at  $\bar{p}$  are pairwise orthogonal.

Conversely, let  $(\bar{p}, N_1, \dots, N_k)$  satisfy the preceding conditions. Define  $\mathbb{R}^{m_i} = V_i \oplus \mathbb{R} a_i$  for  $1 \leq i \leq k$  and set  $\mathbb{R}^{m_0} = \cap_{i=1}^k \mathbb{R}^{m_i \perp}$ . In the case  $c = 0$ , assume without loss of generality that the mean curvature vector  $-a_i$  of  $N_i$  at  $\bar{p}$  vanishes if and only if  $1 \leq i \leq s$ , set  $\bar{q} = \bar{p} - \sum_{i=s+1}^k \|a_i\|^{-2} a_i$ ,  $\mathcal{H} = \bar{q} + \mathbb{R}^{m_0}$ , and let  $\hat{\mathcal{E}}$  be the  $WP$ -net determined by  $(\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m_i}, \mathcal{H}, s)$  on the open dense subset

$$G := \mathbb{R}^N \setminus (\bar{q} + \cup_{i=s+1}^k (\mathbb{R} a_i \oplus V_i)^\perp).$$

In the case  $c > 0$ , let  $\hat{\mathcal{E}}$  be the  $WP$ -net on the open dense subset

$$G := \mathbb{Q}_c^N \setminus \cup_{i=1}^k (\mathbb{R} a_i \oplus V_i)^\perp,$$

determined by the orthogonal decomposition  $\mathbb{R}^{N+1} = \oplus_{i=0}^k \mathbb{R}^{m_i}$ . Finally, for  $c = 0$  define

$$N_0 := \bar{q} + \{p \in V_0 : \langle a_i, p \rangle > 0 \text{ for all } s+1 \leq i \leq k\}$$

and  $\sigma_i: N_0 \rightarrow \mathbb{R}_+$  by  $\sigma_i(p) = \langle a_i, p - \bar{q} \rangle$  for  $1 \leq i \leq k$ . In the case  $c > 0$  define

$$N_0 := \mathbb{Q}_c^N \cap \{p \in \mathbb{R}\bar{p} \oplus V_0 : \langle a_i, p \rangle > 0 \text{ for all } 1 \leq i \leq k\}$$

and  $\sigma_i: N_0 \rightarrow \mathbb{R}_+$  by  $\sigma_i(p) = \langle a_i, p \rangle$  for  $1 \leq i \leq k$ . Note that in both cases  $N_0$  is an open subset of the totally geodesic submanifold of  $\mathbb{Q}_c^N$  that is tangent to  $V_0$  at  $\bar{p}$ . Then we have the following from Nölker's work.

**Proposition 18** *The map  $\Phi: N_0 \times_\sigma \prod_{i=1}^k N_i \rightarrow G \subset \mathbb{Q}_c^N$  given by*

$$\Phi(p_0, \dots, p_k) = p_0 + \sum_{i=1}^k \sigma_i(p_0)(p_i - \bar{p})$$

*is an isometric net isomorphism with respect to the product net of  $\prod_{i=0}^k N_i$  and  $\hat{\mathcal{E}} := \hat{\mathcal{E}}_\Phi$ .*

The map  $\Phi$  is called the warped product representation of  $\mathbb{Q}_c^N$  determined by  $(\bar{p}, N_1, \dots, N_k)$ . In the case  $c = 0$  (respectively,  $c > 0$ ), we say that  $\mathcal{H}$  (respectively,  $\mathbb{R}^{m_0}$ ) is the *axis* of  $\Phi$ . Clearly, if  $\hat{\mathcal{E}}$  is the *WP*-net determined by  $(\mathbb{R}^N = \oplus_{i=0}^k \mathbb{R}^{m_i}, \mathcal{H}, s)$  if  $c = 0$  or by the orthogonal decomposition  $\mathbb{R}^{N+1} = \oplus_{i=0}^k \mathbb{R}^{m_i}$  if  $c > 0$ , and  $N_i = \hat{S}_i(\bar{p})$  is the leaf of  $\hat{E}_i$  through some fixed point  $\bar{p} \in \mathbb{Q}_c^N$  for  $1 \leq i \leq k$ , then  $\hat{\mathcal{E}} = \hat{\mathcal{E}}_\Phi$  for the warped product representation  $\Phi$  of  $\mathbb{Q}_c^N$  determined by  $(\bar{p}, N_1, \dots, N_k)$ .

Given a warped product representation  $\Phi: N_0 \times_\sigma \prod_{i=1}^k N_i \rightarrow \mathbb{Q}_c^N$  and immersions  $f_i: M_i \rightarrow N_i$ ,  $0 \leq i \leq k$ , set  $\rho_i = \sigma_i \circ f_0$  for  $1 \leq i \leq k$ . Then

$$f := \Phi \circ (f_0 \times \dots \times f_k): M := \prod_{i=0}^k M_i \rightarrow \mathbb{Q}_c^N$$

is an immersion whose induced metric is the warped product of the metrics induced by  $f_0, \dots, f_k$  with warping function  $\rho = (\rho_1, \dots, \rho_k)$ . It is called the (extrinsic) warped product of  $f_0, \dots, f_k$ . If all functions  $\rho_1, \dots, \rho_k$  are identically one, then the metric induced by  $f$  is the Riemannian product of the metrics induced by  $f_0, \dots, f_k$ , and  $f$  is said to be the *extrinsic Riemannian product* of  $f_0, \dots, f_k$ .

An important special case is the following. Let  $\bar{p} \in \mathbb{Q}_c^N$ ,  $N \geq 2$ , let  $V_0, \dots, V_k$  be nontrivial mutually orthogonal vector subspaces of  $T_{\bar{p}}\mathbb{Q}_c^N$  and let  $z_0, \dots, z_k \in T_{\bar{p}}\mathbb{Q}_c^N$  be vectors that are orthogonal to  $\oplus_{i=0}^k V_i$  and satisfy  $\langle z_i, z_j \rangle = -c$  for  $i \neq j$ . For  $0 \leq i \leq k$  let  $N_i$  be the complete spherical submanifold of  $\mathbb{Q}_c^N$  such that  $T_{\bar{p}}N_i = V_i$  and whose mean curvature vector at  $\bar{p}$  is  $z_i$ . Then, the map  $\Psi: \prod_{i=0}^k N_i \rightarrow \mathbb{Q}_c^N$  given by

$$\Psi(p_0, \dots, p_k) = \bar{p} + \sum_{i=0}^k (p_i - \bar{p})$$

is an isometric embedding called the *extrinsic Riemannian product* of the complete spherical submanifolds  $N_0, \dots, N_k$ . This embedding can be described as an extrinsic Riemannian product of isometric immersions as follows. Let  $\Phi: \tilde{N}_0 \times_\sigma \prod_{i=1}^k N_i \rightarrow \mathbb{Q}_c^N$  be the warped product representation determined by  $(\bar{p}, N_1, \dots, N_k)$ . Then  $N_0 \subset \tilde{N}_0$  and the functions  $\sigma_1|_{N_0}, \dots, \sigma_k|_{N_0}$  are identically one. Hence  $\Psi$  is the extrinsic Riemannian product of  $i_0, \dots, i_k$  determined by  $\Phi$ , where  $i_0$  is the inclusion of  $N_0$  into  $\tilde{N}_0$  and  $i_j: N_j \rightarrow N_j$  is the identity map for  $1 \leq j \leq k$ .

## 6.2 Nölker's and Molzan's theorems

We now state Nölker's result and give a geometric proof of it in Euclidean space, from which the case of the sphere can be easily reduced. We say that the second fundamental form of an immersion  $f: M^n := \prod_{i=0}^k M_i \rightarrow \mathbb{Q}_c^N$  of a product manifold is *adapted to the product net*  $\mathcal{E} = (E_i)_{i=0, \dots, k}$  of  $M^n$  if

$$\alpha(X_i, X_j) = 0, \quad \text{for all } X_i \in \Gamma(E_i), X_j \in \Gamma(E_j), \quad i \neq j.$$

**Theorem 19 [No]** *Let  $f: M^n := M_0 \times_\rho \prod_{i=1}^k M_i \rightarrow \mathbb{Q}_c^N$  be an isometric immersion of a warped product whose second fundamental form is adapted to the product net of  $M^n$ . For a fixed point  $\bar{p} \in M^n$ , set  $f_i = f \circ \tau_i^{\bar{p}}: M_i \rightarrow \mathbb{Q}_c^N$  for  $0 \leq i \leq k$  and let  $N_i$  be the spherical hull of  $f_i$  for  $1 \leq i \leq k$ . Then  $f_0$  is an isometric immersion,  $f_i$  is a homothetical immersion for each  $1 \leq i \leq k$  with homothety factor  $\rho_i(\bar{p})$  (hence it is an isometric immersion if the metric of  $M$  is normalized with respect to  $\bar{p}$ ), and  $(f(\bar{p}); N_1, \dots, N_k)$  determines a warped product representation  $\Phi: N_0 \times_\sigma \prod_{i=1}^k N_i \rightarrow \mathbb{Q}_c^N$  such that  $f_0(M_0) \subset N_0$ ,  $\rho_i = \rho_i(\bar{p})(\sigma_i \circ f_0)$  and  $f = \Phi \circ (f_0 \times \dots \times f_k)$ , where  $f_i$  is regarded as a map into  $N_i$  for  $0 \leq i \leq k$ .*

*Proof:* By Proposition 3, the product net  $\mathcal{E} = (E_i)_{i=0, \dots, k}$  of  $M^n$  is a  $WP$ -net. Therefore, we obtain from Proposition 15 that for each  $p \in M^n$  and for each  $1 \leq i \leq k$  there exists an  $(N - n_i)$ -dimensional affine subspace  $H_i(p)$  that contains the image  $f(L_i^\perp(p))$  of the leaf  $L_i^\perp(p)$  of  $E_i^\perp$  through  $p$ , and whose normal space along  $f(L_i^\perp(p))$  is  $f_*E_i$ . We assume without loss of generality that there exists  $r \in \{1, \dots, k\}$  such that the warping function  $\rho_i$  is constant if and only if  $1 \leq i \leq r$ . Then, we also have from Proposition 15 that for each  $p \in M^n$  and for each  $1 \leq i \leq r$  (respectively,  $r + 1 \leq i \leq k$ ), there exists an  $(N - n + n_i)$ -dimensional affine subspace (respectively, sphere or affine subspace)  $\tilde{S}_i(p)$  that contains the image  $f(L_i(p))$  of the leaf  $L_i(p)$  of  $E_i$  through  $p$ , and whose normal space along  $f(L_i(p))$  is  $f_*E_i^\perp$ .

If  $k = 1$ , then the families  $\mathcal{F}_1 = (H_1(p))_{p \in M}$  and  $\mathcal{F}_0 = (\tilde{S}_1(p))_{p \in M}$  satisfy the assumptions of Lemma 14 with  $\Lambda_i = M_i$ . For  $k \geq 2$ , it follows that the families  $\mathcal{F}_i = (H_i(p))_{p \in M}$ ,  $1 \leq i \leq k$ , satisfy the assumptions of Lemma 13 with  $\Lambda_i = M_i$ . In any case, after reordering the families if necessary if  $k \geq 2$ , there exist  $s \in \{0, \dots, r\}$ , an orthogonal decomposition  $\mathbb{R}^N = \bigoplus_{i=0}^k \mathbb{R}^{m_i}$  with  $m_i \geq n_i$  for  $1 \leq i \leq s$  and  $m_i \geq n_i + 1$  for  $s + 1 \leq i \leq k$ , and affine  $(N - m_i)$ -dimensional subspaces  $\mathcal{H}_i$  parallel to  $\mathbb{R}^{m_i^\perp}$ ,  $s + 1 \leq i \leq k$ , such that, for  $1 \leq i \leq s$  (respectively,  $s + 1 \leq i \leq k$ ),  $\mathcal{F}_i$  is a light-like (respectively, space-like) family of affine subspaces with axis  $\mathcal{H}_{\mathcal{F}_i} = \mathbb{R}^{m_i^\perp}$  (respectively,  $\mathcal{H}_{\mathcal{F}_i} = \mathcal{H}_i$ ).

Let  $\hat{\mathcal{E}} = (\hat{E}_i)_{i=0, \dots, k}$  be the  $WP$ -net on the open dense subset  $G = \mathbb{R}^N / \cup_{i=s+1}^k \mathcal{H}_i$  determined by  $(\mathbb{R}^N = \bigoplus_{i=0}^k \mathbb{R}^{m_i}, \mathcal{H} = \cap_{i=s+1}^k \mathcal{H}_i, s)$ . We claim that  $f$  is a net morphism with respect to the product net  $\mathcal{E} = (E_i)_{i=0, \dots, k}$  of  $M$  and  $\hat{\mathcal{E}}$ . Given a leaf  $L_i(p)$  of  $E_i$  through  $p \in M$ , we have that the affine normal space of  $f|_{L_i(p)}$  at any  $q \in L_i(p)$  is precisely  $H_i(q)$ . It follows from Proposition 9 that the spherical hull of  $f(L_i(p))$  is the leaf  $\hat{S}_i(f(p))$  of  $\hat{E}_i$  through  $f(p)$  for  $1 \leq i \leq k$ . Thus, in order to prove the claim it remains to prove that  $f$  takes each leaf  $L_0$  of  $E_0$  into a leaf of  $\hat{E}_0$ , or equivalently, that  $f_*E_0(p) \subset \hat{E}_0(f(p))$  for every  $p \in M$ . Notice that, for  $s + 1 \leq i \leq k$ , the normal space at  $p$  of  $f|_{L_i^\perp(p)}$  is  $T_{f(p)}\tilde{S}_i(p)$ , that is,  $T_{f(p)}\tilde{S}_i(p) = (f_*E_i^\perp(p))^\perp$ . Since  $\tilde{S}_i(p)$  contains  $f(L_i(p))$ , it also contains the spherical hull  $\hat{S}_i(f(p))$  of  $f(L_i(p))$ . Thus

$$T_{f(p)}\hat{S}_i(f(p)) \subset T_{f(p)}\tilde{S}_i(p) = (f_*E_i^\perp(p))^\perp,$$

and hence

$$f_*E_i^\perp(p) \subset T_{f(p)}^\perp \hat{S}_i(f(p)) = \mathbb{R}^{m_i^\perp} \oplus \text{span}\{a_i(f(p))\}.$$

Using that  $E_0(p) = \cap_{i=1}^k E_i^\perp(p)$ , we conclude that

$$f_*E_0(p) \subset \mathbb{R}^{m_0} \oplus \text{span}\{a_i(f(p)) : s+1 \leq i \leq k\} = \hat{E}_0(f(p)).$$

For a fixed  $\bar{p} \in M$ , the map  $\tau_0^{\bar{p}}: M_0 \rightarrow M$  is an isometric immersion and  $\tau_i^{\bar{p}}: M_i \rightarrow M$  is a homothetical immersion with homothety factor  $\rho_i(\bar{p})$  for every  $1 \leq i \leq k$ . Hence the same holds for  $f_0 = f \circ \tau_0^{\bar{p}}: M_0 \rightarrow \mathbb{R}^N$  and  $f_i = f \circ \tau_i^{\bar{p}}: M_i \rightarrow \mathbb{R}^N$ ,  $1 \leq i \leq k$ . Moreover, the spherical hull  $N_i$  of  $f_i$  is the leaf of  $\hat{\mathcal{E}}$  through  $f(\bar{p})$  for  $1 \leq i \leq k$ , thus  $(f(\bar{p}); N_1, \dots, N_k)$  determines a warped product representation  $\Phi: N_0 \times_\sigma \prod_{i=1}^k N_i \rightarrow G$ , with  $f_0(M_0) \subset N_0$ , which is an isometric net morphism with respect to the product net of  $\prod_{i=0}^k N_i$  and  $\hat{\mathcal{E}}$ . Since  $f$  is a net morphism with respect to the product net of  $M$  and  $\hat{\mathcal{E}}$ , it follows that  $\Phi^{-1} \circ f$  is also a net morphism with respect to the product nets of  $M$  and  $\prod_{i=0}^k N_i$ . We conclude from Lemma 2 that  $\Phi^{-1} \circ f = f_0 \times \dots \times f_k$ . ■

**Remark 20.** It follows from the proof that, for  $c = 0$  (resp.,  $c > 0$ ), the axis of the warped product representation  $\Phi$  is given by  $\mathcal{H} = \cap_{i \in A} \mathcal{H}_i$  (resp.,  $\mathcal{H} = \mathbb{R}^{m_0} = \cap_{i=1}^k \mathbb{R}^{m_i^\perp}$ ) where  $A$  is the subset of indexes  $i \in \{0, \dots, k\}$  for which  $N_i \subset \mathbb{R}^N$  is a sphere and  $\mathcal{H}_i$  (resp.,  $\mathbb{R}^{m_i^\perp}$ ) is the axis of the space-like family of affine normal spaces of  $f_i$ .

For later use we state the following consequence of Nölker's theorem, which is an extension to nonflat ambient space forms due to Molzan of Moore's decomposition theorem of isometric immersions of Riemannian products into Euclidean space.

**Theorem 21 [MI]** *Let  $f: M^n := \prod_{i=0}^k M_i \rightarrow \mathbb{Q}_c^N$  be an isometric immersion of a Riemannian product whose second fundamental form is adapted to the product net of  $M^n$ . For a fixed  $\bar{p} \in M$ , set  $f_i = f \circ \tau_i^{\bar{p}}: M_i \rightarrow \mathbb{Q}_c^N$ , and let  $N_i$  be the spherical hull of  $f_i$ . Then  $f_i$  is an isometric immersion for each  $0 \leq i \leq k$  and there exists an isometric embedding  $\Psi: \prod_{i=0}^k N_i \rightarrow \mathbb{Q}_c^N$  as an extrinsic Riemannian product such that  $f = \Psi \circ (f_0 \times \dots \times f_k)$ , where  $f_i$  is regarded as a map into  $N_i$  for  $0 \leq i \leq k$ .*

## 7 Conformal immersions of Riemannian products

We are now in a position to consider the main theme of this article.

**Definition 22.** An isometric immersion  $f: M^n \rightarrow \mathbb{R}^{n+p}$  is a *k-isothermic submanifold*, or simply an *isothermic submanifold*, if  $M^n$  carries a *k-CP-net* with respect to which the second fundamental form of  $f$  is adapted.



In this section we prove our main result, namely, we obtain a local classification of all isothermic submanifolds of dimension  $n \geq 3$  of Euclidean space. Equivalently, in view of Theorem 7, we give a complete description of all conformal immersions of a Riemannian product of dimension  $n \geq 3$  into Euclidean space whose second fundamental forms are adapted to the product net of the manifold. We follow with several applications of this result. First we prove two basic lemmas.

**Lemma 23** *Let  $\Phi: N_0^m \times_\sigma N_1^{N-m} \rightarrow \mathbb{R}^N \setminus \mathcal{H}^{m-1}$  be a warped product representation. Then there exist conformal diffeomorphisms  $\Theta_\Phi: \mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N \setminus \mathcal{H}^{m-1}$  and  $i_0: N_0^m \rightarrow \mathbb{H}^m(-c)$  with conformal factors  $\sigma \circ \pi_0$  and  $\sigma$ , respectively, and an isometry  $i_1: N_1^{N-m} \rightarrow \mathbb{S}^{N-m}(c)$  such that  $\Phi = \Theta_\Phi \circ (i_0 \times i_1)$ .*

$$\begin{array}{ccc} N_0^m \times_\sigma N_1^{N-m} & \xrightarrow{\Phi} & \mathbb{R}^N \\ i_0 \downarrow & i_1 \downarrow & \nearrow \Theta_\Phi \\ \mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c) & & \end{array}$$

*Proof:* We assume for simplicity that  $\Phi: N_0^m \times_\sigma N_1^{N-m} \rightarrow \mathbb{R}^N$  is the warped product representation determined by  $(\bar{p}, N_1)$ , where  $\bar{p}_i = c^{-1/2}\delta_{mi}$ ,  $c > 0$ ,  $N_1^{N-m} = \mathbb{S}^{N-m}(c)$  is the sphere through  $\bar{p}$  whose tangent space at  $\bar{p}$  is  $\mathbb{R}^{m\perp}$  and whose mean curvature vector  $-a$  at  $\bar{p}$  is given by  $a_i = c^{1/2}\delta_{mi}$ , and  $N_0^m = \mathbb{R}_+^m := \{X = (x_1, \dots, x_m) \in \mathbb{R}^m : x_m > 0\}$ . Then  $\sigma: \mathbb{R}_+^m \rightarrow \mathbb{R}$  is given by  $\sigma(x) = c^{1/2}x_m$ , and  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N \setminus \mathbb{R}^{m-1}$ , where  $\mathbb{R}^{m-1} = \{X = (x_1, \dots, x_m) \in \mathbb{R}^m : x_m = 0\}$ , is the isometry

$$(X, Y) \mapsto (x_1, \dots, x_{m-1}, c^{1/2}x_m Y), \quad \text{for } X \in \mathbb{R}_+^m \text{ and } Y \in \mathbb{S}^{N-m}(c). \quad (12)$$

Endowing  $\mathbb{R}_+^m$  with the metric  $\sigma^{-2}\langle \cdot, \cdot \rangle$ , where  $\langle \cdot, \cdot \rangle$  denotes the Euclidean metric, we obtain the half-space model of  $\mathbb{H}^m(-c)$ , and the metric induced by (12) is conformal to the product metric of  $\mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c)$  with conformal factor  $\sigma \circ \pi_0$ . ■

**Lemma 24** *Let  $\langle \cdot, \cdot \rangle^\sim = \sum_{i=0}^k \varphi_i^2 \pi_i^* \langle \cdot, \cdot \rangle_i^\sim$  and  $\langle \cdot, \cdot \rangle = \sum_{i=0}^k \rho_i^2 \pi_i^* \langle \cdot, \cdot \rangle_i$  be twisted product metrics on a product manifold  $M^n := \prod_{i=0}^k M_i^{n_i}$ . If  $\langle \cdot, \cdot \rangle^\sim = \psi^2 \langle \cdot, \cdot \rangle$  for some  $\psi \in C^\infty(M)$ , then there exist  $\psi_i \in C^\infty(M_i)$  such that  $\psi \rho_i / \varphi_i = \psi_i \circ \pi_i$  and  $\langle \cdot, \cdot \rangle_i^\sim = \psi_i^2 \langle \cdot, \cdot \rangle_i$  for  $0 \leq i \leq k$ .*

*Proof:* Compute both sides of the equality  $\langle \cdot, \cdot \rangle^\sim = \psi^2 \langle \cdot, \cdot \rangle$  for the lift  $\tilde{X}_i$  to  $M$  of a local unit vector field  $X_i \in \Gamma(TM_i)$ . ■

Given a product manifold  $M = \prod_{i=0}^k M_i$ , we denote by  $T_j: M \rightarrow M_j \times \prod_{i \neq j} M_i$  the map given by  $T_j(p_0, \dots, p_k) = (p_j, (p_0, \dots, \hat{p}_j, \dots, p_k))$ , where the hat indicates a missing element.

**Theorem 25** Let  $f: M^n := \Pi_{i=0}^k M_i^{n_i} \rightarrow \mathbb{R}^N$ ,  $n \geq 3$ , be a conformal immersion of a Riemannian product whose second fundamental form is adapted to the product net of  $M^n$ . Then one of the following holds:

- (i) There exist an isometric embedding  $\Psi: \Pi_{i=0}^k N_i \rightarrow \mathbb{Q}_c^N \subset \mathbb{R}^{N+\epsilon}$  of a product of complete spherical submanifolds of  $\mathbb{Q}_c^N$  as an extrinsic product, where  $c \geq 0$ ,  $\epsilon = 1$  if  $c > 0$  and  $\epsilon = 0$  if  $c = 0$ , and isometric immersions  $f_i: M_i \rightarrow N_i$ ,  $0 \leq i \leq k$ , such that

$$f = \mathcal{P} \circ H \circ \Psi \circ (f_0 \times \cdots \times f_k),$$

where  $H$  is a homothety in  $\mathbb{R}^{N+\epsilon}$  and  $\mathcal{P}$  is either a stereographic projection of  $H(\mathbb{Q}_c^N)$  onto  $\mathbb{R}^N$  if  $c > 0$  or an inversion in  $\mathbb{R}^N$  otherwise.

$$\begin{array}{ccccc} M_0 \times \cdots \times M_k & \xrightarrow{f} & \mathbb{R}^N & & \\ f_0 \downarrow & & & \uparrow \mathcal{P} & \\ N_0 \times \cdots \times N_k & \xrightarrow{\Psi} & \mathbb{Q}_c^N \subset \mathbb{R}^{N+\epsilon} & \xrightarrow{H} & H(\mathbb{Q}_c^N) \subset \mathbb{R}^{N+\epsilon} \end{array}$$

- (ii)  $k \leq N - n + 1$  and for exactly one  $j \in \{0, \dots, k\}$  there exist an isometric embedding  $\Psi: \Pi_{i=0, i \neq j}^k \mathbb{S}^{m_i}(c_i) \rightarrow \mathbb{S}^{N-m}(c)$  of a product of complete spherical submanifolds of  $\mathbb{S}^{N-m}(c)$  as an extrinsic product, isometric immersions  $f_j: M_j \rightarrow \mathbb{H}^m(-c)$  and  $f_i: M_i \rightarrow \mathbb{S}^{m_i}(c_i)$ ,  $i \neq j$ , and an inversion  $I$  in  $\mathbb{R}^N$  such that

$$f = I \circ \Theta \circ (f_j \times (\Psi \circ (f_0 \times \cdots \times \hat{f}_j \times \cdots \times f_k)) \circ T_j).$$

$$\begin{array}{ccccc} M_0 \times M_1 \times \cdots \times M_k & \xrightarrow{f} & \mathbb{R}^N & & \\ f_0 \downarrow & f_1 \downarrow & & & \uparrow I \\ & \mathbb{S}_1(c_1) \times \cdots \times \mathbb{S}_k(c_k) & & & \\ & \downarrow \Psi & & & \\ \mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c) & \xrightarrow{\Theta} & \mathbb{R}^N & & \end{array}$$

Case (ii)  
for  $j = 0$

*Proof:* By Proposition 6, the product net  $\mathcal{E} = (E_i)_{0, \dots, k}$  is a  $CP$ -net with respect to the metric  $\langle \cdot, \cdot \rangle^*$  induced by  $f$ . If  $k \geq 2$  then  $E_i^\perp$  has dimension at least 2 for every  $0 \leq i \leq k$ . It follows from Proposition 16 that  $E_i^\perp$  is spherical, and hence the same holds for  $E_i$  by Proposition 5-(ii). If  $k = 1$  then either  $E_0$  or  $E_1$  has dimension at least 2 by our assumption that  $n \geq 3$ , and again Proposition 16 together with Proposition 5-(ii) imply that both  $E_0$  and  $E_1$  are spherical distributions. Thus, in any case  $E_i$  and  $E_i^\perp$  are spherical for every  $0 \leq i \leq k$ .

By Proposition 15, for each  $0 \leq i \leq k$  and for each  $p \in M^n$  there exists an  $(N - n_i)$ -dimensional sphere or affine subspace  $S_i(p)$  that contains the image  $f(L_i^\perp(p))$  by  $f$  of the leaf of  $E_i^\perp$  through  $p$  and whose normal space along  $f(L_i^\perp(p))$  is  $f_*E_i$ . Therefore, the families  $\mathcal{F}_i = (S_i(p))_{p \in M}$ ,  $0 \leq i \leq k$ , satisfy the assumptions of Lemma 11 with  $\Lambda_i = M_i$ .

Let us assume first that the statement in either part (a) or (b) of Lemma 11 holds. Denote by  $\mathcal{P}$  either an inversion in  $\mathbb{R}^N$  or a stereographic projection of a punctured sphere  $\mathbb{S}^N(\tilde{c}) \setminus \{z\} \subset \mathbb{R}^{N+1}$  onto  $\mathbb{R}^N$  as in part (a) or (b) of Lemma 11, respectively. Set  $g = \mathcal{P}^{-1} \circ f$ . In case (a) (resp., in case (b)), we have that  $\mathcal{P}^{-1}(S_i(p))$  is an affine subspace in  $\mathbb{R}^N$  (resp., a great sphere in  $\mathbb{S}^N(\tilde{c})$ ) that contains  $g(L_i^\perp(p))$  and whose normal space along  $g(L_i^\perp(p))$  is  $g_*E_i$ ,  $0 \leq i \leq k$ . It follows from Proposition 15 that  $L_i^\perp(p)$  is totally geodesic with respect to the metric  $\langle \cdot, \cdot \rangle^\sim$  induced by  $g$ , and hence  $E_i^\perp$  is a totally geodesic distribution with respect to  $\langle \cdot, \cdot \rangle^\sim$ . Therefore  $\langle \cdot, \cdot \rangle^\sim = \sum_{i=0}^k \pi_i^* \langle \cdot, \cdot \rangle_i^\sim$  is a Riemannian product metric on  $M$ , and thus Molzan's theorem can be applied to  $g: (M^n, \langle \cdot, \cdot \rangle^\sim) \rightarrow \mathbb{Q}_{\tilde{c}}^N$ , where  $\tilde{c} = 0$  in case (a) and  $\tilde{c} > 0$  in case (b). For a fixed  $\bar{p} \in M$ , let  $f_i = g \circ \tau_i^{\bar{p}}: M_i \rightarrow \mathbb{Q}_{\tilde{c}}^N$  and let  $N_i$  be the spherical hull of  $f_i$ . Then  $f_i: (M_i, \langle \cdot, \cdot \rangle_i^\sim) \rightarrow \mathbb{Q}_{\tilde{c}}^N$  is an isometric immersion for every  $0 \leq i \leq k$  and there exists an isometric embedding  $\Phi: \prod_{i=0}^k N_i \rightarrow \mathbb{Q}_{\tilde{c}}^N$  as an extrinsic Riemannian product such that  $g = \Phi \circ (f_0 \times \cdots \times f_k)$ .

Since  $\langle \cdot, \cdot \rangle^\sim$  is conformal to the Riemannian product metric  $\langle \cdot, \cdot \rangle = \sum_{i=0}^k \pi_i^* \langle \cdot, \cdot \rangle_i$  of  $M$ , it follows from Lemma 24 that they must differ by a constant factor. Hence, there exists a homothety  $H$  in  $\mathbb{R}^{N+\epsilon}$ , where  $\epsilon = 0$  in case (a) and  $\epsilon = 1$  in case (b), such that  $H^{-1} \circ g: (M, \langle \cdot, \cdot \rangle) \rightarrow \mathbb{R}^{N+\epsilon}$  is an isometric immersion. Setting  $H^{-1}(\mathbb{S}^N(\tilde{c})) = \mathbb{S}^N(c)$  in case (b), we conclude that the statement in part (i) holds.

Now assume that the alternative (c) in Lemma 11 holds. Arguing as in the preceding case, we obtain that  $E_j^\perp$  is spherical and that  $E_i^\perp$  is totally geodesic for  $0 \leq i \neq j \leq k$  with respect to the metric  $\langle \cdot, \cdot \rangle^\sim$  induced by  $g = I \circ f$ . Thus, also  $E_j = \cap_{i \neq j} E_i^\perp$  is totally geodesic. By Proposition 3,  $\langle \cdot, \cdot \rangle^\sim$  is a warped-product metric

$$\langle \cdot, \cdot \rangle^\sim = \pi_j^* \langle \cdot, \cdot \rangle_j^\sim + (\rho \circ \pi_j)^2 \pi_{M_j^\perp}^* \langle \cdot, \cdot \rangle_{\perp_j}^\sim$$

on  $M$  regarded as a product  $M = M_j \times M_j^\perp$ , where  $M_j^\perp := M_0 \times \cdots \times \hat{M}_j \times \cdots \times M_k$ , which we can assume to be normalized with respect to some point  $\bar{p} \in M$ . Denote  $\tilde{g}_j = g \circ \tau_j^{\bar{p}}: M_j \rightarrow \mathbb{R}^N$  and  $\bar{g}_j = g \circ \tau_{\perp_j}^{\bar{p}}: M_j^\perp \rightarrow \mathbb{R}^N$ , where  $\tau_{\perp_j}^{\bar{p}}: M_j^\perp \rightarrow M$  is the canonical inclusion of  $M_j^\perp$  into  $M$  as the leaf of  $E_j^\perp$  through  $\bar{p}$ . Then  $\tilde{g}_j: (M_j, \langle \cdot, \cdot \rangle_j^\sim) \rightarrow \mathbb{R}^N$  and  $\bar{g}_j: (M_j^\perp, \langle \cdot, \cdot \rangle_{\perp_j}^\sim) \rightarrow \mathbb{R}^N$  are isometric immersions. Moreover, denoting by  $\tilde{N}_j$  the spherical hull of  $\tilde{g}_j$ , it follows from Theorem 19 that  $(\bar{p}, \tilde{N}_j)$  determines a warped product representation  $\Phi: \tilde{N}_j \times_\sigma \tilde{N}_j \rightarrow \mathbb{R}^N$  such that  $\tilde{g}_j(M_j) \subset \tilde{N}_j$ ,  $\rho = \sigma \circ \tilde{g}_j$  and  $g = \Phi(\tilde{g}_j \times \bar{g}_j)$ . Since  $\langle \cdot, \cdot \rangle^\sim$  is conformal to the product metric  $\langle \cdot, \cdot \rangle = \pi_j^* \langle \cdot, \cdot \rangle_j + \pi_{M_j^\perp}^* \langle \cdot, \cdot \rangle_{\perp_j}$ , where  $\langle \cdot, \cdot \rangle_{\perp_j} = \sum_{i=0, i \neq j}^k \pi_i^* \langle \cdot, \cdot \rangle_i$ , it follows from Lemma 24 that there exists a constant  $a > 0$  such that  $\langle \cdot, \cdot \rangle_j^\sim = a^2(\sigma \circ \tilde{g}_j)^2 \langle \cdot, \cdot \rangle_j$  and  $\langle \cdot, \cdot \rangle_{\perp_j}^\sim = a^2 \langle \cdot, \cdot \rangle_{\perp_j}$ . By renormalizing the

metric  $\langle \cdot, \cdot \rangle^\sim$ , we may assume that  $a = 1$  and hence  $\bar{g}_j: (M_j^\perp, \langle \cdot, \cdot \rangle_{\perp_j}) \rightarrow \bar{N}_j$  is an isometric immersion and  $\tilde{g}_j: (M_j, \langle \cdot, \cdot \rangle_j) \rightarrow \tilde{N}_j$  is a conformal immersion with conformal factor  $(\sigma \circ \tilde{g}_j)$ , where we now regard  $\bar{g}_j$  and  $\tilde{g}_j$  as maps into  $\bar{N}_j$  and  $\tilde{N}_j$ , respectively. By Lemma 23, there exist conformal diffeomorphisms  $\Theta_\Phi: \mathbb{H}^m(-c) \times \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N \setminus \mathcal{H}^{m-1}$  and  $i_0: \tilde{N}_j \rightarrow \mathbb{H}^m(-c)$  with conformal factors  $\sigma \circ \pi_0$  and  $\sigma$ , respectively, and an isometry  $i_1: \bar{N}_j \rightarrow \mathbb{S}^{N-m}(c)$  such that  $\Phi = \Theta_\Phi \circ (i_0 \times i_1)$ . Set  $f_j = i_0 \circ \tilde{g}_j$  and  $\hat{g}_j = i_1 \circ \bar{g}_j$ . Then  $f_j: (M_j, \langle \cdot, \cdot \rangle_j) \rightarrow \mathbb{H}^m(-c)$  and  $\hat{g}_j: (M_j^\perp, \langle \cdot, \cdot \rangle_{\perp_j}) \rightarrow \mathbb{S}^{N-m}(c)$  are isometric immersions and

$$g = \Theta \circ (f_j \times \hat{g}_j). \quad (13)$$

Finally, we apply Molzan's theorem to  $\hat{g}_j: (M_j^\perp, \langle \cdot, \cdot \rangle_{\perp_j}) \rightarrow \mathbb{S}^{N-m}(c)$ . We obtain that  $f_i = g \circ \tau_i^{\bar{p}} = \hat{g}_j \circ \tau_i^{\bar{p}}: M_i \rightarrow \mathbb{S}^{N-m}(c)$  is an isometric immersion for  $0 \leq i \neq j \leq k$  and, denoting by  $\mathbb{S}^{m_i}(c_i)$  the spherical hull of  $f_i$ , there exists an isometric embedding  $\Psi: \prod_{i=0, i \neq j}^k \mathbb{S}^{m_i}(c_i) \rightarrow \mathbb{S}^{N-m}(c)$  as an extrinsic product such that

$$\hat{g}_j = \Psi \circ (f_0 \times \cdots \times \hat{f}_j \times \cdots \times f_k), \quad (14)$$

We conclude from (13) and (14) that the statement in part (ii) holds. ■

**Remarks 26.** (a) The proof shows that part (ii) could be stated in the following alternative way: (ii')  $k \leq N - n + 1$  and for exactly one  $j \in \{0, \dots, k\}$  there exist a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N$ , an isometric embedding  $\Psi: \prod_{i=0, i \neq j}^k \mathbb{S}^{m_i}(c_i) \rightarrow \mathbb{S}^{N-m}(c)$  of a product of complete spherical submanifolds of  $\mathbb{S}^{N-m}(c)$  as an extrinsic product, a conformal immersion  $f_j: M_j \rightarrow \mathbb{R}_+^m$  with conformal factor  $(\sigma \circ f_j)$ , isometric immersions  $f_i: M_i \rightarrow \mathbb{S}^{m_i}(c_i)$ ,  $i \neq j$ , and an inversion  $I$  in  $\mathbb{R}^N$  such that

$$f = I \circ \Phi \circ (f_j \times (\Psi \circ (f_0 \times \cdots \times \hat{f}_j \times \cdots \times f_k))) \circ T_j.$$

(b) A conformal immersion  $f: M^n := \prod_{i=0}^k M_i^{n_i} \rightarrow \mathbb{R}^N$  given as in part (i) of the statement with  $c > 0$  can occur only if  $k \leq N - n$ . This follows from the fact that this is the case in which the families  $\mathcal{F}_i$ ,  $0 \leq i \leq k$ , in the proof are as in Lemma 11-(b). Moreover, it follows from Remark 12 and the proof of part (ii) that in this case also the description in that part holds for any value of  $j \in \{0, \dots, k\}$ .

(c) If  $k = 1$ , that is,  $M^n := M_0^{n_0} \times M_1^{n_1}$  is a Riemannian product with only two factors, then part (ii) of the statement simplifies considerably:

(ii) For some  $j \in \{0, 1\}$  there exist isometric immersions  $f_j: M_j \rightarrow \mathbb{H}^m(-c)$  and  $f_i: M_i \rightarrow \mathbb{S}^{N-m}(c)$ ,  $0 \leq i \neq j \leq 1$ , and an inversion  $I$  such that  $f = I \circ \Theta \circ (f_j \times f_i) \circ T_j$ .

Alternatively, in view of the first remark:

(ii') there exist  $j \in \{0, 1\}$ , a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m}(c) \rightarrow \mathbb{R}^N$ , an inversion  $I$  in  $\mathbb{R}^N$ , a conformal immersion  $f_j: M_j \rightarrow \mathbb{R}_+^m$  with conformal factor  $(\sigma \circ f_j)$ , and an isometric immersion  $f_i: M_i \rightarrow \mathbb{S}^{N-m}(c)$ ,  $0 \leq i \neq j \leq 1$ , such that  $f = I \circ \Phi \circ (f_j \times f_i) \circ T_j$ .

The assumption that  $n \geq 3$  in Theorem 25 was used only in the first paragraph of the proof in order to insure that the product net  $\mathcal{E} = (E_i)_{0,\dots,k}$  of  $M$  has the property that  $E_i$  and  $E_i^\perp$  are spherical for every  $0 \leq i \leq k$ . For a surface  $f: M^2 \rightarrow \mathbb{R}^N$ , the existence of an orthogonal net  $\mathcal{E} = (E_0, E_1)$  with both  $E_0$  and  $E_1$  spherical and such that the second fundamental form of  $f$  is adapted to  $\mathcal{E}$  is equivalent to the assumptions that the surface has flat normal bundle and that the curvature lines have constant geodesic curvature. Therefore, we obtain the following in the surface case.

**Corollary 27** *Let  $f: M^2 \rightarrow \mathbb{R}^N$  be a surface with flat normal bundle without umbilic points. Assume that the curvature lines of both families have constant geodesic curvature. Then one of the following holds:*

- (i) *There exist local isothermic parameterizations by curvature lines  $\psi: J_1 \times J_2 \rightarrow M^2$ , an isometry  $\Phi: \mathbb{R}^{N_1} \times \mathbb{R}^{N_2} \rightarrow \mathbb{R}^N$ , an inversion  $I$  in  $\mathbb{R}^N$  and unit speed curves  $\alpha: J_1 \rightarrow \mathbb{R}^{N_1}$  and  $\beta: J_2 \rightarrow \mathbb{R}^{N_2}$  such that  $f \circ \psi = I \circ \Phi \circ (\alpha \times \beta)$ .*
- (ii) *There exist local isothermic parameterizations by curvature lines  $\psi: J_1 \times J_2 \rightarrow M^2$ , unit speed curves  $\alpha: J_1 \rightarrow \mathbb{H}^m(-c)$  and  $\beta: J_2 \rightarrow \mathbb{S}^{N-m}(c)$ , and an inversion  $I$  in  $\mathbb{R}^N$  such that  $f \circ \psi = I \circ \Theta \circ (\alpha \times \beta)$ .*

In the case  $N = 3$ , this is a classical result of Bonnet ([Bo]). In fact, the geometric proof of Bonnet's theorem given later by Ribaucour ([Ri], cf. [Da<sub>2</sub>], vol. III, p. 121) served as an inspiration for the proof of Theorem 25.

We now derive several consequences of Theorem 25. First, we obtain a classification of all *conformal representations* of Euclidean space as a Riemannian product, which contains as a special case Liouville's theorem on the classification of conformal mappings between open subsets of  $\mathbb{R}^N$ ,  $N \geq 3$ . It also provides a geometric proof of the classification of conformally flat Riemannian products (cf. [La], Section D, Proposition 2).

**Corollary 28** *Let  $f: M^N := \Pi_{i=0}^k M_i^{N_i} \rightarrow \mathbb{R}^N$ ,  $N \geq 3$ , be a conformal local diffeomorphism of a Riemannian product. Then one of the following holds:*

- (i) *There exist an isometry  $\Phi: \Pi_{i=0}^k \mathbb{R}^{N_i} \rightarrow \mathbb{R}^N$ , local isometries  $f_i: M_i^{N_i} \rightarrow \mathbb{R}^{N_i}$ , an inversion  $I$  and a homothety  $H$  in  $\mathbb{R}^N$  such that*

$$f = I \circ H \circ \Phi \circ (f_0 \times \cdots \times f_k).$$

- (ii)  *$k = 1$  and for some  $j \in \{0, 1\}$  there exist local isometries  $f_j: M_j^{N_j} \rightarrow \mathbb{H}^{N_j}(-c)$  and  $f_i: M_i^{N_i} \rightarrow \mathbb{S}^{N_i}(c)$ ,  $0 \leq i \neq j \leq 1$ , and an inversion  $I$  in  $\mathbb{R}^N$  such that*

$$f = I \circ \Theta \circ (f_j \times f_i) \circ T_j.$$

Corollary 28 yields a classification of all *CP*-nets on open subsets of Euclidean space.

**Corollary 29** *Let  $\mathcal{E} = (E_i)_{0,\dots,k}$  be a CP-net on an open subset  $U \subset \mathbb{R}^N$ . Then one of the following holds:*

- (i) *There exists an inversion  $I$  in  $\mathbb{R}^N$  such that  $\hat{\mathcal{E}} = (I_*E_i)_{0,\dots,k}$  is the restriction to  $U$  of the RP-net determined by some orthogonal decomposition  $\mathbb{R}^N = \bigoplus_{i=0}^k \mathbb{R}^{m_i}$ .*
- (ii)  *$k = 1$  and there exist an inversion  $I$  in  $\mathbb{R}^N$ , an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^m \oplus \mathbb{R}^{N-m}$ ,  $0 \leq m \leq N-1$ , and an affine subspace  $\mathcal{H}^m \subset \mathbb{R}^N$  parallel to  $\mathbb{R}^m$  (which reduces to a point  $\bar{q}$  if  $m = 0$ ), such that  $\hat{\mathcal{E}} = (I_*E_0, I_*E_1)$  is the restriction to  $U$  of the WP-net on  $\mathbb{R}^N \setminus \mathcal{H}^m$  determined by  $(\mathbb{R}^N = \mathbb{R}^m \oplus \mathbb{R}^{N-m}, \mathcal{H}^m, s = 0)$ .*

The remaining applications deal with products with two factors, in which case one should have in mind Remark 26-(c). The next corollary follows by using the observation that any warped product metric  $\langle \cdot, \cdot \rangle = \pi_0^* \langle \cdot, \cdot \rangle_0 + (\rho \circ \pi_0)^2 \pi_1^* \langle \cdot, \cdot \rangle_1$  is conformal to the Riemannian product metric  $\langle \cdot, \cdot \rangle^\sim = \pi_0^* \rho^{-2} \langle \cdot, \cdot \rangle_0 + \pi_1^* \langle \cdot, \cdot \rangle_1$  with conformal factor  $\rho \circ \pi_0$ .

**Corollary 30** *Let  $f: M^n := M_0^{n_0} \times_\rho M_1^{n_1} \rightarrow \mathbb{R}^N$ ,  $n \geq 3$ , be a conformal immersion of a warped product whose second fundamental form is adapted to the product net of  $M^n$ . Then the conclusion of Theorem 25 holds for  $f: (M^n, \langle \cdot, \cdot \rangle^\sim) \rightarrow \mathbb{R}^N$ , where  $\langle \cdot, \cdot \rangle^\sim$  is the product metric  $\langle \cdot, \cdot \rangle^\sim = \pi_0^* \rho^{-2} \langle \cdot, \cdot \rangle_0 + \pi_1^* \langle \cdot, \cdot \rangle_1$ .*

In particular, we obtain a classification of conformal representations of open subsets of Euclidean space as a warped product with two factors.

**Corollary 31** *Let  $f: M^N := M_0^{N_0} \times_\rho M_1^{N_1} \rightarrow \mathbb{R}^N$ ,  $N \geq 3$ , be a local conformal diffeomorphism of a warped product. Then one of the following holds:*

- (i) *There exist an isometry  $\Phi: \mathbb{R}^{N_0} \times \mathbb{R}^{N_1} \rightarrow \mathbb{R}^N$ , a local conformal diffeomorphism  $f_0: M_0^{N_0} \rightarrow \mathbb{R}^{N_0}$  with conformal factor  $\rho^{-1}$ , a local isometry  $f_1: M_1^{N_1} \rightarrow \mathbb{R}^{N_1}$ , an inversion  $I$  and a homothety  $H$  in  $\mathbb{R}^N$  such that  $f = I \circ H \circ \Phi \circ (f_0 \times f_1)$ .*
- (ii) *There exist either a local conformal diffeomorphism  $f_0: M_0^{N_0} \rightarrow \mathbb{H}^{N_0}(-c)$  with conformal factor  $\rho^{-1}$  and a local isometry  $f_1: M_1^{N_1} \rightarrow \mathbb{S}^{N_1}(c)$  such that*

$$f = I \circ \Theta \circ (f_0 \times f_1)$$

*or a local conformal diffeomorphism  $f_0: M_0^{N_0} \rightarrow \mathbb{S}^{N_0}(c)$  with conformal factor  $\rho^{-1}$  and a local isometry  $f_1: M_1^{N_1} \rightarrow \mathbb{H}^{N_1}(c)$  such that*

$$f = I \circ \Theta \circ (f_1 \times f_0) \circ T_1.$$

**Remark 32.** Corollaries 29 and 31 (resp., Corollary 30) are also valid for  $N = 2$  (resp.,  $n=2$ ) if the product net  $\mathcal{E} = (E_0, E_1)$  of  $M^2$  is assumed to satisfy the condition that both  $E_0$  and  $E_1$  be spherical distributions (by Proposition 5-(ii) it suffices to assume that

this holds for either  $E_0$  or  $E_1$ ), or equivalently, that their integral curves have constant geodesic curvature. In the case of Corollary 29 (resp., Corollary 31), this is equivalent to assuming that the leaves of  $E_0$  and  $E_1$  are (resp., are mapped by  $f$  into) segments of circles or straight lines in  $\mathbb{R}^2$ . Thus, for  $N = 2$  Corollary 29 reduces to the classification of orthogonal nets on  $\mathbb{R}^2$  for which the curves of both families are segments of circles or straight lines (cf. [Da<sub>2</sub>], vol I, p. 228). On the other hand, Corollary 31 reduces to an intrinsic version of the problem in cartography solved as early as 1779 by Lagrange (cf. [Da<sub>2</sub>], vol I, p. 236), of determining all *tracés géographiques* (conformal mappings into the plane) of a rotation surface in  $\mathbb{R}^3$  such that meridians and parallels are mapped into circles or straight lines.

We have also the following consequence of Theorem 25 for isometric immersions of twisted products into Euclidean space.

**Corollary 33** *Let  $f: (M^n, \langle \cdot, \cdot \rangle) := M^{n_0} \times M^{n_1} \rightarrow \mathbb{R}^N$ ,  $n \geq 2$ , be an isometric immersion of a twisted product whose second fundamental form is adapted to the product net  $\mathcal{E} = (E_0, E_1)$  of  $M^n$ . If  $n_i = 1$ ,  $0 \leq i \leq 1$ , suppose further that the leaves of  $E_i$  are (extrinsic) circles in  $M^n$ . Then  $\langle \cdot, \cdot \rangle$  is conformal to a product metric  $\langle \cdot, \cdot \rangle^\sim = \pi_0^* \langle \cdot, \cdot \rangle_0^\sim + \pi_1^* \langle \cdot, \cdot \rangle_1^\sim$  and the conclusion of Theorem 25 holds for  $f: (M^n, \langle \cdot, \cdot \rangle) \rightarrow \mathbb{R}^N$ .*

*Proof:* By Proposition 3 the product net  $\mathcal{E} = (E_0, E_1)$  of  $M^n$  is a *TP*-net. Then, Proposition 16 and the assumption for the case when  $n_i = 1$  imply that both  $E_0$  and  $E_1$  are spherical distributions. Therefore  $\mathcal{E}$  is a *CP*-net by Proposition 5-(i), and hence  $\langle \cdot, \cdot \rangle$  is conformal to a product metric  $\langle \cdot, \cdot \rangle^\sim$  by Proposition 6. The proof is completed by taking into account the observation before Corollary 27. ■

Corollary 33 immediately implies the (local part of) the main result of [DFT], an alternative statement of which is given below. Several consequences are obtained therein of that result, which contains as a particular case the classification of the cyclides of Dupin of arbitrary dimension up to conformal transformations due to Cecil ([Ce<sub>1</sub>], [Ce<sub>2</sub>]). In fact, trying to understand the conformal nature of those results was one of the motivations of this work.

Recall that a smooth normal vector field to an isometric immersion  $f: M^n \rightarrow \mathbb{R}^N$  is called a *principal normal* with multiplicity  $m \geq 1$  if the tangent subspaces  $\mathcal{E}_\eta = \ker(\alpha_f - \langle \cdot, \cdot \rangle_\eta)$  have constant dimension  $m$ . If, in addition,  $\eta$  is parallel in the normal connection of  $f$  along the distribution  $\mathcal{E}_\eta$ , then it is said to be *Dupin*. This condition is automatic when  $m \geq 2$  (see [Re] or Proposition 8 of [DFT]). If  $\eta$  is nonvanishing, it is well-known that  $\mathcal{E}_\eta$  is an involutive distribution whose leaves are round  $m$ -dimensional spheres in  $\mathbb{R}^N$ ; see [Re] or [DFT] for details. In particular,  $\mathcal{E}_\eta$  is a spherical distribution. When  $\eta$  vanishes identically, the distribution  $\mathcal{E}_\eta = \mathcal{E}_0$  is known as the relative nullity distribution, in which case the leaves are open subsets of affine subspaces of  $\mathbb{R}^N$ .

**Corollary 34.** [DFT] *Let  $f: M^n \rightarrow \mathbb{R}^N$  be an isometric immersion carrying a Dupin principal normal  $\eta$  of multiplicity  $m$  such that the distribution  $E_\eta^\perp$  is umbilical on  $M^n$ . If  $m = n - 1$ , suppose further that the integral curves of  $E_\eta^\perp$  are extrinsic circles in  $M^n$ . Then the following holds:*

- i) there exist a Riemannian manifold  $M^{n-m}$ , a conformal diffeomorphism  $\psi$  of an open subset  $V \subset M^{n-m} \times \mathbb{R}^m$  onto  $M^n$ , an inversion  $I$ , a homothety  $H$ , an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^{N-m} \times \mathbb{R}^m$ , and an isometric immersion  $f_0: M^{n-m} \rightarrow \mathbb{R}^{N-m}$ , such that  $f \circ \psi = H \circ I \circ (f_0 \times i)$ , where  $i: \mathbb{R}^m \rightarrow \mathbb{R}^m$  is the identity map.*
- ii) there exist a Riemannian manifold  $M^{n-m}$ , a conformal diffeomorphism  $\psi$  of an open subset  $V \subset \mathbb{H}^m(-c) \times M^{n-m}$  onto  $M^n$ , an inversion  $I$ , and an isometric immersion  $f_1: M^{n-m} \rightarrow \mathbb{S}^{N-m}(c)$  such that  $f \circ \psi = I \circ \Theta \circ (i \times f_1)$ , where  $i: \mathbb{H}^m(-c) \rightarrow \mathbb{H}^m(-c)$  is the identity map.*
- iii) there exist a Riemannian manifold  $M^{n-m}$ , a conformal diffeomorphism  $\psi$  of an open subset  $V \subset M^{n-m} \times \mathbb{S}^m(c)$  onto  $M^n$ , an inversion  $I$ , and an isometric immersion  $f_0: M^{n-m} \rightarrow \mathbb{H}^{N-m}(-c)$  such that  $f \circ \psi = I \circ \Theta \circ (f_0 \times i)$ , where  $i: \mathbb{S}^m(c) \rightarrow \mathbb{S}^m(c)$  is the identity map.*

*Proof:* The net  $\mathcal{E} = (E_\eta, E_\eta^\perp)$  is a  $TP$ -net. Thus, by Theorem 4 there exist local product representations  $\psi: M_0 \times M_1 \rightarrow U \subset M$  such that the metric induced by  $f \circ \psi$  is a twisted product metric. By the assumption for the case where  $q = n - 1$  and the fact that  $E_\eta$  is a spherical distribution, we may apply Corollary 33 to  $f \circ \psi$ . For the global conclusion we refer to the end of the proof of Theorem 1 in [DFT], where one argues that different possible local structures can not be glued together. ■

Even in the case of the cyclides of Dupin we believe that some new insight is provided by the following description. Recall that a hypersurface  $f: M^n \rightarrow \mathbb{R}^{n+1}$  is a cyclide of Dupin of characteristic  $(m, n - m)$  if it has everywhere two distinct principal curvatures of multiplicities  $m$  and  $n - m$ , respectively.

**Corollary 35.** *Let  $f: M^n \rightarrow \mathbb{R}^{n+1}$  be a cyclide of Dupin of characteristic  $(m, n - m)$ . Then there exist a conformal diffeomorphism  $\psi$  of an open subset  $V \subset \mathbb{Q}_c^{n-m} \times \mathbb{S}^m$  onto  $M^n$ ,  $c > -1$ , an inversion  $I$  and a homothety  $H$  in  $\mathbb{R}^{n+1}$  such that  $f \circ \psi = H \circ I \circ \Theta \circ (f_0 \times i)$ , where  $f_0: \mathbb{Q}_c^{n-m} \rightarrow \mathbb{H}^{n-m+1}$  is an umbilical inclusion and  $i: \mathbb{S}^m \rightarrow \mathbb{S}^m$  is the identity map. Moreover, the classes of conformally congruent cyclides of Dupin are parameterized by their characteristic and the value of  $c$ .*

**Remark 36.** If  $c > 0$  then the following alternative description holds:

There exist an isometric embedding  $\Psi$  of  $\mathbb{S}^{n-m}(c) \times \mathbb{S}^m$  into  $\mathbb{S}^{n+1}(\tilde{c})$ ,  $\tilde{c} = c/(c + 1)$ , a homothety  $H$  in  $\mathbb{R}^{n+1}$  and a stereographic projection  $\mathcal{P}$  of  $\mathbb{S}^{n+1}(\tilde{c}) = H(\mathbb{S}^{n+1}(\tilde{c}))$  onto  $\mathbb{R}^{n+1}$  such that  $f \circ \psi = \mathcal{P} \circ \Psi$ .



## 8 Christoffel problem and Darboux transformations

In this section we classify submanifolds that are solutions of a generalized Christoffel problem, as well as submanifolds that admit nontrivial conformal sphere congruences.

### 8.1 Codazzi tensors

We summarize some facts on Codazzi tensors that will be needed in the sequel. Recall that a symmetric tensor  $\mathcal{S}$  is a *Codazzi tensor* if

$$(\nabla_X \mathcal{S})Y = (\nabla_Y \mathcal{S})X \text{ for all } X, Y \in \Gamma(TM),$$

where  $(\nabla_X \mathcal{S})Y = \nabla_X \mathcal{S}Y - \mathcal{S}(\nabla_X Y)$ . The following basic result is due to Reckziegel [Re] (cf. Proposition 13 of [To<sub>1</sub>]).

**Proposition 37** *Let  $\mathcal{S}$  be a Codazzi tensor on a Riemannian manifold  $M$ , and let  $\lambda \in C^\infty(M)$  be an eigenvalue of  $\mathcal{S}$  such that  $E_\lambda = \ker(\lambda I - \mathcal{S})$  has constant rank  $r$ . Then the following hold:*

- (i) *If  $r \geq 2$  then  $\lambda$  is constant along  $E_\lambda$ .*
- (ii)  *$E_\lambda$  is an umbilical distribution (in fact spherical if  $\lambda$  is constant along  $E_\lambda$ ) with mean curvature normal  $\eta$  given by  $(\lambda I - \mathcal{S})\eta = (\nabla \lambda)_{E_\lambda^\perp}$ .*

The remaining results deal with Codazzi tensors that have two distinct eigenvalues  $\lambda$  and  $\mu$  of constant multiplicities. The next one is contained in Theorem 14 of [To<sub>1</sub>].

**Proposition 38** *Let  $M$  be a Riemannian manifold and let  $\mathcal{S}$  be a Codazzi tensor on  $M$  with exactly two distinct eigenvalues  $\lambda$  and  $\mu$  of constant multiplicities. Let  $E_\lambda$  and  $E_\mu$  be the corresponding eigenbundles. Then  $(E_\lambda, E_\mu)$  is a CP-net if and only if*

$$2\rho X(\psi)Y(\psi) + \psi X(\psi)Y(\rho) + \psi Y(\psi)X(\rho) - \rho\psi \text{Hess}\psi(X, Y) = 0,$$

where  $\lambda = \psi(1 - \rho)$  and  $\mu = \psi(1 + \rho)$ .

The following is a special case of Corollary 17 of [To<sub>1</sub>]. For a product manifold  $M = M_0 \times M_1$  with product net  $(E_0, E_1)$ , we denote by  $\Pi_i: TM \rightarrow E_i$  the canonical projection onto  $E_i$ ,  $0 \leq i \leq 1$ .

**Proposition 39.** *Let  $M$  be a Riemannian manifold and let  $\mathcal{S}$  be a Codazzi tensor on  $M$  with eigenvalues  $\lambda$  and  $-\lambda$ , where  $\lambda \neq 0$  everywhere. Let  $E_+$  and  $E_-$  be the corresponding eigenbundles and assume that  $\lambda$  is constant along  $E_-$ . Then one of the following holds:*

- (i)  $\lambda$  is constant along  $E_+$  and for every point  $p \in M$  there exists a local product representation  $\psi: M_0 \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , which is an isometry with respect to a Riemannian product metric  $\langle \cdot, \cdot \rangle$  on  $M_0 \times M_1$ . Moreover,  $\psi_*^{-1} \mathcal{S} \psi_* = a(\Pi_1 - \Pi_0)$  for some  $a \neq 0$ .
- (ii) for every point  $p \in M$  there exists a local product representation  $\psi: I \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , where  $I \subset \mathbb{R}$  is an open interval, which is an isometry with respect to a warped product metric  $\langle \cdot, \cdot \rangle$  on  $I \times M_1$  with warping function  $\rho$ . Moreover,  $\psi_*^{-1} \mathcal{S} \psi_* = a(\rho \circ \pi_0)^{-2}(\Pi_1 - \Pi_0)$  for some  $a \neq 0$ .

We will also need the following fact on Codazzi tensors on warped products.

**Lemma 40** *Let  $M^n = M_0 \times_\rho M_1$  be a warped product of dimension  $n \geq 3$ , let  $(E_0, E_1)$  be its product net and let  $\mathcal{S} = \lambda \Pi_0 + \mu \Pi_1$  be a Codazzi tensor on  $M^n$  with  $\lambda \neq \mu$  everywhere. Then one of the following holds:*

- (i)  $\lambda = A \in \mathbb{R}$  and  $\mu = B(\rho \circ \pi_0)^{-1} + A$  for some  $B \neq 0$ .
- (ii)  $M_1$  is an open interval  $J \subset \mathbb{R}$ , and there exist  $A \in \mathbb{R}$  and  $\phi \in C^\infty(J)$  such that  $\lambda = A \in \mathbb{R}$  and  $\mu = (\phi \circ \pi_1)(\rho \circ \pi_0)^{-1} + A$ .
- (iii)  $M_0$  is an open interval  $I \subset \mathbb{R}$  and there exists  $\tilde{\lambda} \in C^\infty(I)$  such that  $\lambda = \tilde{\lambda} \circ \pi_0$  and  $\mu = (\rho^{-1} \int \tilde{\lambda} \rho') \circ \pi_0$ .

*Proof:* By Propositions 3 and 37, the distributions  $E_0$  and  $E_1$  are umbilical with mean curvature normals given, respectively, by

$$\eta = (\lambda - \mu)^{-1}(\nabla \lambda)_{E_1} = 0 \quad \text{and} \quad \zeta = (\mu - \lambda)^{-1}(\nabla \mu)_{E_0} = -\nabla \log \circ \rho \circ \pi_0. \quad (15)$$

Thus there exists  $\tilde{\lambda} \in C^\infty(M_0)$  such that  $\lambda = \tilde{\lambda} \circ \pi_0$ . If  $\tilde{\lambda} = A$  for some  $A \in \mathbb{R}$  and  $\mu$  is constant along  $M_1$ , then  $\nabla \mu = (A - \mu) \nabla \log \circ \rho \circ \pi_0$  by the second equation in (15). This implies that  $\nabla(\mu(\rho \circ \pi_0)) = A \nabla \rho \circ \pi_0$ , and (i) follows. Otherwise, either  $M_0$  or  $M_1$  is an open interval by Proposition 37-(iii). In the first case, we have that  $\mu = \tilde{\mu} \circ \pi_0$  for some  $\tilde{\mu} \in C^\infty(I \subset \mathbb{R} = M_0)$ , and the second equation in (15) implies that  $\tilde{\mu}' = (\tilde{\lambda} - \tilde{\mu})\rho^{-1}\rho'$ . Thus  $(\tilde{\mu}\rho)' = \tilde{\lambda}\rho'$ , which yields (iii). Finally, if  $M_1$  is an open interval  $J \subset \mathbb{R}$ , then  $\lambda = A \in \mathbb{R}$  and the second equation in (15) gives

$$(\nabla \log \circ (\mu - A))_{E_0} = -\nabla \log \circ \rho \circ \pi_0.$$

Thus there exists  $\phi \in C^\infty(J)$  such that  $\log \circ (\mu - A) = \log \circ (\rho \circ \pi_0)^{-1} + \log \circ \phi \circ \pi_1$ , and (ii) follows. ■

**Corollary 41** *Let  $M^n = M_0 \times M_1$  be a Riemannian product of dimension  $n \geq 3$ , let  $(E_0, E_1)$  be its product net and let  $\mathcal{S} = \lambda \Pi_0 + \mu \Pi_1$  be a Codazzi tensor on  $M^n$  with  $\lambda \neq \mu$  everywhere. Then either  $\lambda$  and  $\mu$  are constants or one of the factors, say,  $M_0$ , is an open interval  $I \subset \mathbb{R}$  and there exist  $\tilde{\lambda} \in C^\infty(I)$  and  $A \in \mathbb{R}$  with  $\tilde{\lambda} \neq A$  everywhere such that  $\lambda = \tilde{\lambda} \circ \pi_0$  and  $\mu = A$ .*

## 8.2 Christoffel problem

The classical proof that isothermic surfaces are precisely the ones that admit a *dual* surface (or Christoffel transform) starts by showing that if two surfaces are mapped conformally onto each other with parallel tangent planes at corresponding points, then either they are a pair of minimal surfaces with the same conformal structures and orientations, or the correspondence between them necessarily preserves curvature lines. Then, the proof proceeds by showing that in the latter case either the surfaces differ by a homothety and a translation or they are a Christoffel pair of isothermic surfaces.

Classically, two surfaces  $f: M^2 \rightarrow \mathbb{R}^3$  and  $\mathcal{F}: M^2 \rightarrow \mathbb{R}^3$  that can be mapped onto each other with preservation of curvature lines and with parallel tangent planes at corresponding points are said to be related by a *Combescure transformation* (cf. [Bi], v.II-1, p. 108). Therefore, Christoffel's characterization of isothermic surfaces implies that a surface is isothermic if and only if it admits a nontrivial conformal Combescure transform. By nontrivial we mean that the surfaces do not differ by a composition of a homothety and a translation.

In this section we classify Euclidean submanifolds of arbitrary dimension and codimension that admit a nontrivial conformal Combescure transform, the latter being understood in the sense of the following natural extension of the classical definition developed in [DT<sub>2</sub>].

Given an isometric immersion  $f: M^n \rightarrow \mathbb{R}^N$ , a map  $\mathcal{F}: M^n \rightarrow \mathbb{R}^N$  is said to be a *Combescure transform* of  $f$  determined by a symmetric tensor  $\mathcal{S}$  on  $M$  if  $\mathcal{F}_* = f_* \circ \mathcal{S}$ . This implies (see Proposition 1 of [DT<sub>2</sub>]) that  $\mathcal{S}$  is a *commuting* Codazzi tensor on  $M^n$ , i.e.,

$$\alpha_f(X, \mathcal{S}Y) = \alpha_f(\mathcal{S}X, Y) \quad \text{for all } X, Y \in \Gamma(TM).$$

If  $\mathcal{S}$  is invertible, then  $\mathcal{F}$  is an immersion with the same Gauss map as  $f$  into the Grassmann manifold of nonoriented  $n$ -planes in  $\mathbb{R}^N$ . Moreover, the requirement that the tensor  $\mathcal{S}$  be symmetric reduces in the surface case to the assumption that  $f$  and  $\mathcal{F}$  have the same curvature lines, since their second fundamental forms are related by

$$\alpha_{\mathcal{F}}(X, Y) = \alpha_f(\mathcal{S}X, Y) \quad \text{for all } X, Y \in \Gamma(TM). \quad (16)$$

If  $M^n$  is simply-connected and  $\mathcal{F}: M^n \rightarrow \mathbb{R}^N$  is any Combescure transform of  $f$  determined by a symmetric tensor  $\mathcal{S}$ , it was shown in Proposition 3 of [DT<sub>2</sub>] that there exist  $\varphi \in C^\infty(M)$  and  $\beta \in T_f^\perp M$  satisfying

$$\alpha_f(\nabla\varphi, X) + \nabla_X^\perp \beta = 0 \quad \text{for all } X \in \Gamma(TM), \quad (17)$$

such that  $\mathcal{S}$  and  $\mathcal{F}$  are given by

$$\mathcal{S} = \mathcal{S}_{\varphi, \beta} := \text{Hess } \varphi - A_\beta^f \quad \text{and} \quad \mathcal{F} = \mathcal{F}_{\varphi, \beta} := f_* \nabla\varphi + \beta. \quad (18)$$

Conversely, if  $\varphi \in C^\infty(M)$  and  $\beta \in T_f^\perp M$  satisfy (17) then  $\mathcal{F}_{\varphi, \beta}$  given by (18) defines a Combescure transform of  $f$  with  $\mathcal{S}_{\varphi, \beta}$  as the corresponding commuting Codazzi tensor.

As in the surface case, trivial Combescure transforms of a given isometric immersion  $f: M^n \rightarrow \mathbb{R}^N$  are compositions of  $f$  with a homothety and a translation, whose correspondent commuting Codazzi tensors are constant multiples of the identity tensor.

We say that an immersion  $\mathcal{F}: M^n \rightarrow \mathbb{R}^N$  is a *Christoffel transform* of  $f$  if it is a nontrivial *conformal* Combescure transform of  $f$ . In the following result we classify Euclidean submanifolds that admit Christoffel transforms.

**Theorem 42.** *Let  $f: M^n \rightarrow \mathbb{R}^N$  be an isometric immersion that admits a Christoffel transform  $\mathcal{F}: M^n \rightarrow \mathbb{R}^N$ . Then  $(f, \mathcal{F})$  is a pair of 2-isothermic submanifolds. More precisely, there exists a 2-CP-net  $(E_+, E_-)$  with respect to which the second fundamental forms of both  $f$  and  $\mathcal{F}$  are adapted. Moreover, if  $n \geq 3$  then one of the following holds:*

- (i) *for every point  $p \in M$  there exists a local product representation  $\psi: M_0 \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , which is an isometry with respect to a Riemannian product metric  $\langle \cdot, \cdot \rangle$  on  $M_0 \times M_1$ , such that  $f \circ \psi = f_0 \times f_1$  is an extrinsic Riemannian product of isometric immersions with respect to an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^{N_0} \times \mathbb{R}^{N_1}$ . Moreover,  $\mathcal{F} \circ \psi = a(-f_0 \times f_1) \circ \psi + v$  for some  $a \neq 0$  and  $v \in \mathbb{R}^N$ .*
- (ii) *for every point  $p \in M$  there exist a local product representation  $\psi: I \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , which is an isometry with respect to a warped product metric  $\langle \cdot, \cdot \rangle$  on  $(I \subset \mathbb{R}) \times M_1$  with warping function  $\rho$ , a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m} \rightarrow \mathbb{R}^N$ , a unit speed curve  $\gamma: I \rightarrow \mathbb{R}_+^m$  and an isometric immersion  $g: M_1 \rightarrow \mathbb{S}^{N-m}$  such that  $\gamma_m = \rho = \sigma \circ \gamma$  and  $f \circ \psi = \Phi \circ (\gamma \times g)$ . Moreover,  $\mathcal{F} \circ \psi = \Phi \circ (a\tilde{\gamma} \times g) + v$ , where  $a \neq 0$ ,  $v \in \mathbb{R}^N$  and  $\tilde{\gamma} = \int \gamma_m^{-2}(\tau) \gamma'(\tau) d\tau$ .*

*Proof:* Since the metric induced by  $\mathcal{F}$  is  $\langle X, Y \rangle_* = \langle \mathcal{F}_* X, \mathcal{F}_* Y \rangle = \langle \mathcal{S}X, \mathcal{S}Y \rangle$  for all  $X, Y \in \Gamma(TM)$ , the symmetry of  $\mathcal{S}$  and the assumption that  $f$  and  $\mathcal{F}$  are conformal imply that  $\mathcal{S}^2 = \lambda^2 I$  for some  $\lambda \in C^\infty(M)$ . Therefore, either  $\mathcal{S} = \pm \lambda I$  or  $TM$  splits as an orthogonal direct sum  $TM = E_+ \oplus E_-$ , where  $E_+$  and  $E_-$  are the eigenbundles of  $\mathcal{S}$  correspondent to the eigenvalues  $\lambda$  and  $-\lambda$ , respectively. In the first case Proposition 37-(i) implies that  $\lambda$  must be constant, say,  $\mathcal{S} = aI$  for some  $a \neq 0$ , and hence  $\mathcal{F} = af + v$  for some  $v \in \mathbb{R}^N$ . In the latter case, it follows from Proposition 38 that  $(E_+, E_-)$  is a CP-net. Moreover, since  $\mathcal{S}$  is commuting the second fundamental form of  $f$  is adapted to  $(E_+, E_-)$  and, because of (16), the same holds for the second fundamental form of  $\mathcal{F}$ .

Now assume that  $n \geq 3$ . Then either  $E_+$  or  $E_-$ , say, the latter, has dimension at least two, and hence  $\lambda$  must be constant along  $E_-$  by Proposition 37-(i). Thus Proposition 39 applies. In case (i), since the second fundamental form of  $f$  is adapted to  $(E_+, E_-)$  it follows from Moore's theorem that  $f \circ \psi = f_0 \times f_1$  splits as an extrinsic Riemannian product of isometric immersions with respect to an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^{N_0} \times \mathbb{R}^{N_1}$ . Moreover, integrating  $(\mathcal{F} \circ \psi)_* = (f \circ \psi)_* \psi_*^{-1} \mathcal{S} \psi_*$  with  $\psi_*^{-1} \mathcal{S} \psi_* = a(\Pi_1 - \Pi_0)$

for some  $a \neq 0$  yields  $\mathcal{F} \circ \psi = a(-f_0 \times f_1) \circ \psi + v$  for some  $v \in \mathbb{R}^N$ . In case (ii), Nölker's theorem implies that there exist a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m} \rightarrow \mathbb{R}^N$ , a unit speed curve  $\gamma: I \rightarrow \mathbb{R}_+^m$  and an isometric immersion  $g: M_1 \rightarrow \mathbb{S}^{N-m}$  such that  $\gamma_m = \rho = \sigma \circ \gamma$  and  $f \circ \psi = \Phi \circ (\gamma \times g)$ . Finally, integrating  $(\mathcal{F} \circ \psi)_* = (f \circ \psi)_* \psi_*^{-1} \mathcal{S} \psi_*$  with  $\psi_*^{-1} \mathcal{S} \psi_* = a(\rho \circ \pi_0)^{-2}(\Pi_1 - \Pi_0)$  for some  $a \neq 0$  implies that  $\mathcal{F} \circ \psi$  is as stated. ■

### 8.3 Conformal sphere congruences

Classically, two surfaces in Euclidean space are said to be related by a Ribaucour transformation when there exists a curvature-lines-preserving diffeomorphism between them such that the normal lines at corresponding points intersect at a point that is equidistant to both points. It was shown by Darboux [Da<sub>1</sub>] that a surface is isothermic if and only if it is related to another surface by a conformal Ribaucour transformation that reverts orientation. In the sequel we study whether a similar characterization holds for isothermic Euclidean submanifolds of arbitrary dimension. For that we make use of the following extension of the notion of Ribaucour transformation developed in [DT<sub>2</sub>].

**Definition 43.** An immersion  $\tilde{f}: M^n \rightarrow \mathbb{R}^N$  is said to be a *Ribaucour transform* of an isometric immersion  $f: M^n \rightarrow \mathbb{R}^N$  when  $f - \tilde{f} \neq 0$  everywhere and there exist a vector bundle isometry  $\mathcal{P}: f^*T\mathbb{R}^N \rightarrow \tilde{f}^*T\mathbb{R}^N$ , a symmetric tensor  $D$  on  $M$  and a nowhere vanishing smooth vector field  $\delta \in f^*T\mathbb{R}^N$  such that

- (a)  $\mathcal{P}Z - Z = \langle \delta, Z \rangle (f - \tilde{f})$  for all  $Z \in f^*T\mathbb{R}^N$ ;
- (b)  $\tilde{f}_* = \mathcal{P} \circ f_* \circ D$

Geometrically, condition (a) means that for any  $Z \in T_{f(x)}\mathbb{R}^N$  with  $\langle \delta, Z \rangle \neq 0$  the lines in  $\mathbb{R}^N$  through  $f(x)$  and  $\tilde{f}(x)$  tangent to  $Z$  and  $\mathcal{P}(Z)$ , respectively, intersect at a point that is at a common distance  $|Z|/\langle \delta, Z \rangle$  to  $f(x)$  and  $\tilde{f}(x)$ . When  $\langle \delta, Z \rangle = 0$  the lines are parallel.

The classical parameterization of the Ribaucour transforms of a given surface was extended as follows in Theorem 17 of [DT<sub>2</sub>].

**Theorem 44.** Let  $f: M^n \rightarrow \mathbb{R}^N$  be an isometric immersion of a simply connected Riemannian manifold and let  $\tilde{f}: M^n \rightarrow \mathbb{R}^N$  be a Ribaucour transform of  $f$  with data  $(\mathcal{P}, D, \delta)$ . Then there exists  $(\varphi, \beta)$  satisfying (17) such that

$$\tilde{f} = f - 2\nu\varphi\mathcal{F}, \quad (19)$$

where  $\mathcal{F} = f_* \text{grad} \varphi + \beta$  and  $\nu^{-1} = \langle \mathcal{F}, \mathcal{F} \rangle$ . Moreover,

$$\mathcal{P}Z = Z - 2\nu\langle \mathcal{F}, Z \rangle \mathcal{F}, \quad D = I - 2\nu\varphi\mathcal{S}_{\varphi, \beta} \quad \text{and} \quad \delta = -\varphi^{-1}\mathcal{F}. \quad (20)$$

Conversely, given  $(\varphi, \beta)$  satisfying (17) and such that  $\varphi\mathcal{F} \neq 0$  everywhere, let  $\mathcal{P}, D$  and  $\delta$  be given by (20) on an open subset  $U \subset M^n$  where  $D$  is invertible. Then  $\tilde{f}: U \rightarrow \mathbb{R}^N$  given by (19) is a Ribaucour transform of  $f|_U$  with data  $(\mathcal{P}, D, \delta)$ .

By Theorem 44, a Ribaucour transform  $\tilde{f}$  of  $f$  is completely determined by a pair  $(\varphi, \beta)$  satisfying (17), or equivalently, by  $\varphi$  and  $\mathcal{F} = f_* \text{grad } \varphi + \beta$ . We denote  $\tilde{f} = \mathcal{R}_{\varphi, \beta}(f)$ .

**Remark 45.** For later use we observe the following invariance property of the Ribaucour transformation. Let  $f: M^n \rightarrow \mathbb{R}^N$  be an isometric immersion, let  $\tilde{f}: M^n \rightarrow \mathbb{R}^N$  be a Ribaucour transform of  $f$  with data  $(\mathcal{P}, D, \delta)$  and let  $\psi: \tilde{M}^n \rightarrow M^n$  be a diffeomorphism. Endow  $\tilde{M}^n$  with the metric induced by  $\psi$ . Then the tensor  $\bar{D}$  on  $\tilde{M}^n$  defined by  $\psi_* \circ \bar{D} = D \circ \psi_*$  is symmetric, and  $\tilde{f} \circ \psi$  is a Ribaucour transform of the isometric immersion  $f \circ \psi: \tilde{M}^n \rightarrow \mathbb{R}^N$  with data  $(\bar{\mathcal{P}}, \bar{D}, \bar{\delta})$ , where  $\bar{\mathcal{P}} = \mathcal{P} \circ \psi$  and  $\bar{\delta} = \delta \circ \psi$ . Moreover, if  $\tilde{f} = \mathcal{R}_{\varphi, \beta}$  for a pair  $(\varphi, \beta)$  satisfying (17), then  $\tilde{f} \circ \psi = \mathcal{R}_{\bar{\varphi}, \bar{\beta}}(f \circ \psi)$  for  $\bar{\varphi} = \varphi \circ \psi$  and  $\bar{\beta} = \beta \circ \psi$ .

Next we summarize some basic relations between a submanifold  $f: M^n \rightarrow \mathbb{R}^N$  and one of its Ribaucour transforms  $\tilde{f}: M^n \rightarrow \mathbb{R}^N$  that will be needed in the sequel. We denote by  $\tilde{M}^n$  the manifold  $M^n$  endowed with the metric induced by  $\tilde{f}$  and mark with a tilde any object that refers to  $\tilde{f}$ .

**Proposition 46.** *The metrics and Levi-Civita connections of  $M^n$  and  $\tilde{M}^n$ , and the second fundamental forms of  $f$  and  $\tilde{f}$  are related by:*

$$\langle X, Y \rangle^\sim = \langle DX, DY \rangle, \quad (21)$$

$$D\tilde{\nabla}_X Y = \nabla_X DY + 2\nu \langle SX, DY \rangle \text{grad } \varphi - 2\nu \langle \text{grad } \varphi, DY \rangle SX, \quad (22)$$

$$\tilde{\alpha}(X, Y) = \mathcal{P}(\alpha(DX, Y) + 2\nu \langle SX, DY \rangle \beta). \quad (23)$$

It follows from (21) and the fact that  $D$  is a symmetric tensor that if  $f$  and  $\tilde{f}$  induce conformal metrics on  $M$  then  $D^2 = \rho^2 I$  for some  $\rho \in C^\infty(M)$ . Therefore, either (up to a sign change)  $D = \rho I$  or  $TM$  splits as an orthogonal direct sum  $TM = E_+ \oplus E_-$ , where  $E_+$  and  $E_-$  are the eigenbundles of  $D$  correspondent to the eigenvalues  $\rho$  and  $-\rho$ , respectively. Since  $D = I - 2\nu\varphi\mathcal{S}$  by (20), in the first case  $\mathcal{S}$  must be a multiple of the identity tensor by Proposition 37-(i), in which case the proof of Corollary 32 of [DT<sub>2</sub>] implies that there exists an inversion  $I$  in  $\mathbb{R}^N$  such that  $L'(\tilde{f}) = I(L(f))$ , where  $L$  and  $L'$  are compositions of a homothety and a translation. We say that  $\tilde{f}$  is a *Darboux transform* of  $f$  if the second possibility holds, in which case  $E_+$  and  $E_-$  are also the eigenbundles of  $\mathcal{S}$  correspondent to its distinct eigenvalues  $\lambda = \psi(1 - \rho)$  and  $\mu = \psi(1 + \rho)$ , respectively, where  $\psi = (2\nu\varphi)^{-1}$ . Thus, the condition for  $f$  and  $\tilde{f}$  to induce conformal metrics on  $M^n$  is that

$$(\lambda + \mu)\varphi = \nu^{-1} = \langle \mathcal{F}, \mathcal{F} \rangle. \quad (24)$$

**Theorem 47.** *Let  $f: M^n \rightarrow \mathbb{R}^N$  be an isometric immersion that admits a Darboux transform  $\tilde{f} = \mathcal{R}_{\varphi, \beta}(f): M^n \rightarrow \mathbb{R}^N$ . Then  $(f, \tilde{f})$  is a pair of 2-isothermic submanifolds. More precisely, there exists a 2-CP-net  $(E_+, E_-)$  with respect to which the second fundamental forms of both  $f$  and  $\tilde{f}$  are adapted. Moreover, if  $n \geq 3$  then for every  $p \in M$  there exist a local product representation  $\psi: M_0 \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , a homothety  $H$  and an inversion  $I$  in  $\mathbb{R}^N$  such that one of the following holds:*

- (i)  *$\psi$  is a conformal diffeomorphism with respect to a Riemannian product metric on  $M_0 \times M_1$  and  $f \circ \psi = H \circ I \circ g$ , where  $g = g_0 \times g_1: M_0 \times M_1 \rightarrow \mathbb{R}^{N_0} \times \mathbb{R}^{N_1} = \mathbb{R}^N$  is an extrinsic product of isometric immersions. Moreover, there exists  $i \in \{0, 1\}$  such that either  $M_i$  is one-dimensional or  $g_i(M_i) \subset \mathbb{S}^{N_i-1}(P_i; r_i)$  for some sphere  $\mathbb{S}^{N_i-1}(P_i; r_i) \subset \mathbb{R}^{N_i}$ .*
- (ii)  *$\psi$  is a conformal diffeomorphism with respect to a warped product metric on  $M_0 \times M_1$  and  $f \circ \psi = H \circ I \circ g$ , where  $g = \Phi \circ (g_0 \times g_1): M_0 \times M_1 \rightarrow \mathbb{R}^N$  is a warped product of isometric immersions determined by a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m} \rightarrow \mathbb{R}^N$ .*

*Conversely, any such isometric immersion admits a Darboux transform.*

*Proof:* Let  $TM = E_+ \oplus E_-$  be the orthogonal splitting of  $TM$  by the eigenbundles of  $D$  correspondent to the eigenvalues  $\rho$  and  $-\rho$ , respectively, where  $\rho \in C^\infty(M)$ . Since  $E_+$  and  $E_-$  are also the eigenbundles of  $\mathcal{S}$  correspondent to the eigenvalues  $\lambda = \psi(1 - \rho)$  and  $\mu = \psi(1 + \rho)$ , respectively, where  $\psi = (2\nu\varphi)^{-1}$ , it follows from Proposition 37 that  $E_+$  and  $E_-$  are umbilical distributions with mean curvature normals given by

$$\eta^+ = (\lambda - \mu)^{-1}(\text{grad } \lambda)_{E_-} \quad \text{and} \quad \eta^- = (\mu - \lambda)^{-1}(\text{grad } \mu)_{E_+}.$$

We claim that  $(E_+, E_-)$  is a CP-net. By Proposition 38, this is the case if and only if

$$2\rho X_+(\psi)X_-(\psi) + \psi X_+(\psi)X_-(\rho) + \psi X_-(\psi)X_+(\rho) - \rho\psi \text{Hess } \psi(X_+, X_-) = 0. \quad (25)$$

Here and in the sequel,  $X_+$  and  $X_-$  will always denote sections of  $E_+$  and  $E_-$ , respectively. We now compute  $\text{Hess } \psi(X_+, X_-)$ . First notice that

$$X(\nu^{-1}) = X\langle \mathcal{F}, \mathcal{F} \rangle = 2\langle \mathcal{F}_* X, \mathcal{F} \rangle = 2\langle f_* \mathcal{S} X, \mathcal{F} \rangle = 2\langle \mathcal{S} X, \text{grad } \varphi \rangle,$$

where in the last equality we have used the second equation in (18). Thus

$$\text{grad } \nu = -2\nu^2 \mathcal{S} \text{grad } \varphi,$$

and

$$\text{grad } \psi^{-1} = 2(\varphi \text{grad } \nu + \nu \text{grad } \varphi) = 2(-2\nu^2 \varphi \mathcal{S} \text{grad } \varphi + \nu \text{grad } \varphi) = 2\nu D \text{grad } \varphi.$$

We obtain

$$\text{grad } \psi = -2\psi^2 \nu D \text{ grad } \varphi, \quad (26)$$

and hence

$$\nabla_X \text{grad } \psi = -4\psi X(\psi) \nu D \text{ grad } \varphi - 2\psi^2 X(\nu) D \text{ grad } \varphi - 2\psi^2 \nu \nabla_X D \text{ grad } \varphi.$$

Therefore,

$$\begin{aligned} \text{Hess } \psi(X_+, X_-) &= \langle \nabla_{X_-} \text{grad } \psi, X_+ \rangle = -4\psi X_-(\psi) \nu \langle D \text{ grad } \varphi, X_+ \rangle - \\ &\quad 2\psi^2 X_-(\nu) \langle D \text{ grad } \varphi, X_+ \rangle - 2\psi^2 \nu \langle \nabla_{X_-} D \text{ grad } \varphi, X_+ \rangle. \end{aligned} \quad (27)$$

We compute each of the three terms in the right-hand-side of (27). By (26) we have

$$X_-(\psi) = 2\psi^2 \nu \rho X_-(\varphi) \quad \text{and} \quad X_+(\psi) = -2\psi^2 \nu \rho X_+(\varphi),$$

or, setting  $\rho = -\varphi/\tau$ ,

$$X_-(\psi) = -\frac{\psi}{\tau} X_-(\varphi) \quad \text{and} \quad X_+(\psi) = \frac{\psi}{\tau} X_+(\varphi).$$

Using that

$$\langle D \text{ grad } \varphi, X_+ \rangle = \rho X_+(\varphi) = \frac{\rho\tau}{\psi} X_+(\psi),$$

we obtain

$$-4\psi X_-(\psi) \nu \langle D \text{ grad } \varphi, X_+ \rangle = -4\rho\tau\nu X_-(\psi) X_+(\psi) = 2\psi^{-1} X_-(\psi) X_+(\psi). \quad (28)$$

Moreover, from

$$X_-(\nu) = -2\nu^2 \langle \mathcal{S} \text{ grad } \varphi, X_- \rangle = -2\nu^2 \psi(1 + \rho) X_-(\varphi) = 2\nu^2(1 + \rho) \tau X_-(\psi),$$

it follows that

$$-2\psi^2 X_-(\nu) \langle D \text{ grad } \varphi, X_+ \rangle = -4\psi\nu^2 \rho(1 + \rho) \tau^2 X_-(\psi) X_+(\psi) = -\frac{1 + \rho}{\rho\psi} X_-(\psi) X_+(\psi). \quad (29)$$

It remains to compute  $\langle \nabla_{X_-} D \text{ grad } \varphi, X_+ \rangle$ . By (22) we have

$$D \tilde{\nabla}_X Y = \nabla_X DY + 2\nu \langle \mathcal{S} X, DY \rangle \text{grad } \varphi - 2\nu \langle \text{grad } \varphi, DY \rangle \mathcal{S} X.$$

On the other hand, since  $\langle \cdot, \cdot \rangle^\sim = \rho^2 \langle \cdot, \cdot \rangle$ , the connections  $\nabla$  and  $\tilde{\nabla}$  are also related by

$$\tilde{\nabla}_X Y = \nabla_X Y + \rho^{-1} (\langle \text{grad } \rho, X \rangle Y + \langle \text{grad } \rho, Y \rangle X - \langle X, Y \rangle \text{grad } \rho).$$



Thus,

$$\begin{aligned}\nabla_X D \operatorname{grad} \varphi &= D \nabla_X \operatorname{grad} \varphi + \rho^{-1} \langle \operatorname{grad} \rho, X \rangle D \operatorname{grad} \varphi + \rho^{-1} \langle \operatorname{grad} \rho, \operatorname{grad} \varphi \rangle DX - \\ &\quad \rho^{-1} \langle X, \operatorname{grad} \varphi \rangle D \operatorname{grad} \rho - 2\nu \langle \mathcal{S}X, D \operatorname{grad} \varphi \rangle \operatorname{grad} \varphi + 2\nu \langle \operatorname{grad} \varphi, D \operatorname{grad} \varphi \rangle \mathcal{S}X.\end{aligned}$$

We obtain

$$\begin{aligned}\langle \nabla_{X_-} D \operatorname{grad} \varphi, X_+ \rangle &= X_-(\rho)X_+(\varphi) - X_-(\varphi)X_+(\rho) + 2\nu\psi(1+\rho)\rho X_-(\varphi)X_+(\varphi) \\ &= \frac{\tau}{\psi}(X_-(\rho)X_+(\psi) + X_+(\rho)X_-(\psi)) + \tau(1+\rho)X_-(\psi)X_+(\psi).\end{aligned}$$

Hence,

$$-2\psi^2\nu\langle \nabla_{X_-} D \operatorname{grad} \varphi, X_+ \rangle = \rho^{-1}(X_-(\rho)X_+(\psi) + X_+(\rho)X_-(\psi)) + \frac{1+\rho}{\rho\psi}X_-(\psi)X_+(\psi). \quad (30)$$

It follows from (27), (28), (29) and (30) that

$$\operatorname{Hess} \psi(X_-, X_+) = 2\psi^{-1}X_-(\psi)X_+(\psi) + \rho^{-1}(X_-(\rho)X_+(\psi) + X_+(\rho)X_-(\psi)),$$

and (25) follows.

Since  $\mathcal{S}$  is commuting then the second fundamental form of  $f$  is adapted to  $(E_+, E_-)$ . Because of (23), the same holds for the second fundamental form of  $\tilde{f}$ .

Now assume that  $n \geq 3$ . By Theorem 7, for every point  $p \in M$  there exist a local product representation  $\psi: M_0 \times M_1 \rightarrow U$  of  $(E_+, E_-)$  with  $p \in U \subset M$ , which is a conformal diffeomorphism with respect to a Riemannian product metric  $\langle \cdot, \cdot \rangle = \pi_0^* \langle \cdot, \cdot \rangle_0 + \pi_1^* \langle \cdot, \cdot \rangle_1$  on  $M_0 \times M_1$ . Therefore Theorem 25 can be applied to  $f \circ \psi$ . We obtain (see Remark 26-(c)) that one of the following holds:

- (a) there exist an orthogonal decomposition  $\mathbb{R}^N = \mathbb{R}^{N_0} \times \mathbb{R}^{N_1}$ , isometric immersions  $f_i: M_i \rightarrow \mathbb{R}^{N_i}$ ,  $0 \leq i \leq 1$ , a homothety  $H$  and an inversion  $I$  in  $\mathbb{R}^N$  such that  $f \circ \psi = H \circ I \circ (f_0 \times f_1)$ .
- (b) there exist a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m} \rightarrow \mathbb{R}^N$ , a conformal immersion  $f_j: M_j \rightarrow \mathbb{R}_+^m$ ,  $0 \leq j \leq 1$ , with conformal factor  $(\sigma \circ f_j)$ , an isometric immersion  $f_i: M_i \rightarrow \mathbb{S}^{N-m}$ ,  $0 \leq i \neq j \leq 1$ , a homothety  $H$  and an inversion  $I$  in  $\mathbb{R}^N$  such that  $f \circ \psi = H \circ I \circ \Phi \circ (f_j \times f_i) \circ T_j$ .

In the latter case, we may assume without loss of generality that  $j = 0$ . Define  $\langle \cdot, \cdot \rangle_0^\sim = \rho^2 \langle \cdot, \cdot \rangle_0$ , where  $\rho = \sigma \circ f_0$ . Then  $g_0 := f_0: (M_0, \langle \cdot, \cdot \rangle_0^\sim) \rightarrow \mathbb{R}_+^m$  becomes an isometric immersion, and  $\psi: (M_0 \times M_1, \langle \cdot, \cdot \rangle^\sim) \rightarrow U$  is still a conformal diffeomorphism with respect to the warped product metric  $\langle \cdot, \cdot \rangle^\sim = \pi_0^* \langle \cdot, \cdot \rangle_0^\sim + (\rho \circ \pi_0)^2 \pi_1^* \langle \cdot, \cdot \rangle_1$ . Thus, the statement in part (ii) holds by setting  $g_1 = f_1$  and  $g = \Phi \circ (g_0 \times g_1)$ .

In order to complete the proof of the direct statement, it remains to show that the restriction in case (i) holds. First notice that, by Remark 45 and the fact that  $\psi$  is conformal, we have that  $\tilde{f} \circ \psi$  is a Darboux transform  $\tilde{f} \circ \psi = \mathcal{R}_{\bar{\varphi}, \bar{\beta}}(f \circ \psi)$  of  $f \circ \psi$ , where  $\bar{\varphi} = \varphi \circ \psi$  and  $\bar{\beta} = \beta \circ \psi$ . Thus  $\mathcal{S}_{\bar{\varphi}, \bar{\beta}}$  is given by  $\psi_* \circ \mathcal{S}_{\bar{\varphi}, \bar{\beta}} = \mathcal{S}_{\varphi, \beta} \circ \psi_*$ , and therefore the eigenbundle net of  $\mathcal{S}_{\bar{\varphi}, \bar{\beta}}$  is the product net of  $M_0 \times M_1$ . It now follows from Proposition 31 in [DT<sub>2</sub>] that  $\tilde{f} \circ \psi = H \circ I \circ \tilde{g}$ , where  $\tilde{g} = \mathcal{R}_{\hat{\varphi}, \hat{\beta}}(g)$  is a Darboux transform of  $g = g_0 \times g_1$ , with  $\mathcal{S}_{\hat{\varphi}, \hat{\beta}} = -2\bar{\varphi}I + \|g\|^2 \mathcal{S}_{\bar{\varphi}, \bar{\beta}}$ . In particular, the eigenbundle net of  $\mathcal{S}_{\hat{\varphi}, \hat{\beta}}$  is also the product net of  $M_0 \times M_1$ . By Corollary 41, either there exist  $a_0, a_1 \in \mathbb{R}$  such that  $\mathcal{S}_{\hat{\varphi}, \hat{\beta}} = a_0 \Pi_0 + a_1 \Pi_1$  with  $a_0 \neq a_1$  or one of the factors, say,  $M_0$ , is an open interval  $I \subset \mathbb{R}$  and there exist  $A \in \mathbb{R}$  and  $\lambda \in C^\infty(I)$  with  $\lambda \neq A$  everywhere such that  $\mathcal{S}_{\hat{\varphi}, \hat{\beta}} = (\lambda \circ \pi_0) \Pi_0 + A \Pi_1$ . Thus, it suffices to prove that in the first case there must exist  $i \in \{0, 1\}$  such that  $g_i(M_i) \subset \mathbb{S}^{N_i-1}(P_i; r_i)$  for some sphere  $\mathbb{S}^{N_i-1}(P_i; r_i) \subset \mathbb{R}^{N_i}$ . Integrating

$$\mathcal{F}_{\hat{\varphi}, \hat{\beta}*} = g_* \mathcal{S}_{\hat{\varphi}, \hat{\beta}} \quad \text{and} \quad X(\hat{\varphi}) = \langle \mathcal{F}_{\hat{\varphi}, \hat{\beta}}, g_* X \rangle \quad \text{for all } X \in \Gamma(TM)$$

implies that

$$\mathcal{F}_{\hat{\varphi}, \hat{\beta}} = a_0(g_0 - P_0) \times a_1(g_1 - P_1) \quad \text{and} \quad 2\hat{\varphi} = a_0\|g_0 \circ \pi_0 - P_0\|^2 + a_1\|g_1 \circ \pi_1 - P_1\|^2 + C,$$

for some  $P_0 \in \mathbb{R}^{N_0}$ ,  $P_1 \in \mathbb{R}^{N_1}$  and  $C \in \mathbb{R}$ . Using that  $\hat{\varphi}(a_0 + a_1) = \hat{\nu}^{-1} = \langle \mathcal{F}_{\hat{\varphi}, \hat{\beta}}, \mathcal{F}_{\hat{\varphi}, \hat{\beta}} \rangle$ , as follows from (24), we obtain

$$a_0(a_1 - a_0)\|g_0 \circ \pi_0 - P_0\|^2 + a_1(a_1 - a_0)\|g_1 \circ \pi_1 - P_1\|^2 + C(a_0 + a_1) = 0,$$

which implies that  $a_0\|g_0 - P_0\|^2$  and  $a_1\|g_1 - P_1\|^2$  must be constants  $a_0 r_0^2$  and  $a_1 r_1^2$ , respectively. Since  $a_0$  and  $a_1$  can not be both zero, we conclude that  $\|g_i - P_i\|^2 = r_i^2$  for at least one  $i \in \{0, 1\}$ .

We now prove the converse. It suffices to show that if  $g$  is either an extrinsic product of isometric immersions  $g = g_0 \times g_1: M^n = M_0 \times M_1 \rightarrow \mathbb{R}^{N_0} \times \mathbb{R}^{N_1} = \mathbb{R}^N$  as in (i) or a warped product of isometric immersions  $g = \Phi \circ (g_0 \times g_1): M^n = M_0 \times_\rho M_1 \rightarrow \mathbb{R}^N$  determined by a warped product representation  $\Phi: \mathbb{R}_+^m \times_\sigma \mathbb{S}^{N-m} \rightarrow \mathbb{R}^N$ , then  $g$  admits a Darboux transform  $\tilde{g}$ . In effect, if this is the case and  $f = H \circ I \circ g$  for such a  $g$ , then  $H \circ I \circ \tilde{g}$  is a Darboux transform of  $f$  by Proposition 31 in [DT<sub>2</sub>].

Assume first that  $g = g_0 \times g_1: M^n = M_0 \times M_1 \rightarrow \mathbb{R}^{N_0} \times \mathbb{R}^{N_1} = \mathbb{R}^N$  is an extrinsic product of isometric immersions with  $\|g_1 - P_1\|^2 = r_1^2$  for some  $r_1 > 0$ ,  $P_1 \in \mathbb{R}^{N_1}$ . Define  $\mathcal{F} = g_1 \circ \pi_1 - P_1$  and  $\varphi = r_1^2$ . Then  $X(\varphi) = 0 = \langle \mathcal{F}, g_* X \rangle$  for all  $X \in \Gamma(TM)$ , and  $\mathcal{F}_* = g_* \Pi_1$ . Since  $\mathcal{S} = \Pi_1$  is a commuting Codazzi tensor, it follows that  $\mathcal{F}, \varphi$  determine a Ribaucour transform  $\tilde{g}$  of  $g$ . Moreover, since the eigenvalues of  $\mathcal{S}$  are  $\lambda = 0$  and  $\mu = 1$ , we have that (24) is satisfied, and hence  $\tilde{g}$  is in fact a Darboux transform of  $g$ .

Now suppose that, say,  $M_0$  is an open interval  $I \subset \mathbb{R}$ , so that  $\alpha := g_0: I \rightarrow \mathbb{R}^{N_0}$  is a unit speed curve. We may assume that the Frenet curvatures  $k_1, \dots, k_{N_0-1}$  of  $\alpha$  are

nowhere vanishing and consider the linear first order system of ODE's

$$\begin{cases} (i) \lambda' = \beta, & (ii) \beta' = B^{-1}\lambda + B^{-1}k_1V_2, \quad B \neq 0, \\ (iii) V_2' = -Bk_1\beta + k_2V_3, & (iv) V_{N_0}' = -k_{N_0-1}V_{N_0-1}, \\ (v) V_j' = -k_{j-1}V_{j-1} + k_jV_{j+1}, & 3 \leq j \leq N_0 - 1, \end{cases} \quad (31)$$

which has the first integral

$$B\lambda^2 - B^2\lambda'^2 - \sum_{j=2}^{N_0} V_j^2 = K \in \mathbb{R}. \quad (32)$$

Let  $(\lambda, \beta = \lambda', V_2, \dots, V_{N_0})$  be a solution of (31) with initial conditions chosen so that the constant  $K$  in the right-hand-side of (32) vanishes. Let  $\gamma: I \rightarrow \mathbb{R}^{N_0}$  be defined by

$$\gamma = B\lambda'\alpha' + \sum_{j=2}^{N_0} V_j e_j,$$

where  $e_1 = \alpha', e_2, \dots, e_{N_0}$  is the Frenet frame of  $\alpha$ . Using (31) we obtain that  $\gamma' = \lambda\alpha'$ , that is,  $\gamma$  is Combescure transform of  $\alpha$ . Now set

$$\mathcal{F} = \gamma \circ \pi_0 \quad \text{and} \quad \varphi = B\lambda \circ \pi_0.$$

Since  $B\lambda' = \langle \gamma, \alpha' \rangle$ , we have that  $X(\varphi) = \langle \mathcal{F}, g_*X \rangle$  for all  $X \in \Gamma(TM)$ . Moreover,  $\mathcal{F}_* = g_*(\lambda \circ \pi_0)\Pi_0$ . Since  $\mathcal{S} = (\lambda \circ \pi_0)\Pi_0$  is a commuting Codazzi tensor, it follows that  $\varphi, \mathcal{F}$  determine a Ribaucour transform  $\tilde{g}$  of  $g$ . Furthermore, since (32) holds with  $K = 0$ , it follows that (24) is satisfied, for

$$\varphi(\lambda \circ \pi_0) = B(\lambda \circ \pi_0)^2 = \|\gamma \circ \pi_0\|^2 = \langle \mathcal{F}, \mathcal{F} \rangle.$$

Hence  $\tilde{g}$  is in fact a Darboux transform of  $g$ .

Finally, we prove that any warped product  $g = \Phi \circ (g_0 \times g_1): M^n = M_0 \times_\rho M_1 \rightarrow \mathbb{R}^N$ ,  $n \geq 3$ , of isometric immersions admits a Darboux transform. We may assume that  $g$  is parameterized by

$$g = (h_1, \dots, h_{m-1}, h_m k),$$

where  $g_0 = (h_1, \dots, h_{m-1}, h_m)$ ,  $\rho = h_m$  and  $g_1 = (k_1, \dots, k_{N-m+1})$  has unit length. Define

$$\mathcal{F} = B(0, \dots, 0, k_1, \dots, k_{N-m+1}) \quad \text{and} \quad \varphi = Bh_m, \quad B \neq 0.$$

Then  $X(\varphi) = BX(h_m) = \langle \mathcal{F}, g_*X \rangle$  for all  $X \in \Gamma(TM)$ , and  $\mathcal{F}_* = g_*\mathcal{S}$ , where  $\mathcal{S} = B(h_m \circ \pi_0)^{-1}\Pi_1$ . Since  $\mathcal{S}$  is a commuting Codazzi tensor, it follows that  $\mathcal{F}, \varphi$  determine a Ribaucour transform  $\tilde{g}$  of  $g$ . Moreover, since the eigenvalues of  $\mathcal{S}$  are  $\lambda = 0$  and  $\mu = (h_m \circ \pi_0)^{-1}$ , we have that  $\varphi(\lambda + \mu) = B^2 = \langle \mathcal{F}, \mathcal{F} \rangle$ . Therefore (24) is satisfied, and hence  $\tilde{g}$  is a Darboux transform of  $g$ . ■

**Remark 48.** In Theorem 47 we have restricted ourselves to determine the Euclidean submanifolds that *admit* a Darboux transform. By making use of Proposition 40, one might in fact proceed and go into the problem of determining *all* Darboux transforms of such submanifolds.

## References

- [Bi] BIANCHI, L.: *Lezioni di Geometria Differenziale*, Bologna, 1927.
- [Bo] BONNET, O.: Mémoire sur la théorie des surfaces applicables. *J. de l'École Polytechnique XLII (1867)*.
- [BHPP] BURSTALL, F., HERTRICH-JEROMIN, U., PEDIT, F., PINKALL, U.: Curved flats and isothermic surfaces, *Math. Z.* **225** (1997), 199-209.
- [Bu] BURSTALL, F.: Isothermic surfaces: conformal geometry, Clifford algebras and integrable systems, preprint, math-dg/0003096.
- [BDPT] BRÜCK, M., DU, X., PARK, J., TERNG, C-L: Submanifold geometry of real Grassmannian Systems, *The Memoirs, vol. 155, AMS*, **735** (2002), 1-95.
- [Ce<sub>1</sub>] CECIL, T. E.: *Lie Sphere Geometry*, Springer-Verlag (1992).
- [Ce<sub>2</sub>] CECIL, T. E.: Lie sphere geometry and Dupin submanifolds, *Geometry and Topology of Submanifolds, III*, ed. L. Verstraelen and A. West, 90-107, World Scientific, Singapore, etc., 1991.
- [Ci] CIESLINSKI, J.: The Darboux-Bianchi transformation for isothermic surfaces. Classical results versus the soliton approach, *Diff. Geom. Appl.* **7** (1997), 1-28.
- [CGS] CIESLINSKI, J.; GOLDSTEIN, P.; SYM, A.: Isothermic surfaces in  $E^3$  as soliton surfaces, *Physics Letters A* **205** (1995), 37-43.
- [Ch] CHRISTOFFEL, E.: Ueber einige allgemeine Eigenschaften der Minimumsflächen, *Crelle's J.* **67** (1867), 218-228.
- [DFT] DAJCZER, M.; FLORIT, L.; TOJEIRO, R.: On a class of submanifolds carrying an extrinsic umbilic foliation, *Israel J. of Math.* **125** (2001), 203-220.
- [DT<sub>1</sub>] DAJCZER, M.; TOJEIRO, R.: An extension of the classical Ribaucour transformation, *Proc. London Math. Soc.* **85** (2002), 211-232.
- [DT<sub>2</sub>] DAJCZER, M.; TOJEIRO, R.: Commuting Codazzi tensors and the Ribaucour transformation for submanifolds. To appear in *Result. Math.*

- [Da<sub>1</sub>] DARBOUX, G.: Sur les surfaces isothermiques, *C.R.Acad. Sci. Paris* **128** (1899), 1483–1487.
- [Da<sub>2</sub>] DARBOUX, G.: *Leçons sur la théorie des surfaces* (Reprinted by Chelsea Pub. Co., 1972), Paris 1914.
- [HTD] HERTRICH-JEROMIN, U., TJADEN, E., ZÜRCHER, M.: *On Guichard's nets and cyclic systems*, preprint dg-ga/9704003.
- [La] LAFONTAINE, J.: Conformal geometry from the Riemannian point of view. Aspects of Mathematics, E 12, Vieweg, Braunschweig, 1988.
- [Mi] MOLZAN, R.: “Extrinsische Produkte und symmetrische Untermannigfaltigkeiten in Standardräumen konstanter und konstanter holomorpher Krümmung”, Doctoral Thesis, Köln, 1983.
- [MRS] MEUMERTZHEIM, M.; RECKZIEGEL, H.; SCHAAF, M.: Decomposition of twisted and warped product nets, *Result. Math.* **36** (1999), 297–312.
- [Mo] MOORE, J. D.: Isometric immersions of Riemannian products, *J. Diff. Geom.* **5** (1971), 159–168.
- [No] NÖLKER, S., Isometric immersions of warped products, *Diff. Geom. Appl.* **6** (1996), 31–50.
- [RS] RECKZIEGEL, H.; SCHAAF, M.: De Rham decomposition of netted manifolds, *Result. Math.* **35** (1999), 175–191.
- [Re] RECKZIEGEL, H.: Krümmungsflächen von isometrischen immersionen in raume konstanter krummung, *Math. Ann.* **223** (1976), 169–181.
- [Ri] RIBAUCCOUR, M.: *Sur la théorie des surfaces*. Bull. Soc. Philomathique (1870).
- [To<sub>1</sub>] TOJEIRO, R.: Conformal de Rham decomposition of Riemannian manifolds. Preprint.
- [To<sub>2</sub>] TOJEIRO, R.: Conformal immersions of warped products into Euclidean space. In preparation.

Universidade Federal de São Carlos  
 Via Washington Luiz km 235  
 13565-905 – São Carlos – Brazil  
 email: tojeiro@dm.ufscar.br