

A Characterization of Norm Compactness in the Bochner Space $L^p(G; B)$ For an Arbitrary Locally Compact Group G

Josh Isralowitz

*Department of Mathematical Sciences, New Jersey Institute of Technology,
Newark, NJ 07102-1982 U.S.A.* ¹

Abstract

In this paper, we generalize a result of N. Dinculeanu which characterizes norm compactness in the Bochner space $L^p(G; B)$ in terms of an approximate identity and translation operators, where G is a locally compact abelian group and B is a Banach space. Our characterization includes the case where G is nonabelian, and we weaken the hypotheses on the approximate identity used, providing new results even for the case $B = \mathbb{C}$ and $G = \mathbb{R}^n$.

Key words: Bochner Integral, Vector Valued Integration, Function Spaces, Compactness

1991 MSC: 46G10

1 Introduction

The classical Kolmogorov-Riesz-Tamarkin theorem characterizes norm compact subsets Γ of $L^p(\mathbb{R}^n)$ in terms of the uniform convergence (in the L^p norm) of families of convolution and translation operators, where a translation operator T^h is defined by $(T^h f)(x) = f(x - h)$ and a convolution operator J is given by $(Jf)(x) = \int_{\mathbb{R}^n} j(y) f(x - y) dy$ for an appropriate kernel j (for a brief history of this theorem see [2] or [8]). Many authors have generalized this theorem to $L^p(X)$ where X is more general than \mathbb{R}^n and for function spaces that are more general than the classical Lebesgue spaces L^p . One such generalization is the following result, due to N. Dinculeanu [2, Sect. 5] :

Email address: jbi2@buffalo.edu (Josh Isralowitz).

¹ Current Address: Mathematics Department, University at Buffalo, Buffalo, NY 14260-2900

Theorem 1 *Let G be a locally compact abelian group with Haar measure μ and B be a Banach space. A bounded subset Γ of the Bochner space $L^p(G; B)$, $1 \leq p < \infty$, is relatively norm compact iff*

1) *for any $A \subseteq G$ with $\mu(A) < \infty$, the set $\{\int_A f(y) d\mu(y) : f \in \Gamma\} \subseteq B$ is relatively norm compact in B ,*

2) *for any $\epsilon > 0$, there exists a compact $S \subseteq G$ such that*

$$\forall f \in \Gamma, \quad \|f\|_{p;G \setminus S} < \epsilon,$$

and one of the following is true:

3a) *let \mathfrak{V} be a basis of relatively compact neighborhoods of the identity in G , and for each $V \in \mathfrak{V}$ let u_V be a positive, bounded, symmetric function which vanishes outside V where $\|u_V\|_{1;G} = 1$. Then $\lim_{V \in \mathfrak{V}} \|u_V * f - f\|_{p;G} = 0$,*

3b) *for any $\epsilon > 0$, there exists a neighborhood Θ of the identity such that*

$$\forall f \in \Gamma, \quad h \in \Theta \implies \|T^h f - f\|_{p;G} < \epsilon.$$

Here we let

$$\|f\|_{p;A} = \left(\int_A \|f(x)\|_B^p d\mu(x) \right)^{\frac{1}{p}}$$

where $\|\cdot\|_B$ is the Banach space norm.

We generalize the above result in two ways. First, we generalize the theorem to include the case where G is nonabelian. Second, we weaken the hypotheses in property 3a). More specifically, we show that property 3a) can be replaced by the following: for any $\epsilon > 0$ and compact $S \subseteq G$ provided by property 2), there exists $j \in L^{p'}(G)$ such that $\forall f \in \Gamma, \|j * f - f\|_{p;S} < \epsilon$, where p' is the conjugate exponent of p and where j has compact support if G is nonabelian or $p = 1$. We note that our weakening of these hypotheses appears to be new even for the classical cases $B = \mathbb{C}$ and $G = \mathbb{R}^n$.

2 Preliminary Results

In this section we provide some preliminary results that will be needed to prove our main theorem. We start by introducing some relevant definitions and notation. In all of the rest of the paper, G will denote a locally compact group with left invariant Haar measure μ . For $1 \leq p < \infty$ and a Banach space B , $L^p(G; B)$ will denote the space of μ measurable functions $f : G \rightarrow B$ satisfying $\|f\|_B^p \in L^1(G)$, where $L^1(G)$ denotes the standard Lebesgue space of complex valued functions. The integral $\int_A f(x) d\mu(x)$ over a set A will also be written as $\int_A f(x) dx$. We refer the reader to [4, Chap. 3] for further properties of the Bochner integral.

We let Δ denote the continuous homomorphism from G into \mathbb{R}^+ defined by $\Delta(x)\mu(A) = \mu(Ax^{-1})$, which by the definition of the Bochner integral implies that $\int_G f(yx) dy = \Delta(x) \int_G f(y) dy$.

If $j \in L^p(G)$ and $f \in L^p(G, B)$, then the convolution product $(j * f)(x)$ is defined by $\int_G j(xy) f(y^{-1}) dy$. The convolution can be defined equivalently as $\int_G j(y) f(y^{-1}x) dy$, since the left invariance of μ implies that $\int_G h(xy) dy = \int_G h(y) dy$ for any $x \in G$ and any Bochner integrable function h . [2, Sect. 3]

If we require that j have compact support when either G is nonabelian or $p = 1$, then the continuity of Δ and Hölder's inequality implies that $j * f$ gives us a well defined function from G to B . Moreover, Lemma 7 and the fact that G is locally compact will imply that $j * f$ is continuous on G so that $j * f \in L^r_{\text{loc}}(G; B)$ for any $r \geq 1$, where $L^r_{\text{loc}}(G; B)$ denotes the space of μ measurable functions $f : G \rightarrow B$ such that $\|f\|_{r;K} < \infty$ for every compact $K \subseteq G$.

The main component in the proof of our main result will be Lemma 7. Before proving this lemma, however, we need an elementary result regarding the integration of functions in Γ over sets of small measure.

Definition 2 A subset $\Gamma \subseteq L^1(G; B)$ is called **Uniformly Integrable** on $S \subseteq G$ if for any $\epsilon > 0$, there exists $\delta > 0$ such that

$$\forall f \in \Gamma, \quad A \subseteq S \text{ and } \mu(A) < \delta \implies \|f\|_{1;A} < \epsilon.$$

Lemma 3 Let $\Gamma \subseteq L^1(G; B)$ be bounded and let $S \subseteq G$ be compact. For any $\epsilon > 0$, assume that there exists $j \in L^\infty(G)$ with compact support V such that

$$\forall f \in \Gamma, \quad \|f - j * f\|_{1;S} < \epsilon.$$

Then Γ^- is uniformly integrable on S where $\Gamma^- = \{f^- : f \in \Gamma\}$ and f^- denotes the function defined by $f^-(x) = f(x^{-1})$.

PROOF. First we show that for fixed j , the set $j * \Gamma = \{j * f : f \in \Gamma\}$ is uniformly integrable on G . Note that as V is compact, there exists $m > 0$ where $\Delta(y^{-1}) > m$ for all $y \in V$ since Δ has a (nonzero) minimum on V^{-1} . Let $\|f\|_{\infty;V}$ denote the essential supremum $\|f\|_B$ over V .

Thus, we have

$$\begin{aligned}
\|j * f\|_{1;A} &\leq \int_A \left(\int_V |j(xy)| \|f(y^{-1})\|_B dy \right) dx \\
&\leq (\|j\|_{\infty;V}) \mu(A) \int_V \|f(y^{-1})\|_B dy \\
&= (\|j\|_{\infty;V}) \mu(A) \int_V \left(\frac{\Delta(y^{-1})}{\Delta(y^{-1})} \right) \|f(y^{-1})\|_B dy \\
&< \frac{1}{m} (\|j\|_{\infty;V}) \mu(A) \int_V \Delta(y^{-1}) \|f(y^{-1})\|_B dy \\
&\leq \frac{1}{m} (\|j\|_{\infty;V}) \mu(A) \int_G \|f(y)\|_B dy \\
&\leq M' \mu(A)
\end{aligned}$$

where M' does not depend on $f \in \Gamma$, and so $j * \Gamma$ is uniformly integrable on G .

Now we show that Γ is uniformly integrable on G . By hypothesis, for any $\epsilon > 0$, we can choose $j \in L^\infty(G)$ with compact support, satisfying

$$\forall f \in \Gamma, \quad \|f - j * f\|_{1;G} \leq \frac{\epsilon}{2}.$$

Since $\{j * f : f \in \Gamma\}$ is uniformly integrable on G , choose $\delta > 0$ so that

$$\mu(A) < \delta \implies \forall f \in \Gamma, \quad \|j * f\|_{1;A} < \frac{\epsilon}{2},$$

and so

$$\mu(A) < \delta \implies \forall f \in \Gamma, \quad \|f\|_{1;A} \leq \|f - j * f\|_{1;A} + \|j * f\|_{1;A} < \epsilon.$$

Finally, let $S \subseteq G$ be compact and let $A \subseteq S$. By compactness, there exist m and M such that $0 < m < \Delta(y^{-1}) < M$ for all $y \in S$. Thus, if $f \in \Gamma$, we have

$$\begin{aligned}
\int_A \|f(y^{-1})\|_B dy &< \frac{1}{m} \int_A \Delta(y^{-1}) \|f(y^{-1})\|_B dy \\
&\leq \frac{1}{m} \int_G \Delta(y^{-1}) (\chi_{A^{-1}}(y^{-1}) \|f(y^{-1})\|_B) dy \\
&= \frac{1}{m} \int_G (\chi_{A^{-1}}(y)) \|f(y)\|_B dy \\
&= \frac{1}{m} \int_{A^{-1}} \|f(y)\|_B dy.
\end{aligned}$$

However,

$$\begin{aligned}
\mu(A^{-1}) &= \int_G \chi_{A^{-1}}(x) dx \\
&= \int_G \chi_{A^{-1}}(x^{-1}) \Delta(x^{-1}) dx \\
&< M \int_G \chi_{A^{-1}}(x^{-1}) dx \\
&= M\mu(A).
\end{aligned}$$

Therefore, since Γ is uniformly integrable on G , we have that Γ^- is uniformly integrable on S .

□

Before we state and prove Lemma 7, which is crucial to our main result, we will need two preliminary results, the first of which is a norm compactness criteria for general normed spaces and was proven in [3, Lemma 3].

Lemma 4 *Let S be any set, F be a normed space with norm $\|\cdot\|_F$, and $\{f_\alpha\}$ be a generalized sequence of functions with $f_\alpha : S \rightarrow F$.*

If $f : S \rightarrow F$ is a function such that $\lim_\alpha \|f_\alpha(s) - f(s)\|_F = 0$ uniformly for $s \in S$, and if each $f_\alpha(S)$ is relatively norm compact in F , then $f(S)$ is relatively norm compact in F .

The second preliminary result is the following form of the Ascoli-Arzelà theorem, which is a special case of the general theorem [6, Sect. 47]. First we introduce some relevant notation and definitions.

Let S be a subspace of G . We let $C_B(S)$ denote the Banach space of all continuous B valued functions on S endowed with the “sup” norm $\sup_{x \in S} \|f(x)\|_B$.

Definition 5 *If S is a subspace of G , then a subset $K \subseteq C_B(S)$ is said to be **equi-continuous at x** if for any $\epsilon > 0$, there exists a relative neighborhood $N_x \subseteq S$ of x such that*

$$x' \in N_x \implies \forall f \in K, \quad \|f(x) - f(x')\|_B < \epsilon.$$

If K is equi-continuous on all of S , we say K is **equi-continuous** on S

Lemma 6 Let S be a compact Hausdorff space and B a Banach space. A subset $K \subseteq C_B(S)$ is relatively norm compact iff

1) K is equi-continuous on S ,

2) for every $s \in S$, the set $K(s) = \{f(s) : f \in K\}$ is relatively norm compact in B .

Lemma 7 Let $\Gamma \subseteq L^p(G; B)$ be bounded and let $j \in L^{p'}(G)$ where j has compact support if either G is nonabelian or $p = 1$. Moreover, assume that when $A \subseteq G$ and $\mu(A) < \infty$, then $\{\int_A f^-(y) dy : f \in \Gamma\} \subseteq B$ is relatively norm compact in B .

Suppose that $S \subseteq G$ is compact and contains the identity. Then $(j * \Gamma)|_S$ is relatively norm compact in $C_B(S)$ where $(j * \Gamma)|_S = \{(j * f)|_S : f \in \Gamma\}$ and $(j * f)|_S$ is the restriction of $j * f$ to S .

PROOF. We first remark that we will show the existence of each $\{\int_A f^-(y) dy : f \in \Gamma\}$. We now check that $(j * \Gamma)$ is equi-continuous on S . Assume that $p > 1$, G is nonabelian, and $\text{supp } j = V$ is compact. Let SV denote the set $\{sv : s \in S, v \in V\}$ so that $V \subseteq SV$ since S contains the identity. Since Γ is bounded and SV is compact, we can choose $M > 0$ such that

$$\forall f \in \Gamma, \quad \left(\int_{SV} \|f(y^{-1})\|_B^p dy \right)^{\frac{1}{p}} < M.$$

Then using Hölder's equality, we compute that

$$\begin{aligned} \|(j * f)(x) - (j * f)(x')\|_B &\leq \int_V |j(xy) - j(x'y)| \|f(y^{-1})\|_B dy \\ &\leq \left(\int_V |j(xy) - j(x'y)|^{p'} dy \right)^{\frac{1}{p'}} \left(\int_V \|f(y^{-1})\|_B^p dy \right)^{\frac{1}{p}} \\ &\leq M \left(\int_V |j(xy) - j(x'y)|^{p'} dy \right)^{\frac{1}{p'}}. \end{aligned}$$

This clearly shows equi-continuity, since j is translation continuous in the $L^{p'}$ norm. If G is abelian and $p > 1$, then unimodularity implies that j does not need compact support.

If $p = 1$ and G is not necessarily abelian, choose any $\epsilon > 0$. Since Lemma 1

states that Γ^- is uniformly integrable on SV , we can choose $\delta > 0$ so that

$$A \subseteq SV \text{ and } \mu(A) < \delta \implies \forall f \in \Gamma, \quad \|f^-\|_{1;A} < \frac{\epsilon}{4 \|j\|_{\infty;A}}.$$

By Lusin's theorem, we may choose a compact $F \subseteq SV$ where

$$\mu(SV) - \mu(F) < \frac{\delta}{2}$$

and $j|_F$ is continuous. Moreover, by the Tietze extension theorem, we may extend j to a continuous function g on SV , which provides us with a continuous $g : SV \rightarrow \mathbb{C}$ and a compact $F \subseteq SV$ such that $j = g$ on F , where

$$\mu(SV) - \mu(F) < \frac{\delta}{2}.$$

Since g has compact support, for any $x \in G$, there exists a relative neighborhood $N_x \subseteq S$ of x such that

$$x' \in N_x \implies |j(xy) - j(x'y)| = |g(xy) - g(x'y)| < \frac{\epsilon}{2M}$$

for all $y \in (x^{-1}F \cap x'^{-1}F)$.

Thus fixing x and fixing any such x' , we find that

$$\begin{aligned} \|(j * f)(x) - (j * f)(x')\|_B &\leq \int_V |j(xy) - j(x'y)| \|f(y^{-1})\|_B \, dy \\ &\leq \int_{SV \setminus (x^{-1}F \cap x'^{-1}F)} |j(xy) - j(x'y)| \|f(y^{-1})\|_B \, dy \\ &\quad + \int_{(x^{-1}F \cap x'^{-1}F)} |j(xy) - j(x'y)| \|f(y^{-1})\|_B \, dy \\ &< \frac{\epsilon}{2} + \int_{(x^{-1}F \cap x'^{-1}F)} |j(xy) - j(x'y)| \|f(y^{-1})\|_B \, dy \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2M} \int_{(x^{-1}F \cap x'^{-1}F)} \|f(y^{-1})\|_B \, dy < \epsilon. \end{aligned}$$

Therefore, we have

$$\forall f \in \Gamma, \quad \|(j * f)(x) - (j * f)(x')\|_B < \epsilon.$$

Since $x \in S$ was arbitrary, it follows that $j * \Gamma$ is equi-continuous on S .

Now we check that the set $(j * \Gamma)(s) = \{(j * f)(s) : f \in \Gamma\}$ is relatively norm compact in B for every $s \in S$. Although Dinculeanu proved this for G abelian,

and although the proof is nearly identical for G non-abelian, we nevertheless present the proof for the sake of completeness. Assume that $\mu(A) < \infty \implies \{\int_A f^-(y) dy : f \in \Gamma\}$ is relatively norm compact in B and that G is not necessarily abelian.

Let j be a simple function of the form $j(y) = \sum \alpha_i \chi_{A_i}(y)$ where each A_i has finite measure and where the sum is taken over finitely many i . Then we have

$$(j * f)(x) = \sum \alpha_i \int_G \chi_{A_i}(xy) f(y^{-1}) dy = \sum \alpha_i \int_{x^{-1}A_i} f(y^{-1}) dy.$$

Thus, since each $\{\int_{x^{-1}A_i} f(y^{-1}) dy : f \in \Gamma\}$ is relatively norm compact in B , we have that $(j * \Gamma)(x) = \{\sum \alpha_i \int_{x^{-1}A_i} f(y^{-1}) dy : f \in \Gamma\}$ is relatively norm compact in B .

If j is not a simple function, then let $j_n \rightarrow j$ in the $L^{p'}$ norm where j_n is a simple function. We show that $(j_n * f)(x) \rightarrow (j * f)(x)$ uniformly for $f \in \Gamma$. To that end, again let $0 < m < \Delta(y^{-1})$ on V , so from Hölder's inequality we have

$$\begin{aligned} \|(j_n * f)(x) - (j * f)(x)\|_B &\leq \int_G |j_n(y) - j(y)| \|f(y^{-1}x)\| dy \\ &\leq \|j_n - j\|_{p';V} \left(\int_V \|f(y^{-1}x)\|_B^p dy \right)^{\frac{1}{p}} \\ &< \frac{(\Delta(x))^{\frac{1}{p}}}{m} \|j_n - j\|_{p';V} \left(\int_V \|f(y)\|_B^p dy \right)^{\frac{1}{p}} \\ &\leq \frac{M}{m} (\Delta(x))^{\frac{1}{p}} \|j_n - j\|_{p';V} \end{aligned}$$

where Γ is norm bounded by M .

Thus, $(j_n * f)(x) \rightarrow (j * f)(x)$ uniformly for $f \in \Gamma$. Note that if G is abelian, then unimodularity implies that j does not need compact support. Also note that if j has compact support V , then we may assume that $A \subseteq V$ so that A is relatively compact, and therefore each $\int_A f^-(y) dy$ exists.

Finally, by Lemma 4, each $(j * \Gamma)(x)$ for $x \in K$ is norm compact in B . Therefore, by Lemma 6, we are done.

□

We note that if G is abelian or compact, then unimodularity implies that the following two statements are equivalent:

1) $A \subseteq G$ and $\mu(A) < \infty \implies \{\int_A f^-(y) dy : f \in \Gamma\}$ is relatively norm compact in B .

2) $A \subseteq G$ and $\mu(A) < \infty \implies \{\int_A f(y) dy : f \in \Gamma\}$ is relatively norm compact in B .

3 Main Result

Now we may state our principal theorem.

Theorem 8 *Let G be a locally compact group with left invariant Haar measure μ and B be a Banach space. A bounded subset Γ of the Bochner space $L^p(G; B)$ is relatively norm compact iff*

1) *for any $A \subseteq G$ with $\mu(A) < \infty$, the set $\{\int_A f^-(y) dy : f \in \Gamma\} \subseteq B$ is relatively norm compact in B ,*

and one of the following is true:

2a) *for any $\epsilon > 0$, there exists a compact $S \subseteq G$ and $j \in L^{p'}(G)$ such that*

$$\forall f \in \Gamma, \quad \|f\|_{p; G \setminus S} < \epsilon$$

and

$$\forall f \in \Gamma, \quad \|f - j * f\|_{p; S} < \epsilon$$

where j has compact support if either G is nonabelian or $p = 1$,

2b) *for any $\epsilon > 0$, there exists a compact $S \subseteq G$ and a neighborhood Θ of the identity such that*

$$\forall f \in \Gamma, \quad \|f\|_{p; G \setminus S} < \epsilon$$

and

$$\forall f \in \Gamma, \quad h \in \Theta \implies \|T^h f - f\|_{p; S} < \epsilon$$

where T^h is the translation operator defined by $(T^h f)(x) = f(h^{-1}x)$.

PROOF. First we prove necessity for the general case of a locally compact group and $1 \leq p < \infty$. We show that both 2a) and 2b) hold. To that end, let $\epsilon > 0$. Since Γ is totally bounded, there exists $\{f_1, \dots, f_N\} \subseteq \Gamma$ such that

$$\Gamma \subseteq \bigcup_{i=1}^N B\left(f_i, \frac{\epsilon}{3}\right)$$

where $B\left(f_i, \frac{\epsilon}{3}\right)$ denotes a ball of radius $\frac{\epsilon}{3}$ at f_i in $L^p(G, B)$.

Choose a compact $S \subseteq G$ such that

$$\|f_i\|_{p;G \setminus S} < \frac{\epsilon}{2}$$

for all $i \in \{1, \dots, N\}$. Then, we have

$$\forall f \in \Gamma, \quad \|f\|_{p;G \setminus S} \leq \|f - f_k\|_{p;G \setminus S} + \|f_k\|_{p;G \setminus S} < \epsilon,$$

where $k \in \{1, \dots, N\}$ corresponds to f .

For property 2a), let \mathfrak{V} be a basis of relatively compact neighborhoods of the identity of G . For each $V \in \mathfrak{V}$, let $u_V = \mu(V)^{-1}\chi_V$. We show that $\lim_{V \in \mathfrak{V}} \|u_V * f - f\|_{p;G} = 0$ uniformly in Γ . This holds for any $f \in \{f_1, \dots, f_N\}$, so choose V' such that

$$\forall f \in \{f_1, \dots, f_N\}, \quad V \subseteq V' \implies \|u_V * f - f\|_{p;G} < \frac{\epsilon}{3}.$$

Since $\|u_V\|_{1;G} = 1$, a standard computation using Hölder's inequality and left invariance shows that $\|u_V * f - u_V * f_k\|_{p;G} \leq \|f - f_k\|_{p;G}$. Therefore, we have

$$\begin{aligned} \forall f \in \{f_1, \dots, f_N\}, \quad V \subseteq V' &\implies \|u_V * f - f\|_{p;G} \\ &\leq \|u_V * f - u_V * f_k\|_{p;G} + \|u_V * f_k - f_k\|_{p;G} \\ &\quad + \|f_k - f\|_{p;G} \\ &\leq 2\|f - f_k\|_{p;G} + \|f_k - j * f_k\|_{p;G} \\ &< \epsilon, \end{aligned}$$

where $k \in \{1, \dots, N\}$ corresponds to f .

For property 2b), choose a neighborhood Θ of the identity with the property

$$\forall k \in \{1, \dots, N\}, \quad y \in \Theta \implies \int_G \|f_k(x) - f_k(y^{-1}x)\|_B^p dx < \left(\frac{\epsilon}{3}\right)^p.$$

Thus, we have

$$\begin{aligned} \forall f \in \Gamma, \quad y \in \Theta &\implies \left(\int_G \|f(x) - f(y^{-1}x)\|_B^p dx\right)^{1/p} \\ &\leq \left(\int_G \|f(x) - f_k(x)\|_B^p dx\right)^{1/p} + \left(\int_G \|f_k(x) - f_k(y^{-1}x)\|_B^p dx\right)^{1/p} \\ &\quad + \left(\int_G \|f_k(x) - f(y^{-1}x)\|_B^p dx\right)^{1/p} \\ &< \epsilon \end{aligned}$$

by left invariance.

Finally, if $\mu(A) < \infty$ and G is abelian, then the continuity of the map $f \mapsto \int_A f^-(x) dx$ of $L^p(G; B)$ into B implies that $\{\int_A f^-(x) dx : f \in \Gamma\} \subseteq B$ is norm compact in B . If G is nonabelian, then as we noted in the proof of Lemma 7, we only need to consider the case where A is relatively compact. Therefore, the continuity of the map $f \mapsto \int_A f^-(x) dx$ easily follows.

Now we prove sufficiency. Assume that properties 1) and 2a) hold. Let $\{f_n\} \subseteq \Gamma$ and choose any $\epsilon > 0$. Choose a compact $S \subseteq G$ and $j \in L^{p'}(G)$ so that

$$\forall f_n \in \Gamma, \quad \|f_n\|_{p; G \setminus S} < \frac{\epsilon}{6}$$

and

$$\forall f_n \in \Gamma, \quad \|f_n - j * f_n\|_{p; S} < \frac{\epsilon}{3}.$$

Moreover, since G is locally compact, we may assume that S contains the identity. Therefore, Lemma 7 implies that $(j * \Gamma)|_S$ is relatively compact in $C_B(S)$, and so passing to a subsequence if necessary (and assuming without loss of generality that each $j * f_n$ is restricted to S), we have that $j * f_n \rightarrow f$ in $C_B(S)$ for some $f \in C_B(S)$. Define f to be zero on $G \setminus S$.

By our hypothesis and the fact that $j * f_n \rightarrow f$ in $C_B(S)$, we have that for all large enough $n \in \mathbb{N}$,

$$\begin{aligned} \forall f \in \Gamma, \quad \|f_n - f\|_{p; G} &\leq \|f_n - f\|_{p; G \setminus S} + \|f_n - f\|_{p; S} \\ &\leq \epsilon/3 + \|f_n - f\|_{p; S} \\ &\leq \epsilon/3 + \|f_n - j * f_n\|_{p; S} + \|j * f_n - f\|_{p; S} \\ &\leq \epsilon \end{aligned}$$

Hence, Γ is relatively norm compact.

Finally, we show that property 2b) implies property 2a). Choose a neighborhood Θ' of the identity such that

$$y \in \Theta' \implies \int_S \|f(x) - f(y^{-1}x)\|_B^p dx < \epsilon^p.$$

Since G is locally compact, we may assume Θ' is relatively compact, and so Θ' has nonzero finite measure. Let $j = \mu(\Theta')^{-1} \chi_{\Theta'}$.

Then for any $f \in \Gamma$, we find, using Hölder's inequality and Fubini's theorem, that

$$\begin{aligned}
\forall f \in \Gamma, \quad \|f - j * f\|_{p;S}^p &= \int_S \left\| f(x) - \int_{\Theta'} j(y) f(y^{-1}x) dy \right\|_B^p dx \\
&\leq \int_S \left(\int_{\Theta'} j(y)^{\frac{1}{p'}} j(y)^{\frac{1}{p}} \|f(x) - f(y^{-1}x)\|_B dy \right)^p dx \\
&\leq \int_S \left(\int_{\Theta'} j(y) \|f(x) - f(y^{-1}x)\|_B^p dy \right) dx \\
&= \int_{\Theta'} \left(\int_S \|f(x) - f(y^{-1}x)\|_B^p dx \right) j(y) dy.
\end{aligned}$$

But for all $y \in \Theta'$, we have $\int_S \|f(x) - f(y^{-1}x)\|_B^p dx < \epsilon^p$, and so $\|f - j * f\|_{p;S} < \epsilon$.

□

4 Final Comments

We make two final comments. First, Theorem 1 was proven in [7] for the case where $B = \mathbb{C}$, G is an arbitrary locally compact group, and each $u_V = \mu(V)^{-1} \chi_V$ where V is a nonempty neighborhood of the identity, so that our theorem generalizes this result in two different directions.

Also, in the case $B = \mathbb{C}$ and $G = \mathbb{R}^n$, Shurenkova and Buldygin [1] proved Theorem 1 for the case when j is a function of the form ω_h for some $h > 0$, where $\{\omega_h : h \in \mathbb{R}^+\}$ is any family of real functions in $\subseteq L^1(\mathbb{R}^n) \cap L^{p'}(\mathbb{R}^n)$ satisfying the following:

- 1) $h > 0$, $\|\omega_h\|_{1;\mathbb{R}^n} = 1$, and ω_h is continuous a.e. on \mathbb{R}^n ,
- 2) for any $\delta > 0$, $\lim_{h \rightarrow 0} \|\omega_h\|_{p;\{|x|>\delta\}} = 0$,
- 3) $\|\omega_h\|_{1;\mathbb{R}^n} = 1$

(here of course, we do not consider the property 3b) in the statement of Theorem 1.)

Thus, considering the well-known fact that $L^p(\mathbb{R}^n)$ functions can be arbitrarily approximated (in the L^p norm) by such functions [9, chap. 9, Sect. 2], it is clear that our theorem is a significant generalization of Shurenkova and Buldygin's results.

Acknowledgements

The author would like to thank the entire NJIT mathematics department, and especially Professor Denis Blackmore, for their time and very generous help.

References

- [1] BULDYGIN, V. and SHURENKOVA, A. *On a Criterion of Compactness in the Space L_p* , Theory of Probability and Mathematical Statistics, No. 52, 1996, 33 - 37.
- [2] DINCULEANU, N. *On Kolmogorov-Tamarkin and M. Riesz Compactness Criteria in Function Spaces Over a Locally Compact Group*, Journal of Mathematical Analysis and Applications, No. 89, 1982, 67 - 85.
- [3] DINCULEANU, N. *Strong Additivity, Absolute Continuity, and Compactness in Spaces of Measures*, Journal of Mathematical Analysis and Applications, No. 45, 1974, 172 - 188.
- [4] HILLE, E. and PHILLIPS, R. *Functional Analysis and Semi-Groups*, American Mathematical Society., 1957
- [5] LOOMIS, L. *An Introduction to Abstract Harmonic Analysis*, Van Nostrand Company, inc., 1953.
- [6] MUNKRES, J. *Topology* Prentice Hall, 1999.
- [7] TELEMANN, S. *Sur Les Ensembles Compacts de Fonctions Sommables*, Rev. Math. Pures Appl. No. 6, 1961, 659-684.
- [8] WELLAND, R. and GOES, S. *Compactness Criteria for Köthe Spaces* Math Ann. No. 188, 1970, 251 - 269.
- [9] WHEEDEN, R. and ZYGMUND, A. *Measure and Integral*, Marcel Dekker Inc., 1977.