

# A volume formula for hyperbolic tetrahedra in terms of edge lengths

Jun MURAKAMI

Department of Mathematical Science  
School of Science and Engineering  
Waseda University  
Tokyo 169-8555, JAPAN  
murakami@waseda.jp

Akira USHIJIMA

Mathematics Institute  
University of Warwick  
Coventry CV4 7AL, UK  
ushijima@maths.warwick.ac.uk

## Abstract

We give a closed formula for volumes of generic hyperbolic tetrahedra in terms of edge lengths. The cue of our formula is by the volume conjecture for the Turaev-Viro invariant of closed 3-manifolds, which is defined from the quantum  $6j$ -symbols. This formula contains the dilogarithm functions, and we specify the adequate branch to get the actual value of the volumes.

**Keywords:** hyperbolic tetrahedron, quantum  $6j$ -symbol, volume formula.

**2000 Mathematics Subject Classifications:** primary: 52A38; secondary: 51M09.

## 1 Introduction

To determine the volume of a hyperbolic tetrahedron is one of the fundamental problem of the hyperbolic 3-space. For orthoschemes, a closed formula is known at the very beginning of the hyperbolic geometry. However, closed formulas for the volumes of generic hyperbolic tetrahedra are given rather recently by [CK] and is modified to more symmetric form in [MY]. These formulas are given in terms of dihedral angles of the tetrahedron.

In this paper, we give a closed formula in terms of edge lengths instead of dihedral angles. The background of this formula is the volume conjecture for

hyperbolic 3-manifolds. In [Ka], R. Kashaev conjectured in a relation between certain quantum invariants of hyperbolic knots and the volumes of the knot complements of them, and checked it numerically for some examples. In [Mu], H. Murakami observed that such relation also holds for some closed hyperbolic 3-manifolds. These observations suggest that there is a relation between the volume of a hyperbolic 3-manifold and its Turaev-Viro invariant (see [TV]), which is a state-sum invariant given by quantum  $6j$ -symbols associated to the tetrahedra of a simplicial decomposition of the manifold. Indeed, applying Kashaev's method to investigating the asymptotics of the quantum  $6j$ -symbols, we get the formula for the volume of a hyperbolic tetrahedron. The formula by dihedral angles in [MY] also comes from the  $6j$ -symbols in the same way, and it resembles so much the formula given in this paper.

Such similarity comes from the relations in [Mi, p. 294] and [SA, p. 311], by which the volume function in terms of edge lengths implicitly comes from the volume function in terms of dihedral angles. We can apply this method to the formula in [MY]. However, to get the actual volumes, we have to specify the adequate branches of the dilogarithm functions in the formula. Here we give a direct proof of our formula, and show that we can actually compute the volume of any hyperbolic tetrahedron from it.

**Acknowledgement.** We would like to thank Professor Yana Mohanty for informing the relations in [Mi] and [SA]. The second author would like to thank the support from the JSPS Postdoctoral Fellowships for Research Abroad 2003.

## 2 Volume formulae for hyperbolic tetrahedra

For complex numbers  $z, a_1, a_2, \dots, a_6$ , we first define a complex-valued function  $U = U(a_1, a_2, a_3, a_4, a_5, a_6, z)$  as follows:

$$U := \text{Li}_2(z) + \text{Li}_2(a_1 a_2 a_4 a_5 z) + \text{Li}_2(a_1 a_3 a_4 a_6 z) + \text{Li}_2(a_2 a_3 a_5 a_6 z) \\ - \text{Li}_2(-a_1 a_2 a_3 z) - \text{Li}_2(-a_1 a_5 a_6 z) - \text{Li}_2(-a_2 a_4 a_6 z) - \text{Li}_2(-a_3 a_4 a_5 z),$$

where  $\text{Li}_2(\cdot)$  is the dilogarithm function defined by the analytic continuation of the following integral:

$$\text{Li}_2(x) := - \int_0^x \frac{\log(1-t)}{t} dt \quad \text{for a real number } x < 1.$$

Secondly we define two complex values  $z_-$  and  $z_+$  as follows:

$$z_- := \frac{-q_1 - \sqrt{q_1^2 - 4 q_0 q_2}}{2 q_2} \quad \text{and} \quad z_+ := \frac{-q_1 + \sqrt{q_1^2 - 4 q_0 q_2}}{2 q_2},$$

where

$$\begin{aligned}
q_0 &:= 1 + a_1 a_2 a_3 + a_1 a_5 a_6 + a_2 a_4 a_6 + a_3 a_4 a_5 \\
&\quad + a_1 a_2 a_4 a_5 + a_1 a_3 a_4 a_6 + a_2 a_3 a_5 a_6, \\
q_1 &:= -a_1 a_2 a_3 a_4 a_5 a_6 \left\{ (a_1 - a_1^{-1}) (a_4 - a_4^{-1}) \right. \\
&\quad \left. + (a_2 - a_2^{-1}) (a_5 - a_5^{-1}) + (a_3 - a_3^{-1}) (a_6 - a_6^{-1}) \right\}, \\
q_2 &:= a_1 a_2 a_3 a_4 a_5 a_6 (a_1 a_4 + a_2 a_5 + a_3 a_6 \\
&\quad + a_1 a_2 a_6 + a_1 a_3 a_5 + a_2 a_3 a_4 + a_4 a_5 a_6 + a_1 a_2 a_3 a_4 a_5 a_6).
\end{aligned}$$

These  $z_-$  and  $z_+$  are originally defined in [MY] as two non-trivial solutions of  $\Re(z \frac{\partial U}{\partial z}) = 0$ , where  $\Re$  means the real part. We will see it (and other properties of  $z_-$  and  $z_+$ ) in Section 4.

Finally we define a complex-valued function  $V(a_1, a_2, a_3, a_4, a_5, a_6)$  as follows:

$$\begin{aligned}
V(a_1, a_2, a_3, a_4, a_5, a_6) \\
&:= \frac{\sqrt{-1}}{4} \left\{ \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_-) - z_- \frac{\partial U}{\partial z} \Big|_{z=z_-} \log z_- \right) \right. \\
&\quad \left. - \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_+) - z_+ \frac{\partial U}{\partial z} \Big|_{z=z_+} \log z_+ \right) \right\}.
\end{aligned}$$

Then the following theorem has been proved in [MY]:

**Theorem 2.1.** *Let  $T = T(A_1, A_2, A_3, A_4, A_5, A_6)$  be a tetrahedron with dihedral angles as shown in Figure 1, and let  $\text{Vol}(T)$  be its volume. Then the following equations hold:*

1. *if  $T$  is hyperbolic, then  $\text{Vol}(T) = -V_A$ ,*
2. *if  $T$  is Euclidean, then  $V_A = 0$ , and*
3. *if  $T$  is spherical, then  $\text{Vol}(T) = -\sqrt{-1} V_A$ ,*

where  $V_A := V(e^{\sqrt{-1} A_1}, e^{\sqrt{-1} A_2}, e^{\sqrt{-1} A_3}, e^{\sqrt{-1} A_4}, e^{\sqrt{-1} A_5}, e^{\sqrt{-1} A_6})$ .

We can easily see that both  $z_-$  and  $z_+$  have the following simpler presentations in this case:

$$\begin{aligned}
z_- &= -\frac{2}{q_A} \left( \sin A_1 \sin A_4 + \sin A_2 \sin A_5 + \sin A_3 \sin A_6 - \sqrt{\det G_A} \right), \\
z_+ &= -\frac{2}{q_A} \left( \sin A_1 \sin A_4 + \sin A_2 \sin A_5 + \sin A_3 \sin A_6 + \sqrt{\det G_A} \right),
\end{aligned}$$




Figure 1: Dihedral angles and edge lengths of a tetrahedron

where

$$\begin{aligned}
 q_A := & e^{\sqrt{-1}(A_1+A_4)} + e^{\sqrt{-1}(A_2+A_5)} + e^{\sqrt{-1}(A_3+A_6)} \\
 & + e^{\sqrt{-1}(A_1+A_2+A_6)} + e^{\sqrt{-1}(A_1+A_3+A_5)} + e^{\sqrt{-1}(A_2+A_3+A_4)} + e^{\sqrt{-1}(A_4+A_5+A_6)} \\
 & + e^{\sqrt{-1}(A_1+A_2+A_3+A_4+A_5+A_6)}
 \end{aligned}$$

and  $G_A$  is the Gram matrix in terms of dihedral angles defined as follows (precise definition will be seen in Section 3):

$$G_A := \begin{pmatrix} 1 & -\cos A_1 & -\cos A_2 & -\cos A_6 \\ -\cos A_1 & 1 & -\cos A_3 & -\cos A_5 \\ -\cos A_2 & -\cos A_3 & 1 & -\cos A_4 \\ -\cos A_6 & -\cos A_5 & -\cos A_4 & 1 \end{pmatrix}. \quad (2.1)$$

The original proof of Theorem 2.1 was to show the equivalence to Cho-Kim's formula in [CK], and now there are several different proofs and formulations (see [DL, Us]).

The main purpose of this paper is to give a proof of the following theorem :

**Theorem 2.2.** *Let  $l_i$  be the length of the edge corresponding to a dihedral angle  $A_i$  (see Figure 1 again), and we denote by  $T = T(l_1, l_2, l_3, l_4, l_5, l_6)$  a hyperbolic tetrahedron with edge lengths  $l_i$ . Then the following equation holds:*

$$\text{Vol}(T) = V_l - \sum_{i=1}^6 l_i \frac{\partial V_l}{\partial l_i},$$

where  $V_l := V(-e^{l_4}, -e^{l_5}, -e^{l_6}, -e^{l_1}, -e^{l_2}, -e^{l_3})$ .

We here note that, in this case, both  $z_-$  and  $z_+$  have the following simpler presentations:

$$\begin{aligned} z_- &= \frac{2}{q_l} \left( \sinh l_1 \sinh l_4 + \sinh l_2 \sinh l_5 + \sinh l_3 \sinh l_6 - \sqrt{\det G_l} \right), \\ z_+ &= \frac{2}{q_l} \left( \sinh l_1 \sinh l_4 + \sinh l_2 \sinh l_5 + \sinh l_3 \sinh l_6 + \sqrt{\det G_l} \right), \end{aligned}$$

where

$$\begin{aligned} q_l &:= e^{(l_1+l_4)} + e^{(l_2+l_5)} + e^{(l_3+l_6)} \\ &\quad - e^{(l_1+l_2+l_3)} - e^{(l_1+l_5+l_6)} - e^{(l_2+l_4+l_6)} - e^{(l_3+l_4+l_5)} + e^{(l_1+l_2+l_3+l_4+l_5+l_6)} \end{aligned}$$

and  $G_l$  is the Gram matrix in terms of edge lengths defined as follows (precise definition will be seen in Section 3):

$$G_l := \begin{pmatrix} -1 & -\cosh l_4 & -\cosh l_5 & -\cosh l_3 \\ -\cosh l_4 & -1 & -\cosh l_6 & -\cosh l_2 \\ -\cosh l_5 & -\cosh l_6 & -1 & -\cosh l_1 \\ -\cosh l_3 & -\cosh l_2 & -\cosh l_1 & -1 \end{pmatrix}. \quad (2.2)$$

### 3 A relationship between dihedral angles and edge lengths of a hyperbolic simplex

Unlike Euclidean one, in hyperbolic geometry the dihedral angles and the lengths of the edges of a simplex is closely related each other. In this section we review two well-known aspects of the relationship, namely we review how to obtain dihedral angles from edge lengths, and vice verse. See, for example, [AVS, Lu] for details.

For any  $n \geq 3$ , let  $(\mathbb{H}^n, \langle \cdot, \cdot \rangle)$  be the *hyperboloid model of an  $n$ -dimensional hyperbolic space*. For a hyperbolic  $n$ -simplex  $\triangle \subset \mathbb{H}^n$ , we denote by  $\{\mathbf{v}_i\}_{i=1}^{n+1}$  the set of vectors of the vertices, and by  $\rho_{ij}$  the length of the edge with end points  $\mathbf{v}_i$  and  $\mathbf{v}_j$ . Then the following relation holds:

$$\langle \mathbf{v}_i, \mathbf{v}_j \rangle = -\cosh \rho_{ij}. \quad (3.1)$$

We denote by  $\mathbf{u}_i$  the outward normal vector to the facet (i.e.,  $n-1$ -dimensional face) opposite to  $\mathbf{v}_i$ , and by  $\theta_{ij}$  the dihedral angle between two facets with normal vectors  $\mathbf{u}_i$  and  $\mathbf{u}_j$  (i.e., two facets opposite to  $\mathbf{v}_i$  and  $\mathbf{v}_j$ ). We also define  $\theta_{ii} := \pi$ . Then the following relation holds:

$$\langle \mathbf{u}_i, \mathbf{u}_j \rangle = -\cos \theta_{ij}. \quad (3.2)$$

## From dihedral angles to edge lengths

Let  $G_A := (\langle \mathbf{u}_i, \mathbf{u}_j \rangle)_{i,j=1}^{n+1}$  be an  $(n+1) \times (n+1)$  real matrix of signature  $(n, 1)$ . This matrix is called the *Gram matrix in terms of dihedral angles*, and determined independent of the position in  $\mathbb{H}^n$  via (3.2). If  $n = 3$  and dihedral angles are named as in Figure 1, then you can easily check that  $G_A$  becomes as in (2.1).

Let  $M$  be the diagonal matrix with diagonal entries  $\langle \mathbf{u}_i, \mathbf{v}_i \rangle$ , and let  $(c_{Aij})_{i,j=1}^{n+1} := (\det G_A) G_A^{-1}$  be the cofactor matrix of  $G_A$ , namely  $c_{Aij} := (-1)^{i+j} \det G_{ij}$ , where  $G_{ij}$  is a submatrix of order  $n$  obtained from  $G_A$  by removing the  $i$ th row and  $j$ th column. Then we have  $G_l = M G_A^{-1} M$ . So, using (3.1) and  $\langle \mathbf{u}_i, \mathbf{v}_i \rangle = -\sqrt{(\det G_A)/c_{Aii}}$  for any  $i$ , we have

$$\begin{aligned} \cosh \rho_{ij} &= -\sqrt{\frac{\det G_A}{c_{Aii}}} \frac{c_{Aij}}{\det G_A} \sqrt{\frac{\det G_A}{c_{Ajj}}} \\ &= \frac{c_{Aij}}{\sqrt{c_{Aii} c_{Ajj}}}. \end{aligned} \quad (3.3)$$

## From edge lengths to dihedral angles

The method is just the same as in the previous subsection. Let  $G_l := (\langle \mathbf{v}_i, \mathbf{v}_j \rangle)_{i,j=1}^{n+1}$  be an  $(n+1) \times (n+1)$  real matrix of signature  $(n, 1)$ . This matrix is called the *Gram matrix in terms of edge lengths*, and determined independent of the position in  $\mathbb{H}^n$  via (3.1). If  $n = 3$  and edge lengths are named as in Figure 1, then you can easily check that  $G_l$  becomes as in (2.2).

Let  $(c_{lij})_{i,j=1}^{n+1} := (\det G_l) G_l^{-1}$  be the cofactor matrix of  $G_l$ . Then, using (3.2) and  $\langle \mathbf{u}_i, \mathbf{v}_i \rangle = -\sqrt{(\det G_l)/c_{lii}}$  for any  $i$  in this case, we have the following relationship between  $\theta_{ij}$  and edge lengths:

$$\cos \theta_{ij} = \frac{c_{lij}}{\sqrt{c_{lii} c_{ljj}}}. \quad (3.4)$$

*Remark.* We already know a volume formula Theorem 2.1 in terms of dihedral angles. So combine it with (3.4) and we can theoretically obtain a formula in terms of edge lengths. However the formula in Theorem 2.2 does not come from this procedure, and its proof given in the next section is also independent of Theorem 2.1.

## 4 Proof of the main theorem

We start this section with checking the details of  $z_-$  and  $z_+$ , most of which are already known in [MY].

**Proposition 4.1.** *For each  $j = "+"$  or  $"-"$ , we have  $z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \equiv 0$  modulo  $2\pi\sqrt{-1}$ .*

*Proof.* Using  $\frac{d}{dz} \text{Li}_2(z) = -\frac{1}{z} \log(1-z)$ , we have

$$\begin{aligned} z \frac{\partial U}{\partial z} &= -\log(1-z) - \log(1-z a_1 a_2 a_4 a_5) - \log(1-z a_1 a_3 a_4 a_6) \\ &\quad - \log(1-z a_2 a_3 a_5 a_6) + \log(1+z a_1 a_2 a_3) + \log(1+z a_1 a_5 a_6) \\ &\quad + \log(1+z a_2 a_4 a_6) + \log(1+z a_3 a_4 a_5). \end{aligned}$$

So the following are equivalent:

$$\begin{aligned} \Re(z \frac{\partial U}{\partial z}) &= 0 \\ \iff \log \left| \frac{(1+z a_1 a_2 a_3)(1+z a_1 a_5 a_6)(1+z a_2 a_4 a_6)(1+z a_3 a_4 a_5)}{(1-z)(1-z a_1 a_2 a_4 a_5)(1-z a_1 a_3 a_4 a_6)(1-z a_2 a_3 a_5 a_6)} \right| &= 0 \\ \iff (1+z a_1 a_2 a_3)(1+z a_1 a_5 a_6)(1+z a_2 a_4 a_6)(1+z a_3 a_4 a_5) \\ &\quad \pm (1-z)(1-z a_1 a_2 a_4 a_5)(1-z a_1 a_3 a_4 a_6)(1-z a_2 a_3 a_5 a_6) = 0. \end{aligned}$$

When we choose “ $-$ ” part of the above equation, the following equation holds:

$$\begin{aligned} &(1+z a_1 a_2 a_3)(1+z a_1 a_5 a_6)(1+z a_2 a_4 a_6)(1+z a_3 a_4 a_5) \\ &- (1-z)(1-z a_1 a_2 a_4 a_5)(1-z a_1 a_3 a_4 a_6)(1-z a_2 a_3 a_5 a_6) \\ &= z(q_2 z^2 + q_1 z + q_0). \end{aligned}$$

Thus  $z_j$  is a solution of  $\Re(z \frac{\partial U}{\partial z}) = 0$ , namely  $\Re(z_j \frac{\partial U}{\partial z} \Big|_{z=z_j}) = 0$ .

On the other hand, by the definition of log, the following holds modulo  $2\pi$ :

$$\begin{aligned} &\Im(z_j \frac{\partial U}{\partial z} \Big|_{z=z_j}) \\ &\equiv \arg \frac{(1+z_j a_1 a_2 a_3)(1+z_j a_1 a_5 a_6)(1+z_j a_2 a_4 a_6)(1+z_j a_3 a_4 a_5)}{(1-z_j)(1-z_j a_1 a_2 a_4 a_5)(1-z_j a_1 a_3 a_4 a_6)(1-z_j a_2 a_3 a_5 a_6)} \\ &= 0, \end{aligned}$$

where  $\Im$  means the imaginary part. We have thus proved Proposition 4.1.  $\square$

Proposition 4.1 means that  $z_j \frac{\partial U}{\partial z} \Big|_{z=z_j}$  can take discrete values. But, by the definition,  $z_j \frac{\partial U}{\partial z} \Big|_{z=z_j}$  is an analytic function in terms of  $a_1, a_2, \dots, a_6$ . So

$\frac{\partial}{\partial a_i} \left( z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \right)$  must be zero. Thus we have the following corollary:

**Corollary 4.2.** *For any  $i = 1, 2, \dots, 6$ , we have  $\frac{\partial}{\partial a_i} \left( z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \right) = 0$ .  $\square$*

Next we prepare the following proposition:

**Proposition 4.3.** *For each  $i = 1, 2, \dots, 6$ , we have  $2 \frac{\partial V_l}{\partial l_i} \equiv A_i$  modulo  $\pi$ .*

*Proof.* Here we only show that  $2 \frac{\partial V_l}{\partial l_1} \equiv A_1$  modulo  $\pi$ . Other cases can be shown similarly. Using Corollary 4.2, the following holds for each  $j = “+”$  or “ $-$ ”:

$$\begin{aligned}
& \frac{\partial}{\partial l_1} \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_j) - z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \log z_j \right) \\
&= \frac{\partial}{\partial a_4} \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_j) - z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \log z_j \right) \frac{da_4}{dl_1} \\
&= a_4 \left\{ \frac{\partial U}{\partial a_4} \Big|_{z=z_j} + \frac{\partial U}{\partial z} \Big|_{z=z_j} \frac{\partial z_j}{\partial a_4} \right. \\
&\quad \left. - (\log z_j) \frac{\partial}{\partial a_4} \left( z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \right) - \left( z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} \right) \left( \frac{1}{z_j} \frac{\partial z_j}{\partial a_4} \right) \right\} \\
&= a_4 \frac{\partial U}{\partial a_4} \Big|_{z=z_j} \\
&\equiv \log \frac{\varphi(z_j)}{\psi(z_j)} \pmod{2\pi\sqrt{-1}},
\end{aligned}$$

where  $\varphi(z)$  and  $\psi(z)$  are defined as follows:

$$\begin{aligned}
\varphi(z) &:= (1 + z a_2 a_4 a_6) (1 + z a_3 a_4 a_5), \\
\psi(z) &:= (1 - z a_1 a_2 a_4 a_5) (1 - z a_1 a_3 a_4 a_6).
\end{aligned}$$

Thus the following holds:

$$\begin{aligned}
2 \frac{\partial V_l}{\partial l_1} &= \frac{\sqrt{-1}}{2} \frac{\partial}{\partial l_1} \left\{ \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_-) - z_- \frac{\partial U}{\partial z} \Big|_{z=z_-} \log z_- \right) \right. \\
&\quad \left. - \left( U(a_1, a_2, a_3, a_4, a_5, a_6, z_+) - z_+ \frac{\partial U}{\partial z} \Big|_{z=z_+} \log z_+ \right) \right\} \\
&\equiv \frac{\sqrt{-1}}{2} \left( \log \frac{\varphi(z_-)}{\psi(z_-)} - \log \frac{\varphi(z_+)}{\psi(z_+)} \right) \pmod{\pi} \\
&\equiv \frac{\sqrt{-1}}{2} \log \frac{\varphi(z_-) \psi(z_+)}{\varphi(z_+) \psi(z_-)} \pmod{2\pi}.
\end{aligned}$$



Now

$$\begin{aligned}\varphi(z_-)\psi(z_+) &= \frac{4a_1a_2^2a_3^2a_4^4a_5^2a_6^2}{q_2^2}(a_1a_2+a_3)(a_2+a_1a_3) \times \\ &\quad (a_1a_5+a_6)(a_5+a_1a_6)\left(c_{l_{12}}-\sqrt{-1}\sqrt{c_{l_{11}}c_{l_{22}}-c_{l_{12}}^2}\right), \\ \varphi(z_+)\psi(z_-) &= \overline{\varphi(z_-)\psi(z_+)},\end{aligned}$$

so we have

$$\begin{aligned}2\frac{\partial V_l}{\partial l_1} &\equiv \frac{\sqrt{-1}}{2}\log\frac{\varphi(z_-)\psi(z_+)}{\varphi(z_+)\psi(z_-)} \pmod{\pi} \\ &= -\arg\varphi(z_-)\psi(z_+) \\ &= \arctan\frac{\sqrt{c_{l_{11}}c_{l_{22}}-c_{l_{12}}^2}}{c_{l_{12}}} \\ &= A_1,\end{aligned}$$

where we use (3.4) at the last equality. Thus we have proved Proposition 4.3.  $\square$

We finally recall the so-called Schläfli's differential formula:

**Proposition 4.4** (see, for example, [Mi]). *Let  $T$  be a hyperbolic tetrahedron with dihedral angle  $A_i$  and edge length  $l_i$ . Then the following equation holds:*

$$d\text{Vol}(T) = -\frac{1}{2}\sum_{i=1}^6 l_i dA_i.$$

*Proof of Theorem 2.2.* Let  $W = W(T)$  be the right-hand side of the formula. Since  $W(T)$  is represented by the edge lengths symmetrically, and the edge lengths can be represented by the dihedral angles symmetrically as in (3.3), all we need to show are, by the Schläfli's differential formula, the following two facts:

1.  $\frac{\partial W}{\partial A_1} = -\frac{l_1}{2},$
2. At some value  $T$  of a hyperbolic tetrahedron,  $W(T)$  is actually its volume.

Using Proposition 4.3, the former one can be shown as follows:

$$\begin{aligned}
\frac{\partial W}{\partial A_1} &= \frac{\partial V_l}{\partial A_1} - \sum_{i=1}^6 \frac{\partial}{\partial A_1} \left( l_i \frac{\partial V_l}{\partial l_i} \right) \\
&= \frac{\partial V_l}{\partial A_1} - \sum_{i=1}^6 \left( \frac{\partial l_i}{\partial A_1} \frac{\partial V_l}{\partial l_i} + l_i \frac{\partial}{\partial A_1} \frac{\partial V_l}{\partial l_i} \right) \\
&= \frac{\partial V_l}{\partial A_1} - \sum_{i=1}^6 \frac{\partial V_l}{\partial l_i} \frac{\partial l_i}{\partial A_1} - \sum_{i=1}^6 \frac{l_i}{2} \frac{\partial}{\partial A_1} (A_i + k_i \pi) \quad (k_i \in \mathbb{Z}) \\
&= -\frac{l_1}{2}.
\end{aligned}$$

Since both  $W(T)$  and  $\text{Vol}(T)$  are analytic function with variables  $l_1, l_2, \dots, l_6$ , the result obtained above means that the difference between  $W(T)$  and  $\text{Vol}(T)$  is a constant independent of  $l_1, l_2, \dots, l_6$ .

For the proof of the later one, let  $T_\rho$  be a hyperbolic regular tetrahedron with all edge lengths being  $\rho$ , and we consider a sequence of such tetrahedra shrinking to a point. Then clearly  $\lim_{\rho \rightarrow 0} \text{Vol}(T_\rho) = 0$ .

On the other hand, we have the following equations in this case:

$$\begin{aligned}
z_\pm &= \frac{3(\cosh \rho + 1) \pm \sqrt{-1} \sqrt{(\cosh \rho - 1)(3 \cosh \rho + 1)}}{2 e^{4\rho} (2 \cosh \rho - \sinh \rho + 1)}, \\
U &= \text{Li}_2(z_j) + 3 \text{Li}_2(e^{4\rho} z_j) - 4 \text{Li}_2(e^{3\rho} z_j), \\
z_j \frac{\partial U}{\partial z} \Big|_{z=z_j} &= -\sqrt{-1} \{ \arg(1 - z_j) + 3 \arg(1 - e^{4\rho} z_j) - 4 \arg(1 - e^{3\rho} z_j) \}, \\
\frac{\partial V_l}{\partial \rho} &= \frac{1}{2} \left( \arctan \frac{\sqrt{(\cosh \rho + 1)(3 \cosh \rho + 1)}}{\cosh \rho} + k \pi \right) \quad (k \in \mathbb{Z}),
\end{aligned}$$

where the third equation follows from Proposition 4.1, and the last one from Proposition 4.3. Thus the following equation holds:

$$\begin{aligned}
W(T_\rho) &= \frac{\sqrt{-1}}{4} \left\{ \operatorname{Li}_2(z_-) - \operatorname{Li}_2(z_+) \right\} + 3 \left\{ \operatorname{Li}_2(e^{4\rho} z_-) - \operatorname{Li}_2(e^{4\rho} z_+) \right\} \\
&\quad - 4 \left\{ \operatorname{Li}_2(e^{3\rho} z_-) - \operatorname{Li}_2(e^{3\rho} z_+) \right\} \\
&\quad + \sqrt{-1} \left\{ \arg(1 - z_-) + 3 \arg(1 - e^{4\rho} z_-) - 4 \arg(1 - e^{3\rho} z_-) \right\} \log z_- \\
&\quad - \sqrt{-1} \left\{ \arg(1 - z_+) + 3 \arg(1 - e^{4\rho} z_+) - 4 \arg(1 - e^{3\rho} z_+) \right\} \log z_+ \Big\} \\
&\quad - 3\rho \left( \arctan \frac{\sqrt{(\cosh \rho + 1)(3 \cosh \rho + 1)}}{\cosh \rho} + k\pi \right) \\
&= -\frac{1}{2} \left\{ \Im \left\{ \operatorname{Li}_2(z_-) + 3 \operatorname{Li}_2(e^{4\rho} z_-) - 4 \operatorname{Li}_2(e^{3\rho} z_-) \right\} \right. \\
&\quad \left. + \left\{ \arg(1 - z_-) + 3 \arg(1 - e^{4\rho} z_-) - 4 \arg(1 - e^{3\rho} z_-) \right\} \log |z_-| \right\} \\
&\quad - 3\rho \left( \arctan \frac{\sqrt{(\cosh \rho + 1)(3 \cosh \rho + 1)}}{\cosh \rho} + k\pi \right),
\end{aligned}$$

where we used the fact  $z_+ = \overline{z_-}$  in the last equation.

When  $\rho$  converges to 0, then  $z_-$  converges to 1, a branch point of  $\operatorname{Li}_2(z)$ . However this convergence is not spiral because  $\Im z_- \neq 0$  for any  $\rho > 0$ . So we have

$$\lim_{\rho \rightarrow 0} \Im \left\{ \operatorname{Li}_2(z_-) + 3 \operatorname{Li}_2(e^{4\rho} z_-) - 4 \operatorname{Li}_2(e^{3\rho} z_-) \right\} = 0.$$

Furthermore we have

$$\begin{aligned}
\arg(1 - z_-) &= \arctan \frac{-\sqrt{3e^{2\rho} + 2e^\rho + 3}}{2e^{5\rho} + 6e^{4\rho} + 12e^{3\rho} + 12e^{2\rho} + 9e^\rho + 3}, \\
\arg(1 - e^{4\rho} z_-) &= \arctan \frac{\sqrt{3e^{2\rho} + 2e^\rho + 3}}{e^\rho + 3}, \\
\arg(1 - e^{3\rho} z_-) &= \arctan \frac{-\sqrt{3e^{2\rho} + 2e^\rho + 3}}{2e^{2\rho} + 3e^\rho + 3}.
\end{aligned}$$

Thus we conclude as follows:

$$\lim_{\rho \rightarrow 0} \left\{ \arg(1 - z_-) + 3 \arg(1 - e^{4\rho} z_-) - 4 \arg(1 - e^{3\rho} z_-) \right\} \log |z_-| = 0.$$

Finally we have

$$\lim_{\rho \rightarrow 0} \frac{\sqrt{(\cosh \rho + 1)(3 \cosh \rho + 1)}}{\cosh \rho} = 2\sqrt{2}$$

and  $\arctan 2\sqrt{2}$  is actually the dihedral angle of the Euclidean regular tetrahedron.

Thus we have  $\lim_{\rho \rightarrow 0} W(\rho) = 0$  and finished the proof of Theorem 2.2.  $\square$

## References

- [AVS] D. V. Alekseevskij, È. B. Vinberg and A. S. Solodovnikov, *Geometry of Spaces of Constant Curvature*, in *Geometry II*, 1–138, edited by E. B. Vinberg, Encyclopaedia of Mathematical Sciences **29**, Springer-Verlag, Berlin, 1993.
- [CK] Yunhi Cho and Hyuk Kim, *On the Volume Formula for Hyperbolic Tetrahedra*, Discrete & Computational Geometry **22** (1999), 347–366.
- [DL] Peter Doyle and Gregory Leibon, *23040 symmetries of hyperbolic tetrahedra*, arXiv:math.GT/0309187.
- [Ka] R. M. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Letters in Mathematical Physics **39** (1997), 269–275.
- [Lu] Feng Luo, *On a Problem of Fenchel*, Geometriae Dedicata **64** (1997), 277–282.
- [Mi] John Milnor, *The Schläfli differential equality*, in *John Milnor Collected Papers Volume 1, Geometry*, 281–295, Publish or Perish, Inc., Houston and Texas, 1994.
- [Mu] Hitoshi Murakami, *Optimistic calculations about the Witten-Reshetikhin-Turaev invariants of closed three-manifolds obtained from the figure-eight knot by integral Dehn surgeries*, in *Recent progress towards the volume conjecture*, Sūrikaiseikikenkyūsho Kōkyūroku **1172** (2000), 70–79.
- [MY] Jun Murakami and Masakazu Yano, *On the volume of a hyperbolic and spherical tetrahedron*, available at <http://www.f.waseda.jp/murakami/papers/tetrahedronrev3.pdf>.
- [SA] Luis A. Santaló, *Integral geometry and geometric probability*, Encyclopedia of Mathematics and its Applications **1** (1976), Addison-Wesley Publishing Co.
- [TV] V. G. Turaev and O. Ya. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology **31** (1992), 865–902.
- [Us] Akira Ushijima, *A volume formula for generalized hyperbolic tetrahedra*, arXiv:math.GT/0309216.