

Fujita's very ampleness conjecture for singular toric varieties

Sam Payne

February 28, 2019

Abstract

We prove a generalization of Fujita's effective very ampleness conjecture for toric varieties with arbitrary singularities, using combinatorial methods.

Given an ample divisor D and any other Cartier divisor D' on an algebraic variety, we can choose t sufficiently large so that $tD + D'$ is basepoint free or very ample. In either case, it is not easy say how large we must choose t in general. However, for $D' = K_X$, Fujita made the following conjectures.

Fujita's Conjectures *Let X be an n -dimensional projective algebraic variety, smooth or with mild singularities, D an ample divisor on X .*

- (i) *For $t \geq n + 1$, $tD + K_X$ is basepoint free.*
- (ii) *For $t \geq n + 2$, $tD + K_X$ is very ample.*

The example $X \cong \mathbb{P}^n$, $D \sim H$ shows that Fujita's conjectured effective bounds are best possible.

For smooth varieties, the corresponding statements with "basepoint free" and "very ample" replaced by "nef" and "ample", respectively, are consequences of Mori's Cone Theorem [Fuj]. For divisors on smooth *toric* varieties, nefness and ampleness are equivalent to freeness and very ampleness, respectively, so Fujita's conjectures follow immediately for smooth toric varieties. One can also deduce Fujita's conjectures for smooth toric varieties by general (non-toric) cohomological arguments of Ein and Lazarsfeld in characteristic zero [EL], and Smith in positive characteristic [Sm1] [Sm2], again using the fact that ample divisors on smooth toric varieties are very ample.

For toric varieties with arbitrary singularities, a strong generalization of Fujita's basepoint freeness conjecture was proved by Fujino [Fu]. We follow the usual toric convention fixing $K_X = -\sum D_i$, the sum of the T -invariant prime divisors each with coefficient -1, as a convenient representative of the canonical class.

Fujino's Theorem [Fu] *Let X be a projective n -dimensional toric variety not isomorphic to \mathbb{P}^n . Let D, D' be \mathbb{Q} -Cartier divisors such that $0 \geq D' \geq K_X$,*

$D + D'$ is Cartier, and $D \cdot C \geq n$ for all T -curves C . Then $D + D'$ is basepoint free.

Fujita's basepoint freeness conjecture for toric varieties is the special case of Fujino's Theorem when X is Gorenstein, $D' = K_X$, and $D = tL$ for some ample Cartier divisor L and some integer $t \geq n + 1$. In fact, Fujino's Theorem shows that, for toric varieties, Fujita's conjectured bound can be improved by excluding the extremal case $X \cong \mathbb{P}^n$.

Of course, the case of \mathbb{P}^n can be analyzed separately. The canonical divisor on \mathbb{P}^n is linearly equivalent to $(-n-1)H$, so $D' \sim sH$ for some $0 \geq s \geq (-n-1)$. Any \mathbb{Q} -Cartier divisor D on \mathbb{P}^n is linearly equivalent to tH for some $t \in \mathbb{Q}$. Then $D + D'$ is Cartier exactly when $t + s$ is an integer and basepoint free exactly when $t + s$ is a nonnegative integer.

The main purpose of this paper is to prove an analogous strong generalization of Fujita's very ampleness conjecture for toric varieties with arbitrary singularities.

Theorem 1 *Let X be a projective n -dimensional toric variety not isomorphic to \mathbb{P}^n . Let D, D' be \mathbb{Q} -Cartier divisors such that $0 \geq D' \geq K_X$, $D + D'$ is Cartier, and $D \cdot C \geq n + 1$ for all T -curves C . Then $D + D'$ is very ample.*

The statement of Fujino's Theorem can be strengthened by removing the assumption that $D + D'$ is Cartier. A Cartier divisor on a toric variety is basepoint free if and only if it is nef, i.e. if and only if D intersects every curve nonnegatively [La, Proposition 1.5]. Without the hypothesis that $D + D'$ is Cartier, the sharper statement of Fujino's theorem is then:

Fujino's Theorem⁺ [Fu] *Let X be a projective n -dimensional toric variety, not isomorphic to \mathbb{P}^n . Let D, D' be \mathbb{Q} -Cartier divisors such that $0 \geq D' \geq K_X$ and $D \cdot C \geq n$ for all T -curves C . Then $D + D'$ is nef.*

Similarly, the statement of Theorem 1 can be strengthened by using a toric characterization of very ampleness to remove the hypothesis that $D + D'$ is Cartier. Since every divisor on a toric variety is linearly equivalent to a T -invariant divisor, we may assume D and D' are \mathbb{Q} -linear combinations of T -invariant divisors. To state the stronger theorem, we need some notation from toric geometry.

Let $D = \sum d_i D_i$ be a T - \mathbb{Q} -Cartier divisor on a complete toric variety X . For each maximal cone σ in the fan defining X , we have a point $u_\sigma \in M_{\mathbb{Q}}$ determined by the conditions $\langle u_\sigma, v_i \rangle = -d_i$ for each of the primitive generators v_i of the rays of σ . When D and D' denote T - \mathbb{Q} -Cartier divisors, we will write u_σ and u'_σ for the points of $M_{\mathbb{Q}}$ associated to D and D' , respectively. The association $D \rightsquigarrow u_\sigma$ is linear, i.e. $tD \rightsquigarrow tu_\sigma$ and $D + D' \rightsquigarrow \{u_\sigma + u'_\sigma\}$. A T - \mathbb{Q} -Cartier divisor D is Cartier if and only if $u_\sigma \in M$ for all maximal cones σ . Also associated to D is a polytope $P_D \subset M_{\mathbb{Q}}$ cut out by the inequalities $\langle u, v_i \rangle \geq -d_i$ for all of the primitive generators v_i of the rays of the fan. When D is T - \mathbb{Q} -Cartier and nef, the $\{u_\sigma\}$ are the vertices of P_D . If we translate P_D

so that the vertex u_σ is at the origin, then all of P_D sits inside the dual cone σ^\vee . Write P_D^σ for this translation, i.e. $P_D^\sigma := P_D - u_\sigma$. In case D is Cartier, then D is very ample if and only if $P_D^\sigma \cap M$ generates the semigroup $\sigma^\vee \cap M$ for all maximal cones σ . Without the hypothesis that $D + D'$ is Cartier, the stronger version of Theorem 1 we will prove is then:

Theorem 2 *Let X be a projective n -dimensional toric variety not isomorphic to \mathbb{P}^n . Let D, D' be T - \mathbb{Q} -Cartier divisors such that $0 \geq D' \geq K_X$ and $D \cdot C \geq n + 1$ for all T -curves C . Then $P_{D+D'}^\sigma \cap M$ generates $\sigma^\vee \cap M$ for all maximal cones σ .*

Theorem 1 is the special case of Theorem 2 when $D + D'$ is Cartier.

Our approach starts from an observation made by Laterveer in [La]: if D is ample, then the lattice length of the edge of P_D corresponding to a T -curve C is precisely $D \cdot C$. This fact can be seen as a consequence of Riemann-Roch for toric varieties [Ful, p.112]. Adding D' corresponds to moving the faces of P_D inward at most a unit distance with respect to the dual lattice. When all of the edges of P_D have lattice length at least $n + 1$, we show that $P_{D+D'}^\sigma$ contains an explicit generating set for $\sigma^\vee \cap M$ for all maximal cones σ . The computations are straightforward in the simplicial case, as can be seen in the example at the end of the introduction.

In the proof of Fujino's Theorem, there is a simple reduction to the simplicial case, or even to the smooth case (see [La, Lemma 2.4], [Fu, 1.12, Step 2], or [P]). The reduction works via a partial, or full, projective resolution of singularities corresponding to a triangulation of the fan defining X . This type of reduction seems not to work for very ampleness. Instead, for each nonsimplicial maximal cone σ , we make a canonical subdivision of the dual cone σ^\vee .

Remarks – The earliest results on Fujita's freeness conjecture for singular toric varieties of which the author is aware are due to Laterveer. In [La], Laterveer proved Fujino's Theorem for \mathbb{Q} -Gorenstein toric varieties when $D' = K_X$ using toric Mori theory, as developed in [Re]. Our statement of Theorem 1, like the statement of Fujino's Theorem, is influenced by Mustață's formulations in [Mu]. In particular, Mustață stated and proved Fujino's Theorem and Theorem 1 for smooth toric varieties when D and D' are Cartier as consequences of a characteristic-free vanishing theorem for toric varieties. For proofs of Fujino's Theorem that do not use vanishing theorems or toric Mori theory, see also [Lin] and [P].

The only previous result on Fujita's very ampleness conjecture for singular toric varieties of which the author is aware is due to Lin,¹ who proved the conjecture for simplicial Gorenstein toric varieties in dimension ≤ 6 [Lin]. In [La], Laterveer also claimed to prove a generalization of Fujita's very ampleness conjecture for arbitrary \mathbb{Q} -Gorenstein toric varieties. As noted by Lin, there

¹Another effective very ampleness result for singular toric varieties, due to Ewald and Wessels [EW], may be stated as follows: let X be an n -dimensional projective toric variety and D a T - \mathbb{Q} -Cartier divisor on X such that $D \cdot C \geq n - 1$ for all T -curves C . Then $P_D^\sigma \cap M$ generates $\sigma^\vee \cap M$ for all maximal cones σ . In particular, if D is Cartier, then D is very ample.

is an error in the proof of this claim. In particular, it is not true in general that P_{tD+K_X} contains $P_{(t-1)D}$. The case $X \cong \mathbb{P}^n, D \sim H$ is a counterexample. Nevertheless, Laterveer's approach to Fujita's very ampleness conjecture for singular toric varieties contains fruitful insights, in particular, the realization that \mathbb{P}^n is the only toric extremal case and the characterization of the intersection numbers $D \cdot C$ as the lattice lengths of the edges of P_D . The results we prove here are strong enough to imply all of the very ampleness results claimed in [La].

Example We illustrate the essential techniques of this paper in a concrete simplicial example. Let u_1, \dots, u_n be linearly independent primitive vectors in M . Let P be the simplex with vertices $\{0, u_1, \dots, u_n\}$. Associated to P , there is a projective toric variety X_P with an ample divisor D such that $P = P_D$ [Ful, Section 1.5]. The vertex 0 of P corresponds to a maximal cone σ of the fan defining X_P whose dual cone σ^\vee is spanned by $\{u_1, \dots, u_n\}$. Let D_1, \dots, D_n be the divisors corresponding to the rays of σ , and let $D' = -D_1 - \dots - D_n$. We will show that, for $t \geq n + 1$, $P_{tD+D'}^\sigma \cap M$ generates $\sigma^\vee \cap M$.

Every point in M can be written uniquely as an integer linear combination of the $\{u_i\}$ plus a fractional part. So the semigroup $\sigma^\vee \cap M$ is generated by $\{0, u_1, \dots, u_n\}$ together with $\{(a_1 u_1 + \dots + a_n u_n) \in M : 0 \leq a_i < 1\}$. For $t \geq n + 1$, we will show that $P_{tD+D'}^\sigma$ contains this generating set.

Define a linear function λ on $M_\mathbb{Q}$ by

$$\lambda(a_1 u_1 + \dots + a_n u_n) = a_1 + \dots + a_n.$$

Note that $P_{tD} = \{u \in \sigma^\vee : \lambda(u) \leq t\}$. In other words, if $\{v_i\}$ are the primitive generators of the rays of σ , then P_{tD} is cut out by the conditions $\langle u, v_i \rangle \geq 0$ and the condition $\lambda(u) \leq t$. Similarly, $P_{tD+D'} = \{u'_\sigma + u : u \in \sigma^\vee, \lambda(u + u'_\sigma) \leq t\}$, i.e. $P_{tD+D'}$ is cut out by the conditions $\langle u, v_i \rangle \geq \langle u'_\sigma, v_i \rangle = 1$ and the condition $\lambda(u) \leq t$. It follows that any lattice point in P_{tD} that is in the interior of σ^\vee is contained in $P_{tD+D'}$. Indeed, if u is in P_{tD} then $\lambda(u) \leq t$, and if u is a lattice point in the interior of σ^\vee then $\langle u, v_i \rangle$ is a positive integer.

Suppose $t \geq n + 1$. Then $u_1 + \dots + u_n$ is a lattice point in P_{tD} that is in the interior of σ^\vee , so $u_1 + \dots + u_n$ is contained in $P_{tD+D'}$. Note that u'_σ is the point of $P_{tD+D'}$ for which λ achieves its minimum. In particular, $\lambda(u'_\sigma) \leq \lambda(u_1 + \dots + u_n) = n$.

For each u_i , we have $\lambda(u'_\sigma + u_i) = \lambda(u'_\sigma) + 1 \leq n + 1 \leq t$. Therefore $u'_\sigma + u_i \in P_{tD+D'}$, i.e. $u_i \in P_{tD+D'}^\sigma$. Given a lattice point p of the form $p = a_1 u_1 + \dots + a_n u_n$ with $0 \leq a_i < 1$, we have another lattice point $p' = (1 - a_1)u_1 + \dots + (1 - a_n)u_n$ in P_{tD} that is in the interior of σ^\vee . So p' is contained in $P_{tD+D'}$, and therefore

$$\lambda(u'_\sigma) \leq \lambda(p') = n - \lambda(p).$$

So $\lambda(u'_\sigma + p) \leq n < t$, and hence $p \in P_{tD+D'}^\sigma$, as required.

I wish to thank M. Hering, P. Horja, R. Lazarsfeld, and M. Mustața for helpful conversations related to this work. I am especially grateful to W. Fulton

for his encouragement on this project and for his comments and suggestions on earlier drafts of this paper.

1 Preliminaries

As a first step to proving Theorem 2, we have:

Lemma 1 *Let $X, D,$ and D' satisfy the hypotheses of Theorem 2. Let σ be a maximal cone, and let $\{u_1, \dots, u_s\}$ be the primitive generators of the rays of σ^\vee . Then $P_{D+D'}^\sigma$ contains $\{0, u_1, \dots, u_s\}$.*

Proof: By Fujino's Theorem⁺, $\frac{n}{n+1}D + D'$ is nef. Therefore, for any T -curve C ,

$$(D + D') \cdot C = \frac{1}{n+1}(D \cdot C) + \left(\frac{n}{n+1}D + D'\right) \cdot C \geq 1.$$

By Laterveer's observation, this means that every edge of $P_{D+D'}$ has lattice length at least 1. Translating the vertex u_σ to the origin, it follows that $P_{D+D'}^\sigma$ contains 0 and the primitive generators of each of the rays of σ^\vee . \square

If σ is regular, then $s = n$ and $\{0, u_1, \dots, u_n\}$ generates $\sigma^\vee \cap M$, so the conclusion of Theorem 2, i.e. the fact that $P_{D+D'}^\sigma \cap M$ generates $\sigma^\vee \cap M$, follows immediately. In general, if we let $\Delta = \text{conv}\{0, u_1, \dots, u_s\}$, then Lemma 1 says that $P_{D+D'}^\sigma$ contains Δ . If σ is not regular, then Δ may not contain a generating set for $\sigma^\vee \cap M$. The following example, due to Ewald and Wessels [EW], illustrates this possibility.

Example Let $M = \mathbb{Z}^3$; $u_1 = (1, 0, 0)$, $u_2 = (0, 1, 0)$, and $u_3 = (1, 1, 2)$. Let σ^\vee be the cone spanned by $\{u_1, u_2, u_3\}$, so $\Delta = \text{conv}\{0, u_1, u_2, u_3\}$. Then $\Delta \cap M = \{0, u_1, u_2, u_3\}$, so the semigroup generated by $\Delta \cap M$ only contains lattice points whose third coordinate is even. In particular, the lattice point $(1, 1, 1) = \frac{1}{2}(u_1 + u_2 + u_3)$ is in σ^\vee , but not in the semigroup generated by $\Delta \cap M$.

Although Δ may not contain a generating set for $\sigma^\vee \cap M$, we will show that $P_{D+D'}^\sigma$ contains a dilation of Δ that does contain a generating set. Let $m = \min\{(D + D') \cdot V(\sigma \cap \tau)\}$, where τ varies over all maximal cones adjacent to σ , so that m is the minimum of the lattice lengths of the edges of $P_{D+D'}^\sigma$ incident to the vertex 0. Note that $m\Delta$ is the largest rational dilation of Δ contained in $P_{D+D'}^\sigma$. We will show that $m\Delta$ does contain a generating set for $\sigma^\vee \cap M$.

In preparation for proving this, we develop a few preliminaries. First, we generalize Laterveer's observation on the lattice lengths of the edges of P_D to the case where D is not necessarily ample.

Lemma 2 *Let X be a complete toric variety, D a T - \mathbb{Q} -Cartier divisor on X . Let σ, τ be adjacent maximal cones in the fan defining X , and let u be the primitive generator of the ray of σ^\vee perpendicular to $\sigma \cap \tau$. Then*

$$u_\tau = u_\sigma + (D \cdot V(\sigma \cap \tau))u.$$

Proof: Since u_σ and u_τ agree on $\sigma \cap \tau$, their difference must vanish on $\sigma \cap \tau$, i.e. $u_\tau - u_\sigma = ku$ for some rational number k . By the toric intersection formulas in [Ful, Section 5.1], for v_j the primitive generator of any ray of τ not contained in σ ,

$$D \cdot V(\sigma \cap \tau) = \frac{\langle u_\sigma - u_\tau, v_j \rangle}{-\langle u, v_j \rangle}.$$

Therefore,

$$D \cdot V(\sigma \cap \tau) = \frac{\langle ku, v_j \rangle}{\langle u, v_j \rangle} = k.$$

□

Now we develop some tools for working with rational cones. Let σ^\vee be a rational cone, and let u_1, \dots, u_s be the primitive generators of the rays of σ^\vee . Define a function λ^{\min} on σ^\vee by

$$\lambda^{\min}(u) = \min\{(a_1 + \dots + a_s) : a_1 u_1 + \dots + a_s u_s = u, a_i \geq 0\}.$$

Define λ^{\max} similarly. A few combinatorial properties of λ^{\min} and λ^{\max} , all of them immediate from the definitions, will be useful in what follows.

First, λ^{\min} and λ^{\max} are anticonvex and convex, respectively. In other words, for any $u, u' \in \sigma^\vee$,

$$\lambda^{\min}(u + u') \leq \lambda^{\min}(u) + \lambda^{\min}(u'),$$

and similarly $\lambda^{\max}(u + u') \geq \lambda^{\max}(u) + \lambda^{\max}(u')$.

Second, suppose $D' = \sum d'_i D_i$ is minus-effective, i.e. $d'_i \leq 0$ for all i . Then for any primitive generator v_i of a ray of σ , $\langle u'_\sigma, v_i \rangle = -d'_i \geq 0$. So u'_σ is in the dual cone σ^\vee . In particular, $\lambda^{\min}(u'_\sigma)$ and $\lambda^{\max}(u'_\sigma)$ are well-defined.

Finally, with $\Delta = \text{conv}\{u_1, \dots, u_s\}$, note that

$$m\Delta = \{u \in \sigma^\vee : \lambda^{\min}(u) \leq m\}.$$

The distinction between λ^{\min} and λ^{\max} is meaningful only in the nonsimplicial case; when σ is simplicial, then the primitive generators of the rays of σ^\vee are linearly independent, so the expression $u = a_1 u_1 + \dots + a_n u_n$ is unique.

In order to show that $m\Delta$ contains a generating set for $\sigma^\vee \cap M$, one seeks lower bounds for m . To get a rough idea of how one might get such bounds, imagine that P_D is very large, as it will be under the hypotheses of Theorem 2. When we add a small, minus-effective divisor D' to D , we get $P_{D+D'}$ by moving the faces of P_D inward a small distance. In particular, the new vertex $u_\sigma + u'_\sigma$ of $P_{D+D'}$ will be inside P_D and close to the old vertex u_σ of P_D . We can measure this closeness by $\lambda^{\min}(u'_\sigma)$. Assume that for some large positive t , P_D contains $u_\sigma + t\Delta$ (i.e. $D \cdot V(\sigma \cap \tau) \geq t$ for all maximal cones τ adjacent to σ). Then $t\Delta$ contains u'_σ and $u'_\sigma + (t - \lambda^{\min}(u'_\sigma))u_i$ for each of the primitive generators u_i of the rays of σ^\vee . Therefore, if the faces of P_D not containing u_σ did not move inwards at all (i.e. if D' were supported on the divisors corresponding to

the rays of σ), then $P_{D+D'}$ would contain $u_\sigma + u'_\sigma + (t - \lambda^{\min}(u'_\sigma))\Delta$. Taking into account the movement of the other faces, we would see that $P_{D+D'}$ still contains $u_\sigma + u'_\sigma + (t - \lambda^{\min}(u'_\sigma) - r)\Delta$, where r is the maximum distance with respect to the dual lattice that the other faces move inward. The key to giving lower bounds for m will be the following proposition, which makes the essence of the preceding discussion precise.

Proposition *Let X be a complete toric variety, and let σ be a maximal cone in the fan defining X . Let D, D' be T - \mathbb{Q} -Cartier divisors such that D is nef and $0 \geq D' \geq rK_X$, for some positive rational number r . Let $t = \min\{D \cdot V(\sigma \cap \tau)\}$ and $m = \min\{(D + D') \cdot V(\sigma \cap \tau)\}$, where τ varies over all maximal cones adjacent to σ . Suppose $t \geq \lambda^{\min}(u'_\sigma)$. Then*

$$m \geq t - \lambda^{\min}(u'_\sigma) - r.$$

Proof: Let τ be a maximal cone adjacent to σ . Let $k = (D + D') \cdot V(\sigma \cap \tau)$, and let $c = t - \lambda^{\min}(u'_\sigma)$. In order to prove the proposition, we must show that $c - k \leq r$. Recall that $u_\sigma + u'_\sigma$ and $u_\tau + u'_\tau$ are the points of $M_{\mathbb{Q}}$ associated to $D + D'$ for σ and τ , respectively. Let u be the primitive generator of the ray of σ^\vee perpendicular to $\sigma \cap \tau$. By Lemma 2

$$u_\tau + u'_\tau = u_\sigma + u'_\sigma + ku.$$

Let v_j be the primitive generator of any ray of τ not in σ . Say $D = \sum d_i D_i$ and $D' = \sum d'_i D_i$. Then

$$\langle u_\sigma + u'_\sigma + ku, v_j \rangle = \langle u_\tau + u'_\tau, v_j \rangle = -d_j - d'_j. \quad (1)$$

Now c is nonnegative by hypothesis, so $cu \in \sigma^\vee$. Therefore, since λ^{\min} is anticonvex and $\lambda^{\min}(cu) = c$,

$$\lambda^{\min}(u'_\sigma + cu) \leq \lambda^{\min}(u'_\sigma) + c = t.$$

Hence $u'_\sigma + cu$ is contained in $t\Delta$.

Since D is nef, P_D contains $u_\sigma + t\Delta$. In particular, P_D contains $u_\sigma + u'_\sigma + cu$. Therefore,

$$\langle u_\sigma + u'_\sigma + cu, v_j \rangle \geq -d_j. \quad (2)$$

Subtracting (2) from (1) we see that $(k - c)\langle u, v_j \rangle \leq -d'_j$. Now, since $D' \geq rK_X$, $-d'_j$ is at most r . Since $\langle u, v_j \rangle \leq -1$, it follows that $c - k \leq r$. \square

Remark – The conclusion of the proposition is false in general if $t < \lambda^{\min}(u'_\sigma)$. Consider, for example, the complete toric surface X whose fan is spanned by three rays, the primitive generators of which satisfy $v_1 + v_2 + 2v_3 = 0$ (i.e. X is the weighted projective plane $\mathbb{P}(1, 1, 2)$). Let σ be the cone spanned by v_2 and v_3 . Taking Cartier divisors $D = 2D_1$ and $D' = 2K_X$, one computes $t = D \cdot D_2 = 1$, $\lambda^{\min}(u'_\sigma) = 4$, and $(D + D') \cdot D_2 = -6$, which is strictly less than $1 - 4 - 2$.

The proposition gives good lower bounds for m , provided we can give good upper bounds for $\lambda^{\min}(u'_\sigma)$. We get sufficient bounds indirectly by using convexity to bound $\lambda^{\max}(u'_\sigma)$.

Lemma 3 *Let D' be T - \mathbb{Q} -Cartier, with $0 \geq D' \geq K_X$. Then $\lambda^{\max}(u'_\sigma) \leq \lambda^{\max}(u)$ for any lattice point u in the interior of σ^\vee .*

Proof: For any lattice point u in the interior of σ^\vee , and for the primitive generator v_j of any ray of σ , $\langle u, v_j \rangle$ is a positive integer, and in particular is at least 1. On the other hand, $\langle u'_\sigma, v_j \rangle = -d'_j$, which, since $D' \geq K_X$, is at most 1. Therefore $u - u'_\sigma$ is in σ^\vee . Since λ^{\max} is convex and nonnegative on σ^\vee , it follows that $\lambda^{\max}(u'_\sigma) \leq \lambda^{\max}(u)$. \square

2 Proof of Theorem 2

Let X, D , and D' satisfy the hypotheses of Theorem 2, and let σ be a maximal cone in the fan defining X . Let $m = \min\{(D+D') \cdot V(\sigma \cap \tau)\}$, where τ varies over all maximal cones adjacent to σ . Let $\{u_1, \dots, u_s\}$ be the primitive generators of the rays of σ^\vee , and let $\Delta = \text{conv}\{0, u_1, \dots, u_s\}$. To prove Theorem 2, it will suffice to show that $m\Delta$ contains a generating set for $\sigma^\vee \cap M$. To prove this, we will give a canonical subdivision of σ^\vee and show that, for each maximal cone γ of the subdivision, $\gamma \cap m\Delta$ contains a generating set for $\gamma \cap M$.

We claim that λ^{\max} is piecewise-linear and therefore defines a canonical subdivision of σ^\vee : the subdivision whose maximal cones are the maximal subcones of σ^\vee on which λ^{\max} is linear. This subdivision can also be realized by looking at $Q = \text{conv}\{u_1, \dots, u_s\}$ and taking the cones over the ‘‘lower faces’’ of Q , i.e. the faces of Q visible from the vertex 0 of σ^\vee . Indeed, for any $t > 0$, tQ is the set of u in σ^\vee that can be written $u = a_1u_1 + \dots + a_su_s$ with $a_i \geq 0$ and $a_1 + \dots + a_s = t$. Now the points in the lower faces of Q are precisely those points that are not contained in tQ for any $t > 1$. So the restriction of λ^{\max} to the lower faces of Q is identically 1. Since $\lambda^{\max}(cu) = c\lambda^{\max}(u)$ for any $c \geq 0$, it follows that λ^{\max} is linear precisely on the cones over the lower faces of Q .

Let γ be the cone over a maximal lower face of Q , and let $\gamma^{(1)} \subset \{u_1, \dots, u_s\}$ denote the set of primitive generators of the rays of γ . We must show that $\gamma \cap m\Delta$ contains a generating set for $\gamma \cap M$. Every point of γ can be written as a nonnegative linear combination:

$$u = a_1u_{i_1} + \dots + a_nu_{i_n},$$

where $a_j \geq 0$, and $\{u_{i_j}\} \subset \gamma^{(1)}$ is linearly independent. This expression can be decomposed as a nonnegative integer combination of the $\{u_{i_j}\}$ plus a nonnegative fractional part. So $\gamma \cap M$ is generated by 0 and $\gamma^{(1)}$ together with $\{(a_1u_{i_1} + \dots + a_nu_{i_n}) \in M : 0 \leq a_j < 1, \{u_{i_j}\} \subset \gamma^{(1)} \text{ linearly independent}\}$. By Lemma 1, $m\Delta$ contains 0 and $\gamma^{(1)}$. It will therefore suffice to show that any lattice point p that is a nonnegative fractional linear combination of some

independent set $\{u_{i_j}\} \subset \gamma^{(1)}$ is contained in $m\Delta$. For this, it will suffice to show that $m \geq \lambda^{\max}(p)$.

Suppose $p = a_1 u_{i_1} + \cdots + a_n u_{i_n} \in M$, where $0 \leq a_j < 1$, and $\{u_{i_j}\} \subset \gamma^{(1)}$ is linearly independent. Then $p' = (1 - a_1)u_{i_1} + \cdots + (1 - a_n)u_{i_n}$ is a lattice point in the interior of σ^\vee . By Lemma 3, $\lambda^{\max}(u'_\sigma) \leq \lambda^{\max}(p')$, and since λ^{\max} is linear on γ and $\lambda^{\max}(u_{i_j}) = 1$, we have

$$\lambda^{\max}(p') = (1 - a_1) + \cdots + (1 - a_n) = n - \lambda^{\max}(p).$$

Therefore,

$$\lambda^{\max}(u'_\sigma) \leq n - \lambda^{\max}(p).$$

Let t be as in the proposition, i.e. $t = \min\{D \cdot V(\sigma \cap \tau)\}$, where τ varies over all maximal cones adjacent to σ . Then $t \geq n + 1 > \lambda^{\min}(u'_\sigma)$, so we can apply the proposition with $r = 1$ to obtain

$$\begin{aligned} m &\geq n + 1 - \lambda^{\min}(u'_\sigma) - 1. \\ &\geq n - \lambda^{\max}(u'_\sigma). \\ &\geq \lambda^{\max}(p). \end{aligned}$$

□

Remark – The collection of cones over the lower faces of Q is an example of what is called a “regular subdivision”. In general, a regular subdivision of a cone is constructed by choosing a nonzero point on each of the rays of the cone, and perhaps specifying some additional rays inside the cone with nonzero points on them as well. One looks at the convex hull of all of these points and then takes the cones over all of the lower faces. For more details on regular subdivisions of convex polytopes, see [Lee] and [Zi]. The translation from polytopes to cones is straightforward.

In the toric literature, regular subdivisions have generally been applied to the fan defining a toric variety, and sometimes to the polytope defining an ample line bundle. See, for instance, [OP] and [GKZ]. The regular subdivisions that we have used in this paper are of the dual cones $\{\sigma^\vee\}$. The author is not aware of any significant geometric interpretations for these subdivisions.

The subdivisions of a fan Σ correspond naturally and bijectively to the proper birational toric morphisms $\tilde{X} \rightarrow X(\Sigma)$ [Ful, Section 2.5], and the regular subdivisions of Σ are precisely those for which the corresponding morphism is projective. A regular subdivision of Σ is obtained by specifying a continuous function Ψ on the support of Σ that is convex and piecewise-linear on each cone. By subdividing each cone of Σ into the maximal subcones on which Ψ is linear, we get a projective birational morphism for which Ψ is the piecewise-linear function associated to a relatively ample T - \mathbb{Q} -Cartier divisor on \tilde{X} . In particular, if $D = \sum d_i D_i$ is a T - \mathbb{Q} -Weil divisor on X , and if we define Ψ_D^{\max} on each maximal cone σ by

$$\Psi_D^{\max}(v) = \max \left\{ \sum_{v_i \in \sigma} -a_i d_i : \sum_{v_i \in \sigma} a_i v_i = v, a_i \geq 0 \right\},$$

then we get the unique projective birational morphism $\pi : \tilde{X} \rightarrow X$ such that π^*D is \mathbb{Q} -Cartier and relatively ample, and π is an isomorphism in codimension 1. If D is effective (resp. minus effective) then, for each maximal cone σ , the same subdivision is obtained by looking at $\text{conv}\{\frac{1}{d_i}v_i : v_i \in \sigma\}$ (resp. $\text{conv}\{-\frac{1}{d_i}v_i : v_i \in \sigma\}$) and taking the cones over the upper faces (resp. lower faces). Note that, for a subdivision of a fan to be regular, it is not enough for the subdivision to be regular on each cone. This is the toric manifestation of the fact that quasiprojectivity is not local on the base (see [EGA, II.5.3]).

References

- [EGA] Grothendieck, A. and Dieudonné, J. “Eléments de Géométrie Algébrique II,” Publ. Math. IHES 8 (1961).
- [EL] Ein, L. and Lazarsfeld, R. “Syzygies and Koszul Cohomology of Smooth Projective Varieties of Arbitrary Dimension,” Invent. Math. **111** 1993, 51-67.
- [EW] Ewald, G. and Wessels, U. “On the ampleness of invertible sheaves on complete projective toric varieties,” Results in Math. **19** 1991, 275-278.
- [Fu] Fujino, O. “Notes on toric varieties from Mori theoretic viewpoint,” Preprint 2001. To appear in Tohoku Math J. math.AG/0112090.
- [Fuj] Fujita, T. “On polarized manifolds whose adjoint bundles are not semi-positive,” in *Algebraic Geometry, Sendai 1985*, Adv. Stud. Pure Math. **10**, North-Holland, Amsterdam 1987.
- [Ful] Fulton, W. *Introduction to Toric Varieties*, Princeton University Press 1993.
- [GKZ] Gelfand, I., Kapranov, M., and Zelevinsky, A. *Discriminants, Resultants, and Multidimensional Determinants*, Birkhauser 1994.
- [La] Laterveer, R. “Linear systems on toric varieties,” Tôhoku Math J. **48** 1996, 451-458.
- [Lee] Lee, C. “Regular triangulations of convex polytopes,” in *Applied Geometry and Discrete Mathematics, the Victor Klee Festschrift*, DIMACS Series in Discrete Math. and Theoretical Comp. Science **4**, AMS 1991, 443-456.
- [Lin] Lin, H.-W. “Combinatorial method in adjoint linear systems on toric varieties,” Preprint 2002. To appear in Michigan Math. J.
- [Mu] Mustață, M. “Vanishing theorems on toric varieties,” Tohoku Math. J. (2) **54** 2002, 451-470.

- [OP] Oda, T. and Park, H. S. “Linear Gale transformations and Gelfand-Kapranov-Zelevinskij decompositions,” *Tôhoku Math J.* **43** 1991, 375-399.
- [P] Payne, S. “Combinatorial proofs of Fujita’s freeness conjecture for singular toric varieties,” in preparation.
- [Re] Reid, M. “Decomposition of toric morphisms,” in *Arithmetic and Geometry II*, Progress in Math. **36**, Birkhauser 1983, 395-418.
- [Sm1] Smith, K. “Fujita’s freeness conjecture in terms of local cohomology,” *J. Algebraic Geometry* **6** 1997, 417-429.
- [Sm2] Smith, K. “A tight closure proof of Fujita’s freeness conjecture for very ample line bundles,” *Math. Annalen* **317** 2000, 285-293.
- [Zi] Ziegler, G. *Lectures on Polytopes*, Springer-Verlag 1995.