

ON THE SPLITTING-TYPE OF THE PRINCIPAL PARTS ON THE PROJECTIVE LINE

HELGE MAAKESTAD

ABSTRACT. We develop techniques to split explicitly the sheaf of principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left and right module on the projective line over an arbitrary ring. We then apply the techniques developed to split the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ for all $n \in \mathbf{Z}$ and $k \geq 1$ as left and right \mathcal{O} -module on the projective line over any field of characteristic zero giving a complete description of the principal parts on the projective line.

CONTENTS

1. Introduction	1
2. Sheaves of principal parts	2
3. Calculation of transition matrices over \mathbf{Z}	4
4. Maps of sheaves and finitely generated rings	7
5. Splitting over fields of characteristic zero	14
6. Appendix: calculations of determinants	18
References	19

1. INTRODUCTION

The sheaf of principal parts $\mathcal{P}_X^k(\mathcal{E})$ of an \mathcal{O}_X -module \mathcal{E} is an important sheaf used in many situations in algebraic geometry (see [8], [10], [12],[13] and [15]). For any smooth scheme X/S and any locally free sheaf \mathcal{E} there exists an exact sequence

$$(1.0.1) \quad 0 \rightarrow \Omega_X^1 \otimes \mathcal{E} \rightarrow \mathcal{P}_X^1(\mathcal{E}) \rightarrow \mathcal{E} \rightarrow 0$$

called the *Atiyah sequence* (see [1]). The sequence 1.0.1 is split by a connection

$$\nabla : \mathcal{E} \rightarrow \Omega_X^1 \otimes \mathcal{E},$$

and a connection ∇ gives rise to the *Chern-character* of \mathcal{E} . The Atiyah sequence is defined in terms of the first order sheaf of principal parts $\mathcal{P}_X^1(\mathcal{E})$, hence the first order principal parts contain all information about the Chern-character of \mathcal{E} . There exist higher order analogies of the sequence 1.0.1 and in the paper [7] splittings of higher order sequences is considered for holomorphic vector bundles on Kähler manifolds. It is shown that existence of holomorphic splittings of higher order fundamental exact sequences puts strong restrictions on \mathcal{E} and X . Not much

Date: Spring 2004.

Key words and phrases. Principal parts, sheaves of bi-modules, splitting type.

is known on splittings of higher order fundamental exact sequences in characteristic p .

A first step in the direction of considering splitting obstructions defined by higher fundamental exact sequences in characteristic p is to consider splitting formulas for the principal parts in characteristic p . Splitting formulas on projective space over a field of characteristic zero as left module has been studied by many authors (see [9], [11] and [14]). In the paper [5] explicit techniques was developed to split the principal parts on the projective line over any field as left module. It was also observed ([5], Theorem 7.1) that the splitting type as left and right module differed in general and that the splitting type depended on the characteristic of the base field.

In this paper the techniques from [5] are generalized to study the splitting type of the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ for all $k \geq 1$ and all $n \in \mathbf{Z}$ as left and right module on the projective line over any ring. The techniques are powerful enough to enable us to give a complete classification of the splitting type of the principal parts on the projective line over any field of characteristic zero. The main result of the paper is the following:

Theorem 1.1. *Let F be a field of characteristic zero. Consider the principal parts $\mathcal{P}^k(\mathcal{O}(d))$ on the projective line \mathbf{P}_F^1 . The structure of the principal parts as left module is as follows: If $k \geq 1$ and $n < 0$ or $n \geq k$ there exist an isomorphism*

$$(1.1.1) \quad \mathcal{P}^k(\mathcal{O}(n))^l \cong \oplus^{k+1} \mathcal{O}(n-k)$$

of left \mathcal{O} -modules. If $0 \leq n < k$ there exist an isomorphism

$$(1.1.2) \quad \mathcal{P}^k(\mathcal{O}(n))^l \cong \mathcal{O}^{n+1} \oplus \mathcal{O}(-k-1)^{k-n}$$

of left \mathcal{O} -modules. The structure of the principal parts as right module is as follows: For all $n \in \mathbf{Z}$ and $k \geq 1$ there exist an isomorphism

$$(1.1.3) \quad \mathcal{P}^k(\mathcal{O}(n))^r \cong \mathcal{O}(n) \oplus \mathcal{O}(n-k-1)^k$$

of right \mathcal{O} -modules.

Proof. See Theorem 4.1, 4.2 and 4.3. □

It is hoped that the techniques developed in the paper will shed light on the problem of classifying the structure of the principal parts as left and right module on the projective line over any field.

The paper is organized as follows: In section one we define and prove general properties of the sheaf of principal parts. In section two we calculate the transition matrices of the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ for all $k \geq 1$ and $n \in \mathbf{Z}$ on the projective line over the integers. In section three we develop techniques to split the principal parts as left and right module over any ring. In section four we apply the techniques developed in section three to give a complete classification of the splitting type of the principal parts on the projective line over any field of characteristic zero.

2. SHEAVES OF PRINCIPAL PARTS

In this section we define and give general properties of the sheaves of principal parts following [2], chapter 16.

Notation. Let in the following X/S be a separated scheme, and consider the two projection maps $p, q : X \times X \rightarrow X$. The diagonal embedding $\Delta : X \rightarrow X \times X$

is a closed immersion, and we get an exact sequence of sheaves of $\mathcal{O}_{X \times X}$ -modules on $X \times X$

$$(2.0.4) \quad 0 \rightarrow \mathcal{I}^{k+1} \rightarrow \mathcal{O}_{X \times X} \rightarrow \mathcal{O}_{\Delta^k} \rightarrow 0,$$

where \mathcal{I} is the sheaf of ideals defining the diagonal in $X \times X$.

Definition 2.1. Let \mathcal{E} be an quasi-coherent \mathcal{O}_X -module. We define the k 'th order sheaf of principal parts of \mathcal{E} , to be

$$\mathcal{P}_X^k(\mathcal{E}) = p_*(\mathcal{O}_{\Delta^k} \otimes_{X \times X} q^* \mathcal{E}).$$

We write \mathcal{P}_X^k for the module $\mathcal{P}_X^k(\mathcal{O}_X)$.

When it is clear from the context which scheme we are working on, we write $\mathcal{P}^k(\mathcal{E})$ instead of $\mathcal{P}_X^k(\mathcal{E})$.

Proposition 2.2. Let X/S be smooth and let \mathcal{E} be a locally free \mathcal{O}_X -module. There exists an exact sequence

$$0 \rightarrow S^k(\Omega_X^1) \otimes \mathcal{E} \rightarrow \mathcal{P}_X^k(\mathcal{E}) \rightarrow \mathcal{P}_X^{k-1}(\mathcal{E}) \rightarrow 0$$

of left \mathcal{O}_X -modules, where $k = 1, 2, \dots$

Proof. See [5], Proposition 2.2. \square

From Proposition 2.2 it follows that for a smooth morphism $X \rightarrow S$ of relative dimension n , and \mathcal{E} a locally free sheaf on X of rank e , the principal parts $\mathcal{P}^k(\mathcal{E})$ is locally free of rank $e \binom{n+k}{n}$.

Proposition 2.3. Let $f : X \rightarrow Y$ be a map of smooth schemes over S , and let \mathcal{E} be a locally free \mathcal{O}_Y -module. There exists a commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & S^k(f^*\Omega_Y^1) \otimes f^*\mathcal{E} & \longrightarrow & f^*\mathcal{P}_Y^k(\mathcal{E}) & \longrightarrow & f^*\mathcal{P}_Y^{k-1}(\mathcal{E}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & S^k(\Omega_X^1) \otimes f^*\mathcal{E} & \longrightarrow & \mathcal{P}_X^k(f^*\mathcal{E}) & \longrightarrow & \mathcal{P}_X^{k-1}(f^*\mathcal{E}) \longrightarrow 0 \end{array}$$

of left \mathcal{O}_X -modules for all $k = 1, 2, \dots$

Proof. See [5], Proposition 2.3. \square

Note that it follows from Proposition 2.3 that for any open set $U \subseteq X$ of a scheme X smooth over S there is an isomorphism

$$(2.3.1) \quad \mathcal{P}_X^k(\mathcal{E})|_U \cong \mathcal{P}_U^k(\mathcal{E}|_U)$$

of left \mathcal{O}_U -modules: Let $i : U \rightarrow X$ be the inclusion map. It follows that i is smooth. By induction and applying the 5-lemma Proposition 2.3 proves the isomorphism 2.3.1. Hence the sheaf of principal parts can be studied using local calculations.

Given any \mathcal{O}_X -modules \mathcal{F} and \mathcal{G} one may define the sheaf of differential operators of order k from \mathcal{F} to \mathcal{G} following [2] section 16.8, denoted $\mathcal{D}_X^k(\mathcal{F}, \mathcal{G})$, and there exists an isomorphism

$$(2.3.2) \quad \text{Hom}_X(\mathcal{P}_X^k(\mathcal{F}), \mathcal{G}) \cong \mathcal{D}_X^k(\mathcal{F}, \mathcal{G})$$

of sheaves of abelian groups. The sheaf of abelian groups $\mathcal{D}_X^k(\mathcal{F}, \mathcal{G})$ is naturally a sheaf of \mathcal{O}_X -bimodules and it follows from the isomorphism (2.3.2) that the sheaf of principal parts $\mathcal{P}_X^k(\mathcal{F})$ is also a sheaf of \mathcal{O}_X -bimodules. This means that for

any open set U of X , and local sections $a, b \in \mathcal{O}(U)$ there exists a left and right multiplication on the module $\mathcal{P}^k(\mathcal{E})(U)$ with the property that for all w in $\mathcal{P}^k(\mathcal{E})(U)$ we have

$$a(wb) = (aw)b.$$

Restriction-maps of the sheaf $\mathcal{P}_X^k(\mathcal{E})$ satisfy obvious criteria. Note that for X smooth over S and \mathcal{E} locally free of finite rank, it follows that $\mathcal{P}_X^k(\mathcal{E})$ is locally free of finite rank as left and right \mathcal{O}_X -module separately.

3. CALCULATION OF TRANSITION MATRICES OVER \mathbf{Z}

In this section we calculate the transition-matrices L_n^k and R_n^k of the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ on the projective line over \mathbf{Z} for all $n \in \mathbf{Z}$ and all $k \geq 1$. The structure-matrices defining the principal parts on the projective line are matrices with entries being Laurent-polynomials with binomial coefficients, hence the principal parts are naturally defined over \mathbf{Z} . The matrices L_n^k and R_n^k are elements of the group $\mathrm{GL}(k+1, \mathbf{Z}[t, t^{-1}])$. The matrix L_n^k define the left \mathcal{O} -module structure, and the matrix R_n^k define the right \mathcal{O} -module structure. Note that in this section all tensor products are over \mathbf{Z} .

Notation. Consider the projective line $\mathbf{P} = \mathbf{P}_{\mathbf{Z}}^1$ over the integers \mathbf{Z} . By definition \mathbf{P} equals $\mathrm{Proj} \mathbf{Z}[x_0, x_1]$. Let $U_i = \mathrm{Spec} \mathbf{Z}[x_0, x_1]_{(x_i)}$. Let furthermore $t = \frac{x_1}{x_0} \otimes 1$ and $u = 1 \otimes \frac{x_1}{x_0}$ under the isomorphism

$$\mathbf{Z}\left[\frac{x_1}{x_0}\right] \otimes_{\mathbf{Z}} \mathbf{Z}\left[\frac{x_1}{x_0}\right] \cong \mathbf{Z}[u, t].$$

It follows from Definition 2.1 and Proposition 2.3.1 that the sheaf of principal parts $\mathcal{P}^k(\mathcal{O}(n))|_{U_0}$ as a left \mathcal{O}_{U_0} -module equals

$$\mathbf{Z}[t]\{1 \otimes x_0^n, dt \otimes x_0^n, \dots, dt^k \otimes x_0^n\}$$

as a free $\mathbf{Z}[t]$ -module where $dt^p = (u - t)^p$. Let $\frac{1}{t} = \frac{x_0}{x_1} \otimes 1$ and $\frac{1}{u} = 1 \otimes \frac{x_0}{x_1}$ under the isomorphism

$$\mathbf{Z}\left[\frac{x_0}{x_1}\right] \otimes_{\mathbf{Z}} \mathbf{Z}\left[\frac{x_0}{x_1}\right] \cong \mathbf{Z}\left[\frac{1}{u}, \frac{1}{t}\right].$$

It follows from Definition 2.1 and Proposition 2.3.1 that $\mathcal{P}^k(\mathcal{O}(n))|_{U_1}$ as a left \mathcal{O}_{U_1} -module equals

$$\mathbf{Z}[s]\{1 \otimes x_1^n, ds \otimes x_1^n, \dots, ds^k \otimes x_1^n\}$$

as a free $\mathbf{Z}[s]$ -module, where $s = \frac{1}{t}$ and $ds^p = (\frac{1}{u} - \frac{1}{t})^p$. Let

$$C = \{1 \otimes x_0^n, dt \otimes x_0^n, \dots, dt^k \otimes x_0^n\}$$

and

$$C' = \{1 \otimes x_1^n, ds \otimes x_1^n, \dots, ds^k \otimes x_1^n\}.$$

We aim to describe the principal parts by calculating the transition-matrix

$$L_n^k = [I]_C^{C'}$$

between the bases C' and C as an element of the group $\mathrm{GL}(k+1, \mathbf{Z}[t, t^{-1}])$. The following theorem gives the structure-matrix L_n^k describing the left \mathcal{O} -module structure of the principal parts on the projective line over \mathbf{Z} .

Theorem 3.1. Consider $\mathcal{P}^k(\mathcal{O}(n))$ as left \mathcal{O} -module on \mathbf{P} . In the case $n < 0$ the structure-matrix L_n^k is given by the following formula:

$$(3.1.1) \quad ds^p \otimes x_1^n = \sum_{j=p}^k (-1)^j \binom{j-n-1}{p-n-1} t^{n-p-j} dt^j \otimes x_0^n,$$

where $0 \leq p \leq k$. If $n \geq 0$ the matrix L_n^k is given as follows:

Case $1 \leq k \leq n$:

$$(3.1.2) \quad ds^p \otimes x_1^n = \sum_{j=p}^k (-1)^p \binom{n-p}{j-p} t^{n-p-j} dt^j \otimes x_0^n,$$

where $0 \leq p \leq k$.

Case $n < k$:

$$(3.1.3) \quad ds^p \otimes x_1^n = \sum_{j=p}^n (-1)^p \binom{n-p}{j-p} t^{n-p-j} dt^j \otimes x_0^n,$$

if $1 \leq p \leq n$, and

$$(3.1.4) \quad ds^p \otimes x_1^n = \sum_{j=p}^k (-1)^j \binom{j-n-1}{p-n-1} t^{n-p-j} dt^j \otimes x_0^n,$$

if $n < p \leq k$.

Proof. Case 3.1.1: let $0 \leq p \leq k$.

$$ds^p \otimes x_1^n = \left(\frac{1}{u} - \frac{1}{t}\right)^p u^n \otimes x_0^n = (-1)^p \frac{1}{t^p} dt^p (t + dt)^{n-p} \otimes x_0^n.$$

We get

$$\sum_{i=0}^{k-p} (-1)^{i+p} \binom{i+p-n-1}{p-n-1} t^{n-2p-i} dt^{i+p} \otimes x_0^n.$$

Changing index with $j = i + p$ we get

$$\sum_{j=p}^k (-1)^j \binom{j-n-1}{p-n-1} t^{n-p-j} dt^j \otimes x_0^n,$$

and equation 3.1.1 is proved.

We prove equation 3.1.2: let $0 \leq p \leq k$. We get

$$\begin{aligned} ds^p \otimes x_1^n &= \left(\frac{1}{u} - \frac{1}{t}\right)^p \otimes u^n x_0^n = (-1)^p \frac{1}{t^p} dt^p u^{n-p} \otimes x_0^n = \\ &= (-1)^p \frac{1}{t^p} dt^p (t + dt)^{n-p} \otimes x_0^n = (-1)^p \frac{1}{t^p} dt^p \sum_{i=0}^{n-p} \binom{n-p}{i} t^{n-p-i} dt^i \otimes x_0^n = \\ &= \sum_{i=0}^{n-p} (-1)^p \binom{n-p}{i} t^{n-2p-i} dt^{i+p} \otimes x_0^n. \end{aligned}$$

Changing index with $j = i + p$ we get

$$\sum_{j=p}^k (-1)^p \binom{n-p}{j-p} t^{n-p-j} dt^j \otimes x_0^n,$$

and equation 3.1.2 is proved.

We prove equation 3.1.3: If $0 \leq p \leq n$ we get

$$ds^p \otimes x_1^n = \left(\frac{1}{u} - \frac{1}{t}\right)^p u^n \otimes x_0^n = (-1)^p \frac{1}{t^p} dt^p (t + dt)^{n-p} \otimes x_0^n =$$

$$\sum_{i=0}^{n-p} (-1)^p \binom{n-p}{i} t^{n-2p-i} dt^{i+p} \otimes x_0^n.$$

Changing index with $j = i + p$ we get

$$\sum_{j=p}^n (-1)^p \binom{n-p}{j-p} t^{n-p-j} dt^j \otimes x_0^n,$$

and equation 3.1.3 is proved.

Let $n < p \leq k$. We prove equation 3.1.4: We get

$$ds^p \otimes x_1^n = (-1)^p \frac{1}{t^p} dt^p u^{n-p} \otimes x_0^n = (-1)^p \frac{1}{t^p} dt^p (t + dt)^{n-p} \otimes x_0^n.$$

It follows that

$$(-1)^p \frac{1}{t^p} dt^p \sum_{i \geq 0} (-1)^i \binom{i+p-n-1}{p-n-1} \frac{1}{t^i} dt^i \otimes x_0^n =$$

$$\sum_{i \geq 0} (-1)^{i+p} \binom{i+p-n-1}{p-n-1} \frac{1}{t^{i+2p-n}} dt^{i+p} \otimes x_0^n.$$

We change index with $j = i + p$ and get

$$\sum_{j=p}^k (-1)^j \binom{j-n-1}{p-n-1} t^{n-p-j} dt^j \otimes x_0^n,$$

and equation 3.1.4 is proved, and the theorem follows. \square

We use similar calculations as in the Theorem 3.1 to study the structure-matrix R_n^k defining the principal parts as right \mathcal{O} -module on the projective line over \mathbf{Z} .

Notation. It follows from Definition 2.1 and Proposition 2.3.1 that the principal parts $\mathcal{P}^k(\mathcal{O}(n))|_{U_0}$ as a right \mathcal{O}_{U_0} -module equals

$$\mathbf{Z}[u]\{1 \otimes x_0^n, du \otimes x_0^n, \dots, du^k \otimes x_0^n\}$$

as a free $\mathbf{Z}[u]$ -module, where $du^p = (t - u)^p$. It follows that $\mathcal{P}^k(\mathcal{O}(n))|_{U_1}$ as a right \mathcal{O}_{U_1} -module equals

$$\mathbf{Z}[v]\{1 \otimes x_1^n, dv \otimes x_1^n, \dots, dv^k \otimes x_1^n\}$$

as a free $\mathbf{Z}[v]$ -module, where $v = \frac{1}{u}$ and $dv^p = (\frac{1}{t} - \frac{1}{u})^p$. Let

$$D = \{1 \otimes x_0^n, du \otimes x_0^n, \dots, du^k \otimes x_0^n\}$$

and

$$D' = \{1 \otimes x_1^n, dv \otimes x_1^n, \dots, dv^k \otimes x_1^n\}.$$

We wish to calculate the transition-matrix

$$R_n^k = [I]_D^{D'}$$

describing the right module structure on the principal parts. The following theorem gives the structure-matrix R_n^k :

Theorem 3.2. *Consider $\mathcal{P}^k(\mathcal{O}(n))$ as right \mathcal{O} -module on \mathbf{P} . If $n \in \mathbf{Z}$ and $k \geq 1$ the following formulas holds:*

$$(3.2.1) \quad 1 \otimes x_1^n = u^n \otimes x_0^n$$

and

$$(3.2.2) \quad dv^p \otimes x_1^n = \sum_{j=p}^k (-1)^j \binom{j-1}{p-1} u^{n-p-j} du^j \otimes x_0^n,$$

where $1 \leq p \leq k$.

Proof. Equation 3.2.1 is trivial hence we prove equation 3.2.2: Let $1 \leq p \leq k$. We get

$$\begin{aligned} dv^p \otimes x_1^n &= \left(\frac{1}{t} - \frac{1}{u}\right)^p u^n \otimes x_0^n = (-1)^p u^{n-p} du^p \frac{1}{(u+du)^p} \otimes x_0^n = \\ &= (-1)^p u^{n-2p} du^p \frac{1}{\left(1 + \frac{1}{u} du\right)^p} \otimes x_0^n = \\ &= (-1)^p u^{n-2p} du^p \sum_{i=0}^k (-1)^i \binom{i+p-1}{p-1} \frac{1}{u^i} du^i \otimes x_0^n. \end{aligned}$$

We get

$$\sum_{i=0}^{k-p} (-1)^{i+p} \binom{i+p-1}{p-1} u^{n-2p-i} du^{i+p} \otimes x_0^n.$$

Let $j = i + p$ we get

$$\sum_{j=p}^k (-1)^j \binom{j-1}{p-1} u^{n-p-j} du^j \otimes x_0^n,$$

and equation 3.2.2 is proved, and the theorem follows. \square

The Theorems 3.1 and 3.2 describe completely the matrices L_n^k and R_n^k for all $n \in \mathbf{Z}$ and $k \geq 1$. By [3] Theorem 2.1 and [4] Theorem 3.1 we know that any finite rank locally free sheaf on \mathbf{P}^1 over any field split into a direct sum of invertible sheaves. The formation of principal parts commute with direct sums, hence it follows from Theorem 3.1 and 3.2 that we have calculated the transition matrix of $\mathcal{P}^k(\mathcal{E})$ for any locally free finite rank sheaf \mathcal{E} as left and right \mathcal{O} -module on the projective line over any field.

4. MAPS OF SHEAVES AND FINITELY GENERATED RINGS

In this section we develop techniques to define maps of sheaves of left and right $\mathcal{O}_{\mathbf{P}^1_A}$ -modules, where A is any ring. We define for all $n \in \mathbf{Z}$ and $k \geq 1$ finitely generated rings $B_{k,n}^L$ and $B_{k,n}^R$ over the integers, with the property that any ring-homomorphism $f : B_{k,n}^L \rightarrow A$ give rise to maps of left modules

$$\phi_{k,n}^L : \oplus_{i=0}^k \mathcal{O}(d_i(k,n)) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

on the projective line over A . Furthermore, any ring-homomorphism $g : B_{k,n}^R \rightarrow A'$ give rise to a map of right modules

$$\phi_{k,n}^R : \oplus_{j=0}^k \mathcal{O}(e_j(k,n)) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

on the projective line over A' . This is a generalization of the systems of linear equations $A_r \mathbf{x}_r = b_r$ introduced in [5], chapter 5 where only the left module structure in the case $1 \leq k \leq n$ was considered.

Notation. We consider in the following discussion the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left \mathcal{O} -module with $n < 0$ and $k \geq 1$. Let $0 \leq l \leq k$, and let $x_j^{i,1}$ be independent variables over \mathbf{Z} for all $0 \leq j \leq i \leq k$. and let $A_1 = \mathbf{Z}[x_j^{i,1}]$. Consider the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ on $\mathbf{P}_{A_1}^1$. We seek to define a map

$$\phi_l : \mathcal{O}(n-k) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_{B_l^1}^1}$ -modules, where B_l^1 is an A_1 -algebra. Define on U_1 the map

$$(4.0.3) \quad \phi_l^1(x_1^{n-k}) = \sum_{p=0}^l x_{l-p}^{l,1} \frac{1}{t^{l-p}} ds^p \otimes x_1^n.$$

We want to define a map of $\mathcal{O}_{\mathbf{P}^1}$ -modules, and since $x_0^{n-k} = t^{k-n} x_1^{n-k}$ we define on U_0 the map

$$\phi_l^0(x_0^{n-k}) = \frac{1}{t^{n-k}} \sum_{p=0}^l x_{l-p}^{l,1} t^{p-l} ds^p \otimes x_1^n = \sum_{p=0}^l x_{l-p}^{l,1} t^{k-n-l+p} ds^p \otimes x_1^n.$$

By Theorem 3.1, equation 3.1.1 we get

$$\begin{aligned} & \sum_{p=0}^l x_{l-p}^{l,1} t^{k-n-l+p} \sum_{j=p}^k (-1)^j \binom{j-n-1}{p-n-1} t^{n-p-j} dt^j \otimes x_0^n = \\ & \sum_{p=0}^l \sum_{j=0}^k (-1)^j x_{l-p}^{l,1} \binom{j-n-1}{p-n-1} t^{k-l-j} dt^j \otimes x_0^n = \\ & \sum_{j=0}^k \sum_{p=0}^l (-1)^j x_{l-p}^{l,1} \binom{j-n-1}{p-n-1} t^{k-l-j} dt^j \otimes x_0^n = \\ & \sum_{j=0}^k (-1)^j \left[\sum_{p=0}^l x_{l-p}^{l,1} \binom{j-n-1}{p-n-1} \right] t^{k-l-j} dt^j \otimes x_0^n = \\ & \sum_{j=0}^k (-1)^j c_j^{l,1} t^{k-l-j} dt^j \otimes x_0^n, \end{aligned}$$

where

$$(4.0.4) \quad c_j^{l,1} = \sum_{p=0}^l x_{l-p}^{l,1} \binom{j-n-1}{p-n-1}.$$

We get

$$\begin{aligned} & \sum_{j=0}^{k-l-1} (-1)^j c_j^{l,1} t^{k-l-1} dt^j \otimes x_0^n + (-1)^{k-l} c_{k-l}^{l,1} dt^{k-l} \otimes x_0^n + \\ & (-1)^{k-l+1} c_{k-l+1}^{l,1} \frac{1}{t} dt^{k-l+1} \otimes x_0^n + \cdots + (-1)^k c_k^{l,1} \frac{1}{t^l} dt^k \otimes x_0^n. \end{aligned}$$

Definition 4.1. Let U_i^1 be the ideal generated by the variables $x_j^{i,1}$ with $0 \leq j \leq i$. Let also $U^{l,1} = \sum_{i \neq l} U_i^1$. Let I_l be the ideal in A^1 generated by the polynomials

$$c_k^{l,1}, c_{k-1}^{l,1}, \dots, c_{k-l+1}^{l,1}, c_{k-l}^{l,1} - 1,$$

and let B_l^1 be the A^1 -algebra $B_l^1 = A^1/I_l + U^{l,1}$.

Note that B_l^1 is finitely generated over \mathbf{Z} . The A -valued points of the ring B_l^1 has a relation to maps of left modules:

Proposition 4.2. *For all $l = 0, \dots, k$ the ring B_l^1 has the following property: Any B_l^1 -algebra A determines a map of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules*

$$\phi_l^1 : \mathcal{O}(n-k) \rightarrow \mathcal{P}^k(\mathcal{O}(n)).$$

Proof. Let $f : B_l^1 \rightarrow A$ be a ring-homomorphism. Define a map ϕ_l as follows:

$$(4.2.1) \quad \phi_l^{1,0}(x_0^{n-k}) = \sum_{j=0}^{k-l} (-1)^j f(\bar{c}_j^{l,1}) t^{k-l-1} dt^j \otimes x_0^n$$

and

$$(4.2.2) \quad \phi_l^{1,1}(x_1^{n-k}) = \sum_{p=0}^l f(\bar{x}_{l-p}^{l,1}) \frac{1}{t^{l-p}} ds^p \otimes x_1^n,$$

where $\bar{c}_j^{l,1}$ and $\bar{x}_{l-p}^{l,1}$ are the equivalence-classes of $c_j^{l,1}$ and $x_{l-p}^{l,1}$ in B_l^1 . By construction of the coefficients $c_j^{l,1}$ the maps 4.2.1 and 4.2.2 glue to a well-defined map ϕ_l of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules, and the proposition is proved. \square

We proceed to consider $\mathcal{P}^k(\mathcal{O}(n))$ as left module when $0 \leq n < k$.

Notation. Consider the sheaf \mathcal{O} and let $0 \leq l \leq n$. Pick a basis e for $\mathcal{O}(U_0)$ as a free left $\mathbf{Z}[t]$ -module. Let f be a basis for $\mathcal{O}(U_1)$ as a free left $\mathbf{Z}[s]$ -module. Let $x_j^{i,2}$ be independent variables over \mathbf{Z} with $0 \leq j \leq i \leq n$, and let $A^2 = \mathbf{Z}[x_j^{i,2}]$.

Consider $\mathbf{P}_{A^2}^1$ and define the map on U_1 :

$$\phi_l^1(f) = \sum_{p=0}^l x_{l-p}^{l,2} t^{p-l} ds^p \otimes x_1^n.$$

By Theorem 3.1 and equation 3.1.3 we can define a map on U_0 as follows:

$$\begin{aligned} \phi_l^0(e) &= \sum_{p=0}^l x_{l-p}^{l,2} t^{p-l} \sum_{j=0}^n (-1)^p \binom{n-p}{j-p} t^{n-p-j} dt^j \otimes x_0^n = \\ &= \sum_{p=0}^l \sum_{j=0}^n (-1)^p x_{l-p}^{l,2} \binom{n-p}{j-p} t^{n-l-j} dt^j \otimes x_0^n = \\ &= \sum_{j=0}^n \left[\sum_{p=0}^l (-1)^p x_{l-p}^{l,2} \binom{n-p}{j-p} \right] t^{n-l-p} dt^j \otimes x_0^n = \\ &= \sum_{j=0}^n c_j^{l,2} t^{n-l-j} dt^j \otimes x_0^n, \end{aligned}$$

where

$$(4.2.3) \quad c_j^{l,2} = \sum_{p=0}^l (-1)^p x_{l-p}^{l,2} \binom{n-p}{j-p}.$$

Definition 4.3. Let U_i^2 be the ideal generated by the monomials $x_j^{i,2}$ where $0 \leq j \leq i$ and let $U^{l,2} = \sum_{i \neq l} U_i^2$. Let for all $l = 0, \dots, n$ the ideal I_l^2 be generated by the polynomials

$$c_{n-l+1}^{l,2}, \dots, c_n^{l,2}, c_{n-l}^{l,2} - 1.$$

Put $B_l^2 = A^2/I_l^2 + U^{l,2}$.

Note that B_l^2 is finitely generated over the integers. The A -valued points of the ring B_l^2 is related to maps of left modules in the following way:

Proposition 4.4. Consider $\mathcal{P}^k(\mathcal{O}(n))$ as left module for $0 \leq n < k$. For all $l = 0, \dots, n$ the ring B_l^2 has the following property: any B_l^2 -algebra A determines a map

$$\phi_l^2 : \mathcal{O} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Proof. Let $f : A \rightarrow B_l^2$ be an A -valued point. Define a map on U_0 as follows:

$$(4.4.1) \quad \phi_l^{2,0}(e) = \sum_{j=0}^{n-l} f(\bar{c}_j^{l,2}) t^{n-l-j} dt^j \otimes x_0^n$$

and on U_1 :

$$(4.4.2) \quad \phi_l^{2,1}(f) = \sum_{p=0}^l f(\bar{x}_{l-p}^{l,2}) t^{p-l} ds^p \otimes x_1^n,$$

where $\bar{c}_j^{l,2}$ and $\bar{x}_{l-p}^{l,2}$ are the equivalence-classes of $c_j^{l,2}$ and $x_{l-p}^{l,2}$ in B_l^2 . By construction the maps 4.4.1 and 4.4.2 glue to a well-defined map ϕ_l^2 , and the proposition is proved. \square

We furthermore want to define a map

$$\phi_m^3 : \mathcal{O}(-k-1) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left \mathcal{O} -modules with $0 \leq m \leq k-n-1$.

Notation. Let $x_j^{i,3}$ be independent variables over \mathbf{Z} with $0 \leq j \leq i \leq k-n-1$. Let $A^3 = \mathbf{Z}[x_j^{i,3}]$ and consider $\mathbf{P}_{A^3}^1$. On the open set U_1 we define

$$\phi_m^{3,1}(x_1^{-k-1}) = \sum_{i=0}^m x_{m-i}^{m,3} t^{i-m} ds^{n+1+i} \otimes x_1^n.$$

By Theorem 3.1 and equation 3.1.4 we can define on U_0 the following map:

$$\begin{aligned} \phi_m^{3,0}(x_0^{-k-1}) &= t^{k+1} \sum_{i=0}^m x_{m-i}^{m,3} t^{i-m} \sum_{j=i+n+1}^k (-1)^j \binom{j-n-1}{i} t^{-1-i-j} dt^j \otimes x_0^n = \\ &= t^{k+1} \sum_{i=0}^m \sum_{j=i+n+1}^k (-1)^j x_{m-i}^{m,3} \binom{j-n-1}{i} t^{-j-m-1} dt^j \otimes x_0^n = \end{aligned}$$

$$\begin{aligned}
& t^{k+1} \sum_{j=n+1}^k \sum_{i=0}^m (-1)^j x_{m-i}^{m,3} \binom{j-n-1}{i} t^{-j-m-1} dt^j \otimes x_0^n = \\
& t^{k+1} \sum_{j=n+1}^k (-1)^j \left[\sum_{i=0}^m x_{m-i}^{m,3} \binom{j-n-1}{i} \right] t^{-j-m-1} dt^j \otimes x_0^n = \\
& \sum_{j=k+1}^k (-1)^j c_j^{m,3} t^{k-m-j} dt^j \otimes x_0^n,
\end{aligned}$$

where

$$(4.4.3) \quad c_j^{m,3} = \sum_{i=0}^m x_{m-i}^{m,3} \binom{j-n-1}{i}.$$

We have defined a map on U_0 as follows:

$$\begin{aligned}
\phi_m^{3,0}(x_0^n) &= \sum_{j=n+1}^{k-n-1} (-1)^j c_j^{m,3} t^{k-m-j} dt^j \otimes x_0^n + (-1)^{k-m} c_{k-m}^{m,3} dt^{k-m} \otimes x_0^n + \\
& (-1)^{k-m+1} c_{k-m+1}^{m,3} \frac{1}{t} dt^{k-m+1} \otimes x_0^n + \cdots + (-1)^k c_k^{m,3} \frac{1}{t^m} dt^k \otimes x_0^n,
\end{aligned}$$

Definition 4.5. Let the ideal U_i^3 be generated by the monomials $x_j^{i,3}$ where $0 \leq j \leq i$ and let $U^{m,3} = \sum_{i \neq m} U_i^3$. Let $0 \leq m \leq k-n-1$. Let I_m^3 be the ideal in A^3 defined by the polynomials

$$c_{k-m+1}^{m,3}, \dots, c_k^{m,3}, c_{k-m}^{m,3} - 1.$$

Define the A^3 -algebra B_m^3 to be $B_m^3 = A^3/I_m^3 + U^{m,3}$.

Note that B_m^3 is finitely generated over the integers. The A -valued points of B_m^3 has a relation to maps of left modules:

Proposition 4.6. Consider $\mathcal{P}^k(\mathcal{O}(n))$ as left module with $0 \leq n < k$. Let $0 \leq m \leq k-n-1$. The ring B_m^3 has the property that any B_m^3 -algebra A give rise to a map

$$\phi_m^3 : \mathcal{O}(-k-1) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Proof. Let $f : B_m^3 \rightarrow A$ be a ring-homomorphism. The map ϕ_m^3 looks as follows on U_0 :

$$(4.6.1) \quad \phi_m^{3,0}(x_0^{-k-1}) = \sum_{j=k+1}^{k-m} (-1)^j f(\bar{c}_j^{m,3}) t^{k-m-j} dt^j \otimes x_0^n$$

On U_1 :

$$(4.6.2) \quad \phi_m^{3,1}(x_1^{-k-1}) = \sum_{i=0}^m f(\bar{x}_{m-i}^{m,3}) t^{i-m} ds^{n+1+i} \otimes x_1^n.$$

where $\bar{c}_j^{m,3}$ and $\bar{x}_{m-i}^{m,3}$ are the equivalence-classes of $c_j^{m,3}$ and $x_{m-i}^{m,3}$ in B_m^3 . By construction the maps 4.6.1 and 4.6.2 glue to give a well-defined map ϕ_m^3 , and the proposition is proved. \square

In the following we consider the principal parts as right \mathcal{O} -module and construction of maps of right \mathcal{O} -modules using the same techniques as introduced in the previous discussion.

Notation. Recall that the principal parts $\mathcal{P}^k(\mathcal{O}(n))|_{U_0}$ as a right \mathcal{O}_{U_0} -module equals

$$\mathbf{Z}[u]\{1 \otimes x_0^n, du \otimes x_0^n, \dots, du^k \otimes x_0^n\}$$

as a free $\mathbf{Z}[u]$ -module where $du^p = (u - t)^p$. It also follows that $\mathcal{P}^k(\mathcal{O}(n))|_{U_1}$ as a right \mathcal{O}_{U_1} -module equals

$$\mathbf{Z}[v]\{1 \otimes x_1^n, dv \otimes x_1^n, \dots, dv^k \otimes x_1^n\}$$

as a free $\mathbf{Z}[v]$ -module, where $v = \frac{1}{u}$ and $dv^p = (\frac{1}{t} - \frac{1}{u})^p$. Let

$$D = \{1 \otimes x_0^n, du \otimes x_0^n, \dots, du^k \otimes x_0^n\}$$

and

$$D' = \{1 \otimes x_1^n, dv \otimes x_1^n, \dots, dv^k \otimes x_1^n\}.$$

We want to define maps of sheaves

$$\psi_l : \mathcal{O}(n - k - 1) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of right \mathcal{O} -modules. Let in the following $y_j^{i,1}$ be independent variables over \mathbf{Z} with $0 \leq j \leq i \leq k - 1$ and let $0 \leq l \leq k - 1$. Put $C^1 = \mathbf{Z}[y_j^{i,1}]$ and consider $\mathbf{P}_{C^1}^1$. Define a map on U_1 :

$$\psi_l^{1,1}(x_1^{n-k-1}) = \sum_{i=0}^l y_{l-i}^{l,1} u^{i-l} dv^{i+1} \otimes x_1^n.$$

By Theorem 3.2 and equations 3.2.1 and 3.2.2 we can define a map on U_0 as follows:

$$\begin{aligned} \psi_l^{1,0}(x_0^n) &= u^{k+1-n} \sum_{i=0}^l y_{l-i}^{l,1} u^{i-l} dv^{i+1} \otimes x_1^n = \\ &= u^{k+1-n} \sum_{i=0}^l y_{l-i}^{l,1} u^{i-l} \sum_{j=i+1}^k (-1)^j \binom{j-1}{i} u^{n-i-j-1} du^j \otimes x_0^n = \\ &= \sum_{i=0}^l \sum_{j=i+1}^k (-1)^j y_{l-i}^{l,1} \binom{j-1}{i} u^{k-l-j} du^j \otimes x_0^n = \\ &= \sum_{j=0}^k (-1)^j \left[\sum_{i=0}^l y_{l-i}^{l,1} \binom{j-1}{i} \right] u^{k-l-j} du^j \otimes x_0^n = \\ &= \sum_{j=0}^k (-1)^j d_j^{l,1} u^{k-l-j} du^j \otimes x_0^n, \end{aligned}$$

where

$$d_j^{l,1} = \sum_{i=0}^l y_{l-i}^{l,1} \binom{j-1}{i}.$$

We get

$$\psi_l^{1,0}(x_0^n) = \sum_{j=0}^{k-l-1} (-1)^j d_j^{l,1} u^{k-l-j} du^j \otimes x_0^n + (-1)^{k-l} d_{k-l}^{l,1} du^{k-l} \otimes x_0^n +$$

$$(-1)^{k-l+1} d_{k-l+1}^{l,1} \frac{1}{u} du^{k-l+1} \otimes x_0^n + \cdots + (-1)^k d_k^{l,1} \frac{1}{u} du^k \otimes x_0^n,$$

Definition 4.7. Define the ideal V_i^1 to be generated by the monomials $y_j^{i,1}$ where $0 \leq j \leq i$ and let $V^{l,1} = \sum_{i \neq l} V_i^1$. Let $0 \leq l \leq k-1$. Define J_l^1 to be the ideal generated by the polynomials

$$d_k^{l,1}, \dots, d_{k-l+1}^{l,1}, d_{k-l}^{l,1} - 1.$$

Let D_l^1 be the C^1 -algebra defined by $D_l^1 = C^1/J_l^1 + V^{l,1}$.

The ring D_l^1 is finitely generated over \mathbf{Z} . We get a proposition on the A -valued points of the ring D_l^1 :

Proposition 4.8. Consider $\mathcal{P}^k(\mathcal{O}(n))$ as right module for all $n \in \mathbf{Z}$ and $k \geq 1$. Let $0 \leq l \leq k-1$. The ring D_l^1 has the property that any D_l^1 -algebra A gives rise to a map

$$\psi_l : \mathcal{O}(n-k-1) \rightarrow \mathcal{P}^l(\mathcal{O}(n))$$

of right $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Proof. Let $f : D_l^1 \rightarrow A$ be a ring-homomorphism. Define a map ψ_l^0 on U_0 as follows:

$$(4.8.1) \quad \psi_l^0(x_0^{n-k-1}) = \sum_{j=0}^{k-l} (-1)^j f(\bar{d}_j^{l,1}) u^{k-l-j} du^j \otimes x_0^n.$$

Define a map on U_1 :

$$(4.8.2) \quad \psi_l^1(x_1^{n-k-1}) = \sum_{i=0}^l f(\bar{y}_{l-i}^{l,1}) u^{i-l} dv^{i+1} \otimes x_1^n,$$

where $\bar{d}_j^{l,1}$ and $\bar{y}_{l-i}^{l,1}$ are the equivalence-classes of the elements $d_j^{l,1}$ and $y_{l-i}^{l,1}$ in D_l^1 . By construction the maps 4.8.1 and 4.8.2 glue to a map ψ_l and the proposition is proved. \square

We sum up the results in the preceding propositions on existence of maps of left and right modules in a concluding theorem. We first need some definitions.

Definition 4.9. Let I^1 be the sum of the ideals I_l from definition 4.1 for $l = 0, \dots, k$, and let for $n < 0$ and $k \geq 1$ $B_{k,n}^L = A^1/I^1$. Let I^2 be the ideal $\sum_{i=0, \dots, n} I_i^2 + \sum_{j=0, \dots, k-n-1} I_j^3$ in the ring $\mathbf{Z}[x_j^{i,2}, x_v^{u,3}]$ where $0 \leq j \leq i \leq n$ and $0 \leq v \leq u \leq k-n-1$. Let for $0 \leq n \leq k$ $B_{k,n}^L = \mathbf{Z}[x_j^{i,2}, x_v^{u,3}]/I^2$. Let finally for all $n \in \mathbf{Z}$ and $k \geq 1$ $B_{k,n}^R = C^1 / \sum_{l=0, \dots, k-1} J_l^1$.

Note that the rings defined in 4.9 are finitely generated over \mathbf{Z} . We sum up the results in this section in a concluding remark:

Theorem 4.10. Consider in the following the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left module.

Let $n < 0$ and $k \geq 1$. The ring $B_{k,n}^L$ has the following property: Any ring-homomorphism $f_1 : B_{k,n}^L \rightarrow A$ gives rise to a map

$$(4.10.1) \quad \phi_{f_1} : \mathcal{O}(n-k)^{k+1} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Let $0 \leq n \leq k$ and consider the ring $B_{k,n}^L$: Any ring-homomorphism $f_2 : B_{k,n}^L \rightarrow A$ gives rise to a map

$$(4.10.2) \quad \phi_{f_2} : \mathcal{O}^{n+1} \oplus \mathcal{O}(-k-1)^{k-n} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Let $n \in \mathbf{Z}$ and $k \geq 1$ and consider the ring $B_{k,n}^R$: Any ring-homomorphism $f_3 : B_{k,n}^R \rightarrow A$ gives rise to a map

$$(4.10.3) \quad \phi_{f_3} : \mathcal{O}(n) \oplus \mathcal{O}(n-k-1)^k \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of right $\mathcal{O}_{\mathbf{P}_A^1}$ -modules.

Proof. The theorem follows from the constructions in this section. \square

5. SPLITTING OVER FIELDS OF CHARACTERISTIC ZERO

In this section we look at the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left and right \mathcal{O} -module on the projective line for all $n \in \mathbf{Z}$ and $k \geq 1$ over any field F of characteristic zero. We use the results of section 2 and techniques developed in section 3 to give the splitting-type of the principal parts as left and right module over any field of characteristic zero, settling completely the problem of describing the structure of the principal parts on the projective line. We first consider the splitting-type of the principal parts as left module.

Theorem 5.1. *Consider the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left module on \mathbf{P}_F^1 with $n < 0$ and $k \geq 1$ or $1 \leq k \leq n$. There exists an isomorphism*

$$\mathcal{O}(n-k)^{k+1} \cong \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_F^1}$ -modules.

Proof. The case where $1 \leq k \leq n$ is proved in [5] Proposition 6.3. Consider the case where $n < 0$ and $k \geq 1$. Let $0 \leq l \leq k$. According to Theorem 4.10, there exists a ring $B_{k,n}^L$ with the property that maps of rings

$$f : B_{k,n}^L \rightarrow F$$

give rise to a map

$$\phi : \mathcal{O}(n-k)^{k+1} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_F^1}$ -modules. To define a map of rings $f : B_{k,n}^L \rightarrow F$ is by definition of $B_{k,n}^L$ equivalent to the following: We seek solutions in F to the following system of equations $A_l^1 \mathbf{x}_1^1 = b_l^1$:

$$\begin{pmatrix} \binom{k+m-l-1}{m-1} & \binom{k+m-l-1}{m} & \binom{k+m-l-1}{m+1} & \cdots & \binom{k+m-l-1}{m+l-1} \\ \binom{k+m-l}{m-1} & \binom{k+m-l}{m} & \binom{k+m-l}{m+1} & \cdots & \binom{k+m-l}{m+l-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{k+m-2}{m-1} & \binom{k+m-2}{m} & \binom{k+m-2}{m+1} & \cdots & \binom{k+m-2}{m+l-1} \\ \binom{k+m-1}{m-1} & \binom{k+m-1}{m} & \binom{k+m-1}{m+1} & \cdots & \binom{k+m-1}{m+l-1} \end{pmatrix} \begin{pmatrix} x_l^l \\ x_{l-1}^l \\ \vdots \\ x_1^l \\ x_0^l \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$

for all $l = 0, \dots, k$. By Lemma 6.1 The determinant of A_l^1 is

$$|A_l^1| = \frac{\binom{k-l+m-1}{m-1} \binom{k-l+m}{m} \cdots \binom{k+m-1}{m-1+l}}{\binom{k-l}{0} \binom{k-l+1}{1} \cdots \binom{k}{l}}$$

which is non-zero for all l since the characteristic of F is zero, hence there exists a map

$$\phi : \mathcal{O}(n-k)^{k+1} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left \mathcal{O} -modules. According to the proof of Proposition 4.2 we get for all $l = 0, \dots, k$ maps defined as follows:

$$(5.1.1) \quad \phi_l^0(x_0^{n-k}) = \sum_{j=0}^{k-l} (-1)^j f(\bar{c}_j^{l,1}) t^{k-l-1} dt^j \otimes x_0^n$$

and

$$(5.1.2) \quad \phi_l^1(x_1^{n-k}) = \sum_{p=0}^l f(\bar{x}_{l-p}^{l,1}) \frac{1}{t^{l-p}} ds^p \otimes x_1^n,$$

Let $\phi = \oplus_{i=0}^k \phi_i$. Consider the maps $\phi_i = \phi|_{U_i}$ for $i = 0, 1$. The determinant of ϕ_0 equals

$$|\phi_0| = \pm f(c_k^0 c_{k-1}^1 c_{k-2}^2 \cdots c_0^k)$$

which is nonzero by construction, hence ϕ_0 is an isomorphism. The determinant $|\phi_1|$ equals

$$|\phi_1| = f(x_0^0 x_0^1 x_0^2 \cdots x_0^k)$$

which is nonzero by induction. It follows that ϕ_1 is an isomorphism, and the theorem follows. \square

Theorem 5.2. *Consider the principal parts $\mathcal{P}^k(\mathcal{O}(n))$ as left $\mathcal{O}_{\mathbf{P}_F^1}$ -module on \mathbf{P}_F^1 with $0 \leq n < k$: There exists an isomorphism*

$$(5.2.1) \quad \mathcal{O}^{n+1} \oplus \mathcal{O}(-k-1)^{k-n} \cong \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_F^1}$ -modules.

Proof. Consider the ring $B_{k,n}^L$ from Theorem 4.10. To give a ring homomorphism

$$f : B_{k,n}^L \rightarrow F$$

is by definition of $B_{k,n}^L$ equivalent to the following: We seek solutions in F to the following systems of linear equations: $A_l^2 \mathbf{x}_1^2 = b_l^2$

$$\begin{pmatrix} \binom{n}{n} & -\binom{n-1}{n-1} & \binom{n-2}{n-2} & \cdots & (-1)^l \binom{n-l}{n-l} \\ \binom{n}{n-1} & -\binom{n-1}{n-2} & \binom{n-2}{n-3} & \cdots & (-1)^l \binom{n-l}{n-l-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{n}{n-l+1} & -\binom{n-1}{n-l} & \binom{n-2}{n-l-1} & \cdots & (-1)^l \binom{n-l}{n-2l+1} \\ \binom{n}{n-l} & -\binom{n-1}{n-l-1} & \binom{n-2}{n-l-2} & \cdots & (-1)^l \binom{n-l}{n-2l} \end{pmatrix} \begin{pmatrix} x_l^{l,2} \\ x_{l-1}^{l,2} \\ \vdots \\ x_1^{l,2} \\ x_0^{l,2} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}$$

and the system $B_m^2 \mathbf{y}_m^2 = c_m^2$

$$\begin{pmatrix} \binom{k-n-1}{0} & \binom{k+n-1}{k-n-2} & \binom{k-n-1}{2} & \cdots & \binom{k-n-1}{k-n-2} \\ \binom{k-n-2}{0} & \binom{k+n-1}{1} & \binom{k-n-1}{2} & \cdots & \binom{k-n-1}{m} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{k-n-m}{0} & \binom{k-n-m}{1} & \binom{k-n-m}{2} & \cdots & \binom{k-n-m}{m} \\ \binom{k-n-m-1}{0} & \binom{k-n-m-1}{1} & \binom{k-n-m-1}{2} & \cdots & \binom{k-n-m-1}{m} \end{pmatrix} \begin{pmatrix} x_m^{m,3} \\ x_{m-1}^{m,3} \\ \vdots \\ x_1^{m,3} \\ x_0^{m,3} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$

where $l = 0, \dots, n$ and $m = 0, \dots, k - n - 1$. The determinant of A_l^2 is by Lemma 6.2 equal to $(-1)^{l+1}$, hence the system $A_l^2 \mathbf{x}_l^2 = b_l^2$ has solutions in any field F of characteristic zero. The determinant of the matrix B_m^2 equals

$$|B_m^2| = \frac{\binom{k-n-1}{0} \binom{k-n}{1} \dots \binom{k-n+m-1}{m}}{\binom{k-n-1}{0} \binom{k-n}{1} \dots \binom{k-n+m-1}{m}} = 1$$

which is different from zero in F . It follows that the system $B_m^2 \mathbf{y}_m^2 = c_m^2$ always has solutions in F , hence we get a map

$$\phi : \mathcal{O}^{n+1} \oplus \mathcal{O}(-k-1)^{k-n} \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_F^1}$ -modules defined as follows: According to the proof of Proposition 4.4 and 4.6 we define maps ϕ_l^i for $l = 0, \dots, n$ on U_0 as follows:

$$(5.2.2) \quad \phi_l^0(e) = \sum_{j=0}^{n-l} f(\bar{c}_j^{l,2}) t^{n-l-j} dt^j \otimes x_0^n.$$

On U_1 we define:

$$(5.2.3) \quad \phi_l^1(f) = \sum_{p=0}^l f(\bar{x}_{l-p}^{l,2}) t^{p-l} ds^p \otimes x_1^n.$$

We define for $0 \leq l \leq k - n - 1$ maps ψ_l^i as follows:

$$(5.2.4) \quad \psi_l^0(x_0^{-k-1}) = \sum_{j=k+1}^{k-l} (-1)^j f(\bar{c}_j^{l,3}) t^{k-l-j} dt^j \otimes x_0^n$$

On U_1 :

$$(5.2.5) \quad \psi_l^1(x_1^{-k-1}) = \sum_{i=0}^l f(\bar{x}_{l-i}^{l,3}) t^{i-l} ds^{n+1+i} \otimes x_1^n.$$

Let $\phi = \bigoplus_{i=0}^n \phi_l \oplus \bigoplus_{i=0}^{k-n-1} \psi_i$. We claim that ϕ is an isomorphism. We consider the map $\phi|_{U_0}$: Its determinant equals

$$|\phi|_{U_0} = \pm f(c_n^{0,2} c_{n-1}^{1,2} \dots c_0^{n,2} c_k^{0,3} c_{k-1}^{1,3} \dots c_{n+1}^{k-n-1,3})$$

which is different from zero by construction, hence $\phi|_{U_0}$ is an isomorphism. We next consider the map $\phi|_{U_1}$. Its determinant equals

$$|\phi|_{U_1} = f(x_0^{0,2} x_0^{1,2} \dots x_0^{n,2} x_0^{0,3} x_0^{1,3} \dots x_0^{k-n-1,3})$$

which is nonzero by induction, hence $\phi|_{U_1}$ is an isomorphism, and the theorem is proved. \square

Note that the determinants of the matrices A_l^2 and B_m^2 are ± 1 , hence the statement of the theorem is valid over any field.

We get a similar theorem on the splitting-type of the principal parts as right module:

Theorem 5.3. *Let F be a field of characteristic zero. Consider $\mathcal{P}^k(\mathcal{O}(n))$ as right \mathcal{O} -module on \mathbf{P}_F^1 for all $n \in \mathbf{Z}$ and $k \geq 1$. There exists an isomorphism*

$$\mathcal{P}^k(\mathcal{O}(n)) \cong \mathcal{O}(n) \oplus \mathcal{O}(n-k-1)^k$$

of right \mathcal{O} -modules.

Proof. Existence of a map

$$\phi_{-1} : \mathcal{O}(n) \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

is trivial. According to Theorem 4.10 there exists a ring $B_{k,n}^R$ with the property that a ring homomorphism $f : B_{k,n}^R \rightarrow F$ gives rise to a map

$$\mathcal{O}(n) \oplus \mathcal{O}(n-k-1)^k \rightarrow \mathcal{P}^k(\mathcal{O}(n))$$

of left $\mathcal{O}_{\mathbf{P}_F^1}$ -modules. Existence of the map f is by definition equivalent to the existence of solutions in F of the systems of linear equations $A_l^3 \mathbf{x}_1^3 = b_l^3$

$$\begin{pmatrix} \binom{k-1}{0} & \binom{k-1}{k-2} & \binom{k-1}{1} & \binom{k-1}{2} & \dots & \binom{k-1}{l} \\ \binom{k-2}{0} & \binom{k-2}{k-3} & \binom{k-2}{1} & \binom{k-2}{2} & \dots & \binom{k-2}{l} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{k-l}{0} & \binom{k-l}{k-l-1} & \binom{k-l}{1} & \binom{k-l}{2} & \dots & \binom{k-l}{l} \end{pmatrix} \begin{pmatrix} y_{l-1}^{l,1} \\ y_l^{l,1} \\ \vdots \\ y_1^{l,1} \\ y_0^{l,1} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix},$$

where $l = 0, \dots, k-1$. The determinant of the matrix A_l^3 equals

$$|A_l^3| = \frac{\binom{k-1}{0} \binom{k}{1} \dots \binom{k+l-1}{l}}{\binom{k-1}{0} \binom{k}{1} \dots \binom{k+l-1}{l}}$$

which is different from zero, hence the map ϕ exists over the field F . According to the proof of Proposition 4.8 we may define for $0 \leq l \leq k-1$ maps ψ_l as follows: On U_0 we define

$$(5.3.1) \quad \phi_l^0(x_0^{n-k-1}) = \sum_{j=0}^{k-l} (-1)^j f(\bar{d}_j^{l,1}) u^{k-l-j} du^j \otimes x_0^n.$$

Define a map on U_1 :

$$(5.3.2) \quad \phi_l^1(x_1^{n-k-1}) = \sum_{i=0}^l f(\bar{y}_{l-i}^{l,1}) u^{i-l} dv^{i+1} \otimes x_1^n,$$

Define $\phi = \oplus_{i=-1}^{k-1} \phi_i$. The determinant of $\phi|_{U_0}$ equals

$$|\phi|_{U_0}| = \pm f(\bar{d}_k^{0,1} \bar{d}_{k-1}^{1,1} \dots \bar{d}_j^{k-1,1})$$

which is non-zero by construction, hence $\phi|_{U_0}$ is an isomorphism. The determinant of $\phi|_{U_1}$ equals

$$|\phi|_{U_1}| = f(\bar{y}_0^{0,1} \bar{y}_0^{1,1} \dots \bar{y}_0^{k-1,1}),$$

which is non-zero by induction, hence $\phi|_{U_1}$ is an isomorphism, and the theorem is proved. \square

Note that the determinant of the matrix A_l^3 from the proof of Theorem 3.2.1 is one, hence the result of the theorem is valid over an arbitrary field.

The Theorems 5.1, 5.2 and 5.3 generalize Proposition 6.3 and Theorem 7.1 in [5], settling completely the problem of describing the principal parts on the projective line over a field of characteristic zero. Since formation of principal parts commute with direct sums and because on the projective line over any field any finite rank locally free sheaf splits into invertible sheaves ([3],[4]), we have described the splitting-type of the principal parts of any locally free finite rank sheaf as left and right module over any field of characteristic zero, generalizing results obtained

in [5],[9],[11] and [14]. Note that the ring $B_{k,n}^L$ contains all information necessary to split the sheaf of principal parts over any field for all $n \leq 0$, $k \geq 1$ or $1 \leq k \leq n$. This problem will be studied in future papers.

6. APPENDIX: CALCULATIONS OF DETERMINANTS

In this section we calculate some determinants appearing in section 3 and 4.

Lemma 6.1. *Assume a, b are integers with $b \leq a$. Then the following is true for all $l \geq 1$: The determinant of the matrix*

$$(6.1.1) \quad M = \begin{pmatrix} \binom{a}{b} & \binom{a}{b+1} & \binom{a}{b+2} & \cdots & \binom{a}{b+l} \\ \binom{a+1}{b} & \binom{a+1}{b+1} & \binom{a+1}{b+2} & \cdots & \binom{a+1}{b+l} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{a+l-1}{b} & \binom{a+l-1}{b+1} & \binom{a+l-1}{b+2} & \cdots & \binom{a+l-1}{b+l} \end{pmatrix}$$

equals

$$(6.1.2) \quad |M| = \frac{\prod_{i=0}^l \binom{a+i}{b+i}}{\prod_{i=0}^l \binom{a-b+i}{i}}$$

Proof. One easily shows the following formula:

$$(6.1.3) \quad \binom{a+i}{b+j} - \frac{a+i+1-b-l}{a+i+1} \binom{a+i+1}{b+j} = \frac{l-j}{b+j} \binom{a+i}{b-1+j}.$$

Apply equation 6.1.3 to the matrix 6.1.1 to get the matrix

$$(6.1.4) \quad \begin{pmatrix} \frac{l}{b} \binom{a}{b-1} & \frac{l-1}{b+1} \binom{a}{b} & \frac{l-2}{b+2} \binom{a}{b+1} & \cdots & \frac{1}{b+l-1} \binom{a}{b+l-2} & 0 \\ \frac{l}{b} \binom{a+1}{b-1} & \frac{l-1}{b+1} \binom{a+1}{b} & \frac{l-2}{b+2} \binom{a+1}{b+1} & \cdots & \frac{1}{b+l-1} \binom{a+1}{b+l-2} & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & 0 \\ \frac{l}{b} \binom{a+l-1}{b-1} & \frac{l-1}{b+1} \binom{a+l-1}{b} & \frac{l-2}{b+2} \binom{a+l-1}{b+1} & \cdots & \frac{1}{b+l-1} \binom{a+l-1}{b+l-2} & 0 \\ \binom{a+l}{b} & \binom{a+l}{b+1} & \binom{a+l}{b+2} & \cdots & \binom{a+l}{b+l-1} & \binom{a+l}{b+l} \end{pmatrix}$$

We get the following determinant

$$\frac{l!}{b(b+1)(b+2) \cdots (b+l-2)} \binom{a+l}{b+l} |N|,$$

where N is the matrix

$$(6.1.5) \quad \begin{pmatrix} \binom{a}{b-1} & \binom{a}{b} & \binom{a}{b+1} & \cdots & \binom{a}{b+l-2} \\ \binom{a+1}{b-1} & \binom{a+1}{b} & \binom{a+1}{b+1} & \cdots & \binom{a+1}{b+l-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{a+l-1}{b-1} & \binom{a+l-1}{b} & \binom{a+l-1}{b+1} & \cdots & \binom{a+l-1}{b+l-2} \end{pmatrix}$$

By induction we get the following formula:

$$\frac{l!}{b(b+1)(b+2) \cdots (b+l-2)} \binom{a+l}{b+l} \frac{\binom{a}{b-1} \cdots \binom{a+l-1}{b+l-2}}{\binom{a-b+2}{1} \binom{a-b+3}{2} \cdots \binom{a-b+l}{l-1}},$$

and the formula 6.1.2 follows. \square

Lemma 6.2. *Let a be an integer with $a \geq 0$. Then the following is true: The determinant of the matrix*

$$(6.2.1) \quad P = \begin{pmatrix} \binom{a}{a} & \binom{a-1}{a-1} & \binom{a-2}{a-2} & \cdots & \binom{a-l}{a-l} \\ \binom{a}{a-1} & \binom{a-1}{a-2} & \binom{a-2}{a-3} & \cdots & \binom{a-l}{a-l-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{a}{a-l+1} & \binom{a-1}{a-l} & \binom{a-2}{a-l-1} & \cdots & \binom{a-l}{a-2l+1} \\ \binom{a}{a-l} & \binom{a-1}{a-l-1} & \binom{a-2}{a-l-2} & \cdots & \binom{a-l}{a-2l} \end{pmatrix}$$

equals

$$(6.2.2) \quad |P| = (-1)^{l+1}$$

Proof. The matrix P equals the matrix

$$(6.2.3) \quad P = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ a & a-1 & a-2 & \cdots & a-l \\ \frac{a(a-1)}{2!} & \frac{(a-1)(a-2)}{2!} & \frac{(a-2)(a-3)}{2!} & \cdots & \frac{(a-l)(a-l-1)}{2!} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{a}{a-l} & \binom{a-1}{a-l-1} & \binom{a-2}{a-l-2} & \cdots & \binom{a-l}{a-2l} \end{pmatrix}.$$

We get the Vandermonde-matrix

$$(6.2.4) \quad \frac{1}{1!2!\cdots l!} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ a & a-1 & a-2 & \cdots & a-l \\ a^2 & (a-1)^2 & (a-2)^2 & \cdots & (a-l)^2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a^l & (a-1)^l & (a-2)^l & \cdots & (a-l)^l \end{pmatrix},$$

which is easily seen to have determinant $(-1)^{l+1}$, and the lemma follows. \square

Acknowledgments. This paper was written in the period autumn 2003- spring 2004 and I want to thank Dan Laksov for comments on the problems discussed in this paper. Thanks also to Andrei Zelevinsky. It is also a great pleasure to thank colleagues and staff at the Royal Institute of Technology for providing a friendly and inspiring atmosphere during its preparation.

REFERENCES

- [1] M. F. Atiyah, Complex analytic connections in fiber bundles, *Trans. AMS* no. 85 (1957)
- [2] A. Grothendieck, Éléments de géométrie algébrique IV₄, Étude locale des schémas et des morphismes de schémas, *Publ. Math. IHES* 32 (1967)
- [3] A. Grothendieck, Sur la classification des fibrés holomorphes sur la sphère de Riemann, *Amer. J. Math.* 79 (1957)
- [4] G. Harder, Halbeinfache Gruppenschemata über vollständigen Kurven, *Invent. Math.* 6 (1968)
- [5] H. Maakestad, Modules of principal parts on the projective line, accepted for publication in *Arkiv för Matematik*
- [6] H. Maakestad, A note on principal parts on projective space and linear representations, accepted for publication in *Proc. Amer. Math. Soc.*
- [7] P. Jahnke, I. Radloff, Splitting jet sequences, *preprint math.AG/0210454*
- [8] D. Laksov, A. Thorup, Weierstrass points on schemes, *J. Reine Angew. Math* no. 460 (1995)
- [9] D. Perkinson, Principal parts of line bundles on toric varieties, *Compositio Math.* 104 (1996)
- [10] D. Perkinson, Curves in Grassmannians, *Trans. Amer. Math. Soc.* no. 347 (1995)
- [11] R. Piene, G. Sacchiero, Duality for rational normal scrolls, *Comm. Algebra* 12 (1984)

- [12] R. Piene, Numerical characters of a curve in projective n -space, *Real and complex singularities*, *Proc. ninth nordic summer school Oslo* (1976)
- [13] W. F. Pohl, Differential geometry of higher order, *Topology* no. 1 (1962)
- [14] S. di Rocco, A. J. Sommese, Line bundles for which a projectivized jet bundle is a product, *Proc. Amer. Math. Soc.* 129, (2001)
- [15] C. Widland, R. Lax, Weierstrass points on Gorenstein curves, *Pacific J. Math* no. 142 (1990)

DEPT. OF MATHEMATICS, KTH, STOCKHOLM
E-mail address: `makestad@math.kth.se`