

ON DIRECTED ZERO-DIVISOR GRAPHS OF FINITE RINGS

TONGSUO WU

Department of Mathematics, Shanghai Jiaotong
University, Shanghai 200030, P. R. China

ABSTRACT. For a finite ring R , the directed zero-divisor graph $\Gamma(R)$ is connected if and only if there is no end vertex (sink and source); sinks and sources are characterized and clarified. This article also shows that $\Gamma(R)$ can not be a network for any ring R .

1. INTRODUCTION

For any noncommutative ring R , let $Z(R)$ be the set of (one-sided) zero-divisors of R . The directed zero-divisor graph of R is a directed graph $\Gamma(R)$ with vertices $Z(R)^* = Z(R) - \{0\}$, where for distinct vertices x and y of $Z(R)^*$ there is a directed edge from x to y if and only if $xy = 0$ ([6]). This is a generalization of zero-divisor graph of commutative rings. The concept of a zero-divisor graph of a commutative ring was introduced in [4], and it was mainly concerned with colorings of rings there. In [2], the vertex set of $\Gamma(R)$ is chosen to be $Z(R)^*$ and the authors study interplay between the ring-theoretic properties of commutative ring R and the graph-theory property of $\Gamma(R)$. The zero-divisor graph of a commutative ring is also studied by several other authors, see [3] for a list of references. The zero-divisor graph has been also introduced and studied for semigroups in [5].

In this paper, we study the directed zero-divisor graph of noncommutative rings and we focus our attention on finite rings. In section 2, we prove that a finite ring R has connected zero-divisor graph if and only if one-sided identity of R (if exists)

1991 *Mathematics Subject Classification.* 16N60, 05C15.

Key words and phrases. finite ring, zero-divisor graph, connectedness, sink, source.

is two-sided, if and only if $\Gamma(R)$ contains no end vertex (i.e., sinks and sources). For any distinct vertices x, y of a finite ring with proper one-sided identity, the directed distance from x to y is less than 7, if a directed path exists from x to y . In section 3 and 4, we study sinks and sources of finite and infinite ring. We proved that for *any* ring R , if there exists a source b in $\Gamma(R)$ with $b^2 = 0$, then $|R| = 4$ and $R = \{0, a, b, c\}$, where a and c are left identity elements and $ba = 0 = bc$. Such a ring R is also the only ring such that $\Gamma(R)$ has exactly one source. The dual result for sink vertex is also true. This result is key to clarify sinks and sources in $\Gamma(R)$. In Section 3, we show that for a finite ring R with at least five elements, sink and source can not coexist in $\Gamma(R)$. For finite ring R with at least five elements, sinks (sources) are characterized by strongly right (left) invertible elements. In section 4, we show that for any ring R with at least five elements, there are only four possibilities for the semigroups (when non-empty) $Sink(R)$, the set of sinks of $\Gamma(R)$, and $Sour(R)$, the semigroup of all sources:

- (1) $Sink(R) = \emptyset$ and $Sour(R) = \emptyset$;
- (2) $Sink(R) = \emptyset$ and $|Sour(R)| \geq 2$;
- (3) $Sour(R) = \emptyset$ and $|Sink(R)| \geq 2$;
- (4) $|Sour(R)| = \infty = |Sink(R)|$.

Therefore $\Gamma(R)$ can not be a network for any ring R .

2. CONNECTEDNESS AND DIAMETERS

Lemma 2.1. *Suppose that every right identity element of a finite ring R is a two-sided identity. We have*

- (1) *Each left zero-divisor is also a right zero-divisor;*
- (2) *If in addition $|R| \geq 5$, then for distinct $a, b \in R^*$ with $ab = 0$, there exists $c \in R^*$ such that $ca = 0$ and $c \neq a$.*

Proof. (1) If $ab = 0$ and $xa \neq 0$ for all $x \in R$, then $Ra = R$ and there is a right identity element $e \in R$. Then e is also a left identity. Let $ga = e$, then $0 = gab = eb = b$, a contradiction.

(2) Suppose that $ann_l(a) = \{a, 0\}$, then for distinct $b, c, d \in R^* - \{a\}$, we have $daa = 0, da \neq 0$. Thus $da = a$. Then we have $b - c = b - d$ and therefore, $c = d$. Then $|R| \leq 4$. This completes the proof. QED

The following lemma is a special case of a recent result of [7, Section 4]. For

completeness we present our original proof here:

Lemma 2.2. *Suppose $|R| \leq 4$ and every proper one-sided identity element is a two-sided identity. Then for distinct $a, b \in R^*$ with $ab = 0$, we have $ba = 0$.*

Proof. If R has three elements, then for $a \neq b$ with $ab = 0$, one has $(a + b)b = 0$. Thus $ba = b(a + b) = 0$.

Now suppose that $R = \{0, a, b, a + b\}$ has four elements. Without loss, we assume $ab = 0$. If R has identity element 1, assume $a + b = 1$. Then $a^2 = a, b^2 = b$. So if $ba = 1$, then $a = aba = 0$; if $ba = a$, then $a = ba = b^2a = 0$; If $ba = b$, then $b = b^2a = 0$. There is contradiction in each case. Hence $ba = 0$. In the remaining part of the proof, assume that R has no one-sided identity element. Without loss, we can assume $2a = 2b = 0$ (The only other case is that the additive group of R is cyclic of order four.). Now we show that $ab = 0$ implies $ba = 0$:

(1) $ba \neq a$: If $ba = a$, then $a^2 = 0$. In this case, we assert $b^2 = 0$. Actually, $b^2 \neq b$ since otherwise b is a left identity element of R ; $b^2 \neq a$ since otherwise $a = ba = b^2a = a^2 = 0$; $b^2 \neq a + b$ since otherwise, $b^2 = (a + b)b = b^3 = b(a + b) = a + (a + b) = b$. But $b^2 = 0$ implies $a = ba = b^2a = 0$, a contradiction.

(2) $ba \neq b$: If $ba = b$, then $b^2 = 0$. Then $a^2 \neq a$ since otherwise $a(a + b) = a$, $b(a + b) = b$, $(a + b)^2 = a + b$, i.e., $a + b$ is a right identity element of R ; $a^2 \neq 0$ since $b = ba^2$; $a^2 \neq b$ since otherwise $b = ba = ba^2 = b^2 = 0$. Finally, $a^2 = a + b$ and we have $a^2 = a + b = a^3 = a^2 + ba = a^2 + b$, contradicting with the assumption $b \neq 0$.

(3) $ba \neq a + b$: If $ba = a + b$, then $a^2 = 0$, $0 = ba^2 = a^2 + ba$, $0 = ba = a + b$, another contradiction. This completes the proof of $ba = 0$, for rings without one-sided identity element. QED

Let K_i be the complete directed graph with i vertices. By Lemma 2.2, we have a list of all possibilities of $\Gamma(R)$ for rings R with $|R| \leq 4$:

- (1) $|R| = 2$: $K_i, i = 0, 1$
- (2) $|R| = 3$: $K_i, i = 0, 2$
- (3) $|R| = 4$: K_i ($i = 0, 1, 2, 3$); $\circ \rightleftharpoons \circ \rightleftharpoons \circ$; $\circ \rightarrow \circ \leftarrow$; $\circ \leftarrow \circ \rightarrow \circ$.

We remark that every graph in the list can be realized as the zero-divisor graph of some ring R with $|R| \leq 4$.

Proposition 2.3. *Suppose that R is a finite ring with the property that every one-sided identity element is a two-sided identity in R . Then for any path $a \rightarrow b$ in $\Gamma(R)$, there is a walk $c \rightarrow a \rightarrow b \rightarrow d$, where $c \neq a$ and $b \neq d$.*

A vertex g in a directed graph G is called a *sink*, if the in-degree of g is positive and the out-degree of g is zero. The dual concept of sink is called *source*.

Let $Z_r(R)$ be the set of right zero divisors. In [6], it is proved that for any ring R , $\Gamma(R)$ is connected if and only if $Z_r(R) = Z_l(R)$. It is also proved that $\Gamma(R)$ is connected for all finite rings with two sided-identity element. We now characterize all finite rings R whose directed zero-divisor graph $\Gamma(R)$ is connected:

Theorem 2.4. *For any finite ring R , the following statements are equivalent:*

- (1) *The directed zero-divisor graph $\Gamma(R)$ is connected;*
 - (2) *Every one-sided identity element of R is the two-sided identity of R ;*
 - (2') *Either R has two-sided identity or R has no proper one-sided identity;*
 - (3) *There exists no end-vertex (sink or source) in $\Gamma(R)$.*
- If $\Gamma(R)$ is connected, then $\text{diam}(\Gamma(R)) \leq 3$.*

Proof. (1) \implies (2) and (3) \implies (2). If R has a left identity e that is not a right identity element of R , then there exists $a \in R$ such that $ae - a \neq 0$. In this case $ae - a \rightarrow b$ for all $b \in R^*$. Then there exist at least two left identity elements in R , say e and $f = e + ae - a$. Then e and f are sink vertices in $\Gamma(R)$. So the directed graph $\Gamma(R)$ is not connected.

(2) \implies (3) and (2) \implies (1). Assume that every one-sided identity element of R is the two-sided identity of R . If $|R| < 5$, then by Lemma 2.2, we know that there is no sink (source) vertex in $\Gamma(R)$ and that $\Gamma(R)$ is connected. In what follows, we assume $|R| \geq 5$. Then by Lemma 2.1, if $a \rightarrow b$ for distinct vertices a, b , then there exist $c, d \in R^*$ such that $c \neq a, d \neq b$ and in $\Gamma(R)$ there is a walk $c \rightarrow a \rightarrow b \rightarrow d$. So there is no sink or source vertex in $\Gamma(R)$. This proves (3). Now we use the proof of corresponding result in [2] to finish our proof: For any distinct $x, y \in Z(R)^*$, if $xy = 0$, then $d(x, y) = 1$. If $xy \neq 0$, then again by Lemma 2.1, there exists $a \neq x, b \neq y$ such that $xa = 0 = by$. If $a = b$, then $x \rightarrow a \rightarrow y$ and $d(x, y) = 2$; If $a \neq b$ and $ab = 0$, then we have $x \rightarrow a \rightarrow b \rightarrow y$ and $d(x, y) \leq 3$; If $a \neq b$ and $ab \neq 0$, then $ab \neq x, ab \neq y$ since $xy \neq 0$, and there is a path $x \rightarrow ab \rightarrow y$. In all cases, $d(x, y) \leq 3$. QED

Now let R be a ring with proper left identity element e . Denote

$$I_e = \{a \in R | ae = 0\}, R_e = \{a \in R | a = ae\}.$$

Then (1) $I_e = \text{ann}_l(e)$ and I_e is a two-sided ideal of R with at least two elements;

- (2) R_e is a subring of R with identity e ;
- (3) $R = R_e \oplus I_e$ as left R -module;
- (4) There is a ring isomorphism $R_e \cong R/I_e$.

For any subset M, N of $Z(R)^*$, let $\Gamma(M, N)$ be the induced bipartite subgraph of $\Gamma(R)$ and denote its edge set as $E(M, N)$. For the graph $\Gamma(R)$, denote its edge set as $E(R)$. If there is a cycle, we add 2 to the number of the edges of $\Gamma(R)$. Then the graph $\Gamma(R)$ is completely determined by the following four induced subgraphs: $\Gamma(R_e)$, complete directed graph $\Gamma(I_e)$, bipartite graphs $\Gamma(I_e^*, R_e^*)$ and $\Gamma(R_e^*, R_e * \oplus I_e^*)$. We have

Proposition 2.5. *For a ring R with proper left identity element e , the graph $\Gamma(R)$ has the following properties:*

- (1) For any $a \in I_e$, the out-degree of a is $|R| + 1$;
- (2) The number of vertices of $\Gamma(R)$ is $|R| - 1 = |R_e||I_e| - 1$;
- (3) $|E(R)| = |I|(|I| - 1 + E(R_e^*, I_e^*) + E(R_e^*, R_e^* \oplus I_e^*) + (2 - |I|)E(R_e))$.

Proof. (1) The number of directed edges from I_e^* to R^* is $|R|(|I| - 1)$;

(2) The number of directed edges from $R^* - I_e^*$ to I_e^* is

$$|I|(|E(R_e^*, I_e^*)| - (|I| - 1)(|R_e| - 1));$$

(3) An element of $R^* - I_e^*$ has the form of $a_i + x$, where $x = 0$ or $x = b_j$. Besides, $(a_i + x)(a_j + y) = a_i(a_j + y)$. Thus the number of directed edges from $R^* - I_e^*$ to $R^* - I_e^*$ is

$$|I|(|E(R_e)| + |E(R_e^*, R_e^* \oplus I_e^*)| - (|I| - 1)|E(R_e)|).$$

Note that $|R| = |R_e||I_e|$, then we obtain our formula. QED

Proposition 2.6. *Let R be a finite ring with proper one-sided identity. For distinct elements x, y in R , either $d(x, y) = \infty$ or $d(x, y) \leq 6$.*

Proof. Let R be a ring with proper left identity and assume that e is a left identity of R . Then $R^* = K \cup I_e^* \cup R_e^*$ and this is a disjoint union where we assume

$$R_e^* = \{a_i | i = 1, 2, \dots, m\}, I_e^* = \{b_j | j = 1, 2, \dots, n\}$$

and $K = I_e^* \oplus R_e^*$. For distinct $x, y \in R^*$, assume that there exists a directed path from x to y in $\Gamma(R)$, say, $x \rightarrow x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_r \rightarrow y$.

(1) If $x \in I_e$, then $x \rightarrow z$ for any z . In this case, $d(x, y) = 1$.

(2) Assume $x \in R_e$. If $0 \in xI_e^*$, then we have a path $x \rightarrow I_e^* \rightarrow y$, $d(x, y) \leq 2$; If $0 \notin xI_e^*$, then $0 \notin xK$. Without loss, we can assume $x_i \in R_e$ for $i = 1, 2, 3, 4$. Since R_e is a ring with two-sided identity, $\text{diam}(R) \leq 3$. So there exists a path $x \rightarrow x_1 \rightarrow x_2 \rightarrow x_3 \rightarrow I_e^* \rightarrow y$. In this case, $d(x, y) \leq 5$.

(3) The final case is $x \in K$. Assume $x = a_1 + b_1$. If $0 \in xI_e^*$, then $d(x, y) \leq 2$; If $0 \notin xI_e^*$, then $x_1 \in R_e^* \cup K$. If $x_1 \in K$, then we have $0 = (a_1 + b_1)(a_s + b_t) = a_1a_s + a_1b_t$. Then we have a path $x \rightarrow b_t \rightarrow y$, and $d(x, y) \leq 2$; Hence we can assume $x_i \in R_e^*$ for $i = 1, 2, 3, 4$. and $x_4 \rightarrow I_e^*$. We have a path $x \rightarrow x_1 \rightarrow x_2 \rightarrow x_3 \rightarrow x_4 \rightarrow I_e^* \rightarrow y$ and hence $d(x, y) \leq 6$. This completes the proof. QED

Corollary 2.7. *For any ring R with proper one-sided identity element e , R_e is a subring of R and e is a two-sided identity of R_e , and $\text{diam}(R) \leq 3 + \text{diam}(R_e)$.*

We end this section with the following examples:

Example 2.8 For any field F , let R be the n by n full matrix ring over F ($n > 1$). Then the diameter of $\Gamma(R)$ is 2.

We need only to prove the following facts: for any $A, B \in Z(R)^*$, there exists $C \in Z(R)^*$ such that $AC = 0$ and $CB = 0$. First, there exist invertible matrices $P, Q \in R$ such that the last column of AP is zero and the first row of QB is zero. Second, let

$$C = P \begin{pmatrix} 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ 1 & 0 & \cdots & 0 \end{pmatrix} Q,$$

then we have $C \neq 0$, $AC = 0$, $CB = 0$.

Example 2.9 Let R be the n by n full matrix ring over $\mathbb{Z}/2\mathbb{Z}$. Let S be the non-unitary subring of R consisting of those matrices all of whose rows are zero except the first row. Then $\text{Sink}(S) = \left\{ \begin{pmatrix} 1 & \alpha \\ 0 & 0 \end{pmatrix} \mid \alpha \right\}$, and $\text{Sink}(S)$ consists of all left identity elements of S . $I_e = \left\{ \begin{pmatrix} 0 & \alpha \\ 0 & 0 \end{pmatrix} \mid \alpha \right\}$ for $e = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. $|S| = 2^n$, $|I_e| = |\text{Sink}(S)| = 2^{n-1}$. $\Gamma(S)$ is a star-like directed graph whose kernel is the complete graph $K_{2^{n-1}-1}$, each vertex of which also connects to 2^{n-1} sinks.

Example 2.10 Let $R = \mathbb{Z}/n\mathbb{Z}$ and denote $S = \left\{ \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \mid a, b \in R \right\}$. Then $Sink(S) = \left\{ \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \mid a, b \in R, a = \overline{m}, (m, n) = 1 \right\}$. $I_e = \left\{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \mid b \in R \right\}$ for $e = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. Thus $|Sink(S)| = n\varphi(n)$, $|I_e| = n$, where $\varphi(n)$ is the Eulerian number of n . When $n \geq 3$, there is no source in $\Gamma(R)$. The clique number of $\Gamma(R)$ is $n - 1$.

3. SINKS AND SOURCES OF FINITE RINGS

Proposition 3.1. (1) For any ring R , if there exists a source b in $\Gamma(R)$ with $b^2 = 0$, then $R = \{0, a, b, c\}$, where a and c are left identity elements, $ba = 0 = bc$;

(2) For any ring R , if there exists a sink b in $\Gamma(R)$ with $b^2 = 0$, then $R = \{0, a, b, c\}$, where a and c are right identity elements, $ab = 0 = cb$.

Proof. (1) Let b be a source in $\Gamma(R)$ with $b^2 = 0$. Then consider the left R -module epimorphism $\eta : R \rightarrow Rb, r \mapsto rb$. Notice that $\ker(\eta) = ann_l(b) = \{0, b\}$ is a submodule of R . For any $r \in R - \{0, b\}$, we have $rb = b$. We conclude $|R| \leq 4$, since otherwise, let $a, c, d \in R - \{0, b\}$ be distinct elements. Then we have $(a - c)b = b - b = 0 = (a - d)b$. Therefore, $a - c = a - d$ and thus $c = d$, a contradiction.

Now by the proof of Lemma 2.2, R has proper left identity element. So R contains at least two left identity element, say a and c . Thus $R = \{0, a, b, c\}$, where a and c are left identity elements, $ba = 0 = bc$. In this case, we have $a \leftarrow b \rightleftharpoons b \rightarrow c$.

(2) The proof is dual to the above proof.

The structure of rings R with $|R| \leq 4$, as well as the related graph $\Gamma(R)$, is rather clear (see Lemma 2.2). As for finite rings R with $|R| \geq 5$, we have the following:

Proposition 3.2. Let R be a finite ring with proper left identity elements. If $|R| \geq 5$, then

(1) $\Gamma(R)$ contains at least two sinks;

(2) For any sink r in $\Gamma(R)$, $r^2 \neq 0$;

(3) $\Gamma(R)$ contains no sources.

Proof. Since every left identity element of R is a sink in $\Gamma(R)$, thus $\Gamma(R)$ has at least two sinks. Let e be any left identity element of R . Then I_e^* is not empty, and for any $x \in I_e^*, y \in R^*$, we have $xy = 0$. Thus every non-zero element of R is a vertex of $\Gamma(R)$ and, if $\Gamma(R)$ has source, the source vertex must lie in I_e . Of course, $\Gamma(R)$ has source vertex if and only if $|I_e| = 2$ and there is no directed edge from $R^* - I_e^*$ to I_e^* . This is equivalent to saying that for the left identity element e , there exists a non-zero element b of R , such that $\text{ann}_l(e) = \{0, b\}$ and $\text{ann}_l(b) = \{0, b\}$. So if $\Gamma(R)$ has a source, this source is b . Then we have $b^2 = 0$, contradicting with the assumption and Proposition 3.1. With the same reason, sinks $r \in \Gamma(R)$ must satisfy $r^2 \neq 0$. QED

Corollary 3.3. *Let R be a finite ring with proper left identity elements. Assume that R has at least five elements. For any left identity e of R , if $\text{ann}_l(e) = \{0, b\}$ for some non-zero element b , then in $\Gamma(R)$, the out-degree of b is $|R| - 1$ and the in-degree of b is positive.*

Similarly, we have

Proposition 3.4. *Let R be a finite ring with proper right identity elements. If R has at least five elements, then*

- (1) $\Gamma(R)$ contains at least two sources;
- (2) For any source r in $\Gamma(R)$, $r^2 \neq 0$;
- (3) $\Gamma(R)$ contains no sink.

Corollary 3.5. *Let R be a finite ring with proper right identity elements. If R has at least five elements, then for any right identity e of R with $\text{ann}_r(e) = \{0, b\}$ for some non-zero element b , the in-degree of b in $\Gamma(R)$ is $|R| - 1$ and the out-degree of b is positive.*

Recall that a *network* N is a directed graph with exactly one sink vertex k and a unique source vertex c such that c connects to every vertex of N and every vertex of N connects to k . By Lemma 2.2, Theorem 2.5, Propositions 3.1, 3.5 and 3.7, we immediately have,

Corollary 3.6. *For any finite ring R with at least five elements,*

- (1) $\Gamma(R)$ can not contain sink and source at the same time;*
- (2) ([7, Cor. 3.2]) $\Gamma(R)$ is not a network for any finite ring.*

Proof. (2) is an obvious consequence of (1). To prove (1), we list all our previous results on rings with proper one-sided identity in a single place as follows:

- (a) If $|R| \leq 4$, then $\Gamma(R)$ is one of the following

$$\circ \leftarrow \circ \rightarrow \circ, \circ \rightarrow \circ \leftarrow \circ.$$

- (b) If $|R| \geq 5$ and R has proper left identity elements, then $\Gamma(R)$ contains at least two sinks, but in it there is no source.

- (c) If $|R| \geq 5$ and R has proper right identity elements, then $\Gamma(R)$ contains at least two sources, but in it there is no sink.

Definition 3.7 Suppose that R has proper left (respectively, right) identity element. An element $r \in R$ is called **strongly right (left) invertible**, if for *any* left (right) identity element e of R , r has unique right (left) inverse s relative to this e . In this case, such s has more than one left (right) inverse relative to the same left identity e . Obviously, r is strongly right invertible if for *some* left identity element e of R , r has a unique right inverse s relative to this e .

Proposition 3.8. *For an element r in a finite ring R satisfying $r^2 \neq 0$, r is a sink vertex (source vertex) in $\Gamma(R)$, if and only if r is strongly right (respectively, left) invertible in R .*

Proof. (1) Suppose r is a sink vertex of $\Gamma(R)$. We have $ry \neq 0$ for all $y \in R^*$ since $r^2 \neq 0$. Since R is a finite ring, we have $rR = R$. Assume $re = r$. Then from $res = rs \neq 0$ for all $s \in R^*$, we obtain $es = s$ for all $s \in R$. Thus e is a left identity of R . For any left identity $f \in R$, let $ru = f$. Then u is unique relative to the f , since r is a sink vertex. For the same reason, there is $r \neq v \in R^*$ such that $vr = 0$. Hence $0 = vru = vf$. Thus f is not a right identity element of R . In this case, we have $(r+v)u = ru + v(fu) = f$, where $r+v \neq r$. Thus this u has at least two left inverses (say, r and $r+v$) relative to the left identity f . Thus r is strongly right invertible in R . Notice that R has at least two sink vertex in this case.

(2) Conversely, Suppose that r is strongly right invertible in R . Then by definition, there is left, but not right, identity element in R , and for any left identity $e \in R$, r has a unique right inverse u relative to this e . If there is a path $r \rightarrow v$, then we have $r(u + v) = e$ and this implies $v + u = u$, a contradiction. Finally, suppose $x = ae - a \neq 0$, then $x \neq r$ and $xr = 0$. Thus r is a sink vertex of $\Gamma(R)$.

The proof of the other case is similar.

QED

Corollary 3.9. *Let R be a finite ring.*

(1) *If $\Gamma(R)$ contains exactly one source (respectively, sink), then $|R| = 4$, and R^* has the form of*

$$a \leftarrow b \rightarrow b \rightarrow c;$$

$$(respectively, a \rightarrow b \rightarrow b \leftarrow c);$$

(2) *Let $|R| \geq 5$. Then an element r of R^* is a sink (source) if and only if r is strongly right (left) invertible in R . In this case, $r^2 \neq 0$ and in $\Gamma(R)$ there are at least two sinks (sources).*

4. SINK(R), SOUR(R) AND NETWORK

All rings in this section have at least five elements.

Definition 4.1 For any ring R , denote

$$Sink(R) = \{\text{sinks in } \Gamma(R)\}, Sour(R) = \{\text{sources in } \Gamma(R)\},$$

$$Inv_r(R) = \{\text{strongly right invertible elements of } R \text{ relative to some proper left identity}\},$$

$$Inv_l(R) = \{\text{strongly left invertible elements of } R \text{ relative to some proper right identity}\}.$$

Proposition 4.2. *Let R be any ring with at least five elements.*

(1) *If $Sink(R)$ (respectively, $Sour(R)$) is not empty, then it is a left (right) cancellative multiplicative semigroup;*

(2) *If $Inv_r(R)$ ($Inv_l(R)$) is not empty, then it is also a left (right) cancellative multiplicative semigroup and $Inv_r(R) \subseteq Sink(R)$ ($Inv_l(R) \subseteq Sour(R)$);*

$$(3) Sink(R) = Z_r(R) - Z_l(R), Sour(R) = Z_l(R) - Z_r(R);$$

(4) $Z(R)^*$ has a disjoint decomposition

$$Z(R)^* = Sour(R) \cup (Z_r(R) \cap Z_l(R)) \cup Sink(R).$$

Proof. (1) Assume that $Sink(R)$ is nonempty. For any $a, b \in Sink(R)$, there exists $x \in R^*$ such that $x \neq a$ and $xa = 0$. By Proposition 3.1, $ann_r(a) = 0$. So $Sink(R)$ is left cancellative and $Sink(R) = Z_r(R) - Z_l(R)$. We assert $x \neq ab$, since otherwise, we would have $abx = 0$, which implies $x = 0$. Thus $ab \in Sink(R)$, since $xab = 0$. Hence $Sink(R)$ is a semigroup under the multiplication of R .

(2) Suppose that $Inv_r(R)$ is not empty. For any proper left identity elements e, f of R and any $a, b \in Inv_r(R)$, let $ax = e, f = by$. Then $(ab)(yx) = afx = ax = e$. Thus ab is right invertible. If $(ab)c = e = (ab)d$, then $bc = bd$, and $c = d$ since $Inv_r(R) \subseteq Sink(R)$. Thus ab is strongly right invertible, i.e., $ab \in Inv_r(R)$ and hence $Inv_r(R)$ is a semigroup.

The other case is dual to the sink case.

QED

Remark For some fixed proper left identity e , let $Inv_{r_e}^{-1}(R) = \{u \in R | au = e \text{ for some } a \in Inv_r(R)\}$. Then it is easy to verify that $Inv_{r_e}^{-1}(R)$ is a multiplicative semigroup with identity e .

Proposition 4.3. *For any ring R with at least five elements, R contains a proper left identity if and only if the following two conditions hold: (1) $Sink(R) \neq \emptyset$; and (2) There exists an $x \in Sink(R)$ such that $x[Sink(R)] = Sink(R)$.*

In this case, $\Gamma(R)$ contains no source while $Sink(R)$ contains at least two elements.

Proof. If R contains a proper left identity e , then $e \in Sink(R)$ and $e[Sink(R)] = Sink(R)$.

Conversely, assume that $Sink(R) \neq \emptyset$ and $x[Sink(R)] = Sink(R)$ for some $x \in Sink(R)$. Let $xe = x, e \in Sink(R)$. Then $x(ey - y) = 0$ for all $y \in R^*$. Then by Proposition 3.1, $ey = y$ and hence, e is a left identity of R . Since e is a zero-divisor, it is a proper left identity.

If there is a source in $\Gamma(R)$, then it must lie in I_e . Then $I_e = \{0, b\}$ for some nonzero b . Then b is a source of $\Gamma(R)$ with $b^2 = 0$. Then by Proposition 3.1, $|R| = 4$ and there is no sink in $\Gamma(R)$, a contradiction. So in this case, there is no source in $\Gamma(R)$.

QED

Since $Sink(R)$ is left cancellative, we immediately have

Corollary 4.4. *For any ring R with at least five elements, if $0 < |Sink(R)| < \infty$, then $Sink(R) = Inv_r(R)$ and there is no source in $\Gamma(R)$.*

The following two results are duals of 4.3 and 4.4

Proposition 4.5. *For any ring R with at least five elements, R contains a proper right identity, if and only if the following two conditions hold: (1) $\text{Sour}(R) \neq \emptyset$; and (2) There exists an $y \in \text{Sour}(R)$ such that $[\text{Sour}(R)]y = \text{Sour}(R)$.*

In this case, $\Gamma(R)$ contains no sink but it contains at least two sources.

Corollary 4.6. *For any ring R with at least five elements, if $0 < |\text{Sour}(R)| < \infty$, then $\text{Sour}(R) = \text{Inv}_l(R)$ and there is no sink in $\Gamma(R)$.*

As a combination of Propositions 4.2, 4.3 and 4.5, we have

Proposition 4.7. *For any ring R , one-sided identity of R are two-sided identity if and only if $\Gamma(R)$ satisfies one of the following conditions:*

- (1) $\text{Sink}(R) = \emptyset$ and $\text{Sour}(R) = \emptyset$;
- (2) $\text{Sink}(R) = \emptyset$, $\text{Sour}(R) \neq \emptyset$ and for any $t \in \text{Sour}(R)$, $[\text{Sour}(R)]t \subset \text{Sour}(R)$; (In this case, $\Gamma(R)$ has infinitely many sources);
- (3) $\text{Sour}(R) = \emptyset$, $\text{Sink}(R) \neq \emptyset$ and for any $s \in \text{Sink}(R)$, $s[\text{Sink}(R)] \subset \text{Sink}(R)$; (In this case, $\Gamma(R)$ has infinitely many sinks);
- (4) $\text{Sink}(R) \neq \emptyset$ and for any $s \in \text{Sink}(R)$, $s[\text{Sink}(R)] \subset \text{Sink}(R)$. At the same time, $\text{Sour}(R) \neq \emptyset$ and for any $t \in \text{Sour}(R)$, $[\text{Sour}(R)]t \subset \text{Sour}(R)$. (In this case, $\Gamma(R)$ has infinitely many sinks and infinitely many sources.)

Corollary 4.8. *Suppose that in a ring R , one-sided identity element is two-sided identity. If in addition, R satisfies descending chain condition on principal left (respectively, right) ideals, then $\Gamma(R)$ contains no source (sink). In particular, if R is a left and right artinian ring, and one-sided identity element in R is two-sided identity, then $\Gamma(R)$ contains neither source no sink.*

Proof. It is easy to verify that for $K = \text{Sink}(R)$ and any $t \in K$, $tK \subset K$ if and only if for any (or some) $s \in K$, $stK \subset sK$, if and only if $stR \subset sR$. So, if R has DCC on right principal ideals, then in Proposition 4.7, cases (3) and (4) could not occur. Thus $\Gamma(R)$ contains no sinks. QED

Finally, as a corollary of Propositions 3.1, 4.3, 4.5 and 4.7, we have

Corollary 4.9. *For any ring R , $\Gamma(R)$ is not a network.*

REFERENCES

1. D. D. Anderson and M. Naseer, *Beck's coloring of a commutative ring*, J. Algebra **159** (1993), 500-514.
2. D. F. Anderson and P. S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434-447.
3. D. F. Anderson, Ron Levy, J. Shapiro, *Zero-divisor graphs, von Neumann regular rings, and Boolean algebras*, J. Pure Applied Algebra **180** (2003), 221-241.
4. I. Beck, *Coloring of commutative rings*, J. Algebra **116** (1988), 208-226.
5. F. R. DeMeyer, T. McKenzie, and K. Schneider, *The zero-divisor graph of a commutative semigroup*, Semigroup Forum **65** (2002), 206-214.
6. S. P. Redmond, *The zero-divisor graph of a non-commutative ring*, Internat. J. Commutative Rings **1**(4) (2002), 203-211.
7. S. P. Redmond, *Structure in the zero-divisor graph of a non-commutative ring*, Preprint (2003).
8. R. J. Wilson, *Introduction to Graph Theory*, Longman Inc., New York, Third Edition 1985.