Cobordism independence of Grassmann manifolds

ASHISH KUMAR DAS

Department of Mathematics, North-Eastern Hill University, Permanent Campus, Shillong 793 022, India E-mail: akdas@nehu.ac.in

MS received 11 April 2003; revised 9 October 2003

Abstract. This note proves that, for $F = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , the bordism classes of all nonbounding Grassmannian manifolds $G_k(F^{n+k})$, with k < n and having real dimension d, constitute a linearly independent set in the unoriented bordism group \mathfrak{N}_d regarded as a \mathbb{Z}_2 -vector space.

Keywords. Grassmannians; bordism; Stiefel-Whitney class.

1. Introduction

This paper is a continuation of the ongoing study of cobordism of Grassmann manifolds. Let *F* denote one of the division rings \mathbb{R} of reals, \mathbb{C} of complex numbers, or \mathbb{H} of quaternions. Let $t = \dim_{\mathbb{R}} F$. Then the Grassmannian manifold $G_k(F^{n+k})$ is defined to be the set of all k-dimensional (left) subspaces of F^{n+k} . $G_k(F^{n+k})$ is a closed manifold of real dimension *nkt*. Using the orthogonal complement of a subspace one identifies $G_k(F^{n+k})$ with $G_n(F^{n+k})$.

In [8], Sankaran has proved that, for $F = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , the Grassmannian manifold $G_k(F^{n+k})$ bounds if and only if v(n+k) > v(k), where, given a positive integer m, v(m)denotes the largest integer such that $2^{\nu(m)}$ divides *m*.

Given a positive integer d, let $\mathscr{G}(d)$ denote the set of bordism classes of all nonbounding Grassmannian manifolds $G_k(F^{n+k})$ having real dimension d such that k < n. The restriction k < n is imposed because $G_k(F^{n+k}) \approx G_n(F^{n+k})$ and, for k = n, $G_k(F^{n+k})$ bounds. Thus, $\mathscr{G}(d) = \{ [G_k(F^{n+k})] \in \mathfrak{N}_* \mid nkt = d, k < n, \text{ and } \nu(n+k) \le \nu(k) \} \subset \mathfrak{N}_d.$ The purpose of this paper is to prove the following:

Theorem 1.1. $\mathscr{G}(d)$ is a linearly independent set in the \mathbb{Z}_2 -vector space \mathfrak{N}_d .

Similar results for Dold and Milnor manifolds can be found in [6] and [1] respectively.

2. The real Grassmannians — a Brief review

The real Grassmannian manifold $G_k(\mathbb{R}^{n+k})$ is an *nk*-dimensional closed manifold of *k*planes in \mathbb{R}^{n+k} . It is well-known (see [3]) that the mod-2 cohomology of $G_k(\mathbb{R}^{n+k})$ is given by

$$H^*(G_k(\mathbb{R}^{n+k}); \mathbb{Z}_2) \cong \mathbb{Z}_2[w_1, w_2, \dots, w_k, \bar{w}_1, \bar{w}_2, \dots, \bar{w}_n]/\{w.\bar{w} = 1\},\$$

34 Ashish Kumar Das

where $w = 1 + w_1 + w_2 + \dots + w_k$ and $\bar{w} = 1 + \bar{w}_1 + \bar{w}_2 + \dots + \bar{w}_n$ are the total Stiefel– Whitney classes of the universal *k*-plane bundle γ_k and the corresponding complementary bundle γ_k^{\perp} , both over $G_k(\mathbb{R}^{n+k})$, respectively.

For computational convenience in this cohomology one uses the flag manifold $\operatorname{Flag}(\mathbb{R}^{n+k})$ consisting of all ordered (n+k)-tuples $(V_1, V_2, \ldots, V_{n+k})$ of mutually orthogonal one-dimensional subspaces of \mathbb{R}^{n+k} with respect to the 'standard' inner product on \mathbb{R}^{n+k} . It is standard (see [4]) that the mod-2 cohomology of $\operatorname{Flag}(\mathbb{R}^{n+k})$ is given by

$$H^*(\operatorname{Flag}(\mathbb{R}^{n+k}); \mathbb{Z}_2) \cong \mathbb{Z}_2[e_1, e_2, \dots, e_{n+k}] / \left\{ \prod_{i=1}^{n+k} (1+e_i) = 1 \right\}$$

where $e_1, e_2, \ldots, e_{n+k}$ are one-dimensional classes. In fact each e_i is the first Stiefel–Whitney class of the line bundle λ_i over $\operatorname{Flag}(\mathbb{R}^{n+k})$ whose total space consists of pairs, a flag $(V_1, V_2, \ldots, V_{n+k})$ and a vector in V_i .

There is a map π_{n+k} : Flag $(\mathbb{R}^{n+k}) \longrightarrow G_k(\mathbb{R}^{n+k})$ which assigns to $(V_1, V_2, \dots, V_{n+k})$, the *k*-dimensional subspace $V_1 \oplus V_2 \oplus \dots \oplus V_k$. In the cohomology, $\pi_{n+k}^* : H^*(G_k(\mathbb{R}^{n+k}); \mathbb{Z}_2) \longrightarrow H^*(\text{Flag}(\mathbb{R}^{n+k}); \mathbb{Z}_2)$ is injective and is described by

$$\pi_{n+k}^*(w) = \prod_{i=1}^k (1+e_i), \quad \pi_{n+k}^*(\bar{w}) = \prod_{i=k+1}^{n+k} (1+e_i)$$

In [9], Stong has observed, among others, the following facts:

Fact 2.1. The value of the class $u \in H^*(G_k(\mathbb{R}^{n+k}); \mathbb{Z}_2)$ on the fundamental class of $G_k(\mathbb{R}^{n+k})$ is the same as the value of

$$\pi_{n+k}^*(u)e_1^{k-1}e_2^{k-2}\dots e_{k-1}e_{k+1}^{n-1}e_{k+2}^{n-2}\dots e_{n+k-1}$$

on the fundamental class of $Flag(\mathbb{R}^{n+k})$.

Fact 2.2. In $H^*(\operatorname{Flag}(\mathbb{R}^{n+k}); \mathbb{Z}_2)$ one has

$$e_{i_1}^{n+k-(r-1)}e_{i_2}^{n+k-(r-1)}\dots e_{i_{r-1}}^{n+k-2}e_{i_r}^{n+k-1} = 0$$

if $1 \le r \le n+k$ and the set $\{i_1, i_2, \dots, i_r\} \subset \{1, 2, \dots, n+k\}$. In particular $e_i^{n+k} = 0$ for each $i, 1 \le i \le n+k$.

Fact 2.3. In the top dimensional cohomology of $\operatorname{Flag}(\mathbb{R}^{n+k})$, a monomial $e_1^{i_1}e_2^{i_2}\dots e_{n+k}^{i_{n+k}}$ represents the non-zero class if and only if the set $\{i_1, i_2, \dots, i_{n+k}\} = \{0, 1, \dots, n+k-1\}$.

The tangent bundle τ over $G_k(\mathbb{R}^{n+k})$ is given (see [5]) by

$$\tau \oplus \gamma_k \otimes \gamma_k \cong (n+k)\gamma_k.$$

In particular, the total Stifel–Whitney class $W(G_k(\mathbb{R}^{n+k}))$ of the tangent bundle over $G_k(\mathbb{R}^{n+k})$ maps under π_{n+k}^* to

$$\prod_{1 \le i \le k} (1+e_i)^{n+k} \cdot \prod_{1 \le i < j \le k} (1+e_i+e_j)^{-2}$$

Choosing a positive integer α such that $2^{\alpha} \ge n + k$, we have, using Fact 2.2,

$$\pi_{n+k}^*(W(G_k(\mathbb{R}^{n+k}))) = \prod_{1 \le i \le k} (1+e_i)^{n+k} \cdot \prod_{1 \le i < j \le k} (1+e_i+e_j)^{2^{\alpha}-2}$$

Thus, the *m*th Stiefel–Whitney class $W_m = W_m(G_k(\mathbb{R}^{n+k}))$ maps under π_{n+k}^* to the *m*th elementary symmetric polynomial in e_i , $1 \le i \le k$, each with multiplicity n+k, and e_i+e_j , $1 \le i < j \le k$, each with multiplicity $2^{\alpha} - 2$. Therefore, if $S_p(\sigma_1, \sigma_2, ..., \sigma_p)$ denotes the expression of the power sum $\sum_{m=1}^{q} y_m^p$ as a polynomial in elementary symmetric polynomials σ_m 's in q 'unknowns' $y_1, y_2, ..., y_q, q \ge p$, we have (see [8])

$$S_p(\pi_{n+k}^*(W_1), \pi_{n+k}^*(W_2), \dots, \pi_{n+k}^*(W_p)) = \sum_{1 \le i \le k} (n+k)e_i^p.$$

Thus we have a polynomial

$$S_p(G_k(\mathbb{R}^{n+k})) = S_p(W_1, W_2, \dots, W_p) \in H^p(G_k(\mathbb{R}^{n+k}); \mathbb{Z}_2)$$

of Stiefel–Whitney classes of $G_k(\mathbb{R}^{n+k})$ such that

$$\pi_{n+k}^*(S_p(G_k(\mathbb{R}^{n+k}))) = \begin{cases} \sum_{1 \le i \le k} e_i^p, & \text{if } n+k \text{ is odd and } p < n+k \\ 0, & \text{otherwise.} \end{cases}$$
(2.4)

3. Proof of Theorem 1.1

It is shown in [2] that

$$[G_{2k}(\mathbb{R}^{2n+2k})] = [G_k(\mathbb{R}^{n+k})]^4 \quad \text{in } \mathfrak{N}_{4nk}$$

From this, we have, in particular,

$$[G_k(F^{n+k})] = [G_k(\mathbb{R}^{n+k})]^t \quad \text{in } \mathfrak{N}_{nkt}$$

For this one has to simply observe that the mod-2 cohomology of the \mathbb{F} -Grassmannian is isomorphic as ring to that of the corresponding real Grassmannian by an obvious isomorphism that multiplies the degree by *t*. On the other hand, since \mathfrak{N}_* is a polynomial ring over the field \mathbb{Z}_2 , we have the following:

Remark 3.1. A set $\{[M_1], [M_2], \dots, [M_m]\}$ is linearly independent in \mathfrak{N}_d if and only if the set $\{[M_1]^{2^{\beta}}, [M_2]^{2^{\beta}}, \dots, [M_m]^{2^{\beta}}\}$ is linearly independent in $\mathfrak{N}_{d,2^{\beta}}, \beta \ge 0$.

Therefore, noting that t = 1, 2, or 4, it is enough to prove Theorem 1.1 for real Grassmannians only. Thus, from now onwards, we shall take

$$\mathscr{G}(d) = \{ [G_k(\mathbb{R}^{n+k})] \mid nk = d, k < n, \text{ and } \nu(n+k) \le \nu(k) \}.$$

If $G_k(\mathbb{R}^{n+k})$ is an odd-dimensional real Grassmannian manifold then both *n* and *k* must be odd, and so v(n+k) > v(k). This means that $G_k(\mathbb{R}^{n+k})$ bounds and so it follows that $\mathscr{G}(d) = \emptyset$ if *d* is odd. Therefore we assume that *d* is even.

Ashish Kumar Das

Lemma 3.2. In $H^*(\operatorname{Flag}(\mathbb{R}^{n+k}); \mathbb{Z}_2)$ one has, for $1 \leq j \leq k$,

$$\begin{pmatrix} \sum_{1 \le i \le k} e_i^{n+k-(2j-1)} \end{pmatrix} \cdot e_1^{k-1} e_2^{k-2} \dots e_{k-j}^j \cdot e_{k-(j-1)}^{j-1} \cdot e_{k-(j-2)}^{n+k-(j-1)} \dots e_k^{n+k-1} \\ = e_1^{k-1} e_2^{k-2} \dots e_{k-j}^j \cdot e_{k-(j-1)}^{n+k-j} \cdot e_{k-(j-2)}^{n+k-(j-1)} \dots e_k^{n+k-1}.$$

Proof. Note that

(a) if $i \neq k - (j - 1)$ then the exponent of e_i in the product

$$e_1^{k-1}e_2^{k-2}\dots e_{k-j}^j\cdot e_{k-(j-1)}^{j-1}\cdot e_{k-(j-2)}^{n+k-(j-1)}\dots e_k^{n+k-1}$$

is greater than or equal to *j*, and

(b) $\{n+k-(2j-1)\}+j=n+k-(j-1).$

Therefore, invoking Fact 2.2, the lemma follows.

PROPOSITION 3.3.

Let $\mathcal{O}(d) = \{ [G_k(\mathbb{R}^{n+k})] \in \mathcal{G}(d) \mid n+k \text{ is odd } \}.$ Then $\mathcal{O}(d)$ is linearly independent in \mathfrak{N}_d .

Proof. Arrange the members of $\mathcal{O}(d)$ in descending order of the values of n + k, so that

$$\mathscr{O}(d) = \{ [G_{k_1}(\mathbb{R}^{n_1+k_1})], \ [G_{k_2}(\mathbb{R}^{n_2+k_2})], \ \dots, \ [G_{k_s}(\mathbb{R}^{n_s+k_s})] \} \}$$

where $n_1 + k_1 > n_2 + k_2 > \cdots > n_s + k_s$. Note that $n_1 = d$ and $k_1 = 1$.

For a *d*-dimensional Grassmannian manifold $G_k(\mathbb{R}^{n+k})$, consider the polynomials

$$f_{\ell}(G_k(\mathbb{R}^{n+k})) = \prod_{\substack{1 \le j \le k_{\ell} + k_{\ell} - (2j-1)}} S_{n_{\ell}+k_{\ell}-(2j-1)}(G_k(\mathbb{R}^{n+k})) \in H^d(G_k(\mathbb{R}^{n+k});\mathbb{Z}_2)$$

of Stiefel–Whitney classes of $G_k(\mathbb{R}^{n+k})$, where $1 \le \ell \le s$. Then, for each ℓ , $1 \le \ell \le s$, we have, using (2.4),

$$\begin{split} \pi_{n_{\ell}+k_{\ell}}^{*}(f_{\ell}(G_{k_{\ell}}(\mathbb{R}^{n_{\ell}+k_{\ell}})))e_{1}^{k_{\ell}-1}e_{2}^{k_{\ell}-2}\dots e_{k_{\ell}-1}e_{k_{\ell}+1}^{n_{\ell}-1}e_{k_{\ell}+2}^{n_{\ell}-2}\dots e_{n_{\ell}+k_{\ell}-1} \\ &=\left(\prod_{1\leq j\leq k_{\ell}}\left(\sum_{1\leq i\leq k_{\ell}}e_{i}^{n_{\ell}+k_{\ell}-(2j-1)}\right)\right)e_{1}^{k_{\ell}-1}e_{2}^{k_{\ell}-2}\dots e_{k_{\ell}-1}e_{k_{\ell}+1}^{n_{\ell}-2}e_{k_{\ell}+2}^{n_{\ell}-2} \\ \dots e_{n_{\ell}+k_{\ell}-1} \\ &=e_{1}^{n_{\ell}}e_{2}^{n_{\ell}+1}\dots e_{k_{\ell}}^{n_{\ell}+k_{\ell}-1}e_{k_{\ell}+1}^{n_{\ell}-2}e_{k_{\ell}+2}^{n_{\ell}-2}\dots e_{n_{\ell}+k_{\ell}-1}, \end{split}$$

applying Lemma 3.2 repeatedly for successive values of *j*.

Thus, in view of Facts 2.1 and 2.3, the Stiefel–Whitney number

$$\langle f_\ell(G_{k_\ell}(\mathbb{R}^{n_\ell+k_\ell})), [G_{k_\ell}(\mathbb{R}^{n_\ell+k_\ell})] \rangle \neq 0$$

for each ℓ , $1 \le \ell \le s$. On the other hand, using (2.4), it is clear that

$$\langle f_\ell(G_{k_h}(\mathbb{R}^{n_h+k_h})), [G_{k_h}(\mathbb{R}^{n_h+k_h})] \rangle = 0$$

36

for each $h > \ell$, since $n_{\ell} + k_{\ell} - 1 \ge n_h + k_h$. Therefore, it follows that the $s \times s$ matrix

$$[\langle f_\ell(G_{k_h}(\mathbb{R}^{n_h+k_h})), [G_{k_h}(\mathbb{R}^{n_h+k_h})]
angle]_{1 \le \ell \le s, \ 1 \le h \le s}$$

is non-singular; being lower triangular with 1's in the diagonal. This completes the proof.

Now we shall complete the proof of Theorem 1.1 using induction on d. First note that

$$\begin{aligned} \mathscr{G}(2) &= \{ [G_1(\mathbb{R}^{2+1})] \} = \{ [\mathbb{R}P^2] \}, \\ \mathscr{G}(4) &= \{ [G_1(\mathbb{R}^{4+1})] \} = \{ [\mathbb{R}P^4] \}, \end{aligned}$$

and so both are linearly independent in \mathfrak{N}_2 , \mathfrak{N}_4 respectively. Assume that the theorem holds for all dimensions less than d.

We have $\mathscr{G}(d) = \mathscr{E}(d) \bigcup \mathscr{O}(d)$, where

$$\mathscr{E}(d) = \{ [G_k(\mathbb{R}^{n+k})] \in \mathscr{G}(d) | n+k \text{ is even} \}$$

and

$$\mathscr{O}(d) = \{ [G_k(\mathbb{R}^{n+k})] \in \mathscr{G}(d) | n+k \text{ is odd} \}$$

Observe that if $[G_k(\mathbb{R}^{n+k})] \in \mathscr{E}(d)$ then both *n* and *k* are even with $v(k) \neq v(n)$. On the other hand, $[G_2(\mathbb{R}^{\frac{d}{2}+2})] \in \mathscr{E}(d)$ if $d \equiv 0 \pmod{8}$. Thus, $\mathscr{E}(d) \neq \emptyset$ if and only if $d \equiv 0 \pmod{8}$.

In view of Proposition 3.3, we may assume without any loss that $\mathscr{E}(d) \neq \emptyset$. Then, by the above observation and by Theorem 2.2 of [8] every member of $\mathscr{E}(d)$ is of the form $[G_{\frac{k}{2}}(\mathbb{R}^{\frac{n}{2}+\frac{k}{2}})]^4$, where $[G_{\frac{k}{2}}(\mathbb{R}^{\frac{n}{2}+\frac{k}{2}})] \in \mathscr{G}(\frac{d}{4})$. By induction hypothesis, $\mathscr{G}(\frac{d}{4})$ is linearly independent in $\mathfrak{N}_{\underline{4}}$.

So, by Remark 3.1,

$$\mathscr{E}(d)$$
 is linearly independent in \mathfrak{N}_d . (3.4)

Again note that if $[G_k(\mathbb{R}^{n+k})] \in \mathscr{E}(d)$, then, by (2.4), the polynomial $S_p(G_k(\mathbb{R}^{n+k})) = 0$, $\forall p \ge 1$. So, for each of the polynomials f_ℓ , $1 \le \ell \le s$, considered in Proposition 3.3, we have

$$\langle f_{\ell}(G_k(\mathbb{R}^{n+k})), [G_k(\mathbb{R}^{n+k})] \rangle = 0.$$

Therefore, writing

$$\mathscr{E}(d) = \{ [G_{k_{s+1}}(\mathbb{R}^{n_{s+1}+k_{s+1}})], [G_{k_{s+2}}(\mathbb{R}^{n_{s+2}+k_{s+2}})], \dots, [G_{k_{s+q}}(\mathbb{R}^{n_{s+q}+k_{s+q}})] \},$$

.

where $n_{s+1} + k_{s+1} > n_{s+2} + k_{s+2} > \cdots > n_{s+q} + k_{s+q}$, we see that the $s \times (s+q)$ matrix

$$[\langle f_{\ell}(G_{k_h}(\mathbb{R}^{n_h+k_h})), [G_{k_h}(\mathbb{R}^{n_h+k_h})]\rangle]_{1 \le \ell \le s, \ 1 \le h \le s+q}$$

is of the form

$$\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ \star & 1 & 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ \star & \star & 1 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ - & - & - & \cdots & - & - & - & - & \cdots & - \\ - & - & - & \cdots & - & - & - & - & \cdots & - \\ \star & \star & \star & \cdots & 1 & 0 & 0 & 0 & \cdots & 0 \\ \ell(d) & & & & \mathcal{E}(d) \end{bmatrix}.$$
(3.5)

38 Ashish Kumar Das

Thus, no non-trivial linear combination of members of $\mathcal{O}(d)$ can be expressed as a linear combination of the members of $\mathscr{E}(d)$. This, together with (3.4) and Proposition 3.3, proves that the set $\mathscr{G}(d) = \mathscr{E}(d) \bigcup \mathscr{O}(d)$ is linearly independent in \mathfrak{N}_d . Hence, by induction, Theorem 1.1 is completely proved.

Remark 3.6. Using the decomposition of the members of $\mathscr{E}(d)$, and the polynomials f_{ℓ} , in the lower dimensions together with the *doubling homomorphism* defined by Milnor [7], one can obtain a set of polynomials of Stiefel–Whitney classes which yield, as in Proposition 3.3, a lower triangular matrix for $\mathscr{E}(d)$ with 1's in the diagonal. Thus using (3.5) we have a lower triangular matrix, with 1's in the diagonal, for the whole set $\mathscr{G}(d)$.

Acknowledgement

Part of this work was done under a DST project

References

- Dutta S and Khare S S, Independence of bordism classes of Milnor manifolds, J. Indian Math. Soc. 68(1–4) (2001) 1–16
- [2] Floyd E E, Steifel–Whitney numbers of quaternionic and related manifolds, *Trans. Am. Math. Soc.* 155 (1971) 77–94
- [3] Hiller H L, On the cohomology of real Grassmannians, *Trans. Am. Math. Soc.* 257 (1980) 521–533
- [4] Hirzebruch F, Topological Methods in Algebraic Geometry (New York: Springer-Verlag) (1966)
- [5] Hsiang W-C and Szczarba R H, On the tangent bundle for the Grassmann manifold, *Am. J. Math.* 86 (1964) 698–704
- [6] Khare S S, On Dold manifolds, Topology Appl. 33 (1989) 297-307
- [7] Milnor J W, On the Stiefel–Whitney numbers of complex manifolds and of spin manifolds, *Topology* 3 (1965) 223–230
- [8] Sankaran P, Determination of Grassmann manifolds which are boundaries, *Canad. Math. Bull.* **34** (1991) 119–122
- [9] Stong R E, Cup products in Grassmannians, Topology Appl. 13 (1982) 103-113