# arXiv:math/0403505v2 [math.CO] 9 Aug 2004

# A GRAPHIC GENERALIZATION OF ARITHMETIC

BILAL KHAN \*, KIRAN R. BHUTANI <sup>†</sup>, AND DELARAM KAHROBAEI <sup>‡</sup>

Abstract. In this paper, we extend the classical arithmetic defined over the set of natural numbers  $\mathbb{N}$ , to the set of all finite directed connected multigraphs having a pair of distinguished vertices. Specifically, we introduce a model  $\mathcal{F}$  on the set of such graphs, and provide an interpretation of the language of arithmetic  $\mathcal{L} = \{0, 1, \leq, +, \times\}$  inside  $\mathcal{F}$ . The resulting model exhibits the property that the standard model on  $\mathbb{N}$  embeds in  $\mathcal{F}$  as a submodel, with the directed path of length n playing the role of the standard integer n. We will compare the theory of the larger structure  $\mathcal{F}$  with classical arithmetic statements that hold in  $\mathbb{N}$ . For example, we explore the extent to which  $\mathcal{F}$  enjoys properties like the associativity and commutativity of + and  $\times$ , distributivity, cancellation and order laws, and decomposition into irreducibles.

Key words. arithmetic, graphs.

AMS subject classifications. 05C99, 11U10

**1. Introduction.** The *language of arithmetic*  $\mathcal{L}$  consists of two 0-ary relations **0** and **1**, one binary relation  $\leq$ , and two ternary relations + and  $\times$ . In this paper, we generalize classical arithmetic defined over the natural numbers  $\mathbb{N} = \{0, 1, 2, \ldots\}$ , to the set F consisting of all *flow graphs*: finite directed connected multigraphs in which a pair of distinguished vertices designated as the *source* and *target* vertex. We give natural interpretation for  $\mathcal{L}$  on the set F. To avoid confusion with the standard model of arithmetic, the corresponding operations in F are denoted with a circumscribed circle. The new model  $\mathcal{F} = \langle F, \odot, \odot, \odot, \odot, \odot, \otimes \rangle$  is a natural extension of the standard model  $\mathcal{N} = \langle \mathbb{N}, 0, 1, \leq +, \times \rangle$ . Specifically, we exhibit an embedding  $i: \mathcal{N} \stackrel{i}{\to} \mathcal{F}$  satisfying:

$$\begin{split} i(0) &= & \textcircled{0}, \\ i(1) &= & \textcircled{1}, \\ \forall x, y \in \mathbb{N}, \ x \leqslant y \Rightarrow i(x) \bigotimes i(y), \\ \forall x, y \in \mathbb{N}, \ i(x+y) &= i(x) \bigoplus i(y), \\ \forall x, y \in \mathbb{N}, \ i(x \times y) &= i(x) \bigotimes i(y). \end{split}$$

**Objective:** Compare the theory  $Th(\mathcal{F}) = \{\phi \mid \mathcal{F} \models \phi\}$  with true arithmetic  $TA = \{\phi \mid \mathcal{N} \models \phi\}^1$ .

There have been other attempts to define algebraic and metric structures on the set of all graphs. The classical operations on graphs [5] (including extensive literature on graph products [4]) have yielded deep results and a profound mathematical theory. However, to date, these operations have not provided an interpretation of the

<sup>\*</sup>Department of Mathematics and Computer Science, City University of New York John Jay College of Criminal Justice, 899 Tenth Avenue, New York, NY 10019 (grouptheory@hotmail.com).

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, The Catholic University of America, Washington DC 20064 (bhutani@cua.edu).

<sup>&</sup>lt;sup>‡</sup>Department of Mathematics, City University of New York, Graduate Center, New York, NY 10016 (dkahrobaei@gc.cuny.edu).

<sup>&</sup>lt;sup>1</sup>Following standard model theory, here  $\phi$  is a first-order sentence in the language  $\mathcal{L}$ .

language of arithmetic on graphs. This paper presents results and open questions in this direction. In [1, 2, 3], the authors used graph embeddings to define a metric on the set of all simple connected graphs of a given order. This work differs from those investigations in that it considers an infinite collection of graphs in order to extend the standard model of arithmetic, and in doing so does not seek to establish a metric structure.

DEFINITION 1.1 (Flow graph). A flow graph A is a triple  $(G_A, s_A, t_A)$ , where  $G_A$  is a finite directed connected multigraph and  $s_A, t_A \in V[G_A]$  are called the source and the target vertex of A, respectively. The set of all flow graphs is denoted F. The unique flow graph for which  $|V[G_A]| = 1$  and  $|E[G_A]| = 0$  is called the trivial flow graph; all other flow graphs are considered non-trivial. If  $s_A = t_A$  and A is non-trivial then A is called an infinitesimal flow graph. The set of all infinitesimal flow graphs is denoted I. Given two flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ , a map  $\phi : A \to B$  is called an injective morphism of flow graphs if (as a graph embedding)  $\phi$  maps  $G_A$  injectively into  $G_B$  and additionally satisfies  $\phi(s_A) = s_B$ ,  $\phi(t_A) = t_B$ . Flow graphs A and B are considered isomorphic if there is an injective morphism  $\phi : A \to B$  for which  $Im(\phi) = B$ .

DEFINITION 1.2 (Graphical natural number). We represent the natural number n as a directed chain of length n, having n + 1 vertices. More formally, let  $P_n$  be a directed chain of length n (having n + 1 vertices) where each vertex has in-degree  $\leq 1$ and out-degree  $\leq 1$ . Denote by  $s_n$ , the unique vertex in  $P_n$  having in-degree 0, and let  $t_n$  be the unique vertex in  $P_n$  having out-degree 0. The flow graph  $F_n = (P_n, s_n, t_n)$ is referred the graphic natural number n. Define the map  $i : \mathcal{N} \to \mathcal{F}$  as

$$i:n\mapsto F_n.$$

**1.1. Addition.** In Definition 1.1, we represented the natural number n by the flow graph  $F_n$ . It follows that we interpret the addition of two numbers  $n_1$  and  $n_2$  inside  $\mathcal{F}$  as "concatenating"  $F_{n_1}$  with  $F_{n_2}$ . Consider, for example, the addition of 3 and 2 depicted in Figure 1.1.



FIG. 1.1. Interpreting addition of natural numbers inside  $\mathcal{F}$ .

To extend this definition of  $\oplus$  to all of F, we define general addition of flow graphs as follows: Given two flow graphs A and B, define  $A \oplus B$  to be the flow graph

obtained by identifying  $t_A$  with  $s_B$  and defining  $s_{A\oplus B} = s_A$  and  $t_{A\oplus B} = t_B$ . An example of such an addition is shown in Figure 1.2.



FIG. 1.2. General addition of flow graphs.

To make this formal we define the following operation on connected directed multigraphs: Given directed graphs  $G_1$  and  $G_2$ , and vertices  $u_1 \in V[G_1]$ ,  $u_2 \in V[G_2]$ , we define

$$G_1 \bigoplus_{u_1 \approx u_2} G_2 \stackrel{\text{\tiny def}}{=} (G_1 \sqcup G_2) / (u_1 \approx u_2)$$

to be the graph obtained by taking disjoint copies of  $G_1$  and  $G_2$  and identifying vertex  $u_1$  in  $G_1$  with vertex  $u_2$  in  $G_2$ . Note the obvious and natural injective graph homomorphisms

$$\begin{array}{rcl}
\sigma_{u_1 \approx u_2}^{\oplus} : G_1 & \hookrightarrow & G_1 \bigoplus_{u_1 \approx u_2} G_2 \\
\tau_{u_1 \approx u_2}^{\oplus} : G_2 & \hookrightarrow & G_1 \bigoplus_{u_1 \approx u_2} G_2.
\end{array}$$
(1.1)

DEFINITION 1.3. Given two flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ , we define

$$A \bigoplus B \stackrel{\text{\tiny def}}{=} (G_A \bigoplus {}_{t_A \approx s_B} G_B, s_A, t_B).$$

REMARK 1.4. Note that if A is a flow graph with  $p_A$  vertices and  $q_A$  edges, and B is a flow graph with  $p_B$  vertices and  $q_B$  edges, then  $A \oplus B$  is a flow graph having  $p_A + p_B - 1$  vertices and  $q_A + q_B$  edges.

The next lemma follows immediately from Definitions 1.2 and 1.3.

LEMMA 1.5. Let m, n be natural numbers. Then  $i(n+m) = i(n) \oplus i(m)$ .

LEMMA 1.6.  $\textcircled{0} \stackrel{\text{def}}{=} F_0$  is the unique two-sided identity with respect to  $\bigoplus$ . That is, for all flow graphs  $A, G \in \mathcal{F}$ ,

$$A \oplus G = A \iff G = \bigcirc \iff G \oplus A = A.$$

**Proof.** If  $G = F_0$  then  $A \oplus G = G \oplus A = A$ . For the reverse, we appeal to Remark 1.4, noting that  $A \oplus G = A$  implies  $p_A + p_G - 1 = p_A$  and  $q_A + q_G = q_A$ . Hence  $p_G = 1$  and  $q_G = 0$ , so  $G = F_0$ . An analogous argument shows that  $G \oplus A = A$ implies  $G = F_0$ .  $\square$ 

OBSERVATION 1.7. We note that the sum of two infinitesimals is again an infinitesimal. On the other hand, if at least one summand is a non-trivial non-infinitesimal flow graph then the summation evaluates to a non-trivial non-infinitesimal flow graph.

DEFINITION 1.8 (Scalar multiplication of flow graphs). Given a flow graph A, and a positive natural number k in  $\mathbb{N}$ , we define left-multiplication inductively as follows:

$$1A = A$$
$$kA = (k-1)A \bigoplus A$$

Right-multiplication is defined analogously. However, as we will see,  $\oplus$  is associative, and so the two notions coincide. We shall subsequently consider only left-multiplication by integer scalars.

**1.2.** Multiplication. In the previous section, we presented an interpretation of addition in  $\mathcal{F}$  that is a natural extension of addition on the natural numbers. In this section, we give an interpretation of multiplication in  $\mathcal{F}$ . In doing this, we must respect the fact that for each pair of natural numbers  $n_1, n_2$ , the following identity holds in  $\mathcal{N}$ :

$$I_{n_1,n_2}: \underbrace{n_2 + n_2 + \dots + n_2}_{n_1 \text{ times}} = n_1 n_2 = \underbrace{n_1 + n_1 + \dots + n_1}_{n_2 \text{ times}}.$$

So, in particular, the definition of  $\otimes$  in  $\mathcal{F}$  must satisfy

$$n_1 F_{n_2} = F_{n_1} \otimes F_{n_2} = n_2 F_{n_1}. \tag{1.2}$$

Given that we represent the natural number n by the flow graph  $F_n$ , the product of two graphical numbers  $F_{n_1}$  and  $F_{n_2}$  can be made to satisfy relation (1.2) if we take multiplication to be the act of replacing each edge of  $F_{n_1}$  with a copy of  $F_{n_2}$ . Consider the multiplication of graphical natural numbers  $F_3$  and  $F_2$ , as depicted in Figure 1.3.

To extend this definition of  $\otimes$  to all of F, we define general multiplication of flow graphs as follows: Given two flow graphs A and B, define  $A \otimes B$  to be the flow graph obtained by replacing every edge e (from  $E[G_A]$ ) with a copy of B as follows: For each edge e = (u, v) in A, we remove e and replace it with a graph  $B_e$  isomorphic to B, by identifying u with  $s_{B_e}$ , and v with  $t_{B_e}$ . An example of such a multiplication is shown in Figure 1.4.

To make this formal we define the following operation on connected directed multigraphs: Given directed graphs  $G_1$  and  $G_2$ , an edge  $e = (u_1, v_1) \in E[G_1]$  and vertices  $u_2, v_2 \in V[G_2]$ , we define

$$G_1 \otimes_{e \approx (u_2, v_2)} G_2 \stackrel{\text{def}}{=} [(G_1 \setminus e) \sqcup G_2] / (u_1 \approx u_2, v_1 \approx v_2)$$

to be the graph obtained by removing e from  $G_1$  and attaching a copy of  $G_2$  to the resulting graph by gluing  $u_1$  with  $u_2$  and  $v_1$  with  $v_2$ . Note the obvious and natural injective maps

$$\begin{aligned} & \sigma_{e\approx(u_2,v_2)}^{\otimes} : G_1 \setminus e \quad \hookrightarrow \quad G_1 \otimes_{e\approx(u_2,v_2)} G_2 \\ & \tau_{e\approx(u_2,v_2)}^{\otimes} : G_2 \quad \hookrightarrow \quad G_1 \otimes_{e\approx(u_2,v_2)} G_2. \end{aligned} \tag{1.3}$$



FIG. 1.3. Standard multiplication of natural numbers in  $\mathcal{F}$  (represented as flow graphs).

DEFINITION 1.9. Given flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ . Define the directed graph  $K_0 = G_A$ ,  $E_0 = E[G_A]$ , and let  $\bar{\sigma}_0, \sigma_0 : K_0 \to K_0$  be the identity isomorphisms. Fix any enumeration  $\eta$  of the edges  $E[G_A]$ , say  $\eta = e_1, e_2, \ldots, e_m$ . Inductively, for  $i = 1, 2, \ldots, m$  we define

$$K_{i} = K_{i-1} \otimes_{\bar{\sigma}_{i-1}(e_{i})\approx(s_{B},t_{B})} G_{B}$$
  

$$\sigma_{i} = \sigma_{\bar{\sigma}_{i-1}(e_{i})\approx(s_{B},t_{B})} : K_{i-1} \setminus e_{i} \to K_{i}$$
  

$$E_{i} = E_{i-1} \setminus \{e_{i}\}$$
  

$$\bar{\sigma}_{i} = (\sigma_{i})_{|E_{i}}(\sigma_{i-1})_{|E_{i}} \cdots (\sigma_{1})_{|E_{i}}(\sigma_{0})_{|E_{i}}.$$

Informally,  $K_i$  is the directed graph obtained after edges  $e_1, \ldots, e_i$  have been deleted from  $G_A$  and replaced by copies of  $G_B$ . Finally, we put

$$A \otimes {}_{n}B \stackrel{{}_{def}}{=} (K_m, \bar{\sigma}_m(s_A), \bar{\sigma}_m(t_A)).$$

The reader may verify that the operation  $\otimes_{\eta}$  is well-defined, and that in particular, it is independent of the chosen enumeration  $\eta$  of the edges  $E[G_A]$ .

REMARK 1.10. Let A be a flow graph with  $p_A$  vertices and  $q_A$  edges, and B be a flow graph having  $p_B$  vertices and  $q_B$  edges. Then  $A \otimes B$  has  $q_A q_B$  edges. If B is either trivial or infinitesimal then  $A \otimes B$  has  $1 + q_A(p_B - 1)$  vertices. If B is non-trivial and not infinitesimal then  $A \otimes B$  has  $p_A + q_A(p_B - 2)$  vertices.

The next lemma follows immediately from Definitions 1.2 and 1.9.

LEMMA 1.11. Let m, n be natural numbers. Then  $i(n \times m) = i(n) \otimes i(m)$ .

LEMMA 1.12. Given flow graphs G and H, and a non-trivial, non-infinitesimal flow graph A:

$$A \otimes G = A \Leftrightarrow G = (1) \Leftrightarrow G \otimes A = A, G \otimes H = (0) \Leftrightarrow H = (0) \quad or \quad G = (0).$$



FIG. 1.4. General multiplication of flow graphs.

*Proof.* If  $G = F_1$  then  $A \otimes G = G \otimes A = A$ . For the reverse, we appeal to Remark 1.10, noting that  $A \otimes G = A$  implies  $p_A + q_A(p_G - 2) = p_A$  and  $q_Aq_G = q_A$ . Hence  $p_G = 2$  and  $q_G = 1$ , so  $G = F_1$ . An analogous argument shows that  $G \otimes A = A$ implies  $G = F_1$ .

If  $G = F_0$  then  $G \otimes H = H \otimes G = F_0$ . For the reverse, we appeal to Remark 1.10, noting that  $G \otimes H = \textcircled{0}$  implies  $q_G q_H = 0$ , so either  $q_G = 0$  or  $q_H = 0$ . It follows that either H = 0 or G = 0.  $\square$ 

Note that if we remove the hypothesis that A is non-infinitesimal in Lemma 1.12, then  $A \otimes G = A$  and  $G \otimes A = A$  do not necessarily imply G = (1). The simplest counterexample is seen by taking G = A to be the flow graph consisting of one vertex and one loop edge. We denote this graph as  $C_1$ . Indeed, it is easy to see that this is the only counterexample. Suppose  $A \otimes G = A$ , for some infinitesimal A. By Remark 1.10, this implies  $1 + q_A(p_G - 1) = p_A$  and  $q_Aq_G = q_A$ . Hence  $q_G = 1$  and  $p_G = 1 + (p_A - 1)/q_A$ . If  $q_G = 1$  then either  $p_G = 2$  or 1. If  $p_G = 2$ , then  $G = F_1$ , and this is not a counterexample. If  $p_G = 1$ , then G is a graph consisting of one vertex and one loop edge. Moreover, since  $q_A$  is finite,  $1 + q_A(p_G - 1) = p_A$  implies that  $p_A = 1$ . It follows that  $A = q_A C_1$ .

OBSERVATION 1.13. Suppose G and H are non-trivial flow graphs of which at least one, say H, is infinitesimal. Then  $s_{G \otimes H} = t_{G \otimes H}$  in  $G \otimes H$  and  $s_{H \otimes G} = t_{H \otimes G}$  in  $H \otimes G$ . Hence  $G \otimes H$  and  $H \otimes G$  are both infinitesimal.

On the other hand, suppose G and H are non-trivial flow graphs that are both non-infinitesimal. Then  $s_{G \otimes H} \neq t_{G \otimes H}$  in  $G \otimes H$  and  $s_{H \otimes G} \neq t_{H \otimes G}$  in  $H \otimes G$ . Hence  $G \otimes H$  and  $H \otimes G$  are both non-infinitesimal.

It follows that if G and H are non-trivial flow graphs, then  $G \otimes H$  is infinitesimal if and only if at least one of the two factors is infinitesimal. The reader may wish to compare this with the second assertion of Lemma 1.12.

DEFINITION 1.14 (Scalar exponentiation of flow graphs). Given a flow graph A, and a positive natural number k in  $\mathbb{N}$ , we define right-exponentiation inductively as follows:

$$A^{1} = A$$
$$A^{k} = A^{k-1} \otimes A.$$

Left-exponentiation is defined analogously. However, as we will see shortly,  $\otimes$  is associative, and so the two notions coincide. We shall subsequently consider only right-exponentiation by integer scalars.

**1.3.** Order. Given our representation of the natural number n by the flow graph  $F_n$  in Definition 1.2, comparing the order of two numbers  $n_1$  and  $n_2$  amounts to simply comparing the lengths of the corresponding chain graphs  $F_{n_1}$  and  $F_{n_2}$ . To generalize this to all of  $\mathcal{F}$ , however, we cannot refer to "length". In what follows, we present two possible interpretations of  $\leq$  in  $\mathcal{F}$ . To avoid confusion, we refer to these interpretations as  $\otimes$  and  $\otimes$ .

**1.3.1. Weak Order**  $\bigotimes$  . Suppose we are given two flow graphs A and B. Informally, we say that  $A \bigotimes B$  iff there is a way to partition A into edge-disjoint neighborhoods of the source/target of vertices of A in such a way that these neighborhoods can be mapped into disjoint neighborhoods of the source/target vertices of B. To make this more precise we define the following operation on connected directed multigraphs.

DEFINITION 1.15 ((s,t)-splitting). Given a connected directed multigraph G = (V, E) and two vertices s and t in V, an (s,t)-splitting of G is a pair of graphs  $(H_1, H_2)$  with the following properties:

- $H_1$  and  $H_2$  are connected subgraphs of G.
- s is in  $V[H_1]$  and t is in  $V[H_2]$ .
- $\{E[H_1], E[H_2]\}$  is a partition of E. While this implies  $V[H_1] \cup V[H_2] = V[G]$ , we remark that  $V[H_1] \cap V[H_2]$  need not be empty.

We can now give a precise definition of the weak ordering.

DEFINITION 1.16 (Weak order). Given two flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ , we say that  $A \otimes B$  if there is an  $(s_A, t_A)$ -splitting  $(H_1, H_2)$  of  $G_A$  and graph embeddings  $\phi_1 : H_1 \to G_B$ ,  $\phi_2 : H_2 \to G_B$  such that  $\phi_1(s_A) = s_B$  and  $\phi_2(t_A) = t_B$  and  $\phi_1(E[H_1]) \cap \phi_2(E[H_2]) = \emptyset$ .

Consider the comparison of  $F_3$  and  $F_5$  in Figure 1.5 which illustrates the assertion that  $F_3 \otimes F_5$ .

The proof of the following lemma is immediate.

LEMMA 1.17. Let m, n be natural numbers. Then  $n \leq m \Leftrightarrow i(n) \otimes i(m)$ .

Figure 1.6 illustrates a more general example in which weak order is used to compare two elements of  $\mathcal{F}$  which are *not* graphical natural numbers.

The next Proposition follows immediately from Lemmas 1.5, 1.6, 1.11, 1.12, and 1.17.

PROPOSITION 1.18. Under the embedding  $i: n \mapsto F_n$ , the standard model  $\mathcal{N} = \langle \mathbb{N}, 0, 1, \leq , +, \times \rangle$  is a submodel of  $\mathcal{F} = \langle F, @, @, @, \otimes \rangle$ ,  $\psi \in , \otimes \rangle$ , where  $@ = F_0$ ,



FIG. 1.5. Standard weak ordering of natural numbers (represented as flow graphs).



FIG. 1.6. General weak ordering of flow graphs.

 $(1) = F_1$ , and the relations  $\oplus$ ,  $\otimes$  and  $\otimes$  reinterpret  $+, \times$  and  $\leq$  inside  $\mathcal{F}$ .

**1.3.2.** Strong Order S. We now give an alternate, strengthened ordering on  $\mathcal{F}$ . Given two flow graphs A and B, informally, we say that A S B iff a copy of  $G_A$  appears as a neighborhood of both  $s_B$  and  $t_B$  in  $G_B$ . The next definition makes this statement precise.

DEFINITION 1.19 (Strong order). Given two flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ , we say  $A \otimes B$  iff there are graph embeddings  $\phi_s : G_A \to G_B$  and  $\phi_t : G_A \to G_B$  which satisfy  $\phi_s(s_A) = s_B$  and  $\phi_t(t_A) = t_B$ .

Consider the comparison of  $F_3$  and  $F_5$  depicted in Figure 1.7; clearly  $F_3 \otimes F_5$ . The proof of the following lemma is immediate.

LEMMA 1.20. Let m, n be natural numbers. Then  $n \leq m \Leftrightarrow i(n) \otimes i(m)$ .

Figure 1.8 illustrates a more general example in which strong order is used to compare two elements of  $\mathcal{F}$  which are *not* graphical natural numbers.

The next Proposition follows immediately from Lemmas 1.5, 1.6, 1.11, 1.12, and 1.20.

PROPOSITION 1.21. Under the embedding  $i : n \mapsto F_n$ , the standard model  $\mathcal{N} = \langle \mathbb{N}, 0, 1, \leq, +, \times \rangle$  is a submodel of  $\mathcal{F} = \langle F, @, @, @, \otimes, \oplus, \otimes \rangle$ , where  $@ = F_0$ ,  $@ = F_1$ , and the relations  $\oplus$ ,  $\otimes$  and  $\otimes$  reinterpret  $+, \times$  and  $\leq$  inside  $\mathcal{F}$ .



FIG. 1.7. Standard strong ordering of natural numbers (represented as flow graphs).



FIG. 1.8. General strong ordering of flow graphs.

The next proposition and example show that ordering by  $\bigotimes$  is indeed strictly stronger than ordering by  $\bigotimes$ .

**PROPOSITION 1.22.** Given flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ 

$$A \otimes B \Rightarrow A \otimes B.$$

Proof. Since  $A \otimes B$ , there are graph embeddings  $\phi_s : G_A \to G_B$  and  $\phi_t : G_A \to G_B$  which satisfy  $\phi_s(s_A) = s_B$  and  $\phi_t(t_A) = t_B$ . Let  $E_1 = E[G_A]$ ,  $V_1 = V[G_A]$ ; take  $E_2 = \emptyset$ ,  $V_2 = \{t_A\}$ . Put  $H_1 = (V_1, E_1)$  and  $H_2 = (V_2, E_2)$ . Then  $(H_1, H_2)$  is an  $(s_A, t_A)$ -splitting of  $G_A$ . We take graph embeddings  $\phi_1 = \phi_s|_{H_1} : H_1 \to G_B$ , and  $\phi_2 = \phi_t|_{H_2} : H_2 \to G_B$ . Then  $\phi_1(s_A) = s_B$  and  $\phi_2(t_A) = t_B$  and  $\phi_1(E[H_1]) \cap \phi_2(E[H_2]) = \emptyset$ . Thus,  $A \otimes B$ .  $\square$ 

The converse of Proposition 1.22 is false, as the following example indicates.

EXAMPLE 1.23. Take A and B to be the flow graphs depicted on page 8, where Figure 1.6 illustrates that  $A \otimes B$ . Note that  $G_A$  contains a vertex of degree 3, while  $G_B$  does not, hence no neighborhood of  $s_B$  or  $t_B$  can be isomorphic to  $G_A$ . Thus  $A \otimes B$ .

**2. Results.** We begin by considering properties of  $\oplus$  in Section 2.1. We show that  $\oplus$  is an associative, non-commutative operation, and provide a natural criterion for a flow graph to be irreducible as a proper sum. We prove that every flow

graph is canonically decomposable as a sum of irreducibles. Using this canonical decomposition, we deduce left and right cancellation laws for  $\oplus$ , and show that if two flow graphs A and B commute with respect to  $\oplus$  then they are necessarily scalar multiples of some flow graph C. Then, in Section 2.2 we show that  $\otimes$  is an associative, non-commutative operation and that it right-distributes over  $\oplus$  (but does not left-distribute). We define left and right divisibility of flow graphs, and use this to introduce the notion of a prime flow graph, and show that the concept of left-prime and right-prime coincide. We describe the canonical  $\oplus$  decomposition of flow graph products in terms of the  $\oplus$  decompositions of each of the  $\otimes$  factors. Finally, in Section 2.3, we explore the relationship between strong order (denoted by  $(\mathbf{s})$ ) and weak order (denoted by  $\otimes$ ), describing the interaction between these orders and the operations of  $\oplus$  and  $\otimes$ . We show that while the two orders coincide on the graphical natural numbers, neither order is anti-symmetric on all of  $\mathcal{F}$ , and only  $\bigotimes$  is transitive. On the other hand, many of the laws that govern the relationship between  $\leq$ , + and × in  $\mathcal{N}$  continue to hold for  $\leq$ ,  $\oplus$  and  $\otimes$  in  $\mathcal{F}$ , but these laws are violated under the ordering  $\bigotimes$ .

**2.1.** Additive Properties. In this section we present some properties of  $\oplus$ . LEMMA 2.1 (Associativity of  $\oplus$ ). The operation  $\oplus$  is associative. Proof. Given flow graphs A, B, C,

$$(A \oplus B) \oplus C = (G_A \oplus {}_{t_A \approx s_B}G_B, s_A, t_B) \oplus C$$
$$= ((G_A \oplus {}_{t_A \approx s_B}G_B) \oplus {}_{t_B \approx s_C}G_C, s_A, t_C)$$
$$= (G_A \oplus {}_{t_A \approx s_B}(G_B \oplus {}_{t_B \approx s_C}G_C), s_A, t_C)$$
$$= A \oplus (G_B \oplus {}_{t_B \approx s_C}G_C, s_B, t_C)$$
$$= A \oplus (B \oplus C).$$

L		

EXAMPLE 2.2. Let A be the flow graph consisting of a directed cycle of length 3 and let source and target vertices be any two vertices on this cycle. Then it is easy to check that  $A \oplus F_2$  is not equal to  $F_2 \oplus A$ , that is to say, there is no flow graph isomorphism between  $A \oplus F_2$  and  $F_2 \oplus A$  (see Figure 2.1).



FIG. 2.1. Example showing the non-commutativity of addition in  $\mathcal{F}$ .

The previous example proves the next lemma.

LEMMA 2.3. The operation  $\oplus$  is not commutative.

DEFINITION 2.4 ( $\oplus$ -Irreducible). A flow graph A is called  $\oplus$ -reducible if there is decomposition of A as sum of non-trivial flow graphs B and C. Otherwise A is called  $\oplus$ -irreducible.

DEFINITION 2.5 (s-standard and t-standard). Let A be a flow graph. A is called s-standard if it can not be decomposed as  $A = B \bigoplus C$ , where B is infinitesimal, and C is an arbitrary (possibly trivial) flow graph. A is called t-standard if it cannot be decomposed as  $A = B \bigoplus C$ , where B is an arbitrary (possibly trivial) flow graph, and and C is an infinitesimal flow graph.

Note that  $\bigoplus$ -irreducible non-infinitesimal flow graphs are both *s*-standard and *t*-standard. On the other hand,  $\bigoplus$ -irreducible infinitesimal flow graphs are neither *s*-standard nor *t*-standard. There are no restrictions on the *s*-standardness and *t*-standardness properties of general  $\bigoplus$ -reducible flow graphs.

We would like to devise a graph-theoretic characterization of  $\oplus$  -irreducibility. Towards this, the next definition is the flow graph analogue of a cut vertex in standard graphs.

DEFINITION 2.6 (Splitting vertex for a flow graph). We say that w is a splitting vertex for flow graph  $A = (G_A, s_A, t_A)$  if  $w \neq s_A, t_A$  and the deletion of w from  $G_A$  produces at least two non-trivial components, with  $s_A$  and  $t_A$  lying in distinct components. We denote the component containing  $s_A$  as  $G_A^s(w)$ , the one containing  $t_A$  as  $G_A^t(w)$ , and the remaining components as  $G_A^e(w)$ . Note that  $G_A^e(w)$  may be the union of several disjoint components, and hence is not necessarily connected. Let  $i_s : G_A^s(w) \hookrightarrow G_A$ ,  $i_t : G_A^t(w) \hookrightarrow G_A$ , and  $i_{\epsilon} : G_A^e(w) \hookrightarrow G_A$  denote the natural subgraph injections. Since w is a splitting vertex for A,  $s_A \in Im(i_s)$ ,  $t_A \in Im(i_t)$ . The pair of graphs  $(G_A^s(w), G_A^t(w))$  is called the  $(s_A, t_A)$  splitting of  $G_A$  induced by w.

REMARK 2.7. If w is a splitting vertex in flow graph  $A = (G_A, s_A, t_A)$ , then by Definition 2.6, w is a cut vertex in  $G_A$ . The converse is false, however, since not every cut vertex in  $G_A$  is a splitting vertex in A.

For a concrete example, the reader may wish to consider the flow graph A in Figure 1.2 on page 3, where the radius 1 sphere of  $t_A$  contains two cut vertices for  $G_A$ , only one of which is a splitting vertex for A.

As a more general example, consider an infinitesimal flow graph  $A = (G_A, s_A, t_A)$ , for which  $G_A$  is 1-connected (as a graph). Since A is infinitesimal,  $s_A = t_A$ , so A does not possess a splitting vertex. By 1-connectedness, however,  $G_A$  contains a cut vertex.

DEFINITION 2.8 (Flow graph splitting). Suppose vertex w is a splitting vertex for flow graph  $A = (G_A, s_A, t_A)$ . Take  $t_{A_s^w}$  to be a new vertex (not present in  $V[G_A^s(w)] \cup V[G_A^{\epsilon}(w)])$ , and define flow graph  $A_s^w = (G_{A_s^w}, s_{A_s^w}, t_{A_s^w})$  as follows:

$$\begin{split} V[G_{A_s^w}] &= V[G_A^s(w)] \cup V[G_A^\epsilon(w)] \cup \{t_{A_s^w}\}, \\ E[G_{A_s^w}] &= E[G_A^s(w)] \cup E[G_A^\epsilon(w)] \\ & \cup \{(t_{A_s^w}, u) \mid (w, u) \in E[G_A], u \in (V[G_A^s(w)] \cup V[G_A^\epsilon(w)])\} \\ & \cup \{(u, t_{A_s^w}) \mid (u, w) \in E[G_A], u \in (V[G_A^s(w)] \cup V[G_A^\epsilon(w)])\}, \\ s_{A_s^w} &= i_s^{-1}(s_A). \end{split}$$

Analogously, let  $s_{A_t^w}$  be a vertex not present in  $V[G_A^t(w)]$ . Define  $A_t^w$  to be the flow

graph  $(G_{A_t^w}, s_{A_t^w}, t_{A_t^w})$  as follows.

$$\begin{split} V[G_{A_t^w}] &= V[G_A^t(w)] \cup \{s_{A_t^w}\},\\ E[G_{A_t^w}] &= E[G_A^t(w)]\\ & \cup \{(s_{A_t^w}, u) \mid (w, u) \in E[G_A], u \in V[G_A^t(w)]\} \\ & \cup \{(u, s_{A_t^w}) \mid (u, w) \in E[G_A], u \in V[G_A^t(w)]\},\\ t_{A_t^w} &= i_t^{-1}(t_A). \end{split}$$

The pair of flow graphs  $(A_s^w, A_t^w)$  is referred to as the splitting of A induced by w.

LEMMA 2.9 ( $\oplus$ -Irreducibility Lemma). Flow graph  $A = (G_A, s_A, t_A)$  is  $\oplus$ -reducible if and only if  $V[G_A]$  contains a splitting vertex for A.

*Proof.* If  $A = B \oplus C$ , then

$$w = \sigma_{t_B \approx s_C}^{\oplus -1}(t_B) = \tau_{t_B \approx s_C}^{\oplus -1}(s_C)$$

is a splitting vertex for A (see expression (1.1) on page 3 for definitions of the  $\sigma$  and  $\tau$  injections). Conversely, if w is a splitting vertex for A, then  $A = A_s^w \bigoplus A_t^w$ .  $\square$ 

Suppose  $A = A_0 \bigoplus A_1$  and  $B = B_0 \bigoplus B_1$  are two flow graphs. If  $A_i = B_i$  (i = 0, 1) then A = B. We would like to investigate the extent to which the converse is true. Towards this, we introduce the following property.

DEFINITION 2.10. (Undirected Form) Let G = (V, E) be a directed multigraph. Then the undirected form  $U(G) = (V, \overline{E})$  is defined to be the undirected multigraph on V, in which there is an undirected edge (u, v) in  $\overline{E}$  for each directed edge from u to v or from v to u in E. For any directed edge  $e \in E$ , its corresponding undirected representative in  $\overline{E}$  is denoted as U(e). In short, U is an operation on directed edges which "forgets" their orientations. We extend U to act on sequences of edges from E in the obvious manner. For any undirected edge  $e' \in \overline{E}$ , its corresponding directed representative in E is denoted as  $D_G(e')$ . In short,  $D_G$  is an operation on undirected edges which assigns orientations according to the orientations of edges in G. We extend  $D_G$  to act on sequences of edges from  $\overline{E}$  in the obvious manner.

DEFINITION 2.11. Let  $A = (G_A, s_A, t_A)$  be an arbitrary flow graph. An edge e in  $E[G_A]$  is said to have the ST Property if in the undirected graph  $U(G_A)$  there is a (non self-intersecting) path from  $s_A$  to  $t_A$  which traverses U(e). We say A is an ST-flow graph if every edge e in  $E[G_A]$  has the ST Property. The set of all ST-flow graphs will be denoted  $F^{ST} \subset F$ .

Clearly, no ST-flow graph is infinitesimal. Hence if the source and target vertices of an ST-flow graph coincide, then it must be trivial. ST-flow graphs have several nice features which will be useful.

Lemma 2.12.  $F^{ST}$  is closed under  $\oplus$ .

Proof. If A and B are ST-flow graphs, then any edge e in  $G_{A \oplus B}$  is either in  $Im(\sigma_{t_A \approx s_B}^{\oplus})$  or in  $Im(\tau_{t_A \approx s_B}^{\oplus})$ . Suppose e is in  $Im(\sigma_{t_A \approx s_B}^{\oplus})$ . Since A is an ST-flow graph, there is a path p from  $s_A$  to  $t_A$  in  $U(G_A)$  which contains  $U(\sigma_{t_A \approx s_B}^{\oplus -1}(e))$ ; and since B is an ST-flow graph, there is a path q in  $U(G_B)$  from  $s_B$  to  $t_B$ . Then p concatenated with q is a path in  $U(G_{A \oplus B})$  which contains U(e) and connects  $s_{A \oplus B}$  to  $t_{A \oplus B}$ . The case when e is in  $Im(\tau_{t_A \approx s_B}^{\oplus})$  is analogous.  $\square$ 

The next lemma considers the reverse implication.

LEMMA 2.13. Let  $A = (G_A, s_A, t_A)$  and  $B = (G_A, s_A, t_A)$  be flow graphs. If  $A \oplus B$  is an ST-flow graph, then A and B are ST-flow graphs.

*Proof.* Suppose, towards contradiction, that A is not an ST-flow graph. Then there is an edge e that does not have the ST Property in A. So e is not on any path from  $s_A$  to  $t_A$  in  $G_A$ .

Suppose, towards contradiction, that the edge  $\sigma_{t_A \approx s_B}^{\oplus}(e)$  is on a path p from  $s_{A \oplus B}$  to  $t_{A \oplus B}$  in  $G_{A \oplus B}$ . Then, the restriction of this p to  $Im(\sigma^{\oplus})$  contradicts the fact that e does not have the ST Property. So there is no path from  $s_{A \oplus B}$  to  $t_{A \oplus B}$  in  $G_{A \oplus B}$  containing  $\sigma_{t_A \approx s_B}^{\oplus}(e)$ .

Thus  $\sigma_{t_A \approx s_B}^{\oplus}(e)$  does not have the ST Property in  $A \oplus B$ . This is a contradiction, hence A must be an ST-flow graph. The case where we assume B is not an ST-flow graph leads to a similar contradiction.  $\Box$ 

Let Let  $A = (G_A, s_A, t_A)$  be an arbitrary flow graph. Consider two edges e and e' in  $G_A$ . Suppose that e does not have the ST Property. Then e cannot lie inside any path from s to t in U(G).

- Suppose e' has the ST Property in G. Then there is path p from s to t in U(G) containing e'. Certainly p cannot contain e, since e does not have the ST Property. So p survives in  $G \setminus e$ , and hence e' has the ST Property in  $G \setminus e$ .
- Suppose e' does not have the ST Property in G. Then there is no path p from s to t in U(G) containing e'. But then the deletion of e from G does not change this fact, since deletion of edges cannot create new paths. It follows that in e' does not have the ST Property in  $G \setminus e$ .

We have shown that if an edge e does not have the ST Property, then its deletion does not affect other edges e', for whom the ST Property stays unchanged from G to  $G \setminus e$ . It follows that sequentially deleting all edges which violate the ST Property of G yields a unique directed connected graph whose vertex set contains the source and target vertices of the original flow graph. This leads to the next definition, by which we can approximate general flow graphs using suitable large ST-flow subgraphs.

DEFINITION 2.14. Let  $A = (G_A, s_A, t_A)$  be a general flow graph. We define the ST-core of A to be the largest ST-flow graph  $\hat{A} = (\hat{G}_A, s_A, t_A)$  contained in A, where  $\hat{G}_A = (\hat{V}[G_A], \hat{E}[G_A])$  is obtained by deleting all edges from  $G_A$  which do not have the ST Property and considering the connected subgraph induced by the remaining edges. Note that  $\hat{A}$  is an ST-flow graph on a vertex set  $\hat{V}[G_A] \supseteq \{s_A, t_A\}$ .

Lemma 2.15. The st-core operation distributes over  $\oplus$ .

*Proof.* We would like to show that  $\hat{A} \oplus \hat{B}$  is isomorphic to the st-core $(A \oplus B)$ . Let e be an arbitrary edge in the st-core $(A \oplus B)$ . Then there is a path p from  $s_{A \oplus B}$  to  $t_{A \oplus B}$  in  $U(G_{A \oplus B})$ , such that p contains U(e). But p must pass through the cut vertex  $w = \sigma_{t_A \approx s_B}^{\oplus}(t_A)$  in  $G_{A \oplus B}$ .

- (i) If e is in Im(σ<sup>⊕</sup><sub>t<sub>A</sub>≈s<sub>B</sub></sub>), let p' be the initial segment of p which connects s<sub>A⊕B</sub> to w in U(G<sub>A⊕B</sub>); then U(σ<sup>⊕-1</sup><sub>t<sub>A</sub>≈s<sub>B</sub></sub>(D<sub>G<sub>A⊕B</sub></sub>(p'))) is a path containing U(e) connecting s<sub>A</sub> to t<sub>A</sub> in U(G<sub>A</sub>), so e' = σ<sup>⊕-1</sup><sub>t<sub>A</sub>≈s<sub>B</sub></sub>(e) is in Â.
  (ii) If e is in Im(τ<sup>⊕</sup><sub>t<sub>A</sub>≈s<sub>B</sub></sub>), let p' be the final segment of p which connects w to
- (ii) If e is in  $Im(\tau_{t_A \approx s_B}^{\oplus})$ , let p' be the final segment of p which connects w to  $t_{A \oplus B}$  in  $U(G_{A \oplus B})$ ; then  $U(\tau_{t_A \approx s_B}^{\oplus -1}(D_{G_A \oplus B}(p')))$  is a path containing U(e) connecting  $s_B$  to  $t_B$  in  $U(G_B)$ , so  $e' = \tau_{t_A \approx s_B}^{\oplus -1}(e)$  is in  $\hat{B}$ .

This shows that  $\hat{A} \oplus \hat{B}$  contains st-core $(A \oplus B)$ .

Conversely, let e' be an edge in A. Then there is a path p' containing e' which connects  $s_A$  to  $t_A$  in  $U(G_A)$ . It follows that  $U(\sigma_{t_A \approx s_B}^{\oplus}(D_{G_A}(p')))$  connects  $s_{A \oplus B}$ to the cut vertex  $w = \sigma_{t_A \approx s_B}^{\oplus}(t_A)$  in  $U(G_{A \oplus B})$ . Let p'' be a path from  $s_B$  to  $t_B$ in  $U(G_B)$ . Then by concatenating  $U(\sigma_{t_A \approx s_B}^{\oplus}(D_{G_A}(p')))$  with  $U(\tau_{t_A \approx s_B}^{\oplus}(D_{G_B}(p'')))$ we obtain a path p connecting  $s_{A \oplus B}$  to  $t_{A \oplus B}$  through  $\sigma_{t_A \approx s_B}^{\oplus}(e)$  in  $U(G_{A \oplus B})$ . A similar argument can be carried out when e' is an edge in  $\hat{B}$ . This shows that the ST-core $(A \oplus B)$  contains  $\hat{A} \oplus \hat{B}$ .

The next lemma stands in contrast to Remark 2.7.

LEMMA 2.16. Let  $A = (G_A, s_A, t_A)$  be an ST-flow graph, and let  $w \neq s_A, t_A$  be a cut vertex of  $G_A$ . Then w is a splitting vertex of A.

*Proof.* Let w be a cut vertex of  $G_A$ . Delete w and denote the union of those components that contain neither  $s_A$  nor  $t_A$  as  $G_A^{\epsilon}(w)$  (see Definition 2.6). Towards contradiction, suppose  $V[G_A^{\epsilon}(w)] \neq \emptyset$ ; let v be any vertex therein. Then any walk from  $s_A$  to  $t_A$  in  $U(G_A)$  through v must visit w twice and hence self-intersect. It follows that edges incident to w violate the assumption that A has Property ST (see Definition 2.11). Hence it must be that  $V[G_A^{\epsilon}(w)] = \emptyset$ .

Since w is a cut vertex of  $G_A$  it's deletion generates at least two components. Towards contradiction, let  $s_A$  and  $t_A$  lie in the same component. Then there is at least one component which contains neither  $s_A$  nor  $t_A$ , and so  $V[G_A^{\epsilon}(w)] \neq \emptyset$ . We have shown that the deletion of w generates precisely two components, and that  $s_A$ and  $t_A$  lie in distinct components. Hence w is a splitting vertex of A.  $\square$ 

LEMMA 2.17 (Crossing Summands Lemma). Let  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$  be flow graphs. Let  $C = (G_C, s_C, t_C)$  be an ST-flow graph with

$$\alpha: G_C \hookrightarrow G_{A \oplus B}$$

be a graph embedding satisfying  $\alpha(s_C) = s_{A \oplus B}$  and  $\alpha(G_C) \not\subset Im(\sigma_{t_A \approx s_B}^{\oplus})$ . Then

$$\alpha^{-1}\sigma_{t_A\approx s_B}^{\oplus}(t_A) = \alpha^{-1}\tau_{t_A\approx s_B}^{\oplus}(s_B)$$

is a splitting vertex for C.

*Proof.* Let  $w = \sigma_{t_A \approx s_B}^{\oplus}(t_A) = \tau_{t_A \approx s_B}^{\oplus}(s_B)$ . Clearly, w is a splitting vertex in  $A \oplus B$  and hence a cut vertex in  $G_{A \oplus B}$ . Let v be a vertex in  $\alpha(G_C)$  that is not in  $Im(\sigma_{t_A \approx s_B}^{\oplus})$ . Since w is a cut vertex in  $G_{A \oplus B}$ , every path from  $s_{A \oplus B}$  to v in  $U(G_{A \oplus B})$  must pass through w. Given such a path

$$p = (p_0 = s_{A \oplus B}, p_1, \dots, p_{l-1}, p_l = w, p_{l+1}, \dots, p_{|p|-1}, p_{|p|} = v),$$

which lies entirely in  $U(Im(\alpha))$ , we define the pullback  $\alpha^{-1}(p)$  to be a path in  $U(G_C)$  in the obvious manner. By considering pullbacks of paths from  $s_{A \oplus B}$  to v in  $U(G_{A \oplus B})$ , we see that every path from  $s_C$  to  $\alpha^{-1}(v)$  in  $U(G_C)$  must pass through  $\alpha^{-1}(w)$ . Hence the deletion of  $\alpha^{-1}(w)$  must generate at least two components, one of which contains  $s_C$  while another contains  $\alpha^{-1}(v)$ . This shows that  $\alpha^{-1}(w)$  is a cut vertex in  $U(G_C)$ . Since C is an ST-flow graph, Lemma 2.16 holds and  $\alpha^{-1}(w)$  is a splitting vertex for C.  $\Box$ 

Consider the flow graphs A, B, C shown in Figure 2.2. Note that C is not an ST-flow graph, and  $\alpha : G_C \hookrightarrow G_{A \oplus B}$  embeds as shown. Since  $t_C$  is not a splitting vertex for C, the Crossing Summands Lemma is seen to fail when C is a general flow graph.

PROPOSITION 2.18 (Component-wise decomposition of isomorphisms under  $\oplus$ ). Suppose A and B flow graphs, expressed as sums of  $\oplus$ -irreducible ST-flow graphs as follows:

$$A = A_0 \bigoplus A_1 \bigoplus \cdots \bigoplus A_{m-1},$$
  
$$B = B_0 \bigoplus B_1 \bigoplus \cdots \bigoplus B_{n-1}.$$



FIG. 2.2. An example where the Crossing Summands Lemma fails for general flow graphs.

Then

$$A = B \Leftrightarrow m = n \text{ and } A_i = B_i \text{ for all } i = 0, \dots, m-1$$

*Proof.* [ $\Leftarrow$ ] Let  $\phi_i : A_i \to B_i$  be given component-wise isomorphisms, for  $i = 0, \ldots, m-1$ . Define  $\phi : A \to B$  by defining  $\phi|_{A_i} = \phi_i$ . Since  $t_{B_i} = \phi_i(t_{A_i}) = \phi_{i+1}(s_{A_{i+1}}) = s_{B_{i+1}}$  for  $i = 0, \ldots, m-2$ , this provides a well-defined isomorphism between A and B. Indeed, this implication holds for arbitrary flow graphs.

[⇒] We prove the statement by induction on  $\max(m, n)$ . In the case when m = n = 1, the claim is trivial. For the inductive step, let  $\phi : A \to B$  be an isomorphism. Consider  $\phi(A_0)$ , and take k to be the smallest integer in  $\{0, \ldots, n-1\}$  for which  $\phi(A_0)$  is a subgraph of  $B_0 \oplus B_1 \oplus \cdots \oplus B_k$ . Since  $A_0$  is  $\oplus$  -irreducible it contains no splitting vertices. Since  $A_0$  is an ST-flow graph, it must be that k = 0; otherwise by Lemma 2.17  $A_0$  would contain a splitting vertex, and so (by Lemma 2.9) be  $\oplus$  -reducible. Since k = 0, we have shown that  $\phi(A_0)$  is a subgraph of  $B_0$ . Now, repeating the argument for  $B_0$  using  $\phi^{-1}$ , we see that  $\phi^{-1}(B_0)$  is a subgraph of  $A_0$ . It follows that  $A_0$  is isomorphic to  $B_0$  under a suitable restriction of  $\phi$ . Now, since  $\phi(A) = B$  and  $\phi(A_0) = B_0$  it follows that  $\phi(A \setminus A_0) = B \setminus B_0$ , or more specifically  $\phi(A_1 \oplus \cdots \oplus A_{m-1}) = B_1 \oplus \cdots \oplus B_{n-1}$ . By inductive hypothesis, this implies that m = n and  $A_i = B_i$  for all  $i = 0, \ldots, m - 1$ .  $\square$ 

Because the Crossing Summands Lemma fails (and lies at the heart of the proof of Proposition 2.18) the latter Proposition also fails for general flow graphs. The reader can verify for example, from Figure 2.2, that  $A \oplus B = C \oplus A$  but  $A \neq C$  and  $B \neq A$ .

DEFINITION 2.19 (Splitting vertex ranking). Given flow graph  $A = (G_A, s_A, t_A)$ , let  $\chi(A) \subset V[G_A]$  be the set of all splitting vertices for A. We define the s-ranking and t-ranking functions  $r_s^A, r_t^A : \chi(A) \to \mathbb{N}$  as follows:

$$\begin{split} r^A_s(w) &= |V[G^s_A(w)] \cap \chi(A)|, \\ r^A_t(w) &= |V[G^t_A(w)] \cap \chi(A)|. \end{split}$$

When it is clear from the context, we denote  $r_s(w) = r_s^A(w)$  and  $r_t(w) = r_t^A(w)$ .

LEMMA 2.20. Let  $w \in \chi(A)$  be a splitting vertex for flow graph  $A = (G_A, s_A, t_A)$ . Then for all  $u \in V[G^s_A(w)] \cap \chi(A)$ :

$$r_s(u) < r_s(w),$$
  

$$r_t(u) > r_t(w);$$

and for all  $u \in V[G_A^t(w)] \cap \chi(A)$ :

$$r_s(u) > r_s(w),$$
  
$$r_t(u) < r_t(w).$$

*Proof.* First, note that for any u in  $(V[G_A^s(w)] \cup V[G_A^t(w)]) \cap \chi(A)$ 

$$r_s(u) + r_t(u) + 1 = |\chi(A)|$$
(2.1)

Now if  $u \in V[G_A^s(w)] \cap \chi(A)$ , then since  $w \in (V[G_A^t(u)] \setminus V[G_A^t(w)]) \cap \chi(A)$ , so it follows that  $V[G_A^t(w)] \subsetneq V[G_A^t(u)]$ . But then  $r_t(w) < r_t(u)$ . By expression 2.1 above, it follows that  $r_s(w) > r_s(u)$ . The proof for the case when  $u \in V[G_A^t(w)] \cap \chi(A)$  is analogous.  $\square$ 

LEMMA 2.21. Given a flow graph  $A = (G_A, s_A, t_A)$ , for each  $i = 0, 1, ..., |\chi(A)| - 1$  there is a unique vertex  $v_i$  in  $\chi(A)$  with the property that  $r_s(v_i) = i$ .

*Proof.* First we note that one cannot have two distinct vertices v, v' having  $r_s(v) = r_s(v')$ , since either  $v \in V[G_A^s(v')]$  or  $v' \in V[G_A^s(v)]$ , and so by Lemma 2.20 it follows that  $r_s(v) \neq r_s(v')$ . Base case: i = 0. Let  $w_0$  be any vertex in  $\chi(A)$ . If  $r_s(w) > 0$ , then  $V[G_A^s(w)] \cap \chi(A)$  is not empty. So let  $w_1$  be any vertex in  $V[G_A^s(w)] \cap \chi(A)$ . By Lemma 2.20,  $r_s(w_1) < r_s(w_0)$ . Repeating in this fashion, after finitely many steps  $w_0 \rightsquigarrow w_1 \rightsquigarrow \ldots$  we find some vertex  $v_0$  for which  $r_s(v_0) = 0$ . Inductive step i + 1: Let  $v_i$  be the unique vertex in  $\chi(A)$  having  $r_s(v_i) = i$ . Define  $v_{i+1}$  to be the vertex in  $V[G_A^s(v_i)] \cap \chi(A)$  for whose s-rank is minimal. Since

$$V[G_{A}^{s}(v_{i+1})] \cap \chi(A) = [V[G_{A}^{s}(v_{i})] \cap \chi(A)] \cup \{v_{i}\},$$

it follows that  $r_s(v_{i+1}) = r_s(v_i) + 1 = i + 1$ , hence the result.  $\Box$ 

DEFINITION 2.22 (Canonical  $\bigoplus$  -decomposition). Let  $A = (G_A, s_A, t_A)$  be a flow graph. Take  $\chi(A) = \{v_0, v_1, \ldots, v_{|\chi(A)|-1}\}$  to be the set of splitting vertices for A, ordered according to the indexing scheme postulated in Lemma 2.21. Define  $A^{(0)} = A_s^{v_0}, \bar{A}^{(0)} = A_t^{v_0}$ , and then for each  $i = 1, 2, \ldots, |\chi(A)| - 1$ , put

$$A^{(i)} = (\bar{A}^{(i-1)})_s^{v_i},$$
  
$$\bar{A}^{(i)} = (\bar{A}^{(i-1)})_t^{v_i}.$$

We shall denote  $\bar{A}^{(|\chi(A)|-1)}$  as  $A^{(|\chi(A)|)}$ . The canonical  $\oplus$  -decomposition of A is defined to be the sequence

$$\langle A \rangle \stackrel{\text{def}}{=} (A^{(0)}, A^{(1)}, \dots, A^{(|\chi(A)|-1)}, A^{(|\chi(A)|)}).$$

Note that the effectiveness of this definition guarantees uniqueness of the decomposition.

LEMMA 2.23 ( $\oplus$ -decompositions for sums). Given  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ , two flow graphs with their respective canonical  $\oplus$ -decompositions

 $\langle A \rangle$  and  $\langle B \rangle$ . If B is s-standard, then the canonical  $\oplus$  -decomposition of  $A \oplus B$  is  $\langle A \rangle \langle B \rangle$ , the concatenation of  $\langle A \rangle$  with  $\langle B \rangle$ .

*Proof.* First, note that  $|\chi(A \oplus B)| = |\chi(A)| + |\chi(B)| + 1$ . More specifically, if

$$\chi(A) = \{u_0, u_1, \dots, u_{|\chi(A)|-1}\}, \text{ and}$$
  
$$\chi(B) = \{v_0, v_1, \dots, v_{|\chi(B)|-1}\}$$

are the sets of splitting vertices for A and B respectively, ordered by ascending srank, according to the indexing scheme postulated in Lemma 2.21, then  $A \oplus B$  has splitting vertices:

$$\chi(A \oplus B) = \left\{ \sigma_{t_A \approx s_B}^{\oplus}(u_0), \sigma_{t_A \approx s_B}^{\oplus}(u_1), \dots, \sigma_{t_A \approx s_B}^{\oplus}(u_{|\chi(A)|-1}), \\ \sigma_{t_A \approx s_B}^{\oplus}(t_A) = \tau_{t_A \approx s_B}^{\oplus}(s_B), \\ \tau_{t_A \approx s_B}^{\oplus}(v_0), \tau_{t_A \approx s_B}^{\oplus}(v_1), \dots, \tau_{t_A \approx s_B}^{\oplus}(v_{|\chi(B)|-1}) \right\}.$$

But since  $\sigma_{t_A \approx s_B}^{\oplus}$  and  $\tau_{t_A \approx s_B}^{\oplus}$  are injections,

$$(A \oplus B)^{(i)} = \begin{cases} A^{(i)} & \text{for } 0 \leq i < |\chi(A)|, \\ \bar{A}^{(|\chi(A)|-1)} & \text{for } i = |\chi(A)|, \\ B^{(i-|\chi(A)|-1)} & \text{for } |\chi(A)| < i \leq |\chi(A)| + |\chi(B)| \\ \bar{B}^{(|\chi(B)|-1)} & \text{for } i = |\chi(A)| + |\chi(B)| + 1. \end{cases}$$

It follows that  $\langle A \oplus B \rangle = \langle A \rangle \langle B \rangle$ .

We note that if B is not s-standard then  $(A \oplus B)^{(|\chi(A)|)}$  is not equal to  $\overline{A}^{(|\chi(A)|-1)}$ and the Lemma 2.23 fails to hold. The reader can verify this by considering the flow graphs A and B shown in Figure 2.2.

PROPOSITION 2.24 (Correctness of the  $\oplus$  -decomposition). Consider the canonical  $\oplus$  -decomposition of A as given in Definition 2.22:

$$\langle A^{(0)}, A^{(1)}, \dots, A^{(|\chi(A)|)} \rangle$$

Then  $A = A^{(0)} \bigoplus A^{(1)} \bigoplus A^{(2)} \cdots \bigoplus A^{(|\chi(A)|-1)} \bigoplus A^{(|\chi(A)|)}$ , and every summand is  $\bigoplus$  -irreducible.

Proof. Since  $r_s(v_i) = i$ , it follows that  $\chi(A^{(i)}) = \emptyset$  for all  $i = 0, 1, \ldots, |\chi(A)|$ . Since each summand has no splitting vertices, by Lemma 2.9, each is  $\oplus$  -irreducible. We prove the Proposition by induction on  $|\chi(A)|$ . The base case when  $|\chi(A)| = 1$ is straightforward, since Definition 2.22 specified  $A^{(0)} = A_s^{v_0}$  and  $\bar{A}^{(0)} = A_t^{v_0}$ . Then, since  $A_s^{v_0} \oplus A_t^{v_0} = A$  for any splitting vertex  $v_0$ , the result follows. Suppose the Proposition has been proved for all flow graphs B which enjoy  $|\chi(B)| \leq k$ . Let Abe a flow graph with  $|\chi(A)| = k + 1$ . Unravelling Definition 2.22 yields  $A^{(k+1)} = \bar{A}^{(k)} = (\bar{A}^{(k-1)})_t^{v_k} = A_t^{v_k}$ . Then since  $A = A_s^{v_k} \oplus A_t^{v_k}$  and  $|\chi(A_s^{v_k})| = k$ , by inductive hypothesis and Lemma 2.23,

$$A = A_s^{v_k} \bigoplus A_t^{v_k}$$
  
=  $\langle A_s^{v_k} \rangle \bigoplus A_t^{v_k}$   
=  $(A_s^{v_k})^{(0)} \bigoplus \dots (A_s^{v_k})^{(k)} \bigoplus A_t^{v_k}$   
=  $A^{(0)} \bigoplus A^{(1)} \bigoplus A^{(2)} \dots \bigoplus A^{(k)} \bigoplus A^{(k+1)}.$ 

The result follows.  $\Box$ 

LEMMA 2.25 (Left-cancellation law for  $\oplus$ ). Let A, B, C be st-flow graphs.

$$A \oplus B = A \oplus C \Rightarrow B = C.$$

*Proof.* Let  $\langle A \oplus B \rangle = \langle A \rangle \langle B \rangle$  be the canonical  $\oplus$  -decomposition of  $A \oplus B$ , and  $\langle A \oplus C \rangle = \langle A \rangle \langle C \rangle$  be the canonical  $\oplus$  -decomposition of  $A \oplus C$  respectively.

Fix an isomorphism  $\phi$  between  $A \bigoplus B \to A \bigoplus C$ . By Proposition 2.18,  $\phi$  maps the elements of  $\langle A \rangle \langle B \rangle$  componentwise to the elements of  $\langle A \rangle \langle C \rangle$ . It follows that  $\phi$ maps the elements of  $\langle B \rangle$  componentwise to the elements of  $\langle C \rangle$ . Hence a suitable restriction of  $\phi$  maps B injectively onto C, proving the claim.  $\Box$ 

The next lemma is proved in a manner analogous to Lemma 2.25.

LEMMA 2.26 (Right-cancellation law for  $\oplus$ ). Let A, B, C be flow graphs.

$$B \oplus A = C \oplus A \Rightarrow B = C.$$

PROPOSITION 2.27 (Commutativity condition for  $\oplus$ ). Given ST-flow graphs  $A = (G_A, s_A, t_A)$  and  $B = (G_B, s_B, t_B)$ ,

$$A \oplus B = B \oplus A$$

iff there exists a flow graph C and integers  $k_1, k_2$  in  $\mathbb{N}$  such that

$$A = k_1 C, \text{ and} \\ B = k_2 C.$$

*Proof.*  $[\Leftarrow]$  If  $A = k_1C$  and  $B = k_2C$ , then  $A \oplus B = (k_1 + k_2)C = B \oplus A$ .

 $[\Rightarrow]$  The proof is carried by induction on max $(|\chi(A)|, |\chi(B)|)$ . Consider the canonical decompositions of A and B,

$$A = A^{(0)} \bigoplus A^{(1)} \bigoplus A^{(2)} \bigoplus \cdots \bigoplus A^{(|\chi(A)|-1)} \bigoplus A^{(|\chi(A)|)},$$
  
$$B = B^{(0)} \bigoplus B^{(1)} \bigoplus B^{(2)} \bigoplus \cdots \bigoplus B^{(|\chi(B)|-1)} \bigoplus B^{(|\chi(B)|)}.$$

If  $|\chi(A)| = |\chi(B)|$  then Proposition 2.18 tells us that an isomorphism  $\phi : A \bigoplus B \rightarrow B \bigoplus A$  restricts on the first summand A to yield an isomorphism from A to B. So in this case, we can take C = A = B and  $k_1 = k_2 = 1$ . This proves the case  $\max(|\chi(A)|, |\chi(B)|) = 0$ , which forms the basis of the induction.

Suppose that  $\max(|\chi(A)|, |\chi(B)|) > 0$ , and  $|\chi(A)| \neq |\chi(B)|$ . Without loss of generality, suppose  $|\chi(A)| < |\chi(B)|$ . Then  $A^{(i)} = B^{(i)}$  for  $i = 0, \ldots |\chi(A)|$ . It follows that

$$B^{i+(|\chi(A)|+1)} = B^{(i)} \text{ for } i = 0, \dots, |\chi(B)| - (|\chi(A)| + 1),$$
(2.2)

$$B^{i-(|\chi(A)|+1)} = B^{(i)} \text{ for } i = (|\chi(A)|+1), \dots, |\chi(B)|.$$
(2.3)

If  $(|\chi(B)| + 1)$  is divisible by  $(|\chi(A)| + 1)$ , then expressions (2.2) and (2.3) above are in fact equivalent, and in this setting, we take C = A,  $k_1 = 1$  and  $k_2 = \frac{(|\chi(B)|+1)}{(|\chi(A)|+1)}$  in order to satisfy the proposition. Suppose now that  $(|\chi(B)| + 1)$  is *not* divisible by  $(|\chi(A)| + 1)$ . Put

$$d = \left\lfloor \frac{(|\chi(B)| + 1)}{(|\chi(A)| + 1)} \right\rfloor, \text{ and} r = (|\chi(B)| + 1) \mod (|\chi(A)| + 1).$$

and define

$$X = B^{(0)} \bigoplus B^{(1)} \cdots \bigoplus B^{(r-1)},$$
  

$$Y = B^{(r)} \bigoplus B^{(r+1)} \cdots \bigoplus B^{(|\chi(A)|)}$$

Note that  $B = dA \bigoplus X$  and

$$A = A^{(0)} \bigoplus A^{(1)} \bigoplus A^{(2)} \bigoplus \cdots \bigoplus A^{(|\chi(A)|-1)} \bigoplus A^{(|\chi(A)|)}$$
  
=  $B^{(0)} \bigoplus B^{(1)} \bigoplus \cdots B^{(r-1)} \bigoplus B^{(r)} \bigoplus B^{(r+1)} \cdots \bigoplus B^{(|\chi(A)|-1)} \bigoplus B^{(|\chi(A)|)}$   
=  $X \bigoplus Y.$ 

It follows that  $B = d(X \oplus Y) \oplus X$ . On the other hand,  $X \oplus Y = A = Y \oplus X$  (see Figure 2.3), since

$$X \bigoplus Y = A$$
  
=  $A^{(0)} \bigoplus A^{(1)} \bigoplus \cdots \bigoplus A^{(|\chi(A)|)}$   
=  $B^{(0)} \bigoplus B^{(1)} \bigoplus \cdots \bigoplus B^{(r)} \bigoplus \cdots \bigoplus B^{(|\chi(A)|-1)} \bigoplus B^{(|\chi(A)|)}$   
=  $B^{(r)} \bigoplus \cdots \bigoplus B^{(|\chi(B)|-1)} \bigoplus B^{(|\chi(B)|)} \bigoplus B^{(0)} \bigoplus \cdots \bigoplus B^{(r-1)}$   
=  $Y \bigoplus X$ .



FIG. 2.3. Inductive step showing  $X \oplus Y = A = Y \oplus X$ .

Since  $r \neq 0$  the inductive hypothesis applies to the flow graphs X, Y, i.e. there exists some flow graph Z and suitable integers  $l_1$ ,  $l_2$  so that  $X = l_1Z$ ,  $Y = l_2Z$ . It follows that

$$A = X \bigoplus Y = l_1 Z \bigoplus l_2 Z = (l_1 + l_2) Z \text{ and}$$
  
$$B = dA \bigoplus X = d(X \bigoplus Y) \bigoplus X = d(l_1 + l_2) Z \bigoplus l_1 Z = ((d+1)l_1 + l_2) Z$$

So taking C = Z,  $k_1 = l_1 + l_2$  and  $k_2 = (d+1)l_1 + l_2$ , the Proposition is proved.

Note that Proposition 2.27 fails if A and B are both not sT-flow graphs. To see this, consider the example shown in Figure 2.4.



FIG. 2.4. An example illustrating that the Commutativity Condition fails for general flow graphs.

### **2.2.** Multiplicative Properties. In this section we present properties of $\otimes$ .

LEMMA 2.28. Given flow graphs A and B, there is a natural bijective correspondence

$$\Lambda_{A,B}: E[A \otimes B] \to E[G_A] \times E[G_B]$$

Proof. Fix an edge e in  $E[G_A \otimes G_B]$ . Then e appears in  $(A \otimes_{\eta} B)$  at some stage i where  $1 \leq i \leq |E[G_A]|$  (where  $\eta$  is the enumeration specified in Definition 1.9). We define  $\lambda_A(e)$  to be the edge  $e_i \in E[G_A]$ . At stage i we effectively replace edge  $e_i = (u_i, v_i)$  with a new disjoint copy of  $G_B$ -by gluing  $s_B$  with  $u_i$  and  $t_B$  with  $v_i$ . Thus the edge e corresponds to some edge  $\lambda_B(e)$  in this new disjoint copy of  $G_B$ . The desired bijection  $e \mapsto \Lambda_{A,B}(e) = (\lambda_A(e), \lambda_B(e))$  is thus obtained. Note that the bijection  $\Lambda_{A,B}$  is independent of the enumeration  $\eta$  of the edges of  $E[G_A]$  which appears in the definition of  $A \otimes B$ .  $\square$ 

Lemma 2.29 (Associativity of  $\otimes$ ). The operation  $\otimes$  is associative.

*Proof.* Given flow graphs  $A = (G_A, s_A, t_A)$ ,  $B = (G_B, s_B, t_B)$ ,  $C = (G_C, s_C, t_C)$ , we want to show:

$$(A \otimes B) \otimes C = A \otimes (B \otimes C).$$

By Lemma 2.28, the map  $\Lambda_A \bigotimes_{B,C}$  is a bijective correspondence between the edges of  $(A \otimes B) \otimes C$  and  $(E[G_A] \times E[G_B]) \times E[G_C]$ . Likewise, the edges of  $A \otimes (B \otimes C)$  are in bijective correspondence with  $E[G_A] \times (E[G_B] \times E[G_C])$ , via  $\Lambda_{A,B} \bigotimes_{C}$ . Obviously  $(E[G_A] \times E[G_B]) \times E[G_C]$  is in bijective correspondence with  $E[G_A] \times (E[G_B] \times E[G_C])$ , by the map  $\pi : ((e_1, e_2), e_3) \mapsto (e_1, (e_2, e_3))$ . Then the composite map

$$\mu = \Lambda_{A,B} \bigotimes_C \circ \pi \circ \Lambda_A \bigotimes_{B,C}$$

is an isomorphism of flow graphs which carries  $(A \otimes B) \otimes C$  to  $A \otimes (B \otimes C)$ .

EXAMPLE 2.30. Let A be the flow graph consisting of a directed cycle of length 3 and let source and target vertices be any two vertices on this cycle. Then it is easy to check that there is no flow graph isomorphism between  $A \otimes F_2$  and  $F_2 \otimes A$  (see Figure 2.5).



FIG. 2.5. Example showing the non-commutativity of multiplication in  $\mathcal{F}$ .

The previous example proves the next lemma.

LEMMA 2.31. The operation  $\otimes$  is not commutative.

LEMMA 2.32 (Right-distributivity of  $\otimes$  over  $\oplus$ ). For any flow graphs A, B, C,

$$(A \oplus B) \otimes C = (A \otimes C) \oplus (B \otimes C)$$

*Proof.* Fix  $e \in E[(A \oplus B) \otimes C]$ . Then define  $\beta_0(e) = \Lambda_{A \oplus B,C}(e)$ . Note that  $\beta_0(e) = (e', f)$ , where e' is an edge in  $E[G_{A \oplus B}]$  and f is one in  $E[G_C]$ . Define  $\beta_1 : E[G_{A \oplus B}] \to E[G_A] \cup E[G_B]$  so that

$$\beta_1(e) = \begin{cases} \sigma_{t_A \approx s_B}^{\oplus -1}(e) & \text{if} \quad e \in Im(\sigma_{t_A \approx s_B}^{\oplus -1}) \\ \tau_{t_A \approx s_B}^{\oplus -1}(e) & \text{if} \quad e \in Im(\tau_{t_A \approx s_B}^{\oplus -1}). \end{cases}$$

Then  $\beta_1 \circ \beta_0$  maps  $E[(A \oplus B) \otimes C]$  injectively into  $(E[G_A] \times E[G_C]) \cup (E[G_B] \times E[G_C])$ . Define  $\beta_2$  by taking

$$\beta_2(e) = \begin{cases} \Lambda_{A,C}^{-1}(e) & \text{if} \quad e \in E[G_A \bigotimes c] \\ \Lambda_{B,C}^{-1}(e) & \text{if} \quad e \in E[G_B \bigotimes c] \end{cases}$$

Then  $\beta_2$  maps  $(E[G_A] \times E[G_C]) \cup (E[G_B] \times E[G_C])$  into  $E[G_A \bigotimes _C] \cup E[G_B \bigotimes _C]$  injectively. Finally, define  $\beta_3$  by taking

$$\beta_{3}(e) = \begin{cases} \sigma_{t_{A}\bigotimes_{C}}^{\oplus} \approx s_{B}\bigotimes_{C}}(e) & \text{if} \quad e \in E[G_{A\bigotimes_{C}}] \\ \tau_{t_{A}\bigotimes_{C}}^{\oplus} \approx s_{B}\bigotimes_{C}}(e) & \text{if} \quad e \in E[G_{B\bigotimes_{C}}]. \end{cases}$$

Then  $\beta_3$  maps  $E[G_A \bigotimes_C] \cup E[G_B \bigotimes_C]$  injectively into  $E[G_{(A \bigotimes C) \oplus (B \times C)}]$ . The composite map  $\beta_3 \circ \beta_2 \circ \beta_1 \circ \beta_0$  maps the edges of  $(A \bigoplus B) \otimes C$  injectively into the

edges of  $(A \otimes C) \oplus (B \otimes C)$ , and is the desired flow graph isomorphism demonstrating the claimed equality.  $\Box$ 

Let A be the flow graph consisting of a directed cycle of length 3 taking source and target vertices to be any two vertices on this cycle. Observe that  $A \otimes (F_1 \oplus F_1) = A \otimes F_2$ , while  $(A \otimes F_1) \oplus (A \otimes F_1) = A \oplus A = 2A = F_2 \otimes A$ . Referring to Figure 2.5 again, we see that  $A \otimes F_2 \neq F_2 \otimes A$ . Thus, we have shown

LEMMA 2.33 (Non Left-distributivity of  $\otimes$  over  $\oplus$ ). There exist flow graphs A, B, C,

$$A \otimes (B \oplus C) \neq (A \otimes B) \oplus (A \otimes C)$$

DEFINITION 2.34. Given flow graphs A, B at least one of which is non-trivial, and a flow graph C, we say

$$A/B = C \text{ iff } A = C \otimes B$$
$$A \setminus B = C \text{ iff } A = B \otimes C.$$

If there is no C for which A/B = C, we say that A/B does not exist and A is not right-divisible by B. If there is no C for which  $A \setminus B = C$ , we say that  $A \setminus B$  does not exist, and A is not left-divisible by B. By convention, we say that  $(\bigcirc) / (\bigcirc)$  and  $(\bigcirc) \setminus (\bigcirc)$  are undefined.

Clearly if m and n are standard integers then  $F_m/F_n$  iff  $F_m\setminus F_n$  iff m is divisible by n.

LEMMA 2.35. For all flow graphs A, B, C

$$A/C = (A/B) \otimes (B/C)$$
$$A \setminus C = (B \setminus C) \otimes (A \setminus B)$$

whenever these graphs exist.

Proof. Suppose  $A/C = K_1$  and  $B/C = K_2$ . By definition,  $A = K_1 \otimes B$  and  $B = K_2 \otimes C$ . Thus,  $A = K_1 \otimes (K_2 \otimes C)$ , which by Lemma 2.29 is  $(K_1 \otimes K_2) \otimes C$ . Thus A/C exists and equals  $K_1 \otimes K_2 = (A/B) \otimes (B/C)$ . Suppose  $A \setminus C = K_1$  and  $B \setminus C = K_2$ . Then by definition,  $A = B \otimes K_1$  and  $B = C \otimes K_2$ . Thus,  $A = (C \otimes K_2) \otimes K_1$ , which by Lemma 2.29 is  $C \otimes (K_2 \otimes K_1)$ . Thus  $A \setminus C$  exists and equals  $K_2 \otimes K_1 = (B \setminus C) \otimes (A \setminus B)$ .  $\Box$ 

LEMMA 2.36 (Distributivity of right-divisibility over  $\oplus$ ). For all flow graphs A, B, C,

$$A/B \oplus C/B = (A \oplus C)/B$$

Proof. Suppose  $A/B = K_1$  and  $C/B = K_2$ . Then by definition,  $A = K_1 \otimes B$ and  $C = K_2 \otimes B$ . Thus  $A \oplus C = (K_1 \otimes B) \oplus (K_2 \otimes B)$  which by Lemma 2.32, equals  $(K_1 \oplus K_2) \otimes B$ . It follows that  $(A \oplus C)/B$  equals  $K_1 \oplus K_2$ , which is  $A/B \oplus C/B$ .  $\square$ 

OBSERVATION 2.37 (Non-distributivity of left-divisibility over  $\oplus$ ). Note that Lemma 2.33 can be used to construct examples that demonstrate non-distributivity of left-divisibility over  $\oplus$ . For example, let B be a directed cycle of length 3 with any two vertices as  $s_B$  and  $t_B$ . Take  $A = B \otimes F_2$ . Then  $A \setminus B = F_2$ . Now take C = B. Then  $C \setminus B = F_1$  and so  $(A \setminus B) \oplus (C \setminus B) = F_2 \oplus F_1 = F_3$ . Since  $A \oplus C \neq B \otimes F_3$ , we see that  $(A \oplus C) \setminus B \neq (A \setminus B) \oplus (C \setminus B)$ .

In Definition 2.6, we introduced the notion of a splitting vertex. Now, in unravelling information about  $\otimes$ , we require the notion of a splitting edge.

DEFINITION 2.38 (Splitting edge for a flow graph). Let  $A = (G_A, s_A, t_A)$  be a flow graph. A splitting edge of A is an edge  $e \in E[G_A]$  with the property that  $G \setminus e$ has precisely two components, one of which contains  $s_A$  and the other contains  $t_A$ . We denote the set of all splitting edges in A as  $\Delta(A)$ .

LEMMA 2.39. For any flow graph A, if A is  $\oplus$  -irreducible and  $A \neq (1)$ , then  $\Delta(A) = \emptyset$ .

*Proof.* If  $A \neq F_1$  and e = (u, v) is a splitting edge then either u or v or both must be a splitting vertex. Hence A is  $\oplus$  -reducible.  $\square$ 

LEMMA 2.40 (Splitting edges in  $\oplus$  -decompositions). Given a flow graph A, let  $\langle A \rangle$  be its  $\oplus$  -decomposition. Then there exists a map  $i : \Delta(A) \to \{0, 1, \ldots, \chi(A)\}$  which injectively associates to every splitting edge an  $\oplus$  -irreducible component in the  $\oplus$  -decomposition of A, such that  $A^{i(e)}$  is a component consisting only of edge e, and is isomorphic to  $F_1$ .

**Proof.** Appealing to Proposition 2.24, fix  $\phi$  an isomorphism from A to the component decomposition of  $\langle A \rangle$ . By Proposition 2.24, every  $A^{(i_e)}$  is  $\bigoplus$ -irreducible. By Lemma 2.39 it either has no splitting edges or it is  $F_1$ . Suppose e is a splitting edge in A. Then  $\phi(e)$  is a splitting edge inside  $A^{(i_e)}$  for some  $i_e$  in  $\{0, \ldots, |\chi(A)|\}$ . It follows that  $A^{(i_e)} = F_1$  and the map  $i: e \mapsto i_e$  enjoys the property claimed in the lemma.  $\square$ 

REMARK 2.41. Given flow graphs A and B, a splitting vertex in  $A \otimes B$  comes either from a splitting vertex of A or from a splitting vertex in (a copy of) B which lies on a splitting edge of A.

OBSERVATION 2.42 (  $\oplus$  -decompositions for products). Given flow graphs A, B,

Since  $A^{(i)}$  is  $\oplus$ -irreducible, by Lemma 2.9 we know that  $\chi(A^{(i)}) = \emptyset$ . Case (i).  $A^{(i)} \neq F_1$ . Then by Lemma 2.24,  $\Delta(A^{(i)}) = \emptyset$ . It follows from Remark 2.41 that  $A^{(i)} \otimes B$  has no splitting vertices, so by Lemma 2.9, it is  $\oplus$ -irreducible. Hence,  $\langle A^{(i)} \otimes B \rangle$  is a one-element sequence consisting of  $A^{(i)} \otimes B$ . Case (ii).  $A^{(i)} = F_1$ . Then  $A^{(i)} \otimes B = B$ , so  $A^{(i)} \otimes B$  is  $\oplus$ -irreducible iff B is  $\oplus$ -irreducible. In this case  $\langle A^{(i)} \otimes B \rangle = \langle B \rangle$  is a  $(|\chi(B)| + 1)$ -element subsequence consisting of the  $\oplus$ -decomposition of B.

LEMMA 2.43 (  $\oplus$  -decomposition length for products). Given flow graphs A, B,

$$|\chi(A \otimes B)| = |\chi(A)| + |\Delta(A)| \cdot |\chi(B)|.$$

Proof. Consider each  $\oplus$ -irreducible component  $A^{(i)}$  in  $\langle A \rangle$ . If  $A^{(i)} = F_1$  then by Observation 2.42, it contributes  $(|\chi(B)| + 1)$  components in  $\langle A \oplus B \rangle$ . There are  $|\Delta(A)|$  components in  $\langle A \rangle$  which are isomorphic to  $F_1$ , so these together account for  $|\Delta(A)| \cdot (|\chi(B)| + 1)$  components in  $\langle A \otimes B \rangle$ . The remaining  $|\chi(A)| + 1 - |\Delta(A)|$  components in  $\langle A \rangle$  (again by Observation 2.42) each contribute 1 component to  $\langle A \otimes B \rangle$ . It follows that the total number of components in  $\langle A \otimes B \rangle$  is

$$\begin{aligned} |\chi(A)| + 1 - |\Delta(A)| + |\Delta(A)|(|\chi(B)| + 1) = \\ |\chi(A)| + |\Delta(A)||\chi(B)| + 1. \end{aligned}$$

This shows that  $|\chi(A \otimes B)| = |\chi(A)| + |\Delta(A)||\chi(B)|$  as desired.

PROPOSITION 2.44 ( $\oplus$ -irreducibility for products). Let  $A \neq (1)$  be a flow graph. Then, A is  $\oplus$ -irreducible iff for all flow graphs  $B, A \otimes B$  is  $\oplus$ -irreducible. Proof. [ $\Leftarrow$ ] Taking B = (1), we see that A is  $\oplus$ -irreducible.

 $[\Rightarrow]$  If A is  $\oplus$  -irreducible, then  $\chi(A) = \emptyset$ . Since  $A \neq (1)$ , it follows that  $\Delta(A) = \emptyset$ . So by Lemma 2.43, for any flow graph B,  $\chi(A \otimes B) = \emptyset$ . It follows that  $A \otimes B$  is  $\oplus$  -irreducible.  $\square$ 

DEFINITION 2.45. A flow graph A is called right-prime if A/B exists only for B = (1) or B = A. Similarly a flow graph A is called left-prime if  $A \setminus B$  exists only for B = (1) or B = A.

Note that a natural number n is prime iff the flow graph  $F_n$  is prime.

LEMMA 2.46. For all flow graphs A, A is right-prime iff A is left-prime.

*Proof.* Suppose A is right-prime. Assume that  $A \setminus B$  exists. We want to show that  $B = F_1$  or B = A. Since  $A \setminus B$  exists, it follows  $A = B \otimes K$  for some K. Thus A/K exists, and since A is right-prime it follows that  $K = F_1$  or K = A. If K = A then  $B = F_1$ . If  $K = F_1$ , then B = A.  $\square$ 

**2.3. Order Properties.** In this section we explore the relationship between strong ordering by  $\otimes$  and weak ordering by  $\otimes$ . While the two orders coincide on the graphical natural numbers, neither order is anti-symmetric on all of  $\mathcal{F}$ , and only  $\otimes$  is transitive. On the other hand, many of the laws that govern the relationship between  $\leq$ , + and × in  $\mathcal{N}$  continue to hold for  $\otimes$ ,  $\oplus$  and  $\otimes$  in  $\mathcal{F}$ , but these laws are violated under the ordering  $\otimes$ .

LEMMA 2.47 (Strong Order Preservation). For flow graphs A, B, C, if  $A \otimes B$  then

$$(A \otimes C) \otimes (B \otimes C).$$

*Proof.* Let  $A = (G_A, s_A, t_A)$ ,  $B = (G_B, s_B, t_B)$ . Since  $A \otimes B$  there are graph embeddings  $\phi_s : G_A \to G_B$  and  $\phi_t : G_A \to G_B$  which satisfy  $\phi_s(s_A) = s_B$  and  $\phi_t(t_A) = t_B$ . Define  $\gamma_s : E[G_A] \times E[G_C] \to E[G_B] \times E[G_C]$  by

$$(e, f) \mapsto (\phi_s(e), f).$$

Then the composite map

$$\Phi_s^C : E[A \otimes C] \xrightarrow{\Lambda_{A,C}} E[G_A] \times E[G_C] \xrightarrow{\gamma_s} E[G_B] \times E[G_C] \xrightarrow{\Lambda_{B,C}^{-1}} E[B \otimes C]$$

defines an embedding of  $G_A \bigotimes_C \to G_B \bigotimes_C$  which takes  $s_A \bigotimes_C$  to  $s_B \bigotimes_C$ . An analogous construction can be carried out to produce a map  $\Phi_t^C$  which embeds  $G_A \bigotimes_C \to G_B \bigotimes_C$  and sends  $t_A \bigotimes_C$  to  $t_B \bigotimes_C$ .  $\square$ 

LEMMA 2.48 (Strong Order Violations). There exist ST-flow graphs A, B and C for which which

$$(i)(A \oplus C) \otimes (B \oplus C)$$
$$(ii)(C \oplus A) \otimes (C \oplus B)$$
$$(iii)(C \otimes A) \otimes (C \otimes B).$$



FIG. 2.6. Strong order violations: (i).  $(A \oplus C_1) \otimes (B \oplus C_1)$ , (ii).  $(C_2 \oplus A) \otimes (C_2 \oplus B)$ , and (iii).  $(C_3 \otimes A) \otimes (C_3 \otimes B)$ .

*Proof.* See Figure 2.6. □

We consider possible anti-symmetry of S. Suppose A S B and B S A. There is a graph embedding  $\phi_s : G_A \to G_B$  which satisfies  $\phi_s(s_A) = s_B$ . Hence  $|V[G_A]| =$  $|V[\phi_s(G_A)]| \leq |V[G_B]|$  and  $|E[G_A]| = |E[\phi_s(G_A)]| \leq |E[G_B]|$ . Since B S A, there is a graph embedding  $\psi_s : G_B \to G_A$  which satisfies  $\psi_s(s_B) = s_A$ . So  $|V[G_B]| =$  $|V[\psi_s(G_B)]| \leq |V[G_A]|$  and  $|E[G_B]| = |E[\psi_s(G_B)]| \leq |E[G_A]|$ . It follows that  $\phi_s$  is actually an isomorphism from  $G_A$  to  $G_B$  satisfying  $\phi_s(s_A) = s_B$ . A similar argument shows that there is an isomorphism  $\phi_t$  from  $G_A$  to  $G_B$  satisfying  $\phi_t(t_A) = t_B$ . To conclude that A = B requires a single flow graph isomorphism  $\pi$  from A to B, satisfying both  $\pi(s_A) = s_B$  and  $\pi(t_A) = t_B$ . Indeed in some cases, no such isomorphism may exist.

EXAMPLE 2.49. Let  $G_A$  be a directed cycle of length 4, and take  $s_A, t_A$  to be any two vertices in  $V[G_A]$  that are distance 2 apart. Put  $G_B$  isomorphic to  $G_A$ , taking  $s_B, t_B$  to be two vertices in  $V[G_B]$  that are distance 1 apart. Then it is easy to verify that  $(G_A, s_A, t_A) = A \otimes B = (G_B, s_B, t_B)$  and  $B \otimes A$ . Clearly, however,  $A \neq B$  as flow graphs (see Figure 2.7).



FIG. 2.7. An example which demonstrates that the strong order is not antisymmetric.

The previous example proves the next lemma.

LEMMA 2.50 (Non-antisymmetry of strong order  $\bigcirc$ ). There exist flow graphs A and B for which

$$A \otimes B$$
 and  $B \otimes A$  but  $A \neq B$ .

Indeed, since the flow graphs A and B in Example 2.49 are both ST-flow graphs, it is apparent that antisymmetry of strong order  $\bigotimes$  fails even for ST-flow graphs.

LEMMA 2.51 (Transitivity of strong order  $\otimes$ ). For all flow graphs A, B, C

 $A \otimes B$  and  $B \otimes C$  implies  $A \otimes C$ .

Proof.  $A \otimes B$ : i.e. there are graph embeddings  $\phi_s : G_A \to G_B$  and  $\phi_t : G_A \to G_B$ which satisfy  $\phi_s(s_A) = s_B$  and  $\phi_t(t_A) = t_B$ .  $B \otimes C$ : i.e. there are graph embeddings  $\theta_s : G_B \to G_C$  and  $\theta_t : G_B \to G_C$  which satisfy  $\theta_s(s_B) = s_C$  and  $\theta_t(t_B) = t_C$ . We want to show  $A \otimes C$ : i.e. there are graph embeddings  $\alpha_s : G_A \to G_C$  and  $\alpha_t : G_A \to G_C$  which satisfy  $\alpha_s(s_A) = s_C$  and  $\alpha_t(t_A) = t_C$ . Put  $\alpha_s = \theta_s \circ \phi_s$  and  $\alpha_t = \theta_t \circ \phi_t$ .  $\square$ 

LEMMA 2.52 (Weak Order Preservation). For flow graphs  $A,B,C, \ if \ A \bigotimes B$  then

(i)	(A	$\oplus C$ )	$\bigotimes (B$	$\oplus C$
(ii)	(C	$\oplus A)$	$\bigcirc (C$	$\oplus B)$
(iii)	(A	$\otimes C$	$\bigotimes (B$	$\otimes C$ ).

*Proof.* Let  $A = (G_A, s_A, t_A), B = (G_B, s_B, t_B)$  and  $C = (G_C, s_C, t_C)$  be given.  $A \otimes B$  implies that there exists an  $(s_A, t_A)$ -splitting  $(H_1, H_2)$  of  $G_A$  and graph embeddings

$$\phi_1: H_1 \to G_B$$
$$\phi_2: H_2 \to G_B$$

satisfy  $\phi_1(s_A) = s_B$  and  $\phi_2(t_A) = t_B$  and  $\phi_1(E[H_1]) \cap \phi_2(E[H_2]) = \emptyset$ .

(i). Put  $K_1 = H_1$  and define  $K_2$  to be the graph obtained by gluing  $H_2$  and  $G_C$ such that  $t_A$  is identified with  $s_C$ . Now define  $\Phi_1 = \phi_1 : H_1 \to G_B \bigoplus_{t_B \approx s_C} G_C$ ,  $\Phi_2|_{H_2} = \phi_2 : H_2 \to G_B \bigoplus_{t_B \approx s_C} G_C$  and  $\Phi_2|_{G_C} : G_C \to G_B \bigoplus_{t_B \approx s_C} G_C$ . Then  $(K_1, K_2)$  is an  $(s_A, t_C)$ -splitting of  $G_A \bigoplus_{t_A \approx s_C} G_C$  and

$$\Phi_1: K_1 \to G_B \bigoplus_{t_B \approx s_C} G_C,$$
  
$$\Phi_2: K_2 \to G_B \bigoplus_{t_B \approx s_C} G_C$$

are graph embeddings satisfying  $\Phi_1(s_A) = s_B$  and  $\Phi_2(t_C) = t_C$ .

(ii). We define  $L_1$  to be the graph obtained by gluing  $G_C$  and  $H_1$  such that  $t_C$  is identified with  $s_A$  and we put  $L_2 = H_2$ .

$$\begin{aligned} \theta_1|_{G_C} &: G_C \to G_C \bigoplus {}_{t_C \approx s_B} G_B \\ \theta_1|_{H_1} &: H_1 \to G_C \bigoplus {}_{t_C \approx s_B} G_B \\ \theta_2 &= \phi_2 : H_2 \to G_C \bigoplus {}_{t_C \approx s_B} G_B. \end{aligned}$$

Then  $(L_1, L_2)$  is an  $(s_C, t_A)$ -splitting of  $G_C \bigoplus_{t_C \approx s_A} G_A$  and

$$\theta_1: L_1 \to G_C \bigoplus_{t_C \approx s_B} G_B, \\ \theta_2: L_2 \to G_C \bigoplus_{t_C \approx s_B} G_B$$

26

are graph embeddings satisfying  $\theta_1(s_C) = s_C$  and  $\theta_2(t_A) = t_B$ .

(iii). Put

$$M_1 = H_1 \otimes C$$
$$M_2 = H_2 \otimes C.$$

Since  $E[H_1] \cup E[H_2] = E[G_A]$  and  $E[H_1]) \cap E[H_2] = \emptyset$ , it follows that  $E[H_1 \otimes C] \cup E[H_2 \otimes C] = E[G_A \otimes C]$  and  $E[H_1 \otimes C] \cap E[H_2 \otimes C] = \emptyset$ . Thus  $M_1, M_2$  are an (s, t)-splitting of  $G_A \otimes C$ . Now take  $\beta_1 : E[M_1] \to E[G_B \otimes G_C]$  and  $\beta_2 : E[M_2] \to E[G_B \otimes G_C]$  defined by

$$\beta_1 : (e, f) \mapsto (\phi_1 e, f)$$
  
$$\beta_2 : (e', f) \mapsto (\phi_2 e', f)$$

for  $e \in E[H_1]$ ,  $e' \in E[H_2]$  and  $f \in C$ . The injectivity of  $\beta_1$  and  $\beta_2$  follows immediately from injectivity of  $\phi_1$  and  $\phi_2$ . Since  $\phi_1(E[H_1]) \cap \phi_2(E[H_2]) = \emptyset$ , it follows that  $\beta_1(E[H_1 \otimes C]) \cap \beta_2(E[H_2 \otimes C]) = \emptyset$ . Since  $\phi_1(s_A) = s_B$  and  $\phi_2(t_A) = t_B$ , it follows that  $\beta_1(s_A \otimes C) = s_B \otimes C$  and  $\beta_2(t_A \otimes C) = t_B \otimes C$ . Thus, the maps  $\beta_1$ and  $\beta_2$  demonstrate  $A \otimes C \otimes B \otimes C$ .  $\square$ 

LEMMA 2.53 (Weak Order Violations). There exist flow graphs A, B, C for which  $A \otimes B$  but

$$(C \otimes A) \otimes (C \otimes B).$$

*Proof.* See Figure 2.8. □



FIG. 2.8. An example which demonstrates weak order violation:  $(C \otimes A) \otimes (C \otimes B)$ .

LEMMA 2.54 (Non-transitive weak order  $\otimes$ ). There exist flow graphs A, B, C

 $A \otimes B$  and  $B \otimes C$  but  $A \otimes C$ .

*Proof.* See Figure 2.9. □



FIG. 2.9. An example which demonstrates that the weak order is not transitive.

LEMMA 2.55 (Non-antisymmetry of weak order  $\,\,\textcircled{}_{>}$  ). There exist flow graphs A and B for which

$$B \otimes A \text{ and } A \otimes B \text{ but } A \neq B$$

*Proof.* Since strong order implies weak order, Lemma 2.50 and the example in Figure 2.7 immediately yield:  $\Box$ 

Indeed, since the flow graphs A and B in Example 2.49 are both ST-flow graphs, it is apparent that antisymmetry of weak order fails even for ST-flow graphs.

3. Conclusions and Future Work. As we have seen, strikingly many theorems that are true in  $\mathcal{N}$  continue to hold in  $\mathcal{F}$ , though some fail. Our future research program will proceed on two tracks.

Informally, for each "classical" theorem  $\phi$  in  $TA \setminus Th(\mathcal{F})$ :

- (1) We shall consider the structure of maximal subsets  $X_{\phi}$  which have the property that the submodel  $\mathcal{X}_{\phi} \stackrel{def}{=} (\mathcal{F}|_{X_{\phi}}) \models \phi$ . Of particular interest are sets  $X_{\phi}$  which properly contain  $i(\mathbb{N})$ .
- (2) We shall describe a corresponding theorem  $\phi'$  in  $Th(\mathcal{F})$ , such that  $\phi' \equiv \phi$  when restricted to  $i(\mathbb{N})$ .

## Examples of specific questions include:

- i. Characterize flow graph pairs for which antisymmetry of strong order holds.
- ii. Characterize  $\otimes$  -commuting pairs, i.e. under what conditions on flow graphs A and B does  $A \otimes B = B \otimes A$ ?
- iii. Does  $A \otimes B = A \otimes C$  imply B = C? Does  $B \otimes A = C \otimes A$  imply B = C? In other words, does  $\otimes$  satisfy a left/right cancellation law?
- iv. *Graph Prime Factorization Conjecture*. Every flow graph is uniquely expressible (up to some well-defined reordering) as the product of prime flow graphs.
- v. Describe solution sets (in  $\mathcal{F}$ ) for one-variable equations having the form p(x) = q(x), where p and q are polynomials with coefficients from  $\mathcal{F}$ .

Acknowledgements. The authors would like to thank Martin Kassabov for his incisive reading of this paper, insightful comments, and for bringing to our attention that the Crossing Summands Lemma used in the proof of Proposition 2.18) fails to hold for general flow graphs. The example demonstrating this (in Figure 2.2) is also due to him.

The first author would like to thank the Center for Computational Science at the Naval Research Laboratory, Washington DC and ITT Industries for supporting ongoing research efforts in mathematics and computer science.

### REFERENCES

- [1] K. R. BHUTANI AND B. KHAN, A metric representing dilation between connected simple graphs of a given order. Submitted to Discrete Applied Mathematics.
- [2] —, Distance between graphs using graph labelings, Ars Combinatoria, (2003). Accepted for publication.
- [3] , A metric on the class of connected simple graphs of given order, Aequationes Mathematicae, 66 (2003), pp. 232–240.
- W. IMRICH AND S. KLAVZAR, Product Graphs, Structure and Recognition, John Wiley & Sons Inc., 2000.
- [5] D. B. WEST, Introduction to Graph Theory, Prentice Hall, 2nd ed., 2001.

(s, t)-splitting, 7  $F^{ST}, 12$ (1), 5 $\oplus$ associativity, 10 commutativity condition, 18 identity, 3 irreducibile, 11 Irreducibility Lemma, 12 left-cancellation law, 18 non-commutativity, 11 right-cancellation law, 18  $\oplus$  -decomposition, 16  $\oplus$  -irreducibility for products, 24  $\otimes$ associativity, 20 identity, 5 non-commutativity, 21 not left-distributivity over  $\oplus$ , 22 right-distributivity over  $\oplus$ , 21 (0), 3s standard, 11 t standard, 11 ST Property, 12 ST-Core, 13 st-flow graph, 12 antisymmetry strong order does not satisfy, 25 weak order does not satisfy, 28 canonical  $\oplus$  -decomposition, 16 for products, 23 for sums, 16 length for products, 23 component-wise decomposition of isomorphisms, 14 decomposition  $\oplus$ , 16 edge splitting — for a flow graph, 23finite flow graph, 2 flow graph, 2 ST Property, 12 addition, 3

edge splitting for a -, 23 finite, 2 infinitesimal, 2 isomorphism, 2 multiplication, 5 prime, 24 scalar exponentiation, 7 scalar multiplication, 4 splitting, 11 vertex splitting for a —, 11 Graph flow, 2 Graphical natural number, 2 infinitesimal flow graph, 2 Irreducibility Lemma  $\oplus$ , 12 irreducible  $\oplus$ , 11 isomorphism flow graph, 2 isomorphisms component-wise, 14 left-cancellation law for  $\oplus$ , 18 Natural number graphical, 2 order strong, 8 weak, 7 prime flow graph, 24 right-cancellation law for  $\oplus$ , 18 Scalar exponentiation of flow graphs, 7 Scalar multiplication of flow graphs, 4 splitting (s,t), 7edges and  $\oplus$  -irreducibles, 23 vertex ranking, 15 edge for a flow graph, 23 flow graph, 11 vertex for a flow graph, 11 standard

 $\begin{array}{c} s,t,\,11\\ {\rm strong\ order,\ 8}\\ {\rm non-antisymmetry,\ 25}\\ {\rm preservation,\ 24}\\ {\rm transitivity,\ 26}\\ {\rm violation,\ 24} \end{array}$ 

vertex

splitting—for a flow graph, 11

weak order, 7 non-antisymmetry, 28 non-transitivity, 28 preservation, 26

violation, 27