

# The Ending Laminations Theorem direct from Teichmüller Geodesics

Mary Rees

August 3, 2018

## Abstract

A proof of the Ending Laminations Theorem is given which uses Teichmüller geodesics directly.

## 1 Introduction

Over 30 years ago, Scott proved the remarkable result [52], [53] that any finitely-generated fundamental group of any three-manifold  $N$  is finitely presented, and, further, that the manifold is homotopy equivalent to a compact three-dimensional submanifold with boundary. This submanifold, known as the *Scott core*, was shown [31] to be unique up to homeomorphism — but not, in general, up to isotopy. If, in addition,  $N$  is a hyperbolic manifold, then it is useful to consider the manifold with boundary  $N_d$  obtained by removing from  $N$  components of the  $\varepsilon_0$ -thin part round cusps, for a fixed Margulis constant  $\varepsilon_0$ . Sullivan [56] proved that there are only finitely many cusps. We shall call  $N_d$  *the horoball deletion of  $N$* . Topologically, it does not depend on the choice of Margulis constant  $\varepsilon_0$ , although metrically it does. There is a relative version of Scott's result [29] which says that  $(N_d, \partial N_d)$  is homotopy equivalent to  $(N_c, N_c \cap \partial N_d)$  where  $N_c \subset N_d$  is the *relative Scott core*. The relative Scott core is, again, unique up to homeomorphism [30]. The closure of each component of  $\partial N_c \setminus \partial N_d$  is an orientable compact surface  $S_d$  with boundary, and is also the boundary in  $N_d$  of  $U(S_d)$ , for a unique component  $U(S_d)$  of  $N_d \setminus N_c$ . The surface  $S_d$  can be any compact orientable surface with boundary apart from the closed disc, annulus or torus. The set  $U(S_d) = U(e)$  is a neighbourhood of a unique end  $e$  of  $N_d$  [5]. The correspondence between end  $e$ , and the surface  $S_d(e)$  bounding its neighbourhood, is thus one-to-one. We shall also write  $S(e)$  for the surface without boundary obtained up to homeomorphism by attaching a punctured disc to each boundary component of  $S_d(e)$ .

In 2004, following some close approaches (e.g. [55]), two proofs of the Tame-ness conjecture were announced [2], [14], that  $N_c$  can be chosen so that each closed end neighbourhood  $S_d(e) \cup U(e)$  is homeomorphic to  $S_d(e) \times [0, \infty)$ , that is, each end of  $N_d$  is *topologically tame*. (See also [54].) The topology of  $N$  is thus uniquely determined by the topology of the pair  $(N_c, N_c \cap \partial N_d)$ .

The history of the geometry of  $N$  runs parallel to this. The concept of geometric tameness was originally developed by Thurston [58]. An end of  $N_d$  is *geometrically finite* if it has a neighbourhood disjoint from the set of closed geodesics in  $N$ . (There are a number of equivalent definitions.) An end of  $N_d$  is *simply degenerate* if each neighbourhood of the end contains a *simple* closed geodesic. These two possibilities are mutually exclusive, and if one or the other holds, the end is said to be *geometrically tame*. It is not clear a priori that one or the other possibilities must hold, but in 1986 Bonahon [5] published a proof that an end  $e$  of  $N_d$  is geometrically tame if the inclusion of  $S_d(e)$  in  $N_c$  is injective on fundamental groups, that is, if  $S_d(e)$  is *incompressible* in  $N_d$ . Importantly for future developments, this made rigorous the end invariant suggested by Thurston. In the geometrically finite case, if  $N = H^3/\pi_1(N)$ , and  $\Omega \subset \partial H^3$  is the domain of discontinuity of  $\pi_1(N)$ , then the closure of  $U(e)$  in  $(H^3 \cup \Omega)/\pi_1(N)$  intersects a unique punctured hyperbolic surface homeomorphic to  $S(e)$ , and contains a subsurface which is isotopic in  $(H^3 \cup \Omega)/\pi_1(N)$  to  $S_d(e)$ . Thus, this surface in  $\Omega/\pi_1(N)$  determines a point  $\mu(e)$  in the Teichmüller space  $\mathcal{T}(S(e))$ . Bonahon's work dealt with the simply degenerate ends. He showed that if  $S(e)$  was endowed with any complete hyperbolic structure, then the Hausdorff limit of any sequence of simple closed geodesics exiting  $e$  was the same arational geodesic lamination, independent, up to homeomorphism, of the hyperbolic metric chosen. A *geodesic lamination* on  $S$  is a closed set of nonintersecting geodesics on  $S$ , and a lamination is *arational* if every simple closed loop intersects a recurrent leaf in the lamination transversally. So in both cases, geometrically finite and infinite, an invariant  $\mu(e)$  is obtained, and if there are  $n$  ends  $e_i$ ,  $1 \leq i \leq n$ , this gives an invariant  $(\mu(e_1), \dots, \mu(e_n))$ . Bonahon also showed that geometric tameness implies topological tameness. Later, Canary [16] extended Bonahon's result about geometric tameness to any end of any topologically tame  $N$ . The resolutions of the Tameness Conjecture mentioned above means that any end of any three-dimensional hyperbolic manifold with finitely generated fundamental group is topologically tame, and hence, by the work of Bonahon and Canary, geometrically tame.

Bonahon's result on an invariant  $(\mu(e_1), \dots, \mu(e_i))$  associated to  $N$  for which  $\partial N_c \setminus \partial N_d$  is incompressible, generalises to any tame hyperbolic manifold  $N$ . Let  $G$  denote the group of orientation-preserving homeomorphisms  $\varphi$  of  $N_c$ , modulo isotopy, and are homotopic in  $N_c$  to the identity. Then  $G$  acts on  $\prod_e \mathcal{T}(S(e)) \cup \mathcal{GL}(S(e))$ , where the product is over ends  $e$  of  $N_d$  and  $\mathcal{GL}(S(e))$  is the space of geodesic laminations on  $S(e)$ . We write  $[\mu_1, \dots, \mu_n]$  to denote an element of the quotient space. Note that  $G$  is trivial if each  $S_d(e)$  is incompressible in  $N_d$ . We also write  $\mathcal{GL}_a(S)$  for the space of arational laminations in  $\mathcal{GL}(S)$ . The ending invariant of a tame hyperbolic manifold is now a point

$$[\mu(e_1), \dots, \mu(e_n)] \in \prod_{i=1}^n (\mathcal{T}(S(e_i)) \cup \mathcal{O}_a(S(e_i), N))/G$$

Here,  $\mathcal{O}(S, N) \subset \mathcal{GL}(S)$  is the *Masur domain* as defined by Otal [45], since the concept arose in the case of handlebodies in work by H. Masur [26]. The precise

definition will be given later, but  $\mathcal{O}(S, N)$  is open in  $\mathcal{GL}(S)$ , invariant under the action of the group  $G$  (as above), contains no closed geodesics in  $S$  which are trivial in  $N$ , and  $\mathcal{O}(S, N) = \mathcal{GL}(S)$  if  $(S_d, \partial S_d)$  is incompressible in  $(N_d, \partial N_d)$ . As for  $\mathcal{GL}(S)$ ,  $\mathcal{O}_a(S, N)$  denotes the set of arational laminations in  $\mathcal{O}(S, N)$ .

The obvious questions to ask about the invariants  $[\mu(e_1), \dots, \mu(e_n)]$  of hyperbolic manifolds of a fixed topological type are: does the invariant  $[\mu(e_1), \dots, \mu(e_n)]$  of a hyperbolic manifold uniquely determine that manifold up to isometry, and which invariants can occur.

There is a natural compactification of  $\mathcal{T}(S)$ , or, more generally, of  $\prod_e \mathcal{T}(S(e))/G$ , in which  $\prod_e \mathcal{T}(S(e))/G$  is the interior and  $\prod_e \mathcal{O}_a(S(e), N)/G$  is contained in the boundary. The topology at the boundary will be specified later. The boundary is actually larger than  $\prod_e \mathcal{O}_a(S(e), N)/G$ . It also includes countably many *split boundary pieces*, one for each multicurve  $\Gamma$  in the Masur domain. (Multicurves are defined in 2.1, and the Masur domain in 3.10.) Given a multicurve  $\Gamma_i$  on  $S(e_i)$  for each end  $e_i$  of  $N$ , we define a topological manifold with boundary  $N_d(\Gamma_1, \dots, \Gamma_n)$  which contains  $N_c$  and is contained in  $N$ , as follows. We can assume that  $\gamma \subset S_d(e_i)$  for all  $\gamma \in \Gamma_i$ . Fix a homeomorphism  $\Phi_i$  from  $S(e_i) \times [0, \infty)$  to the closed neighbourhood of  $e_i$  in  $N$  bounded by  $S(e_i)$ . Choose a collection of disjoint open annulus neighbourhoods  $A(\gamma) \subset S_d(e_i)$  of the loops  $\gamma \in \Gamma_i$ . Then

$$N_d(\Gamma_1, \dots, \Gamma_n) = N_d \setminus \bigcup_{i=1}^n \Phi_i(\bigcup_{\gamma \in \Gamma_i} A(\gamma) \times (0, \infty)),$$

$$N_c(\Gamma_1, \dots, \Gamma_n) = N_c \cap N_d(\Gamma_1, \dots, \Gamma_n).$$

The closures of components of  $\partial N_c(\Gamma_1, \dots, \Gamma_n) \setminus \partial N_d(\Gamma_1, \dots, \Gamma_n)$  are sets  $S_d(e_i, \alpha)$ , with interior homeomorphic to  $\alpha$ , for each component  $\alpha$  of  $S(e_i) \setminus (\bigcup \Gamma_i)$ . We write  $\Sigma(\Gamma_i)$  for the set of such components. We let  $S(e_i, \alpha)$  be the topological surface obtained by adding a punctured disc round each boundary component. Write

$$\mathcal{W}(S, \Gamma) = \prod_{\alpha \in \Sigma(\Gamma)} (\mathcal{T}(S(\alpha)) \cup \mathcal{O}_a(S(\alpha), N)).$$

The boundary of  $\prod_e \mathcal{T}(S(e))/G$  consists of

$$\prod_e (\mathcal{O}_a(S(e), N) \cup \bigcup_{\Gamma} \mathcal{W}(S(e), \Gamma))/G,$$

where the union is over all nonempty multicurves  $\Gamma$  as above. We continue to denote an element of this space by  $\mu$ , so that points in  $\mathcal{T}(S)/G$  and its boundary are denoted by  $\mu$ .

The Ending Laminations Theorem can then be formulated as follows, given the proofs of the Tameness conjecture. It says that the ending lamination invariants are unique. This is the first main result of this paper.

**Theorem 1.1.** *Let  $N$  be any three-dimensional hyperbolic manifold with finitely generated fundamental group such that  $N_d$  has ends  $e_i$ ,  $1 \leq i \leq n$ . Then  $N$  is uniquely determined up to isometry by its topological type and the ending lamination data  $[\mu(e_1), \dots, \mu(e_n)]$ .*

As for existence, we have the following, which has extensive overlap with earlier results in the literature.

**Theorem 1.2.** *Let  $N$  be any geometrically finite three-dimensional hyperbolic manifold with finitely generated fundamental group such that the horoball deletion  $N_d$  has ends  $e_i$ ,  $1 \leq i \leq n$ . Let  $N(\mu'_1, \dots, \mu'_n)$  be the manifold in the quasi-isometric deformations space  $Q(N)$  with ending data  $[\mu'_1, \dots, \mu'_n] \in \prod_i \mathcal{T}(S(e_i))/G$ . Then the map*

$$[\mu'_1, \dots, \mu'_n] \rightarrow N([\mu'_1, \dots, \mu'_n])$$

*extends continuously, with respect to both algebraic and geometric convergence, to map any point  $[\mu_1, \dots, \mu_n]$  of the boundary*

$$\left( \prod_{i=1}^n \mathcal{O}_a(S(e_i), N) \cup \cup_{\Gamma_i} \mathcal{W}(S(e_i), \Gamma_i) \right) / G$$

*to a hyperbolic manifold  $N([\mu_1, \dots, \mu_n]) = N'$ , with  $(N'_c, \partial N'_c \setminus \partial N'_d)$  homeomorphic to  $(N_c, S_d(\Gamma_1, \dots, \Gamma_n))$ , if  $\mu_i \in \mathcal{W}(S(e_i), \Gamma_i)$  for  $1 \leq i \leq n$  except in the following cases.*

1.  $(N_c, \partial N_d \cap N_c) = (S_d \times [0, 1], \partial S_d \times [0, 1])$  up to homeomorphism, so that, if  $n = 2$ , we identify  $S \times \{j\}$  with  $S$  under  $(x, j) \mapsto x$ , and  $(\mu_1, \mu_2) = (\mu, \mu)$  for some  $\mu \in \mathcal{GL}(S)$
2. For some  $i \neq j$ , and nonempty  $\Gamma_i, \Gamma_j$ , on  $S(e_i), S(e_j)$ ,  $\mu_i \in \mathcal{W}(S, -)$ ,  $\mu_j \in \mathcal{W}(S, \Gamma_j)$  and some loops  $\gamma \in \Gamma_i$  and  $\gamma' \in \Gamma_j$  are isotopic in  $N$ .

This overlaps with a substantial and longstanding literature of existence results, dating back to Thurston's proof of algebraic convergence of subsequences in the deformation space with converging ending laminations data, initially for acylindrical manifolds in [59]. The deformation space is the space of discrete faithful representations of  $\pi_1(N)$  in  $\text{Isom}(H^3)$ , modulo conjugation in  $\text{Isom}(H^3)$ , that is, the topology of algebraic convergence. Other examples are [42], [43], and, recently, [24], [44]. This well-developed approach, which often uses a compactification of the deformation space by a space of  $\mathbb{R}$ -trees, is by-passed in 1.2. Another approach, for Masur domain laminations for handlebodies, is given in Namazi's thesis [40] in collaboration with Souto, using the Brock-Canary-Minsky Ending Lamination Theorems of [10].

Thurston's Geometrization Theorem ([46], [39], [32]) says that any compact three-dimensional *pared* manifold  $(M, P)$  with boundary is homeomorphic to  $(H^3 \cup \Omega) / \Delta \setminus P', \partial P'$  for some discrete geometrically finite group of isometries  $\Delta \cong \pi_1(M)$  with domain of discontinuity  $\Omega$  and horoball neighbourhoods of cusps  $P'$ . Let  $N = H^3 / \Delta$  for such a  $\Delta$ . It is then classical, a consequence of the Measurable Riemann-Mapping Theorem [3], that the space  $Q(N)$  of manifolds quasi-isometric to  $N$ , using the topology of algebraic convergence, is homeomorphic to  $\prod_{i=1}^n \mathcal{T}(S(e_i))$ , where  $e_i$  ( $1 \leq i \leq n$ ) are the ends of  $N$ . The map from  $\prod_{i=1}^n \mathcal{T}(S(e_i))$  to  $Q(N)$  is also continuous with respect to the geometric

topology on  $Q(N)$ . Theorem 1.2 then implies that  $Q(N)$  is dense, in the algebraic topology, in the space of hyperbolic manifolds of the topological type of  $N$ . This, then, suggests an alternative proof of the Bers-Sullivan-Thurston Density Conjecture, that is, density of geometrically finite groups. This was proved by Bromberg [11] and Brock-Bromberg [9] in the case of incompressible boundary and without parabolic elements, derived the Ending Laminations Theorem and work of Thurston and Ohshika ([59], [60], [42], [43], [44]) by Brock-Canary-Minsky [10] in their proof of the Ending Laminations Theorem, and can apparently be derived from the work of In Kang Kim, Lecuire and Ohshika ([24], [44]) and the Brock-Canary-Minsky proof of the full version of the Ending Laminations Theorem. Another proof is in preparation by Bromberg and Souto [12]

A proof of 1.1, for both geometrically finite and infinite ends was announced in the Kleinian surface case — when  $N_c$  is homeomorphic to  $S_d \times [0, 1]$ , possibly with more than two ends — in 2002 by Y. Minsky, J. Brock and D. Canary, the first part being due to Minsky alone [38]. They also announced the result in general the following year. Their full proof in the Kleinian surface case [10] became publicly available at the end of 2004, and includes a number of other results including a result related to 1.2. The general case is in preparation.

The Brock-Canary-Minsky proof of the Ending Laminations Conjecture is the culmination of a number of papers of Minsky in which the Ending Laminations Theorem was successively proved in important special cases, especially the Kleinian surface once-punctured torus case [36] and the Kleinian surface bounded geometry case [37], [33]. The proof of the once-punctured torus case was striking because of the strategy of proof, which was then used in other cases of the result. The Teichmüller space  $\mathcal{T}(T)$  of the once-punctured torus  $T$  identifies with the unit disc  $\{z : |z| < 1\}$ , and the set of geodesic laminations with the boundary  $\{z : |z| = 1\}$ . So the ending laminations data identifies an element  $(\mu_+(N), \mu_-(N))$  of  $\{(z, w) : |z|, |w| \leq 1\} \setminus \{(z, z) : |z| = 1\}$ , using a slightly non-standard identification between the two components of the relative Scott core. Minsky constructs a geometric model for a Kleinian surface hyperbolic manifold  $N$  homeomorphic to  $T \times \mathbb{R}$  with given end invariants in terms of the paths in the Farey graph to  $z, w$ , at least in the case when  $|z| = |w| = 1$ . He was then able to show that the geometric model manifold was biLipschitz equivalent to  $N$ , in fact boundedly so, with bounds independent of  $(\mu_+(N), \mu_-(N))$ . From this, the fact that  $N$  was the unique manifold in this homeomorphism class with invariant  $(\mu_+(N), \mu_-(N))$  was deduced.

The bounded geometry case of the Ending Laminations Theorem has an interesting history, [37] being a return to the problem of *bounded geometry ending invariants* (as it is reasonable to call them) 8 years after the results of [33], when the Ending Laminations Theorem was proved for hyperbolic manifolds with bounded geometry, which is a stronger assumption than bounded geometry of the end invariants. Some of the techniques of [37] have wider application, and they are of fundamental importance to the current work. The other big new input to [37] was the theory of the curve complex developed by Minsky and H. Masur ([27], [28]) which is replaced by a different theory in the current

work. The general strategy which Minsky developed in the punctured torus and bounded geometry cases was then used in his resolution, with Brock and Canary for the last part, for the proof of the general case. In summary, the Brock-Canary-Minsky proof of the Ending Laminations Theorem can be considered to consist of three stages:

1. the construction of a geometric model  $M(\mu_1, \dots, \mu_{mn})$  up to quasi-isometry, for given ending lamination data  $(\mu_1, \dots, \mu_n)$  and fixed topological type;
2. given any hyperbolic manifold  $N$  with ending laminations the construction of a map from the geometric model  $M(\mu_1, \dots, \mu_n)$  to  $N$  which is Lipschitz;
3. a proof (with Brock and Canary) that this map is, in fact, biLipschitz.

Minsky's construction of the geometric model and the Lipschitz map, and the final proof of biLipschitz, all use the curve complex and deep and extensive work of Masur and Minsky ([27], [28]) on hierarchies of tight geodesics in the curve complex. One purpose of the current paper is to show that it is possible to carry out the programme working directly with Teichmüller geodesics. The general strategy is that developed by Minsky, but the detail is quite different. The work is built on two planks. One, as already mentioned, is some results of [33] about pleated surfaces, which apply in a wider context than used there. The other is results about Teichmüller geodesics which were developed in [50] for quite another purpose. These Teichmüller geodesic results are used both to define a geometric model  $M = M(\mu_1, \mu_2 \dots \mu_n)$  up to Lipschitz equivalence for a hyperbolic manifold in a given homeomorphism class with end invariant  $(\mu_1, \dots, \mu_n)$ , and to construct a Lipschitz map from the model manifold to a hyperbolic manifold with this ending data, and to show that this map is coarse biLipschitz.

We say that a map  $\Phi : M \rightarrow N$  between complete Riemannian manifolds  $M$  and  $N$  is *coarse biLipschitz* if  $d_1, d_2$  are the lifted metrics on the universal covers  $\tilde{M}, \tilde{N}$  of  $M, N$  and there is  $K$  such that for all  $x, y \in \tilde{M}$ ,

$$K^{-1}d_1(x, y) - K \leq d_2(\tilde{\Phi}(x), \tilde{\Phi}(y)) \leq Kd_1(x, y) + K,$$

where  $\tilde{\Phi}$  is any fixed lift of  $\Phi$ .

We refer to data in the compactification of  $\prod_i \mathcal{T}(S(e_i))$  as *excluded* if it is as in cases 1 or 2 of 1.2 and otherwise it is *permissible*. The exclusion of 2 of 1.2 is trivially necessary, if one examines the condition, because if  $\gamma \in \Gamma_i$  and  $\gamma' \in \Gamma_j$  are isotopic in  $N = N(\mu_1, \dots, \mu_n)$  then they must represent the same parabolic element. It may be that hyperbolic manifolds in different homeomorphism classes are in the boundary of  $Q(N)$  but we do not pursue this. The exclusion of 1 of 1.2 is certainly necessary, but this is nontrivial, being tied up with the deepest arguments in [5]. The theorem which will imply both 1.1 and 1.2 is as follows.

**Theorem 1.3.** *Let  $N_0$  be a complete hyperbolic 3-manifold with finitely generated fundamental group with horoball deletion  $N_{0,d}$  and ends  $e_i$  of  $N_{0,d}$ ,  $1 \leq i \leq n$ .*

*Then for any quotiented  $n$ -tuple of permissible invariants for the ends  $e_i$ ,  $[\mu_1, \dots, \mu_n]$ , there is a manifold with boundary  $M = M([\mu_1, \dots, \mu_n])$ , with a base-*

point  $x_0(M)$  with interior homeomorphic to  $N_0$  under a map  $\Psi_M : M \rightarrow N_0$  and a Riemannian metric  $\sigma_M$  with the following properties.

1. There is a constant  $K_1$  which depends only on the topological type, such that the following holds. Any geometric limit of the structures  $(M', x_0(M'))$ , for  $M' = M([\mu'_1, \dots, \mu'_n])$  converges to  $(M, x_0(M))$  up to  $K_1$ -Lipschitz equivalence, with an isometry between any boundary components, as  $(\mu'_1, \dots, \mu'_n) \rightarrow (\mu_1, \dots, \mu_n)$ , for  $M = M(\mu_1, \dots, \mu_n)$ , and geometrically finite invariants  $[\mu'_1, \dots, \mu'_n]$ , and under a limit of maps homotopic to the maps  $\Psi_M^{-1} \circ \Psi_{M'}$ .
2. There is a constant  $K_2 = K_2([\mu_1, \dots, \mu_n])$  for any permissible  $[\mu_1, \dots, \mu_n]$  which varies continuously with  $[\mu_1, \dots, \mu_n]$ , such that the following holds. Let  $N$  be any hyperbolic manifold with  $N_d$  homeomorphic to  $N_{0,d}$ . Let  $\overline{N}$  denote the closure of  $N$  in  $(H^3 \cup \Omega(\Gamma))/\Gamma$ , where  $M = H^3/\Gamma$  and  $\Omega(\Gamma)$  is the domain of discontinuity. Let  $CH(N)$  denote the convex hull of  $N$ . Then there is a map  $\Phi : M(\mu_1, \dots, \mu_n) \rightarrow \overline{N}$  which is  $K_2$ -coarse-biLipshitz with respect to the hyperbolic metric on  $CH(N)$  on the preimage of  $CH(N)$ , a homeomorphism on the preimage of any component of  $\overline{N} \setminus CH(N)$  and an isometry between any boundary components, using the Poincaré metric on  $\Omega(\Gamma)/\Gamma$ .  
If  $N_c$  has incompressible boundary, then  $K_2$  can be chosen independent of  $(\mu_1, \dots, \mu_n)$ .

A version of this for Kleinian surface groups (or part 2, at least) occurs in [10]. It is stated there that a version holds in the general case, but with  $K_2$  depending on  $N$  in the case of compressible boundary.

Theorem 1.3 will imply that, given hyperbolic manifolds  $N_1$  and  $N_2$  with the same ending data  $[\mu_1, \dots, \mu_n]$ , and maps  $\Phi_1$  and  $\Phi_2$  as in 1.3, the set-valued map  $\tilde{\Phi}_2 \circ \tilde{\Phi}_1^{-1}$  extends to a quasiconformal map of  $\partial H^3$ , invariant with respect to the actions of the covering groups  $\pi_1(N_1)$  and  $\pi_1(N_2)$  on  $\partial H^3$  which is conformal on the (possibly empty) domain of discontinuity. This is fairly standard in the case of no geometrically finite ends. There is slightly more to do if there are geometrically finite ends. Details will be given later. Such a quasiconformal map must be conformal by Sullivan's result [57] that there are no  $\pi_1(N_1)$ -invariant line fields on  $\partial H^3$ , and hence  $N_1$  and  $N_2$  are isometric. So Theorem 1.1 follows directly from 1.3. The deduction of 1.2 from 1.3 is almost as direct, once we have defined the topology on the compactification of  $\mathcal{T}(S)$  and the model manifolds.

The paper proceeds as follows.

**Section 2.** Teichmüller space.

**Section 3.** Pleated surfaces and geodesic laminations.

**Section 4.** More on pleated surfaces.

**Section 5.** Teichmüller geodesics: long thick and dominant definitions.

**Section 6.** Long thick and dominant ideas.

**Section 7.** Geometric model manifolds.

**Section 8.** Model-adapted families of pleated surfaces.

**Section 9.** Proof of 1.3 in the combinatorial bounded geometry Kleinian surface case.

**Section 10.** Lipschitz bounds.

**Section 11.** BiLipschitz bounds.

**Section 12.** Proofs of Theorems 1.1, 1.2, 1.3.

Sections 2, 5 and 6 are concerned with the Teichmüller space  $\mathcal{T}(S)$  of a finite type surface  $S$ , with no reference to 3-dimensional hyperbolic geometry. In Section 3, pleated surfaces are introduced, Bonahon's work [5] recalled and its use to define the ending laminations, and the work of Canary in the general tame case. Minsky's rather astonishing result [37], that there is a bounded homotopy between pleated surfaces in  $N$  whose pleating loci are related by an elementary move, at least in the thick part of  $N$ , is discussed, reinterpreted and extended in Section 4. This extension of Minsky's pleated surface result is one of two main inputs into the current paper. The other main input is the theory of Teichmüller geodesics from [50], of which relevant features are described in Section 5 and 6. The most important result (although not the most difficult) is probably 5.5, which states how to decompose  $\ell \times S$ , for any geodesic segment  $\ell$  in the Teichmüller space  $\mathcal{T}(S)$ , into disjoint product sets which are either *bounded* or *long, thick and dominant* (ltd). Definitions are given in Section 5. For the moment, suffice it to say that this decomposition is vital to all three parts of the strategy. From Section 7 onwards we work explicitly towards proving the main theorems. I am following a fairly direct suggestion of the second reader of an earlier version in the Kleinian surface case, and shall highlight the proof of the Kleinian surface combinatorial bounded geometry case at each stage of the proof. Some specialisation to the case of combinatorial bounded geometry occurs even in Section 6. Subsections are devoted to this case in sections 7 and 8 and as the title of the section given above indicates, section 9 proves Theorem 1.3 in this case. Section 7 goes into considerable detail about the model manifolds, especially about the model of the Scott core. The geometric convergence of model manifolds — the first part of Theorem 1.3 — is proved at the end of Section 7. Section 8 includes estimates on the geometry of pleated surfaces given certain purely combinatorial information about their pleating loci. This is used to obtain information about the geometry of the Scott core. In the case of incompressible boundary, the geometry of the non-interval-bundle part of the Scott core is eventually shown to be independent of the ending laminations data, up to bounded Lipschitz equivalence. As their titles indicate, explicit work on proving the main theorems is carried out in Sections 10 to 12, but a lot of groundwork is done before this.



Other approaches to the Ending Laminations Theorem are available, or are in preparation. One, by Bowditch, has been emerging over the last few years ([6], [7], [8]). A “prehistoric approach” is in preparation by Bromberg and Souto (to which [12] is relevant), with contributions from Evans, including his paper [19], and Brock.

The idea of the approach in this paper emerged during the course of other work. I first contacted Minsky directly about it in 2001. I thank Yair Minsky for some helpful discussion of these ideas over the last five years. I also thank his collaborators, and, in particular, Dick Canary, for his generosity as organiser/secretary of the Ahlfors Bers Colloquium in 2005. I am indebted to the referee and second reader of the earlier paper on the Kleinian surface case for their careful reading and (extensive) detailed and useful criticisms, and the editors of the Newton Institute Proceedings (Kleinian Groups 2003) for their fair and tactful handling of the earlier submission. In particular, I should also like to thank Caroline Series for her very helpful and pertinent questioning, and for facilitating discussion of this work. I also thank Marc Lackenby in this respect, and Brian Bowditch for recent comments. I thank Kasra Rafi — and also Misha Kapovich — for their contributions, and for fruitful interchange on further developments. This is not the end of the story.

## 2 Teichmüller space.

### 2.1 Very basic objects in surfaces

Unless otherwise stated, in this work,  $S$  always denotes an oriented finite type surface without boundary, that is, obtained from a compact oriented surface by removing finitely many points, called *punctures*. One does not of course need an explicit realisation of  $S$  as a compact minus finitely many points. One can define a puncture simply to be an end of  $S$ . A *multicurve*  $\Gamma$  on  $S$  is a union of simple close nontrivial nonperipheral loops on  $S$ , which are isotopically distinct, and disjoint. A multicurve is *maximal* if it is not properly contained in any other multicurve. Of course, this simply means that the number of loops in the multicurve is  $3g - 3 + b$ , where  $g$  is the genus of  $S$  and  $b$  the number of punctures. A *gap* is a subsurface  $\alpha$  of a given surface  $S$  such that the topological boundary  $\partial\alpha$  of  $\alpha$  in  $S$  is a multicurve. A *gap* of a multicurve on a surface  $S$  is simply a component of  $S \setminus (\cup\Gamma)$ . If  $\alpha$  is any gap,  $\Gamma$  is a *multicurve in  $\alpha$*  if it satisfies all the above conditions for a closed surface, and, in addition,  $\cup\Gamma \subset \alpha$  and no loops in  $\Gamma$  are homotopic to components of  $\partial\alpha$ . A positively oriented Dehn twist round a loop  $\gamma$  on an oriented surface  $S$  will always be denoted by  $\tau_\gamma$ .

### 2.2 Teichmüller space

We consider Teichmüller space  $\mathcal{T}(S)$  of a surface  $S$ . If  $\varphi_i : S \rightarrow S_i = \varphi_i(S)$  is an orientation preserving homeomorphism, and  $S_i$  is a complete hyperbolic surface with constant curvature  $-1$ , then we define the equivalence relation  $\varphi_1 \sim \varphi_2$  if

and only if there is an orientation-preserving isometry  $\sigma : S_1 \rightarrow S_2$  such that  $\sigma \circ \varphi_1$  is isotopic to  $\varphi_2$ . We define  $[\varphi]$  to be the equivalence class of  $\varphi$ , and  $\mathcal{T}(S)$  to be the set of all such  $[\varphi]$ , this being regarded as sufficient since definition of a function includes definition of its domain. We shall often fix a complete hyperbolic metric of constant curvature  $-1$  on  $S$  itself, which we shall also refer to as “the” Poincaré metric on  $S$ .

Complete hyperbolic structure in dimension two is equivalent to complex structure, for any orientable surface  $S$  of finite topological type and negative Euler characteristic, by the Riemann mapping theorem. So endowing such a surface  $S$  with a complex structure defines an element of the Teichmüller space  $\mathcal{T}(S)$ . More generally, the Measurable Riemann Mapping Theorem [3] means that supplying a bounded measurable conformal structure for  $S$  is enough to define an element of  $\mathcal{T}(S)$ , and indeed  $\mathcal{T}(S)$  is often (perhaps usually) defined in this way.

### 2.3 Teichmüller distance

We shall use  $d$  to denote Teichmüller distance, so long as the Teichmüller space  $\mathcal{T}(S)$  under consideration is regarded as clear. Moreover a metric  $d$  will always be Teichmüller metric unless otherwise specified. If more than one space is under consideration, we shall use  $d_S$  to denote Teichmüller distance on  $\mathcal{T}(S)$ . The distance is defined as

$$d([\varphi_1], [\varphi_2]) = \inf \left\{ \frac{1}{2} \log \|\chi\|_{qc} : [\chi \circ \varphi_1] = [\varphi_2] \right\}$$

where

$$\|\chi\|_{qc} = \|K(\chi)\|_{\infty}, \quad K(\chi)(z) = \lambda(z)/\mu(z),$$

where  $\lambda(z)^2 \geq \mu(z)^2 \geq 0$  are the eigenvalues of  $D\chi_z^T D\chi_z$ , and  $D\chi_z$  is the derivative of  $\chi$  at  $z$  (considered as a  $2 \times 2$  matrix) and  $D\chi_z^T$  is its transpose. The infimum is achieved uniquely at a map  $\chi$  which is given locally in terms of a unique quadratic mass 1 differential  $q(z)dz^2$  on  $\varphi_1(S)$ , and its *stretch*  $p(z)dz^2$  on  $\varphi_2(S)$ . The local coordinates are

$$\zeta = x + iy = \int_{z_0}^z \sqrt{q(t)} dt,$$

$$\zeta' = \int_{z'_0}^{z'} \sqrt{p(t)} dt.$$

With respect to these local coordinates,

$$\chi(x + iy) = \lambda x + i \frac{y}{\lambda}.$$

So the distortion  $K(\chi)(x + iy) = \lambda$  is constant. The singular foliations  $x = \text{constant}$  and  $y = \text{constant}$  on  $\varphi(S)$  given locally by the coordinate  $x + iy$  for  $q(z)dz^2$  are known as the *stable and unstable foliations* for  $q(z)dz^2$ . We also say that  $q(z)dz^2$  is the *quadratic differential at  $[\varphi_1]$*  for  $d([\varphi_1], [\varphi_2])$ , and  $p(z)dz^2$  is its *stretch* at  $[\varphi_2]$ .

## 2.4 Thick and thin parts

Let  $\varepsilon$  be any fixed Margulis constant for dimension two, that is, for any hyperbolic surface  $S$ , if  $S_{<\varepsilon}$  is the set of points of  $S$  through which there is a nontrivial closed loop of length  $< \varepsilon$ , then  $S_{<\varepsilon}$  is a (possibly empty) union of cylinders with disjoint closures. Then  $(\mathcal{T}(S))_{<\varepsilon}$  is the set of  $[\varphi]$  for which  $(\varphi(S))_{<\varepsilon}$  contains at least one nonperipheral cylinder. The complement of  $(\mathcal{T}(S))_{<\varepsilon}$  is  $(\mathcal{T}(S))_{\geq\varepsilon}$ . We shall sometimes write simply  $\mathcal{T}_{<\varepsilon}$  and  $\mathcal{T}_{\geq\varepsilon}$  if it is clear from the context what is meant. We shall also write  $\mathcal{T}(\gamma, \varepsilon)$  for the set of  $[\varphi]$  such that  $(\varphi(S))_{<\varepsilon}$  contains a loop homotopic to  $\varphi(\gamma)$ . If  $\Gamma$  is a set of loops, we write

$$\mathcal{T}(\Gamma, \varepsilon) = \cup \{ \mathcal{T}(\gamma, \varepsilon) : \gamma \in \Gamma \}.$$

## 2.5 Length and the interpretation of Teichmüller distance

We fix a surface  $S$ . It will sometimes be convenient to fix a hyperbolic metric on  $S$ , in which case we shall use  $|\gamma|$  to denote length of a geodesic path  $\gamma$  with respect to this metric. With abuse of notation, for  $[\varphi] \in \mathcal{T}(S)$  and a nontrivial nonperipheral closed loop  $\gamma$  on  $S$ , we write  $|\varphi(\gamma)|$  for the length, with respect to the Poincaré metric on the hyperbolic surface  $\varphi(S)$ , of the geodesic homotopic to  $\varphi(\gamma)$ . We write  $|\varphi(\gamma)|'$  for a modification of this, obtained as follows. We change the metric in  $\varepsilon_0$ -Margulis tube of  $\varphi(S)$ , for some fixed Margulis constant  $\varepsilon_0$ , to the Euclidean metric for this complex structure in the  $\varepsilon_0/2$ -Margulis tube, so that the loop round the annulus is length  $\sqrt{|\varphi(\gamma)|}$ , and a convex-linear combination with the Poincaré metric between the  $\varepsilon_0$ -Margulis tubes and  $\varepsilon_0/2$ -Margulis tubes. Then we take  $|\varphi(\gamma)|'$  to be the length of the geodesic isotopic to  $\varphi(\gamma)$  with respect to this modified metric. If the geodesic homotopic to  $\varphi(\gamma)$  does not intersect any Margulis tube, then, of course,  $|\varphi(\gamma)| = |\varphi(\gamma)|'$ . Then for a constant  $C$  depending only on  $S$  and  $\varepsilon_0$ .

$$|\text{Max}\{|\log |\varphi_2(\gamma)|' - \log |\varphi_1(\gamma)|' : \gamma \in \Gamma\} - d([\varphi_1], [\varphi_2])| \leq C. \quad (2.5.1)$$

Here,  $\Gamma$  can be taken to be the set of all simple closed nonperipheral nontrivial closed loops on  $S$ . This estimate on Teichmüller distance derives from the fact that  $|\varphi(\gamma)|'$  is inversely proportional to the largest possible square root of modulus of an embedded annulus in  $S$  homotopic to  $\varphi(\gamma)$ . See also 14.3, 14.4 and 14.7 of [50] (although the square root of modulus was mistakenly left out of [50]) but this estimate appears in other places, for example [28]. We can simply take  $\Gamma$  to be any set of simple closed nontrivial nonperipheral loops on  $S$  such that that every component of  $S \setminus (\cup \Gamma)$  is a topological disc with at most one puncture. We shall call such a loop set *cell-cutting*

Fix a Margulis constant  $\varepsilon_0$ . We define

$$|\varphi(\gamma)|'' = |\varphi(\gamma) \cap (\varphi(S))_{\geq\varepsilon_0}| + \sum \{ |n_{\zeta, \gamma}([\varphi])| : |\varphi(\zeta)| < \varepsilon_0 \},$$

where  $n_{\zeta, \gamma}([\varphi])$  is such that  $|\tau_{\zeta}^m(\gamma)|$  is minimised when  $m = n$ , for  $\tau_{\zeta}$  as in 2.1. (See 15.9 of [50] and 2.6.)

There is  $L(\varepsilon_0)$  such that a cell-cutting loop set  $\Gamma_i$  can always be chosen with  $|\varphi_i(\Gamma_i)|'' \leq L(\varepsilon_0)$ . Having fixed such loop sets, there is a constant  $C(\varepsilon_0)$  such that

$$|\text{Max}\{\log(\#(\gamma_1 \cap \gamma_2)) : \gamma_1 \in \Gamma_1, \gamma_2 \in \Gamma_2\} - d([\varphi_1], [\varphi_2])| \leq C(\varepsilon_0). \quad (2.5.2)$$

## 2.6 Projections to subsurface Teichmüller spaces

For any gap  $\alpha \subset S$ , we define a topological surface  $S(\alpha)$  without boundary and a continuous map  $\pi_\alpha : \mathcal{T}(S) \rightarrow \mathcal{T}(S(\alpha))$ . We define  $\varphi_\alpha(S(\alpha))$  by defining its conformal structure. After isotopy of  $\varphi$ , we can assume that all the components of  $\varphi(\partial\alpha)$  are geodesic. We now write  $\overline{\varphi(\alpha)}$  for the compactification of  $\varphi(\alpha)$  obtained by cutting along  $\varphi(\partial\alpha)$  and adding boundary components, each one isometric to some component of  $\varphi(\partial\alpha)$ . Then we form the Riemann surface  $\varphi_\alpha(S(\alpha))$  by attaching a once-punctured disc  $\{z : 0 < |z| \leq 1\}$  to  $\overline{\varphi(\alpha)}$  along each component of  $\varphi(\partial\alpha)$ , taking the attaching map to have constant derivative with respect to length on the geodesics  $\varphi(\partial\alpha)$  and length on the unit circle. Then we define  $\varphi_\alpha = \varphi$  on  $\alpha$  and then extend the map homeomorphically across each of the punctured discs. Then  $[\varphi_\alpha]$  is a well-defined element of  $\mathcal{T}(S(\alpha))$ .

Now let  $\alpha$  be a nontrivial nonperipheral simple closed loop. Fix an orientation on  $\alpha$ . Then we define

$$S(\alpha) = \overline{\mathbb{C}} \setminus \{\pm 2, \pm \frac{1}{2}\}.$$

Now we define an element  $[\varphi_\alpha] = \pi_\alpha([\varphi]) \in \mathcal{T}(S(\alpha))$ , for each  $[\varphi] \in \mathcal{T}(S)$ , as follows. Fix a Margulis constant  $\varepsilon$ . If  $\varphi(\alpha) \leq \varepsilon$ , let  $A$  be the closed  $\varepsilon$ -Margulis tube in  $\varphi(S)$  homotopic to  $\varphi(\alpha)$ . If  $|\varphi(\alpha)| > \varepsilon$ , let  $A$  be the closed  $\eta$ -neighbourhood of the geodesic homotopic to  $\varphi(\alpha)$  where  $\eta$  is chosen so that  $A$  is an embedded annulus, and thus can be chosen bounded from 0 if  $|\varphi(\alpha)|$  is bounded above. Fix a simple closed geodesic  $\beta(\alpha)$  which intersects  $\alpha$  at most twice and at least once, depending on whether or not  $\alpha$  separates  $S$ . We can assume after isotopy that  $\varphi(\alpha)$  and  $\varphi(\beta(\alpha))$  are both geodesic, and we fix a point  $x_1(\alpha) \in \varphi(\alpha \cap \beta(\alpha))$ . We make a Riemann surface  $S_1$  homeomorphic to the sphere, by attaching a unit disc to each component of  $\partial A$ , taking the attaching maps to have constant derivative with respect to length. Then we define  $\varphi_\alpha$  to map  $\overline{\mathbb{C}}$  to  $S_1$  by mapping  $\{z : |z| = 1\}$  to  $\varphi(\alpha)$ , 1 to  $x_1(\alpha)$ ,  $\{z : \frac{1}{2} \leq |z| \leq 2\}$  to  $A$  and  $\{z \in \mathbb{R} : \frac{1}{2} \leq z \leq 2\}$  to the component of  $\varphi(\beta(\alpha)) \cap A$  containing  $\alpha$ . Then  $\varphi_\alpha(S(\alpha))$  is a four-times punctured sphere and so we have an element  $[\varphi_\alpha] \in \mathcal{T}(S(\alpha))$ . Now the Teichmüller space  $\mathcal{T}(S(\alpha))$  is isometric to the upper half plane  $H^2$  with metric  $\frac{1}{2}d_P$ , where  $d_P$  denotes the Poincaré metric  $(1/y)(dx^2 + dy^2)$ . This is well-known. We now give an identification. Let  $n_\alpha([\varphi]) = n_{\alpha, \beta(\alpha)}([\varphi])$  be the integer assigning the minimum value to

$$m \rightarrow |\varphi(\tau_\alpha^m(\beta(\alpha)))|.$$

If there is more than one such integer then we take the smallest one. There is a bound on the number of such integers of at most two consecutive ones. We

see this as follows. Let  $\ell$  be a geodesic in the hyperbolic plane and let  $g$  be a Möbius involution such that  $g.\ell$  is disjoint from, and not asymptotic, to  $\ell$ , and such that the common perpendicular geodesic segment from  $\ell$  to  $g.\ell$  meets them in points  $x_0, g.x_0$ , for some  $x_0 \in \ell$ . Then the complete geodesics meeting both  $\ell$  and  $g.\ell$  and crossing them both at the same angle, are precisely those that pass through points  $x$  and  $g.x$  for some  $x \in \ell$ , and the hyperbolic length of the segment joining  $x$  and  $g.x$  increases strictly with the length between  $x_0$  and  $x$ . This implies the essential uniqueness of  $n$ , as follows. We take  $\ell$  to be a lift of  $\varphi(\alpha)$  to the universal cover, and let  $\ell_1$  be another lift of  $\varphi(\alpha)$ , such that some lift of  $\varphi(\beta(\alpha))$  has endpoints on  $\ell$  and  $\ell_1$ . Then  $g$  is determined by making  $\ell_1 = g.\ell$  for  $g$  as above. But also  $\ell_1 = g_2.\ell$ , where  $g_2$  is the element of the covering group corresponding to  $\varphi(\beta(\alpha))$ . We also have an element  $g_1$  of the covering group corresponding to  $\varphi(\alpha)$ , which preserves  $\ell$  and orientation on  $\ell$ . Then  $|\varphi(\tau_\alpha^m(\beta(\alpha)))|$  is the distance between  $x$  and  $g.x$  for the unique  $x$  such that some lift of a loop freely homotopic to  $\varphi(\tau_\alpha^m(\beta(\alpha)))$  has endpoints at  $x$  and  $g.x$ . The endpoints are also  $g_1^{-m}.y$  and  $g_2.y$  for  $y$  such that  $x = g_1^{-m}.y$ . So  $x$  is determined by the  $y = y_m$  such that  $g.x = g_2.g_1^m.x$ . So then  $d(x, x_0) = \frac{1}{2}d(x_0, g^{-1}g_2g_1^m.x_0)$ , which takes its minimum at either one, or two adjacent, values of  $m$ .

Then the isometric identification with  $H^2$  can be chosen so that, if we use the identification to regard  $\pi_\alpha$  as a map to  $H^2$ ,

$$\pi_\alpha([\varphi]) = n_\alpha([\varphi]) + i|\varphi(\alpha)|^{-1} + O(1). \quad (2.6.1)$$

If  $\alpha$  is either a gap or a loop we now define a semimetric  $d_\alpha$  by

$$d_\alpha([\varphi_1], [\varphi_2]) = d_{S(\alpha)}(\pi_\alpha([\varphi_1]), \pi_\alpha([\varphi_2])).$$

## 2.7 Use of the semimetrics to bound metric distance

Using 2.5.1, we see that for a constant  $C = C(L_0)$ , if  $\alpha$  is a gap and  $|\varphi(\partial\alpha)| \leq L_0$  then for all  $[\varphi_1], [\varphi_2] \in \mathcal{T}(S)$ ,

$$d_\alpha([\varphi_1], [\varphi_2]) \leq d([\varphi_1], [\varphi_2]) + C(L_0). \quad (2.7.1)$$

This is simply because, for any simple closed nontrivial nonperipheral loop  $\gamma \subset \alpha$ , an annulus homotopic to  $\varphi(\gamma)$  with modulus boundedly proportional to the maximum possible is contained in the surface homotopic to  $\varphi(\alpha)$ , and bounded by the geodesics homotopic to  $\varphi(\partial\alpha)$ . But (2.7.1) also holds if  $\alpha$  is a loop with  $|\varphi(\alpha)| \leq L_0$ , by considering (2.5.1) applied to the loops  $\alpha$  and  $\tau_\alpha^n(\beta(\alpha))$  ( $n \in \mathbb{Z}$ ).

There is a converse to (2.7.1), again using (2.5.1), which as noted works for a restricted set of loops. Suppose that we have a set  $\Gamma \subset S$  of simple closed nontrivial nonperipheral loops which are all isotopically distinct and disjoint and such that

$$|\varphi(\Gamma)| \leq L_0.$$

Let  $\Sigma(\Gamma)$  denote the set of gaps of  $\Gamma$ . Then for a constant  $C(L_0)$ ,

$$d_S([\varphi_1], [\varphi_2]) \leq \text{Max}\{d_\alpha([\varphi_1], [\varphi_2]) : \alpha \in \Gamma \cup \Sigma(\Gamma)\} + C(L_0). \quad (2.7.2)$$

## 2.8

In 11.1 of [50], a projection  $\pi_\alpha$  was defined differently in the case of marked (equivalently punctured) spheres, the projection being done by simply deleting some of the punctures. So in those cases the condition  $d_\alpha \leq d$  was automatic. The identification of the image of the projection with the  $H^2$  in the case of  $\alpha$  a loop was done in 15.8 of [50]. If we denote the projection above by  $\pi_{1,\alpha}$  and the projection of [50] by  $\pi_{2,\alpha}$ , then for  $L'_0 = L'_0(L_0)$ , if  $|\varphi_1(\partial\alpha)| \leq L_0$ ,

$$d_{S(\alpha)}(\pi_{1,\alpha}([\varphi_1]), \pi_{2,\alpha}([\varphi_1])) \leq L'_0.$$

This is proved simply by constructing a bounded distortion homeomorphism between the surfaces given by  $\pi_{1,\alpha}([\varphi_1])$  and  $\pi_{2,\alpha}([\varphi_1])$ .

## 2.9 $d'_\alpha, d'_{\alpha_1, \alpha_2}$ .

The quantity  $d'_\alpha$  was defined 14.10 in [50], and is an extension of the definition of  $d_\alpha$ . Here,  $\alpha$  is either a nontrivial nonperipheral simple closed loop, or is a subsurface of  $S$  bounded by such loops, all isotopically disjoint and distinct. We use this concept when  $|\varphi_1(\alpha)| \leq L_0$ , or  $|\varphi_1(\partial\alpha)| \leq L_0$  for some fixed constant  $L_0$ . We fix a Margulis constant  $\varepsilon_0$ . If  $\alpha$  is a loop, we take  $[\varphi_3]$  to be the first point on the geodesic segment  $[[\varphi_1], [\varphi_2]]$  for which  $|\varphi_3(\alpha)| \geq \varepsilon_0$ , and

$$d'_\alpha([\varphi_1], [\varphi_2]) = d_\alpha([\varphi_1, \varphi_3]) + |\log |\varphi_2(\alpha)'||.$$

If  $\alpha$  is a subsurface, then

$$d'_\alpha([\varphi_1], [\varphi_2]) = \text{Max}\{|\log |\varphi_1(\gamma)'| - \log |\varphi_2(\gamma)'||\},$$

where the maximum is taken over multicurves in  $\alpha$  which are *cell-cutting in  $\alpha$* , that is, every component of  $\alpha \setminus (\cup \Gamma)$  is either a topological disc with at most one puncture, or an annulus parallel to the boundary. As in 2.5, we can take the maximum over a restricted set of multicurves  $\Gamma_1$  which are cell-cutting in  $\alpha$ , with  $|\varphi_1(\Gamma_1)|'' \leq L(\varepsilon_0)$ , at the expense of changing  $d'_\alpha([\varphi_1], [\varphi_2])$  by an additive constant. We can then also take a multicurve  $\Gamma_2$  which is cell-cutting in  $\alpha$  with  $|\varphi_2(\Gamma_2)| \leq L(\varepsilon_0)$ , and we then have an analogue of 2.5.2, which is more symmetric in  $[\varphi_1]$  and  $[\varphi_2]$ :

$$|\text{Max}\{\log(\#(\gamma_1 \cap \gamma_2)) : \gamma_1 \in \Gamma_1, \gamma_2 \in \Gamma_2\} - d'_\alpha([\varphi_1], [\varphi_2])| \leq C(\varepsilon_0, L_0). \quad (2.9.1)$$

It therefore makes sense to define  $d'_{\alpha_1 \alpha_2}([\varphi_1], [\varphi_2])$  if  $|\varphi_i(\partial\alpha_i)| \leq L_0$ , by taking loop sets  $\Gamma_i$  relative to  $\alpha_i$  like  $\Gamma_1$  relative to  $\alpha_1$  above, with  $|\varphi_i(\Gamma_i)|'' \leq L(\varepsilon_0)$ , and then defining

$$d'_{\alpha_1, \alpha_2}([\varphi_1], [\varphi_2]) = \text{Max}\{\log(\#(\gamma_1 \cap \gamma_2)) : \gamma_1 \in \Gamma_1, \gamma_2 \in \Gamma_2\}.$$

This is symmetric in  $\alpha_1$  and  $\alpha_2$ , and changing the loop sets  $\Gamma_1, \Gamma_2$  only changes the quantity  $d'_{\alpha_1, \alpha_2}([\varphi_1], [\varphi_2])$  by an additive constant. Also, if we write  $S$  as a

disjoint union of gaps and loops  $\alpha_2$ , then  $d'_{\alpha_1}([\varphi_1], [\varphi_2])$  is the maximum of all  $d'_{\alpha_1, \alpha_2}([\varphi_1], [\varphi_2])$  up to an additive constant.

If  $|\varphi_3(\partial\alpha)| \leq L_0$  then it is clear from 2.7 that

$$|d'_{\alpha}([\varphi_1], [\varphi_3]) - d_{\alpha}([\varphi_1], [\varphi_2])| \leq C(L_0).$$

If in addition  $[\varphi_3]$  is on the geodesics segment  $[[\varphi_1], [\varphi_2]]$  then we have from the definitions, and from 2.7:

$$d'_{\alpha}([\varphi_1], [\varphi_2]) \leq d_{\alpha}([\varphi_1], [\varphi_3]) + d'_{\alpha}([\varphi_3], [\varphi_2]) \leq C'(L_0).$$

Actually, a converse inequality holds, and will be discussed in section 6.

## 2.10 Projection is a single point.

Suppose that  $S = \alpha_1 \cup \alpha_2$  is a union of two closed subsurfaces, with disjoint interiors with  $\alpha_2$  not necessarily connected, and suppose that the common boundary consists of nontrivial nonperipheral loops. Fix a homeomorphism  $\varphi_1 : \alpha_1 \rightarrow \alpha'_1 = \varphi_1(\alpha_1)$ , where  $\alpha_1$  is a complete hyperbolic surface with geodesic boundary  $\varphi_1(\alpha_1)$ . Consider the set  $\mathcal{X}$  of  $[\varphi]$  in  $\mathcal{T}(S)$  such that  $\varphi(S) = S'$  is the union of the hyperbolic surface  $\alpha'_1$  and another subsurface joined along the geodesic boundary, and  $\varphi = \varphi_1$  on  $\alpha_1$ . Then the definition of  $\pi_{\alpha_1}$  in 2.6 is such that  $\pi_{\alpha}(\mathcal{X})$  is a single point in  $\mathcal{T}(S(\alpha_1))$ .

## 3 Pleated surfaces and geodesic laminations.

Throughout this section,  $S$  is a finite type surface and  $N$  is a complete hyperbolic 3-manifold with finitely generated fundamental group. We fix a hyperbolic metric on  $S$  and use the length conventions described in 2.5. In later subsections, we shall use the notation for subsets of  $N$  as described in section 1:  $N_d$  for the horoball deletion,  $N_c$  for the relative Scott core,  $U(e)$  for the component of  $N_d \setminus N_c$  which is a neighbourhood of the end  $e$ , and so on. We shall assume the result of [2] and [14] where necessary, that is, that  $N$  is tame.

### 3.1

The powerful tool of pleated surfaces was introduced by Thurston [58]. A basic reference is [15]. A *pleated surface* is a continuous map  $f : S \rightarrow N$  such that peripheral loops are mapped to cusps, and there is a geodesic lamination  $\mu$  on  $S$  with respect to some hyperbolic structure on  $S$ , such that each component of  $f(S \setminus \mu)$  is totally geodesic in  $N$  with boundary consisting of complete geodesics in  $N$ . A *geodesic lamination* is a closed set of nonintersecting geodesics on  $S$ . We then call  $\mu$  the *bending locus* of  $f$ . One could quibble about this because  $\mu$  is then not quite uniquely defined given a map  $f$ : there may be no bending along some leaves of  $\mu$ . But one can at least be sure that there is no bending along any geodesic which intersects  $\mu$  transversally, and it will be convenient in

the present work to assume that the structure of a pleated surface includes a lamination  $\mu$  such that any bending takes place inside  $\mu$  and none outside it.

A pleated surface  $f$  defines an element of  $\mathcal{T}(S)$  which we call  $[f]$ . This is done as follows, basically just pulling back the hyperbolic structure from  $f(S)$ . Of course,  $f$  is not an embedding in general, but it is a local embedding restricted to each component of  $S \setminus \mu$ , whose image is a complete geodesic triangle in  $N$ , and we take the new hyperbolic structure on  $S$  so that  $f$  is an isometry restricted to each such component. This actually defines the hyperbolic structure uniquely, in the given homotopy class. The transverse length of the geodesic lamination  $\mu$  in the new hyperbolic structure is 0, as it was before: bounded length arc intersecting a geodesic lamination has zero one-dimensional Lebesgue measure intersection with that lamination.

We shall sometimes write  $S(f)$  for the abstract Riemann surface with hyperbolic structure induced by  $f$ , if we want to make clear that the metric on it is that induced from  $f(S)$  locally but not globally. The map from  $S$  to  $S(f)$  is homotopic to a homeomorphism, and we denote the corresponding point of  $\mathcal{T}(S)$  by  $[f]$ . There is then a map defined using  $f$  defined pointwise from  $S(f)$  to  $f(S)$ , which we shall sometimes call  $\text{Imp}_1(f)$ , and which is distance-decreasing from  $S(f)$  to  $N$ , a map up to homotopy from  $S$  to  $f(S)$ , which we shall sometimes call  $\text{Imp}(f)$ . By abuse of notation we shall often denote all three of these maps by  $f$ , although  $\text{Imp}(f) = \text{Imp}_1(f) \circ f$ . We shall sometimes call either  $\text{Imp}(f)$  or  $\text{Imp}_1(f)$  the *impression* of  $f$ .

If  $\gamma$  is a closed loop in the pleating locus of  $f$  then  $f(\gamma) = \gamma_*$ , as we shall always denote the closed geodesic in  $N$  which is freely homotopic to  $f(\gamma)$ , and so, of course,  $|f(\gamma)| = |\gamma_*|$ . Similarly suppose that  $f_1$  and  $f_2$  are two pleated surfaces with the same pleating loci restricted to a subsurface  $\alpha$ , where  $\partial\alpha$  is also in the pleating locus of both. Then the subsets of  $S(f_i)$  bounded by  $f_i(\partial\alpha)$ , and isotopic to  $f_i(\alpha)$  are isometric under a homeomorphism  $\sigma$  with  $[\sigma \circ \varphi_1] = [\varphi_2]$ . This, combined with 2.10, will be of crucial importance later.

### 3.2 Making examples.

Fix a homotopy class  $f : S \rightarrow N$  which maps peripheral loops to peripheral loops. The easiest way to make a pleated surface in this homotopy class is to choose a lamination on  $S$  in which every leaf is either a simple closed geodesic or has each end asymptotic to either a closed geodesic of the lamination, or to a puncture, and so that any complementary component of the lamination is a triangle. There are many such examples. A pleated surface is obtained by *spinning round closed loops* if the nonclosed leaves of the pleating locus are obtained as follows. Take a maximal multicurve  $\Gamma$  on  $S$ . If  $f$  is injective on  $\pi_1$ , no further conditions on  $\Gamma$  are needed. But if  $f$  is not injective on  $\pi_1$ , let  $f_*(\gamma)$  be homotopically nontrivial and nonperipheral for each  $\gamma \in \Gamma$ . This ensures that each component  $P$  of  $S \setminus (\cup \Gamma)$  is homotopic in  $S$  to a closed incompressible pair of pants  $P'$  in  $N$ . We shall say that such a loop set  $\Gamma$  is *noncollapsing* (for  $f$ ).

Fix a decomposition of  $S$  into hexagons, possibly with some sides of zero



length, by taking a set of arcs in each pair of pants  $P$  with endpoints on boundary components or punctures. Fix a homeomorphism  $\varphi$  which is a nontrivial Dehn twist round each loop of  $\Gamma$ . If  $\alpha$  is any of the arcs then as  $n \rightarrow +\infty$ ,  $\varphi^n(\alpha)$  converges to an infinite arc which is asymptotic to a loop of  $\Gamma$  or a puncture of  $S$  at each end and the hexagons converge to ideal triangles. The union of  $\Gamma$  and the limits of the arcs is the pleating locus of a pleated surface *obtained by spinning round*  $\Gamma$ . These are, in fact, the basic examples mentioned in Thurston's notes [58] Chapter 8.

### 3.3 Bounded Diameter and Injectivity Radius.

Bonahon made the following important observation (essentially from 1.8 of [5], although finite simplicial surfaces are considered there).

**Bounded Diameter Lemma** *For a constant  $L(\varepsilon_0)$  and any pleated surface  $f : S \rightarrow N$ , the image in  $N$  of each component of the thick part  $(S(f))_{\geq \varepsilon_0}$  of  $S(f)$  has diameter  $\leq L(\varepsilon_0)$ .*

As for the thin part, there is a simple but important estimate on injectivity radius for pleated surfaces, which is actually part of the Bounded Diameter Lemma as stated by Bonahon [5] or the more general form in 2.1 of [16], where the injective-on- $\pi_1$  condition was also relaxed.

**Radius of Injectivity Lemma** *Fix a Margulis constant  $\varepsilon_0$  for both two and three dimensions. There are constants  $D_0$  and  $C = C(\varepsilon_0) > 1$  such that the following holds, with  $D_0$  depending only on the topological type of  $S$  and  $C(\varepsilon_0)$  depending only on  $\varepsilon_0$ . Let  $f : S \rightarrow N$  be a pleated surface. Let  $\zeta$  be any simple closed geodesic on  $S$  such that  $f(\zeta)$  is homotopically nontrivial and nonperipheral in  $N$ . If  $|f(\zeta)| < \varepsilon \leq \varepsilon_0$ , let  $T_S(f(\zeta), \varepsilon)$  denote the  $\varepsilon$ -Margulis tube in  $S(f)$ , if this is nonempty and let  $T_N(\zeta_*, \varepsilon)$  denote the Margulis tube in  $N$ , where  $\zeta_*$  is the closed geodesic freely homotopic to  $f(\zeta)$ . Then*

$$T_S(f(\zeta), \varepsilon) \subset T_N(\zeta_*, \varepsilon). \quad (3.3.1)$$

*Suppose in addition that  $f(\zeta)$  is nontrivial in  $N$  whenever  $\zeta$  is nontrivial in  $S$  and  $|f(\zeta)| \leq D_0$ . Then*

$$(S(f) \setminus T_S(\zeta, \varepsilon)) \cap T_N(\zeta_*, \varepsilon/C) = \emptyset. \quad (3.3.2)$$

(3.3.1) is simply because the pointwise map  $f : S(f) \rightarrow f(S)$  preserves length on paths in  $S$  and one can apply this to closed loops freely homotopic to  $f(\zeta)$ . (3.3.2) is a little more involved, and uses the extra hypothesis. The extra hypothesis, together with the Bounded Diameter Lemma, shows that  $f((S(f))_{\geq \varepsilon_0})$  cannot intersect  $N_{< \varepsilon_0/C}$  for a suitable  $C$ , which gives (3.3.2) for  $\varepsilon = \varepsilon_0$ . Then to get the result for a general  $\varepsilon$  we use the fact that  $f : S(f) \rightarrow N$  decreases length of paths joining  $\partial T(f(\zeta), \varepsilon_0)$  and  $\partial T(f(\zeta), \varepsilon)$ .  $\square$

The following suggests why this hypothesis of “no bounded trivial loops” arises.

**Lemma 3.4.** *Given  $L$ , there is  $\Delta$  depending only on  $L$  and  $N$  such that the following holds. Let  $f : S(e) \rightarrow N$  be any pleated surface such that the restriction*

to  $S_d(e)$  is homotopic to the natural inclusion of  $S_d(e)$  in  $N_c$ , and such that with  $f(S(e))$  contains at least one point in  $U(e)$  distance  $\geq \Delta$  from  $N_c$ . Suppose that  $\zeta \subset f(S(e))$  has length  $\leq L$  and is homotopic to a nontrivial nonperipheral loop on  $S(e)$ . Then  $\zeta$  is nontrivial nonperipheral in  $N$ .

*Proof* Choose  $\varepsilon_L$  such that every component of  $N_{<\varepsilon_L}$  is distance  $\geq L$  from  $N_{\geq\varepsilon_0}$  for some fixed Margulis constant  $\varepsilon_0$ . Then let  $N_d$  be the complement of cuspidal components of  $N_{<\varepsilon_L}$ . We can take our original Scott core and extend it to a core for this new  $N_d$ . We continue to call the core  $N_c$ . Fix  $x \subset f(S)$  distance  $\geq \Delta$  from  $N_c$ . By the Bounded Diameter and Injectivity Radius Lemmas, any point  $x' \in f(S)$  can be joined to  $x$  by a path whose intersection with  $N_{\geq\varepsilon_0}$  has length  $\leq \Delta_1(\varepsilon_0)$ , where  $\Delta_1(\varepsilon_0)$  depends only on  $\varepsilon_0$ . If  $\Delta$  is sufficiently large given  $L$  and  $N$ , we can assume that none of these thin parts of  $N$  is within  $2L + \Delta_1(\varepsilon_0)$  of  $N_c$ . First suppose that  $\zeta$  is trivial in  $N$ . Then  $\zeta$  lifts to a closed loop  $\tilde{\zeta}$  in the universal cover  $H^3$ , and since it has length  $\leq L$  it also has diameter  $\leq L$ , and we can find a continuous map of the disc  $D$  into  $H^3$  with  $\partial D$  mapped to  $\tilde{\zeta}$ , by joining all points on  $\tilde{\zeta}$  to some fixed point on  $\tilde{\zeta}$  by geodesic segments. The image of this disc in  $H^3$  then has diameter  $\leq 2L$ , as does its projection in  $N$ . Then by the choice of  $\Delta$ , this disc does not intersect  $N_c$ , which is impossible. If  $\zeta$  is peripheral in  $N$ , then argument is similar. This time, we have a bound in terms of  $L$  on the diameter of the image of the homotopy between  $\zeta$  and the corresponding element of  $\partial N_d$ . So we again deduce that the homotopy cannot intersect  $N_c$ , if  $\Delta$  is sufficiently large.  $\square$

### 3.5 Short Bridge Arcs.

Here is another result which weakens a common hypothesis of doubly incompressible for pleated surfaces, but *strengthens another hypothesis* to obtain a result in the not-injective-on- $\pi_1$  case. The result which is being generalised is the Short Bridge Arc Lemma 2.2 of [37] or 5.5, Uniform Injectivity, of [59]. There is a Uniform Injectivity result in a somewhat different direction in Otal's thesis [45], and others in [40] and [8]. We need a notion of *badly bent annuli* for  $f_0$  and for a fixed Margulis constant  $\varepsilon_0$ . A *badly bent annulus* (for  $f_0$  and  $\varepsilon_0$ ) is a Margulis tube  $T(f_0(\gamma), \varepsilon(\gamma)) \subset S(f_0)$  for which  $\gamma$  is in the pleating locus of  $f_0$ ,  $|f_0(\gamma)| < \varepsilon_0$ , and  $\varepsilon(\gamma)$  is the largest number  $\leq \varepsilon_0$  such that either  $\varepsilon(\gamma) = |f_0(\gamma)|$  or the images under  $f_0$  two components of  $\partial T(f_0(\gamma), \varepsilon(\gamma))$  are distance  $\leq 1$  apart in  $N$ .

**Lemma** *The following holds for  $L'_0, L_1$  sufficiently large, depending only on the topological type of  $S$  and a given constant  $L_0$ . Let  $N$  be a hyperbolic manifold. Fix a Margulis constant  $\varepsilon_0$ . Let  $f_0 : S(f_0) \rightarrow N$  be a pleated surface. For  $i = 1, 2$  let  $\ell_i = \{x_{i,t} : 0 \leq t \leq T\}$  be geodesic segments in  $S(f_0)$ , such that  $f_0(\ell_i)$  is a geodesic segment in  $N$ , and  $t$  is the length parameter. Let  $\tau \subset S(f_0)$  be a geodesic segment, with respect to the hyperbolic structure on  $S(f_0)$ , joining  $x_{1,0}$  and  $x_{2,0}$ , and let  $\tau_t$  be the continuously varying geodesic segment joining  $x_{1,t}$  and  $x_{2,t}$ .*

*Suppose that  $f_0(\tau_t)$  is homotopic in  $N$ , via homotopy fixing endpoints, to a*

geodesic segment in  $N$  of length  $\leq L_0$ , for all  $t$ .

If  $f_0$  is not injective on  $\pi_1$ , let  $A \subset (S(f_0))_{<\varepsilon_0}$  be the union of badly bent annuli for  $f_0$  and  $\varepsilon_0$  which intersect  $\tau$ , and make two further assumptions.

1. The length of  $\tau$  in  $S(f_0) \setminus A$  is  $\leq L_0$ .
2. For any nontrivial loop  $\gamma \subset S(f_0)$  for which  $f_0(\gamma)$  is trivial in  $N$ ,  $|f_0(\gamma) \setminus A| \geq L_1$ .

Then, after translating the length parameter on one of  $\ell_1, \ell_2$  by  $\leq 2 \log L_0$  if necessary, either  $\tau_t$  has length  $\leq 1$  in  $S(f_0)$  for all  $L'_0 \leq t \leq T - L'_0$ , or  $x_{1,t}$  and  $x_{2,t}$  are in the same badly bent annulus, on opposite sides of the core loop, either for all  $L'_0 \leq t \leq T$ , or for all  $0 \leq t \leq T - L'_0$ .

This has been stated a bit differently from 2.2 of [37], even leaving aside the different hypotheses in the not-injective-on- $\pi_1$  case. It is probably worth saying at this point that the proof does not use geometric limits, in contrast to the proofs of similar results that I am aware of.

*Proof.* Write  $d_2$  and  $d_3$  for the hyperbolic distances in the universal covers  $H^2$  and  $H^3$  of  $S(f_0)$  and  $N$  respectively, where  $H^2$  projects to the lift of  $f_0(S(f_0))$  in  $H^3$ . Then  $\pi_1(S(f_0))$  acts on  $H^2$ . Identify  $x_{i,t}, \tau_t, \ell_i$  with lifts to  $H^2$ . Now there is a lift  $\tilde{f}_0 : H^2 \rightarrow H^3$  of  $f_0$ . By abuse of notation, we also write  $x_{i,t}$  for  $\tilde{f}_0(x_{i,t})$ . Note that

$$d_3(x_{i,t}, x_{i,s}) = |t - s|$$

for all  $s, t$ , but that in general

$$d_3(x_{1,t}, x_{2,t}) \leq d_2(x_{1,t}, x_{2,t}).$$

Assume without loss of generality that  $T \geq 3L_0$ . Also assume without loss of generality that  $x_{1,t}$  and  $x_{2,t}$  are equidistant from the ends of the common perpendicular between  $\ell_1$  and  $\ell_2$  in  $H^3$ , or from the intersection point, if there is one, or are the closest points on their respective geodesics, if these geodesics are asymptotic in  $H^3$ . This can be done by removing length  $\leq \log L_0 + O(1)$  from the  $\ell_i$ , and translating the length parameters by  $\leq$  that amount. We can also assume that  $d_3(x_{1,0}, x_{2,0}) \leq 1$ . This is because two sufficiently long geodesics in  $H^3$  which are distance  $\leq L_0$  apart at the two pairs of endpoints are distance  $\leq 1$  apart in the interiors, apart from within  $\log L_0 + O(1)$  of the endpoints. If  $f_0$  is not injective on  $\pi_1$ , since  $\tau$  changes as a result this translation, we may replace the hypothesis  $|f_0(\tau) \setminus A| \leq L_0$  by  $|f_0(\tau) \setminus A| \leq 2L_0$ .

Now fix the greatest  $t_1 \leq T$  such that  $|\tau_{t_1} \setminus A| \leq 2L_0$ , if such a  $t_1$  exists. If it does not exist, choose  $t_1$  so that  $d_2(x_{1,t}, x_{2,t}) = |\tau_t|$  is minimised at  $t = t_1$ . In both cases,

$$d_2(x_{1,t}, x_{2,t}) = |\tau_t| \geq |\tau_{t_1} \setminus A| \geq 2L_0$$

for all  $t_1 < t \leq T$ . If  $f_0$  is not injective on  $\pi_1$ , we also have a bound of  $3L_0$  on  $|\tau_t \setminus A|$  for  $0 \leq t \leq t_1$ , assuming  $L_0$  is large enough, depending only on a universal constant. We see this as follows. The geodesic segment  $[x_{i,0}, x_{i,T}]$  can intersect at most two components of  $A$ , one at the start and one at the

end, because the core loops of  $A$  are geodesic and cannot be crossed by other geodesics. Also, because of the properties of polygons of geodesics in  $H^2$ ,  $\tau_t$  is in a  $C_0$ -neighbourhood of  $\tau \cup [x_{1,0}, x_{1,t}] \cup [x_{2,0}, x_{2,t}]$ , for a universal constant  $C_0$ . To within a universal constant,  $|\tau_t \setminus A|$  is the sum of the length of 3 segments, on  $\tau \setminus A$ ,  $[x_{1,0}, x_{1,t}] \setminus A$  and on  $[x_{2,0}, x_{2,t}] \setminus A$ , where these 3 segments are maximal with respect to the property that for each segment, no point is distance  $\leq C_0$  in  $H^2$  from any point on either of the other two segments. The length of such maximal segments on  $[x_{1,0}, x_{1,t}] \setminus A$  and  $[x_{2,0}, x_{2,t}] \setminus A$  is essentially increasing with  $t$ : note that  $d_2(x_{1,s}, x_{2,u}) \geq d_3(x_{1,s}, x_{2,u}) \geq |s - u| - 1$ . So  $|\tau_t \setminus A|$  is bounded to within a universal constant by the maximum of  $|\tau_{t_1} \setminus A|$  and  $|\tau \setminus A|$ , for  $t \leq t_1$ . It  $t_1 = T$ ,  $\tau_T \subset A$ , and  $\tau_T$  is contained in a single component of  $A \setminus f_0(\gamma)$ , where  $f_0(\gamma)$  is the core loop in this component of  $A$ , then the bound on  $d_3(x_{1,T}, x_{2,T})$  shows that, as elements of  $N$ ,  $x_{j,T}$ , is in a component of  $\partial N_{\varepsilon_j}$  with  $e^{-2L_0} \leq \varepsilon_1/\varepsilon_2 \leq e^{2L_0}$ . Then by the Radius of Injectivity Lemma, as elements of  $S(f_0)$ ,  $x_{j,T}$  is in a component of  $\partial(S(f_0))_{\geq \varepsilon'_j}$  with  $C^{-2}e^{-2L_0} \leq \varepsilon'_1/\varepsilon'_2 \leq C^2e^{2L_0}$ . Since the components of  $\partial(S(f_0))_{\geq \varepsilon'_j}$  are not separated by a core loop  $f_0(\gamma)$ , we have a bound on  $d_2(x_{1,T}, x_{2,T})$  in this case. So we have a bound on  $d_2(x_{1,t}, x_{2,t})$  for  $t \leq t_1$  in terms of the bound on  $|\tau \setminus A|$ , in the non-injective case, and the proof is completed if we can bound  $T$  above in terms of  $L_0$ . In the case when  $f_0$  is injective on  $\pi_1$ , we need to do a similar procedure for decreasing  $t$ . So in that case, we similarly define  $t_0$  be the least  $t \geq 0$  such that  $|\tau_{t_0} \setminus A| \leq 2L_0$ , if such a  $t_0$  exists, and if not, define  $t_0 = t_1$ . We then need to bound  $t_0$  above in terms of  $L_0$ .

We return to  $t \geq t_1$  and assume that  $t_1 < T$ . We also claim, in both injective and noninjective cases, that the minimum of  $|\tau_t| = d_2(x_{1,t}, x_{2,t})$  occurs at some  $t \leq t_1 + C'_0$  for a universal constant  $C'_0$ , even if we are assuming that  $t = t_1$  is a minimum of  $|\tau_t \setminus A|$ , rather than of  $|\tau_t|$ . This is the same argument as before. To within a universal constant,  $d_2(x_{1,t}, x_{2,t}) = |\tau_t|$  is the sum of the lengths of 3 segments, on  $\tau$ ,  $[x_{1,0}, x_{1,t}]$  and on  $[x_{2,0}, x_{2,t}]$ , where these 3 segments are maximal with respect to the property that for each segment, no point is distance  $\leq C_0$  in  $H^2$  from any point on either of the other two segments. Since we know that  $|\tau_t \setminus A|$  is not decreasing for  $t > t_1$  near  $t_1$   $[x_{1,t_1}, x_{1,t}]$  and  $[x_{2,t_1}, x_{2,t}]$  are bounded from  $\tau$  for  $t \geq t_1$ , and also bounded from each other, because otherwise  $|\tau_{t_1} \setminus A| < 2L_0$ . It follows that, for a universal constant  $C''_0$ , for all  $t \geq t_1$ ,

$$d_2(x_{1,t}, x_{2,t}) \geq d_2(x_{1,0}, x_{2,0}) + 2t - C''_0.$$

Now, to simplify the notation, write  $t_1 = 0$ . Define  $\varepsilon = \varepsilon_0/4C$ , for  $C$  the constant in the Radius of Injectivity Lemma. Fix a constant  $L_2$ . If  $L_1$  is sufficiently large given  $L_0$  and  $L_2$ , and  $T \geq L_2$ , we can assume that  $x_{1,t} \notin (S(f_0))_{<\varepsilon}$  so long as  $0 \leq t \leq L_2$ . For otherwise, since  $d_3(x_{1,t}, x_{2,t}) < 1$ ,  $x_{1,t}$  and  $x_{2,t}$  are in the same lift of the same component of  $N_{<\varepsilon_0/C}$  in  $H^3$ . Assuming that  $L_1$  is large enough for the Radius of Injectivity Lemma to hold, both  $x_{1,t}$  and  $x_{2,t}$  are in  $(S(f_0))_{<\varepsilon_0}$ . Let  $\gamma$  be the core loop or parabolic of the component of  $(S(f_0))_{<\varepsilon_0}$  up to homotopy, whose lift contains  $x_{1,t}$ . If  $x_{1,t}$  and  $x_{2,t}$  are in the same lift in  $H^2$  of the same component of  $(S(f_0))_{<\varepsilon_0}$  in  $H^2$ ,

then  $\tau_t$  is in this component of  $(S(f_0))_{<\varepsilon_0}$ , which means we have  $d_2(x_{1,t}, x_{2,t}) \leq Cd_3(x_{1,t}, x_{2,t}) + O(1)$ , for  $C$  as in the Radius of Injectivity Lemma, and either  $T = 0$ , in which case the proof is finished, or we have a contradiction to our assumption that  $d_2(x_{1,t}, x_{2,t}) \geq 2L_0$  for all  $0 \leq t \leq T$ , assuming only that  $L_0$  is sufficiently large given  $\varepsilon_0$ . If  $x_{1,t}$  and  $x_{2,t}$  are not in the same lift in  $H^2$  of the same component of  $(S(f_0))_{<\varepsilon_0}$ , then we consider the lift to  $H^2$  of  $\tau_t * \beta * \tau_t^{-1}$ , where  $\beta$  is the core loop of the component of  $(S(f_0))_{<\varepsilon_0}$  containing  $x_{2,t}$ . Replacing  $\beta$  by  $\beta^{-1}$  if necessary,  $\gamma$  and  $\tau_t * \beta * \tau_t^{-1}$  represent the same short loop in  $N$  but different loops in  $S(f_0)$ . Then  $\tau_t * \beta * \tau_t^{-1} * \gamma^{-1}$  then gives a nontrivial loop in  $\pi_1(S(f_0))$  with length  $\leq L_1$  in  $S(f_0) \setminus A$ , which is trivial in  $\pi_1(N)$ , if  $L_1$  is sufficiently large given  $L_0$  and  $L_2$ , which contradicts our hypothesis.

So now we assume that for  $t \in [0, L_2]$ ,  $x_{1,t}$  and  $x_{2,t}$  lie in  $(S(f_0))_{\geq \varepsilon}$ , and that  $T = d(x_{1,0}, x_{1,T}) \geq L_2$ . We shall obtain a contradiction for  $T$  and  $L_2$  sufficiently large given a universal constant —  $L_1$  sufficiently large given  $L_0$ , in the case when  $f_0$  is not injective on  $\pi_1$ . So now we can assume that  $[x_{1,0}, x_{1,T}]$  lies in a component of  $(S(f_0))_{\geq \varepsilon}$ , and similarly for  $[x_{1,0}, x_{1,T}]$  — possibly for a different component of  $(S(f_0))_{\geq \varepsilon}$ . So for  $C_1$  depending only on the topological type of  $S$ , and on  $\varepsilon$ , we can find sequences  $g_j$  and  $g'_j$  in  $\pi_1(S(f_0))$  with  $g_{j+1} = g_j h_j$ ,  $g'_{j+1} = g'_j h'_j$ , with

$$\begin{aligned} d_2(x_{1,0}, h_j.x_{1,0}) &\leq C_1, & d_2(x_{2,0}, h'_j.x_{2,0}) &\leq C_1, \\ d_2(x_{1,t_j}, g_j.x_{1,0}) &\leq C_1, & d_2(x_{2,t_j}, g'_j.x_{2,0}) &\leq C_1, \\ 5C_1 &\leq t_{j+1} - t_j \leq 6C_1. \end{aligned}$$

Then

$$d_3(x_{1,0}, g_j^{-1}g'_j.x_{2,0}) = d_3(g_j.x_{1,0}, g'_j.x_{2,0})$$

$$\leq d_2(g_j.x_{1,0}, x_{1,t_j}) + d_3(x_{1,t_j}, x_{2,t_j}) + d_2(x_{2,t_j}, g'_j.x_{2,0}) \leq 2C_1 + 1,$$

while

$$\begin{aligned} d_2(x_{1,0}, g_j^{-1}g'_j.x_{2,0}) &= d_2(g_j.x_{1,0}, g'_j.x_{2,0}) \\ &\geq d_2(x_{1,t_j}, x_{2,t_j}) - 2C_1 \geq 2t_j + d_2(x_{1,0}, x_{2,0}) - C_0'' - 2C_1 \\ &> 2t_{j-1} + d_2(x_{1,0}, x_{2,0}) + 3C_1 - C_0'' \geq d_2(x_{1,t_{j-1}}, x_{2,t_{j-1}}) + 3C_1 - C_0'' \\ &> d_2(g_{j-1}.x_{1,0}, g'_{j-1}.x_{2,0}) + C_1 - C_0'' \\ &> d_2(g_{j-1}.x_{1,0}, g'_{j-1}.x_{2,0}) = d_2(x_{1,0}, g_{j-1}^{-1}g'_{j-1}.x_{2,0}). \end{aligned}$$

So, as points in  $H^3$ , all the points  $g_j^{-1}g'_j.x_2$  lie in a ball in  $H^3$  centred on  $x_1$  of radius  $2C_1 + 1$ , but as points in  $H^2$ , they are distinct. So assuming that  $T$  is sufficiently large given  $C_1$ , we can choose  $j$  and  $k$  so that  $d_2(g_j.x_{1,0}, g_k.x_{1,0})$  and  $d_2(g'_j.x_{2,0}, g'_k.x_{2,0})$  are bounded in terms of  $C_1$  and  $d_3(g_j^{-1}g'_j.x_2, g_k^{-1}g'_k.x_2) = 0$

but  $d_2(g_j^{-1}g'_j.x_2, g_k^{-1}g'_k.x_2) \neq 0$ . Because of the assumptions on  $\tau$ , both  $g_jg_k^{-1}$  and  $g'_j(g'_k)^{-1}$  can be represented, up to free homotopy, by closed loops at both  $x_1$  and  $x_2$  with length outside  $A$  bounded in terms of  $L_0$  and  $C_1$ , in the case of not injective on  $\pi_1$ . So, if  $L_2$  is sufficiently large given  $C_1$ , and  $L_1$  sufficiently large given  $L_0$ , we obtain a loop given by the free homotopy class of  $g_kg_j^{-1}g'_j(g'_k)^{-1}$  which is nontrivial in  $S(f_0)$ , with length  $\leq L_1$  in  $S(f_0) \setminus A$  in the case of  $f_0$  not being injective on  $\pi_1$ , and trivial in  $N$ . This gives the required contradiction.  $\square$

### 3.6 Efficiency of Pleated Surfaces.

Following Thurston in [59], Minsky used his Short Bridge Arc Lemma to prove the following. Actually, he proved more, but we only state what we need.

**Lemma** *The following holds for a suitable  $L_1$  and constant  $C_1(n)$  given  $n$ . Let  $f_0 : S \rightarrow N$  be a pleated surface with pleating locus including a maximal multicurve  $\Gamma$ . Let  $\gamma \subset S$  have  $\leq n$  intersections with closed loops in the pleating locus of  $f_0$ . Let  $A \subset (S(f_0))_{<\varepsilon_0}$  be the union of badly bent annuli for  $f_0$  for a fixed Margulis constant  $\varepsilon_0$  (3.5) which intersect  $f_0(\gamma) \subset S(f_0)$ . Suppose also that for any loop  $\gamma'$  for which  $f_0(\gamma')$  is trivial in  $N$ ,  $|f_0(\gamma') \setminus A| \geq L_1$ . Then*

$$|f_0(\gamma \setminus A)| \leq |\gamma_*| + C_1(n),$$

and  $f_0(\gamma \cap A)$  is a union of  $\leq C_1(n)$  segments, each of which is homotopic in  $N$ , via homotopy fixing endpoints, to a segment of length  $\leq 1$ .

*Proof.* For some  $n' \leq 2n$ , which depends on how the arcs of  $\gamma \setminus \Gamma$  compare with the infinite geodesics outside  $\Gamma$  which are in the pleating locus of  $f_0$ , we choose a connected union of  $4n'$  geodesic segments  $\ell_i$ ,  $1 \leq i \leq 4n'$  in the universal cover  $H^2$  of  $S(f_0)$  which projects to a closed loop homotopic to  $f_0(\gamma)$ . The segments  $\ell_{2i+1}$  project to short segments in  $S(f_0)$ , of length  $< 1$ , say. It is convenient to extend this sequence, using the action of the covering group to a bi-infinite sequence which is homotopic to the lift of  $f_0(\gamma)$  in  $H^2$ . So  $\ell_i$  and  $\ell_{i+4kn'}$  project to the same segment in  $S(f_0)$  for all integers  $k$ . The segments  $\ell_{4i}$  project to loops in  $\Gamma$ , probably not injectively. The geodesic containing  $\ell_{4i+2}$  is asymptotic in  $H^3$  to the geodesic containing  $\ell_{4i}$  at one end, and to the geodesic containing  $\ell_{4i+4}$  at the other, and, again,  $\ell_{4i+2}$  projects to the pleating locus of  $f_0$ . We apply the Short Bridge Arc Lemma 3.5 to each pair of geodesics containing  $\ell_{4i}$  and  $\ell_{4i+2}$  and to each pair containing  $\ell_{4i+2}$  and  $\ell_{4i+4}$ . The role of  $\tau$  is played by  $\ell_{4i+1}$  and  $\ell_{4i+3}$  respectively. By abuse of notation, we write  $A$  for the preimage in  $H^2$  of  $A \subset S(f_0)$ . Then 3.5 implies that  $\ell_{4i}$  and  $\ell_{4i+2}$  spread apart in  $H^2$  only when they spread apart in  $H^3$ . We can then reduce the lengths of even-indexed segments, possibly increasing the lengths of odd-numbered segments by a bounded amount, possibly removing some segments altogether and renumbering, but keeping the segments preserved by the action of the element of the covering group determined by  $f_0(\gamma)$ . Carrying out this procedure a bounded number of times, bounded in terms of  $n'$  and  $n$ , we reach the stage where, for some  $p \leq 4n'$ ,  $\cup_{0 \leq i < p}$  projects to  $f_0(\gamma)$  up to homotopy, and each segment  $\ell_i$  is the union of at most two end segments in  $A$ , one or two

segments of length bounded in terms of  $i$ , and a segment which is bounded from  $\cup_{0 < j < i} \ell_j$  in  $H^2$ . So then either  $\cup_{0 \leq i < p} \ell_i \setminus A$  has bounded length in terms of  $n$ , or every point in  $\cup_{0 \leq i} \ell_i \setminus A$  is a bounded distance from some geodesic in  $H^3$ , which projects to the closed geodesic  $\gamma_*$  in  $N$  which is homotopic to  $f_0(\gamma)$ , with components of  $\cup_{0 \leq i < p} \ell_i \cap A$  having endpoints a bounded distance apart. This gives the required results on  $f_0(\gamma)$ .  $\square$

### 3.7 Recurrent and Minimal Laminations

A point  $x_0$  in a leaf of a lamination is *recurrent* if, for every open set  $U$  containing  $x_0$ , and every  $L > 0$ , there are points  $x, y \in U$  in the same leaf of the lamination and distance  $> L$  apart along that leaf. The set of recurrent points of a lamination is nonempty and closed, as is the case for any compact dynamical system. In fact, all nonisolated points of a geodesic lamination on a finite type hyperbolic surface are recurrent.

A geodesic lamination  $\mu$  on  $S$  is *minimal* if given  $\varepsilon > 0$  there is  $L > 0$  such that for every pair of points  $x, y$  in the lamination, there is a point  $z$  on the same leaf as  $y$ , and distance  $\leq L$  along the leaf from  $y$ , such that the hyperbolic distance between  $x$  and  $z$  is  $< \varepsilon$ . Geodesic laminations are exceptional dynamical systems in that a geodesic lamination is necessarily minimal if it is recurrent and intersects any simple closed geodesic transversally. This is shown in [20], where the language is of measured foliations, but measured foliations and geodesic laminations are basically equivalent concepts. In fact, for measured foliations, minimality and arationality are exactly the same, if minimality is defined in the right way. The method of proof is to endow a recurrent lamination with a finite transverse invariant measure, apply Poincaré recurrence, and then analyse the ways in which return can occur. For a geodesic lamination or measured foliations, the ways in which a return can occur are pretty restricted. A related exceptional property of geodesic laminations is that any recurrent lamination is a union of finitely closed geodesics and minimal laminations on subsurfaces with boundary.

The only recurrent leaves in the lamination in 3.2 are the closed geodesics in  $\Gamma$ .

### 3.8 Intersection number.

Any lamination can be given a finite transverse invariant measure, which is then automatically supported on the recurrent set. If the lamination is minimal, then the support is full. Even if the lamination is minimal, there may be more than one transverse invariant measure up to scalar multiple, but the space of measures is finite dimensional. The simplest examples of measured laminations are simple closed geodesics, or disjoint unions of simple closed geodesics. If a geodesic lamination  $\mu$  is a single closed geodesic then one can assign a finite transverse measure to  $\mu$  by saying that the measure of a traversal  $I$  is  $\#(I \cap \mu)$ . One can of course also do this if  $\mu$  is a finite disjoint union of closed simple geodesics  $\mu_i$ ,  $1 \leq i \leq n$ . More generally one can assign weight  $m_i$  to each  $\mu_i$ ,

and can define a finite transverse invariant measure by defining the measure of a transversal  $I$  to be

$$\sum_{i=1}^n m_i \#(I \cap \mu_i).$$

A geodesic lamination with a transverse invariant measure is called a *measured geodesic lamination*. An *intersection number*  $i(\mu_1, \mu_2)$  is defined for each pair of measured geodesic laminations  $\mu_1, \mu_2$ . If  $\mu_1$  and  $\mu_2$  are simple closed geodesics with transverse measure assigning weight one to each intersection  $\mu_i \cap I$ , then  $i(\mu_1, \mu_2) = \#(\mu_1 \cap \mu_2)$  or 0 if  $\mu_1 = \mu_2$ . We extend this linearly to the case when  $\mu_j$  is a disjoint union of simple closed geodesics for  $j = 1, 2$ . If a disjoint union of simple closed geodesics is being considered as a measured geodesic lamination, we shall always take each of the geodesics to have weight one, unless otherwise stated. More generally, if  $\mu_1$  is any measured geodesic lamination on  $S$  and  $\mu_2$  is still a simple closed geodesic, then  $\mu_1$  and  $\mu_2$  are either disjoint, or  $\mu_2$  is contained in  $\mu_1$ , or  $\mu_2$  is transverse to  $\mu_1$ . In the first two cases,  $i(\mu_1, \mu_2) = 0$ . In the last case,  $i(\mu_1, \mu_2)$  is the measure of the transversal  $\mu_2$  with respect to  $\mu_1$ . This then generalises easily to the case when  $\mu_2$  is a finite disjoint union of weighted simple closed geodesics. In the general case,  $i(\mu_1, \mu_2)$  can be defined using a partition into product rectangles, at least for transverse minimal laminations. But one can also note that the set of measured geodesic laminations has the structure of a piecewise linear manifold [20], using a natural correspondence between the space of measured geodesic laminations  $\mathcal{MGL}(S)$  and the space of measured foliations  $\mathcal{MF}(S)$  of [20], for which the transverse invariant measures are equivalent to Lebesgue measure. The map  $i(\mu_1, \mu_2)$  is uniformly Lipschitz for  $\mu_2$  a finite disjoint union of weighted simple closed geodesics, and so has a unique continuous Lipschitz extension  $\mu_2$  being any measured geodesic lamination (1.10 of [51]).

The natural projection from measured geodesic laminations with nonzero measure to recurrent geodesic laminations is continuous, with respect to the piecewise linear manifold structure on measured geodesic laminations mentioned above and the Hausdorff topology on geodesic laminations. It is not a bijection, trivially so since any transverse measure can be scaled. The natural map from projective measured laminations to geodesic laminations is also to a bijection, even on the inverse image of minimal laminations. Nevertheless, restricted to minimal measured geodesic laminations, the relation described by  $\mu_1 \sim \mu_2$  and only if  $i(\mu_1, \mu_2) = 0$ , is an equivalence relation, and all elements of the equivalence class are the same minimal lamination.

### 3.9 Geodesic laminations, measured foliations and the compactification of Teichmüller space.

There is a topology on the union of Teichmüller space  $\mathcal{T}(S)$  and the space  $\mathcal{PMF}(S)$  of projective measured foliations — or, equivalently, the space  $\mathcal{PMGL}(S)$  of projective measured geodesic laminations — which makes this space compact, homeomorphic to a closed ball, such that the subspace topology



on  $\mathcal{T}(S)$  is the usual topology on  $\mathcal{T}(S)$ , and  $\mathcal{PMF}(S)$  has the piecewise linear topology referred to above [20]. With respect to this topology  $\mathcal{PMF}(S)$  is known as the *Thurston boundary*. The topology is obtained from projectivising an embedding of  $\mathcal{T}(S) \cup \mathcal{MF}(S)$  in  $\mathbb{R}_+^S$  where  $S$  is the set of simple closed nontrivial nonperipheral loops on  $S$ , and the embedding is

$$[\varphi] \mapsto (|\varphi(\gamma)|) \quad ([\varphi] \in \mathcal{T}(S)),$$

$$\mu \mapsto i(\mu, \gamma) \quad (\mu \in \mathcal{MF}(S)).$$

We shall never make direct use of this topology, but it has the property that if  $[\varphi_n]$  is a sequence in  $\mathcal{T}(S)$  converging to a projective class of an arational measured lamination  $\mu$  and  $\gamma_n$  is a sequence of simple closed loops such that  $|\varphi_n(\gamma_n)| \leq L$  for all  $n$ , then the recurrent part of any Hausdorff limit of  $\gamma_n$  is  $\mu$ , and  $i(\gamma_n, \mu)/|\gamma_n| \rightarrow 0$  as  $n \rightarrow \infty$ .

There is a related boundary of  $\mathcal{T}(S)$ , which is of more direct relevance, in which the boundary points are either arational geodesic laminations - with no measure - or in a set  $\mathcal{X}(\Gamma)$  for some multicurve  $\Gamma$  on  $S$ . Let  $\Sigma$  be the set of components of  $S \setminus \Gamma$  which are not topologically 3-holed spheres. If  $\Sigma = \emptyset$  then  $\mathcal{X}(\Gamma)$  is a single point. Otherwise, the elements of  $\mathcal{X}(\Gamma)$  are of the form  $(\mu_1, \dots, \mu_m)$ , where the elements of  $\Sigma$  are numbered,  $\Sigma = \{\alpha_i : 1 \leq i \leq m\}$  and each  $\mu_i$  is either in  $\mathcal{T}(S(\alpha_i))$  or an arational geodesic lamination on  $S(\alpha_i)$ . Here,  $S(\alpha_i)$  is as in 2.6. Convergence of a sequence  $[\varphi_n]$  in  $\mathcal{T}(S)$  to  $(\mu_1, \dots, \mu_m) \in \mathcal{X}(\Gamma)$  (including  $\Gamma = \emptyset$ ) is then defined as follows. We must have  $|\varphi_n(\Gamma)| \rightarrow 0$  as  $n \rightarrow \infty$ , which of course is an empty condition if  $\Gamma = \emptyset$ . We also have  $\pi_{\alpha_i}([\varphi_n]) \rightarrow \mu_i$  as  $n \rightarrow \infty$  if  $\mu_i \in \mathcal{T}(S(\alpha_i))$ , and, if  $\mu_i$  is an arational lamination on  $S(\alpha_i)$ , any limit of  $\pi_{\alpha_i}([\varphi_n])$  in the Thurston compactification  $\mathcal{T}(S(\alpha_i)) \cup \mathcal{PMGL}(S(\alpha_i))$  is  $\mu_i$ , endowed with some transverse measure.

### 3.10 The Masur Domain.

Let  $S \subset N$  be an embedded surface. The *Masur domain*  $\mathcal{O}(S, N)$  according to the original definition ([45], [16], [25]) is a set of measured geodesic laminations, but in fact the property of being in the Masur domain is independent of the transverse invariant measure chosen, simply because a statement  $i(\mu, \mu') = 0$  about measured geodesic laminations is purely topological, provided each minimal component of  $\mu, \mu'$  is in the support of the measure.

The definition we shall use is:  $\mu \in \mathcal{O}(S, N)$  if and only if there is a constant  $c = c(\mu) > 0$  such that the following holds, for any transverse invariant measure on  $\mu$ . For each simple nontrivial  $\gamma \subset S$  which bounds a disc in  $N$ , and any geodesic laminations  $\mu'$  and  $\mu''$  with normalised transverse invariant measures,

$$i(\mu, \mu') \geq c \text{ whenever } i(\mu'', \mu') = 0 \text{ and } i(\gamma, \mu'') \leq c|\gamma|. \quad (3.10.1)$$

In particular, if we take  $\mu'' = \mu' = \gamma/|\gamma|$ , we obtain

$$i(\mu, \gamma) \geq c(\mu)|\gamma|. \quad (3.10.2)$$

For large compression bodies, as they are called, (3.10.2) actually implies (3.10.1), for a different constant  $c(\mu)$  in (3.10.1) from that in (3.10.2). It also does so for arational geodesic laminations in all cases. This is simply because, if  $i(\mu, \mu') = 0$  and  $\mu$  is arational,  $\mu' = \mu$ , and because intersection number is uniformly Lipschitz in each variable. According to [25], (3.10.2) can be weakened further for arational laminations, and the term  $|\gamma|$  can be omitted.

### 3.11 Bonahon's far-out pleated surfaces and the invariants.

Let  $e$  be any end of  $N_d$ , and  $S = S(e)$  the corresponding surface as in Section 1. It was shown by Bonahon [5] in the case of  $S_d$  incompressible, and by Canary [16] in the compressible (tame) case, that there is a sequence  $\zeta_n$  of simple closed geodesics on  $S$  with  $\zeta_n$  nontrivial nonperipheral in  $N$  such that if  $(\zeta_n)_*$  is the geodesic in  $N$  freely homotopic to  $\zeta_n$  then  $\zeta_n^*$  converges to  $e$  in  $N$ . By extending  $\zeta_n$  to a maximal multicurve, possibly with some boundary components of zero length, there is a pleated surface  $g_n$  in the homotopy class of inclusion of  $S$  in  $N$  with pleating locus including  $\zeta_n$ . There are only finitely many Margulis tubes intersecting any compact set, and given any Margulis tube  $T$ , one can find a neighbourhood of  $e$  disjoint from  $T$  — the same proof as in 3.4. So by the Bounded Diameter and Injectivity Radius Lemmas in 3.3,  $g_n(S) \rightarrow e$ .

In the compressible case,  $g_n$  still exists with  $g_n(S) \rightarrow e$ . The following lemma shows the existence of a pleated surface with pleating locus including  $\zeta_n$ . Every short loop on this pleated surface which is nontrivial nonperipheral in  $S(g_n)$  is also nontrivial nonperipheral in  $N$  by 3.4, and then  $g_n(S) \rightarrow e$  as in the incompressible case.

**Lemma** *Any nontrivial nonperipheral simple loop  $\gamma$  on  $S$  which is nontrivial in  $N$  is contained in a noncollapsing maximal multicurve.*

*Proof.* If not, we have a multicurve  $\Gamma$  containing  $\gamma$ , and such that some complementary component  $Q$  of  $S \setminus (\cup \Gamma)$  is not a pair of pants, and such that every simple closed loop in the interior of  $Q$ , which is not homotopic to a boundary component, is trivial in  $N$ , but the loops of  $\Gamma$  are nontrivial in  $N$ . Take any pair  $\gamma_1, \gamma_2$  of boundary components of  $Q$ , positively oriented with as elements of  $\partial Q$ . Then  $\gamma_1 * \gamma_2$  is trivial. If this is true for all pairs then there are at most two boundary components, and  $Q$  must have genus at least one. Except in the case when  $Q$  has genus one and one boundary component, we can find two simple loops  $\zeta_1, \zeta_2$  in  $Q$  which are not homotopic to the boundary but such that  $\zeta_1 * \zeta_2$  is homotopic to a boundary component, which must then be trivial, a contradiction. If  $Q$  is a torus with one boundary component we can again find  $\zeta_1, \zeta_2$  on  $S$  such that  $\zeta_1 * \zeta_2 * \zeta_1^{-1} * \zeta_2^{-1}$  is trivial, again a contradiction.  $\square$

Bonahon and Canary also showed, in the incompressible and compressible cases respectively, that any such sequence  $\zeta_n$  converging to  $e$  converged in the Hausdorff topology to a geodesic lamination  $\mu$ . It was shown further (Section 5 of [5] and 10.1 of [16]) that  $\mu$  was minimal, and, Canary showed that  $\mu$  is in the Masur domain (10.2 of [16]). His proof was said to be for the non-cusp case only but does in fact work in general. The proofs in [5], which are also part of

the argument in [16], are inextricably linked with the proof of the main result, the existence of the sequence  $\zeta_n$ , and thus very delicate.

We shall deal with the case of geometrically finite ends in Section 4.

## 4 More on pleated surfaces.

### 4.1 Removing badly bent annuli.

The following lemma shows that it is possible to avoid badly bent annuli.

**Lemma** *The following holds for a suitable constant  $D_0$  and, given any integer  $k$  and Margulis constant  $\varepsilon_0$ , an integer  $r(\varepsilon_0, k)$ . Let  $f : S \rightarrow N$  be continuous. Let  $\gamma$  be a simple nontrivial nonperipheral loop on  $S$  such that  $f(\gamma)$  is nontrivial with  $|\gamma_*| < \varepsilon_0$ . We allow  $f(\gamma)$  to be a cusp. Let  $\zeta$  be a simple loop with  $\leq k$  intersections with  $\gamma$ . Suppose that the loop between each two consecutive returns of  $\zeta$  to  $\gamma$  is not homotopic in  $N$  a multiple of  $\gamma$ . Let  $\zeta_n = \tau_\gamma^n(\zeta)$*

*Then  $|(\zeta_n)_*| < \varepsilon_0$  for at most  $r$  integers  $n$ , which can be taken to be consecutive. Indeed,  $r$  can be chosen so that  $\zeta_n$  intersects  $T(\gamma_*, \varepsilon_0)$  for all but an interval of  $r$  consecutive  $n$ , and  $(\zeta_n)_* \neq \gamma_*$ .*

*A similar result holds if  $\gamma$  is a multicurve  $(\gamma_1, \dots, \gamma_s)$ ,  $|(\gamma_i)_*| < \varepsilon_0$  for all  $i$ , all the  $(\gamma_i)_*$  are distinct, and the loop between each two consecutive returns of  $\zeta$  to  $\gamma_i$  is not homotopic in  $N$  a multiple of  $\gamma_i$ . In this case, write  $\tau_{\underline{n}} = \tau_{\gamma_1}^{n_1} \circ \dots \circ \tau_{\gamma_s}^{n_s}$ ,  $\underline{n} = (n_1, \dots, n_s)$ . Then the lower bound on  $|(\tau_{\underline{n}}(\zeta))_*|$ , and the intersection with the  $T((\gamma_i)_*, \varepsilon_0)$ , hold for each  $n_i$  excluded from an interval of length  $\leq r(\varepsilon_0, k)$ .*

*Proof.* We use the following fundamental fact. There are constants  $C_d$  and  $\Delta_d$  such that the following holds. Let  $\gamma'_i$  be any sequence of geodesics in  $H^d$  such that positive end on  $\partial H^d$  of  $\gamma'_i$  coincides with the negative end of  $\gamma'_{i+1}$ , and such that there is a segment on  $\gamma'_i$  of length  $\geq \Delta_d$  which is distance  $\geq 1$  from both  $\gamma'_{i+1}$  and  $\gamma'_{i-1}$ . Then the union of the  $\gamma'_i$  comes within distance  $\leq C_d$  of a unique geodesic, on each segment of each  $\gamma'_i$  which is distance  $\geq 1$  from both  $\gamma'_{i-1}$  and  $\gamma'_{i+1}$ . Also  $\gamma'_i$  comes within  $\varepsilon_d$  of the geodesic on this segment, if the segment has length  $\geq \Delta'_d$ , where  $\varepsilon_d \rightarrow 0$  as  $\Delta'_d \rightarrow \infty$ .

Up to free homotopy in  $N$ , we can make  $\zeta$  out of  $k$  infinite geodesics, such that the positive and negative ends of each one are asymptotic to  $\gamma$ . This gives a bi-infinite sequence of geodesics in  $H^3$  which we call  $\gamma_i$  such that the positive end of  $\gamma_i$  is asymptotic to the negative end of  $\gamma_{i+1}$ , and this common endpoint is an endpoint of a lift of  $\gamma$ . The endpoint is fixed by  $g_i$ , whose conjugacy class represents  $\gamma$ . The sequence for a lift of  $\zeta_n$  is obtained by taking as an adjacent pair  $h_{i,n} \cdot \gamma'_i$  and  $h_{i,n} \cdot g_i^n \cdot \gamma'_{i+1}$ , where  $h_{i,n} = g_1^n \cdot \dots \cdot g_{i-1}^n$  if  $i \geq 1$ , with modifications if  $i < 1$ .

Distinct Margulis tubes, and components of  $N_{<\varepsilon_0}$  round cusps, are a definite distance apart. So the hypothesis on consecutive returns means the following. Let  $d$  denote hyperbolic distance in  $H^3$ . Let  $X_i$  be the set of points  $x$  on  $\gamma_i$  with  $d(g_i \cdot x, x) \leq \varepsilon_0$ , and  $X'_i$  the set of points on  $\gamma_{i+1}$  with  $d(g_i \cdot x, x) \leq \varepsilon_0$ . Then the distance between  $\cup_n g_i^n \cdot (X_i \cup X'_i)$  and  $\cup_{j \neq i} \cup_n g_j^n \cdot (X_j \cup X'_j)$  is  $\geq C(\varepsilon_0) > 0$ , where  $C(\varepsilon_0)$  can be taken arbitrarily large by choice of sufficiently small  $\varepsilon_0$ . So,

from considering  $j = i + 1$ , the biinfinite sequence for  $\zeta_n$  is a distance  $\leq C_3$  from a lift of the geodesic representing it provided that the sets  $\gamma_i \setminus X_i$  and  $g_i^n(\gamma_{i+1} \setminus X_i')$  are a distance  $\geq (2\Delta_3 - C_3)$  apart. Since points of  $g_i^n(\gamma_{i+1} \setminus X_i')$  and  $g_i^m(\gamma_{i+1} \setminus X_i')$  are a distance  $\geq |m - n|\varepsilon_0$  apart, this is true for all but a bounded interval of  $n$ . For the statement about  $\zeta_n$  entering  $T(\gamma_*, \varepsilon_0)$ , we simply need  $\gamma_i \setminus X_i$  and  $g_i^n(\gamma_{i+1} \setminus X_i')$  somewhat further apart.

The statement with  $\gamma$  replaced by a multicurve is proved in exactly the same way.  $\square$

**Corollary 4.2.** *The following holds for a suitable constant  $L_0$ ,  $L_1$ , and integers  $k_0$  and  $r$ . Here,  $k_0$ ,  $r$  and  $L_1$  depend on  $L_0$ . Suppose that  $f : S \rightarrow N$  is a pleated surface, homotopic to an embedding. Let  $\Gamma \subset S$  be a maximal multicurve such that for any simple nontrivial loop  $\gamma' \subset S$  with  $\leq D_0$  intersections with  $\Gamma$ ,  $f(\gamma')$  is nontrivial in  $N$ . Let  $|\varphi(\Gamma)| \leq L_0$  for some  $[\varphi] \in \mathcal{T}(S)$ .*

*Then we can choose  $\Gamma'$  with  $|\varphi(\Gamma')| \leq L_1$  and  $\leq r$  intersections with  $\Gamma$  such that  $|\zeta_*| \geq \varepsilon_0$  for all  $\zeta \in \Gamma'$ .*

*Proof.* Let  $\Gamma'' = \{\gamma_i : 1 \leq i \leq s\}$  be the set of loops in  $\Gamma$  for which the corresponding geodesics in  $N$  which are of length  $< \varepsilon_0$ . If  $\Gamma'' = \emptyset$ , there is nothing to prove. Otherwise  $\Gamma'$  be a multicurve such that  $\Gamma''$  does not intersect  $\Gamma \setminus \Gamma'$ , and such that each loop of  $\Gamma'$  is intersected by a loop of  $\Gamma''$ , and  $\Gamma'' \cup (\Gamma \setminus \Gamma')$  is a maximal multicurve. We can choose  $\Gamma'$  so that  $|\varphi(\Gamma')| \leq L'_0$ , where  $L'_0$  depends only on  $L_0$ . Then  $\#(\Gamma' \cap \Gamma'')$  is bounded in terms of  $L'_0$ . We only need to show that the other hypotheses of 4.1, about distinct Margulis tubes and nontrivial loops, hold for the multicurve  $(\gamma_1, \dots, \gamma_s)$  and each  $\zeta \in \Gamma''$ , if  $k_0$  is sufficiently large. If Margulis tubes round  $(\gamma_i)_*$  and  $(\gamma_j)_*$  coincide, where  $\gamma_i$  and  $\gamma_j$  are both intersected by  $\zeta$ , with an arc  $\zeta_1$  from  $\gamma_i$  to  $\gamma_j$ , then consider the loop  $[\gamma_i * \zeta_1 * \gamma_j * \overline{\zeta_1}]$ . This loop is nontrivial in  $S$  but trivial in  $N$ , and has a bounded number of intersections with  $\Gamma$  with bound in terms of  $L_0$ . By the Loop Lemma, we can then find a simple loop with these properties. The same argument works if  $i = j$ : a closed loop formed from  $\zeta_1$  by adding an arc along  $\gamma_i$  is a multiple of  $\gamma_i$ .  $\square$

### 4.3 Bounded homotopy distance between pleated surfaces.

In this subsection we rework Minsky's remarkable estimate in 4.1 and 4.2 of [37] on the distance between two pleated surfaces related by elementary moves. Note that we do not require our pleating loci to be related by elementary moves, just to have bounded intersections. We also drop the condition of injective-on- $\pi_1$ , although, of course, we do need something to replace it. By 4.2, the assumption of no badly bent annuli on one of the surfaces is not much of a restriction. Furthermore, we refine the concept of badly bent annuli that was used in Section 3. Let  $C = C(\varepsilon_0)$  be the constant of the Radius of Injectivity Lemma of 3.3. Given homotopic pleated surfaces  $f_0, f_1 : S \rightarrow N$ , a *badly bent annulus* for  $(f_0, f_1)$  is a Margulis tube  $T(\gamma, \varepsilon(\gamma)) \subset S(f_0)$  round a loop  $\gamma$  which is in the pleating locus of  $f_0$ , such that either  $\varepsilon(\gamma) = \varepsilon_0$ , or both components of  $\partial T(\gamma, \varepsilon(\gamma))$  are mapped by  $f_0$  to within  $2C$  of the same point on a single

geodesic in the pleating locus of  $f_1$ . Moreover, the latter is not true for  $\varepsilon(\gamma)$  replaced by  $\varepsilon(\gamma)/C$ .

**Lemma** *The following holds given an integer  $r$  for any Margulis constant  $\varepsilon_0 > 0$ ,  $C > 1$  as in 3.4, and some constants  $D_0$  and  $L(\varepsilon_0, r)$ . Let  $N$  be any complete hyperbolic 3-manifold. Let  $\Gamma_0, \Gamma_1$  be noncollapsing maximal multicurves on  $S$  with  $\#(\Gamma_0 \cap \Gamma_1) \leq r$ . Let  $f_0, f_1 : S \rightarrow N$  be homotopic pleated surfaces with pleating loci  $\Gamma_0$  and  $\Gamma_1$  respectively. Suppose that there are no badly bent annuli for one of  $(f_0, f_1)$ ,  $(f_1, f_0)$ . Let  $A \subset S(f_1)$  (or  $A \subset S(f_0)$ ) be the union of badly bent annuli for  $(f_1, f_0)$  (or  $(f_0, f_1)$ ). Suppose that for any homotopically nontrivial loop  $\gamma \subset S$  with  $f_0(\gamma)$  trivial in  $N$ ,  $|f_0(\gamma)| \geq D_0$  (or  $|f_0(\gamma) \setminus A| \geq D_0$ ).*

*Then composing  $f_1$  on the right with a homeomorphism isotopic to the identity if necessary, there is a homotopy  $(x, t) \mapsto f_t(x) : S \times [0, 1] \rightarrow N$  between  $f_0$  and  $f_1$  whose homotopy tracks have length  $\leq L(\varepsilon_0, r)$  outside  $f_1^{-1}(A) \times [0, 1]$  (or  $f_0^{-1}(A) \times [0, 1]$ ). Moreover, the image under the homotopy of  $f_j^{-1}(A) \times [0, 1]$  is contained in the union of the corresponding Margulis tubes, and the image under  $f_0$  of each component of  $f_1^{-1}(A)$  (or under  $f_1$  of each component of  $f_0^{-1}(A)$ ) has bounded diameter.*

This lemma is true without any restriction on the length of loops in  $(\Gamma_0)_*$ ,  $(\Gamma_1)_*$ . The assumption on no badly bent annuli for one of  $(f_0, f_1)$ ,  $(f_1, f_0)$  is not really necessary. It simplifies the proof slightly – and even the statement. A general statement and proof can be deduced by applying the statement as given, twice.

*Proof.*

We start by assuming that there are no badly bent annuli for  $(f_0, f_1)$ .

Using the version of Efficiency of pleated surfaces which has been reformulated from Minsky's version, and proved, in 3.6, we see that, for a suitable  $C_0 = C_0(r, \varepsilon_0)$  and  $C_1(r, \varepsilon_0)$  as in 3.6, for any  $\gamma \in \Gamma_1$  such that  $|f_1(\gamma)| \geq \varepsilon_0/C$ ,  $f_0(\gamma)$  is, up to homotopy in  $S(f_0)$ , a distance  $\leq C_0$  from  $\gamma_* = f_1(\gamma)$  and

$$||f_0(\gamma)| - |\gamma_*|| \leq C_1.$$

If  $|\gamma_*| \leq \varepsilon_0/C$ , we first deduce that  $|f_0(\gamma)| \leq C_0$  and then, by considering a loop  $\zeta$  with nonzero intersection with  $\gamma$ , and  $\leq r$  intersections with  $\Gamma_0 \cup \Gamma_1$ , if  $A(\gamma, f_1) = \emptyset$ ,

$$|f_0(\gamma)| \leq C_1|\gamma_*|.$$

Let  $A(\gamma)$  denote the badly bent annulus for  $(f_1, f_0)$  round  $\gamma$ , if this exists. If the modulus in  $S(f_1)$  of either component of  $T(f_1(\gamma), \varepsilon_0) \setminus A$  is  $L$ , and  $\varepsilon$  is  $\text{Min}(1/L, 1)$ , then

$$|f_0(\gamma)| \leq C_1\varepsilon.$$

Regarding the domain of  $f_1$  as being  $S(f_0)$  and composing on the right with a homeomorphism isotopic to the identity if necessary, we can assume that  $f_1(\gamma) = \gamma_*$  for all  $\gamma \in \Gamma_1$ . So then we have a bound  $C_2(r)$  on the distance between  $f_0(\gamma)$  and  $f_1(\gamma)$  for all  $\gamma \in \Gamma_1$  with  $|\gamma_*| \geq \varepsilon_0$ . If  $|\gamma_*| \leq \varepsilon_0$ , then we have a bound on  $|f_0(\gamma)|$ , and on the distance between  $f_0(\gamma)$  and  $f_1(A(\gamma))$ , and between  $f_0(\gamma)$  and  $f_1(\gamma)$  if there is no badly bent annulus round  $\gamma$ . Now we

are going to regard the homotopy as having domain  $S(f_0) \times [0, 1]$ . So far, the homotopy has been constructed on  $f_0(\Gamma_1)$ . We can extend it to the boundary of a bounded modulus annulus round  $f_0(\gamma)$  for each  $\gamma \in \Gamma_1$  for which  $A(\gamma)$  exists. Let  $A'$  be the union of these annuli in  $S(f_0)$ . The homotopy tracks on  $f_0^{-1}((f_0(\Gamma_1) \setminus A') \cup \partial A') \subset S$  are bounded, composing  $f_1$  on the right with a homeomorphism isotopic to the identity if necessary, so that the image under  $f_1$  in  $S(f_1)$  is  $(f_1(\Gamma_1) \setminus A) \cup \partial A$ . Let  $A''$  be the union of components of  $(S(f_0))_{<\varepsilon_0} \setminus A'$  homotopic to loops  $f_0(\gamma)$  for which  $\gamma \in \Gamma_1$ ,  $|f_1(\gamma)| = |\gamma_*| < \varepsilon_0/2C$ . Here,  $C$  is, again, the constant of 3.4. The corresponding components of  $(S(f_1))_{<\varepsilon_0} \setminus A$  have moduli which are boundedly proportional. The sets  $f_0(S(f_0))$  and  $f_1(S(f_1))$  are a bounded distance apart in  $N$ . So the homotopy can be extended, with bounded tracks, to  $f_0^{-1}((f_0(\Gamma_1) \setminus A') \cup A'') \subset S$ , composing  $f_1$  on the right with a homeomorphism isotopic to the identity if necessary, so that the image under  $f_1$  in  $S(f_1)$  is

$$f_1(\Gamma_1) \setminus \cup \{T(f_1(\gamma), \varepsilon_0) : \gamma \in \Gamma_1, |\gamma_*| < \varepsilon_0/2C\}.$$

So now, after suitably right-composing  $f_1$  the homotopy with bounded tracks is defined on a set whose complement of this set consists of the annuli  $A'$  and a union of pairs of pants whose images under  $f_0$  in  $S(f_0)$ , and under  $f_1$  in  $S(f_1)$ , have boundaries which have length  $\geq \varepsilon_0/2C$ , and similarly for the images in  $N$ . So we now need to define the homotopy on these pairs of pants. For this, we again follow Minsky [37], but care is needed, because we can only apply the Short Bridge Arc Lemma 3.5 to  $f_0$ , not to  $f_1$ . So, for  $j = 0, 1$ , we foliate the pairs of pants in  $S(f_j)$  by arcs between boundary components, of bounded length, apart from a tripod, of which the arms are of bounded length, one arm ending on each boundary component, the other arms meeting at the centre of the tripod, as we shall call it. Map the foliation plus tripod into  $N$  by  $f_j : S(f_j) \rightarrow N$ . Lift to the universal cover. The images are a bounded distance from geodesics in  $N$  with lifts which are geodesics in  $H^3$ . These three geodesics in  $H^3$  come within a bounded distance of the lift of the image of the centre of the tripod defined using  $S(f_0)$ , and similarly for the tripod defined using  $S(f_1)$ . We claim that for any  $C_2 > 0$  the set of points which are distance  $\leq C_2$  from all three of these geodesics has diameter bounded in terms of  $C_2$ , if  $D_0$  is large enough. This will imply that the images of the tripod are a bounded distance apart, and the homotopy can be extended to match up the images of the foliations. To prove the claim, apply the short Bridge Arc Lemma 3.5 to each of the pairs of geodesics or components of  $\partial(S(f_0))_{<\varepsilon_0}$  in  $S(f_0)$ . The set of points on each pair whose lifts are a bounded distance  $\leq C_2$  apart in  $H^3$  corresponds, up to bounded distance, to the set of points whose lifts in the universal cover  $H^2$  of  $S(f_0)$  are a distance  $\leq C_2$  apart, for  $C_2$  sufficiently large (but universally bounded). In  $H^2$ , the set of points which are distance  $\leq C_2$  from all three geodesics is nonempty and of diameter bounded in terms of  $C_2$ . So the set in  $H^3$  is bounded also, as required. So the images of the tripods are a bounded distance apart, and the homotopy can be extended to match up the images of the foliations, after suitable right-composition of  $f_1$ .

The proof when  $f_0$ , rather than  $f_1$ , is allowed to have badly bent annuli, is essentially exactly the same. The original homotopy is, after right-composition with a homeomorphism, between  $f_0$  on  $f_0^{-1}(f_0(\Gamma_1) \setminus A)$  and  $f_1$  on  $f_1^{-1}(f_1(\Gamma_1) \setminus A')$ , where  $A$  is the union of badly bent annuli round loops of  $f_0(\Gamma_0)$  in  $S(f_0)$  and  $A'$  is the corresponding set in  $S(f_1)$ . The set  $(f_0(\Gamma_1) \setminus A) \cup A''$  is then a union of pairs of pants minus rectangles in  $A$  between boundary components, with sides between boundary components being of bounded length. The foliations between boundary components can then be taken to include  $\partial A$  and to foliate  $A$ , but the homotopy does not extend with bounded tracks across  $f_0^{-1}(A) \times [0, 1]$ .  $\square$

#### 4.4 Bounded Teichmüller distance between pleated surfaces.

We have the following, which is an extension of 4.3 and essentially a reinterpretation of 4.1 and 4.2 of [37].

**Lemma** *Fix a Margulis constant  $\varepsilon_0$ . The following holds for a suitable constant  $L_0$ . Let  $N$ ,  $S$ ,  $\Gamma_0$ ,  $\Gamma_1$ ,  $f_0$  and  $f_1$  satisfy the hypotheses of 4.3.*

*Let  $\alpha$  be the surface which is the complement in  $S$  of loops whose images under  $f_0$  (or  $f_1$ ) are cores of  $A$ . Then*

$$|f_j(\partial\alpha)| \leq L_0$$

*for  $j = 0, 1$  and*

$$d_\alpha([f_0], [f_1]) \leq L_0.$$

*Proof.* There is a constant  $L'(\varepsilon_0)$  such that the following holds. We use the homotopy whose existence is given by 4.3 to relate the maps  $f_0$  and  $f_1$  pointwise (not just up to homotopy). For simplicity, we assume that there are no badly bent annuli for  $(f_0, f_1)$ . As a result of 4.3, we can, in any case, interchange  $f_0$  and  $f_1$  and get the first hypotheses of 4.3. For any path  $\zeta \subset \alpha$  and such that  $f_1(\zeta)$  is in the  $\varepsilon_0$ -thick part of  $S(f_1)$ , assuming that  $f_1(\zeta)$  is geodesic,

$$|f_0(\zeta)| \leq L'(\varepsilon_0)(|f_1(\zeta)| + 1). \quad (4.4.1)$$

We see this as follows. The image under  $f_1$  of  $(S(f_1))_{\geq \varepsilon_0}$  cannot intersect  $N_{< \varepsilon_0/C'}$ , for a suitably large  $C'$  depending only on the topological type of  $S$  and  $L(\varepsilon_0)$ , that is, only on the topological type of  $S(f_0)$ . For if it does so intersect, the image under the homotopy to  $f_0$  means that there is a map  $f_0 : U \rightarrow N_{\varepsilon_0/C}$ , where  $U \subset S(f_0)$  is connected and bounded and carries a nonabelian subgroup of  $\pi_1(S)$ , contradicting the assumptions on  $f_0$ . So  $f_1(\zeta)$  is in  $N_{\geq \varepsilon_0/C'}$ . Then  $f_0(\zeta)$  is in  $N_{\geq \varepsilon_0/C''}$  for a suitable  $C''$ . Then by the Injectivity Radius Lemma,  $f_0(\zeta)$  is contained in  $(S(f_0))_{\geq \varepsilon_0/C_3}$ , where, again, both  $C''$  and  $C_3$  depend only on the topological type of  $S$ . Then it suffices to show that if  $|f_1(\zeta)|$  is bounded, then  $|f_0(\zeta)|$  is also, with a less good bound. Suppose it is not so. Then we can find a large number of points in a single orbit under the covering group of  $S(f_0)$  within a bounded distance of the lift of  $f_1(\zeta)$ . In fact these points can be put in

a sequence  $\{x_n\}$  such that the distance between  $x_n$  and  $x_{n+1}$  is  $\leq C_1$ , where  $C_1$  depends only on the Margulis constant. Then, regarding this as a subset of the universal cover of  $N$ , we have a large number of points in a single orbit of the covering group of  $N$  within a bounded distance of the lift of  $f_0(\zeta)$  — which is a bounded set in the thick part of  $N_{\geq \varepsilon_0/C_3}$ . Then for some  $n$  depending only on the Margulis constant, and the various constants involved, that is, just on the Margulis constant and the topological type of  $S$ , two of the orbit points  $x_m$  and  $x_p$ , for  $m < p \leq m + n$ , must be identified in the universal cover of  $N$ , giving a nontrivial loop in  $f_0(S)$  which is trivial in  $N$  and of length  $\leq D_0$ , assuming  $D_0$  is large enough given the Margulis constant and the topological type of  $S$ , which contradicts our hypothesis (stated at the start of 4.3).

Now if  $|f_1(\gamma)| \leq \varepsilon/2$ ,  $f_1(\gamma)$  can be realised up to free homotopy by the union of two geodesic segments of bounded length in  $(S(f_1))_{\geq \varepsilon_0/2}$ , by taking segments with endpoints in  $\partial(S(f_1))_{\geq \varepsilon_0}$ . So then  $f_0(\gamma)$  has length bounded above, and  $\leq D_0$  for  $D_0$  large enough. So then  $f_0(\gamma)$  is nontrivial. So we now assume that the Injectivity Radius Lemma of 3.3 holds for both  $f_0$  and  $f_1$ . Then the bounded homotopy given by 4.3 implies that for some  $C_4$ , if  $|f_0(\gamma)|, |f_1(\gamma)| < \varepsilon_0/C_4$ , then both have length  $< \varepsilon_0/2$ , and the images under  $f_0, f_1$ , of the boundaries of a Margulis tube round  $\gamma$  in  $S(f_0), S(f_1)$  are a bounded distance apart. This includes loops of  $\partial\alpha$ . It follows that, for a suitable constant  $C_5$ , if  $\gamma$  is not homotopic to  $\partial\alpha$ , the shortest paths in  $(S(f_j))_{< \varepsilon_0}$  between boundary components differ by a constant, and hence

$$\frac{1}{C_5}|f_0(\gamma)| \leq |f_1(\gamma)| \leq C_5|f_0(\gamma)|.$$

It then follows that we can extend (4.4.1) to all closed paths in  $S(f_1)$ , that is, for a constant  $C_6$ , whenever  $f_1(\zeta)$  is a closed geodesic,

$$|f_0(\zeta)| \leq C_6|f_1(\zeta)|. \quad (4.4.2)$$

It now follows that if  $D_0$  is large enough, given  $D_1$  depending only on the topological type of  $S$ , for every nontrivial loop  $\gamma$  in  $\alpha \subset S(f_1)$  of length  $\leq D_1$ ,  $f_1(\gamma)$  is nontrivial in  $N$ . Now that we know that  $|f_0(\partial\alpha)| \leq C_5\varepsilon_0$ , we can apply the above arguments with  $f_1$  and  $f_0$  reversed, and we obtain (4.4.2) with  $f_0$  and  $f_1$  interchanged.

Now to bound  $d_\alpha([f_0], [f_1])$  we want to use 2.7. This means that we need to bound  $|f_1(\zeta)|'/|f_0(\zeta)|'$  for a cell-cutting set of loops  $\zeta$  for which  $|f_0(\zeta)|''$  is bounded. Here,  $|\cdot|'$  and  $|\cdot|''$  are as in 2.7. So  $|f_0(\zeta)|' = |f_0(\zeta)|'' = |f_0(\zeta)|$  unless  $|f_0(\zeta)| < \varepsilon_0$  or  $\zeta$  is transverse to a loop  $\gamma$  for which  $|f_0(\gamma)| < \varepsilon_0$ . If  $|f_0(\zeta)| < \varepsilon_0$  then  $|f_0(\zeta)|'$  is boundedly proportional to  $\sqrt{|f_0(\zeta)|}$  and we already have the bound on  $|f_1(\zeta)|/|f_0(\zeta)|$ , which suffices. So it remains to obtain an estimate for one  $\zeta$  crossing each loop  $\gamma$  which is not the core of a badly bent annulus for  $(f_1, f_0)$ , with  $|f_0(\gamma)| < \varepsilon_0$ ,  $\gamma$  and with  $|f_0(\zeta)|''$  bounded. For such a loop, by the definition of  $|\cdot|'$ ,  $|f_0(\zeta)|'$  is boundedly proportional to  $\exp|f_0(\zeta)|/2$  and to  $1/\sqrt{|f_0(\gamma)|}$ . If we can show that  $\exp|f_1(\zeta)|$  is boundedly proportional to  $1/|f_1(\gamma)|$ , then  $|f_1(\zeta)|'$  is also boundedly proportional to  $1/|f_1(\gamma)|$  and the



result follows. So fix  $\zeta$ . We write  $f_0(\zeta)$  as a union of finitely many components: one or two components in  $(S(f_0))_{\geq \varepsilon_0}$ , not necessarily in the pleating locus of  $f_0$ , but of bounded length, and four or two components — which are in the pleating locus — in the Margulis tube  $T(f_0(\gamma), \varepsilon_0)$ , depending on whether or not  $\gamma$  is in the pleating locus of  $f_0$ . We make a similar decomposition of  $f_1(\zeta)$ . The components in  $(S(f_1))_{\geq \varepsilon_0}$  are again bounded, by 4.4.1 (or 4.4.2).

First, suppose that for at least one of  $j = 0, 1$ ,  $\gamma$  is not in the pleating locus of  $f_j$ . Then these unions of geodesic segments are a bounded distance from the geodesic  $\zeta_*$  in  $N$ , and from the geodesic representatives in  $S(f_0)$ ,  $S(f_1)$  respectively, assuming that  $\varepsilon_0$  is small enough. This uses the fact mentioned in 4.1. The long geodesic segments used here are bounded apart along most of their length, in the lift of  $N$  to  $H^3$ , and the lifts of  $S(f_0)$  and  $S(f_1)$  to  $H^2$ , because the different lifts of Margulis tubes are distinct. So this means we have a bound on  $|f_j(\zeta)| - |\zeta_*|$  for  $j = 0, 1$ , and hence a bound on  $\exp |f_1(\zeta)| / \exp |f_0(\zeta)|$ . Since we already know that  $|f_1(\gamma)| / |f_0(\gamma)|$  is bounded, the result follows. Now suppose that  $\gamma$  is in the pleating locus of both  $f_0$ , and  $f_1$ . We can no longer deduce that the four or two components of  $f_j(\zeta)$  in  $T(f_j(\gamma), \varepsilon_0)$  are within a bounded distance of  $\zeta_*$ . But the Short Bridge Arc Lemma 3.4 implies that a point  $x_0$  in  $f_0(\zeta)$  is within a bounded distance of  $f_0(\gamma) = \gamma_*$  precisely when the same is true for a point  $x_1 \in f_1(\zeta)$  for which  $d_3(x_0, x_1)$  is bounded. So in this case, also, we have a bound on  $||f_0(\zeta)| - |f_1(\zeta)||$ , and, hence, upper and lower bounds on  $|f_0(\zeta)|' / |f_1(\zeta)|'$ .

□

## 4.5

We have the following extension of the ideas of 4.4, 4.1. For simplicity, we remove as many badly bent annuli as possible.

**Lemma** *The following holds for a constant  $L_0$ . Let  $\Gamma_0$  be a maximal multicurve. Let  $\zeta \in \Gamma_0$  and let  $\gamma$  be a simple loop intersecting  $\zeta$  finitely many times but no other loop in  $\Gamma_0$ . Let  $f : S \rightarrow N$  have pleating locus including  $\Gamma = (\Gamma \setminus \{\zeta\}) \cup \{\gamma\}$ . Let  $|\gamma_*| < \varepsilon_0$  but  $|\beta_*| \geq \varepsilon_0$  for all  $\beta \in \Gamma_0 \setminus \{\zeta\}$ . Let  $f_n : S \rightarrow N$  be a pleated surface with pleating locus including  $\Gamma_n$  and homotopic to  $f$ . Let  $\zeta_n = \tau_\gamma^n(\zeta)$  for all  $n$ , or  $\tau_\gamma^{-n}(\zeta)$  for all  $n$ . Suppose that both  $f_0(\zeta)$  and  $f_0(\zeta_1)$  are nontrivial in  $N$ . Let  $\Gamma_n = (\Gamma \setminus \{\gamma\}) \cup \{\zeta_n\}$ .*

*Suppose that either  $f$  satisfies the hypothesis of  $f_0$  in 4.3 (and 4.4), or that  $f_j$  does, for all  $0 \leq j \leq n$ . Then so long as  $|(\zeta_j)_*| \geq \varepsilon_0$  for  $j = 0$  and  $j = n$ ,*

$$d([f_0], [f_n]) \leq L_0(\log |n| + 1),$$

*with similar bounds on the length of homotopy tracks of some homotopy between the impressions of  $f_0$  and  $f_n$ , and if  $\alpha = S \setminus \gamma$ , for all  $n$ ,*

$$d_\alpha([f_0], [f_n]) \leq L_0,$$

$$d_\alpha([f_0], [f]) \leq L_0.$$

*Proof.*

From 4.4, since the hypotheses are satisfied, we have, for suitable  $L_0$ ,

$$d_\alpha([f], [f_j]) \leq L_0/2 \text{ whenever } |f_j(\zeta_j)| \geq \varepsilon_0.$$

This then gives  $d_\alpha([f_0], [f_n]) \leq L_0$ .

Let  $\tilde{T}$  be a lift of  $T(\gamma_*, \varepsilon_0)$  to the universal cover  $H^3$  of  $N$ , and let  $\tilde{\zeta}_i$  be a lift of  $(\zeta_i)_*$  such that any intersection  $\tilde{\zeta}_i \cap \tilde{T}$  corresponds to the intersection between  $f_i(\zeta_i)$  and  $f_i(\gamma)$  in  $S(f_i)$ . Then, by the proof of 4.1, we can assume that  $\tilde{\zeta}_i \cap \tilde{T} \neq \emptyset$  and  $|\tilde{\zeta}_i \cap \tilde{T}|$  is increasing in  $i$ . Moreover, for a constant  $L_1$ , for  $i \geq 0$ , for some constant  $s_0 \geq 0$ ,

$$||\tilde{\zeta}_i \cap \tilde{T}| - \log(|i + s_0| + 1)| \leq L_1. \quad (4.5.1)$$

This can be seen by considering  $\tilde{\zeta}_i$ , , translating one component of  $\tilde{\zeta}_i \cap \tilde{T}$  using the covering element of  $\gamma$  and putting in a new connecting segment. Since the Radius of Injectivity Lemma of 3.3 holds for  $f_j$  for all  $j$ , the image under  $f_i$  of the Margulis tube in  $S(f_i)$  round  $f_i(\gamma)$  is, to within bounded distance, the intersection with  $T(\gamma_*, \varepsilon_0)$ . So it follows from (4.5.1) that, for a constant  $L_2$ , for  $i$  between 0 and  $n$ ,

$$|f_i(\gamma)| \leq \frac{L_2}{i+1} \quad (4.5.2)$$

Now suffices to show that for a suitable  $L_3$ , for all  $0 < i \leq n$ , and assuming that the pleating lamination for  $f_{i-1}$  maps under  $\tau_\gamma$  to the pleating locus for  $f_i$  (as it can be chosen to do)

$$d([f_{i-1}], [f_i]) \leq L_3 |f_i(\gamma)|. \quad (4.5.3)$$

For then

$$d([f_0], [f_n]) \leq L_4 \sum_{i=1}^n \frac{1}{|i|} \leq L_4 (\log |n| + 1),$$

as required. So it remains to prove (4.5.3). We do this by constructing  $\chi_i : S(f_{i-1}) \rightarrow S(f_i)$  with  $[\chi_i \circ f_{i-1}] = [f_i]$  and with  $\chi_i$  of distortion  $1 + O(|f_i(\gamma)|)$ . We already have the bound by 4.4, if  $|f_i(\gamma)|$  is bounded from 0. So now we assume that  $|f_i(\gamma)| < \varepsilon_0/2$ , and similarly for  $i-1$  replacing  $i$ . Then we construct  $\chi_i$  to map  $\partial T(f_{i-1}(\gamma), \varepsilon_0)$  to  $\partial T(f_i(\gamma), \varepsilon_0)$ . Outside these sets the pleating loci are homeomorphic and we simply map them across. Inside the Margulis tubes, we choose  $\chi_i$  to map the geodesic segments in  $\partial T(f_{i-1}(\gamma), \varepsilon_0)$ , geodesic with respect to the hyperbolic metric on  $S(f_{i-1})$ , whose images under  $f_{i-1}$  are homotopic to the segments of  $f_i(\zeta_i)$  in  $\partial T(f_i(\gamma), \varepsilon_0)$ , to those segments. In all cases, geodesic side lengths differ by  $O(|f_i(\gamma)|)$ . So we can take constant derivative with respect to length on each geodesic segment, with derivative  $1 + O(|f_i(\gamma)|)$ . The union of the polygons has full measure. Then we extend across polygons and we get the required bound on distortion.

□

## 4.6 Pleated Surfaces near the domain of discontinuity.

We now return to a topic left open at the end of Section 3. 3.11 is concerned with geometrically infinite ends. We shall also need a corresponding result for geometrically finite ends. So now, let  $e$  be a geometrically finite end of  $(N_d, \partial N_d)$ . Write  $S(e) = S$  and  $S_d(e) = S_d$ . Then there is a component  $S_1$  of the boundary of the convex hull bounding a neighbourhood of  $e$ , whose intersection with  $N_d$  is homeomorphic to  $S_d$ , under a homeomorphism of  $N$  which is isotopic to the identity. The neighbourhood of  $e$  can also be compactified by adding a component  $S_2$  of  $\Omega/\pi_1(N)$ , where  $\Omega$  is the domain of discontinuity of  $\pi_1(N)$ , where we are identifying  $\pi_1(N)$  with the covering group of hyperbolic isometries of  $N$ . These two surfaces  $S_1$  and  $S_2$  both have hyperbolic structures, or equivalently complex structures, and thus give points of  $\mathcal{T}(\mathcal{S})$ , which for the moment we call respectively  $[f_1]$  and  $[f_2]$ . It is proved in [18] that there is a natural bounded distortion map between the two surfaces, and that the two points in Teichmüller space are a bounded distance apart. Actually, there is a fairly direct proof of the following. But note the assumption that  $\Gamma$  is noncollapsing.

**Lemma** *Fix a 2- and 3-dimensional Margulis constant  $\varepsilon_0$ . Given  $L_1$  there is  $L_2$  such that the following hold. Let  $\Gamma \subset S$  be a noncollapsing maximal multicurve with  $|f_2(\gamma)| \leq L_1$  for all  $\gamma \in \Gamma$ .*

*Then  $|\gamma_*| \leq L_2$  for all  $\gamma \in \Gamma$ , where  $\gamma_*$  denotes the geodesic in  $N$  homotopic to  $f_2(\gamma)$ .*

*There is an embedded surface  $S_3$  in  $N$  whose preimage in  $H^3$  bounds a convex subset of  $H^3$ , such that inclusion  $f_3 : S \rightarrow N$  is homotopic to inclusion  $f_2 : S \rightarrow \bar{N}$ , such that, using the metric on  $S_3$  induced by the hyperbolic metric on  $H^3$ ,*

$$L_2^{-1} \leq \frac{|f_3(\gamma)|}{|f_2(\gamma)|} \leq L_2$$

*for all  $\gamma \in \Gamma$ .*

*There is a pleated surface  $f_4$  with pleating locus a maximal multicurve  $\Gamma'$ , such that*

- $\#(\Gamma \cap \Gamma') \leq L_2$ ,
- $n_{\gamma, \beta}([f_2]) \leq L_2 |f_2(\gamma)|^{-1}$  for all  $\beta \in \Gamma'$  transverse to  $\gamma \in \Gamma$  (where  $n_{\gamma, \beta}(\cdot)$  is as in 2.6),
- $f_4$  has no badly bent annuli,
- there is a homotopy between  $f_3$  and  $f_4$  with homotopy tracks of hyperbolic length  $\leq L_2$ ,
- 

$$d([f_2], [f_4]) \leq L_2.$$

#### 4.7 Proof of 4.6 in the case of bounded geometry on $S_2$ .

To start with, we assume that  $[f_2] \in (\mathcal{T}(\mathcal{S}))_{\geq \nu}$  for some  $\nu > 0$ , and obtain estimates in terms of  $\nu$ . We can cover  $S_2$  by  $\leq n(\nu)$  topological discs, which are round discs up to bounded distortion in the hyperbolic metric on  $S_2$  (independent of  $\nu$ ), all with boundedly proportionate radii in terms of  $\nu$ , and lift to Euclidean discs in  $\Omega$ .

We can also assume that Euclidean discs of half the Euclidean radii still cover  $\Omega$ . The discs cover  $\Omega$  with index  $\leq n(\nu)$ , assuming the discs have radius less than half the radius of injectivity. Each Euclidean disc is the base of a Euclidean half-ball in  $H^3$ , using the half-space model for  $H^3$ . The intersection of all the complementary half-balls is a convex set. Fix one boundary component,  $U$ . Then the stabiliser of  $U$  in  $\pi_1(N)$  is, up to conjugacy in  $\pi_1(N)$ ,  $i_*(\pi_1(S))$ , where  $i : S \rightarrow N$  denotes inclusion. One component  $\Omega_2$  of  $\Omega$  is separated by  $U$  from the convex hull of  $\pi_1(N)$ , is stabilised by  $i_*(\pi_1(S))$ , and covers  $S_2$ . Because Euclidean discs of half the radius still cover, each hemisphere boundary of a half-ball which intersects  $U$  does so in a set of bounded hyperbolic diameter, and each bounded loop on  $S_2$  has a lift to the upper boundary of the hemispheres such that the projection to  $U/i_*(\pi_1(S))$  is of bounded hyperbolic length. This is our surface  $S_3$ . We write  $f_3 : S \rightarrow S_3$  for the inclusion map.

Now let  $\Gamma$  be given. For each  $\gamma \in \Gamma$ , if we write  $\gamma$  also for the corresponding element of  $i_*(\pi_1(S))$ , there is  $x = x(\gamma) \in U$  such that the hyperbolic distance between  $x$  and  $\gamma.x$  in  $U$  is  $\leq L_2 = L_2(L_1, \nu)$ . Hence,  $|\gamma_*| \leq L_2$ . It could be that  $|\gamma_*| < \varepsilon_0$ , where  $\varepsilon_0$  is a fixed Margulis constant. But if so, we can find a loop  $\beta(\gamma)$  such that  $\beta(\gamma)$  intersects  $\gamma$  at most twice, but no other loop in  $\Gamma$ , and  $|f_2(\beta(\gamma))| \leq L'_1$  where  $L'_1$  depends only on  $L_1$ . If our first choice of  $\beta(\gamma)$  is trivial in  $N$ , then we can make it nontrivial, just by composing with a single twist round  $\gamma$ . This must be nontrivial, because otherwise  $\gamma$  is trivial. Then  $\beta(\gamma)$  also has a lift to  $U$  of length  $\leq L_2$ , assuming that  $L_2$  large enough given  $L_1$ . Also, the points  $x(\gamma)$  and  $x(\beta(\gamma))$  can be chosen a bounded hyperbolic distance apart. Then  $|(\beta(\gamma))_*| \leq L_2$ . If both  $|\gamma_*|$  and  $|(\beta(\gamma))_*|$  are small, then their  $\varepsilon_0$ -Margulis tubes are disjoint, and both come within a bounded distance of the lifts of  $\gamma, \beta(\gamma)$  respectively on  $U$ . So they cannot both be above  $U$ . But they do not intersect  $U$ . But the region above  $U$  is convex, and its closure contains the limit set. So both  $\gamma_*$  and  $(\beta(\gamma))_*$  must be above  $U$ . The loop  $(\beta(\gamma))_*$  can also not be separated from  $U$  by the Margulis tube of  $\gamma_*$ , if there is one, because it is a bounded distance from  $U$ . So if  $\gamma$  is short,  $\beta(\gamma)$  is not, and is not separated from  $U$  by the orbit of  $\gamma$ . Now we change  $\Gamma$  to  $\Gamma'$  through a sequence of maximal multicurves  $\Gamma_i$ ,  $0 \leq i \leq r$ , with  $\Gamma_0 = \Gamma$  and  $\Gamma_r = \Gamma'$ . At each stage, we replace some short loop  $\gamma \in \Gamma_i \cap \Gamma$  by a loop  $\beta(\gamma)$  disjoint from all loops of  $\Gamma_i \setminus \{\gamma\}$ , such that  $|f_2(\beta(\gamma))|$  is bounded and bounded from 0. We can choose the  $\beta(\gamma)$  at each stage so that  $\#(\Gamma_i \cap \Gamma_j) \leq L_2$  for all  $i, j$ . In particular,  $\#(\Gamma \cap \Gamma') \leq L_2$ . We then take  $f_4$  to be a pleated surface with pleating locus containing  $\Gamma'$ . We have bounds on  $|f_4(\gamma)|$  for all  $\gamma \in \cup_i \Gamma_i$  because either  $\gamma \in \Gamma'$  or  $\#(\gamma \cap \Gamma')$  is bounded, in which case we can apply 3.6. We also have bounds on  $|f_4(\beta(\gamma))|$

for all  $\gamma \in \Gamma \cap \Gamma'$ , again by applying 3.6. So we have

$$d([f_2], [f_4]) \leq L_2.$$

In order to construct a homotopy with bounded tracks between  $f_3$  and  $f_4$ , use the same procedure as in 4.3, which is itself derived from [37]. The loop set  $\Gamma'$  has the property that  $f_3(\gamma)$  and  $f_4(\gamma)$  are a bounded hyperbolic distance apart, and  $\Gamma'$  is a maximal multicurve. We are assuming, here, that  $f_3(\gamma)$  and  $f_4(\gamma)$  are geodesics in the respective Poincaré metrics. So we take the homotopy to homotope  $f_3(\gamma)$  to  $f_4(\gamma)$ , for all  $\gamma \in \Gamma'$ . Then, as in 4.3, we foliate each complementary pairs of pants by arcs, except for two tripods. Because of bound on  $d([f_3], [f_4])$ , bounded arcs on  $f_3(S) = S_3$  are homotopic to bounded arcs on  $f_4(S)$ . Then we homotope tripods to tripods, and arcs in between as dictated by the endpoints of arcs.

#### 4.8 Proof of 4.6 in general.

Now suppose that  $|f_2(\gamma)|$  is small for some loop  $\gamma \in \Gamma$ . Then we construct  $U$  over the preimage in  $\Omega_2$  of  $(S_2)_{\geq \varepsilon_1}$  in the same way as before. But over  $(S_2)_{< \varepsilon_1}$ , for a sufficiently small  $\varepsilon_1$ , we change the construction. As before, let  $\gamma$  denote both a loop with  $|f_2(\gamma)| < \varepsilon_1$ , and a corresponding element of the covering group, which leaves invariant a component of the preimage of  $T(f_2(\gamma), \varepsilon_0) \subset S_2$  in  $\partial H^3$ . We identify  $\partial H^3$  with  $\mathbb{C}$  in the usual way. Let  $\lambda$  be the complex length of  $\gamma$ . Now  $\gamma$  embeds in a 1-parameter subgroup  $\gamma_t$  of  $PSL(2, \mathbb{C})$  which acts on  $\mathbb{C}$ . The eigenvalues of  $\gamma_t$ , considered as an element of  $PSL(2, \mathbb{C})$ , are  $e^{\pm t\lambda/2}$ . The action of the group  $\gamma_t$  has two fixed points  $x_1, x_2$ , the endpoints of the geodesic  $\gamma_*$  in  $H^3$  lifting the loop  $\gamma$ . Let  $V$  be the connected component of the preimage of  $T(f_2(\gamma), \varepsilon_0) \subset S_2$  in  $\mathbb{C}$  whose closure contains the points  $x_1$  and  $x_2$ . The closure of  $V$  in  $\mathbb{C}$  is then the union of the lift of  $T(f_2(\gamma), \varepsilon_0) \subset S_2$  and  $x_1, x_2$ . Normalise  $\mathbb{C}$  so that  $\overline{V} \subset \mathbb{C}$ . In fact, we can, and shall, normalise so that the diameters are both boundedly proportional to 1. The orbits of the  $\gamma_t$  action are spirals connecting  $x_1$  and  $x_2$ .

If  $C_1$  is sufficiently large, a ball of Poincaré radius  $\varepsilon_0$  centred on a point  $x$  of  $(S_2)_{\leq \varepsilon_0/C_1}$  lifts to  $V$  in such a way that the disc  $D(\tilde{x})$  of Euclidean radius  $2|\gamma.\tilde{x} - \tilde{x}|$  centred on any lift  $\tilde{x} \in V$  of  $x$  is contained in the lift. Assuming  $C_1$  is sufficiently large, the section of spiral between  $\tilde{x}$  and  $\gamma.\tilde{x}$  is contained in this disc. By invariance, the whole spiral must be contained in  $V$ , as claimed. One interesting feature of this is that the spiral only intersects  $D(\tilde{x})$  in one arc. If it intersected in more than one arc, then one endpoint of the geodesic  $\gamma$  would be an isolated point in the closure of  $V$ , which is impossible, because the limit set is a perfect set. So now let  $V_1$  be the connected union of spirals. Then the closure  $\overline{V}_1$ , apart from the endpoints  $x_1, x_2$  is contained in  $V$ . We can further normalise  $\mathbb{C}$  so that the two components of  $\partial V_1 \setminus \{x_1, x_2\}$ , as for  $\partial V$ , have proportional Euclidean diameters, and so that the Euclidean diameter of  $V_1$  is 1. The modulus of the annulus  $V_1 / < \gamma >$  is boundedly proportional to the modulus of  $T(f_2(\gamma), \varepsilon_0)$ , and hence inversely proportional, for a multiplicative constant depending on  $\varepsilon_0$ , to  $|f_2(\gamma)|$ . Now let  $\tilde{\gamma}$  be the spiral  $\{\gamma_t.\tilde{x} : t \in \mathbb{R}\}$  for some  $\tilde{x} \in V_1$

which is in the lift of  $f_2(\gamma)$  in  $V$ . The Euclidean distance between spirals for different choices of  $\tilde{x}$  is bounded by  $C_0|x_1 - x_2|$ , for a universal constant  $C_0$ . The modulus of  $V_1 / \langle \gamma \rangle$  is also boundedly proportional to  $\text{Max}(1/d_1, 1/d_2)$ , where  $d_1$  is the maximum Euclidean distance between  $\tilde{x}$  and  $\gamma.\tilde{x}$ , for  $\tilde{x} \in \partial V_1$ , and  $d_2$  is similarly defined for  $\tilde{x} \in \tilde{\gamma}$ . In fact,  $\text{Max}(1/d_1, 1/d_2)$  is always boundedly proportional to  $1/d_2$ , and  $d_1$  and  $d_2$  are boundedly proportional if and only if  $|x_1 - x_2|$  is bounded from 0.

Now we claim that there is a  $\gamma$ -invariant set  $V'_1$  with  $V'_1 \subset V_1$ , such that the Poincaré distance between  $V'_1$  and  $V_1$  is bounded, and  $V'_1$  is the  $\gamma$ -orbit of between one and five Euclidean discs, where two of the discs have Euclidean radius which are bounded and bounded to 0, and the smallest has Euclidean radius boundedly proportional to  $d_2$ . We see this as follows. We can transfer back under a Möbius transformation  $\sigma$  for the moment, to the situation when  $x_1 = 0$  and  $x_2 = \infty$ . So under this transformation,  $V_1$  transforms to

$$V_0 = \sigma(V_1) = \{e^{\lambda t + i\theta} : \theta \in [0, \theta_1]\}$$

for some  $0 < \theta_1 < 2\pi$ . We can assume without loss of generality that  $\text{Re}(\lambda) < 0$  (interchanging 0 and  $\infty$  if necessary). The element  $\gamma$  transfers to multiplication by  $e^\lambda$ , and  $\lambda$  is small. We only need to produce an orbit of up to five for this set, because our normalisation of  $V_1$  has  $V_1$  bounded, and so discs will transfer to discs under  $\sigma$ . If we can do so, we choose just one disc in  $\sigma(V_1)$  which is of Euclidean diameter which is bounded and bounded from 0, and tangent to both components of  $\sigma(\partial V_1) \setminus \{0, \infty\}$ . This is possible if  $\theta_1 < \pi/3$ , and also for any  $\theta_1 < 2\pi$ , if  $\text{Im}(\lambda)/\text{Re}(\lambda)$  is bounded from 0. In general, we can always choose a connected union  $V_0'' \leq 5$  discs of Euclidean diameters which are bounded, and the outer two bounded from 0, and each tangent to one of the components of  $\sigma(\partial V_1) \setminus \{0, \infty\}$ , at points which are bounded from 0. We can choose  $V_0''$  so that  $V_0 \setminus (V_0'' \cup e^\lambda V_0'')$  has at most two components whose closure does not include 0,  $\infty$ , at most one intersecting each component of  $\partial V_0 \setminus \{0, \infty\}$ . If more than two discs are needed, we can also choose all but the outer two so that each one intersects the two nearest discs in two sets, each of diameter proportional to its own diameter. Let  $V'_0 = \sigma(V'_1)$  be the orbit of the union of these discs under multiplication by  $e^\lambda$ . Then  $V_1 \setminus V'_1$  is contained in a neighbourhood of  $\partial V_1$  of bounded Poincaré diameter, with respect to the Poincaré metric on  $\Omega_2$ .

Then we take  $U$  to be the surface bounded by the union of the hemispheres over the lift of  $(S_2)_{\geq \varepsilon_1}$  and over  $V'_1$ , for a suitable  $\varepsilon_1$  with  $\varepsilon_0/\varepsilon_1$  bounded, so that the sets  $V'_1(\tilde{\gamma})$ , for varying  $\tilde{\gamma}$ , and the lifts of  $(S_2)_{\geq \varepsilon_1}$ , cover  $\Omega_2$ . Then  $S_3 = U/i_*(\pi_1(S))$  is an embedded surface, with full preimage in  $H^3$  bounding a convex subset of  $H^3$ , with corresponding map  $f_3 : S \rightarrow S_3$  homotopic, as a map from  $S$  to  $\overline{N}$ , to  $f_2 : S \rightarrow S_2$ . To find the shortest loop in  $S_3$  homotopic to  $f_3(\gamma)$ , we can draw paths on the tops of the hemispheres. As before, we can estimate hyperbolic length by the ratio of the Euclidean length on the top of a hemisphere to the Euclidean radius. The paths on tops of all but the outer hemispheres over  $V'_1$  are bounded, because of the conditions we imposed on the intersections of the base hemispheres. As for the outer hemispheres (if  $V'_1$  is the orbit of more

than one hemisphere) a path across the top of an outer hemisphere  $H_1$  also has bounded hyperbolic length. But the distance to the nearest hemisphere over  $(S_2)_{\geq \varepsilon_1}$  might be much greater, if  $H_1$  has a much larger Euclidean diameter than the nearby hemispheres over  $(S_2)_{\geq \varepsilon_1}$ . In any case, the hyperbolic length of  $f_3(\gamma)$  is boundedly proportional to  $d_2$ , and to  $|f_2(\gamma)|$ . Also by taking paths on the tops of hemispheres, we can choose  $\beta(\gamma)$  transverse to  $\gamma$  such that  $f_3(\beta(\gamma))$  is, up to homotopy in  $S_3$ , a union of boundedly finitely many paths of bounded length and, at most two long geodesic segments in the hemispheres over  $V'_1$  for each crossing of  $\gamma$ . We can ensure that the long segments do not cancel by adjusting by a Dehn twist round  $\gamma$ , if necessary, but so that  $n_{\gamma, \beta(\gamma)}([f_2])$  is bounded by  $O(|f_2(\gamma)|^{-1})$ . So then  $f_3(\gamma)$  is, up to homotopy, a bounded distance from  $(\beta(\gamma))_*$ . So then, as before, we can form  $\Gamma'$  with  $\#(\Gamma \cap \Gamma') \leq L_2$ , from a sequence  $\Gamma_i$  with  $\Gamma_0 = \Gamma$ ,  $\Gamma_r = \Gamma'$ , and such that  $\Gamma_{i+1}$  is obtained from  $\Gamma_i$  by replacing some loop  $\gamma$  with  $|f_2(\gamma)|$  small by a loop  $\beta(\gamma)$ . Then there is a bounded track homotopy between  $f_3$  and  $f_4$ . The homotopy is defined restricted to the set with image in  $N_{\geq \varepsilon_0}$  exactly as before, and the sets  $V_2$  have been constructed so as to ensure that there is a bounded track homotopy on the set  $S_3 \cap N_{< \varepsilon_0}$ . It follows that the moduli of  $T(f_4(\gamma), \varepsilon_0) \subset S(f_4)$  and  $T(f_2(\gamma), \varepsilon_0) \subset S_2$  are boundedly proportional, and that  $|f_2(\beta(\gamma))|'$  and  $|f_4(\beta(\gamma))|'$  are boundedly proportional. The bound on  $d([f_2], [f_4])$  follows.  $\square$

#### 4.9 Generalised pleated surfaces.

A pleated surface  $f : S \rightarrow N$  derives its metric from the hyperbolic metric on  $N$ . It will sometimes be useful to use the metric on  $\partial N$ , where  $\partial N$  is the boundary if  $N$  obtained by projecting the Poincaré metric from the domain of discontinuity  $\Omega \subset \partial H^3$ . This is, in fact, the case for  $f_2 : S \rightarrow S_2$  in 4.6. Write  $\overline{N} = N \cup \partial N$ . We consider maps  $f : S \rightarrow N$ , where  $f$  maps the loops of a multicurve  $\Gamma$  to cusps in  $N$ , and, for each component  $\alpha$  of  $S \setminus \cup \Gamma$ , either  $f|_\alpha$  is a pleated surface, or  $f|_\alpha$  is a homeomorphism onto a component of  $\partial N$ . Then  $[f|_\alpha]$  is an element of  $\mathcal{T}(S(\alpha))$  for each  $\alpha$ . If  $f(\alpha) \subset \partial N$ , we use the Poincaré metric on  $S(\alpha)$  to define the element of  $\mathcal{T}(S(\alpha))$ . We shall refer to such a map  $f$  as a *generalised pleated surface*. It defines an element  $[f] \in \mathcal{T}(S(\omega))$ , where  $\omega = S \setminus (\cup \Gamma)$ . Applying 4.6 to each  $S(\alpha)$  with  $f(\alpha) \subset \partial N$ , there is a genuine pleated surface a bounded  $d_\omega$  distance away.

### 5 Teichmüller geodesics: long thick and dominant definitions.

In this section we explain and expand some of the ideas of long thick and dominant (ltd) segments of geodesics in Teichmüller space  $\mathcal{T}(\mathcal{S})$  which were used in [50]. The theory of [50] was explicitly for marked spheres only, because of the application in mind, but in fact the theory works without adjustment for any finite type surface, given that projections  $\pi_\alpha$  to smaller Teichmüller

spaces  $\mathcal{T}(\mathcal{S}(\alpha))$  for subsurfaces  $\alpha$  of  $S$  have been defined in 2.6. For proofs, for the most part, we refer to [50]. The basic idea is to get into a position to apply arguments which work along geodesics which never enter the thin part of Teichmüller space, by projecting to suitable subsurfaces  $\alpha$  using the projections  $\pi_\alpha$  of 2.6. A reader who wishes to get to the proof of the Ending Laminations Theorem in the case of combinatorial bounded geometry is advised to read to the end of the basic definition 5.3, and then proceed to a recommended menu from Section 6. In the theory of Teichmüller geodesics which is developed here (and earlier, in [50]) it does not seem to make sense to consider geodesics in the thick part of Teichmüller space — which is what combinatorially bounded geometry means — in strict isolation. We use the basic notation and theory of Teichmüller space  $\mathcal{T}(\mathcal{S})$  from Section 2.

### 5.1 Good position.

Let  $[\varphi] \in \mathcal{T}(\mathcal{S})$ . Let  $q(z)dz^2$  be a quadratic differential on  $\varphi(S)$ . All quadratic differentials, as in 2.3, will be of total mass 1. Let  $\gamma$  be a nontrivial nonperipheral simple closed loop on  $S$ . Then there is a limit of isotopies of  $\varphi(\gamma)$  to *good position* with respect to  $q(z)dz^2$ , with the limit possibly passing through some punctures. If  $\gamma$  is the isotopy limit, then either  $\gamma$  is at constant angle to the stable and unstable foliations of  $q(z)dz^2$ , or is a union of segments between singularities of  $q(z)dz^2$  which are at constant angle to the stable and unstable foliations, with angle  $\geq \pi$  between any two consecutive segments at a singularity, unless it is a puncture. An equivalent statement is that  $\gamma$  is a geodesic with respect to the singular Euclidean metric  $|q(z)|d|z|^2$ . If two good positions do not coincide, then they bound an open annulus in  $\varphi(S)$  which contains no singularities of  $q(z)dz^2$ . See also 14.5 of [50].

The *q-d length*  $|\varphi(\gamma)|_q$  is length with respect to the quadratic differential metric for any homotopy representative in good position. (See 14.5 of [50].) We continue, as in Section 2, to use  $|\varphi(\gamma)|$  to denote the hyperbolic, or Poincaré, length on  $\varphi(S)$  of the geodesic on  $\varphi(S)$  homotopic to  $\varphi(\gamma)$ . If  $[\varphi] \in \mathcal{T}_{\geq \varepsilon}$  then there is a constant  $C(\varepsilon) > 0$  such that for all nontrivial nonperipheral closed loops  $\gamma$ ,

$$\frac{1}{C(\varepsilon)} \leq \frac{|\varphi(\gamma)|_q}{|\varphi(\gamma)|} \leq C(\varepsilon).$$

We also define  $|\varphi(\gamma)|_{q,+}$  to be the integral of the norm of the projection of the derivative of  $\varphi(\gamma)$  to the tangent space of the unstable foliation of  $q(z)dz^2$ , and similarly for  $|\varphi(\gamma)|_{q,-}$ . So these are both majorised by  $|\varphi(\gamma)|_q$ , which is, in turn, majorised by their sum.

### 5.2 Area.

The following definitions come from 9.4 of [50]. For any essential nonannulus subsurface  $\alpha \subset S$ ,  $a(\alpha, q)$  is the area with respect to  $q(z)dz^2$  of  $\varphi(\alpha)$  where  $\varphi(\partial\alpha)$  is in good position and bounds the smallest area possible subject to this restriction. If  $\alpha$  is a loop at  $x$  then  $a(\alpha, q)$  is the smallest possible area of an



annulus of modulus 1 and homotopic to  $\varphi(\alpha)$ . We are only interested in this quantity up to a bounded multiplicative constant and it is also boundedly proportional to  $|\varphi(\alpha)|_q^2$  whenever  $\varphi(\alpha)$  is in good position, and  $|\varphi(\alpha)|$  is bounded. We sometimes write  $a(\alpha, x)$  or even  $a(\alpha)$  for  $a(\alpha, q)$ , if it is clear from the context what is meant.

Generalising from 5.1, there is a constant  $C(\varepsilon)$  such that, if  $\varphi(\alpha)$  is homotopic to a component of  $(\varphi(S))_{\geq \varepsilon}$ , then for all nontrivial nonperipheral non-boundary-homotopic closed loops  $\gamma \in \alpha$ ,

$$\frac{1}{C(\varepsilon)}|\varphi(\gamma)| \leq \frac{|\varphi(\gamma)|_q}{\sqrt{a(\alpha, q) + a(\partial\alpha, q)}} \leq C(\varepsilon)|\varphi(\gamma)|.$$

Now suppose that  $\ell$  is a directed geodesic segment in  $\mathcal{T}(\mathcal{S})$  containing  $[\varphi]$ , and that  $q(z)dz^2$  is the quadratic differential at  $[\varphi]$  for  $d([\varphi], [\psi])$  for any  $[\psi]$  in the positive direction along  $\ell$  from  $[\varphi]$  (see 2.2.) Let  $p(z)dz^2$  be the stretch of  $q(z)dz^2$  at  $[\psi]$ , and let  $\chi$  be the minimum distortion map with  $[\chi \circ \varphi] = [\psi]$ . Then  $\chi$  maps the  $q$ -area element to the  $p$ -area element. Then  $a(\alpha, q) = a(\alpha, p)$  if  $\alpha$  is a gap, but if  $\alpha$  is a loop,  $a(\alpha, y)$  varies for  $y \in \ell$ .

If  $\alpha$  is a loop we also make an extra definition. We define  $a'(\alpha, [\varphi], q)$  (or simply  $a'(\alpha)$  if the context is clear) to be the  $q$ -area of the largest modulus annulus (possibly degenerate) homotopic to  $\varphi(\alpha)$  and with boundary components in good position for  $q(z)dz^2$ . Then  $a'(\alpha)$  is constant along the geodesic determined by  $q(z)dz^2$ .

### 5.3 The long thick and dominant definition

Now we fix parameter functions  $\Delta, r, s : (0, \infty) \rightarrow (0, \infty)$  and a constant  $K_0$ .

Let  $\alpha$  be a gap. Let  $\ell$  be a geodesic segment. We say that  $\alpha$  is *long,  $\nu$ -thick and dominant at  $x$*  (for  $\ell$ , and with respect to  $(\Delta, r, s)$ ) if  $x$  is the centre of a segment  $\ell_1$  in the geodesic extending  $\ell$  of length  $2\Delta(\nu)$  such that  $|\psi(\gamma)| \geq \nu$  for all  $[\psi] \in \ell_1$  and nontrivial nonperipheral  $\gamma \subset \alpha$  not homotopic to boundary components, but  $\ell_1 \subset \mathcal{T}(\partial\alpha, r(\nu))$  and  $a(\partial\alpha, y) \leq s(\nu)a(\alpha, y)$  for all  $y \in \ell_1$ . We shall then also say that  $\alpha$  is long  $\nu$ -thick and dominant along  $\ell_1$ . See 15.3 of [50].

A loop  $\alpha$  at  $x$  is  *$K_0$ -flat at  $x = [\varphi]$*  (for  $\ell$ ) if  $a'(\alpha) \geq K_0 a(\alpha)$ . This was not quite the definition made in [50] where the context was restricted to  $S$  being a punctured sphere, but the results actually worked for any finite type surface. The term arises because if  $\alpha$  is  $K_0$ -flat then the metric  $|q(z)|dz^2$  is equivalent to a Euclidean (flat) metric on an annulus homotopic to  $\varphi(\alpha)$  of modulus  $K_0 - O(1)$ . For fixed  $K_0$  we may simply say *flat* rather than  $K_0$ -flat. In future, we shall often refer to parameter functions as quadruples of the form  $(\Delta, r, s, K_0)$ .

If  $\alpha$  is long  $\nu$ -thick and dominant along a segment  $\ell$ , that is long thick and at all points of  $\ell$ , then  $d_\alpha(x, y)$  is very close to  $d(x, y)$  at all points of  $\ell$ . This is a consequence of the results of Section 11 of [50]. All we care about for the moment is that they differ by some additive constant. It is also probably worth noting (again by the results of Chapter 11 of [50]) that if  $[\varphi] \in \ell$  and

$\pi_\alpha([\varphi]) = [\varphi_\alpha]$ , then  $\varphi_\alpha(S(\alpha))$  and the component  $S(\alpha, r(\nu)[\varphi])$  of  $(\varphi(S))_{\geq r(\nu)}$  homotopic to  $\varphi(\alpha)$  are isometrically very close, except in small neighbourhoods of some punctures, and the quadratic differentials  $q(z)dz^2$  at  $[\varphi]$  for  $d([\varphi], [\psi])$  ( $[\psi] \in \ell$ ) and the quadratic differential  $q_\alpha(z)dz^2$  at  $[\varphi_\alpha]$  for  $d_\alpha([\varphi], [\psi])$ , are very close.

## 5.4

Before starting to describe the usefulness of long thick and dominants, we need to show they exist, in some abundance. This was the content of the first basic result about long thick and dominants in 15.4 of [50], which was stated only for  $S$  being a punctured sphere, but the proof worked for a general finite type surface.

**Lemma** *For some  $\nu_0$  and  $\Delta_0$  depending only on  $(\Delta, r, s, K_0)$  (and the topological type of  $S$ ), the following holds. Any geodesic segment  $\ell$  of length  $\geq \Delta_0$  contains a segment  $\ell'$  for which there is  $\alpha$  such that:*

- *either  $\alpha$  is a gap which is long  $\nu$ -thick and dominant along  $\ell'$  for some  $\nu \geq \nu_0$  and  $a(\alpha) \geq 1/(-2\chi(S) + 1) = c(S)$  (where  $\chi$  denotes Euler characteristic,*
- *or  $\alpha$  is a  $K_0$ -flat loop along  $\ell'$ .*

*More generally there is  $s_0$  depending only on  $(\Delta, r, s, K_0)$  (and the topological type of  $S$ ) such that, whenever  $\omega \subset S$  is such that  $a(\partial\omega) \leq s_0 a(\omega)$ , then we can find  $\alpha$  as above with  $\alpha \subset \omega$  and  $a(\alpha) \geq 1/(-2\chi(\omega) + 1)a(\omega)$  if  $\alpha$  is a gap.*

*Proof.* (See also 15.4 of [50].) We consider the case  $\omega = S$ . Write  $r_1(\nu) = e^{-\Delta(\nu)}r(\nu)$ . Let  $g = -2\chi(S)$  and let  $r_1^g$  denote the  $g$ -fold iterate. We then take

$$\nu_0 = r_1^g(\varepsilon_0)$$

for a fixed Margulis constant  $\varepsilon_0$  and we define

$$\Delta_0 = 2 \sum_{j=1}^g \Delta(r_1^j(\varepsilon_0)).$$

Then for some  $j \leq g$ , we can find  $\nu = r_1^j(\varepsilon_0)$  and  $[\varphi] \in \ell$  such that the segment  $\ell'$  of length  $2\Delta(\nu)$  centred on  $y = [\varphi]$  is contained in  $\ell$  and such that for any nontrivial nonperipheral loop  $\gamma$ , either  $|\varphi'(\gamma)| \geq \nu$  for all  $[\varphi'] \in \ell'$ , or  $|\varphi(\gamma)| \leq r_1(\nu)$  — in which case  $|\varphi'(\gamma)| \leq r(\nu)$  for all  $[\varphi'] \in \ell'$ . Suppose there are no  $K_0$ -flat loops at  $[\varphi]$ , otherwise we are done. For any loop  $\gamma$  with  $|\varphi(\gamma)| < \varepsilon_0$ , if  $\beta$  is a gap such that  $\gamma \subset \partial\beta$  and there is a component of  $(\varphi(S))_{\geq \varepsilon_0}$  homotopic to  $\varphi(\beta)$  and separated from the flat annulus homotopic to  $\varphi(\gamma)$  by an annulus of modulus  $\Delta_1$ , we have, since every zero of  $q(z)dz^2$  has order at most  $2g$ , for a constant  $C_1$  depending only on the topological type of  $S$ ,

$$C_1^{-1}a(\gamma, [\varphi])e^{\Delta_1} \leq a(\beta) \leq C_1 a(\gamma, \varphi)e^{(2g+2)\Delta_1}. \quad (5.4.1)$$

Now let  $\alpha$  be a subsurface such that  $\varphi(\alpha)$  is homotopic to a component  $S(\alpha, \nu)$  of  $(\varphi(S))_{\geq \nu}$  of maximal area. Then by (5.4.1), we have a bound of

$O(e^{(2g^2+2g)/\nu})$  on the ratio of areas of any two components of  $(\varphi(S))_{\geq \varepsilon_0}$  in  $S(\alpha, \nu)$  and assuming  $r(\nu)$  is sufficiently small given  $\nu$ ,

$$a(\partial\alpha, [\varphi]) \leq e^{-1/(3gr(\nu))} a(\alpha),$$

and for all  $y' \in \ell'$ ,

$$a(\partial\alpha, y') \leq e^{\Delta(\nu)} e^{-1/(3gr(\nu))} a(\alpha).$$

Assuming  $r(\nu)$  is sufficiently small given  $s(\nu)$  and  $\Delta(\nu)$ ,  $\alpha$  is long  $\nu$ -thick and dominant along  $\ell'$  for  $(\Delta, r, s)$ , and  $a(\alpha) \geq 1/(g+1)$

The case  $\omega = S$  is similar. We only need  $s_0$  small enough for  $a(\partial\omega)/a(\omega)$  to remain small along a sufficiently long segment of  $\ell$ .  $\square$

Because of this result, we can simplify our notation. So let  $\nu_0$  be as above, given  $(\Delta, r, s, K_0)$ . We shall simply say  $\alpha$  is *ltd* (at  $x$ , or along  $\ell_1$ , for  $\ell$ ) if either  $\alpha$  is a gap and long  $\nu$ -thick and dominant for some  $\nu \geq \nu_0$ , or  $\alpha$  is a loop and  $K_0$ -flat. We shall also say that  $(\alpha, \ell_1)$  is *ltd*.

## 5.5

We refer to Chapters 14 and 15 of [50] for a summary of all the results concerning *ltd*'s, where, as already stated, the context is restricted to  $S$  being a punctured sphere, but the results work for any finite type surface. The main points about *ltd*'s are, firstly, that they are good coordinates, in which arguments which work in the thick part of Teichmüller space can be applied, and secondly that there is only bounded movement in the complement of *ltd*'s. This second fact, together with 4.4, is worth scrutiny. It is, at first sight, surprising. It is proved in 15.14 of [50], which we now state, actually slightly corrected since short interior loops in  $\alpha$  were forgotten in the statement there (although the proof given there does consider short interior loops) and slightly expanded in the case of  $\alpha$  being a loop.

**Lemma** *Fix long thick and dominant parameter functions  $\Delta, r, s, K_0) > 0$ , and let  $\nu_0 > 0$  also be given and sufficiently small. Then there exists  $L = L(\Delta, r, s, K_0, \nu_0)$  such that the following holds. Let  $\ell$  be a geodesic segment and let  $\ell_1 \subset \ell$  and, given  $\ell_1$ , let  $\alpha \subset S$  be a maximal subsurface up to homotopy with the property that  $\alpha \times \ell_1$  is disjoint from all *ltd*'s  $\beta \times \ell'$  such that  $\beta$  is either  $K_0$ -flat along  $\ell'$  or  $\nu$ -thick long and dominant for some  $\nu \geq \nu_0$ , for  $[\varphi] \in \ell_1$ . Suppose also that all components of  $\partial\alpha$  are nontrivial nonperipheral. Then  $\alpha$  is a disjoint union of gaps and loops  $\beta$  such that the following hold.*

$$|\varphi(\partial\beta)| \leq L \text{ for all } [\varphi] \in \ell. \quad (5.5.1)$$

*If  $\beta$  is a gap, then for all  $[\varphi], [\psi] \in \ell$  and nontrivial nonperipheral non-boundary-parallel closed loops  $\gamma$  in  $\beta$ ,*

$$L^{-1} \leq \frac{|\varphi(\gamma)|'}{|\psi(\gamma)|'} \leq L, \quad (5.5.2)$$

$$|\varphi(\gamma)| \geq L^{-1}. \quad (5.5.3)$$

If  $\beta$  is a loop, then for all  $[\varphi], [\psi] \in \ell$ ,

$$|\operatorname{Re}(\pi_\alpha([\varphi]) - \pi_\alpha([\psi]))| \leq L. \quad (5.5.4)$$

Also if  $\gamma$  is in the interior of  $\alpha$ , and  $\ell_1 = [[\varphi_1], [\varphi_2]]$ , then given  $\varepsilon_1 > 0$  there exists  $\varepsilon_2 > 0$  depending only on  $\varepsilon_1$  and the ltd parameter functions and flat constant such that

$$\begin{aligned} &\text{If } |\varphi(\gamma)| < \varepsilon_2, \text{ then} \\ &\operatorname{Min}(|\varphi_1(\gamma)|, |\varphi_2(\gamma)|) \leq \varepsilon_1, \text{ and} \\ &\operatorname{Max}(|\varphi_1(\gamma)|, |\varphi_2(\gamma)|) \leq L. \end{aligned} \quad (5.5.5)$$

If (5.5.1), and either (5.5.2) and (5.5.3), or (5.5.4) hold for  $(\beta, \ell_1)$ , depending on whether  $\beta$  is a gap or a loop, we say that  $(\beta, \ell_1)$  is *bounded* (by  $L$ ). Note that  $L$  depends on the ltd parameter functions, and therefore is probably extremely large compared with  $\Delta(\nu)$  for many values of  $\nu$ , perhaps even compared with  $\Delta(\nu_0)$ .

Here are some notes on the proof. For fuller details, see 5.14 of [50]. First of all, under the assumption that  $\partial\alpha$  satisfies the condition (5.5.1), it is shown that  $\alpha$  is a union of  $\beta$  satisfying (5.5.1) to (5.5.3). First, we show that (5.5.2) holds for all  $\gamma \subset \alpha$  such that  $|\varphi_i(\gamma)|$  is bounded from 0 for  $i = 1, 2$ , and that (5.5.5) holds for  $\alpha$ . This is done by breaking  $\ell$  into three segments, with  $a'(\alpha)$  dominated by  $|\varphi(\partial\alpha)|_q^2$  on the two outer segments  $\ell_-, \ell_+$ , where  $q(z)dz^2$  is the quadratic differential at  $[\varphi]$  for  $\ell$ . The middle segment  $[[\varphi_-], [\varphi_+]]$  has to be of bounded length by the last part of 5.4, since there are no ltd's in  $\alpha$  along  $\ell$ . Then  $|\varphi(\partial\alpha)|_q$  is boundedly proportional to  $|\varphi(\partial\alpha)|_{q,-}$  along  $\ell_-$ , and to  $|\varphi(\partial\alpha)|_{q,+}$  along  $\ell_+$ . We can obtain (5.5.2) along  $\ell_+$ , at least for a nontrivial  $\alpha_1 \subset \alpha$  for which we can “lock” loops  $\varphi(\gamma)$ , for which  $|\varphi(\gamma)|$  is bounded, along stable segments to  $\varphi_+(\partial\alpha)$ . If  $\alpha_1 \neq \alpha$  and  $\gamma' \subset \partial\alpha_1$  is in the interior of  $\alpha$ , then either  $|\varphi_+(\gamma')|$  is small, or  $|\varphi_+(\gamma')|_{q_+}$  is dominated by  $|\varphi_+(\gamma')|_{q_+,-}$ , where  $q_+(z)dz^2$  is the stretch of  $q(z)dz^2$  at  $[\varphi_+]$ . In the case when  $|\varphi_+(\gamma')|$  is small, there is some first point  $[\varphi_{++}] \in \ell_+$  for which  $|\varphi_{++}(\gamma')|_{q_{++}}$  is dominated by  $|\varphi_{++}(\gamma')|_{q_{++},-}$ , where  $q_{++}(z)dz^2$  is the stretch of  $q(z)dz^2$  at  $[\varphi_{++}]$ . For this point,  $|\varphi_{++}(\gamma')|$  is still small, and can be locked to a small segment of  $\varphi_{++}(\partial\alpha)$ . This means that we can deduce that  $|\varphi_2(\gamma')|$  is small, giving (5.5.5). So one proceeds by induction on the topological type of  $\alpha$ , obtaining (5.5.2) and (5.5.5) for  $\alpha$  from that for  $\alpha \setminus \alpha_1$ . Then (5.5.5) and (5.5.2) imply that the set of loops with  $|\varphi_1(\gamma)| < \varepsilon_1$  or  $|\varphi_2(\gamma)| < \varepsilon_1$ , for a sufficiently small  $\varepsilon_1$ , do not intersect transversally. This allows for a decomposition into sets  $\beta$  satisfying (5.5.1), (5.5.2) and (5.5.3). One then has to remove the hypothesis (5.5.1) for  $\partial\alpha$ . This is done by another induction, considering successive gaps and loops  $\alpha'$  disjoint from all ltd's along segments  $\ell'$  of  $\ell$ , with  $|\varphi(\partial\alpha')| \leq \varepsilon_0$  for  $[\varphi] \in \ell'$ , possibly with  $\partial\alpha' = \emptyset$ . One then combines the segments and reduces the corresponding  $\alpha'$ , either combining two at a time, or a whole succession together, if the  $\alpha'$  are the same along a succession of segments. In finitely many steps, one reaches  $(\alpha, \ell)$  finding in the process that  $\partial\alpha$  does satisfy (5.5.1).

As for showing that  $\alpha$  satisfies (5.5.4), that follows from the following lemma — which is proved in 15.13 of [50], but not formally stated. Note that if  $\alpha$  is

a loop,  $a'(\alpha, [\varphi])$  is constant for  $[\varphi]$  in a geodesic segment  $\ell$ , but  $a(\alpha, q)$  is proportional to  $|\varphi(\alpha)|_q^2$  (if  $q(z)dz^2$  is the quadratic differential at  $[\varphi]$  for  $\ell$ ), which has at most one minimum on the geodesic segment and otherwise increases or decreases exponentially with distance along the segment, depending on whether  $|\varphi(\alpha)|_q$  is boundedly proportional to  $|\varphi(\alpha)|_{q,+}$  or  $|\varphi(\alpha)|_{q,-}$ . So for any  $K_0$ , the set of  $[\varphi] \in \ell$  for which  $a'(\alpha) \geq K_0 a(\alpha, [\varphi])$  is a single segment, up to bounded distance. This motivates the following.

**Lemma 5.6.** *Given  $K_0$ , there is  $C(K_0)$  such that the following holds. Let  $\ell$  be any geodesic segment. Suppose that  $a'(\alpha) \leq K_0 a(\alpha, [\varphi])$  for all  $[\varphi] \in \ell$ . Then for all  $[\varphi], [\psi] \in \ell$ ,*

$$|\operatorname{Re}(\pi_\alpha([\varphi]) - \pi_\alpha([\psi]))| \leq C(K_0).$$

*Proof.* The argument is basically given in 15.13 of [50]. Removing a segment of length bounded in terms of  $K_0, \varepsilon_0$  at one end, we obtain a reduced segment  $\ell'$  such that  $a'(\alpha) \leq \varepsilon_0 a(\alpha, [\varphi])$  for all  $[\varphi] \in \ell'$ . We use the quantity  $n_\alpha([\varphi])$  of 2.6, which is  $\operatorname{Re}(\pi_\alpha([\varphi])) + O(1)$  and is given to within length  $O(1)$  by  $m$  minimising  $|\varphi(\tau_\alpha^m(\zeta))|$  for a fixed  $\zeta$  crossing  $\alpha$  at most twice (or a bounded number of times). This is the same to within  $O(1)$  as the  $m$  minimising  $|\varphi(\tau_\alpha^m(\zeta))|_q$  for any quadratic differential  $q(z)dz^2$ . (To see this note that the shortest paths, in the Poincaré metric, across a Euclidean annulus  $\{z : r < |z| < 1\}$  are the restrictions of straight lines through the origin.) Assume without loss of generality that  $|\varphi(\alpha)|_q$  is boundedly proportional to  $|\varphi(\alpha)|_{q,+}$  for  $[\varphi] \in \ell$ , and  $q(z)dz^2$  the quadratic differential at  $[\varphi]$  for  $\ell$ . The good positions of  $\varphi(\tau_\alpha^m(\zeta))$  for all  $m$  are locked together along stable segments whose qd-length is short in comparison with  $|\varphi(\alpha)|_q$ , if  $\varepsilon_0$  is sufficiently small. So  $n_\alpha([\varphi])$  varies by  $< 1$  on  $\ell$ , and is thus constant on  $\ell'$ , if  $\varepsilon_0$  is sufficiently small, and hence varies by at most  $C(K_0)$  on  $\ell$ .  $\square$

## 5.7 Decomposing $S \times \ell$ .

Let  $\alpha_i \subset S$  be a gap or loop for  $i = 1, 2$ , isotoped so that  $\partial\alpha_1$  and  $\partial\alpha_2$  have only essential intersections, or with  $\alpha_1 \subset \alpha_2$  if  $\alpha_1$  is a loop which can be isotoped into  $\alpha_2$ . Then the *convex hull*  $C(\alpha_1, \alpha_2)$  of  $\alpha_1$  and  $\alpha_2$  is the union of  $\alpha_1 \cup \alpha_2$  and any components of  $S \setminus (\alpha_1 \cup \alpha_2)$  which are topological discs with at most one puncture. Then  $C(\alpha_1, \alpha_2)$  is again a gap or a loop. The latter only occurs if  $\alpha_1 = \alpha_2$  is a loop. We are only interested in the convex hull up to isotopy, and it only depends on  $\alpha_1$  and  $\alpha_2$  up to isotopy. It is so called because, if  $\alpha_i$  is chosen to have geodesic boundary, and  $\tilde{\alpha}_i$  denotes the preimage of  $\alpha_i$  in the hyperbolic plane covering  $S$ , then up to isotopy  $C(\alpha_1, \alpha_2)$  is the projection of the convex hull of any component of  $\tilde{\alpha}_1 \cup \tilde{\alpha}_2$ .

The following version of 5.5 will be important in constructing the geometric model. It follows directly from the statement of 5.5.

**Lemma** *Fix ltd parameter functions  $(\Delta, r, s, K_0)$ , and an associated constant  $\nu_0$  as in 5.4, and  $L = L(\Delta, r, s, K_0, \nu_0)$  as in 5.5. Let  $\ell$  be any geodesic segment*

in  $\mathcal{T}(S)$ . Then we can write  $S \times \ell$  as

$$S \times \ell = \cup_{j=1}^R \alpha_j \times \ell_j$$

where each  $\alpha_i \times \ell_j$  is either bounded by  $L$ , or long  $\nu$ -thick and dominant along  $\ell_j$  for  $(\Delta, r, s)$  and some  $\nu \geq \nu_0$ , or  $K_0$  flat along  $\ell_j$ , depending on whether  $\alpha_j$  is a gap or a loop.

In addition the decomposition is vertically efficient in the following sense.

1. If  $\gamma \subset \partial\alpha_j$  or  $\gamma = \alpha_j$  for some  $j$ , then  $\ell_j$  is contained in a connected union  $\ell' = [x, y]$  of segments  $\ell_k$  such that  $\gamma \subset \partial\alpha_k$ , and  $\gamma$  is in the convex hull of those  $\alpha_m$  for which  $(\alpha_m, \ell_m)$  is ltd and  $\ell_m \subset \ell'$ .
2. If  $\ell_j$  and  $\ell_k$  intersect precisely in an endpoint, and  $\alpha_j$  and  $\alpha_k$  have essential intersections, then there is no gap or nontrivial nonperipheral loop  $\beta \subset \alpha_j \cap \alpha_k$ .
3. For any  $\gamma$  and  $\ell' = [x, y]$  as in 1, either  $x$  is an endpoint of  $\ell$ , or  $x \in \ell_p$  for some ltd  $(\alpha_p, \ell_p)$  such that  $\gamma$  intersects  $\alpha_p$  essentially, and similarly for  $y$ .

*Proof.* Choose any disjoint set of  $\alpha_j \times \ell_j$  ( $1 \leq j \leq R_0$ ) such that the complement of the union contains no ltd, and such that for every  $(\alpha_j, \ell_j)$  and  $x \in \ell_j$ ,  $\gamma \subset \partial\alpha_j$ , there is an ltd  $(\alpha_k, \ell_k)$  with  $x \in \ell_k$  and  $\gamma \subset \partial\alpha_k$ . Then condition 1 is satisfied. By 5.5, for every  $(\alpha_j, \ell_j)$  which is not ltd,  $\alpha_j$  is a disjoint union of  $\beta$  such that  $(\beta, \ell_j)$  is bounded. If we can refine this partition to satisfy conditions 2 and 3, then every  $(\beta, \ell')$  in the complement of the ltds will automatically be bounded by 5.5 (especially (5.5.5)), because conditions 2 and 3 will ensure that there is no  $\gamma$  in the interior of  $\beta$  with  $|\varphi(\gamma)| < \varepsilon_2$  for  $[\varphi]$  an endpoint of  $\ell$ , at least if we take  $\varepsilon_1$  small enough given the ltd parameter functions.

Then we modify the partition in finitely many steps, always keeping condition 1, until condition 2 is satisfied. We do this as follows. Suppose we have a partition  $\mathcal{P}$  into ltd and bounded sets, satisfying 1 of the vertically efficient conditions, and there are  $\ell_k$  and  $\ell_m$  intersecting in precisely one point and such that  $(\alpha_k, \ell_k)$  and  $(\alpha_m, \ell_m)$  are bounded and  $\alpha_k \cap \alpha_m$  contains a  $\beta$  as is disallowed in 2 of vertically efficient. Then we can take  $\beta$  to be a maximal union of components of  $C(\partial\alpha_k, \partial\alpha_m)$ .

Then rewrite

$$\alpha_k \times \ell_k \cup \alpha_m \times \ell_m = ((\alpha_k \setminus \beta) \times \ell_k) \cup ((\alpha_m \setminus \beta) \times \ell_m) \cup (\beta \times (\ell_k \cup \ell_m)).$$

By 5.5, the  $(\alpha, \ell')$  arising in this rewriting are still bounded for the same  $L$ . Since rewriting reduces the topological type of the surfaces involved, and no new endpoints of segments  $\ell_j$  are introduced, finitely many rewritings gives a partition satisfying 1 and 2 of vertically efficient. Finally, to get 3 of vertically efficient, if  $\gamma \times \ell'$  as in 1 of vertically efficient does not have endpoints as required by 3, we extend  $\ell'$  through adjacent  $\ell_p$  with  $(\alpha_p, \ell_p)$  bounded, possibly joining up such segments, until endpoints are in ltds intersecting  $\gamma$  essentially, as required.

□

The pairs  $(\alpha_j, \ell_j)$  in the above are not unique. For example, as already noted, it is possible for  $(\alpha_j, \ell_j)$  to be both ltd and bounded, because the constant  $L$  of 5.5 is typically much bigger than  $\Delta(\nu_0)$ , for  $\nu_0$  as in 5.4.

## 6 Long, thick and dominant ideas.

This is a rather long section, which is pure theory of Teichmüller geodesics, with no input from three-dimensional hyperbolic geometry. It does seem necessary to go through some of these results in some detail, where they have not previously appeared in [50], or not in the same forms as given here. For understanding the proof of the Ending Laminations Theorem in the case of combinatorially bounded geometry, the parts most obviously needed are: the first theorem in 6.4, subsection 6.9, and the first lemma in 6.13. We also make use of Lemma 6.2 at one point, in the case of the long  $\nu$ -thick and dominant  $\alpha$  being the whole surface  $S$ . However, we also, at one point, make explicit use of the theorem in 6.5 — which is a deduction from the main theorem in 6.4. These two results are about general Teichmüller geodesics, not confined to the thick part of Teichmüller space. Thus, even the proof in the case of combinatorial bounded geometry explicitly relies, at one point, on the theory of general Teichmüller geodesics. This should not be a surprise, because, historically, the case of combinatorial bounded geometry is highly nontrivial. I regard the most difficult result in the whole paper as 6.14, which like the rest of this section, is purely about Teichmüller geodesics. This may be in some contrast to the experience of others who have worked on the Ending Laminations Theorem. I shall comment on this later.

### 6.1 Fundamental dynamical lemma.

The whole of the theory of ltd gaps and loops is based on a simple dynamical lemma which quantifies density of leaves of the stable and unstable foliations of a quadratic differential. This is basically 15.11 of [50], where three alternative conclusions are given. Here is a statement assuming the gap  $\alpha$  is ltd at  $[\varphi]$ .

**Lemma** *Given  $\delta > 0$ , the following holds for suitable ltd parameter functions  $(\Delta, r, s, K_0)$  and for a suitable function  $L(\delta, \nu)$ . Let  $\alpha$  be a gap which is long  $\nu$ -thick and dominant along a segment  $\ell = [[\varphi_1], [\varphi_2]]$  and let  $[\varphi] \in \ell$  with  $d([\varphi], [\varphi_1]) \geq \Delta(\nu)$ . Let  $q(z)dz^2$  be the quadratic differential at  $[\varphi]$  for  $d([\varphi], [\varphi_2])$  with stable and unstable foliations  $\mathcal{G}_\pm$ . Let  $a = a(\alpha, q)$ . Then there is no segment of the unstable foliation  $\mathcal{G}_+$  of qd-length  $\leq 2L(\nu, \delta)\sqrt{a}$  with both ends on  $\varphi(\partial\alpha)$ , and every segment of the unstable foliation  $\mathcal{G}_+$  of qd-length  $\geq L(\nu, \delta)\sqrt{a}$  in  $\varphi(\alpha)$  intersects every segment of  $\mathcal{G}_-$  of length  $\geq \delta\sqrt{a}$ . Similar statements hold with the role of stable and unstable reversed.*

## 6.2 Loops cut the surface into cells.

Now we give some of the key results about long thick and dominants which we shall need. We start with two fairly simple results, both of which follow directly from 6.1. These properties are used several times in [50], but may never be explicitly stated. The first may be reminiscent of the concept of tight geodesics in the curve complex developed by Masur and Minsky [28], and the point may be that these occur “naturally” in Teichmüller space

**Lemma** *Given  $L > 0$ , there is a function  $\Delta_1(\nu)$  depending only on the topological type of  $S$ , such that the following holds for suitable parameter functions  $(\Delta, r, s, K_0)$ . Let  $\alpha$  be a gap which is long  $\nu$ -thick and dominant along  $\ell$  for  $(\Delta, r, s, K_0)$ , with  $\Delta(\nu) \geq \Delta_1(\nu)$ . Let  $y_1 = [\varphi_1]$ ,  $y_2 = [\varphi_2] \in \ell$  with  $d(y_1, y_2) \geq \Delta_1(\nu)$ . Let  $\gamma_i \subset \alpha$  with  $|\varphi_i(\gamma_i)| \leq L$ ,  $i = 1, 2$ . Then  $\alpha \setminus (\gamma_1 \cup \gamma_2)$  is a union of topological discs with at most one puncture and topological annuli parallel to the boundary. Furthermore, for a constant  $C_1 = C_1(L, \nu)$ ,*

$$\#(\gamma_1 \cap \gamma_2) \geq C_1 \exp d(y_1, y_2).$$

*Proof.* Let  $[\varphi]$  be the midpoint of  $[[\varphi_1], [\varphi_2]]$  and let  $q(z)dz^2$  be the quadratic differential for  $d([\varphi], [\varphi_2])$  at  $[\varphi]$ . Because  $|\psi(\gamma_i)| \geq \nu$  for all  $[\psi] \in [[\varphi_1], [\varphi_2]]$ , by 5.2, the good position of  $\varphi_1(\gamma_1)$  satisfies

$$|\varphi_1(\gamma_1)|_{q,+} \geq C(L, \nu) \sqrt{a(\alpha, q)},$$

and similarly for  $|\varphi_2(\gamma_2)|_{q,-}$ . So

$$|\varphi(\gamma_1)|_{q,+} \geq C(L, \nu) e^{\Delta_1(\nu)/2} \sqrt{a(\alpha, q)},$$

$$|\varphi(\gamma_2)|_{q,2} \geq C(L, \nu) e^{\Delta_1(\nu)/2} \sqrt{a(\alpha, q)},$$

Then 6.1 implies that, given  $\varepsilon$ , if  $\Delta_1(\nu)$  is large enough given  $\varepsilon$ ,  $\varphi(\gamma_1)$  cuts every segment of stable foliation of  $q(z)dz^2$  of qd-length  $\geq \varepsilon \sqrt{a(\alpha)}$  and  $\varphi(\gamma_2)$  cuts every segment of unstable foliation of  $q(z)dz^2$  of qd-length  $\geq \varepsilon \sqrt{a(\alpha)}$ . So components of  $\varphi(\alpha) \setminus (\varphi(\gamma_1) \cup \varphi(\gamma_2))$  have Poincaré diameter  $< \nu$  if  $\Delta_1(\nu)$  is sufficiently large, and must be topological discs with at most one puncture or boundary-parallel annuli.

The last statement also follows from 6.1. If  $d(y_1, y_2) < \Delta_1(\nu)$ , there is nothing to prove, so now assume that  $d(y_1, y_2) \geq \Delta_1(\nu)$ . It suffices to bound below the number of intersections of  $\varphi(\gamma_1)$  and  $\varphi(\gamma_2)$ . Let  $L(\nu, 1)$  be as in 6.1, and assume without loss of generality that  $L(\nu, 1) \geq 1$ . Suppose that  $\Delta_1(\nu)$  is large enough that each of  $\varphi(\gamma_1)$  and  $\varphi(\gamma_2)$  contains at least one segment which is a qd distance  $\leq \sqrt{a}/L(\nu, 1)$  from unstable and stable segments, respectively, of qd-length  $\geq \sqrt{a}L(\nu, 1)$ . Note that the number of singularities of the quadratic differential is bounded in terms of the topological type of  $S$ . So apart from length which is a bounded multiple of  $L(\nu, 1)\sqrt{a}$ , each of  $\varphi(\gamma_1)$  and  $\varphi(\gamma_2)$  is a union of such segments. Then applying 6.1, each such segment of  $\varphi(\gamma_1)$  intersects each such segment on  $\varphi(\gamma_2)$ . So we obtain the result for  $C_1 = c_0 L(\nu, 1)^{-2}$ , for  $c_0$  depending only on the topological type of  $S$ .  $\square$



### 6.3 A partial order on $\text{ltd}(\beta, \ell)$ .

**Lemma** For  $i = 1, 3$ , let  $y_i = [\psi_i] \in \ell_i$ , and let  $\beta_i$  be a subsurface of  $S$  with  $|\psi_i(\partial\beta_i)| \leq L$ . Let  $\text{ltd}$  parameter functions be suitably chosen given  $L$ . Let  $\ell_2 \subset [y_1, y_3]$  and let  $\beta_2 \cap \beta_i \neq \emptyset$  for both  $i = 1, 3$ , and let  $\beta_2$  be  $\text{ltd}$  along  $\ell_2$ . Then  $\beta_1 \cap \beta_3 \neq \emptyset$ , and  $\beta_2$  is in the convex hull of  $\beta_1$  and  $\beta_3$ .

*Proof.* This is obvious unless both  $\partial\beta_1$  and  $\partial\beta_3$  intersect the interior of  $\beta_2$ . So now suppose that they both do this. First suppose that  $\beta_2$  is a gap and long,  $\nu$ -thick and dominant. Let  $y_{2,1} = [\psi_{2,1}]$ ,  $y_{2,3} = [\psi_{2,3}] \in \ell_2$  with  $y_{2,i}$  separating  $\ell_i$  from  $y_2$ , with  $y_{2,i}$  distance  $\geq \frac{1}{3}\Delta(\nu)$  from the ends of  $\ell_2$  and from  $y_2$ . If  $\beta_2$  is a loop, then we can take these distances to be  $\geq \frac{1}{6}\log K_0$ . For  $[\psi] \in [y_-, y_+]$ , let  $\psi(\beta)$  denote the region bounded by  $\psi(\partial\beta)$  and homotopic to  $\psi(\beta)$ , assuming  $\psi(\partial\beta)$  is in good position with respect to the quadratic differential at  $[\psi]$  for  $[y_-, y_+]$ . Then if  $\beta_2$  is a gap,  $\psi_{2,1}(\partial\beta_1 \cap \beta_2)$  includes a union of segments of in approximately unstable direction, of Poincaré length bounded from 0, and similarly for  $\psi_{2,3}(\partial\beta_3 \cap \beta_2)$ , with unstable replaced by stable. Then as in 6.2,  $\psi_2(\partial\beta_3 \cap \beta_2)$  and  $\psi_2(\partial\beta_1 \cap \beta_2)$  cut  $\psi_2(\beta_2)$  into topological discs with at most one puncture and annuli parallel to the boundary. It follows that  $\beta_2$  is contained in the convex hull of  $\beta_1$  and  $\beta_3$ . If  $\beta_2$  is a loop it is simpler. We replace  $\psi(\beta_2)$  by the maximal flat annulus  $S([\psi])$  homotopic to  $\psi(\beta_2)$ , for  $[\psi] \in \ell_2$ . Then  $\psi_2(\partial\beta_1) \cap S([\psi_2])$  is in approximately the unstable direction and  $\psi_2(\partial\beta_3) \cap S([\psi_2])$  in approximately the stable direction. They both cross  $S([\psi_2])$ , so must intersect in a loop homotopic to  $\psi_2(\beta_2)$ .  $\square$

We define  $(\beta_1, \ell_1) < (\beta_2, \ell_2)$  if  $\ell_1$  is to the left of  $\ell_2$  (in some common geodesic segment) and  $\beta_1 \cap \beta_2 \neq \emptyset$ . We can make this definition for any segments in a larger common geodesic segment, and even for single points in a common geodesic segment. So in the same way we can define  $(\beta_1, y_1) < (\beta_2, \ell_2)$  if  $y_1$  is to the left of  $\ell_2$ , still with  $\beta_1 \cap \beta_2 \neq \emptyset$ , and so on. This ordering is transitive restricted to  $\text{ltd}$ 's  $(\beta_i, \ell_i)$  by the lemma.

### 6.4 Triangles of geodesics.

The concept of long thick and dominant was mainly developed in order to formulate results about triangles of geodesics in  $\mathcal{T}(S)$ . The following theorem was proved in 15.8 of [50] in the case of  $S$  being a punctured (or marked) sphere. The proof is in fact completely general, once the approximate product structure of the thin part of  $\mathcal{T}(S)$  has been formalised, as we did in Section 2. Before we state the general theorem, we state it in the special case of a geodesic segment  $[y_0, y_1] \subset \mathcal{T}_{\geq \nu}$ .

**Triangle Theorem** (special case). *There exists a function  $C : (0, \infty) \rightarrow (0, \infty)$  such that the following holds. Let  $[y_0, y_1] \subset \mathcal{T}(S)_{\geq \nu}$ . Let  $y_2 \in \mathcal{T}(S)$ . Then for all  $y \in [y_0, y_1]$ , there exists  $y' \in [y_0, y_2] \cup [y_1, y_2]$  such that  $d(y, y') \leq C(\nu)$ . If  $y' \in [y_0, y_2]$  and  $w \in [y_0, y]$  then the corresponding  $w'$  is in  $[y_0, y']$ , and similarly if  $y' \in [y_1, y_2]$ .*

**Triangle Theorem** (general case). *There are functions  $C : (0, \infty) \rightarrow (0, \infty)$ ,  $\Delta_1 : (0, \infty) \rightarrow (0, \infty)$ ,  $\nu_1 : (0, \infty) \rightarrow (0, \infty)$  and constants  $L_0, L_1$  such that*

the following holds for suitable parameter functions  $(\Delta', r', s', K'_0)$ , and for ltd parameter functions  $(\Delta, r, s, K_0)$  given  $(\Delta', r', s', K'_0)$ .

Let  $y_0, y_1, y_2 \in \mathcal{T}(S)$  with  $y_j = [\varphi_j]$ . Take any  $y = [\varphi] \in [y_0, y_1]$ .

1. Let  $\alpha$  be a loop which is  $K_0$ -flat on  $\ell \subset [y_0, y_1] \subset \mathcal{T}$ . Then  $\ell$  is a union of two disjoint segments  $\ell_0$  and  $\ell_1$ , and there are segments  $\ell'_j \subset [y_j, y_2]$  such that for all  $y \in \ell_j$  there is  $y' = [\varphi']$ ,  $\ell'_j$  such that

$$|\varphi'(\alpha)| \leq L_0,$$

$$|\operatorname{Re}(\pi_\alpha(y)) - \operatorname{Re}(\pi_\alpha(y'))| \leq L_0,$$

and either  $\alpha$  is  $K'_0$ -flat along  $\ell'_j$ , or  $\ell_j$  and  $\ell'_j$  have length  $\leq L_1$ .

2. Let  $\alpha$  be a long  $\nu$ -thick and dominant gap along  $\ell \subset [y_0, y_1] \subset \mathcal{T}$  for  $(\Delta, r, s)$ . Then  $\ell$  is a union of two disjoint segments  $\ell_0$  and  $\ell_1$ , and there are  $\ell'_j \subset [y_j, y_2]$  such that for all  $y \in \ell_j$  there is  $y' = [\varphi'] \in \ell_j$ , with  $y' \in \ell'_j \subset [y_j, y_2]$  such that

$$|\varphi'(\partial\alpha)| \leq L_0,$$

$$d_\alpha(y, y') \leq C(\nu).$$

and either  $\alpha$  is long,  $\nu'$ -thick and dominant along  $\ell'_j$  for  $(\Delta', r', s')$  and some  $\nu' \geq \nu_1(\nu)$  or  $\ell_j$  and  $\ell'_j$  have length  $\leq \Delta_1(\nu)$ .

3. If  $y \in \ell_j$  and  $y'$  are as in either 1 or 2 above and  $\lambda \subset [y, y_j]$ , with  $\beta, \lambda$ , satisfying the conditions of  $\alpha, \ell$ , in 1 or 2 above, and  $\alpha \cap \beta \neq \emptyset$ , then  $\lambda = \lambda_j$ , where  $\lambda_j$  are defined relative to  $\lambda$  as the  $\ell_j$  to  $\ell$ .

This result extends to larger cycles of geodesic segments in  $\mathcal{T}(S)$ . If  $y_i \in \mathcal{T}(S)$  for  $0 \leq i \leq n$ , then we can consider the geodesic segments  $[y_i, y_{i+1}]$  for  $0 \leq i < n$  and  $[y_0, y_n]$ . Then for any ltd  $(\alpha, \ell)$  along  $[y_0, y_1]$ , we get a corresponding result to the above relative to a decomposition of  $\ell$  into sets  $\ell_j$  for  $j = 0$  or  $1 \leq j \leq n$  with  $\ell_j \subset [y_j, y_{j+1}]$  for  $j > 0$  and  $\ell_0 \subset [y_0, y_n]$ . Typically, one expects all but one of the  $\ell_j$  to be empty, but they could all be nonempty. The result is generalised by considering a decomposition into triangles, for example, triangles with vertices at  $y_0, y_j$  and  $y_{j+1}$  for each  $1 \leq j < n$ .

## 6.5 Only coordinates matter.

The ltd's which occur along a geodesic segment, up to bounded distance, are often determined by only some coordinates of the ends of the geodesic segment. This is what the following theorem says.

**Theorem** Given  $L_1$  and suitable ltd parameter functions  $(\Delta, r, s, K_0)$ , there is  $L_2$  such that the following holds. Let  $y_j = [\varphi_j]$ ,  $y'_j = [\varphi'_j] \in \mathcal{T}(S)$ ,  $j = 0, 1$ . Suppose that there are gaps or loops  $\alpha_j$  such that

$$|\varphi_j(\alpha_j)| \leq L_1, \quad |\varphi'_j(\alpha_j)| \leq L_1,$$

and

$$d_{\alpha_j}(y_j, y'_j) \leq L_1 \text{ or } |\operatorname{Re}(\pi_{\alpha_j}(y_j) - \pi_{\alpha_j}(y'_j))| \leq L_1.$$

Then we have the following.

Let  $\alpha$  be a loop or gap which is  $K_0$ -flat or long  $\nu$ -thick and dominant along  $\ell = [z_0, z_1] \subset [y_0, y_1]$ , with essential intersections with both  $\alpha_0$  and  $\alpha_1$  and such that

$$d'_{\alpha_j, \alpha}(y_j, z_j) \geq L_2.$$

Then in the conclusion of 6.4, but considering  $[y_0, y'_0] \cup [y'_0, y'_1] \cup [y'_1, y_1]$  instead of  $[y_0, y_2] \cup [y_2, y_1]$ , and with decompositions  $\ell = \ell_{0,0} \cup \ell_{0,1} \cup \ell_{1,1}$  instead of  $\ell_0 \cup \ell_1$ , we can take  $\ell_{0,0} = \ell_{1,1} = \emptyset$ , so that there is a corresponding segment  $\ell'$  on  $[y'_0, y'_1]$  to all of  $\ell$ .

*Proof.* Suppose that we cannot take  $\ell_{0,0} = \emptyset$ . Then let  $\ell'_{0,0}$  be the corresponding segment on  $[y_0, y'_0]$ . Then by the following lemma, for a constant  $L_3$  depending only on the ltd parameter functions,

$$d_{\alpha_0}(y_0, y'_0) \geq d'_{\alpha_0, \alpha}(y_0, y) + d'_{\alpha_0, \alpha}(y'_0, y) - L_3,$$

giving a contradiction. Similarly  $\ell_{1,1} = \emptyset$ .  $\square$

**Lemma 6.6.** *The following holds for suitable ltd parameter functions  $(\Delta, r, s, K_0)$ , a function  $C : (0, \infty) \rightarrow (0, \infty)$  and constants  $C_0 > 0$ ,  $C'_0 > 0$ . Let  $y_i = [\varphi_i]$ ,  $0 \leq i \leq n$  and  $z_i$ ,  $1 \leq i < n$  be points on a geodesic segment in  $\mathcal{T}(S)$ , in the order  $y_0, y_1, z_1, \dots, y_{n-1}, z_{n-1}, y_n$ . Write  $\ell_i = [y_i, z_i]$  for  $1 \leq i < n$ . Let  $\alpha_i$  be gaps or loops such that  $|\varphi_i(\partial\alpha_i)| \leq L_0$  for all  $i$ . For  $i = 0$  or  $n$ , if  $\alpha_i$  is a gap, let  $|\varphi_i(\gamma)| \geq \varepsilon_0$  for all nontrivial nonperipheral non-boundary-parallel  $\gamma \subset \alpha_i$ . For  $1 \leq i < n$ , let  $\alpha_i$  be either a gap which is long  $\nu_i$ -thick and dominant along  $\ell_i$ , or let  $\alpha_i$  be a loop which is  $K_0$ -flat along  $\ell_i$ . Let  $\alpha_i \cap \alpha_{i+1} \neq \emptyset$  for  $0 \leq i < n$ . Write  $C_i = C(\nu_i)$  for  $0 < i < n$  if  $\alpha_i$  is a gap and  $C_i = 2C_0$  if  $\alpha_i$  is a loop. Then*

$$d'_{\alpha_0, \alpha_n}(z_0, z_n) \geq \sum_{i=0}^{n-1} (d_{\alpha_i}(y_i, z_i) + d'_{\alpha_i, \alpha_{i+1}}(z_i, y_{i+1}) - C_i).$$

*Proof.* This is done by a *locking technique* which is used frequently in [50]. By locking, we mean the following. Suppose that  $\beta$  and  $\beta'$  are two paths in  $\psi(S)$  and some quadratic differential  $q(z)dz^2$  is fixed. Then we say that  $x' \in \beta'$  is *locked* to  $x \in \beta$  (for some fixed  $\delta > 0$ ) if there is an arc of stable foliation between  $x$  and  $x'$  of Poincaré length  $\leq \delta$  times the injectivity radius at  $\psi(S)$ , also measured in Poincaré length. If  $\delta$  is sufficiently small, depending only on the topological type of  $S$ , then we can equally well use the qd-length to measure this ratio. This means there is a constant  $C(S) > 0$  such that if an arc has length  $\leq \delta \leq C(S)^{-1}$  times the injectivity radius, with both measured in the Poincaré metric then the qd-length is  $\leq C(S)\delta$  times the injectivity radius, also measured in the qd-metric, and similarly with Poincaré metric and qd-metric interchanged. Let  $\chi_t$  denote the family of homeomorphisms obtained by scaling unstable and stable length for  $q(z)dz^2$  by  $e^{\pm t}$ . If  $\beta'$  is locked to  $\beta$ , then  $\chi_t(\beta')$  is locked to  $\chi_t(\beta)$  for all  $t > 0$ , because qd-length of locking segments get multiplied

by  $e^{-t}$ , while the injectivity radius, measured in qd-length, cannot decrease by more than  $e^{-t}$ . A technique developed in [50] for showing one loop was much longer than another was to show that one loop had many points locked to each point on the other.

If  $\alpha_i$  is a loop, put  $\gamma_i = \gamma'_i = \alpha_i$ . Now suppose that  $\alpha_i$  is a gap. For some  $L_1(\nu)$  depending only on  $\nu$  (and the topological type of  $S$ ), for each  $[\varphi] \in \ell_i$  there is a loop  $\gamma \subset \alpha_i$  such that

$$\nu_i \leq |\varphi(\gamma)| \leq L_1(\nu_i).$$

By 6.1, given  $\delta > 0$  there are then  $\Delta_1(\nu)$  and  $L_2(\nu_i)$  depending only on  $\nu$  and  $L_1(\nu)$  such that if  $d([\varphi], z_i) \geq \Delta_1(\nu_i)$ , and  $[\varphi] \in \ell_i$ , then for each  $x \in \varphi(\gamma)$  and each point  $x' \in \varphi(\alpha_i)$  where the injectivity radius is  $\geq \nu_i$ , there is a stable segment in  $\varphi(\alpha_i)$  starting from  $x$  and coming within  $\delta\nu_i$  of  $x'$ . From now on we assume that  $\Delta(\nu) > \Delta_1(\nu)$ , and sufficiently large in a sense to be determined.

If  $\alpha_i$  is a loop, then define  $\gamma_i = \gamma'_i = \alpha_i$ . In this case,  $\alpha_{i-1} \neq \alpha_i$ , except possibly if  $i = 1$ , and  $\alpha_{i+1} \neq \alpha_i$ , except possibly if  $i = n - 1$ . If  $\alpha_i$  is a gap, choose  $\gamma_i, \gamma'_i \subset \alpha_i$  such that,

$$\nu_i \leq |\varphi_i(\gamma_i)| \leq L_1(\nu_i),$$

$$\nu_i \leq |\psi_i(\gamma'_i)| \leq L_1(\nu_i),$$

$$C'|\varphi_{i+1}(\gamma'_i \cap \alpha_{i+1})| \geq \exp d'_{\alpha_i, \alpha_{i+1}}(z_i, y_{i+1}).$$

This last is possible by 2.9, and should be done for  $0 \leq i < n$ , for a suitable constant  $C'$ . In addition, choose  $\gamma_0 \subset \alpha_0$  so that

$$C'|\varphi_1(\gamma_0 \cap \alpha_1)| \geq \exp d'_{\alpha_0, \alpha_1}(y_0, y_1).$$

If  $\alpha_i$  is a gap, let  $z'_i = [\psi'_i] \in \ell_i$  with  $d(z'_i, z_i) = \Delta_1(\nu_i)$ , for  $\Delta_1(\nu)$  to be chosen as follows, and if  $\alpha_i$  is a loop, let  $z'_i = [\psi'_i] \in \ell_i$  with  $d(z'_i, z_i) = \Delta_1$ , for  $\Delta_1$  to be chosen as follows. By 7.2, if  $\alpha_i$  is a gap every point on  $\psi'_i(\gamma'_i)$  is locked to  $\geq C_2(\nu_i)^{-1} \exp d(y_i, z'_i)$  points on  $\psi_i(\gamma_i)$  along stable segments of Poincaré length  $\leq L_2(\nu_i)$ , and every point on  $\varphi_i(\gamma_i)$  is locked to  $\geq C'^{-1} \exp d'_{\alpha_{i-1}, \alpha_i}(z_{i-1}, y_i)$  points on  $\varphi_i(\gamma'_{i-1})$  along stable segments of Poincaré length  $\leq L_2(\nu_i)$ . Then assuming  $\Delta_1(\nu)$  is sufficiently large given  $L_2(\nu)$ , and  $\Delta(\nu)$  sufficiently large given  $\Delta_1(\nu)$ , every point on  $\psi_i(\gamma'_i)$  is locked to  $\geq C_2(\nu_i)^{-1} \exp d(y_i, z'_i)$  points on  $\psi_i(\gamma_i)$  along segments of Poincaré length  $\leq C' \cdot L_2(\nu_i) \cdot \exp -d_{\alpha_i}(y_i, z_i) < \delta\nu_i$  and  $\geq (C_2(\nu_i)C')^{-1} \exp d'_{\alpha_{i-1}, \alpha_i}(z_{i-1}, y_i) \cdot \exp d_{\alpha_i}(y_i, z'_i)$  points on  $\psi_i(\gamma'_{i-1})$  along stable segments of Poincaré length  $\leq C' \cdot L_2(\nu_i) \cdot \exp -d_{\alpha_i}(y_i, z_i) \leq \delta\nu_i$ .

If  $\alpha_i$  is a loop,  $\varphi_i(\gamma'_{i-1})$  intersects the flat annulus homotopic to  $\varphi_i(\alpha_i)$  in  $\geq C'^{-1} \cdot C \exp d'_{\alpha_{i-1}, \alpha_i}(z_{i-1}, y_i)$  segments in an approximately stable direction. The corresponding segments of  $\psi'_i(\gamma'_{i-1})$  each contain  $\geq C'^{-1} \exp d_{\alpha_i}(y_i, z'_i)$  disjoint segments which can be locked to  $\psi'_i(\alpha_i)$  along stable segments of Poincaré length  $\leq C' \cdot |\psi'_i(\alpha_i)|$ , for suitable  $C'$ . Then  $\psi_i(\gamma'_{i-1})$  contains  $\geq C'^{-2} \exp d'_{\alpha_{i-1}, \alpha_i}(z_{i-1}, y_i) \cdot \exp d_{\alpha_i}(y_i, z_i)$  segments which can be locked to  $\psi(\alpha_i)$  by stable segments of length  $\leq \delta \cdot |\psi(\alpha_i)|$ , assuming that  $\Delta_1$  is large enough given  $C'$  and  $\delta$ .

Define  $C_i = \log C'$  if  $i = 0$  or  $n$ . If  $0 < i < n$ , define  $C_i = \Delta_1(\nu_i) + \log C_2(\nu_i) + 3 \log C'$  if  $\alpha_i$  is a gap, and  $C_i = C'_0 = 4 \log C' + \log \Delta_1$  if  $\alpha_i$  is a loop. Define

$$\Delta_j = \sum_{i=1}^j (d'_{\alpha_{i-1}\alpha_i}(z_{i-1}, z_i) - C_i),$$

$$\Delta'_j = \sum_{i=1}^j (d'_{\alpha_{i-1}\alpha_i}(z_{i-1}, z_i) - C_i) - d_{\alpha_i}(y_j, z_j).$$

We claim inductively that, assuming the ltd parameter functions are strong enough, for  $j \geq 1$ , each point of  $\psi_j(\gamma'_j)$  is locked to  $\geq \exp \Delta_j$  points on  $\psi_j(\gamma'_0)$  along stable segments of Poincaré  $\leq 2\delta$  times the injectivity radius in the Poincaré metric, and similarly  $\varphi_j(\gamma_j)$  is locked to  $\geq \exp \Delta'_j$  points on  $\varphi_j(\gamma'_0)$  along stable segments of Poincaré  $\leq 2\delta$  times the injectivity radius in the Poincaré metric. As before, for  $\delta$  sufficiently small, it suffices to prove this using the qd-metric. But in the qd-metric,  $\chi_t$  multiplies qd-length along the stable direction by  $e^{-t}$ , while the injectivity radius in the qd-metric decreases by at most a factor  $e^{-t}$ . Relative ratios of lengths of locking segments are preserved up to a bounded proportion. So locking segments between  $\psi_k(\gamma'_i)$  and  $\psi_k(\gamma'_j)$  have Poincaré length  $\leq C'' \exp(\Delta_j - \Delta_k)$ , for  $i < j < k$ , for a suitable constant  $C''$ .

This gives the required estimate, apart from minor adjustments if  $\alpha_0$  or  $\alpha_n$  is a loop. In those cases, given the definition of  $d'_{\alpha_0, \alpha_n}$ , there is nothing to prove if  $\alpha_0 = \alpha_n$ , or if  $\alpha_0$  is not flat along any segment of  $[y_0, y_n]$ . If  $\alpha_0$  is along a segment of  $[y_0, y_n]$ , we can introduce another segment  $\ell_1$  if necessary, renumbering, so that  $\alpha_0$  is flat precisely along  $\ell_1$ . We make similar adjustments near  $y_n$  but otherwise the proof is exactly as above.

□

## 6.7 The graph of the qd-length function.

One of the basic technical considerations in the study of Teichmüller geodesics, as is probably already apparent, is the difference between the qd- and Poincaré metrics. The two metrics are not globally Lipschitz equivalent. But they are Lipschitz-equivalent, up to scalar, on any thick part of a surface. The Lipschitz constant is bounded in terms of the topological type of the surface, but the scalar is completely uncontrollable. This should not be regarded as a problem. One simply has to look at ratios of lengths rather than at absolute lengths. Also, the qd-length function has a rather remarkable property. Fix a Teichmüller geodesic  $\{[\chi_t \circ \varphi_0] : t \in \mathbb{R}\}$ , where  $\chi_t$  minimises distortion and  $d([\chi_t \circ \varphi_0], [\chi_s \circ \varphi_0]) = |t - s|$ . Let  $q_0(z)dz^2$  be the quadratic differential at  $[\varphi_0]$  for  $d([\varphi_0], [\chi_t \circ \varphi_0])$ , for  $t > 0$ , and  $q_t(z)dz^2$  the stretch at  $[\chi_t \circ \varphi_0]$  (2.3). Let  $|\cdot|_t = |\cdot|_{q_t}$ , the qd-length (5.1). For any finite loop set  $\gamma$ , define

$$F(t, \gamma) = \log |\chi_t \circ \varphi_0(\gamma)|_t.$$

By 14.7 of [50] (and I am sure this is well-known), there is a constant  $C_0$  depending only on the topological type of  $S$ , and a bound on the number of loops in  $\gamma$ , and there are  $c(\gamma), t(\gamma) \in \mathbb{R}$  such that

$$|F(t, \gamma) - |t - t(\gamma)| - c(\gamma)| \leq C_0. \quad (6.7.1)$$

The graph of the function  $t \mapsto F(t, \gamma)$ , for any  $\gamma$ , therefore lies within  $C_0$  of a  $V$ , with the slopes of the arms of the  $V$  being  $-1$  on the left and  $1$  on the right, and minimum at  $t(\gamma)$ .

Comparison between Poincaré and qd-length can then be made as follows. Given  $L_1 > 0$  there is  $L_2 \in \mathbb{R}$  such that

$$|\chi_t \circ \varphi_0(\gamma)| \leq L_1 \quad (6.7.2)$$

whenever

$$F(t, \gamma) - F(t, \gamma') \leq L_2 \quad (6.7.3)$$

for all nontrivial nonperipheral  $\gamma'$  intersecting  $\gamma$  transversely. Conversely, given  $L_2 \in \mathbb{R}$ , there is  $L_1$  such that (6.7.2) holds whenever (6.7.3) holds for all  $\gamma'$  intersecting  $\gamma$  transversely. There is a similar characterisation of short loops. Given  $L_2 < 0$ , there is  $L_1 > 0$  (small if  $L_2$  is negatively large) such that, whenever (6.7.2) holds, then (6.7.3) holds for all  $\gamma'$  intersecting  $\gamma$  transversely. Conversely, given  $L_1 > 0$ , there is  $L_2$  (negative if  $L_1$  is small) such that (6.7.2) holds for  $\gamma$ , whenever (6.7.3) holds for  $\gamma$  and all  $\gamma'$  transverse to  $\gamma$ .

If  $\gamma$  satisfies 6.7.3 for all transverse  $\gamma'$  and  $\gamma''$  is another loop, disjoint from  $\gamma$ , with  $|\chi_t \circ \varphi_0(\gamma'')|$  bounded, and  $F(t, \gamma) - F(t, \gamma'') \leq L_2$ , then it is possible that  $\chi_t \circ \varphi_0(\gamma'')$  is short, while  $\chi_t \circ \varphi_0(\gamma)$  is not. However, if  $\Gamma$  is a set of loops  $\gamma$ , satisfying 6.7.3, then any component  $\alpha$  of the convex hull of  $\Gamma$  is such that  $\chi_t \circ \varphi_0(\alpha)$  is contained in a single component of  $(\chi_t \circ \varphi_0(S))_{\geq \varepsilon(L_2)}$ , for a suitable  $\varepsilon > 0$  depending only on  $L_2$ . Conversely, if  $\chi_t \circ \varphi_0(\alpha)$  is a component of  $(\chi_t \circ \varphi_0(S))_{\geq \varepsilon}$  then we can find a set of loops  $\Gamma$  with convex hull  $\alpha$  such that  $\gamma$  satisfies (6.7.3) for all  $\gamma \in \Gamma$ , for a suitable  $L_2 = L_2(\varepsilon)$ .

## 6.8 Ltd's in the projection are the same.

We shall need the following.

**Theorem** *Given  $L_1$ , there is  $L_2$ , and given ltd parameter functions  $(\Delta', r', s', K'_0)$ , there are  $(\Delta, r, s, K_0)$ ,  $C_0$ ,  $\nu_0$ ,  $\nu'_0$  and  $C : (0, \infty) \rightarrow (0, \infty)$  such that the following holds. Let  $y_j = [\varphi_j]$ ,  $j = 0, 1$ , and let  $\alpha$  be a loop or gap with  $|\varphi_j(\alpha)| \leq L_1$  or  $|\varphi_j(\partial\alpha)| \leq L_1$  for  $j = 0, 1$ . Let  $\ell = [z_0, z_1] \subset [\pi_\alpha(y_0), \pi_\alpha(y_1)] \subset \mathcal{T}(\mathcal{S}(\alpha))$ , let  $\beta$  be long  $\nu$ -thick and dominant along  $\ell$  with respect to  $(\Delta, r, s)$  for some  $\nu \geq \nu_0$ , or  $K_0$ -flat along  $\ell$  and let  $\beta, z_j, \pi_\alpha(y_j), L'_1$  satisfy the conditions of 6.5 with  $\pi_\alpha(y_j)$  replacing  $y_j$ . Then there is  $\ell' \subset [y_0, y_1]$  such that  $\beta$  is long,  $\nu'$ -thick and dominant along  $\ell'$  with respect to  $(\Delta', r', s')$  and some  $\nu' \geq \nu'_0$  or  $K'_0$  flat along  $\ell'$ , and for each  $y \in \ell$  there is  $y' \in \ell'$  such that*

$$d_\beta(y, y') \leq C(\nu) \quad \text{or} \quad |\operatorname{Re}(\pi_\beta(y) - \pi_\beta(y'))| \leq C_0.$$

*Proof.* This is proved by similar techniques to 6.4, but since there is no precise statement like this in [50], we had better give some details. We can assume  $L_1$  is bounded and choose the ltd parameter functions relative to it, because we can then get the result for a general  $L_1$  using 6.5

Given  $\gamma \subset S$  and  $[\varphi] \in \mathcal{T}(S)$ , we say that  $\varphi(\gamma)$  is *almost bounded (by  $L$ )* if  $|\varphi(\gamma)|'' \leq L$ , where  $|\varphi(\gamma)|''$  is as in 2.5. A sufficient criterion for a loop to be almost bounded at some point on a geodesic segment  $[[\varphi_0], [\varphi_1]]$  is given by the negation of a necessary condition for a loop to be not bounded at any point of the geodesic segment, as follows. Given  $L$ , there is  $L'$  such that if, for all choices of disjoint simple loops  $\gamma'_0, \gamma'_1$  which both intersect  $\gamma$ , either

$$|\varphi_0(\gamma)| \leq L|\varphi_0(\gamma'_j)| \text{ for } j = 0, 1, \quad (6.8.1)$$

or

$$|\varphi_1(\gamma)| \leq L|\varphi_1(\gamma'_j)| \text{ for } j = 0, 1, \quad (6.8.2)$$

then there is a point  $[\varphi] \in [[\varphi_0], [\varphi_1]]$  such that  $|\varphi(\gamma)|'' \leq L'$ . It is not clear if (6.8.1) or (6.8.2) is a necessary condition for  $[\varphi]$  to be bounded for some  $[\varphi] \in [[\varphi_1], [\varphi_2]]$ , in general. But if  $[\varphi] \in [[\varphi_0], [\varphi_1]]$  and  $\gamma \subset \alpha$ , where  $\beta$  is ltd or flat along a segment of  $[[\varphi_0], [\varphi_1]]$  containing  $[\varphi]$ , and  $|\varphi(\gamma)| \leq L''$ , then for any choice of  $(\gamma'_0, \gamma'_1)$  as above for a suitable  $L$  given  $L''$ , for any choice of  $(\gamma'_0, \gamma'_1)$ , one of (6.8.1) or (6.8.2) holds. This is essentially the content of 15.8 of [50].

Now let  $\beta$  be a gap which is long,  $\nu$ -thick and dominant along  $\ell \subset [\pi_\alpha(y_0), \pi_\alpha(y_1)]$ . So for any loop  $\gamma \subset \beta$  and  $[\varphi] \in \ell$  such that  $|\varphi(\gamma)| \leq L$ , we can find  $[\varphi'] \in [y_0, y_1]$  such that  $|\varphi'(\gamma)|'' \leq L'$ . But we actually want a bound on  $|\varphi'(\gamma)|$ , and that  $\beta$  should be ltd along a segment containing  $[\varphi']$  for suitable parameter functions. This is done as follows. Take any  $w_i = [\psi_i] \in \ell$  and loop sets  $\Gamma_i \subset \beta$  which are cell-cutting in  $\beta$   $i = 0, 1, 2$  with  $|\psi_i(\Gamma_i)| \leq L$  such that  $|\psi_i(\Gamma_i)| \leq L$ , with  $w_1 \in [w_0, w_2]$  and

$$L_3 \leq d(w_i, w_{i+1}) \leq L_4,$$

where  $L_3$  is large enough for  $\zeta_i \cup \zeta_{i+1}$  to be cell-cutting in  $\beta$  for any  $\zeta_j \in \Gamma_j$  for  $i = 0, 1, 2$ , using 6.2. We shall also need  $L_3$  large enough for there to be no loop  $\zeta \subset \beta$  with  $|\psi_i(\zeta)| \leq L''$  for  $i = 0, 1$ , or both  $i = 1, i = 2$ , where  $L''$  depends only on  $L$  and  $L'$ . This is again possible using 6.2, for  $L_3$  depending only on  $L$  and  $L'$ . So now we fix this choice of  $L_3$ , and  $L_4$ . Using (6.8.1) and (6.8.2) as above, we have  $w_i = [\psi'_i] \in [y_0, y_1]$  with  $|\psi'_i(\Gamma_i)|'' \leq L'$ ,  $i = 0, 1, 2$ . Now again using 6.2, for suitable parameter functions  $(\Delta'', r'', s'')$  and flat constant  $K''_0$  strong enough given  $L_4$ , which bounds  $\#(\Gamma_0 \cap \Gamma_1)$  and  $\#(\Gamma_1 \cap \Gamma_2)$ , there cannot be any segment  $\lambda \subset [w'_i, w'_{i+1}]$  and  $\omega$  intersecting both  $\zeta_i$  and  $\zeta_{i+1}$  (any  $\zeta_j \in \Gamma_j$ ) along which  $\omega$  is long thick and dominant for  $(\Delta'', r'', s'')$  or  $K''_0$ -flat. Any  $\omega$  which intersects one of  $\zeta_i, \zeta_{i+1}$  intersects the other, since  $\zeta_i \cup \zeta_{i+1}$  is cell-cutting in  $\beta$ . So then let  $L_5$  be the constant  $L$  given by 5.5 relative to  $(\Delta'', r'', s'')$  and  $K''_0$ . Then there is  $\varepsilon > 0$  bounded below in terms of  $L_5$  so that if  $\zeta \subset \text{Int}(\beta)$  and  $|\psi'_1(\zeta)| < \varepsilon$  then  $|\psi'_i(\zeta)| < \varepsilon_0$  for either  $i = 0$  or  $i = 2$ ,

by (5.5.5). Now suppose there is such a loop  $\zeta$  in the interior of  $\beta$ . Suppose without loss of generality that  $|\psi'_0(\zeta)| \leq \varepsilon_0$ . Then  $\zeta$  has  $\leq L'$  intersections with each of  $\Gamma_0, \Gamma_1$ , by the definition of  $|\cdot|''$ , since the loops of  $\psi'_i(\Gamma_i)$  have  $|\cdot|''$ -length  $\leq L'$  for  $i = 0, 1$ . Since these loop sets are both cell-cutting in  $\beta$ , we deduce that  $|\psi_i(\zeta)| \leq L''$  for  $L''$  depending only on  $L$  and  $L'$ , and  $i = 0, 1$ . By the choice of  $L_3$  this is impossible. So this means we have a bound on  $|\psi'_1(\Gamma_1)|$  and we can take  $[\varphi'] = w'_1$ , if we take  $w_1 = y$ .

So we have  $\ell' \subset [y_0, y_1]$ , and, for each  $y \in \ell$ , we have  $y' \in \ell$  with  $d(y, y') \leq C(\nu)$ . Now we need to show that given  $(\Delta', r', s')$ ,  $\ell'$  is long  $\nu'$ -thick and dominant along  $\ell'$  for some  $\nu' \geq \nu/C(\nu)$ , if  $(\Delta, r, s)$  are suitably chosen. First, we note that because  $\partial\beta$  can be homotoped into  $\Gamma_0$  as above,  $|\psi(\partial\beta)| \leq C_1(\nu')$  for all  $[\psi] \in \ell'$ , and indeed of an extension  $\ell'_1$  of  $\ell'$  at both endpoints, if the ltd parameter functions  $(\Delta, r, s)$  are sufficiently strong. The  $d_\beta$ -lengths of  $\ell$  and  $\ell'$  differ by at most  $2C(\nu)$ , and there are similar properties for  $\ell'_1$ . The Poincaré length of  $\psi(\partial\beta)$  is bounded along  $\ell'_1$ . Now we need to show that the ratio  $a(\partial\beta, [\psi])/a(\beta)$  decreases exponentially in the middle of  $\ell'_1$ . The easiest way to see this is to make use of the functions  $F(t, \gamma)$  of 6.7, for the geodesic  $[y_0, y_1]$ . Write  $\ell'_1 = [[\chi_a \circ \varphi_0], [\chi_b \circ \varphi_0]]$ . Because the Poincaré length of  $\psi(\partial\beta)$  is bounded all along  $\ell'_1$ , and we have a lower bound of  $\nu'$  (with  $\nu' = \nu'(\nu)$ ) on the length of loops  $\psi(\zeta)$  for  $\zeta$  in the interior of  $\beta$ , there is a function  $C_2(\nu)$  such that

$$F(t, \partial\beta) \leq C_2(\nu') + F(t, \gamma) \quad (6.8.3)$$

for all  $\gamma$  in the interior of  $\beta$  and  $t \in [a, b]$ . The function  $F(\cdot, \gamma)$ , for any  $\gamma$  in the interior of  $\beta$  has minimum at most  $C_2(\nu')$  below the minimum of  $F(t, \partial\beta)$ , if the minimum is in  $[a, b]$ . But, by comparing with  $\ell_1$ , for a  $\gamma$  in the interior of  $\beta$  for which  $\chi_t \circ \varphi_0(\gamma)$  is bounded, and for  $T = \min(|t - a|, |t - b|)$ ,

$$|\chi_a \circ \varphi_0(\gamma)| \geq C_3(\nu') \cdot e^T,$$

and similarly for  $a$  replaced by  $b$ . It follows that, for such  $\gamma$ ,

$$F(a, \partial\beta) \leq F(a, \gamma) + C_4(\nu') - T,$$

and similarly with  $a$  replaced by  $b$ . It follows that

$$F(t, \partial\beta) \leq F(t, \gamma) - T + C_4(\nu') + C_0,$$

where  $C_0$  is the constant of (6.7.1). Then the good position of  $\chi_t \circ \varphi_0(\gamma)$  is bounded from the stable and unstable positions. Suppose that this  $t$  is such that the good position of  $\chi_t \circ \varphi_0(\partial\beta)$  is close to the stable foliation, or to the unstable foliation. These happen except on a bounded interval of  $t$ . Then the good position of  $\chi_t \circ \varphi_0(\beta)$  contains a ball of definite Poincaré radius centred on a point of  $\chi_t \circ \varphi_0(\gamma)$ . So  $|\chi_t \circ \varphi_0(\gamma)|_t$  is boundedly proportional to  $\sqrt{a(\beta)}$  and, for this  $t$ ,

$$|\chi_t \circ \varphi_0(\partial\beta)|_t \leq C_5(\nu') e^{-T} \sqrt{a(\beta)}.$$

Then, since this is true except on a bounded interval, it must be true for all  $t \in [a, b]$ , if we adjust the constant. If  $\ell_1$  and  $\ell'_1$  are sufficiently long, that



is, if the parameter functions  $(\Delta, r, s)$  are sufficiently strong, we can ensure that the qd-length of  $\psi(\partial\beta)$  along  $\ell'$  is  $\leq s(\nu)\sqrt{a(\beta)}$ . Then we also have the Poincaré length of  $\psi(\partial\beta)$  is  $< r(\nu)$ , assuming without loss of generality that  $s$  is sufficiently strong given  $r$ . The bound on  $d_\beta(y, y')$  for all  $y$  means that  $\beta$  is  $\nu'$ -thick along  $\ell'$  for some  $\nu'$  depending only on  $\nu$ . So altogether, by suitable choice of  $(\Delta, r, s)$  given  $(\Delta', r', s')$  we can ensure that  $\beta$  is long,  $\nu'$ -thick and dominant along  $\ell'$  for  $(\Delta', r', s')$ .

Finally, let  $\beta$  be a loop. By 5.6, the quantity  $n_\beta([\varphi])$  only changes for  $[\varphi] \in \ell$ . We have

$$n_\beta(z_j) = n_\beta(\pi_\alpha(y_j)) + O(1) = n_\beta(y_j) + O(1)$$

So

$$n_\beta(y_1) - n_\beta(y_0) = n_\beta(z_1) - n_\beta(z_0) + O(1) = |\ell| + O(1).$$

The only way to achieve this is if there is  $\ell' = [w_0, w_1] \subset [y_0, y_1]$  along which  $\beta$  is  $K_0$ -flat and with  $n_\beta(w'_j) = n_\beta(y_j) + O(1)$ . Then assuming  $L_0$  is suitable chosen we do indeed have, for each  $y \in \ell$ , a corresponding  $y' \in \ell'$  with

$$|\operatorname{Re}(\pi_\beta(y') - \pi_\beta(y))| \leq C_0.$$

□

## 6.9 “Orthogonal projection” for geodesics in $\mathcal{T}_{\geq \nu}$ .

Now we describe an analogue of orthogonal projection for a geodesic segment in  $\mathcal{T}(\mathcal{S})$ . In order to describe the idea, we first consider the definition for a geodesic segment

$$[y_-, y_+] \subset \mathcal{T}_{\geq \nu}$$

for a fixed  $\nu > 0$ . In this case we define

$$x = x(., [y_0, y_1]) : \mathcal{T}(\mathcal{S}) \rightarrow [y_0, y_1]$$

as follows. Take any  $z \in \mathcal{T}(\mathcal{S})$ . Then by the special case Triangle Theorem of 6.4, there is  $y \in [y_-, y_+]$ , unique up to moving it a bounded distance in  $\mathcal{T}(\mathcal{S})$ , such that there exist  $y' \in [y_-, z]$  and  $y'' \in [y_+, z]$  such that, for  $C(\nu)$  as in the special case Triangle Theorem

$$d(y, y') \leq C(\nu), \quad d(y, y'') \leq C(\nu).$$

We then choose such a  $y$  for each  $z$  and define

$$x(z) = x(z, [y_-, y_+]) = y.$$

The function  $x$  is not continuous (unless we are more careful with the definition, at least), but it is coarse Lipschitz, and hence, by a coarse Intermediate Value Theorem (since  $x(y_-) = y_-$  and  $x(y_+) = y_+$ ) coarsely surjective onto  $[y_-, y_+]$  along any path in  $\mathcal{T}(\mathcal{S})$  joining  $y_-$  and  $y_+$ .

**Lemma** If  $[y_-, y_+] \subset (\mathcal{T}(S))_{\geq \nu}$  then there are  $L(\nu)$  and  $C_1(\nu)$  such that for any  $y_1, y_2 \in \mathcal{T}(S)$ , if  $d(x(y_1), x(y_2)) \geq L(\nu)$ , with  $x(y_1)$  nearer  $y_-$  than  $x(y_2)$ , then there are  $y'_1, y'_2 \in [y_1, y_2]$  with

$$d(y'_i, x(y_i)) \leq C_1(\nu),$$

giving

$$d(y_1, y_2) \geq d(y_1, x(y_1)) + d(y_2, x(y_2)) + d(x(y_1), x(y_2)) - 4C_1(\nu). \quad (6.9.1)$$

*Proof.* From the definition of  $x(y_i)$ , there are points  $y_{i,1} \in [y_-, y_i]$ ,  $y_{i,2} \in [y_i, y_+]$  and such that, for  $j = 1, 2$ , and  $i = 1, 2$ ,

$$d(y_{i,j}, x(y_i)) \leq C(\nu).$$

Hence, for  $j = 1, 2$

$$d(y_{2,j}, y_{1,j}) \geq L(\nu) - 2C(\nu)$$

and so, assuming  $L(\nu)$  sufficiently large given  $C(\nu)$ ,  $y_{2,1}$  cannot be within  $C(\nu)$  of any point on  $[y_-, y_1]$ . So considering the triangle with vertices at  $y_1, y_2, y_-$ , and again applying the special case of the Triangle Theorem of 6.4, there must be  $y'_2 \in [y_1, y_2]$  with

$$d(y_{2,1}, y'_2) \leq C(\nu).$$

This gives the existence of  $y'_2$ , for  $C_1(\nu) = 2C(\nu)$ , for  $C(\nu)$  as in 6.4. The existence of  $y'_1$  is similar, and (6.9.1) follows.  $\square$

## 6.10 “Orthogonal projection”: two set-valued functions.

Now let  $[y_-, y_+]$  be any geodesic segment in  $\mathcal{T}(S)$ . Before defining a function  $x$  with values in  $\mathcal{T}(S)$ , we shall define two set-valued functions  $T(z, +)$  and  $T(z, -)$  with essentially complementary values in  $S \times [y_-, y_+]$ . We shall sometimes use these rather than the “orthogonal projection” itself.

Fix ltd parameter functions  $(\Delta, r, s, K_0)$  for which the results general case of the Triangle Theorem of 6.4 holds. By 5.7, we can choose a vertically efficient partition of  $S \times [y_-, y_+]$  into sets  $\alpha \times \ell$ , where each  $(\alpha, \ell)$  is either ltd or bounded. Let  $\mathcal{P}$  denote the set of the  $(\alpha, \ell)$  from the partition. We have the ordering of 6.3, which is transitive on  $\mathcal{P}$ . Fix any  $z \in \mathcal{T}(S)$ . By 6.4, for each  $(\alpha, \ell) \in \mathcal{P}$ ,  $\ell$  is the disjoint union  $\ell_- \cup \ell_+$ , where one of these two segments could be empty, such that  $\ell_-$  is a bounded  $d_\alpha$ -distance from a corresponding segment on  $[y_-, z]$ , and similarly for  $(\alpha, \ell_+)$ , with the usual modifications if  $\alpha$  is a loop. Write  $T_{\text{ltd}}(z, -) = T_{\text{ltd}}(z, -, [y_-, y_+])$  for the resulting set of  $(\alpha, \ell_-)$  and  $T_{\text{ltd}}(z, +)$  for the set of  $(\alpha, \ell_+)$ . Then let  $T(z, -) = T(z, -, [y_-, y_+])$  be the union of  $T_{\text{ltd}}(z, -)$  and of all bounded  $(\beta, \ell') \in \mathcal{P}$  with  $(\beta, \ell') \leq (\alpha, \ell_-)$  for some  $(\alpha, \ell_-) \in T_{\text{ltd}}(z, -)$  and of all  $(\alpha', \ell') \in \mathcal{P}$  for which there is no  $(\alpha, \ell) \in T_{\text{ltd}}(z, +)$  with  $(\alpha', \ell') > (\alpha, \ell)$ . We define  $T(z, +) = T(z, +, [y_-, y_+])$ , and similarly with  $T_{\text{ltd}}(z, +) \subset T(z, +)$ . Note that some bounded  $(\alpha, \ell)$  are likely to be in both  $T(z, -)$  and

$T(z, +)$ , maximal elements in  $T(z, -)$  and minimal elements in  $T(z, +)$ . The definitions are such that  $S$  is the disjoint union of those  $\alpha$  such that  $(\alpha, \ell)$  is maximal in  $T(z, -)$ , and similarly for  $T(z, +)$  and the  $(\alpha, \ell)$  minimal in  $T_{z,+}$ . We write  $T_{\max}(z, -)$  for the set of maximal elements in  $T(z, -)$  and  $T_{\min}(z, +)$  for the set of minimal elements in  $T(z, +)$ . The  $\alpha$  with  $(\alpha, \ell) \in T_{\max}(z, -)$  are disjoint and their union is  $S$ , and similarly for  $T_{\min}(z, +)$ .

By construction, the sets  $T(z, +)$  and  $T(z, -)$  are coarse Lipschitz in  $z$ , in a natural sense.

### 6.11 “Orthogonal projection”: Upper and Lower Boundary and $x(\cdot, \cdot)$ .

The sets  $T(z, -)$  and  $T(z, +)$  fit into a more general framework of taking a vertically efficient partition  $\mathcal{P}$  of  $S \times [y_-, y_+]$  into ltd and bounded sets  $(\alpha, \ell)$ , taking a maximal unordered set of ltd's in this partition, splitting the elements of this into two sets  $E_{\text{ltd}}(-)$  and  $E_{\text{ltd}}(+)$  and forming resulting sets  $E(-)$  and  $E(+)$  with maximal and minimal sets  $E_{\max}(-)$  and  $E_{\min}(+)$  respectively. We shall say that such  $E(\pm)$  are obtained from an *order splitting*  $E$  of  $\mathcal{P}$ . As in 6.10, the sets  $E(-)$  and  $E(+)$  are probably not disjoint because they can have some bounded  $(\alpha, \ell)$  in common.

The *upper boundary* of  $E(-)$  is the set of all  $(\alpha, y)$  such that  $(\alpha, \ell) \in E_{\max}(-)$  and  $y$  is the right endpoint of  $\ell$ . The lower boundary of  $E(+)$  is defined similarly.

The upper and lower boundary can be used to define a single element  $x(E(+)) = x(E(-))$  of  $\mathcal{T}(S)$  up to bounded Teichmüller distance. If  $E(\pm) = T(z, \pm)$  then we can regard this element of  $\mathcal{T}(S)$  as the orthogonal projection of  $z$  — which we shall do sometimes, but not always, because it makes for worse constants, which seems an unnecessary complication. The  $\alpha$  with  $(\alpha, y)$  in the upper boundary of  $E_{\max}(-)$  are disjoint, with union  $S$ , and similarly for the lower boundary. Then we can define  $x(E(-))$  up to bounded distance by defining the image under projections  $\pi_\alpha$ . So let  $(\alpha, y)$  be in the upper boundary of  $E(-)$ . If  $\alpha$  is a gap, we define

$$\pi_\alpha(x(E(-))) = \pi_\alpha(y).$$

If  $\alpha$  is a loop, we define

$$\operatorname{Re}(\pi_\alpha(x(E(-)))) = \frac{1}{\varepsilon_0}, \quad \operatorname{Im}(\pi_\alpha(x(E(-)))) = \operatorname{Im}(\pi_\alpha(y)).$$

So we are stipulating that loops  $\alpha$  are not short at  $x(E(-))$ , for  $\alpha$  in the decomposition. We define  $x(E(+))$  similarly using the lower boundary of  $E(+)$ . Then  $d(x(E(+)), x(E(-)))$  is bounded in terms of the ltd parameter functions and flat constant. This follows because if  $(\alpha_+, y_+)$  and  $(\alpha_-, y_-)$  are in the lower and upper boundaries of  $E(+)$ ,  $E(-)$  and  $\alpha_+ \cap \alpha_- \neq \emptyset$ , then  $y_- = y_+$  if  $\alpha_- \neq \alpha_+$ , and  $d_\alpha(y_-, y_+)$  is bounded if  $\alpha_- = \alpha_+ = \alpha$ . So  $x(E) = x(E(-)) = x(E(+))$  is well-defined up to bounded distance.

If  $E(\pm) = T(z, \pm, [y_-, y_+])$ , then we shall denote  $x(E)$  by  $x(z, [y_-, y_+])$ , or sometimes by  $x(z)$ , if the context is clear. So we no longer have  $x(z) \in [y_-, y_+]$ , even up to bounded distance, as in the case of  $[y_-, y_+] \subset \mathcal{T}_{\geq \nu}$ . But we do have this for suitable coordinates.

The following lemma is a general analogue of 6.9.

**Lemma 6.12.** *Choose ltd parameter functions  $(\Delta, r, s, K_0)$  such that the results on this section hold, and sufficiently strong given a first set of ltd functions  $(\Delta', r', s', K'_0)$ . There is a function  $C_1(\nu)$  and constant  $C_1$  depending only on  $(\Delta, r, s, K_0)$  such that the following hold. Let  $[y_-, y_+]$  be any geodesic segment, and take sets  $T(y, \pm) = T(y, \pm, [y_-, y_+])$  with respect to  $(\Delta, r, s, K_0)$ .*

*Let  $y_1, y_2 \in \mathcal{T}(\mathcal{S})$ . Let  $\ell = \ell_1 \cap \ell_2$ , where  $(\beta, \ell_1) \in T(y_1, +)$ ,  $(\beta, \ell_2) \in T(y_2, -)$ , with  $\beta$  long  $\nu$ -thick and dominant or  $K_0$ -flat along  $\ell$ . Then (as in 6.9), there is  $\ell' \subset [y_1, y_2]$  such that  $\beta$  is ltd for  $(\Delta', r', s', K'_0)$ , and such that, for all  $w \in \ell$ , there is  $w' \in \ell'$  with*

$$d_\beta(w, w') \leq C_1(\nu) \text{ or } |\operatorname{Re}(\pi_\beta(w) - \pi_\beta(w'))| \leq C_1.$$

*and for all  $w_1, w_2 \in \ell'$*

$$d(y_1, y_2) \geq d'_\beta(y_1, w_1) + d'_\beta(y_2, w_2) + d_\beta(w_1, w_2) - 4C_1 \quad (6.12.1)$$

*where  $C_1 = C_1(\nu)$  if  $\beta$  is a gap and is a fixed constant if  $\beta$  is a loop.*

*Proof.* The argument is similar to 6.9. There are segments corresponding to  $(\beta, \ell)$  on  $[y_1, y_+]$  and  $[y_2, y_-]$ . Since there is a corresponding segment on  $[y_1, y_+]$ , there cannot be one on  $[y_-, y_1]$ . So considering the triangle with vertices  $y_1, y_2, y_-$ , and using the general Triangle Theorem of 6.4, there must be a corresponding segment on  $[y_1, y_2]$  to the one on  $[y_2, y_-]$ , for the chosen parameter functions  $(\Delta', r', s')$  and  $K'_0$  if  $(\Delta, r, s)$  and  $K_0$  are suitably chosen given these. (6.12.1) then follows as (6.9.1).  $\square$

## 6.13

In hyperbolic geometry a  $\kappa$ -quasi-geodesic path is distance  $O(\kappa)$  from a geodesic. No such precise result is available in Teichmüller geometry. But there is a simple result for geodesic segments in  $\mathcal{T}_{\geq \nu}$  for any fixed  $\nu > 0$ , which are proved analogously to corresponding results in hyperbolic space. Variants on the following lemma are possible, and some of those statements might be slightly simpler than the following, but this is precisely the form we shall need. An earlier proof that a quasigeodesic in  $\mathcal{T}(\mathcal{S})$  with endpoints joined by a geodesic in  $\mathcal{T}_{\geq \nu}$  is a bounded distance from that geodesic appears in 4.2 of [35].

**Lemma** *The following holds for any sufficiently large  $L_1$ . Let  $z_\pm \in \mathcal{T}(\mathcal{S})$  with  $[z_-, z_+] \subset \mathcal{T}_{\geq \nu}$  and  $d(z_-, z_+) \geq 4L_1$ . Let  $x(\cdot) = x(\cdot, [z_-, z_+])$  as in 6.9. Let  $\{y_i : 0 \leq i \leq n\}$  be a sequence in  $\mathcal{T}(\mathcal{S})$  with  $d(x(y_0), z_-) \leq L_1$ ,  $d(x(y_n), z_+) \leq L_1$ , and  $d(y_i, y_{i+1}) \leq L_1$  for all  $i$ . Let  $\{z_i : 0 \leq i \leq n\}$  be a sequence of successive points on  $[z_-, z_+]$  with  $z_0 = z_-$ ,  $z_n = z_+$  and  $L_1^{-1} \leq d(z_i, z_{i+1}) \leq L_1$  for all  $0 \leq i < n$ . Suppose also that there is a function  $K(L)$  such that,*

whenever  $d(z_i, y_j) \leq L$  for some  $i$  and  $j$ , then  $d(z_j, y_j) \leq K(L)$ . Then there is  $L_2$  depending only on  $L_1, \nu$  and the function  $K(L)$  such that for all  $i$ ,

$$d(z_i, y_i) \leq L_2.$$

*Proof.* We have  $x(y, [z_-, z_+]) \in [z_-, z_+]$  for all  $y \in \mathcal{T}(\mathcal{S})$  by the definition of  $x$  in 6.9. Fix  $L_2$ , to be taken sufficiently large given  $L_1$  and  $\nu$ . Take any  $i_1 < i_2$  such that, for  $i_1 < i < i_2$ ,

$$d(y_j, z_i) \geq L_2 \text{ for all } j$$

and either  $i_1 = 0$  or  $d(y_j, z_{i_1}) \leq L_2$  for some  $j$ , and either  $i_2 = n$  or  $d(y_j, z_{i_2}) \leq L_2$  for some  $j$ . To prove the lemma, it suffices to bound  $i_2 - i_1$  in terms of  $L_1$ , if  $L_2$  is suitably defined in terms of  $L_1$ , and to obtain a contradiction for  $L_2$  sufficiently large in terms of  $L_1$ , if  $i_2 - i_1 = n$  and  $d(z_0, y_0) \geq L_2$ .

Since  $i_2 - i_1 \leq L_1 d(z_{i_1}, z_{i_2})$ , we may suppose that

$$d(z_{i_1}, z_{i_2}) \geq 4K(L_2).$$

Then since  $x(\cdot)$  is coarse Lipschitz, we can choose a sequence  $i(j)$ ,  $0 \leq j \leq r$ , such that the points  $x(y_{i(j)})$  occur in strictly increasing order in  $[x(y_{i_1}), x(y_{i_2})] \subset [z_-, z_+]$ , with  $i(0) = i_1$ ,  $i(r) = i_2$ , and

$$2L_1 \leq d(x(y_{i(j)}), x(y_{i(j+1)})) \leq 4L_1, \quad (6.13.1)$$

$$d(z_{i_1}, x(y_{i_1})) \leq K(L_2), \quad d(z_{i_2}, x(y_{i_2})) \leq K(L_2).$$

Then

$$d(x(y_{i_1}), x(y_{i_2})) \geq \frac{1}{2}d(z_{i_1}, z_{i_2}).$$

It follows from (6.13.1) that

$$r \geq \frac{i_2 - i_1}{8L_1^2}.$$

Assuming  $L_1$  is sufficiently large given  $C_1 = C_1(\nu)$  of 6.9, by (6.9.1),

$$d(y_{i(j)}, y_{i(j+1)}) \geq$$

$$\begin{aligned} & d(y_j, x(y_{i(j)})) + d(x(y_{i(j)}), x(y_{i(j+1)})) + d(x(y_{i(j+1)}), y_{i(j+1)}) - 4C_1 \\ & \geq d(y_j, x(y_{i(j)})) + d(x(y_{i(j+1)}), y_{i(j+1)}). \end{aligned}$$

For  $1 \leq j < r$ ,  $d(y_j, x(y_{i(j)})) \geq L_2$ . So

$$L_1(i_2 - i_1) \geq \sum_{j=0}^{r-1} d(y_{i(j)}, y_{i(j+1)})$$

$$\geq d(y_{i_1}, x(y_{i_1})) + \frac{1}{2}(r-1)L_2 \geq \frac{L_2}{16L_1^2}(i_2 - i_1) - \frac{1}{2}L_2$$

Now put  $L_2 = 32L_1^3$ . Then we have

$$(i_2 - i_1) \leq 16L_1^2.$$

So in all cases, we have a bound on  $i_2 - i_1$  in terms of  $L_1$ , if  $L_2 = 32L_1^3$ . Also,

$$d(y_{i_1}, x(y_{i_1})) \leq L_1(i_2 - i_1).$$

So then,

$$d(y_{i_1}, z_{i_1}) \leq 16L_1^3 + K(L_2)$$

and for  $i_1 \leq i \leq i_2$ ,

$$d(y_i, z_i) \leq 48L_1^3 + K(L_2).$$

Since the righthand side is  $> L_2$ , we have this for all  $i$ .

□

The generalisation to  $\mathcal{T}(S)$  that we shall use is somewhat weaker than this — necessarily so.

**Lemma** *The following holds for a constant  $L_0$ ,  $\Delta_0$  given fixed ltd parameter functions  $(\Delta, r, s, K_0)$ , related constant  $\nu_0$  as in 5.5, and given constants  $\kappa_0$ ,  $L_1$ ,  $\lambda_0 > 0$ . Let  $d(y_i, y_{i+1}) \leq L_1$  for all  $i$  and*

$$\sum_{i=0}^{n-1} d(y_i, y_{i+1}) \leq \kappa_0 d(y_0, y_n).$$

*Suppose also that  $\{(\alpha_i, \ell_i) : 1 \leq i \leq m \text{ is a totally ordered set of ltds along } S \times [y_0, y_n], \text{ such that if } \alpha_i \text{ is a gap, then } \alpha_i \text{ is } \nu\text{-thick, long and dominant along } \ell_i, \text{ for some } \nu \geq \nu_0, \text{ and, if } \alpha_i \text{ is a loop, } \alpha_i \text{ is } K_0\text{-flat along } \ell_i. \text{ Then if } [x_1, x_2] \subset [y_0, y_n] \text{ is such that}$*

$$\sum_{j=1}^n |\ell_j \cap [x_1, x_2]| \geq \lambda_0 d(y_0, y_n),$$

*there are at least one  $i, j, y = y_i$  and  $w \in \ell_j \cap [x_1, x_2]$  such that*

$$d'_{\alpha_j}(y, w) \leq L_0.$$

*Proof.* Suppose the lemma is not true for  $[x_1, x_2]$ . Let  $C_1$  be as in 6.12. Removing some of the  $y_i$ , and assuming  $L_1$  is large enough that  $L_1/2 \geq \Delta(\nu)$  for all  $\nu \geq \nu_0$ , we can assume that for each  $0 \leq i < n$ , the sets  $(\alpha_j, \ell_{j,i})$  in  $T(y_i, +) \cap T(y_{i+1}, -)$  for  $\ell_{j,i} \subset \ell_j$  are for  $j \in I(i)$  where

$$L_1 \leq \sum_{j \in I(i)} |\ell_{j,i}| \leq 2L_1.$$

Of course, we no longer have the upper bound on  $d(y_i, y_{i+1})$ , but we still have the same upper bound on  $\sum_i d(y_i, y_{i+1})$ . By our assumption that the lemma is not true, for all  $w \in \ell_j$ , all  $j \in I(i)$ ,

$$d'_{\alpha_j}(w, y_i) \geq L_0, \quad d'_{\alpha_j}(w, y_{i+1}) \geq L_0.$$

Let  $A$  be the set of  $i$  such that  $\ell_{j,i} \subset [x_1, x_2]$  for all  $j \in I(i)$ . Then by 6.12 we have, for all  $w \in \ell_{j,i}$ , and  $i \in A$ ,  $j \in I(i)$ ,

$$d(y_i, y_{i+1}) \geq d'_{\alpha_j}(w, y_i) + d'_{\alpha_j}(w, y_{i+1}) - 2C_1 \geq 2L_0 - 2C_1.$$

Now we also have

$$\#(A) \geq \frac{\lambda_0}{2L_1} d(y_0, y_n).$$

So then we get, assuming  $L_0 \geq 2C_1$ , as we may do,

$$\sum_{i \in A} d(y_i, y_{i+1}) \geq \frac{\lambda_0 L_0}{2L_1} d(y_0, y_n).$$

This gives a contradiction if  $L_0$  is large enough that

$$\frac{\lambda_0 L_0}{2L_1} > \kappa_1.$$

□

Note that we do not get such a strong result as in the case when  $[y_0, y_n] \subset \mathcal{T}_{\geq \nu}$  for some  $\nu$  — and actually, our assumption in 6.12 was a little weaker than this. The reason is that in the case  $[y_0, y_n] \subset \mathcal{T}_{\geq \nu}$  we can deduce a bound on  $d(y_i, y_j) - d(x(y_i), x(y_j))$  from bounds on  $d(y_i, x(y_i))$  and  $d(y_j, x(y_j))$ . But what we have above is (effectively) a bound on  $d_{\alpha}(y_i, x(y_i))$  for some  $i$  and for some subsurface  $\alpha$ . But we used the bound on  $d(y_0, y_n)$  to get this, and would need to bound  $d(y_i, y_j)$  to proceed further. So there is no real analogue of the quasi-geodesic-implies-geodesic result for Teichmüller geodesics in general. Instead, it seems to be possible to obtain results for families of paths following certain rules. The paths through pleated surfaces that will be used do follow such a set of rules, as we shall see.

## 6.14 A Chain of ltd's.

We shall use the following generalization of 5.4. The notation  $(\alpha_j, \ell_j)$  is used, the same notation as in 5.7. In subsequent sections the  $\{(\alpha_j, \ell_j) : 1 \leq j \leq R_1\}$  produced below will be a subset of the set in 5.7.

The proof of the result is different in character from that of 5.4, being, essentially, a construction of a zero measure Cantor set, while 5.4 obtained a set with a lower bound on area. This result is in any case more sophisticated, because it uses 5.5 — and hence also 5.4 — in the course of the proof. This result can be regarded as a parallel to the existence of a tight geodesic in the curve complex used by Minsky et al.. For reasons which are not entirely clear

to me, but which may be significant, this result appears to be much harder to prove.

**Theorem** *Fix long thick and dominant parameter functions and flat constant  $(\Delta, r, s, K_0)$ . Then there exist  $\Delta_0$  and  $\nu_0$  depending only on  $(\Delta, r, s, K_0)$  and the topological type of  $S$  such that the following holds. Let  $[y_0, y_T] = [[\varphi_0], [\varphi_T]]$  be any geodesic segment in  $\mathcal{T}(S)$  of length  $T \geq \Delta_0$ , parametrised by length. Then there exists a sequence  $(\alpha_i, \ell_i)$  ( $1 \leq i \leq R_1$ ) of ltd's with  $(\alpha_i, \ell_i) < (\alpha_{i+1}, \ell_{i+1})$  for  $i < R_0$ , where the ordering  $<$  is as in 6.3 and such that each segment of  $[y_0, y_T]$  of length  $\Delta_0$  intersects some  $\ell_i$ .*

*Start of Proof.*

We shall prove the theorem by showing that, if  $\Delta_0$  is sufficiently large, for some loop  $\gamma \subset S$  and some  $\xi \in \gamma$ , for every segment  $\ell$  of length  $\Delta_0$ , along  $[y_0, y_T]$ , there is an ltd  $(\alpha, \ell')$  with  $\ell' \subset \ell$  and  $\varphi_0(\xi) \in \varphi_0(\alpha)$ . It then follows that the intersection of all such  $\varphi_0(\alpha)$  is nonempty and therefore any two such  $\alpha$  intersect essentially. It is not quite the case that any sequence  $(\alpha_i, \ell_i)$  for  $i \leq j$  is extendable, but, essentially, if  $(\alpha_j, \ell_j)$  is chosen so that  $\alpha_j$  does not intersect any bounded gaps (in the sense of 5.5) for distance  $\Delta_0^{1/n}$  beyond  $\ell_j$ , for a suitable  $n$  depending only on the topological type of  $S$ , then further extension becomes possible.

Let  $y_t = [\chi_t \circ \varphi_0]$  be such that  $d(y_0, y_t) = t$  with stretch  $q_t(z)dz^2$  at  $y_t$  and  $d(y_-, y_+) = t$  and with  $\chi_t$  minimising distortion. We use this parameterisation for the whole geodesic containing  $[y_0, y_T]$ . We shall write  $|\cdot|_t$  for  $|\cdot|_{q_t}$  and  $|\cdot|_{t,+}$  for the unstable length  $|\cdot|_{q_{t,+}}$ . We fix a vertically efficient decomposition of  $S \times [y_0, y_T]$  as in 5.6, in which every  $(\beta, \ell)$  is either ltd or bounded by  $L$  in the sense of 5.5, with  $L$  as in 5.5. In particular, for  $[\varphi] \in \ell$ :

$$|\varphi(\partial\beta)| \leq L. \quad (6.14.1)$$

In fact, (6.14.1) holds for all  $[\varphi] \in \ell'(\partial\beta)$ , where  $\ell'(\partial\beta) = \{[\chi_t \circ \varphi_0] : t_0(\partial\beta) \leq t \leq t'_0(\partial\beta)\}$  is the possibly larger interval containing  $\ell$  such that  $\partial\beta$  does not intersect  $\alpha$  transversally, for any  $\alpha$  which is ltd along a segment of  $\ell'$ . Here,  $t_0(\partial\beta) < 0$  and  $t'_0(\partial\beta) > T$  are allowed. Also, for  $y_t = [\chi_t \circ \varphi_0] \in \ell$ , by the last part of 5.4 we have, if  $(\beta, \ell)$  is bounded by  $L$ ,

$$|\chi_t \circ \varphi_0(\partial\beta)|_t^2 \geq s_0 a'(\beta) \quad (6.14.2)$$

for a constant  $s_0$  depending only on the ltd parameter functions and the topological type of  $S$ . Here,  $a'(\beta)$  is as in 5.2.

Now for any loop or finite set of (not necessarily disjoint) loops  $\gamma$ , let  $F(t, \gamma)$  be the qd-length function as in 6.7, with minimum  $t(\gamma)$ . Now suppose that  $(\beta, \ell)$  is bounded by  $L$  and that  $\ell$  is of length  $\geq \Delta_0$ . Then from (6.14.1), (6.14.2) and (6.7.1) we obtain, for a constant  $C_1$  depending only on the topological type of  $S$ ,

$$a'(\beta) \leq C_1 L^2 e^{-\Delta_0}. \quad (6.14.3)$$

Also, by definition,

$$t_0(\partial\beta) \leq t(\partial\beta) \leq t'_0(\partial\beta).$$



**Lemma 6.15.** *The following holds for a suitable constant  $L_1$  depending only on the topological type of  $S$ , if  $\Delta_0$  is sufficiently large given a constant  $L$ . Let  $\beta$  be bounded by  $L$  along  $\ell \subset [y_0, y_T]$ , where  $\ell$  has length  $\geq \Delta_0$ . Then there is  $(\beta_1, \ell_1)$  bounded by  $L_1$ , such that  $\beta \subset \beta_1$ ,  $\beta$  is properly contained in  $\beta_1$  if  $t(\partial\beta) \leq t_0(\partial\beta) + \sqrt{\Delta_0}$  and  $\ell_1 \subset \ell$  has length at least  $\Delta_0^{1/3}$ .*

By (6.14.3) with  $\Delta_0$  replaced by  $\Delta_0^{1/3}$ , the union of all such  $\beta_1$  must also be properly contained in  $S$ , assuming  $\Delta_0$  is sufficiently large given the ltd parameter functions. Then applying the lemma a number of times which is bounded in terms of the topological type of  $S$ , with  $\Delta_0$  replaced by  $\Delta_0^{1/n}$  for different  $n$ , we obtain the following.

**Corollary 6.16.** *The following holds for a suitable constant  $L_1$  as above, if  $\Delta_0$  is sufficiently large given  $L$ , and for  $k$  which is bounded in terms of the topological type of  $S$ . Let  $\ell \subset [y_0, y_T]$  be any interval of length  $\geq \Delta_0$ . Then there is  $\ell' \subset \ell$  of length  $\Delta_1 \geq \Delta_0^{1/k}$  such that, for any  $\ell'' \subset \ell'$  of length  $\geq \Delta_1^{1/3}$ , the union of  $\beta$  which are bounded by  $L$  along  $\ell'$  is the same as the union of  $\beta$  which are bounded by  $L_1$  along  $\ell''$ , and for all such  $\beta$ ,  $t(\partial\beta) \geq t_0(\partial\beta) + \Delta_1^{1/2}$ .*

## 6.17 Proof of 6.15.

The lemma is proved by using the characterisation of bounded Poincaré Poincaré length of 6.7, in particular, the discussion on comparing length of non-transverse loops. Fix  $L_1$ , depending only on the topological type of  $S$ , so that, for any  $[\varphi] \in \mathcal{T}(S)$ , a maximal multicurve  $\Gamma$  with  $\varphi(\Gamma) \leq L_1$  exists. Also fix a Margulis constant  $\varepsilon_0$ . We consider the function

$$G(t, \partial\beta) = \text{Max}(F(t, \partial\beta) - F(t, \Gamma)) = F(t, \partial\beta) - \text{Min}F(t, \Gamma),$$

where, the minimum is taken over loop sets  $\Gamma$  such that  $\gamma$  satisfies 6.7.3 for all  $\gamma \in \Gamma$ , and with specified convex hull for  $\Gamma$ . The convex hull is a finite union of gaps and loops  $\alpha$  such that, for each  $\alpha$ ,  $\chi_t \circ \varphi_0(\alpha)$  is homotopic to a component of  $(\chi_t \circ \varphi_0(S))_{\geq \varepsilon_0}$ . The closure of the union of the  $\alpha$  contains  $\beta$ , and every such  $\alpha$  either intersects  $\beta$ , or shares a boundary component with  $\beta$ . The possibility that  $\alpha$  is disjoint from  $\beta$  only happens if  $t(\partial\beta) < t_0(\partial\beta) + \sqrt{\Delta_0}$ . Any boundary component  $\gamma_1$  which is shared by  $\beta$  and  $\alpha$  is such that  $F(\gamma_1, t)$  is maximal among  $\gamma_1 \subset \partial\beta$ . Assuming  $L_2$  is large enough given the topological type of  $S$ , for each  $t$ , there is at least one  $\Gamma$  satisfying these conditions, but only boundedly many can give the minimum value of  $G(t, \partial\beta)$ , with bound depending on  $L_2$ . We write  $\Gamma_t$  for a choice of  $\Gamma$  such that

$$|G(\partial\beta, t) - (F(\partial\beta, t) - F(\Gamma_t, t))| \leq C(\varepsilon_0),$$

for a suitable constant  $C(\varepsilon_0)$  depending only on  $\varepsilon_0$  and the topological type of  $S$ .

For  $t$  such that  $|\chi_t \circ f_0(\partial\beta)| \leq L$ , that is,  $t_0(\partial\beta) \leq t \leq t'_0(\partial\beta)$ ,  $G(t, \partial\beta)$  is bounded above in terms of  $L$ . For  $t$  such that  $|\chi_t \circ f_0(\partial\beta)| \geq \varepsilon$ ,  $G(t, \partial\beta)$

is bounded below in terms of  $\varepsilon$ . So by our choice of  $t_0(\beta)$ ,  $t'_0(\beta)$ ,  $G(t, \partial\beta)$  is bounded above and below in terms of  $L$  and  $\varepsilon_0$  (our fixed Margulis constant). By the restrictions we have put on the choice of  $\Gamma$ ,  $G(t, \partial\beta)$  is also bounded below in terms of  $L$  for all  $t \in [t_0(\partial\beta), t'_0(\partial\beta)]$ , if  $t(\partial\beta) \geq t_0(\partial\beta) + \sqrt{\Delta_0}$ .

So in all cases, by the choice of  $\Gamma_t$  for  $t \in [t_0(\partial\beta), t'_0(\partial\beta)]$  there is a bound on the number of different  $\Gamma_t$  in terms of  $L$  and  $\varepsilon_0$  if  $t(\partial\beta) \geq t_0(\partial\beta) + \sqrt{\Delta_0}$ , and if  $t(\partial\beta) \leq t_0(\partial\beta) + \sqrt{\Delta_0}$  then the number of different sets of loops  $\Gamma_t$  for  $t \in [t_0(\partial\beta), t'_0(\partial\beta)]$  is  $\leq C_1(L, \varepsilon_0)\sqrt{\Delta_0}$ . We see this as follows. The graph of  $F(t, \partial\beta)$  is an approximate V shape, with slope +1 and -1 to the right and left of the minimum. The graph of each  $F(t, \Gamma)$  is also such a V shape, and the graph for  $\Gamma = \Gamma_s$ ,  $s \in [t_0(\partial\beta), t'_0(\partial\beta)]$ , must be above that of  $t \mapsto F(t, \partial\beta) - C_2(L, \varepsilon_0)$  for  $t \in [t_0(\partial\beta), t'_0(\partial\beta)]$ . At  $t = t_0(\partial\beta)$ ,  $t = t'_0(\beta)$ ,  $G(t, \partial\beta)$  is bounded above and below. Then minima of the functions  $F(t, \Gamma)$  move progressively to the right, and the right branches of the V's move down, but all of them within a rectangle with sides slopes -1 and +1, where the -1 sides have width  $\leq C_3(L, \varepsilon_0)$  if  $t(\partial\beta) \geq t_0(\partial\beta) + \sqrt{\Delta_0}$ , and width  $\sqrt{\Delta_0} + C_3(L, \varepsilon_0)$  if  $t(\partial\beta) \leq t_0(\partial\beta) + \sqrt{\Delta_0}$ . For each  $t$ , there are only a bounded number of choices for  $\Gamma_t$ . The minima cannot get too close, because then there would be too many choices for  $\Gamma_t$ , for a slightly larger  $L_2$ . So the number of different loop sets  $\Gamma_t$  which arise must be bounded by  $C_1(L, \varepsilon_0)$  or  $C_1(L, \varepsilon_0)\sqrt{\Delta_0}$  in the respective cases. So there must be  $\ell_1 \subset \ell$  of length  $> \Delta_0^{1/3}$  along which the choice of  $\Gamma_t$  can be chosen to be constant. Let  $\beta_1$  be the union of the gaps in the convex hull of  $\Gamma_t$ . Then  $\beta_1$  is bounded by  $L_1$  along  $\ell_1$ , where  $L_1$  is independent of  $L$ , and  $\beta_1$  strictly contains  $\beta$  if  $t(\partial\beta) \leq t_0(\partial\beta) + \sqrt{\Delta_0}$ , as required.  $\square$

## 6.18 More on transfer between Poincaré and qd-length.

So now, we consider  $(\beta, \ell)$  bounded by  $L_1$ , with  $\ell$  of length  $\geq \Delta_0^{1/k}$  and

$$t(\beta) > t_0(\beta) + \Delta_0^{1/2k}.$$

For  $t < t(\beta)$ , the  $q_t$ -length of unstable segments across  $\chi_t \circ f_0(\beta)$  is  $\leq C_1 L_1 e^{t-t(\beta)}$ , for  $C_1$  depending only on the topological type of  $S$ . We now need to interpret and strengthen this statement in terms of Poincaré length, under the assumption that  $\ell$  has length  $\geq \Delta_0^{1/k}$  and that  $\Delta_0$  is sufficiently large given the ltd parameter functions and flat constant, and hence also sufficiently large given  $s_0$  and  $L_1$ . If  $-\frac{1}{2}\Delta_0^{1/2k} < t - t(\beta) < -\frac{1}{4}\Delta_0^{1/2k}$ , then we can be sure that the Poincaré length of unstable segments is  $\leq C_2 e^{t-t(\beta)}$ , because the  $q_{t,+}$ -length is exponentially smaller than the  $q_{t,-}$ -length, and the Poincaré length of  $\chi_t \circ f_0(\partial\beta)$  is bounded. We also have the following.

**Lemma** *For  $p$  depending only on  $L_1$ , whenever  $t - t(\beta) < -\frac{1}{4}\Delta_0^{1/2k}$ , the following holds. Let  $\gamma \subset \chi_t \circ \varphi_0(S)$  be a segment of unstable foliation whose diameter is boundedly proportional to the injectivity radius at any point of  $\gamma$ . Then all components of  $\gamma \cap \chi_t \circ f_0(\beta)$  have Poincaré length  $\leq -\exp(\Delta_0^{1/3k})$  times the injectivity radius at  $\gamma$  and at least one in any  $p$  complementary components in  $\gamma$*

has Poincaré length at least  $\exp(\Delta_0^{1/3k})$  times the length of any unstable segment in  $\gamma \cap \chi_t \circ f_0(\beta)$ .

*Proof.* Because we are restricting to a set whose diameter is boundedly proportional to the injectivity radius we only need to prove this for  $-\frac{1}{2}\Delta_0^{1/2k} < t - t(\beta) < -\frac{1}{4}\Delta_0^{1/2k}$ , if we replace  $\exp(\Delta_0^{1/3k})$  by  $\exp(\Delta_0^{3/7k})$ . For if we can do this, it is true for this range of  $t$  with respect to  $q_t$ -length, with  $\exp(\Delta_0^{3/7k})$  replaced by  $\exp(\Delta_0^{2/5k})$ . Then since unstable length is multiplied by  $e^t$  under application of  $\chi_t$  for  $t < 0$ , and we are restricting to a set whose diameter is boundedly proportional to the injectivity radius, we have the result for all  $t < -\frac{1}{4}\Delta_0^{1/2k}$ , with respect to  $q_t$ -length. But then we also have the result for Poincaré length, if we replace  $\exp(\Delta_0^{2/5k})$  by  $\exp(\Delta_0^{1/3k})$ , because the image under  $\chi_{-t}$  of an unstable segment of short Poincaré length remains inside the injectivity radius of the surface, and so Poincaré length and  $q_t$ -length are still comparable. So now we suppose that  $-\frac{1}{2}\Delta_0^{1/2k} < t - t(\beta) < -\frac{1}{4}\Delta_0^{1/2k}$ . Suppose that  $p > e^{3L'}$ . Then if we take a string of  $p$  successive segments, two of them must intersect  $\chi_t \circ \varphi_0(\partial\beta)$  at Poincaré distance  $< e^{-2L_1}$  apart. The segments of intersection with  $\chi_t \circ \varphi_0(\partial\beta)$  have Poincaré length  $< \exp(-\frac{1}{5}\Delta_0^{1/2k})$ . Then if all the components outside  $\chi_t \circ \varphi_0(\beta)$  have length  $< e^{-4L_1}$  we can make a loop from such segments of Poincaré length  $< e^{-L_1}$  intersecting  $\chi_t \circ \varphi_0(\partial\beta)$  transversally, assuming that  $\Delta_0$  is sufficiently large. This is impossible.  $\square$

### 6.19 Proof of 6.14: construction of the sequences.

For  $1 \leq i \leq R_0$ , some  $R_0$ , we shall find sequences  $t_i$ ,  $\alpha_i$ ,  $\zeta_i$  such that the following hold, for a constant  $C_1$ .

1.  $\alpha_i$  is ltd at  $y_{t_i}$  for  $[y_0, y_T]$ .
2.  $\zeta_{i+1} \subset \zeta_i \subset f_0(\alpha_i)$ .
3. For all  $i \geq 1$ ,  $\chi_{t_i}(\zeta_i)$  is an unstable segment whose Poincaré length is boundedly proportional, with bounds given by  $C_1$ , to the injectivity radius at that point of  $\chi_{t_i} \circ \varphi_0(S)$ . If  $\alpha_i$  is a loop,  $\chi_{t_i}(\zeta_i)$  is an unstable segment in a component of  $(\chi_{t_i} \circ \varphi_0(S))_{<\varepsilon_0}$  homotopic to  $\chi_{t_i} \circ \varphi_0(\alpha_i)$ , in the part which is flat with respect to the  $q_{t_i}$ -metric.
4.  $t_1 \leq \Delta_0$  and  $t_{R_0} \geq T - \Delta_0$ .
5. For all  $i < R_0$ ,  $t_i + \Delta_0^{1/3k} < t_{i+1} \leq t_i + \Delta_0$ .
6. For all  $i$ ,  $\chi_{t_i}(\zeta_i) \cap \chi_{t_i} \circ \varphi_0(\beta) = \emptyset$  for any  $(\beta, \ell)$  such that  $(\beta, \ell)$  is bounded by  $L_1$ , and  $\ell$  is of length  $\geq \Delta_0^{1/3k}$  with  $\ell$  starting ending within  $\frac{1}{2}\Delta_0^{1/k}$  of the right of  $y_{t_i}$ .

This completes the proof of 6.14, since we then have

$$\zeta_j \cap \varphi_0(\alpha_i) \neq \emptyset$$

for all  $i \leq j$ .

To choose  $(\alpha_1, \ell_1)$ , let  $\ell$  be the initial segment of  $[y_0, y_T]$  of length  $\Delta_0$ . Apply 6.16 to find  $\ell'$ ,  $\Delta_1 = \Delta_0^{1/n}$  for some  $n$  and take any  $\ell'' \subset \ell'$  of length  $\Delta_1^{1/3}$  in the first half of  $\ell'$ . Assuming that the ltd parameter functions are sufficiently strong given  $L_1$ , that a component of a  $\partial\beta$  cannot be contained in the interior of an ltd gap  $\alpha$ , and cannot be an ltd loop. Then by 5.5 we can find an ltd  $(\alpha_1, \ell_1)$  with  $\ell_1 \subset \ell''$  and with  $\alpha$  of sufficiently large qd-area not to be contained in any  $\beta$ , and hence disjoint from all the  $\beta$  which are bounded along  $\ell''$ . We can then choose  $\zeta_1 \subset f_0(\alpha_1)$  so that  $\chi_{t_1}(\zeta_1)$  has Poincaré length boundedly proportional to the injectivity radius at any point of  $\chi_{t_1}(\zeta_1)$ , in the flat-metric part of  $\chi_{t_1} \circ f_0(\alpha_1)$  if  $\alpha_1$  is a loop.

So now suppose that we have found  $\zeta_i$ ,  $t_i$  and  $\alpha_i$  and we look for  $\zeta_{i+1}$  by looking for an appropriate subset of  $\chi_{t_i}(\zeta_i)$ . Then we claim that the set of  $I(\beta)$  with  $(\beta, \ell)$  bounded by  $L_1$ ,  $\ell \subset [y_{t_i}, y_T]$  of length  $\geq \Delta_0^{1/k}$  is not all of  $\chi_{t_i}(\zeta_i)$ , if  $\Delta_0$  is sufficiently large in terms of the ltd parameter functions. We only need to consider  $\beta$  with  $t_0(\partial\beta) \geq t_i + \frac{1}{2}\Delta_0^{1/k}$ , by condition 6. Then, as in the case  $i = 1$ , by 6.16, for  $\Delta_1 = \Delta_0^{1/2k^2}$ , we only need to prove this for  $\ell$  of length  $\geq \Delta_1$  and

$$t_i + \Delta_1 \leq t_0(\partial\beta) \leq \Delta_0, \quad (6.19.1)$$

$$t(\partial\beta) \geq t_0(\partial\beta) + \sqrt{\Delta_1}. \quad (6.19.2)$$

First, the number of such  $\beta$  is  $\leq C\Delta_0$  for  $C$  depending only on the topological type of  $S$ . Define

$$I(\beta) = \chi_{t_i}(\zeta_i) \cap \chi_{t_i} \circ \varphi_0(\beta).$$

Then, restricting to this set of  $\beta$  satisfying (6.19.1) and (6.19.1), by 6.18, the sum of the Poincaré lengths of the intervals of the set  $I(\beta)$  is  $\leq C_1\Delta_0pe^{-\Delta_1}$  times the injectivity radius at any point of  $\chi_{t_i}(\zeta_i)$ . Since  $\chi_{t_i}(\zeta_i)$  has Poincaré length  $\geq C_1^{-1}$  times the injectivity radius, the complement of the union of the  $I(\beta)$  is nonempty. Note that this argument would not work without the upper bound in (6.19.1). The calculation just done is the first step in constructing a zero measure Cantor set, albeit of Hausdorff dimension close to 1 if  $\Delta_0$  is large.

Now choose an interval in  $[y_{t_i}, y_{t_i+\Delta_0}]$  as  $\ell''$  in 6.16, with  $\ell''$  in the first half of  $\ell'$ , again for  $\ell'$  as in 6.16. The complement in  $\zeta_i$  of the  $I(\beta)$  for the finitely many  $\beta$  for which  $\beta$  is bounded by  $L_1$  along  $\ell'$  is nonempty, by 6.17, and the complement of this set of  $\beta$  has qd-area bounded from 0 and has nonempty intersection with convex hull of the  $\alpha$  which are ltd along segments of  $\ell''$ , by 5.5, assuming that  $\Delta_0$  is sufficiently large. Assuming that the ltd parameter functions are sufficiently strong given  $L_1$ , that a component of a  $\partial\beta$  cannot be contained in the interior of an ltd gap  $\alpha$ , and cannot be an ltd loop. So the complement of the union of the  $\beta$  is a union of components of the convex hull of the ltd's. Each component  $J$  of the complement in  $\zeta_i$  of these  $I(\beta)$  must intersect  $\chi_{t_i} \circ \varphi_0(\alpha)$  for some ltd  $(\alpha, \ell)$ , because otherwise we can make a nontrivial loop out of the boundaries of the  $\chi_{t_i} \circ \varphi_0(\partial\beta_i)$  and  $J$  which is disjoint from all the  $\chi_{t_i} \circ \varphi_0(\alpha)$ , and homotoping that loop to good position, it remains disjoint from

$J$ , and separates the convex hull from  $J$ . Take any such  $(\alpha, \ell)$  to be  $(\alpha_{i+1}, \ell_{i+1})$ , and  $\zeta_{i+1}$  so that  $\chi_{t_i}(\zeta_{i+1})$  is a component of  $J \cap \chi_{t_i} \circ \varphi_0(\alpha_{i+1})$ . Then  $\chi_{t_{i+1}}(\zeta_{i+1})$  has Poincaré length bounded from 0, even if we restrict to the intersection with  $(\chi_{t_{i+1}} \circ \varphi_0(S))_{\geq \varepsilon_0}$ . Then by the choice of  $\ell''$ ,  $(\alpha_{i+1}, \ell_{i+1})$  and  $\zeta_{i+1}$  have all the required properties.

□

## 7 Geometric model manifolds.

In this section, we construct the geometric model  $M = M(\mu(e_1), \dots, \mu(e_n))$  for any homeomorphism type of relative Scott core with ends  $e_i$ . We start by constructing geometric models in the geometrically finite cases, when  $\mu(e_i) \in \mathcal{T}(S(e_i))$  for all  $i$ . The simplest case of combinatorial bounded geometry is dealt with first, in 7.1. The other geometrically finite cases follow. All models are geometrically finite until 7.16. The geometrically infinite case, when  $\mu(e_i) \in \mathcal{O}_a(S(e_i))$ , for some, or all,  $i$ , is constructed by taking geometric limits, for which, of course, we need to know that if  $\mu^k(e_i) \rightarrow \mu(e_i)$  then the corresponding geometric models  $M(\mu^k(e_1), \dots, \mu^k(e_n))$  converge geometrically, up to Lipschitz equivalence, to a single limit. This is dealt with in 7.16 in the case of combinatorial bounded geometry, and in the general case in 7.17. The geometrical model in the geometrically infinite case is a manifold which is also a metric space, defined up to a bounded Lipschitz equivalence, and boundedly Lipschitz equivalent to a Riemannian manifold whose ends are topological products.

### 7.1 The combinatorially bounded geometry geometrically finite Kleinian surface case.

We assume that the relative Scott core is homeomorphic to  $S_d \times [0, 1]$ , where  $S_d$  is the horodisc deletion of a finite type surface  $S$ , and that the end invariants  $y_0, y_u \in \mathcal{T}(S)$  are such that

$$[y_0, y_u] = \{y_t : t \in [0, u]\} \subset \mathcal{T}_{\geq \nu}$$

for some fixed  $\nu > 0$ . We are parametrising by length, so that  $d(y_s, y_t) = t - s$  for any  $s < t$ . We can, of course, extend the parametrisation, by  $\mathbb{R}$ , to the geodesic in  $\mathcal{T}(S)$  containing  $[y_0, y_u]$ . We let  $S_t$  be the hyperbolic surface for  $y_t$ , with hyperbolic metric  $\sigma_t$  on  $S_t$ . We fix a smooth manifold structure on  $S$ . We take any continuous family of  $C^1$  diffeomorphisms  $\varphi_t : S \rightarrow S_t$  such that  $[\varphi_t] = y_t$  and such that if  $|t - s| < 1$ ,

$$\varphi_t \circ \varphi_s^{-1} : S_s \rightarrow S_t$$

has norm of derivative bounded by  $K$  for some fixed  $K$ . Here, the norm of the derivative is taken with respect to the metrics  $\sigma_t$  and  $\sigma_s$ . We can also choose the family  $\varphi_t$  so that

$$(z, t) \mapsto \varphi_s^{-1} \circ \varphi_t(z) : S \times \mathbb{R} \rightarrow S$$

is  $C^1$  (or even  $C^\infty$ ). Then the model metric  $\sigma_M$  on  $M = S \times [0, u]$  is

$$\sigma_M = \varphi_t^* \sigma_t + dt^2.$$

Fix  $s$  and consider the map

$$(z, t) \mapsto (\varphi_s(z), t) : S \times [0, u] \rightarrow S_s \times [0, u].$$

Then the model metric is transformed to the metric

$$(\varphi_s \circ \varphi_t^{-1})_* \sigma_t + dt^2,$$

which, on  $S_s \times [s-1, s+1]$ , is boundedly equivalent to  $\sigma_s + dt^2$ , because of the bound on the derivative of  $\varphi_s \circ \varphi_t^{-1}$ . It follows that the exact choice of the family  $\varphi_t$  is unimportant. Subject to the above constraints, any choice gives the same geometric model up to Lipschitz equivalence.

In this special case, it may be preferred to use the singular Euclidean metric on  $S_t$  coming from the quadratic differential  $q_t(z)dz^2$  on  $S_t$  for the geodesic and to use  $\varphi_t = \chi_t \circ \varphi_0$  for some  $\varphi_0$  and for  $\chi_t : S_0 \rightarrow S_t$  minimising distortion. In this case, we have a singular Euclidean structure on  $S$  such that the metric on  $S \times [0, u]$  is

$$e^{2t} dx^2 + e^{-2t} dy^2 + dt^2.$$

Although singular, the metric induced by this Riemannian metric is boundedly Lipschitz equivalent to the previous one. See also [37].

## 7.2 Geometrically finite Kleinian surface case: $S_{j,t}$ and properties of $\varphi_t$ .

We fix a Margulis constant  $\varepsilon_0$  for both dimensions 2 and 3.

We again consider a geodesic segment  $[y_0, y_u] = \{y_t : t \in [0, u]\}$  and let  $S_t, \sigma_t$  be as in 7.1. We fix a vertically efficient decomposition as in 5.7 into sets  $\alpha_j \times \ell_j$ ,  $1 \leq j \leq R$ . Write

$$I_j = \{t : y_t \in \ell_j\}.$$

Then the model manifold  $M = M(y_0, y_u)$  is given topologically by

$$M = S \times [0, u] = \cup_{j=0}^R \alpha_j \times I_j,$$

where each  $\alpha_j$  is either a gap — not including the boundary — or a loop. It remains to define the model Riemannian metric  $\sigma_M$ . This is done by decomposing  $M$  into pieces corresponding to each  $(\alpha_j, \ell_j)$ .

Now for each  $j$  and  $t \in I_j$  we define a subsurface  $S_{j,t} \subset S_t$  which is homotopic to  $\varphi_t(\alpha_j)$ , for any  $[\varphi_t] = y_t$ . If  $\alpha_j$  is a loop, then we know from 5.5 that  $|\varphi_t(\alpha_j)| \leq L$ . Then there is  $\varepsilon(L)$ , continuous in  $L$ , such that closed  $\varepsilon(|\varphi_t(\gamma_i)|)$ -neighbourhoods  $T(\varphi_t(\gamma_i))$  of nonintersecting geodesics  $\varphi_t(\gamma_i)$  are disjoint, for any hyperbolic surface ([13] Chapter 4). In fact, we can take  $T(\gamma)$  to be the  $\varepsilon_0$ -Margulis tube  $T(\varphi_t(\gamma), \varepsilon_0)$ , if  $|\varphi_t(\gamma)| \leq \frac{1}{2}\varepsilon_0$ .

We define

$$S_{j,t} = T(\varphi_t(\alpha_j))$$

if  $\alpha_j$  is a loop. If  $\alpha_j$  is a gap, then  $S_{j,t}$  is the closure of the component of the complement of

$$\cup\{T(\varphi_t(\gamma)) : \gamma \subset \partial\alpha_j\}$$

which is homotopic to  $\varphi_t(\alpha_j)$ . Note that the vertically efficient conditions ensure that, if  $\alpha_j$  is a loop,  $S_{j,t}$  has modulus bounded below in terms of the ltd parameter functions for all  $t \in I_j$ . If  $\alpha$  is any loop such that  $\alpha = \alpha_j$  or  $\alpha \subset \partial\alpha_j$  for at least one  $j$ , then let  $I(\alpha) = [s(\alpha), t(\alpha)]$  be the connected union of the  $I_j$  for all such  $j$ . For such a  $j$  for which  $\alpha_j$  is a loop, define  $S_{j,t} = S_{\alpha,t}$ . Then, except when  $s(\alpha) = 0$ , or similarly  $t(\alpha) = u$ ,  $S_{\alpha,s(\alpha)}$  and  $S_{\alpha,t(\alpha)}$  also have modulus bounded above. This is because of the properties of vertically efficient in 5.7. These properties imply that we also have  $s(\alpha) \in I_k$  (or similarly  $t(\alpha) \in I_k$ ) for some ltd gap or loop  $\alpha_k$ . Here,  $\alpha_k$  either contains  $\alpha$  in its interior, or intersects  $\alpha$ , transversely. So there is a loop  $\gamma \in \alpha_k$  which intersects  $\alpha$  transversely, and with  $|\varphi_t(\gamma)|$  bounded in terms of the ltd parameter functions.

Now we choose a continuous family of  $C^1$  diffeomorphisms,  $\varphi_t$ , with  $[\varphi_t] = y_t$ , and satisfying the following properties. In all the following let  $y_s, y_t \in \ell_j$ .

1.  $\varphi_t \circ \varphi_s^{-1}(S_{j,s}) = S_{j,t}$  and  $\varphi_t \circ \varphi_s^{-1}(\partial S_{j,s}) = \partial S_{j,t}$ .
2.  $\varphi_t \circ \varphi_s^{-1} : S_{j,s} \rightarrow S_{j,t}$  has bounded derivative with respect to the norm induced by the metrics  $\sigma_s, \sigma_t$  if  $(\alpha_j, \ell_j)$  is bounded, or if  $|s - t| \leq 1$  and  $(\alpha_j, \ell_j)$  is ltd.
3. For any loop  $\alpha$  which is  $\alpha_j$  for at least one  $j$ ,  $\partial S_{\alpha,s}, \varphi_t \circ \varphi_s^{-1} : \partial S_{\alpha,s} \rightarrow \partial S_{\alpha,t}$  has constant derivative with respect to the length induced by  $\sigma_s, \sigma_t$ .
4. For each loop  $\alpha$  which is  $\alpha_j$  for at least one  $j$ , for a fixed loop  $\beta(\alpha)$  intersecting  $\alpha$  at least once and at most twice, (as in 2.6) and chosen intersection point  $x$ , let  $\beta_t$  be the closed geodesic on  $S_t$  which is isotopic to  $\varphi_t(\beta(\alpha))$ , and let  $x_t$  be the image of  $x$  under such an isotopy which also isotopes  $\varphi_t(\alpha)$  to a closed geodesic. Let  $x_{1,t}$  and  $x_{2,t}$  be the endpoints in  $\partial S_{\alpha,t}$  of the geodesic arc of  $\beta_t$  containing  $x_t$ . Then for all  $s, t \in I_j$ , and  $i = 1, 2$ ,

$$\varphi_t \circ \varphi_s^{-1}(x_{i,s}) = x_{i,t}.$$

There are many such choices of  $\varphi_t$ , because the conditions on  $\varphi_t$  are conditions on  $\varphi_t \circ \varphi_s^{-1}$  restricted to  $S_{j,s}$ , for  $s < t$ ,  $s, t \in I_j$  for some  $j$ . But we cannot, as in the bounded geometry case, take  $\varphi_t \circ \varphi_s^{-1}$  to minimise distortion for all  $s < t$ .

If  $\alpha_j$  is a gap, let

$$W_j = \cup_{t \in I_j} \varphi_t^{-1}(S_{j,t}) \times \{t\}$$

Thus,  $W_j$  is a submanifold of  $S \times [0, u]$  with piecewise smooth boundary which is homeomorphic to  $\alpha_j \times I_j$ . We also define

$$W_{j,t} = \varphi_t^{-1}(S_{j,t}) \times \{t\}.$$

If  $\alpha$  is a loop which is  $\alpha_j$  for at least one  $j$ , let

$$T(\alpha) = \cup_{t \in I(\alpha)} \varphi_t^{-1} S_{\alpha,t} \times \{t\}$$

Then  $T(\alpha)$  is solid torus, called a *model Margulis tube*. No metric has been specified yet.

### 7.3 Geometrically finite Kleinian surface case with two ends: metric on the complement of model Margulis tubes.

Let  $\alpha_j$  be a gap. We define the model Riemannian metric  $\sigma_M$  on  $W_{j,t}$  by

$$\sigma_M = \varphi_t^*(\sigma_t) + (c_j(z, t))^2 dt^2,$$

where, if we do not mind discontinuities in the metric on the union of all such  $W_{j,t}$ , we can take  $c_j(z, t) = c_j$  to be constant, and

$$c_j = \begin{cases} 1 & \text{if } (\alpha_j, \ell_j) \text{ is ltd,} \\ \frac{1}{|\ell_j|+1} & \text{if } (\alpha_j, \ell_j) \text{ is bounded.} \end{cases}$$

If we want the metric to be continuous or smooth, we can adjust  $c_j(z, t)$  in a neighbourhood of  $\partial W_j \cap \partial W_k$  whenever  $\alpha_j$  and  $\alpha_k$  are both gaps. As in the case of combinatorial bounded geometry, the precise definition of  $\varphi_t$  does not change the metric up to Lipschitz equivalence, because the map

$$(z, t) \mapsto (\varphi_s(z), t) : (W_j, \sigma_M) \rightarrow (S_{j,s} \times I_j, \sigma_s + dt^2)$$

is boundedly Lipschitz if  $(\alpha_j, \ell_j)$  is bounded, and boundedly Lipschitz restricted to the set of  $(z, t) \in W_j$  with  $|s - t| \leq 1$  if  $(\alpha_j, \ell_j)$  is ltd.

### 7.4 Margulis tubes.

For any three-dimensional  $\varepsilon_0$ -Margulis tube, the metric induced by the hyperbolic metric on the tube boundary is Euclidean, and Margulis tubes are also parametrised by  $H^2$ , at least if the core loop is sufficiently short compared to  $\varepsilon_0$ . Indeed, if we choose coordinates so that in the upper half-space model of hyperbolic space  $H^3$ , the Margulis tube has lift

$$\{(re^{t+i\theta}, e^t) : 0 \leq r \leq R, \theta, t \in \mathbb{R}\},$$

and the hyperbolic isometry corresponding to the core loop is the map  $(z, u) \mapsto (e^\lambda z, e^{\operatorname{Re}(\lambda)u})$  with  $\lambda = o(\varepsilon_0)$ . Then the metric on the boundary of the tube is

$$R^2 d\theta^2 + (R^2 + 1) dt^2.$$

and the corresponding element of the Teichmüller space of the torus,  $H^2$ , is

$$\frac{2\pi i}{\lambda} (1 + R^{-2})^{1/2},$$



and

$$R = \frac{\varepsilon_0}{|\lambda|}(1 + f(\lambda)),$$

for a  $C^1$  function  $f$  with  $f(\lambda) = 0$ . We write

$$\frac{2\pi i}{\lambda} = B + iA.$$

The length of the core loop is then  $\operatorname{Re}(\lambda)$ , and

$$\operatorname{Re}(\lambda) = \frac{2\pi A}{|B + iA|^2},$$

and

$$R = \frac{\varepsilon_0 |B + iA|}{2\pi}(1 + f(\lambda)).$$

See also [36] Section 6.

## 7.5 Geometrically finite Kleinian surface case with two ends: the metric on model Margulis tubes.

It remains to define the metric  $\sigma_M$  on the model Margulis tube  $T(\alpha)$  if  $\alpha = \alpha_j$  for at least one  $j$ . We have

$$\partial T(\alpha) = \cup_{k=1}^4 \partial_k T(\alpha)$$

where  $\partial_1 T(\alpha)$  and  $\partial_2 T(\alpha)$  are the components of  $\partial_v T(\alpha)$  (vertical boundary), where

$$\partial_v T(\alpha) = \cup_{t \in I(\alpha)} \varphi_t^{-1}(\partial S_{\alpha,t}),$$

and

$$\partial_3 T(\alpha) = \varphi_{s(\alpha)}^{-1} S_{\alpha,s(\alpha)}, \quad \partial_4 T(\alpha) = \varphi_{t(\alpha)}^{-1} S_{\alpha,t(\alpha)}.$$

We define the metric restricted to  $\partial_3 T(\alpha)$  to be  $\varphi_{s(\alpha)}^*(\sigma_{s(\alpha)})$  and similarly for  $\partial_4 T(\alpha)$ . On  $\partial_v T(\alpha)$ , the metric has already been defined from the definitions of the metric on the sets  $W_k$  for gaps  $\alpha_k$ . We choose the numbering of  $\partial_1 T(\alpha)$  and  $\partial_2 T(\alpha)$  so that  $x_{k,t} \in \partial_k T(\alpha)$  for  $t \in I(\alpha)$ , for  $x_{k,t}$  as in 4 of 7.3. Both components  $\partial_k S_{\alpha,t}$  ( $i = 1, 2$ ) of  $\partial S_{\alpha,t}$  (using the index  $k$  in the same way) have the same length  $\lambda_\alpha(t)$  in the  $\sigma_t$  metric, by the definition of  $S_{\alpha,t}$ , since  $\lambda_\alpha(t)$  is a function of  $|\varphi_t(\alpha)|$ . Choose an orientation on  $S_{\alpha,t}$ . For  $x \in \partial_k S_{\alpha,s}$  let  $\lambda_\alpha(x, s)$  denote the length of the shortest positively oriented segment from  $x_{k,s}$  to  $x$ , so that

$$g_\alpha(z) = \exp(2\pi i \lambda_\alpha(x, s) / \lambda_\alpha(s))$$

is smooth on  $\partial_1 S_{\alpha,s}$ ,  $\partial_2 S_{\alpha,s}$ . Then, by the properties of 7.3, in particular 3 and 4 of 7.3, the map

$$(x, t) \mapsto (g_\alpha \circ \varphi_s(x), t) : \partial_v T(\alpha) \rightarrow \{e^{2\pi i \theta} : \theta \in \mathbb{R}\} \times I(\alpha)$$

pushes forward  $\sigma_M$  on  $\partial_k T(\alpha)$  ( $k = 1, 2$ ) to

$$d\theta^2 + (c_{\alpha,k}(t)^2)dt^2$$

on  $S^1 \times I(\alpha)$ , for a suitable function  $c_{\alpha,k}(t)$ . Even if we chose the  $c_k(x, t)$  with some dependence on  $x$  in order to smooth the metric, the choice can be made so that the  $c_{\alpha,k}(t)$  depend only on  $t$ , because for each  $t$ ,  $\partial S_{\alpha,t} \subset \partial S_{m,t}$  for at least one gap  $\alpha_m$ , and intersects no other  $\partial S_{p,t}$  for a gap  $\alpha_p$ , unless  $t$  is an endpoint of  $I_m$ , in which case there is exactly one other gap  $\alpha_p$  with  $\partial S_{\alpha,t} \subset \partial S_{p,t}$ . We can then make a further change of variable

$$C_{\alpha,k}(t) = \int_{s(\alpha)}^t c_{\alpha,k}(u) du.$$

Then

$$(e^{i\theta}, t) \mapsto (e^{i\theta}, C_{\alpha,k}(t)) : S^1 \times I(\alpha) \rightarrow S^1$$

pushes forward  $d\theta^2 + (c_{\alpha,k}(t)^2)dt^2$  to  $d\theta^2 + dt^2$  on  $S^1 \times [0, A_k(\alpha)]$  where  $A_k(\alpha) = C_{\alpha,k}(t(\alpha))$ . So  $\sigma_M$  on  $\partial_k T(\alpha)$  is a Euclidean metric, up to isometry.

The annuli  $\partial_3 T(\alpha) = S_{\alpha,s(\alpha)}$  and  $\partial_4 T(\alpha) = S_{\alpha,t(\alpha)}$  are conformally equivalent to Euclidean annuli  $S^1 \times [0, A_3(\alpha)]$  and  $S^1 \times [0, A_4(\alpha)]$ . The uniformising maps restricted to boundaries have constant derivative with respect to length. If the modulus of  $S_{\alpha,s(\alpha)}$  is bounded above and below then the uniformising map transforms  $\sigma_{s(\alpha)}$  to a metric which is boundedly Lipschitz equivalent to the Euclidean metric  $d\theta^2 + dt^2$  on  $S^1 \times [0, A_3(\alpha)]$ . The modulus is bounded below in terms of  $L$ . It is also bounded above if  $|\varphi_{s(\alpha)}(\alpha)|$  is bounded above, which is true if  $s(\alpha) > 0$  by the definition of vertically efficient, as already noted. Similar statements hold for  $\partial_4 T(\alpha)$ .

So now if we match up boundaries, we have a conformal map

$$\psi_\alpha : \partial T(\alpha) \rightarrow (S^1 \times [0, A]) / \sim,$$

where

$$A = A(\alpha) = \frac{1}{2\pi} \sum_{k=1}^4 A_k(\alpha),$$

and where  $\sim$  is defined by

$$(0, e^{2\pi i\theta}) \sim (e^{2\pi i(B+\theta)}, 2\pi A)$$

for some  $B = B(\alpha) \in \mathbb{R}$ . We now determine  $B(\alpha)$ . The meridian is

$$\beta_{s(\alpha)} \cup \{x_{1,t} : t \in I(\alpha)\} \cup \{x_{2,t} : t \in I(\alpha)\} \cup \beta_{t(\alpha)},$$

for  $x_{i,t}$  and  $\beta_t$  as in 4 of 7.2. Thus,  $\beta_t$  is an arc of the geodesic homotopic to  $\varphi_t(\beta(\alpha))$ , for a fixed loop  $\beta(\alpha)$  intersecting  $\alpha$  at most twice (as in 7.2). Then  $n_\alpha(y_{s(\alpha)})$  is, by definition, the value of  $n$  which minimises  $|\varphi_{s(\alpha)} \tau_\alpha^n(\beta(\alpha))|$ , where here (as earlier)  $\tau_\alpha$  denotes oriented Dehn twist around  $\alpha$ . A similar statement holds for  $t(\alpha)$ . Then because  $S_{\alpha,t}$  has modulus bounded from 0 in terms of

the ltd parameter functions, the number of twists of  $\beta_{s(\alpha)}$  round the geodesic homotopic to  $\varphi_{s(\alpha)}(\alpha)$  is within  $C$  of  $-n_\alpha(y_{s(\alpha)})$ , where  $C$  depends only on the ltd parameter functions. A similar statement holds for  $S_{\alpha,t(\alpha)}$ ,  $\beta_{t(\alpha)}$ ,  $n_\alpha(y_{t(\alpha)})$ . It follows that

$$B = \operatorname{Re}(-\pi_\alpha(y_{t(\alpha)}) + \pi_\alpha(y_{s(\alpha)})) = -n_\alpha(y_{t(\alpha)}) + n_\alpha(y_{s(\alpha)}) + O(1), \quad (7.5.1)$$

where the  $O(1)$  term depends on the ltd parameter functions.

It is natural to take generators of the fundamental group of  $\partial T(\alpha)$  as follows. Take the homotopy class of  $\partial_k T(\alpha)$  for any  $k$ , equivalently of the components of  $\varphi_t^{-1}(S_{\alpha,t})$  (any  $t \in I(\alpha)$ ) as first generator and the meridian as second generator. As usual, identify the Teichmüller space of the torus with the upper half-plane  $H^2$ , using these generators. The corresponding point in  $H^2$  is then  $B + iA$ . Note that  $A$  is bounded from 0 in terms of the ltd parameter functions, since  $A_3$  and  $A_4$  are.

Now suppose that

$$\text{either } s(\alpha) > 0 \text{ or } |\varphi_0(\alpha)| \geq \varepsilon_0,$$

and

$$\text{either } t(\alpha) < u \text{ or } |\varphi_u(\alpha_j)| \geq \varepsilon_0.$$

The map  $\psi_\alpha$  is a bounded Lipschitz equivalence from  $\sigma_M|\partial T(\alpha)$  to the Euclidean metric  $d\theta^2 + dt^2$ . Now we define the metric  $\sigma_M$  on  $T(\alpha)$  using a model Margulis tube  $T(B + iA)$  such that the corresponding point in  $H^2$  for  $\partial T(B + iA)$  is exactly  $B + iA$  whenever  $B + iA$  is sufficiently large given  $\varepsilon_0$ , and within a bounded distance of it otherwise. This is possible, because the map  $f$  of 7.4 is boundedly  $C^1$ . Let  $\sigma$  be the hyperbolic metric on  $T(B + iA)$ . We can assume that map  $\psi_\alpha$  already defined on  $\partial T(\alpha)$  has image  $\partial T(B + iA)$ , and under the current assumption it is boundedly biLipschitz on  $\partial T(\alpha)$ . Then extend diffeomorphically on the interior to

$$\psi_\alpha : T(\alpha) \rightarrow T(B + iA)$$

Then we define

$$\sigma_M = \psi_\alpha^* \sigma \text{ on } \operatorname{Int}(T(\alpha)). \quad (7.5.2)$$

If we wish to keep the metric continuous and smooth we can smooth it in the preimage of a small neighbourhood of  $\partial T(\alpha)$  in  $T(\alpha)$ .

Now suppose that

$$t(\alpha) = u, \quad |\varphi_u(\alpha)| < \varepsilon_0,$$

and

$$\text{either } s(\alpha) > 0 \text{ or } |\varphi_0(\alpha)| \geq \varepsilon_0.$$

Write

$$\varepsilon_2 = |\varphi_u(\alpha)|.$$

Then we identify  $\partial_1 T(\alpha) \cup \partial_2 T(\alpha) \cup \partial_3 T(\alpha)$  with an annulus in the tube boundary of a Margulis tube  $T(B + iA')$  corresponding to a point  $B + iA' \in H^2$  for some  $A' \geq A'' = A_1 + A_2 + A_3$  and for  $B$  as above. The length of the core loop will be  $\varepsilon_1$ . Then, as in 7.4, the relation between  $\varepsilon_1$  and  $B + iA'$  is

$$\varepsilon_1 = \frac{2\pi A'}{|B + iA'|^2}.$$

We want to determine  $\varepsilon_1$  so that the Poincaré metric on  $S_{\alpha, t(\alpha)}$  is the same, up to bounded distortion, as that of an annulus in the Margulis tube with boundary components on the boundary of the Margulis tube, and separated in the boundary of the Margulis tube an annulus of area  $A''$  on one side, and an annulus of area  $A' - A''$  on the other. If

$$\varepsilon_2 \leq \frac{2\pi \cdot 2A''}{|B + 2iA''|^2},$$

then we define  $\varepsilon_1 = \varepsilon_2$ . If

$$\frac{2\pi \cdot 2A''}{|B + 2iA''|^2} \leq \varepsilon_2 \leq \frac{2\pi}{|B + 2iA''|}$$

then we define

$$A' = 2A''.$$

The reason for this is that the radius of injectivity within diameter 1 of a core loop of complex length  $\lambda$  varies between  $|\lambda|$  and  $\operatorname{Re}(\lambda)$ . Finally, if

$$\frac{2\pi}{|B + 2iA''|} \leq \varepsilon_2$$

then  $A'' < A' < 2A''$ , so that  $(A' - A'')/A' < \frac{1}{2}$ , and we define  $A'$  implicitly by the equation

$$\log((A' - A'')/A') + \log|A' + iB| + \log \varepsilon_2 = 0.$$

Now we take an annulus  $Y(B, A', A'')$  in  $\partial T(B + iA')$  which is proportion  $A''/A'$  of the total area and bounded by geodesics in the Euclidean metric in the homotopy class in  $T(\alpha)$  of the core loop and of Euclidean length  $\varepsilon_0(1 + O(\varepsilon_0))$ . The homotopy class of these geodesics in  $\partial T(B + iA')$  is determined uniquely, at least if  $\varepsilon_1 = o(\varepsilon_0)$ , and in the other case it does not matter which homotopy class we choose with these properties. Now we define an annulus  $C(B, A', A'') \subset T(B + iA')$  with the same boundary as  $Y(B, A', A'')$ . If  $\varepsilon_2 \cdot |B + 2iA''| > 2\pi$ , we take  $C(B, A', A'')$  to be the union of the shortest geodesics in  $T(B + iA')$  joining points in different components of  $\partial Y(\alpha)$ . These geodesics all have length  $-2\log(\varepsilon_2) + O(1)$  and their union is indeed an annulus. The annulus comes within a bounded distance of the core loop if  $\varepsilon_2 \cdot |B + 2iA''|$  is bounded. If  $\varepsilon_2 \cdot |B + 2iA''| \leq 2\pi$ , then we take  $C(B, A', A'')$  to be the union of two annuli, formed by taking the union of geodesics in  $T(B + iA')$  from each component

of  $\partial Y(B, A', A'')$  to the core loop of  $T(B + iA')$ , meeting both  $\partial Y(B, A', A'')$  and the core loop perpendicularly. In this case,  $C(B, A', A'')$  is not a smooth manifold if  $A' > 2A''$ , but this is what we want. In this case, the convex hull boundary of the corresponding hyperbolic manifold has a sharp angle at the geodesic  $(\alpha)_*$ , where  $\alpha_*$  is the closed geodesic in the free homotopy class represented by  $\alpha$ .

We also have a piecewise smooth diffeomorphism

$$\psi_\alpha : \partial T(\alpha) \rightarrow C(B, A', A'') \cup Y(B, A', A''),$$

smooth on the interior of  $\partial_k T(\alpha)$  for all  $k$  which is boundedly biLipschitz, which maps  $\partial_4 T(\alpha)$  to  $C(B, A', A'')$  and  $\partial_1 T(\alpha) \cup \partial_2 T(\alpha) \cup \partial_3 T(\alpha)$  to  $Y(B, A', A'')$ . Let  $T(B, A', A'')$  be the part of  $T(B + iA')$  bounded by  $C(B, A', A'') \cup Y(B, A', A'')$ . when we extend to  $\psi_\alpha$  to a diffeomorphism on the interior

$$\psi_\alpha : T(\alpha) \rightarrow T(B, A', A'')$$

and then we again use (7.5.2) to define  $\sigma_M$  on  $T(\alpha)$ .

The case when

$$s(\alpha) = 0, \quad |\varphi_0(\alpha)| < \varepsilon_0,$$

and

$$\text{either } t(\alpha) < u \text{ or } |\varphi_u(\alpha)| \geq \varepsilon_0$$

is treated exactly similarly, with  $s(\alpha)$  and  $t(\alpha)$  interchanged, and also 0 and  $u$ ,  $A_3$  and  $A_4$ , and  $\partial_3 T(\alpha)$  and  $\partial_4 T(\alpha)$ . Finally we consider the case when

$$s(\alpha) = 0, \quad |\varphi_0(\alpha)| < \varepsilon_0,$$

$$t(\alpha) = u, \quad |\varphi_u(\alpha)| < \varepsilon_0.$$

In this case we define

$$\varepsilon_2 = |\varphi_T(\alpha)|,$$

$$\varepsilon_3 = |\varphi_0(\alpha)|,$$

and suppose without loss of generality that  $\varepsilon_2 \leq \varepsilon_3$ . Then we define

$$A'' = A_1 + A_2 + \frac{1}{\varepsilon_3}.$$

Then we define  $\varepsilon_1$  and  $A'$  exactly as before.

The set  $Y(B, A', A'')$  now has two components whose areas are  $A_1$  and  $A_2$  chosen Euclidean distance  $1/\varepsilon_3$  apart in  $\partial T(B + iA')$  with respect to the Euclidean metric on  $\partial T(B + iA')$ . The set  $C(B, A', A'')$  also has two components, one of which is a union of geodesic segments in  $T(B + iA')$  joining the components of  $\partial Y(B, A', A'')$  which are Euclidean distance  $1/\varepsilon_3$  apart, in which case these geodesics have length  $2 \log(\varepsilon_0/\varepsilon_3) + O(1)$ . The other component is defined as before, depending on whether  $\varepsilon_2 \cdot |B + 2iA''| > 2\pi$  or  $\varepsilon_2 \cdot |B + 2iA''| \leq 2\pi$ . Then, as before, we take  $T(B, A', A'')$  to be the subset of  $T(B + iA')$  bounded

by  $Y(B, A', A'')$  and  $C(B, A', A'')$  and pull back the metric using the formula of (7.5.2).

The metric  $\sigma_M$  has now been completely determined, up to bounded distortion, in the case of two geometrically finite ends. If the model Margulis tube is homotopic to  $\{t\} \times \gamma$  then we call the core loop  $\gamma_{**}$  and shall sometimes call the solid torus  $T(\gamma_{**})$  rather than  $T(\gamma)$  as above.

## 7.6 Geometrically finite Kleinian surface case: more than two ends.

We now consider the case of a model relative Scott core  $M_c$  being homeomorphic to  $S_d \times [0, 1]$ , where  $S_d$  is a compact surface with boundary, but where the boundary component homeomorphic to  $S_d \times \{k\}$  might be a union of several components of  $\partial_d M_c$  and of annuli  $\partial M_c \cap \partial M_d$ , for  $k = 0, 1$ . This is an example of something which happens regularly. It is sometimes convenient to group ends of  $M_d$  together, by taking a connected submanifold  $W$  of  $M_c$ , and considering the group of ends in each component of  $M_d \setminus W$ . We only do this for  $W$  with the following properties: each component of  $\partial W \cap \partial M_d$  is an essential annulus in  $\partial M_d$ , a component of  $\partial W \setminus \partial M_d$  cannot be homotoped into  $\partial M_d$ , and if  $S_d$  is a component of  $\partial W \setminus \partial M_d$  bounding the closure  $U$  of a nonempty component of  $M_c \setminus W$ , then  $(U, S_d)$  is homeomorphic to  $(S_d \times [0, 1], S_d)$ . In such cases, as in this case, we define  $M_{d,W}$  to be the union of  $M_d$  and components of  $M \setminus M_d$  which are disjoint from  $W$ . Then if  $U'$  is the closure of the component of  $M_{d,W} \setminus W$  bounded by  $S_d$ ,  $(U', S_d, \partial M_{d,W} \cap U')$  is homeomorphic to  $(S_d \times [0, \infty), S_d, \partial S_d \times [0, \infty))$ . So  $U'$  contains a unique end of  $M_{d,W}$ .

In the current case,  $W$  is homeomorphic to  $S_d \times [0, 1]$ , and  $M_{d,W}$  has two ends, which we call  $e_{\pm}$ . Then  $S_d$  is the horodisc deletion of a finite type surface  $S$ , and we have two isotopically disjoint multicurves  $\Gamma(e_{\pm})$ , one of which could be empty, and the end invariants of  $M_d$  split into two sets  $(y_1, \dots, y_m)$  and  $(y_{m+1}, \dots, y_p)$ , which are the end invariants  $\mu(e_{\pm})$  of  $e_{-}$ ,  $e_{+}$  respectively. Here,  $y_i \in \mathcal{T}(S(\alpha_i))$  and  $\alpha_i$  ( $1 \leq i \leq m$ ) are the gaps of  $\Gamma(e_{-})$ , while  $\alpha_i$  ( $m+1 \leq i \leq p$ ) are the gaps of  $\Gamma(e_{+})$ . In this case we make the model manifold  $M(\mu(e_{-}), \mu(e_{+}))$  by slightly modifying the model  $M(y(e_{-}), y(e_{+}))$  of 7.2 to 7.5, (replacing  $y_0$  and  $y_u$  by  $y(e_{-})$  and  $y(e_{+})$ ) where  $y(e_{-}) \in \mathcal{T}(S)$  is defined using the  $y_i$  for  $1 \leq i \leq m$  and  $y(e_{+})$  is defined using the  $y_i$  for  $m+1 \leq i \leq p$ .

Fix a Margulis constant  $\varepsilon_0$ . We define  $y(e_{-})$  up to bounded distance by defining projections  $\pi_{\alpha_i}(y(e_{-}))$  for  $1 \leq i \leq m$  and  $\pi_{\gamma}(y(e_{-}))$  for  $\gamma \in \Gamma(e_{-})$ . So we take

$$\begin{aligned} \pi_{\alpha_i}(y(e_{-})) &= y_i, \quad 1 \leq i \leq m, \\ \pi_{\gamma}(y(e_{-})) &= \frac{1}{\varepsilon_0}, \quad \gamma \in \Gamma_0. \end{aligned}$$

The definition of the imaginary part of  $\pi_{\gamma}$  needs a choice of loop transverse to  $\gamma$  intersecting it at most twice (see 2.6). But in fact this choice is irrelevant, because the imaginary part only influences the geometry of the Margulis tube, which we are about to remove. If  $y(e_{-}) = [\varphi_0]$  then we have chosen to make

$|\varphi_0(\gamma)| = \varepsilon_0(1 + o(1))$ . We define  $y(e_+)$  similarly, using  $\Gamma(e_+)$  and  $y_i$  for  $m < i \leq p$ .

The model manifold  $M(\mu(e_-), \mu(e_+))$  is then obtained from the model  $M(y(e_-), y(e_+))$  constructed in 7.2 to 7.5 by simply leaving out the models for parts of Margulis tubes  $T(\gamma_{**})$  for  $\gamma \in \Gamma(e_-) \cup \Gamma(e_+)$ , and gluing in, instead, pieces of horoballs with the right boundary piece. We only do this when  $\gamma \in \Gamma(e)$  for just one of  $e = e_+, e_-$ . So we have a single annulus  $A$ , with a Euclidean metric. We simply glue in  $A \times [0, \infty)$  with the product Euclidean metric.

## 7.7 Independence of ltd parameter functions.

The model manifold is independent, up to Lipschitz equivalence with bound depending on the ltd parameter functions, of the decomposition of 5.7, and of the long thick and dominant parameter functions used. To see this, we only need to compare the models given by vertically efficient partitions for two sets of ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$ ,  $(\Delta_2, r_2, s_2, K_2)$  where  $(\Delta_2, r_2, s_2, K_2)$  is sufficiently stronger than  $(\Delta_1, r_1, s_1, K_1)$  for the following to hold. For all  $\nu$ ,  $r_2(\nu) \leq \nu_1$ , where every ltd gap in the decomposition for  $(\Delta_1, r_1, s_1, K_1)$  is long  $\nu$ -thick and dominant for some  $\nu \geq \nu_1$ . We can choose this  $\nu_1$  so that, if  $(\alpha, \ell)$  is bounded for  $(\Delta_1, r_1, s_1, K_1)$ , then  $|\varphi(\gamma)| \geq \nu_1$  for all  $\gamma$  in the interior of  $\alpha$ . Such a  $\nu_1$  exists by 5.4, and the definition of bounded in 5.5 and the results discussed there. So if  $(\alpha_i, \ell_i)$  is in the partition for  $(\Delta_i, r_i, s_i, K_i)$  and  $\ell_1 \cap \ell_2 = \emptyset$ , we have  $\alpha_1 \subset \alpha_2$  or  $\alpha_1 \cap \alpha_2 = \emptyset$ . If  $\alpha_1 \subset \alpha_2$ , then the vertically efficient condition for the second partition implies that  $\ell_1 \subset \ell_2$ . So then the partition for  $(\Delta_1, r_1, s_1, K_1)$  is a refinement of the partition for  $(\Delta_2, r_2, s_2, K_2)$ . We can assume that the same family  $\varphi_t$  is used to define both models, because, as already noted in 7.3, the choice of family, subject to the conditions of 7.2, does not affect the model.

Let  $(\alpha_1, \ell_1)$  be ltd for  $(\Delta_1, r_1, s_1, K_1)$  with  $(\alpha_1 \times \ell_1)$  in the first partition. Let  $\alpha_2 \times \ell_2$  be from the second partition, with  $\alpha_1 \times \ell_1 \subset \alpha_2 \times \ell_2$ . If  $(\alpha_2, \ell_2)$  is bounded for  $(\Delta_2, r_2, s_2, K_2)$  then we have a bound on  $|\ell_1|$  in terms of  $(\Delta_2, r_2, s_2, K_2)$  and also a bound on the number of such  $(\alpha_1, \ell_1)$  in terms of  $(\Delta_2, r_2, s_2, K_2)$ . Write  $W_1, W_2$  for the corresponding subsets of  $S \times [0, u]$ , as defined in 7.2, so that  $W_1 \subset W_2$ . Then the scaling factor  $c_1(t)$  is bounded, since  $|\ell_1|$  is bounded. If  $\alpha_1$  is a loop, the corresponding Margulis tube has bounded geometry, because the quantity  $B_1$  of 7.4 is bounded in terms of  $\text{Re}(\pi_{\alpha_j}(y_{t_j}) - \pi_{\alpha_j}(y_{s_j}))$ , if  $\ell_j = [y_{s_j}, y_{t_j}]$ , by (7.5.1). So, in both models, the model metric on  $W_2$  is of bounded geometry, with bound depending on the ltd parameter functions. Now suppose the same situation, but with  $(\alpha_2, \ell_2)$  ltd with respect to  $(\Delta_2, r_2, s_2, K_2)$ . If  $\alpha_1$  is a gap then the metric on  $W_1$  is defined in exactly the same way in both models. If  $\alpha_1$  is a loop properly contained in  $\alpha_2$ , then  $|\varphi(\alpha)| \geq \nu_2$  for all  $[\varphi] \in \ell_1$ , for  $\nu_2$  such that for every  $(\alpha, \ell)$  for in the second partition with  $\alpha$  a gap,  $\alpha$  is long  $\nu$ -thick and dominant for some  $\nu \geq \nu_2$ . Then  $W_1$  has bounded geometry in both models. So the geometry of all pieces is boundedly equivalent in both models.

## 7.8 Consistency with previous models and examples.

The model metric on Margulis tubes in 7.5 is effectively exactly the same as the model metric on Margulis tubes in Minsky's punctured torus paper [36]. The situation here is slightly more general, because the quantity  $A = A(\alpha)$  of 7.5 can be arbitrarily large, although it is always bounded from 0. In the punctured torus case it is bounded above also, because in the punctured torus case, if  $(\alpha, \ell)$  is ltd, then either  $\alpha$  is the punctured torus itself, or  $\alpha$  is a loop.

This model is also consistent with Rafi's examples [48], [49]. In one of Rafi's examples, adapting to the present notation,  $S$  is a closed surface of genus two, and  $[y_0, y_u]$  is the union of two segments  $\ell_1, \ell_2$  along which loops  $\gamma_1$  and  $\gamma_2$  respectively are  $K_0$ -flat, apart from a bounded segment in the middle. The convex hull of  $\gamma_1$  and  $\gamma_2$  could be (for example)  $S \setminus \gamma_3$  where  $\gamma_3$  is not  $K_0$ -flat along any segment and not adjacent to  $\alpha$  for any ltd  $(\alpha, \ell)$ . In the corresponding hyperbolic manifold with ending data  $(y_-, y_+)$  (or even close to this) the geodesic loop  $(\gamma_3)_*$  is not short. But for  $[\psi] \in \ell_1 \cup \ell_2$  bounded from  $y_\pm$ ,  $|\psi(\gamma_3)|$  is small, and  $\rightarrow 0$  as the length of both  $\ell_1$  and  $\ell_2 \rightarrow \infty$ . Rafi proves this, and there is also a proof of something similar in 15.21 of [50]. However, in the model manifold  $M(y_-, y_+)$ , the geodesic  $(\gamma_3)_{**}$  is not short. A model Margulis tube is inserted, but it has bounded geometry.

## 7.9 Model manifolds for compression bodies.

We now have a geometric model for any hyperbolic 3-manifold  $N$  for which  $(N_d, \partial N_d)$  is homeomorphic to  $(S_d \times \mathbb{R}, \partial S_d \times \mathbb{R})$  for the horodisc deletion of a finite type surface  $S$ . The model in the general case is obtained by gluing together interval bundle models. The data for a model is given by topological type and ending lamination data. The ending lamination data for an end  $e$  in  $\mathcal{T}(S(e)) \cup \mathcal{O}_a(S(e))$  is given relative to an identification of the corresponding boundary component of the relative Scott core with  $S(e)$ . The definition of  $\mathcal{O}_a(S(e))$  is given relative to a base metric on  $S(e)$ . We are now going to discuss this identification and base metric briefly, in the case of relative compression bodies.

There are a number of definitions of compression body in the literature, all of them equivalent. See for example [45], [16], [4]. The relative version which is used here is adapted from the definition of [55]. A relative compression body is, in fact, a compression body in the usual sense, but the decomposition is slightly different.

A *relative compression body*  $(W, \partial W \cap A)$  is a compact manifold with boundary  $\partial W$ , with  $\partial W \setminus A$  a finite disjoint union of surfaces with boundary such that all components of  $A$  are closed annuli, and such that there is a subsurface  $S_{d,0}$  of  $\partial W \setminus A$  with the following properties. All components of  $\partial W \setminus (A \cup S_{d,0})$  are incompressible, and the inclusion  $S_{d,0} \rightarrow W$  is surjective on  $\pi_1$ . The surface  $S_{d,0}$  is called the *exterior surface*. We only consider manifolds which have a chance of being homotopy equivalent to hyperbolic manifolds, so we assume also that  $W$  is  $K(\pi, 1)$ .



Now we assume that  $S_{d,0}$  is compressible in  $W$ , since otherwise we are in the interval bundle case. Let  $S_0$  be the surface of which  $S_{d,0}$  is the horodisc deletion. We can then find a multicurve  $\Gamma_0$  on  $S_0$  such that the loops of  $\Gamma_0$  bound disjoint embedded discs in  $W$ , and such that the closure of each component of the complement of the union of the discs and  $\partial M_c$  is either a three-ball or an interval bundle homeomorphic to  $S_{d,i} \times [0, 1]$  for some component  $S_{d,i}$  of  $\partial W \setminus A$ , with the intersection of  $A$  with the interval bundle identifying with  $\partial S_{d,i} \times [0, 1]$ . It is natural to choose our base metric on  $S_0$  so that, for a fixed choice of  $\Gamma_0$ , the loops of  $\Gamma_0$  are of bounded length, and of length bounded from 0. We shall always do this. We therefore fix  $z_{0,0} = [\varphi_0] \in \mathcal{T}(S_0)$  such that  $|\varphi_0(\gamma)|$  is bounded and bounded from 0 for  $\gamma \in \Gamma_0$ . The gaps of  $\Gamma_0$  are  $\alpha_i$ ,  $1 \leq i \leq p$ , where  $p \geq m$ . In  $M$  we can attach a disjoint disc to each loop of  $\Gamma_0$ . This embeds  $\alpha_i$  in a surface  $S_{d,i}$ , which is a union of  $\alpha_i$  and a number of discs. If  $1 \leq i \leq m$  then  $S_{d,i}$  is homotopic in  $W$  to an incompressible component of  $\partial W \setminus A$ , and  $S_{d,i}$  is the horodisc deletion of a surface  $S_i$ . For  $m < i \leq p$ ,  $\alpha_i$  and the attached discs bound a ball in  $W$ . For  $1 \leq i \leq m$  we define  $z_{0,i} \in \mathcal{T}(S_i)$  to have the same conformal structure as  $\pi_{\alpha_i}(z_{0,0})$ , but forgetting the puncture in each disc.

Now suppose that  $M$  is to be the model for a hyperbolic 3-manifold, with  $M_d$  as the model for the horoball deletion, and that  $W \subset M$  is a submanifold of the model for the relative Scott core, as suggested at the start of 7.6. Let  $M_{d,W}$  be as defined in 7.6. The ends of  $M_{d,W}$  are then sets of ends of  $M_d$ . We fix geometrically finite ending invariants for the ends of  $M_d$ . This gives corresponding ending invariants  $\mu(e_i)$  for  $M_{d,W}$ . The model manifold for this end, with these invariants, is obtained by modifying a model  $M(z_{0,i}, y_i)$  to a model  $M(z_{0,i}, \mu(e_i))$  for a certain point  $y_i \in \mathcal{T}(S_i)$ , as described in 7.2 to 7.6. Then the geometric model  $M(\mu(e_0), \dots, \mu(e_m))$  will be made from the models

$$M_k = M(z_{0,k}, \mu(e_k)),$$

$0 \leq k \leq m$ . It seems simpler to construct the geometric manifold for the unquotiented  $(z_{0,k}, y_k)$ , and to note that geometric models for different choices in a single  $\text{Mod}_0(\partial M_k, M_k)$  orbit are homeomorphic.

It only remains to show how to glue together the different models  $M_k$  together. We can assume that these submanifolds are all disjoint in  $M$ , by moving them apart slightly, and we have to extend the metric to the complement. Let  $D(\gamma)$  be the disc in  $M_c$  attached to the loop in  $\partial M_0$  corresponding to  $\varphi_0(\gamma)$ , for  $\gamma \in \Gamma_0$ . We also assume that these discs are all disjoint, and have interiors disjoint from all  $M_i$ . We define the metric on each of these discs to be the pullback of the Euclidean metric on the unit disc, where the map on the boundary is length-preserving with respect to the metric on  $M_0$ . Then we label the components of the complement in  $M$  of the  $M_k$  ( $0 \leq k \leq m$ ) by  $M'_k$  for  $1 \leq k \leq p$ , where  $M'_k$  is between  $M_k$  and  $M_0$  if  $k \leq m$ , and  $M'_k$  is a ball if  $k > m$ . So now we have a metric on the boundary of each complementary component  $M'_k$ ,  $1 \leq k \leq p$ , and the closure of each  $M'_k$  is homeomorphic to  $S_k \times [0, 1]$  if  $1 \leq j \leq m$ . In the case of  $k \leq m$ , the metric on the boundary component

which is also in  $\partial M_k$  is a hyperbolic metric by construction, corresponding to  $y_k \in \mathcal{T}(S_k)$ . For the other boundary component, the metric determines a conformal structure, and hence an element of  $\mathcal{T}(S_k)$  which we claim is a bounded distance from  $z_{0,k}$ . To see this, we consider the definition of  $\pi_{\alpha_k}$ . The corresponding hyperbolic surface is a punctured surface homeomorphic to  $S(\alpha_k)$ . But the definition in 2.6 initially describes this hyperbolic surface in terms of its conformal structure, attaching a punctured disc to the subset of  $\varphi_0(S_0)$  homotopic to  $\varphi_0(\alpha_k)$  and bounded by the geodesics homotopic to  $\varphi_0(\partial\alpha_k)$ . So we might just as well regard this as a conformal structure on the surface, with the punctures corresponding to  $\partial\alpha_k$  removed. We also have a metric, taking the metric on the component of  $\varphi_0(S_0 \setminus \Gamma_0)$  homotopic to  $\varphi_0(\alpha_k)$  and bounded by the geodesics homotopic to  $\varphi_0(\partial\alpha_k)$  and pushing forward the usual Euclidean unit disc metric on the added discs. This metric is boundedly Lipschitz equivalent, under a piecewise smooth diffeomorphism  $\varphi_k$ , to the unique hyperbolic metric which it is conformally equivalent to, which is the metric corresponding to  $z_{0,k}$ . So we can take a metric on  $M'_k$  which is boundedly equivalent to the pullback of product metric  $\sigma_k + dt^2$  on  $S_k \times [0, 1]$ , extending the metric already defined on the boundary. For  $k > m$ , choose the pullback under a homeomorphism  $\varphi'_k$  of a metric on the round unit ball which is boundedly equivalent to the Euclidean metric, extending the metric already defined on the boundary. The metric has some discontinuities since the sets  $\partial M'_k$  are only piecewise smooth, but we can make the metric smooth and continuous by small local perturbations, without changing the metric up to coarse Lipschitz equivalence.

## 7.10 Model for the Scott core.

Choose  $W \subset M_c$  as in 7.6. We can choose  $W = M_c$ , but other choices might be more useful. An model for  $W$  has already been given in the case when  $W$  is a relative compression body 7.9. As we shall see, the general case is similar, with the model being made by gluing together models for interval bundles. First we need to review standard methods for decomposing  $W$  into simpler pieces. We are assuming that  $M_c$  has nonempty nonspherical boundary, and so the same is true of  $W$ . Therefore it is Haken, and can be decomposed into simpler pieces. A decomposition is given in [23] in Chapter 13, with most of the basic work done in Chapter 6. Here is an adaptation which gives some extra properties relating to the annuli and tori in  $\partial W \cap \partial M_d$ . For the moment, we write  $A$  for the disjoint union of annuli and tori in  $\partial W \cap \partial M_d$ .

**Lemma** *Let  $W$  be a compact connected oriented aspherical 3-manifold in which any torus is parallel to the boundary, and with nonempty boundary. Let  $A$  be a disjoint union of essential annuli in  $\partial W$ , which are all homotopically distinct. Then  $W$  is obtained from a finite sequence  $W_i$ ,  $0 \leq i \leq n$  such that  $W_n = W$ ,  $W_0$  is a disjoint union of balls and interval bundles. Any interval bundle component  $V$  is homeomorphic to  $S \times [0, 1]$  in such a way that  $\partial S \times [0, 1]$  is homeomorphic to a subset of  $A \cap \partial V$  and  $S \times \{0\}$  identifies with an incompressible subsurface of  $W$ . There is a surjective map  $j_i : W_i \rightarrow W_{i+1}$  which is injective restricted to the interior of  $W_i$ , at most two-to one, and pairs up disjoint subsurfaces of  $\partial W_i$*

to form  $W_{i+1}$ . Each component of  $\partial W_i$  contains at least one maximal connected subsurface which is mapped by  $j_i$  to an incompressible surface  $S_1$  in a component  $V$  of  $W_{i+1}$ .

The following properties will hold for  $S_1$ . From now on, by abuse of notation, we identify components of  $W_i$  with their images in  $W_j$ , for any  $j > i$ .

1.  $\partial S_1 \neq \emptyset$ .
2.  $S_1$  is incompressible in  $V$ : a simple loop which is homotopically nontrivial in  $S_1$  does not bound a disc in  $V$ .
3.  $S_1$  is boundary incompressible in the following sense. There is no union of a homotopically nontrivial arc in  $S_1$  between points of  $\partial S_1$ , and an arc in  $\partial V \setminus A$ , which bounds a disc in  $V$ .
4. If  $S_1$  is not a disc, and  $S_2$  is any connected union of components of  $\partial V \setminus A$  and  $A$  which are all intersected by  $\partial S_1$ , then  $S_2$  is incompressible in  $V$ .
5. If  $S_1$  is a disc, and  $i + 1 < n$ , then  $V$  is obtained from some component  $V'$  of  $W_{i+2}$  by attaching incompressible surfaces to  $\partial V'$ , none of which is a disc.
6. If  $S_1$  is a disc, and  $i + 1 < n$ , then  $S_1$  is indecomposable in the following sense. Let  $S_2$  be any component of  $\partial(\partial V \cap \partial V')$ , where  $V'$  is the component of  $W_{i+2}$  containing  $V$ . Isotope  $\gamma = \partial S_1$  to have only essential intersections with  $\partial S_2$ . Then up to homotopy preserving intersections with  $\partial S_2$ ,  $\gamma \neq \gamma_1 * \zeta * \gamma_2 * \bar{\zeta}$ , where  $\gamma_1$  and  $\gamma_2$  are both nontrivial in  $\partial V$  but trivial in  $V$ .

*Proof.* The distinguished set of annuli in  $\partial W$  plays a role. Rules 1 to 3 are straightforward, and probably standard. The other rules are formulated in such a way that we can bound geometry in section 8, and rule 3 will also be used in section 8. Rule 5 says we should use a compressing disc whenever it is possible to do so, in keeping with Rule 4. Rule 5 says we should put in all such discs at once. Rule 6 is a decomposability condition: as indeed is the more standard Rule 3.

Suppose that  $W_j$  has been constructed with the required properties for  $j \geq i + 1$  (by abuse of notation we are assuming  $n$  has already been determined) and let  $V$  be a component of  $W_{i+1}$ .

We start by taking a maximal set of compressing discs attached to each component of  $\partial V$ , subject to Rule 4. We can do this satisfying Rule 6 if  $i + 1 < n$ , since each disc is a sum of indecomposable ones. So now we assume that there are no compressing discs which can be attached to  $\partial V$ .

Then we apply 6.8 and 6.5 of [23]. Rules 2, 4, and 5 is automatically satisfied. We can decompose to satisfy Rule 3 if necessary.

## 7.11 Interval bundles in the Scott core.

In the case we are interested in, when  $W$  is homotopy equivalent to a hyperbolic 3-manifold  $N$ ,  $W$  contains no essential tori except possibly parallel to the

boundary, if  $A$  has toroidal components. The decomposition then gives more information, as the following lemma shows.

**Lemma** *Continue with the notation of 7.10. Suppose that  $W$  has no essential tori, except possibly parallel to components of  $\partial W$  which are in  $\partial M_d$ . Let  $\Gamma_0$  be a union of multicurves from the boundaries of the compressing discs of sets  $W_i \setminus W$  on compressible components of  $\partial W \setminus A$ . Let  $\Gamma'_0$  be the union of a maximal multicurves on these boundary components which are noncollapsing, on these compressible boundary components, such that  $\Gamma_0 \cup \Gamma'_0$  cuts each such component of  $\partial W \setminus A$  into discs with at most one puncture and annuli parallel to the boundary. Let*

$$\Sigma = \Gamma'_0 \cup A \cup \bigcup_{1 \leq j \leq n} \partial(W_j \cap \partial W_{j-1}).$$

*Define  $\Sigma_0 = \Sigma$ . Inductively, define  $\Sigma_{i+1}$  to be the union, over all interval bundle components  $V$  of  $W_0$ , and homeomorphism to  $S \times [0, 1]$ , of annuli and rectangles with alternate sides in  $A$  which are images under the homeomorphism of arcs and loops  $\alpha \times [0, 1]$  where the image of  $\alpha \times \{j\}$  is already such an arc or loop in  $\Sigma_i$ , for some  $j = 0, 1$ , with  $\partial \Sigma_i$  replaced by  $\Sigma$  if  $i = 0$ . Then for some  $r$ , each component  $V$  of  $W_0 \setminus \Sigma_r$  is either a ball, or homeomorphic to an interval bundle  $S_d \times [0, 1]$ , where the sets homeomorphic  $S_d \times \{0, 1\}$  are in incompressible components of  $\partial W \setminus A$ . In the case when  $V$  is a ball  $\partial V \setminus \Sigma_r$  is a union of topological discs and annuli parallel to the boundary.*

In future, we shall write  $\Sigma'$  for the set  $\Sigma_r$ .

*Proof.* Inductively, the component of  $W_0 \setminus \Sigma_i$  have the same properties as those listed for components of  $W_0$ . Thus, each component  $V$  is either a ball or homeomorphic to  $S \times [0, 1]$  under a homeomorphism sending  $\partial V \cap A$  to a subset of  $\partial S \times [0, 1]$ . There is a bounded  $r$  for which nontrivial interval bundle components of  $W \setminus \Sigma_r$  do not decompose. Write  $\Sigma_r = \Sigma'$ . The only way for a component  $V$  of  $W \setminus \Sigma'$  to be such an interval bundle is if it is part of a stack of such interval bundles, such that each such interval bundle  $(V, \partial V \cap A)$  is homeomorphic to  $(S \times [0, 1], \partial S \times [0, 1])$  for some surface  $S$ . The surface  $S$  must be the same throughout the stack, and the boundary component homeomorphic to  $S \times \{1\}$  on one matches up with the boundary component homeomorphic to  $S \times \{0\}$  on the next. Since there are only finitely many components, either the stack closes up to give a  $S^1$ -bundle over  $S$ , or there are two end bundles with boundary in  $\partial W \setminus A$ . The first possibility can only occur if  $S$  has boundary which is not in  $A$ , since  $W$  is connected and has boundary which is not in  $A$ . But then we get a contradiction to  $W$  having no essential tori, except parallel to the boundary. So only the second possibility occurs. In this case there is just one manifold on the stack and the components homeomorphic to  $S \times [0, 1]$  are in  $\partial W \setminus A$ .

The statement about  $\partial \Sigma'$  cutting  $\partial V$  into cells follows, because if not, there is a boundary of a compressing disc in the boundary of some  $W_i$  containing  $V$ , which is disjoint from  $\Sigma_r$ . This is impossible. For all  $i$ , the construction is such that all compressing discs automatically intersect at least one loop in  $\Sigma$  transversally.

□

### 7.12 The model for the non-interval-bundle part of the $W$ .

Lemmas 7.10 and 7.11 give a decomposition of the core  $W$  into a union  $W''$  of interval bundles, and the complement  $W'$ . The common boundary is a union of embedded annuli between components of  $\partial W \setminus A$ . These common annuli are taken to be metrically  $S^1 \times [0, 1]$ , up to bounded distortion. Otherwise, the metrics are constructed completely separately. Here, we treat the noninterval bundle piece  $W'$ . The construction is inductive. We take a metric such that  $\Sigma'$  of 7.11 has bounded length. We transfer this metric to each component of  $\partial V$ , for each component  $V$  of  $W_i$ , for each  $i$ . Because of the cell cutting, this metric is unique up to bounded distortion, and does exist. This means that we have assigned an element  $y = y(S) = [\varphi] \in \mathcal{T}(S)$  to each component  $S$  of  $\partial V$ , for each component  $V$  of  $W_i$ , up to bounded distance. We have a model manifold  $M(y)$  homeomorphic to  $S \times [0, 1]$ , simply by taking  $\varphi_*\sigma + dt^2$  where  $\sigma$  is the Poincaré metric on  $\varphi(S)$ . For the moment, we call this metric  $\sigma_{y(S)}$ . So now we need to obtain a metric on  $W'$  from the metrics on each  $\partial V$ , by gluing together the spaces  $M(y)$ . The idea is exactly the same as in 7.9 — which was one reason for doing that first. Perturb the  $W_i$  so that the components of  $W_{i-1}$  inside  $W_i$  have boundaries in the interior of  $W_i$ . Attach incompressible surfaces to  $\partial W_i$  in  $W_i$ , cutting  $W_i$  up into the component pieces and disjoint from the perturbed components of  $W_{i-1}$  inside. The region between  $V$ , and the components of  $W_{i-1}$  inside, minus the cutting surfaces, is a union of open interval bundles. Each corresponding closed interval bundle is homeomorphic to  $S \times [0, 1]$ , where  $S$  is the corresponding component of  $\partial V'$ , for a component  $V'$  of  $W_{i-1}$  inside  $V$ . We can then map  $S \times [0, 1]$  to this region by a map  $\Phi$  which is a diffeomorphism between the interiors, and take the metric  $\Phi_*(\sigma_{y(S)})$  on the region. This metric has discontinuities at the boundary, but is the right metric up to bounded distortion. We can make it continuous and smooth by perturbing at the boundary. If some of the compressing surfaces are discs, we can make the metric on the surfaces inside by using the conformal structure on the surface outside, as explained in 7.9.

This model is made up of finitely many bounded interval bundles. It might be of interest to vary the topological type, and, instead of gluing together models  $M(y)$ , use models  $M(y_1, y_2)$  for  $[y_1, y_2]$  possibly a long geodesic segment in  $\mathcal{T}(S)$ . It seems likely that the whole theory developed here could carry over to produce geometrical models for hyperbolic manifolds, and locally uniform biLipschitz constants, in this setting, under suitable conditions.

### 7.13 The model manifolds for compressible ends and ends without incompressible interval bundle bridges.

Curiously, a model manifold is easier to construct for a *compressible* end  $e$  of  $M_{d,W}$  than for an incompressible one. In fact, we have touched on the construction in 7.9. Let  $S_d(e)$  be the corresponding component of  $W \setminus \partial M_d$ , the horodisc deletion of  $S(e)$ . Let  $\Gamma_0(e)$  be the multicurve of boundaries of compressing discs in components of  $\partial(S_d(e) \cap \partial W_{n-1})$ . As in 7.9, we simply

choose  $z_{e,0} = [\varphi_{e,0}]$  so that  $|\varphi_{e,0}(\Gamma_0)|$  is bounded, but this is also the same, up to bounded distortion, as requiring that the arcs of  $\Sigma'$  on  $S_d(e)$  (as in 7.11) have bounded length. If the geometrically finite ending invariant for this component of  $M_{d,W} \setminus W$  is  $y(e)$  then the model for this end is  $M(z_{e,0}, y(e))$ .

Now suppose that  $e$  is an incompressible end, and is *not* the end of an interval bundle bridge. Then again, the point  $z_{e,0} \in \mathcal{T}(S)$  is determined by making all arcs in  $\Sigma'$  on  $S$  of bounded length, and we again take the model to be  $M(z_{e,0}, y(e))$ , if the associated end invariant is  $(y(e))$ .

## 7.14 The model for the interval bundle part of $W$ and associated ends

The interval bundle part of the Scott core is a disjoint union of interval bundles each of which forms a bridge between two components of  $\partial W \setminus \partial M_d$ . The model manifold on these interval bundles is dependent on the ending invariants of the associated ends, at least in the case of *incompressible* ends. Also, the model manifolds for the associated ends are interdependent.

We first deal with interval bundles between  $S_1$  and  $S_2$  where at least one of  $S_{d,1}$ ,  $S_{d,2}$  is *compressible*. Suppose that  $S_{d,1}$  is compressible. Suppose that the surface that is identified is  $\alpha$ . Then  $\alpha$  itself must be incompressible. The corresponding model manifold is just the portion  $\alpha \times [0, 1]$  of  $M(z_0)$ , with the model metric, where  $z_0$  is the chosen basepoint on  $S_1$ , as in 7.13. If we choose to use the chosen basepoint on  $S_2$ , it does not matter. The metric is the same up to bounded distortion, depending only on the topological type. The choice of basepoints depends only on the topological type, not on the end invariants.

Now we deal with models for other bridging interval bundles, and the end model manifolds that they bridge between. As in 7.13, we only need to choose a basepoint in  $\mathcal{T}(S(e))$  for each end. We can then define the model manifolds for the ends as in 7.13, and the model for the bridging manifold as above. For each incompressible end  $e'$  let  $\omega(e, e')$  be the maximal subsurface of  $S(e)$  which is homotopic to a subsurface of  $S(e')$ . By this, we homotopic in  $N$  and not in  $S(e)$ , in the case  $e = e'$ . We write  $\omega(e', e)$  for this surface on  $S(e')$ . Let

$$\beta = \beta(e) = S(e) \setminus \cup_{e'} \omega(e, e').$$

We start off with an initial choice  $z_{e,0} = [\varphi_{e,0}] \in \mathcal{T}(S(e))$ . We choose  $\pi_\beta(z_{e,0})$  so that  $|\varphi_{e,0}(\Sigma')|$ , for  $\Sigma'$  as in 7.11, is bounded, as before. Outside of  $\beta$ , the choice of  $z_{e,0}$  is arbitrary for the moment. We also need to fix a geometrically finite ending invariant  $(y(e))$ . We let  $y_{e,+}$  be the element of  $\mathcal{T}(S(e))$  defined using the  $(y(e))$ , in the same way as  $y_-$  in 7.6. We recall that the model for the end  $e$  is defined by removing some pieces of Margulis tube and replacing them by pieces of horoball.

Now we need to define  $z'_{e,0} \in \mathcal{T}(S(e))$ . Fix sufficiently strong ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$  and a vertically efficient ltd-bounded decomposition of  $S(e) \times [z_{e,0}, y_{e,+}]$ . For each incompressible end  $e'$  for which  $\omega(e, e') \neq \emptyset$ , we define  $E_{\text{ltd}}(e, e', -)$  to be set of all ltd  $(\alpha, \ell)$  in the decomposition of  $S(e) \times$

$[z_{e,0}, y_{e,+}]$  with  $\alpha \subset \omega(e, e')$ . Then since  $|\varphi_{e,0}(\partial\omega(e, e'))|$  is bounded in terms of topological type, by 6.2, for sufficiently strong ltd parameter functions, if  $(\alpha', \ell') < (\alpha, \ell)$ ,  $\alpha \subset \omega(e, e')$ , and  $(\alpha', \ell')$  is ltd, we also have  $\alpha' \subset \omega(e, e')$ . It follows that if  $E(e, e', -)$  is the set of all  $(\alpha', \ell')$  with  $(\alpha', \ell') < (\alpha, \ell)$  for some  $(\alpha, \ell) \in E_{\text{ltd}}(e, e', -)$  and  $E(e, e', +)$  is all the other  $(\alpha, \ell)$ , then  $E(e, e') = (E(e, e', -), E(e, e', +))$  is an order splitting of the ltd decomposition of  $S(e) \times [z_{e,0}, y_{e,+}]$  in the sense of 6.10. Then we define

$$z(e, e') = x(E(e, e')).$$

We remark that these conditions are consistent in the case  $e = e'$ .

We define

$$\pi_\beta(z'_{e,0}) = \pi_\beta(z_{e,0}).$$

Then, for  $\omega = \omega(e, e')$ , define

$$\pi_\omega(z'_{e,0}) = x(\pi_\omega(z_{e,0}), [\pi_\omega(z(e, e')), \pi_\omega(z(e', e))]).$$

Here,  $x(\cdot, \cdot)$  is as in 6.11. This definition is such that  $d_{\omega(e, e')}(z'_{e,0}, z'_{e',0})$  is bounded in terms of

$(\Delta_1, r_1, s_1, K_1)$ . Also, a ltd decomposition for  $S(e) \times [z'_{e,0}, y_{e,+}]$  is given by a subset of the decomposition for  $S(e) \times [z_{e,0}, y_{e,+}]$ .

Then our model for the end  $e$  is obtained by altering  $M(z'_{e,0}, y_{e,+})$  just as in 7.6. The model for the bridge interval bundle between the ends  $e$  and  $e'$  is obtained from  $M(\pi_\omega(z_{e,0}), \pi_\omega(z_{e',0}))$  by removing a model horoball along the boundary components of  $\partial\omega$ . That is, if  $[z'_{e,0}, y_{e,+}] = \{y_t : t \in \{0, u\}\}$ , we remove  $\cup_t \varphi_t^{-1}(H(\partial\omega, \varepsilon_0))$ , where  $H(\partial\omega, \varepsilon_0)$  is the union of components of  $(S_t)_{<\varepsilon_0}$  homotopic to  $\varphi_t(\partial\omega)$ . In general, we need to glue in a model Margulis tube in the place of each one — which might or might not be bounded. A bounded strip on the boundary comes from the non-interval bundle part of the model. We determine which Margulis tube to use from the geometry on the boundary, as usual.

From now on we redefine  $z_{e,0}$  so that  $[\pi_\omega(z'_{e',0}), \pi_\omega(z_{e',0})] \cup [\pi_\omega(z_{e,0}), \pi_\omega(z'_{e,0})]$  is a bounded distance, coordinatewise, from  $[\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$ , whenever  $e$  and  $e'$  are incompressible ends with  $\omega = \omega(e, e') \neq \emptyset$ . To do this, we leave  $\pi_\beta(z_{e,0})$  as before. But for each pair  $(e, e')$ , we choose  $x(e, e') = x(e', e) \in [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$ , and define  $z_{e,0}$  by

$$\pi_\omega(z_{e,0}) = \pi_\omega(z_{e',0}) = \pi_\omega(x(e, e')),$$

$$\pi_{\partial\omega}(z_{e,0}) = \pi_{\partial\omega}(z_{e',0}) = \pi_{\partial\omega}(x(e, e')),$$

whenever  $\omega = \omega(e, e') = \omega(e', e)$ .

## 7.15 Model for the whole manifold.

The model for a hyperbolic manifold with core  $N_c$  and geometrically finite invariants  $y(e)$  for each end is obtained by gluing together models for  $N_c$  and

models  $M(z_{e,0}, y(e))$  for each end. The boundaries are the same up to bounded distortion and we can remove discontinuities by perturbation, without changing the metric up to coarse Lipschitz equivalence. The model for the core, as we have seen, is formed by gluing together interval bundles. Rather than using a model for  $N_c$  and each of the ends, it is sometimes more convenient to take a model for some  $W \subset N_c$  and glue this together with models for each end of  $N_{d,W}$ . This is the case, for example, when  $N$  is homeomorphic to  $S \times \mathbb{R}$ , but has more than two ends.

## 7.16 The combinatorially bounded geometry geometrically infinite Kleinian surface case.

Now we consider the combinatorially bounded geometry case of  $(\mu_-, \mu_+) \in \mathcal{T}(S) \cup \mathcal{GL}_a(S)$  and convergence of the Kleinian surface models  $M(y_{n,-}, y_{n,+})$  to  $M(\mu_-, \mu_+)$  as  $y_{n,-} \rightarrow \mu_-$ ,  $y_{n,+} \rightarrow \mu_+$ . From now on in this subsection, we assume that  $\mu_+ \in \mathcal{GL}_a(S)$  and  $\mu_- \in \mathcal{GL}_a(S) \cup \mathcal{T}(S)$ .

Let  $i$  be the intersection number of 3.8. We consider the condition that, for at least one normalised transverse invariant measure on  $\mu_+$ , for some  $c = c(\mu_+) > 0$ ,

$$i(\mu_+, \gamma) \geq c(\mu_+) |\gamma|^{-1} \quad (7.16.1)$$

for all simple closed loops  $\gamma$ , and similarly for  $\mu_-$ . Here,  $|\gamma|$  is measured with respect to a fixed hyperbolic metric on  $S$ , as is used in 3.8. It can be shown that this condition implies that  $\mu_+$  has only one transverse invariant measure up to scalar, and similarly for  $\mu_-$ . It probably helps to regard  $\mu_+$  as a measured foliation with transverse invariant measure equivalent to Lebesgue measure (which it is, up to measure isomorphism). The proof of the fundamental dynamical lemma 6.1 shows that the combinatorially bounded geometry condition implies that every leaf of length  $L$  comes within distance  $c_1/L$  of every point in the measured foliation, for every  $L > 0$ , for  $c_1$  depending only on  $c(\mu_+)$ . It follows that given any invariant set  $E$  under the foliation, every point is a positive density point for  $E$ , and therefore  $E$  has full measure and the transverse invariant measure is ergodic. But if every transverse invariant measure is ergodic, there is only one, up to scalar.

Fix a basepoint  $y_0 = [\varphi_0] \in \mathcal{T}(S)$ . We claim that if  $y_{n,+} = [\varphi_{n,+}] \rightarrow \mu_+$  and, for all  $n$

$$[y_0, y_{n,+}] \subset (\mathcal{T}(S))_{\geq \nu}, \quad (7.16.2)$$

then (7.16.1) holds for a  $c(\mu_+)$  depending only on  $\nu$  and  $y_0$ . For let  $\zeta_{n,+}$  be a loop such that

$$C_1^{-1} |\varphi_0(\zeta_{n,+})| \leq \exp d(y_0, y_{n,+}) \leq C_1 |\varphi_0(\zeta_{n,+})| \quad (7.16.3)$$

and

$$|\varphi_{n,+}(\zeta_{n,+})| \leq C_1. \quad (7.16.4)$$



This is possible for a suitable  $C_1$  by 2.5, and the fact that  $y_{n,+} \in (\mathcal{T}(\mathcal{S}))_{\geq \nu}$ . Enlarging  $C_1$  we then also have, for all  $y = [\varphi] \in [y_0, y_{n,+}]$ ,

$$C_1 |\varphi(\zeta_{n,+})| \geq \exp d(y_{n,+}, y).$$

Then any limit of  $\zeta_{n,+}/|\zeta_{n,+}|$  (taking  $|\cdot|$  with respect to a fixed hyperbolic metric on  $S$ ) has zero intersection with  $\mu_+$ , and must be  $\mu_+$  by arationality. For any simple closed loop  $\gamma$ , choose  $y = [\varphi] \in [y_0, y_{n,+}]$  such that  $|\varphi(\gamma)|$  is minimal over all such  $y$ . Then  $|\varphi(\gamma)|$  is bounded from 0, and the good position of  $\varphi(\gamma)$  is such that along most of its length it is bounded from the stable and unstable foliations of the quadratic differential for  $d(y, y_{n,+})$ . So for suitable  $C_1$ ,

$$C_1 |\varphi_0(\gamma)| \geq \exp d(y_0, y).$$

Using this and 6.2, since  $S$  is ltd along  $[y_0, y_{n,+}]$ ,

$$\begin{aligned} i(\gamma, \zeta_{n,+}) &= \#(\varphi(\gamma) \cap \varphi(\zeta_{n,+})) \geq C_2^{-1} |\varphi(\zeta_{n,+})| \geq (C_1 C_2)^{-1} \exp d(y, y_{n,+}) \\ &\geq C_3 \exp(d(y_0, y_{n,+}) - d(y_0, y)) \geq C_4 \frac{|\zeta_{n,+}|}{|\gamma|}. \end{aligned}$$

So (7.16.2) implies (7.16.1).

We claim that the converse is also essentially true. So suppose that (7.16.1) holds and that  $y_{n,+} \rightarrow \mu_+$ . Choose  $\zeta_{n,+}$  so that (7.16.3) is replaced by

$$C_1^{-1} \frac{|\varphi_0(\zeta_{n,+})|}{|\varphi_{n,+}(\zeta_{n,+})|'} \leq \exp d(y_0, y_{n,+}) \leq C_1 \frac{|\varphi_0(\zeta_{n,+})|}{|\varphi_{n,+}(\zeta_{n,+})|}, \quad (7.16.5)$$

and that (7.16.4) is replaced by

$$|\varphi_{n,+}(\zeta_{n,+})|'' \leq C_1. \quad (7.16.6)$$

Here,  $|\cdot|'$  and  $|\cdot|''$  are as in 2.5. This time we do not have a lower bound on  $|\varphi_{n,+}(\zeta_{n,+})|$ . But choose  $y'_{n,+} = [\varphi'_{n,+}] \in [y_0, y_{n,+}]$  such that (7.16.6) holds for  $\varphi'_{n,+}$  replacing  $\varphi_{n,+}$ , and also

$$|\varphi'_{n,+}(\zeta_{n,+})| \geq C_1^{-1}. \quad (7.16.7)$$

Then we claim that

$$[y_0, y'_{n,+}] \subset (\mathcal{T}(S))_{\geq \nu}. \quad (7.16.8)$$

Suppose this is not true and that there is a loop  $\gamma$  and  $y = [\varphi] \in [y_0, y'_{n,+}]$  with  $|\varphi(\gamma)| \leq \nu$ . We take  $[w_1, w_2] = [[\psi_1], [\psi_2]] \subset [y_0, y'_{n,+}]$  such that

$$|\psi_j(\gamma)| \leq C_1,$$

and

$$d(w_1, w_2) \geq L(\nu),$$

where  $L(\nu) \rightarrow \infty$  as  $\nu \rightarrow 0$ . In fact, we can take  $L(\nu) \geq C_1^{-1} \log(1/\nu)$ . Then, using 2.5.2,

$$\begin{aligned} i(\gamma, \zeta_{n,+}) &\leq C_2 \exp d(w_2, y'_{n,+}) \leq C_2 \exp -L(\nu) \exp(d(y_0, y'_{n,+}) - d(w_1, y_0)) \\ &\leq C_3 \exp -L(\nu) \frac{|\varphi_0(\zeta_{n,+})|}{|\varphi_0(\gamma)|}. \end{aligned}$$

This contradicts (7.16.1)

Similar arguments work for  $\mu_-$  and sequences  $y_{n,-}$ ,  $y'_{n,-}$ , if either  $\mu_- \in \mathcal{GL}_a(S)$  satisfies (7.16.1) or (7.16.2) holds for the  $y_{n,-}$ . So now if we assume that either  $\mu_- \in \mathcal{GL}_a(S)$  and satisfies (7.16.1) and  $y_{n,-}$ ,  $\zeta_{n,-}$ ,  $y_{n,-}$  are defined similarly to  $y_{n,+}$ ,  $\zeta_{n,+}$ ,  $y'_{n,+}$ , or  $\mu_- \in \mathcal{T}(S)$  and  $y_{n,-} = y'_{n,-} = \mu_-$  for all  $n$ . Then for a suitable  $\nu > 0$ ,

$$[y_0, y'_{n,-}] \in (\mathcal{T}(S))_{\geq \nu},$$

and hence applying 6.4 to the triangle with vertices at  $y_0$ ,  $y'_{n,+}$ ,  $y'_{n,-}$ , for suitable  $\nu$ , for all  $n$ ,

$$[y'_{n,-}, y'_{n,+}] \subset (\mathcal{T}(S))_{\geq \nu}.$$

Now we claim that if  $\mu_+$ ,  $\mu_- \in \mathcal{GL}_a(S)$  satisfy (7.16.1) and  $\mu_+ \neq \mu_-$  then for a constant  $C = C(\mu_+, \mu_-)$ , for all sufficiently large  $n$ ,

$$d(y'_{n,-}, y_0) + d(y_0, y'_{n,+}) \leq d(y'_{n,-}, y'_{n,+}) + C. \quad (7.16.9)$$

The reason is simply that  $\mu_+ \neq \mu_-$  means  $i(\mu_+, \mu_-) > 0$  by arationality, and hence for all sufficiently large  $n$  and constants  $C_j$  depending on  $\mu_+$ ,  $\mu_-$ ,

$$\begin{aligned} \exp d(y'_{n,-}, y'_{n,+}) &\geq C_1^{-1} i(\zeta_{n,+}, \zeta_{n,-}) \geq C_2^{-1} |\zeta_{n,+}| \cdot |\zeta_{n,-}| \\ &\geq C_3^{-1} |\varphi_0(\zeta_{n,+})| \cdot |\varphi_0(\zeta_{n,-})| \geq C_4^{-1} \exp(d(y'_{n,-}, y_0) + d(y_0, y'_{n,+})). \end{aligned}$$

It follows from 6.4 and the fact that the geodesic segments joining  $y_0$ ,  $y'_{n,+}$ ,  $y'_{n,-}$  are in  $(\mathcal{T}(S))_{\geq \nu}$  that, for a constant  $C'$  depending only on  $\mu_{\pm}$ , for all sufficiently large  $n$ ,

$$d(y_0, x(y_0)) \leq C', \quad (7.16.10)$$

where  $x(y_0)$  denotes the orthogonal projection (6.9) to  $[y'_{n,-}, y'_{n,+}]$  for any  $n$ , that is,  $y_0$  is distance  $\leq C'$  from some point on  $[y'_{n,-}, y'_{n,+}]$ . It then follows from 6.4 that each point on  $[y'_{n,-}, y_0] \cup [y_0, y'_{n,+}]$  is distance  $\leq C''$  from some point on  $[y'_{n,-}, y'_{n,+}]$ , for  $C''$  independent of  $n$ . So then  $y_0$  is a distance  $\leq C'$  from some point on  $[y_{n,-}, y_{n,+}]$ , for all sufficiently large  $n$ , and any  $y' \in [y'_{n,-}, y'_{n,+}]$  is distance  $\leq C''$  from some point in  $[y_{n,-}, y_{n,+}]$ .

This finally puts us in a position to prove geometric convergence, up to bounded coarse Lipschitz equivalence, of suitably based models  $(M(y_{n,-}, y_{n,+}), x_n)$ ,  $(M(y'_{n,-}, y'_{n,+}), x'_n)$ , if  $y_{n,+} \rightarrow \mu_+$  where  $\mu_+$  satisfies (7.16.1) and either similar properties hold for  $\mu_-$ , or  $y_{n,-} = \mu_- \in \mathcal{T}(S)$  for all  $n$ . In all cases we have  $y_{n,0} \in [y_{n,-}, y_{n,+}]$  and  $y'_{n,0} \in [y'_{n,-}, y'_{n,+}]$  with

$$d(y_0, y_{n,0}) \leq C', \quad d(y_0, y'_{n,0}) \leq C'.$$

We translate the vertical coordinate so that

$$\begin{aligned} M(y_{n,-}, y_{n,+}) &= S \times [-u_{n,-}, u_{n,+}], \quad M(y'_{n,-}, y'_{n,+}) = S \times [-u'_{n,-}, u'_{n,+}], \\ d(y_{n,-}, y_{n,0}) &= u_{n,-}, \quad d(y_{n,+}, y_{n,0}) = u_{n,+}, \\ d(y'_{n,-}, y'_{n,0}) &= u'_{n,-}, \quad d(y'_{n,+}, y'_{n,0}) = u'_{n,+}. \end{aligned}$$

We take  $x_0$  to be any fixed point in  $S \times \{0\}$ ,  $x_0 = x_n = x'_n$  for all  $n$ .

Then to prove geometric convergence of the models up to bounded coarse Lipschitz equivalence, it suffices to prove that, for a constant  $C_0$ , for any  $\Delta > 0$ , for all sufficiently large  $k$  and  $n$ , if  $y \in [y_{k,-}, y_{k,+}]$  and  $d(y_0, y) \leq \Delta$ , there is  $y' \in [y_{n,-}, y_{n,+}]$  with

$$d(y, y') \leq C_0, \quad (7.16.11)$$

and similarly for  $y_{k,\pm}$ ,  $y_{n,\pm}$  replaced by  $y'_{k,\pm}$ ,  $y'_{n,\pm}$ . For suppose we have this. Then replacing  $y_{k,\pm}$  by suitable points in the original geodesic segment  $[y_{k,-}, y_{k,+}]$  if necessary, and replacing  $n$  by a subsequence if necessary, we can assume that (7.16.1) holds for all  $y \in [y_{k,-}, y_{k,+}]$ , and for all  $n \geq k$ . We can also assume that  $d(y_{k,+}, y_{n,+})$  and  $d(y_{k,-}, y_{n,-})$  is bounded from 0 for all  $n > k$ . Then we can construct a boundedly coarse biLipschitz map  $\varphi_{k,n}$ , from  $M(y_{k,-}, y_{k,+})$  to a subset of  $M(y_{n,-}, y_{n,+})$ , fixing  $x_0$ , for any  $k > n$ , and such that the distance, in the Riemannian metrics, between  $\varphi_{k,n}(\partial M(y_{k,-}, y_{k,+}))$  and  $V_{k,n} = \partial M(y_{n,-}, y_{n,+}) \setminus \varphi_{k,n}(\partial M(y_{k,-}, y_{k,+}))$ , is bounded from 0. Then we can make a Riemannian manifold  $V_p$  by gluing together  $M(y_{1,-}, y_{1,+})$  and  $V_{n,n+1}$  for  $1 \leq n < p$ , taking a metric which is the model metric for  $M(y_{1,-}, y_{1,+})$  on  $M(y_{1,-}, y_{1,+})$  and the model metric for  $M(y_{n+1,-}, y_{n+1,+})$  on  $V_{n,n+1}$ , except near  $\partial \varphi_{n,n+1}(M(y_{n,-}, y_{n,+}))$ . on all of  $V_{n,n+1}$ , the metric is taken boundedly equivalent to the metric on  $M(y_{n+1,-}, y_{n+1,+})$ . The manifold  $(V_p, x_0)$  has a based submanifold which is naturally diffeomorphic to  $(M(y_{n,-}, y_{n,+}), x_0)$ , for each  $n \leq p$ , with bounds on the derivative and derivative inverse with respect to the Riemannian metrics, and the geometric limit  $\lim_{p \rightarrow \infty} (V_p, x_0)$  exists as a based Riemannian manifold. Each set  $V_{p+1} \setminus V_p = V_{p,p+1}$  is homeomorphic to  $S \times I$  where  $I$  is the union of one or two intervals, depending on whether  $\mu_- \in \mathcal{T}(S)$  or  $\mu_- \in \mathcal{GL}_a(S)$ . It follows that  $\lim_{p \rightarrow \infty} V_p$  is homeomorphic to  $S \times [0, \infty)$  or  $S \times \mathbb{R}$ . The geometric limit  $\lim_{p \rightarrow \infty} V_p$  depends only on  $\mu_{\pm}$  up to coarse biLipschitz equivalence, not on the precise sequences  $y_{n,\pm}$ , by (7.16.11), and not on the precise choice of metric near  $\partial V_p$  for any  $p$ .

It suffices to prove (7.16.11) for  $y'_{k,\pm}$  and  $y'_{n,\pm}$ , since every point in  $[y'_{k,-}, y'_{k,+}]$  is a bounded distance from a point in  $[y_{k,-}, y_{k,+}]$ , and  $d(y_0, y'_{k,+}) \rightarrow \infty$  as  $k \rightarrow \infty$  (because  $|\zeta_{k,+}| \rightarrow \infty$ ), and similarly for  $y'_{k,-}$  if  $\mu_- \in \mathcal{GL}_a(S)$ . So now let  $y \in [y'_{k,-}, y'_{k,+}]$  with  $d(y_0, y) \leq \Delta$ . Suppose that  $y$  is not a bounded distance  $\leq C_0$  from a point in  $[y'_{n,-}, y'_{n,+}]$ . Then by 6.4 for quadrilaterals, and  $(\mathcal{T}(S))_{\geq \nu}$ , taking vertices  $y'_{n,\pm}$ ,  $y'_{k,\pm}$ , for suitable  $C_0$  (given by 6.4),  $y$  must be a distance  $\leq C_0$  from  $y' \in [y'_{n,+}, y'_{k,+}] \cup [y'_{k,-}, y'_{n,-}]$ . We can assume without loss of generality that  $y' = [\varphi'] \in [y'_{n,+}, y'_{k,+}]$ . If  $n$  and  $k$  are both large given  $\Delta$  then  $d(y'_{k,+}, y')$  and  $d(y'_{n,+}, y')$  are both large. Then by 6.2,

$$i(\zeta_{k,+}, \zeta_{n,+}) = \#(\{\varphi'(\zeta_{k,+}) \cap \varphi'(\zeta_{n,+})\}) \geq C_1^{-1} |\varphi'(\zeta_{k,+})| \cdot |\varphi'(\zeta_{n,+})|$$

$$\geq C_2^{-1} e^{-2\Delta} |\varphi_0(\zeta_{k,+})| \cdot |\varphi_0(\zeta_{n,+})| \geq C_3^{-1} e^{-2\Delta} |\zeta_{k,+}| \cdot |\zeta_{n,+}|.$$

This cannot be true for arbitrarily large  $k$  and  $n$ , because we would then deduce that  $i(\mu_+, \mu_+) > 0$ . So geometric convergence of the models is complete in this case.

## 7.17 Geometric convergence of models in the general geometrically infinite case

All geometric models in the geometrically finite case are obtained by gluing together finitely many models  $M(y_-, y_+)$ , with insignificant modifications to some model Margulis tubes in the ending models with extra cusps in the ends. For the geometrically infinite models, we simply want to take geometric limits of models of the form  $M(y_0, y_{n,+})$  for some exterior models, or a geometric limit of a single sequence of models  $M(y_{n,-}, y_{n,+})$ , for a suitable choice of basepoint and with  $y_{n,+}$  convergent to some point in  $\mu_+ \in \partial\mathcal{T}(S)$ , where  $\partial\mathcal{T}(S)$  is the modification of the Thurston boundary described in 3.9, and either the same is true for  $\mu_-$ , or  $\mu_- \in \mathcal{T}(S)$ .

So let  $y_{n,+} = [\varphi_{n,+}] \rightarrow \mu_+$  and  $y_{n,-} \in \mu_-$ . If  $\mu_{\pm}$  are both arational geodesic laminations on  $S$ , we assume that  $i(\mu_-, \mu_+) > 0$ . If  $\mu'_+$  and  $\mu'_-$  denote the lamination parts of  $\mu_{\pm}$ , we also assume that no closed loop  $\gamma \subset S$  in the closure of the support of  $\mu'_+$  or  $\mu'_-$  satisfies  $i(\gamma, \mu'_+) = i(\gamma, \mu'_-) = 0$ . The latter condition is to ensure that any geometric limit is connected (but is not actually a necessary condition for this). We need to show that, for a suitable choice of base-point  $x_n \in M(y_{n,-}, y_{n,+})$ ,  $(M(y_{n,-}, y_{n,+}), x_n)$  has a single geometric limit up to bounded coarse biLipschitz equivalence.

Fix a basepoint  $y_0 = [\varphi_0]$  as in 7.16. As in 7.16, we can find loops  $\zeta_{n,\pm}$  and  $y'_{n,\pm} = [\varphi'_{n,\pm}]$  converging to  $\mu_{\pm}$  such that (7.16.6) and (7.16.7) hold. If  $\mu_+$  has at least one lamination component, we can also assume that  $\zeta_{n,+}$  has nonempty intersection with the support of at least one minimal component of  $\mu_+$ , and similarly for  $\zeta_{n,-}$ ,  $\mu_-$ . Let  $x(y_0, [y_{n,-}, y_{n,+}])$  be the orthogonal projection of  $y_0$  relative to  $[y_{n,-}, y_{n,+}]$ , as in 6.11. We claim that, in order to show geometric convergence, up to bounded coarse Lipschitz equivalence, it suffices to show that for a suitable constant  $L_1$ , given  $\Delta > 0$ , for all sufficiently large  $k$ ,

$$d(y_0, x(y_0, [y_{k,-}, y_{k,+}])) \leq L_1, \quad (7.17.1)$$

and for all sufficiently large  $k$ , and  $n$ , and all ltd  $(\alpha, \ell)$  for  $[y_{k,-}, y_{k,+}]$  with  $d'_\alpha(y, y_0) \leq \Delta$  for  $y \in \ell$ , there is  $(\alpha, \ell')$  which is ltd along  $\ell' \subset [y_{n,-}, y_{n,+}]$ , possibly with respect to different ltd parameter functions (which is enough, by 7.7), and for all  $y \in \ell$ , there is  $y' \in \ell'$  with

$$d_\alpha(y, y') \leq L_1 \text{ or } |\operatorname{Re}(\pi_\alpha(y) - \pi_\alpha(y'))| \leq L_1, \quad (7.17.2)$$

depending on whether  $\alpha$  is a loop or a gap. We see that these suffice as follows. So suppose that both (7.17.1) and (7.17.2) hold.

We first look for suitable basepoints  $x_0 = (z_0, 0)$  in the models. This means looking for a gap or loop  $\alpha$  such that  $z_0 \in \varphi_0^{-1}(S_{\alpha,0})$ , that is,  $S_{\alpha,0}$  is homotopic to  $\varphi(\alpha)$  and bounded by  $S_{\partial\alpha,0}$  in the terminology of 7.2. We can drop the first few terms of the sequence if necessary, and assume that there is  $(\alpha, \ell)$  which is ltd along  $[y_{1,-}, y_{1,+}]$ ,  $d'_\alpha(y, y_0) \leq \Delta_1$  for all  $y \in \ell$ , for  $\Delta_1$  depending on the ltd parameter functions and  $y_{1,\pm}$ , and (7.17.2) holds for all  $n$ . Similarly, (7.17.2) also holds for  $y'_{n,0} = [\varphi'_{n,0}] \in [y'_{n,-}, y'_{n,+}]$  replacing  $y_{n,0}$ , again because the ltd's along  $[y_0, y'_{n,\pm}]$  are a subset of those along  $[y_0, y_{n,\pm}]$  up to bounded distance and hence the ltd's along  $[y'_{n,-}, y'_{n,+}]$  are a subset along  $[y_{n,-}, y_{n,+}]$ , up to bounded distance. Conversely if we have (7.17.1) and (7.17.2) with  $y'_{n,\pm}$  replacing  $y_{n,\pm}$  and  $y'_{n,0} \in [y'_{n,-}, y'_{n,+}]$  replacing  $y_{n,0}$ , then we have (7.17.1) and (7.17.2) for  $y_{n,\pm}$ . So suppose that we have all of these. Let  $\varphi_{n,0}$  and  $\varphi'_{n,0}$  be homeomorphisms which are part of families  $\varphi_{n,t}$ ,  $\varphi'_{n,t}$ , relative to  $[y_{n,-}, y_{n,+}]$  and  $[y'_{n,-}, y'_{n,+}]$ , satisfying the properties of 7.2. Translate the vertical coordinate of the model as in 7.16, defining  $u_{n,\pm}$ ,  $u'_{n,\pm}$  as there. If  $\alpha$  is a gap and long  $\nu_n$ -thick and dominant at  $y_{n,0}$  then we take  $x_n = (z_n, 0)$  so that  $\varphi_{n,0}(x_n)$  is in the  $\nu_n$ -thick part of  $\varphi_{n,0}(S)$  which is homotopic to  $\varphi'_n(\alpha)$ . Let  $x'_n = (z'_n, 0)$  be similarly defined relative to  $\varphi'_{n,0}$ . We can, and do, choose  $\varphi_n$  and  $\varphi'_n$  so that  $z_n = z'_n = z_0$ . If  $\alpha$  is a loop, we can choose it so that it is transverse to a lamination component of each of  $\mu_\pm$ . Then we claim that we have a lower bound on  $|\varphi(\alpha)|$  for  $[\varphi] \in [y'_{n,-}, y'_{n,+}]$  and for all  $n$ , and hence similarly for  $[y_{n,-}, y_{n,+}]$ . For from (7.17.1) we have, for  $x(y_0) = x(y_0, [y'_{n,-}, y'_{n,+}])$ , for a constant  $C_0$  depending only on the ltd parameter functions,

$$d(y'_{n,-}, x(y_0)) + d(x(y_0), y'_{n,+}) \leq d(y'_{n,-}, y'_{n,+}) + C_0, \quad (7.17.3)$$

which can be seen by using the bound 2.9 for  $d(y'_{n,-}, y'_{n,+})$  in terms of

$$\text{Max}_\beta(d'_\beta(x(y_0), y'_{n,-}) + d_{\beta'}(x(y_0), y'_{n,+})).$$

So then for  $\zeta_{n,\pm}$ ,  $y'_{n,\pm} = [\varphi'_{n,\pm}]$ ,  $y_{n,\pm} = [\varphi_{n,\pm}]$  satisfying (7.16.6) and (7.16.7) as above, suppose that we do not have a lower bound on  $|\varphi(\alpha)|$ . Let  $w_j = [\psi_j]$ ,  $j = 1, 2$ ,  $[w_1, w_2] \subset [y'_{n,-}, y'_{n,+}]$ ,  $|\psi_j(\alpha)| \leq C_1$ ,  $d(w_1, w_2) \geq \Delta_2$ , where  $\Delta_2$  can be taken large if some  $|\varphi(\alpha)|$  is small. We have

$$\begin{aligned} i(\alpha, \zeta_{n,-}) \cdot i(\alpha, \zeta_{n,+}) &\leq C_2^{-1} \exp((d(w_1, y'_{n,-}) + d(w_2, y'_{n,+})) \\ &\leq C_3^{-1} e^{-\Delta_2} \exp((d(y_0, y'_{n,-}) + d(y_0, y'_{n,+})) \leq C_4^{-1} e^{-\Delta_2} |\zeta_{n,-}| |\zeta_{n,+}| \end{aligned}$$

If this is true for arbitrarily large  $\Delta_2$  and hence also arbitrarily large  $n$ , then taking limits, we obtain

$$i(\alpha, \mu'_+) \cdot i(\alpha, \mu'_-) = 0,$$

or  $i(\alpha, \mu_+) = 0$  if  $y_{n,-} = y'_{n,-} = \mu_-$ . By the choice of  $\alpha$ , this is impossible.

Next, we can extend (7.17.2) to the set of all  $(\alpha, \ell)$  in a vertically efficient decomposition for  $S \times [y_{k,-}, y_{k,+}]$  with  $d'_\alpha(y, y_0) \leq \Delta$  for  $y \in \ell$ . For given such an  $(\alpha, \ell)$ , we can find a corresponding set for  $S \times [y_{n,-}, y_{n,+}]$ , if  $n$  is sufficiently large, by taking upper and lower boundaries (6.11) of sets of ltd's below and above  $\alpha \times \ell$

in  $S \times [y_{k,-}, y_{k,+}]$ , and taking the corresponding upper and lower boundaries in  $S \times [y_{n,-}, y_{n,+}]$ . We can thicken slightly so that the upper and lower boundaries are disjoint. Then the two pieces corresponding to  $(\alpha, \ell)$  in  $M(y_{k,-}, y_{k,+})$  and  $M(y_{n,-}, y_{n,+})$  are again boundedly coarse Lipschitz equivalent. Model Margulis tubes in each are determined by the metrics on their boundaries. So each model Margulis tube in  $M(y_{k,-}, y_{k,+})$  which is completely encased by pieces corresponding to ltd or bounded  $(\alpha, \ell)$ , with  $d'_\alpha(y, y_0) \leq \Delta$  for  $y \in \ell$ , is boundedly Lipschitz equivalent to a model Margulis tube in  $M(y_{n,-}, y_{n,+})$ .

Next, in analogy to what was done in 7.16, we change the sequences  $y_{n,\pm}$  so that (7.17.1) and (7.17.2) hold for all  $(\alpha, \ell)$  in the decomposition for  $[y_{k,-}, y_{k,+}]$ , and for all  $n > k$ . For a suitable  $\Delta_n$ , let  $(E_n(-, +), E_n(+, +))$  be an order splitting for  $[y_0, y_{n,+}]$  (6.11) so that  $d'_\alpha(y, y_0) \leq \Delta_n$  for all  $y \in \ell$  and  $(\alpha, \ell) \in E_n(-, +)$  and  $d'_\alpha(y, y_0) \geq \Delta_n$  for  $y \in \ell$  and  $(\alpha, \ell) \in E_n(+, +)$ . Restricting to a subsequence if necessary, we can assume that (7.17.2) holds for  $\Delta = \Delta_k$  and all  $n > k$  and similarly for the sequence  $y_{n,-}$ . Replace  $y_{n,+}$  by  $x(E_n(-, +)) = x(E_n(+, +))$  in the notation of 6.11, and similarly for  $y_{n,-}$ . Then, as in 7.16, we have a sequence of maps  $\varphi_{k,n}$  from  $(M(y_{k,-}, y_{k,+}), x_0)$  into  $(M(y_{n,-}, y_{n,+}), x_0)$  for  $k \leq n$ . We can choose  $\varphi_{k,n}$  to have uniformly bounded derivative and inverse derivative with respect to the Riemannian metrics on  $M(y_{k,-}, y_{k,+})$  and  $M(y_{n,-}, y_{n,+})$ . Then, as in 7.16, we form the sequence  $V_n$  by gluing together pieces  $M(y_{1,-}, y_{1,+})$  and

$$V_{n,n+1} = M(y_{n+1,-}, y_{n+1,+}) \setminus \varphi_{n+1,n}(M(y_{n,-}, y_{n,+})),$$

so that the Riemannian metric on  $V_{n,n+1}$ , except in a neighbourhood of  $\varphi_{n,n+1}(M(y_{n,-}, y_{n,+}))$ , is the Riemannian metric on  $M(y_{n+1,-}, y_{n+1,+})$ , and is boundedly equivalent to this metric everywhere on  $V_{n,n+1}$ .

If  $\mu_+$  is reducible, that is,  $i(\mu_+, \gamma) = 0$  for at least one nontrivial non-peripheral closed loop, then there are some model Margulis tube boundaries which intersect  $\partial V_n$  for all  $n$ . We denote by  $\partial_h V_n$  the complement in  $\partial V_n$  of any such Margulis tube boundaries. In order for the limit to be a Riemannian manifold and a topological product, we need the distance between  $\partial_h V_n$  and  $\partial_h V_{n+1} \setminus \partial_h V_n$  to be bounded from 0, replacing the original sequence for a sufficiently fast increasing subsequence if necessary. We concentrate on the boundary corresponding to  $\mu_+$ . (There is boundary corresponding to  $\mu_-$  only if  $\mu_- \notin \mathcal{T}(S)$ .) Write  $\partial_{h,+} V_n$  for the union of boundary components corresponding to  $y_{n,+}$ . Fix  $k$  so that  $\partial_{h,+} V_k$  contains components corresponding to all Teichmüller space components of  $\mu_+$ . It suffices to show that the distance in the model Riemannian metric of  $\partial_{h,+} V_n \setminus \partial_{h,+} V_k$  from  $x_0 \rightarrow \infty$  as  $n \rightarrow \infty$ , and the same for any model Margulis tubes intersecting  $\partial_{h,+} V_n \setminus \partial_{h,+} V_k$ . We see this as follows. If not, then there is  $\Delta > 0$ , and, for each  $n$ , a path from  $x_0$  to  $\partial_{h,+} V_n \setminus \partial_{h,+} V_k$  of length  $\leq \Delta$ . Restricting to a subsequence if necessary, we can assume that the path always passes through the same model Margulis tubes and the same sets  $W_j$ , in the notation of 7.2, corresponding to sets  $\alpha \times \ell$  in the vertically efficient decomposition for  $S \times [y_{n,-}, y_{n,+}]$ , for some sufficiently large  $n$ . So then the path for  $n$  can be assumed to end in  $(\gamma, u_{n,+})$  for a nontrivial

nonperipheral closed loop  $\gamma \subset S$  such that  $|\varphi_{n,+}(\gamma)| \leq \Delta'$  for all  $n$ . There is no relation between  $\Delta$  and  $\Delta'$ , but  $\Delta' < +\infty$ , because  $\Delta'$  is determined by the  $W_j$  passed through, corresponding to sets  $\alpha \times \ell$ . Then  $\gamma$  lies in a geodesic lamination component of  $\mu_+$ , but  $i(\mu_+, \gamma) = 0$ , which is a contradiction.

Now we prove (7.17.1). We consider the sets of ltds  $T(y_0, +)$  and  $T(y_0, -)$  of 6.10. If (7.17.1) does not hold for a sufficiently large  $L_1$  given  $L_2$ , then, by 5.5, there must be some maximal totally ordered set of ltds  $(\alpha_i, \ell_i)$ ,  $1 \leq i \leq m$  along  $[y_0, x(y_0)]$  with  $\ell_m$  nearest to  $x(y_0)$ , and

$$\sum_i |\ell_i| \geq L_2.$$

Let  $w_m = [\xi]$  be the end of  $\ell_m$  nearer to  $x(y_0)$ , so that  $d'_{\alpha_n}(w_m, x(y_0)) \leq L_0$ , for  $L_0$  depending only on the ltd parameter functions. Let  $\gamma \subset \alpha_m$  be a loop such that  $|\xi(\gamma)|$  is bounded. Then by 6.6, for a constant  $C_5$  bounded in terms of  $|\xi(\gamma)|$  and the ltd parameter functions,

$$i(\gamma, \zeta_{k,+}) \leq C_5 |\xi(\zeta_{k,+} \cap \alpha_m)| \leq C_5^2 e^{-L_2/2} |\varphi_0(\zeta_{n,+})|.$$

So if this is true for arbitrarily large  $k$ ,

$$i(\gamma, \mu'_+) \leq e^{-L_2/4}.$$

But similarly

$$i(\gamma, \mu'_-) \leq e^{-L_2/4}.$$

This contradicts our assumption on  $\mu'_\pm$ . So now we have (7.17.1) for all sufficiently large  $k$ .

So now we need to show (7.17.2), using a generalisation of the technique used to prove (7.16.11). So let  $k$  be sufficiently large that (7.17.1) holds, and let  $(\alpha, \ell)$  be ltd for  $[y_{k,-}, y_{k,+}]$  with  $d'_\alpha(y_0, y) \leq \Delta$  for  $y \in \ell$ . We again use 6.4, but this time the quadrilateral case, for the quadrilateral with vertices at  $y_{k,\pm}$  and  $y_{n,\pm}$ . So fix  $\Delta > 0$ , and let  $(\alpha, \ell)$  be long  $\nu$ -thick and dominant or  $K_0$ -flat (having fixed ltd parameter functions) along  $[y_{k,-}, y_{k,+}]$  within  $d'_\alpha$  distance  $\Delta$  of  $y_0$ . By assumption,  $\alpha$  does not intersect any loop  $\zeta$  with  $i(\mu_+, \zeta) = 0$ , or  $i(\mu_-, \zeta) = 0$ . Let  $y = [\varphi] \in \ell$ . Suppose that (7.17.2) does not hold for  $y' \in [y_{n,-}, y_{n,+}]$  for  $n$  sufficiently large. By 6.4, there must then be  $y' = [\varphi'] \in [y_{k,+}, y_{n,+}] \cup [y_{k,-}, y_{n,-}]$ . We assume without loss of generality that  $y' \in [y_{k,+}, y_{n,+}]$ . Precisely, we have

$$d_\alpha(y, y') \leq C(\nu) \text{ or } |\operatorname{Re}(\pi_\alpha(y) - \pi_\alpha(y'))| \leq K_0.$$

Suppose that  $\alpha$  is a gap which is long  $\nu$ -thick and dominant. Fix a loop  $\gamma \subset \alpha$  with  $|\varphi_0(\gamma)| \leq L_0$ , with  $L_0$  depending only on the topological type of  $S$ . Now by 6.2 since  $\varphi'(\zeta_{n,+} \cap \alpha)$  and  $\varphi'(\zeta_{k,+} \cap \alpha)$  are close to the stable and unstable foliations of the quadratic differential for  $d(y', y_{n,+})$  respectively, for a constant  $C_2(\nu)$ ,

$$i(\zeta_{n,+}, \zeta_{k,+}) \geq \#(\varphi(\alpha \cap \zeta_{k,+} \cap \zeta_{n,+})) \geq (C_2(\nu))^{-1} i(\zeta_{n,+}, \gamma) \cdot i(\gamma, \zeta_{k,+}).$$

If this is true for arbitrarily large  $k$ , and  $n$ , taking limits, this means that

$$0 = i(\mu'_+, \mu'_+) \geq C_2(\nu)^{-1} i(\mu'_+, \gamma)^2 > 0,$$

which is a contradiction. So now we have (7.17.2), and the proof of geometric convergence of the models is finished in this general case.

## 8 Model-adapted families of pleated surfaces.

Let  $N$  be a three-dimensional hyperbolic manifold with finitely generated fundamental group. Let  $\overline{N}$  be the union of  $N$  and the quotients by the covering group of the complement of the limit set in  $\partial H^3$ . Let  $W$  be a submanifold of the relative Scott core  $N_c$  as in 7.6, and  $N_{d,W}$  as defined there. Thus, each component of  $N_{d,W} \setminus W$  is a neighbourhood of a unique end  $e$  of  $N_{d,W}$ , and the closure is homeomorphic to  $S_d(e) \times [0, \infty)$ , where  $S_d(e)$  is the bounding component of  $\partial W \setminus \partial N_d$ . In this section, we construct a family of pleated surfaces in  $N$ , given a map  $f_{e,+} : S(e) \rightarrow \overline{N}$  homotopic to inclusion of  $S(e)$  in  $N$ , for each end  $e$  of  $N_{d,W}$ . Here,  $f_{e,+}$  is either a pleated surface, or a map to  $\overline{N} \setminus N$ , or a mixture of both. This family of pleated surfaces is made up of a sequence of pleated surfaces for each end, and a family of pleated surfaces for  $W$ . The family of pleated surfaces for  $W$  is a family of pleated surfaces for the noninterval bundle part of  $W$  and a sequence for each interval bundle in  $W$ , using the decomposition of 7.10. The family for the noninterval bundle part of  $W$  is actually independent of the choice of ending pleated surfaces. We shall prove that the geometry of this family on the noninterval bundle part depends only on the topological type of  $(W, N)$ , in the case when all ends are incompressible, and on the topological type and a constant  $c_0$  if some ends are compressible. This result was proved by Thurston [61] in the case of incompressible boundary, with comments on what was needed to extend to the case of compressible boundary. The proof of “bounded window frames” given here, in the case of incompressible boundary, is different from that in [61]. The general result is mostly proved in 8.10, with a key hypothesis left to be proved in Section 10.

For the case of  $N$  combinatorial bounded geometry, it is only necessary to read to the end of 8.2, and for the case of  $N$  being an interval bundle, to the end of 8.3.

### 8.1 Sequences of multicurves and pleated surfaces with particular properties.

First, we consider sequences of maximal multicurves and pleated surfaces with certain properties. Suppose that  $[z_0, y_+] \subset \mathcal{T}(S)$ . We suppose that  $S$  is embedded in  $N$ . For each such  $[z_0, y_+]$ , we shall choose an increasing sequence  $\{z_i = [\varphi_i] : 0 \leq i \leq n\}$  of points in  $[z_0, y_+]$  with  $y_+ = z_n$ , with  $d(z_i, z_{i+1}) \leq 1$  and a sequence of maximal multicurves  $\Gamma_j$  on  $S$  such that the following holds for a suitable constant  $\kappa_0$ , and an integer  $r_0$ .



### 8.1.1

$\Gamma_j$  is a noncollapsing maximal multicurve.

### 8.1.2

$|\gamma| \geq \varepsilon_0$  for all  $\gamma \in \Gamma_j$ , unless  $\gamma \in \cap_{k=0}^n \Gamma_k$ .

### 8.1.3

Either  $\#(\Gamma_j \cap \Gamma_{j+1}) \leq r_0$ , or there is  $\zeta_j \in \Gamma_j$ , and a loop  $\gamma_j$  such that  $i(\zeta, \gamma_j) = 0$  for  $\zeta \neq \zeta_j$ ,  $0 < i(\zeta_j, \gamma_j) \leq r_0$ ,  $\Gamma_j \cup \{\gamma_j\} \setminus \{\zeta_j\}$  and  $\Gamma_j \cup \{\tau_{\gamma_j}(\zeta_j)\} \setminus \{\zeta_j\}$  are noncollapsing, and  $\Gamma_{j+1} = (\Gamma_j \setminus \{\zeta_j\}) \cup \{\tau_{\gamma_j}^{\pm n_j}(\zeta_j)\}$ .

### 8.1.4

$$\sum_{p=0}^{n-1} \log(\#(\Gamma_p \cap \Gamma_{p+1}) + 1) \leq \kappa_0 d(z_0, z_n).$$

Then for each  $j$ , we let  $f_j : S \rightarrow N$  be a pleating surface whose pleating locus includes  $\Gamma_j$  and homotopic to the embedding of  $S$  in  $N$ . The extra property which we shall require is the following, which is automatic in the case when  $S$  is incompressible in  $N$ .

### 8.1.5

For  $f = f_j$  or  $f_{j+1}$  (not necessarily both), whenever  $|f(\gamma)| \leq D_0$  for  $\gamma$  nontrivial in  $S$ , then  $f(\gamma)$  is nontrivial in  $N$ .

### 8.1.6 Consequences for the pleated surfaces.

*Bounded distance in Teichmüller space.* If we do have 8.1.1 to 8.1.5, then by 8.1.2, 4.4, 4.5, for a constant  $\kappa'_0$ , depending only on depending the topological type of  $S$ , and a constant  $\kappa_1$  depending only on depending only on  $r_0$  and on the topological type of  $S$ ,

$$d([f_j], [f_{j+1}]) \leq \kappa'_0 (\log(\#(\Gamma_j \cap \Gamma_{j+1}) + 1),$$

$$\sum_{j=0}^{n-1} d([f_j], [f_{j+1}]) \leq \kappa_1 d(z_0, z_n).$$

*Bounded distance between impressions* Also, by 4.3, there is a homotopy in  $N$  between the impressions of  $f_j$  and  $f_{j+1}$  with homotopy tracks of length  $\leq L_0$  in the case when  $i(\Gamma_j, \Gamma_{j+1}) \leq r_0$ , and also in the case  $\Gamma_{j+1} = (\Gamma_j \setminus \{\zeta_j\}) \cup \{\sigma_{\gamma_j}^{n_j}(\zeta_j)\}$ ,

when  $|(\gamma_j)_*|$  is bounded from 0. In general in this case, we can interpolate pleated surfaces  $f_{j,k}$  between  $f_j$  and  $f_{j+1}$ ,  $0 \leq k \leq n_j$ ,  $f_{j,0} = f_j$ ,  $f_{j,n_j} = f_{j+1}$ ,  $f_{j,k}$  has pleating locus  $(\Gamma_j \setminus \{\zeta_j\}) \cup \{\tau_{\gamma_i}^k(\zeta_i)\}$ , and there is a homotopy in  $N$  between  $f_{j,k}$  and  $f_{j,k+1}$  whose homotopy tracks have length  $\leq L_0$ . The hypotheses of 8.1.3) ensure that all the pleating loci are noncollapsing

## 8.2 Sequence of pleated surfaces: combinatorial bounded geometry Kleinian surface case

We show that we can find maximal multicurves  $\Gamma_i$  satisfying 8.1.1-8.1.4 with certain properties in the case of a hyperbolic manifold  $N$  with  $N_c$  homeomorphic to  $S_d \times [0, 1]$  and with ending invariants  $\mu_{\pm}$  of combinatorial bounded geometry, that is, satisfying (7.16.1). We also assume for the moment that both ends are geometrically infinite, that is, that  $\mu_{\pm} \in \mathcal{GL}_a(S)$ . We identify  $S_d$  with a component of  $\partial N_c$ , so that  $S_d$  is embedded in  $N_d$ .

In this case, since  $S_d$  is incompressible in  $N_d$ , 8.1.5 and 8.1.6 are automatically satisfied. We start by choosing homotopic pleated surfaces  $f_{\pm} : S \rightarrow N$  as in 3.11, with  $[f_{\pm}] = y_{\pm}$  close to  $\mu_{\pm}$  and loop sets  $\Gamma_{\pm}$  such that

$$|f_+(\Gamma_+)| \leq L_0, \quad |f_-(\Gamma_-)| \leq L_0. \quad (8.2.1)$$

Then we can choose  $\zeta_{\pm} \subset \Gamma_{\pm}$  similarly to  $\zeta_{n,\pm}$  in 7.16, and then use these to define  $y'_{\pm} = [\varphi'_{\pm}] \in [y_-, y_+]$  as in 7.16, with

$$[y'_-, y'_+] \subset (\mathcal{T}(S))_{\geq \nu}, \quad (8.2.2)$$

and maximal multicurves  $\Gamma'_{\pm}$  such that  $|\varphi'_+(\Gamma'_+)|$  is bounded, and similarly for  $|\varphi'_-(\Gamma'_-)|$ , where  $\Gamma_+ \cap \Gamma'_+$  contains at least one loop, as does  $\Gamma_- \cap \Gamma'_-$ . We then take  $f'_+ : S \rightarrow N$  to be a pleated surface with pleating locus  $\Gamma'_+$ , and similarly for  $f'_-$ . We can extend to cover the case of  $\mu_{\pm} \in \mathcal{T}(S) \cup \mathcal{GL}_a(S)$  by choosing  $f_+$  as in 4.6 if  $\mu_+ \in \mathcal{T}(S)$ , and similarly if  $\mu_- \in \mathcal{T}(S)$ .

Now we shall choose a sequence which satisfies 8.1.1-8.1.6 for  $r_0$  and  $\kappa_0$  depending only on  $\nu$ , with  $z_0 = y'_-$  and  $y_+ = y'_+$ . We choose points  $z_i = [\varphi_i] \in [y'_-, y'_+]$ , and maximal multicurves  $\Gamma'_i$ ,  $0 \leq i \leq n$ , such that  $z_0 = y'_-$ ,  $z_n = y'_+$ ,  $\Gamma'_0 = \Gamma'_-$ ,  $\Gamma'_n = \Gamma'_+$ ,  $d(z_i, z_{i+1})$  is bounded above by 1 and bounded below by  $\frac{1}{2}$  and, for a constant  $L_0 = L_0(\nu)$  (enlarging the previous  $L_0$  if necessary)

$$|\varphi_i(\Gamma'_i)| \leq L_0.$$

Then the bound on  $d(z_i, z_{i+1})$  means that for a constant  $C_1$  depending only on  $\nu$  and the topological type of  $S$ ,

$$\#(\Gamma'_i \cap \Gamma'_j) \leq C_1 e^{C_1|i-j|}.$$

Now by 4.2 we can replace each  $\Gamma'_i$  by a loop set  $\Gamma_i$  such that, enlarging  $L_0(\nu)$  if necessary,

$$|\varphi_i(\Gamma_i)| \leq L_0, \quad (8.2.3)$$

and for all  $\gamma \in \Gamma_i$ ,

$$|\gamma_*| \geq \varepsilon_0. \quad (8.2.4)$$

(8.2.3) implies a bound on  $\#(\Gamma_i \cap \Gamma'_i)$ , and on  $\#(\Gamma_i \cap \Gamma_j)$ , when  $|i-j|$  is bounded. We then take  $f_j$  to be a pleated surface with pleating locus including  $\Gamma_j$ .

Then conditions 8.1.1 to 8.1.6 are satisfied. In condition 8.1.3, only the first alternative,  $\#(\Gamma_j \cap \Gamma_{j+1}) \leq r_0$ , holds. Also, by 4.3, 4.4 applied to  $f_{n+}$  and  $f_+$ ,  $|f_{n+}(\Gamma'_+)| \leq L_0(\nu)$ , assuming that  $L_0(\nu)$  is large enough given  $r_0(\nu)$ , and similarly for  $f_{n-}$  and  $\Gamma'_-$ . Since  $\Gamma_+ \cap \Gamma'_+ \neq \emptyset$ , and assuming  $L_0(\nu)$  is large enough given  $\nu$ , it follows that  $f_{n+}(S)$  either comes within a distance  $L_0(\nu)$  of  $f_+(S)$ , or of a Margulis tube intersected by  $f_+(S)$ .

### 8.3 Sequence of pleated surfaces: interval bundle case.

First we consider the case when  $\partial N_c$  is homeomorphic to an interval bundle  $S_d \times [0, 1]$  and  $N_d$  has just two ends  $e_{\pm}$ . If  $e_+$  is geometrically infinite, we choose  $\Gamma_+$ ,  $[f_+]$  as in 3.11, with

$$|f_+(\Gamma_+)| \leq L_0. \quad (8.3.1)$$

If  $e_+$  is geometrically finite, we choose  $\Gamma_+$  as  $\Gamma$  in 4.6, and  $f_+$  as  $f_2$  of 4.6. So for suitable  $L_0$ , once again, (8.3.1) holds. We choose  $f_-$  similarly. So  $f_{\pm}$  are injective on  $\pi_1$ . We shall now find a sequence of maximal multicurves  $\Gamma_i$ ,  $0 \leq i \leq n$ , satisfying conditions 8.1.1 to 8.1.4 *corresponding to*  $[y_-, y_+]$ . This means that, enlarging  $L_0$  if necessary, depending only on topological type, we shall have

$$d([f_0], [f_-]) \leq L_0, \quad d([f_n], [f_+]) \leq L_0. \quad (8.3.2)$$

In later sections we shall also use this method to define a sequence of maximal multicurves associated to a geodesic segment  $[z_0, y_+] \subset \mathcal{T}(S)$ , where  $y_+$  will be a pleated surface but  $z_0$  will in general not be. There will then be a bound on  $d([f_n], [f_+]) = d([f_n], y_+)$ , but in general there will be no a priori bound on  $d(z_0, [f_0])$  (although such a bound will be obtained eventually). We can then choose a sequence of pleated surfaces  $f_i : S \rightarrow N$  obtained from  $\Gamma_i$  so that 8.1.5 and 8.1.6 are satisfied.

We proceed as follows. We assume that ltd parameter functions have been fixed so that all relevant results in Sections 5 and 6 work. Write  $y_- = z_0$ . We fix a vertically efficient decomposition of  $S \times [z_0, y_+]$  into sets  $\alpha_j \times \ell_j$ . For each  $j$  such that  $\alpha_j$  is a gap, we choose points  $z_{i,j}$  and loop sets  $\Gamma_{i,j}$  on  $S(\alpha_j)$ ,  $0 \leq i \leq n(j)$ , exactly as in 8.2, with  $S(\alpha_j)$  replacing  $S$ . In condition 8.1.3 the first of the two alternatives becomes

$$\#(\Gamma_{i,j} \cap \Gamma_{i,j+1}) \leq r_0(\nu_j),$$

and we can ensure that, if  $z_{i,j} = [\varphi_{i,j}]$ , then for a function  $L_0(\nu)$  independent of the ltd parameter functions,

$$|\varphi_{i,j}(\Gamma_{i,j})| \leq L_0(\nu_j).$$

The vertically efficient conditions ensure that, for each  $j$ , there is  $\nu_j > 0$  bounded below in terms of the ltd parameter functions such that  $|\varphi(\gamma)| \geq \nu_j$  for all loops  $\gamma \subset \alpha_j$  which are nontrivial nonperipheral and not boundary-homotopic, and  $[\varphi] \in \ell$ . If  $(\alpha_j, \ell_j)$  is ltd, then we have in addition that  $\alpha_j$  is long  $\nu_j$ -thick and dominant along  $\ell_j$ . In any case, we have

$$\ell_j \subset (\mathcal{T}(S(\alpha_j)))_{\geq \nu_j},$$

so that the same argument as in 8.2 works. So then we need to use these pieces to produce the points  $z_i$  and multicurves  $\Gamma_i$ . First, we choose a sequence  $z_i^1 = [\varphi_i^1]$  of successive points on  $[y_-, y_+]$ , such that  $d(z_i^1, z_{i+1}^1) \leq L'_0$ , for  $L'_0$  depending only on the topological type, and where  $d_{\alpha_j}(z_i^1, z_{p,j}) \leq L_0(\nu_j)$  for some  $p$ , if  $z_i^1 \in \ell_j$  such that  $\alpha_j$  is a gap, and enlarging  $L_0(\nu)$  if necessary, but still independent of the parameter functions. We then take  $\Gamma_i^1$  to be the union of all the  $\Gamma_{p,j}$  and  $\partial\alpha_j$  such that  $\alpha_j$  is a gap,  $z_i^1 \in \ell_j$  and  $\pi_{\alpha_j}(z_i^1) = z_{p,j}$ . We can also assume that

$$|\varphi_i^1(\Gamma_i^1)| \leq L''_0,$$

where  $L''_0$  depends only on topological type of  $S$  and on the ltd parameter functions.

The properties 8.1.1 to 8.1.4 probably do not hold for this choice of  $\{z_i^1\}$  and  $\{\Gamma_i^1\}$ . In particular, 8.1.2 probably does not hold, because it is likely that  $|(\partial\alpha_j)_*| < \varepsilon_0$ , and that there is a badly bent annulus with core mapping to  $(\partial\alpha_j)_*$  for many of the pleated surfaces in the sequence. These are, however, the only loops in the pleating loci which can be too short, because all others have been removed. If we rectify this, then we need to be careful about 8.1.3 and 8.1.4. At any rate, any modifications necessary concern the loops  $\partial\alpha_j$ . Our new sequence will be obtained from  $\Gamma_i^1$  by replacing the loops  $\partial\alpha_j$  only. So fix a loop  $\gamma$  which is a component of  $\partial\alpha_j$  for at least one  $\alpha_j \times \ell_j$ . Then the  $i$  for which  $\Gamma_{p,j} \subset \Gamma_i^1$  for some  $j, p$  with  $\gamma \subset \partial\alpha_j$  form an interval  $\{i : m_1(\gamma) \leq i \leq m_2(\gamma)\}$ . Let  $n_\gamma(z) = n_{\gamma, \beta(\gamma)}(z)$  (as in 2.7) be defined relative to some fixed loop  $\beta(\gamma)$  with one or two transverse intersections with  $\gamma$ . Then we are going to choose  $z_i^2 = [\varphi^2] = [\varphi^1 \circ \tau]$ . Here,  $\tau$  is a composition of Dehn twists  $\tau_\gamma^{p_{i,\gamma}}$  for a set of disjoint loops  $\gamma$  and integers  $p_{i,\gamma}$ . We choose  $p_i = p_{i,\gamma}$  so that  $n_\gamma(z_i^2)$  is monotone in  $i$  for  $m_1(\gamma) \leq i \leq m_2(\gamma)$ ,  $z_i^2 = z_i^1$  for  $i = m_1(\gamma)$ ,  $m_2(\gamma)$ , and  $|n_\gamma(z_i^2) - n_\gamma(z_{i+1}^2)| \leq r_0$  except for at most one  $i$ . This then gives

$$\sum_{i=m_1(\gamma)}^{m_2(\gamma)-1} |\log(|n_\gamma(z_i^2) - n_\gamma(z_{i+1}^2)| + 1)| \leq \kappa_2 d(z_{m_1(\gamma)}^2, z_{m_2(\gamma)}^2). \quad (8.3.3)$$

So to choose  $\Gamma_i^2$ , choose any loop  $\zeta_i$  such that  $\zeta_i$  has one or two transverse intersections with  $\gamma$ , and

$$|\varphi_i^1(\zeta_i)|'' \leq L'_0$$

where  $|\cdot|''$  is as in 2.5, and  $L'_0$ , as before, depends only on the topological type of  $S$ . It is possible to make such a choice. Then by choice of  $p_i$ , we can ensure the conditions on  $n_\gamma(z_i^2)$ . Since  $n_\gamma([\varphi \circ \tau_\gamma^m], \beta) = n_\gamma([\varphi], \beta) + m$  for any choice

of  $\beta$ , the condition on the  $n_\gamma(z_i^2)$  translates to:  $|p_i - p_{i+1}| \leq r_0$  for all but at most one  $i$ , and (8.3.3) holds with  $p_i$  replacing  $n_\gamma(z_i^2)$ . This then gives 8.1.3 and 8.1.4. To get 8.1.2, note that by 4.1, the bound  $|p_i - p_{i+1}| \leq r_0$  is compatible with  $|(\tau_\gamma^{p_i}(\zeta_i))_*| \geq \varepsilon_0$

We now consider (8.3.2), which involves twists round loops of  $\Gamma_\pm$ . We might not take  $f_0 = f_-$ ,  $f_n = f_+$ , because we want 8.1.2 satisfied for  $\Gamma_0, \Gamma_n$ . By 4.4 we shall have (8.3.2) for  $f_0$  if we choose  $\Gamma_0$  so that there is no badly bent annulus for  $(f_0, f_-)$ . Fix any loop  $\gamma \in \Gamma_- \setminus \Gamma_+$  with  $|\gamma_*| = |f_-(\gamma)| < \varepsilon_0$  and fix  $\beta = \beta(\gamma)$  with one or two intersections with  $\gamma$ , and no other intersections with  $\Gamma$ . By the method used in 4.1, the set of  $k$  for which  $(\tau_\gamma^k(\beta))_*$  does not intersect a bounded neighbourhood of  $\gamma_*$  lie in an interval  $I$  of integers of length  $\leq 1/|\gamma_*|$ . So we choose  $\Gamma_0$  to include  $\tau_\gamma^k(\beta)$  for some  $k$  outside this range, and if possible so that  $|f_0(\tau_\gamma^k(\beta))|''$  is bounded. Then we proceed as for any  $\gamma$  above. Then (8.3.3) will hold for  $\gamma$ , because, for this  $\gamma$ ,  $d(z_{m_1(\gamma)}^2, z_{m_2(\gamma)}^2)$  is bounded below by a multiple of  $-\log |\gamma_*|$  — unless  $m_2(\gamma) = n$ .

If both  $e_\pm$  are geometrically finite, and  $[f_\pm] = \mu(e_\pm)$  are the corresponding invariants, and there is a loop  $\gamma$  such that  $|f_+(\gamma)| \leq \varepsilon_0$ ,  $|f_-(\gamma)| \leq \varepsilon_0$ , then we can choose the loop sets  $\Gamma_0, \Gamma_n$  such that, if  $\beta_\pm$  are elements of  $\Gamma_0, \Gamma_n$  intersecting  $\gamma$ , then

$$n_{\gamma, \beta_+}([f_+]) - n_{\gamma, \beta_-}([f_-]) \leq L_2 |\log |f_+(\gamma)| - |f_-(\gamma)||.$$

We see this from the proof of 4.6. The loops  $\beta_\pm(\gamma)$  simply had to be adjusted by Dehn twists which excluded intervals of integers of lengths  $O(1/|f_\pm(\gamma)|)$ . So if adjustment is necessary, we can choose the two adjustments on the same side of the smaller excluded interval. Then 8.3.3 still holds.

Now we consider modifications for general interval bundles, the case considered in 7.6. In this case,  $f_\pm : S \rightarrow \overline{N}$  are generalised pleated surfaces in the sense of 4.9. We again choose maximal multicurves  $\Gamma_\pm$  and  $f_\pm$  so that all closed loops in the pleating locus of  $f_+$  are in  $\Gamma_+$ , and similarly for  $f_-$ ,  $\Gamma_-$ , and so that (8.3.1) is satisfied. Now  $[f_+]$  is an element of  $\mathcal{T}(S(\alpha_+))$ , where  $\alpha_+ = S \setminus (\cup \Gamma_+)$ , and similarly for  $f_-$ ,  $\Gamma_-$ . In order to completely determine  $y_\pm = [f_\pm] \in \mathcal{T}(S)$ , we also need to define  $\pi_\gamma(y_+)$  for any  $\gamma \in \Gamma_+$  and  $\pi_\gamma(y_-)$  for any  $\gamma \in \Gamma_-$ . It is convenient to make a choice which minimises  $d(y_-, y_+)$  up to an additive constant. Having defined  $y_\pm$ , the rest of the construction is exactly as before.

#### 8.4 General case: sequence of pleated surfaces corresponding to ends, and to bridges between incompressible ends.

let  $N, W, N_{d,W}, \overline{N}$  be as in the introduction to Section 8. Let  $e$  be an end of  $N_{d,W}$ . The sequence for the end  $e$  is determined by  $[f_{e,+}] = y_{e,+}$ ,  $\Gamma_+(e)$ , and by another element  $z_{e,0} = [\varphi_{e,0}] \in \mathcal{T}(S(e))$ , which has to be determined. It is determined as in 7.13 and 7.14, depending on whether  $e$  is compressible or incompressible. The map  $f_{e,+} : S \rightarrow \overline{N}$ , is a generalised pleated surface as in

8.3, and there is a maximal multicurve  $\Gamma_+(e)$  which includes all the closed loops in the pleating locus of  $f_{e,+}$ , and such that

$$|f_{e,+}(\Gamma_+(e))| \leq L_0.$$

If  $f_{e,+}$  maps some nonperipheral loops on  $S$  to cusps in  $N$ , then, similarly to 8.3, we define  $y_{e,+} = [f_{e,+}]$  as an element of  $\mathcal{T}(S(e))$  for each  $\gamma \in \Gamma_+(e)$ . Given a choice of  $z_{e,0}$ , we define  $\pi_\gamma(y_{e,+})$  for  $(\gamma \in \Gamma_+(e))$  so as to minimise  $d(z_{e,0}, y_{e,+})$ , up to a bounded additive constant.

Once we have fixed  $y_{e,+}$  and  $z_{e,0}$ , as in 8.3, if  $e$  is incompressible, we can choose a sequence of maximal multicurves  $\Gamma_{e,j}$ ,  $0 \leq j \leq n_+(e)$  and a sequence of pleated surfaces  $f_{e,j}$  with pleating locus containing  $\Gamma_{e,+}$  so that 8.1.1 to 8.1.6 are satisfied. In analogy to 8.3.2 we shall require

$$d([f_{e,n}], y_{e,+}) \leq L_0.$$

But it is important to note that we do *not* attempt to bound  $d([f_{e,0}], z_{e,0})$ , nor  $|(\Gamma_{e,0})_*|$ . Instead, we choose  $\Gamma_{e,0}$  so that

$$|\varphi_{e,0}(\Gamma_{e,0})|'' \leq L_0,$$

where  $L_0$  depends only on the topological type. We can usually choose  $z_{e,0} = [\varphi_{e,0}] \in (\mathcal{T}(S(e)))_{\geq \varepsilon_0}$  and hence use  $|\cdot|$  rather than  $|\cdot|''$ . In fact, we can always do this if there are no interval bundle bridges between incompressible ends. Even in the case where there are such bridges, We can choose  $z_{e,0}$ , consistent with the choice of  $z_{e,0}$  in 7.14 so that  $z_{e,0} = [\varphi_{e,0}] \in (\mathcal{T}(S(e)))_{\geq \varepsilon_0}$ , unless some loop  $\gamma$  in the common subsurface  $\omega(e, e') = \omega(e', e)$  is such that both  $|\varphi_{e,0}(\gamma)| < \varepsilon_0$  and  $|\varphi_{e',0}(\gamma)| < \varepsilon_0$ .

The choice of  $z_{e,0}, z_{e',0}$  was made in 7.14 so that  $\pi_\omega(z_{e,0}) = \pi_\omega(z_{e',0})$  whenever  $e, e'$  are incompressible ends with  $\omega = \omega(e, e') \neq \emptyset$ . However, we also defined  $z'_{e,0}$  and  $z'_{e',0}$ , essentially with  $[\pi_\omega(z'_{e,0}), \pi_\omega(z'_{e',0})]$  as large as possible. We shall later make use of a sequence of maximal multicurves and pleated surfaces for  $[\pi_\omega(z'_{e,0}), \pi_\omega(z'_{e',0})]$ , that is, for the model manifold  $M(\pi_\omega(z'_{e,0}), \pi_\omega(z'_{e',0}))$ . We call these sequences  $\Gamma_i(e, e'), f_{e,e',i}$  ( $0 \leq i \leq n(e, e')$ ), consisting of maximal multicurves  $\Gamma_i(e, e')$  on  $\omega(e, e')$ ,  $\partial\omega(e, e') \subset \Gamma_i(e, e')$  and corresponding pleated surfaces  $f_{e,e',i}$  with domain  $\omega(e, e')$ . constructed from the interval We can construct such a sequence as before, satisfying 8.1.1 to 8.1.4. We also choose  $\Gamma_{e,0}, \Gamma_{e',0}$  so that

$$\begin{aligned} \Gamma_{e,0} \cap \omega(e, e') &= \Gamma_{e',0} \cap \omega(e, e'), \\ f_{e,0} &= f_{e',0} \text{ on } \omega(e, e'). \end{aligned}$$

At the moment, we have no upper or lower bound on  $|(\partial\omega(e, e'))_*|$ . So it is not yet possible to verify 8.1.6, even though 8.1.5 is automatic. We shall see later in this section that  $|(\partial\omega(e, e'))_*|$  is bounded above. But it may not be bounded below. So since the sequence  $\Gamma_{e,i}$  is chosen so that  $|\gamma_*| \geq \varepsilon_0$  for all  $\gamma \in \Gamma_i(e)$ , it may not be possible to make  $\Gamma_{n(e,e')}(e, e') \subset \Gamma_0(e)$ . But we can ensure that for a suitable  $r_0$ ,

$$\#(\Gamma_{n(e,e')}(e, e') \cap \Gamma_0(e)) \leq r_0.$$

If  $e$  is a compressible end, then we can try to carry out the same construction of sequences of multicurves and pleated sequences as for an incompressible end. But it is not clear that the multicurves are noncollapsing (8.1.1). Nor is it clear that 8.1.5 to 8.1.6 are satisfied. For this, Lemmas 8.5 and 8.8 below are relevant. Lemmas 8.6 and 8.7 will be used to construct the sequence of pleated surfaces for the core.

**Lemma 8.5.** *Let  $N$  be a hyperbolic 3-manifold with finitely generated fundamental group. Let  $S$  be a finite type surface, possibly with boundary, with  $(S_d, \partial S_d)$  embedded in  $(N_d, \partial N_d)$ , and bounding an essential submanifold  $W$  of  $N$ . Fix a Margulis constant  $\varepsilon_0$  and integer  $r$  and a constant  $L_0$ . There is  $L = L(r, (S, N), \varepsilon_0, L_0)$  such that the following holds. Let  $\Gamma_0 \subset S$  be a maximal multicurve. Let  $f_0 : S \rightarrow N$  be a pleated surface homotopic to inclusion, with pleating locus  $\Gamma_0$ , and such that any nontrivial nonperipheral component of  $\partial S$  is mapped to a geodesic of length  $\leq L_0$ . Let  $A$  be the union of badly bent annuli for  $f_0$ . Suppose that  $\gamma' \subset S$  is a simple closed nontrivial loop such that  $\gamma'$  bounds a disc  $D_1$  in  $W$ , with interior disjoint from  $S$  and  $\#(\gamma' \cap \Gamma_0) \leq r$ .*

*Then there is  $\gamma$  such that  $\gamma$  is nontrivial in  $S$  and bounds an embedded disc in  $W$ , and  $|f_0(\gamma) \setminus A| \leq L$  and  $f_0(\gamma)$  is homotopic to the union in  $N$  of  $f_0(\gamma) \setminus A$  and finitely many geodesic segments in  $N$  of length  $\leq L$ .*

*Proof.* Since  $f_0(\gamma')$  has  $\leq r$  intersections with  $f_0(\Gamma_0)$ , we can find a union of  $\leq 8r$  geodesic segments in  $N$  which is homotopic to  $f_0(\gamma')$  and a bounded distance from  $f_0(\gamma')$ . To do this, an arc in a component of  $S \setminus \Gamma_0$ , with endpoints in  $\Gamma_0$ , can be homotoped, keeping endpoints in  $\Gamma_0$ , to either a single segment in the pleating locus of  $f_0$ , which is contained in a geodesic segment homotopic at both ends to loops of  $\Gamma_0$ , and two geodesic arcs which can be taken arbitrarily short in  $S(f_0)$ , or a union of two asymptotic, a segment along a loop of  $\Gamma_0$ , and up to four short arcs joining these up and joining to endpoints in  $\Gamma_0$ . So we can form a loop homotopic to  $f_0(\gamma')$  which is contained in the union of  $\leq 2r$  arcs of  $f_0(\Gamma_0)$ ,  $\leq 2r$  geodesic arcs in the image of the pleating locus of  $f_0$  which are asymptotic to  $f_0(\Gamma_0)$  at each end, and arbitrarily short arcs joining these. Now complete this union of  $\leq 8r$  geodesics to a triangulation of  $D_1$ . So then we have a finite union of  $\leq 8r - 2$  geodesic triangles in  $N$ . This union bounds a disc  $D_1$ , where  $D_1$  is a union of  $\leq 8r - 2$  topological discs of diameter  $\leq L_1$  connected by long thin pieces with two boundary components, which we call “rectangles”. Note that a sequence of short arcs across triangles, avoiding the “thick” parts of the triangles, must have  $\leq 24r - 6$  arcs between intersections with  $\partial D_1$ , because otherwise we can find a closed loop in  $D_1$  whose image in  $N$  can be homotoped to a closed geodesic. This would imply the closed loop was nontrivial, which is impossible since  $S_1$  is a disc. We can choose this decomposition into bounded diameter pieces and thin rectangles so that a rectangle always connects two bounded diameter pieces, simply by adding a bounded diameter piece at an end. We choose one of the bounded diameter topological discs  $D_2$  which has only one boundary component. This is possible: in any partition of a disc by finitely many disjoint arcs, there is at least one complementary component with at most one arc in its boundary. If  $D_2 = D_1$  we are done. If not, there is

exactly one rectangle adjoined to  $D_2$ . If there is a nontrivial loop in  $S(f_0)$  with length  $\leq L_1$  in  $S(f_0) \setminus A$  which is trivial in  $N$ , then we are done. If there is no such loop, then we apply the Short Bridge Arc Lemma of 3.5 to the arc in  $\partial D_2$  which is adjacent to the rectangle. The hypothesis of 3.5 is satisfied, because the length of  $\partial D_2 \cap \partial D_1$  is bounded. Then we can replace the arc in common with the rectangle with an arc in  $S(f_0)$  of bounded length. Then either  $D_2$  is the disc we require, or if its boundary is trivial in  $S(f_0)$ , we can remove it and repeat the process with the remainder of the disc. After repeating the process a bounded number of times, we have a disc with boundary in  $S(f_0)$  of length  $\leq L$  for suitable  $L$ . In both cases, identifying  $S(f_0)$  with  $S$  embedded in  $N$ , we can use the Loop Theorem [23] by another one, with boundary contained in the boundary of the first length (and hence again with image under  $f_0$  of length  $\leq L$ ) and embedded in  $W$ .

□

**Lemma 8.6.** *Take the same hypotheses on  $N$ ,  $S$ ,  $W$ ,  $f_0$ ,  $\Gamma_0$  as in 8.5, but suppose that every component of the surface  $S$ , which is not assumed to be connected, is incompressible in  $W$ , and instead of bounding a disc,  $\gamma'$  bounds a surface  $S_1$  in  $W$  such that  $\#(\partial S_1 \cap \Gamma_0) \leq r$ ,  $S_1$  is incompressible and boundary incompressible (in the sense of 7.10 Rules 2 and 3).*

*Then one of the following holds.*

1.  $|f_0(\partial S_1) \setminus A| \leq L$ , and  $f_0(\partial S_1)$  is homotopic in  $N$  to the union of  $f_0(\partial S_1) \setminus A$  and finitely many geodesic segments in  $N$  of length  $\leq L$ . In particular,  $(\partial S_1)_* \leq L$ .
2. There is an essential embedded annulus  $S_2$  in  $W$  with boundary in  $S$ , such that  $|f_0(\partial S_2) \setminus A| \leq L$ . In particular,  $(\partial S_2)_* \leq L$ .

*It follows that the lengths of the boundary components of maximal interval bundles in  $W$  are bounded.*

*Remark* The final statement leads another proof of Thurston's "bounded window frame" theorem in [61] – but using the full force of the Annulus Theorem. *Proof.* We start as in 8.5: taking a bounded union of  $\leq 8r$  geodesic segments and extending to a triangulation of  $S_1$ . The number of triangles needed is  $\leq 8r + 2k$ , where  $-k$  is the Euler characteristic of  $S_1$ . The corresponding geodesic triangles in  $N$  determine a pleated surface  $f_1 : S_1 \rightarrow N$ , but not the boundary components may consist of finitely many geodesic segments, and not be completely geodesic. This pleated surface gives  $S_1$  the structure of a complete hyperbolic surface  $S(f_1)$ , but the boundary only consists of finitely many geodesic segments, and is only piecewise geodesic. The universal cover of  $S(f_1)$  then identifies a subset with a closed convex subset of the hyperbolic plane, and the covering group is a discrete group of hyperbolic isometries of the hyperbolic plane. So Margulis' Lemma holds [58]. For a suitable Margulis constant  $\varepsilon_0 > 0$ , nontrivial geodesics of length  $< \varepsilon_0$  are disjoint and separated by distance  $-\log \varepsilon_0 - O(1)$  from all closed geodesics of length  $\leq 1$ . But in fact, Margulis' Lemma extends in this case of a hyperbolic surface with piecewise geodesic boundary. Nontrivial



arcs of length  $\leq \varepsilon_0$  between boundary components, and in different homotopy classes, are also disjoint. A nontrivial arc of length  $< \varepsilon_0$  and nontrivial loops of length  $< \varepsilon_0$  are separated by rectangles and cylinders of length  $-\log \varepsilon_0 - O(1)$ . So we have a decomposition of  $S(f_1)$  into a finite union of  $\leq 2k + 8r$  bounded diameter pieces, which, this time, may be connected by cylinders around  $\leq 2k$  short closed loops, as well as  $\leq 2k + 8r$  long thin rectangles. In order to bound  $|(\partial S_1)_*|$ , it suffices to show there are no rectangles over a given length.

Suppose that  $R$  is such a rectangle, with long sides in  $f_0(S(f_0))$ . If  $R$  intersects  $N_{<\varepsilon}$  for a suitable  $T(f, \gamma_*, \varepsilon)$ , and then we deduce from the Radius of Injectivity Lemma 3.4 that  $\gamma$  is freely homotopic in  $N$  to  $f_0(\gamma_1)$  and  $f_0(\gamma_2)$  for loops  $\gamma_1$  and  $\gamma_2$  which are not freely homotopic in  $S$ . Then we can choose the annulus  $f_1(S_2)$  homotopic to  $\gamma$ , giving conclusion 2. So now we assume that the sides of the rectangle do not intersect  $f_0((S(f_0))_{<\varepsilon})$  for a suitable  $\varepsilon$  depending only on the original Margulis constant  $\varepsilon_0$  the constant  $C(\varepsilon_0)$  of the Radius of Injectivity Lemma. Parametrize nearest points on the long sides of the rectangle by  $(\zeta_1(t), \zeta_2(t))$ , where  $t$  is a length parameter, with  $t \in [0, T]$ , say. Then  $\{(\zeta_1(t), \zeta_2(t)) : t \in [0, T]\}$  lies in a compact subset of  $(S(f_0))_{\geq \varepsilon} \times ((S(f_0))_{\geq \varepsilon})$ , of diameter bounded in terms of  $\varepsilon$  using the product of the hyperbolic metric on  $S(f_1)$ . Since  $S$  is incompressible, if the lift of the rectangle has boundary components in the same component of the lift of  $f_0(S)$ , and  $T$  is sufficiently large given  $\varepsilon$ , we can apply the Short Bridge Arc Lemma to deduce that an arc across the rectangle can be homotoped to a short arc in  $S$ . So such rectangles can either be removed (if they are boundary-homotopic in  $S_1$ ), or discounted, using the boundary incompressibility of  $S_1$ .

So now we assume that the lift of the rectangle has boundary components in different lifts of  $f_0(S)$ . So if  $T$  is sufficiently large given  $\varepsilon$ , for any  $t_0 \in [0, T]$ , we can find  $t_1, t_2$  with  $|t_2 - t_1| \geq 1$  but  $|t_i - t_0|$  bounded above in terms of  $\varepsilon$  for  $i = 1, 2$ , and such that the hyperbolic distance in  $S(f_0)$  between  $f_0(\zeta_j(t_1))$  and  $f_0(\zeta_j(t_2))$  is  $< \varepsilon/10$  for  $j = 1, 2$ , and the distance in  $N$  between  $f_0(\zeta_1(t_j))$  and  $f_0(\zeta_2(t_j))$  is also  $< \varepsilon/10$  for  $j = 1, 2$ . Then we can join  $\zeta_j(t_1)$  and  $\zeta_j(t_2)$  by a short arc  $\beta_j$  to give a nontrivial closed loop  $\alpha_j$  on  $S(f_0)$ , which we identify with  $S$ , embedded in  $N$ . Let  $\gamma_j$  denote the short arc across the rectangle from  $\zeta_1(t_j)$  to  $\zeta_2(t_j)$ . Then  $f_0(\gamma_1 * \beta_2 * \overline{\gamma_2} * \beta_1)$  has length  $\leq 2\varepsilon/5$ . Since it is in  $N_{\geq \varepsilon}$  it must be trivial. So  $\gamma_1 * \beta_2 * \overline{\gamma_2} * \beta_1$  is trivial in  $W$ . So the loops  $\alpha_1$  and  $\alpha_2$  are freely homotopic in  $W$  but not in  $S_1$ . By the proof of Waldhausen's Annulus Theorem (Theorem 3) in [17], we can find a boundary incompressible embedded annulus with boundary in  $S(f_0)$  arbitrarily close to a subset of  $\alpha_1 \cup \alpha_2$ . Taking the image of this boundary under  $f_0$  This gives alternative 2 above. If  $T$  is bounded for all such rectangles  $R$ , then we obtain alternative 1.

The bound on the boundary of maximal interval bundles is achieved by applying the above with  $S_1$  an annulus, if there is one, and then replacing by an annulus with bounded boundary length, and then repeating the process until a maximal set of boundary incompressible annuli with bounded boundary lengths has been constructed. This uses the fact that a sufficiently long geodesic segment in  $H^2$  which has endpoints a bounded distance from a segment joining  $x_0$  and  $g^n.x_0$  must project to have self-intersections in  $S(f_0)$ . This needs to be

applied to  $g$  in the conjugacy class of bounded annuli boundaries, in order to construct a next annulus with bounded boundary, if there is another annulus homotopically disjoint from a set already constructed.  $\square$

In 8.5, when  $S_1$  is a disc, we only know that there is some disc, possibly different, whose boundary has bounded length. In the case  $W \neq W_n$ , we can strengthen this, as follows. This will mean that if we have good information on the geometry of  $W = W_n$ , then we can ensure bounds on  $\partial(\partial W_i \cap \partial W_{i+1})$ .

**Lemma 8.7.** *Let  $N, S$  be as in 8.5. Let  $S = S_1 \cup S_2$ , and such that  $\gamma_1, \gamma_2$  and  $\gamma_1 * \gamma_2$  are nontrivial in  $N$ , whenever  $\gamma_i$  is a closed loop with endpoints on  $\partial S_1 = \partial S_2$  which is nontrivial in  $S_i$ . Let a Margulis constant  $\varepsilon_0$  and another constant  $L_1$  be given.*

*Then there exists  $L = L((S, N), \varepsilon_0, L_1)$  such that the following holds. Let  $\gamma \subset S$  bound a disc in  $N$  such that  $\gamma$  is nontrivial in  $S$  and indecomposable, in the sense that  $\gamma$  is not homotopic in  $S$  to  $\gamma_3 * \zeta * \gamma_4 * \bar{\zeta}$ , where  $\gamma_3 * \zeta * \gamma_4 * \bar{\zeta}$  has only essential intersections with  $\partial S_1 = \partial S_2$  and  $\gamma_j$  is homotopically nontrivial in  $S$  and trivial in  $N$  for  $j = 3, 4$ .*

*Let  $f : S \rightarrow N$  be a pleated surface homotopic to the identity, and with  $|f(\partial S_2)| \leq L_1$ . Let  $A$  be the union of badly bent annuli for  $f$ . Then  $|f(\gamma) \setminus A| \leq L$  and  $f(\gamma)$  is homotopic in  $N$  to the union of  $f(\gamma) \setminus A$  and a finite union of geodesic segments in  $N$  of length  $\leq L$ .*

*Proof.* We use the argument of 8.5, splitting  $f(\gamma)$  up into bounded diameter bits and long thin rectangles. If there is a long thin rectangle, between two different discs, there is either one with boundary in  $f(S_i)$  for one of  $i = 1, 2$ , or one with one boundary component in each of  $f(S_1), f(S_2)$ . In the first case, since  $S_i$  is incompressible, we can apply the Short Bridge Arc Lemma of 3.5 to deduce that the arc across the rectangle can be homotoped into  $f(S_i)$ . So we have two discs, neither of which can have boundary completely in  $S_i$ , because  $S_i$  is incompressible. So we get a contradiction to indecomposability, if there are no such rectangles and we have a bound on  $|f(\gamma) \setminus A|$ . The alternating bits in  $N$  are bounded because of the way  $f(\gamma)$  was split up. In the second case, the same argument works as in 8.6. If there is a long thin rectangle then, as in 8.6, there must be a long subrectangle in  $N_{\geq \varepsilon_0}$ . Then, as in 8.6 we can find closed loops on  $S_1$  and  $S_2$  which are not multiples of loops in  $\partial S_1 = \partial S_2$ , and such that the product is trivial in  $N$ , contradicting our assumption.  $\square$

The following lemma is useful for obtaining information about a pleated surface corresponding to a compressible component of  $W$ . It will be applied in conjunction with 8.5.

**Lemma 8.8.** *Let  $N$  be a hyperbolic 3-manifold with finitely generated fundamental group, and let  $W \subset N_c$  be such that each component of  $N_d \setminus W$  is homeomorphic to the interior of an interval bundle. Fix such a component, with corresponding component  $S_d$  of  $\partial_d W$ , where  $S_d$  is the horodisc deletion of  $S$ . Let  $z_0 = [\varphi_0] \in \mathcal{T}(S)$ . Let  $w_0 = [\xi_0] \in \mathcal{T}(S)$ . Let  $\Gamma_+ \subset S$  be a multicurve of*

loops which are all nontrivial in  $N$ , with  $|\xi_0(\Gamma_+)| \leq L_0$ . Let  $\Gamma_+$  satisfy

$$\begin{aligned} i(\zeta, \mu) &\geq c_0|\zeta| \text{ for at least one } \zeta \in \Gamma_+ \\ \text{if } i(\gamma', \mu) &\leq c_0|\gamma'|, \end{aligned} \quad (8.8.1)$$

whenever  $\gamma'$  is a simple closed loop which is nontrivial in  $S$  but trivial in  $N$ ,  $\mu$  is a geodesic lamination on  $S$ , with a normalised transverse invariant measure.

Let  $y_0 = [\psi_0]$  satisfy

$$|\psi_0(\gamma)|'' \leq L_0,$$

where  $\gamma$  is a closed loop which is nontrivial on  $S$  but bounds a disc in  $N$ .

Let  $x(\cdot) = x(\cdot, [y_0, w_0])$ , as in 6.11.

Then for a constant  $D_2 = D_2(L_0, z_0, c_0)$ , which is locally bounded in  $z_0$  and also depends on the topological type of  $(S_d, W)$  and suitable ltd parameter functions,

$$d(z_0, x(z_0)) \leq D_2.$$

*Proof.* Suppose for contradiction that  $d(z_0, x(z_0)) \geq D_2$ . Then we can find  $\alpha$  which is ltd along a segment of  $[x_\alpha(z_0), z_0]$ , and for some  $x_\alpha(z_0) = [\chi_0] \in [y_0, w_0]$ , for  $C_1$  depending only on the ltd parameter functions,

$$|\chi_0(\partial\alpha) \leq C_1,$$

$$d_\alpha(x(z_0), x_\alpha(z_0)) \leq \log C_1,$$

$$d'_\alpha(x_\alpha(z_0), z_0) \geq D_2 - \log C_1.$$

Now for suitable  $L_0$  we can choose  $\zeta_2$  such that  $\zeta_2 \subset \alpha$  and

$$|\chi_0(\zeta_2)| \leq L_0.$$

Then by 6.6, or, at least, by the locking technique employed in 6.6, enlarging the constant  $C_1$  if necessary but still only depending on the ltd parameter functions,

$$|\varphi_0(\gamma)| \geq C_1^{-1} i(\gamma, \zeta_2) \exp d'_\alpha(x_\alpha(z_0), z_0).$$

This is trivially satisfied if  $\gamma$  does not intersect  $\zeta_2$ . It uses the definition of  $x(z_0)$ , that is, that  $x_\alpha(z_0)$  is a bounded  $d_\alpha$  distance, depending only on the ltd parameter functions, from each of  $[z_0, w_0]$  and  $[z_0, y_0]$ . Similarly, for any  $\zeta \in \Gamma_+$ ,

$$|\varphi_0(\zeta)| \geq C_1^{-1} i(\zeta, \zeta_2) \exp d'_\alpha(x_\alpha(z_0), z_0).$$

Now  $|\varphi_0(\zeta)|$  is boundedly proportional to  $|\zeta|$  for any loop  $\zeta$ , with bound depending locally uniformly on  $z_0$ . So we have

$$i(\gamma, \zeta_2) \leq C_1^3 e^{-D_2} |\gamma|,$$

and similarly with  $\zeta$  replacing  $\gamma$  for any  $\zeta \in \Gamma_+$ . Putting  $\gamma = \gamma'$ , and  $\mu = \zeta_2/|\zeta_2|$ , we obtain a contradiction to (8.8.1), if  $D_2$  is large enough given  $L_0, c_0$  and locally on  $z_0$ .

□

## 8.9 Sequence of pleated surfaces corresponding to an incompressible end.

We shall always use the following hypothesis on  $\Gamma_{e,+}$ , which depends on a constant  $c_0$ . The assumption is a bit stronger than required in (8.8.1). We can manage with just (8.8.1) for much of what follows, but not all, so we might as well fix on the stronger assumption now.

$$\begin{aligned} i(\zeta, \mu) &\geq c_0 |\zeta| \text{ for at least one } \zeta \in \Gamma_+ \\ \text{if } i(\mu, \mu') &= 0 \text{ and } i(\mu' \gamma') \leq c_0 |\gamma|, \end{aligned} \quad (8.9.1)$$

whenever  $\gamma'$  is a simple closed loop which is nontrivial in  $S$  but trivial in  $N$ , and  $\mu, \mu'$  are geodesic laminations on  $S$ , with normalised transverse invariant measures. With this assumption, fix any  $z_{e,0} = [\varphi_{e,0}] \in \mathcal{T}(S(e))$ . We can find sequences of multicurves  $\Gamma_{e,j}$  on  $S(e)$  and  $f_{e,j} : S(e) \rightarrow N$  homotopic to inclusion of  $S(e)$  in  $N$  ( $0 \leq j \leq n_+(e)$ ) satisfying 8.1.1 to 8.1.4, with

$$|\varphi_{e,0}(\Gamma_{e,0})| \leq L_0,$$

where  $L_0$  is bounded in terms of  $z_{e,0}$  and  $c_0$ , but does not depend further on  $y_{e,+}$ . We see this as follows. By replacing  $z_{e,0}$  by a point on  $[z_{e,0}, y_{e,+}]$  sufficiently far from  $z_{e,0}$  given  $D_0$ , by 8.5 and 8.8 we can ensure that

$$|\varphi(\gamma)|'' \geq D_0 |\gamma| \quad (8.9.2)$$

for all  $[\varphi] \in [z_{e,0}, y_{e,+}]$  and any  $\gamma$  which is nontrivial on  $S(e)$  but trivial in  $N$ . Now for some constant  $L_0$  depending on an initial choice of ltd parameter functions, we can choose the sequences  $z_i^2 = [\varphi_i^2]$  and  $\Gamma_i = \Gamma_i^2$  as in 8.3 so that for

$$|\varphi_i^2(\Gamma_i^2)|'' \leq L_0.$$

If  $D_0$  is sufficiently large, none of the loop sets can be collapsing. For if some one is, then for some  $i$  and  $\gamma \in \Gamma_i^2$  if we take any loop  $\zeta$  intersecting  $\gamma$  at most twice and disjoint from other elements of  $\Gamma_i^2$ , not separating loops with the same image in  $N$ ,  $\tau_\gamma^n(\zeta)$  is trivial in  $N$  for all  $n$ . Replacing  $\zeta$  by  $\tau_\gamma^n(\zeta)$  for a suitable  $n$ , we can assume that  $|\varphi_i^2(\zeta)|''$  is bounded in terms of  $L_0$ . This contradicts (8.9.2) if  $D_0$  is large enough given  $L_0$ . 8.8 also implies that (8.1.5) and (8.1.6) are satisfied for  $j \geq j_0$ , if  $j_0$  is the largest integer such that

$$d(x([f_j], [z_{e,0}, y_{e,+}]), z_{e,0}) \leq D_2.$$

We shall use this in Section 10.

Then by 8.8, if  $D_0$  is sufficiently large given  $D_2$  and  $D_1$ ,  $|f_j(\gamma)| \geq D_1$  whenever  $T([f_j], +, [z_0, y_+])$  (in the notation of 6.10) does not contain  $(\alpha_1, \ell_1)$ . This is automatically true for  $j = n_+$ .

## 8.10 Family of pleated surfaces for the non-interval bundle part of the core

So far, we have constructed a sequence of maximal multicurves and pleated surfaces for each end  $e$  of  $N_{d,W}$ . Conditions 8.1.1 to 8.1.4 are satisfied for all

ends. Conditions 8.1.5 and 8.1.6 are also satisfied for incompressible ends. It is not clear if 8.1.5 is satisfied for compressible ends. If it is, then 8.1.6 is also satisfied. We are now going to construct a family of pleated surfaces for  $W$ , under a temporary assumption which we shall prove in Section 10. The temporary assumption we make is the following. It is, of course, unnecessary if all ends are incompressible.

### 8.10.1 Assumption on compressible ends

For any compressible end,

$$|f_{e,0}(\Gamma_{e,0})| \leq L_1.$$

**Theorem** *Assume that 8.10.1 holds. Let  $W_i$  be a decomposition of  $W$  satisfying conditions 1 to 7 of 7.10. Then we can find a family of maximal multicurves and corresponding pleated surfaces for the non-interval-bundle part  $W'$  of  $W$ , consisting of sequences of two multicurves and pleated sequences for each component  $S$  of  $\partial V$ , each non-ball component  $V$  of  $W_i$ , each  $i \geq 1$ , such that the following hold, for a suitable constant  $L_0$ , and for  $L_2$  sufficiently large given  $L_0$  and  $L_1$ .*

1. *For the two multicurves  $\Gamma, \Gamma'$  in the sequence for  $S$   $\#(\Gamma \cap \Gamma') \leq L_0$ .*
2. *If  $i = n$  and  $S = S(e)$  and  $e$  is a compressible end then the second multicurve of the sequence for  $S(e)$  is  $\Gamma_0(e)$ , and the corresponding pleated surface is  $f_{e,0}$ . (These are the first elements in the sequence for the end  $e$ .)*
3. *If  $S$  is incompressible, then the first multicurve of the sequence includes  $\partial(S \cap S')$  for any component  $S'$  of  $\partial V'$ , for  $V'$  a component of  $W_{i-1}$ ,  $V' \subset V$ . For any such  $S'$ ,*

$$|\partial(S \cap S')|_* \leq L_2.$$

4. *Whether  $S$  is compressible or incompressible, with  $i < n$ , the second multicurve in the sequence for  $S$  coincides with the first multicurve for  $S''$  on  $S \cap S''$ , where  $S''$  is a component of  $\partial V''$  for a component  $V''$  of  $W_{i+1}$ ,  $V \subset V''$ ,  $S \cap S'' \neq \emptyset$ . If  $V_2$  is another component of  $W_i$  in  $V''$  and  $S_2$  is the component of  $\partial V_2$  meeting  $S$ , then the second multicurves in the sequences for  $S$  and  $S_2$  coincide on  $S \cap S_2$ .*
5. *If  $S$  is compressible, the first multicurve in the sequence for  $S$  can be written in the form  $\Gamma_1 = \Gamma_{1,1} \cup \Gamma_{1,2}$ , where  $\Gamma_{1,1} \cup \Gamma'_{1,2}$  is also a maximal multicurve, and  $\Gamma'_{1,2}$  is the set of distinct isotopy classes in  $S$  of  $\partial(S \cap S')$  for any component  $S'$  of  $\partial V'$ , for  $V'$  a component of  $W_{i-1}$ ,  $V' \subset V$ .*
6. *Let  $\Sigma'$  be the graph in  $W$  of 7.11. All the maximal multicurves in the sequence for  $S$  are made up of  $\leq r_0$  arcs of  $\Sigma'$  on  $S$ . For any pleated surface  $f : S \rightarrow N$  in the sequence, and any  $\gamma \subset S$  made up of  $\leq r_0$  arcs of  $\Sigma'$  if  $A$  is the set of badly bent annuli for  $f$ , then*

$$|f(\gamma) \setminus A| \leq L_2,$$

and  $f(\gamma)$  is homotopic in  $N$  to the union of  $f(\gamma) \setminus A$  and finitely many arcs of total length  $\leq L_2$ .

7. There is a map  $f_{W'} : W' \rightarrow N$  which is homotopic to inclusion, such that  $f|\partial W' \cap S_d(e) = f_{e,0}$  for any end  $e$ , and such that for any arc  $\gamma$  in  $\Sigma'$ , or between arcs of  $\Sigma'$  which are homotopic then  $|f(\gamma)| \leq L_2$ .

### 8.11 Proof of 8.10: 1-5 and a bit.

The construction is inductive, working down. To start with we only aim to satisfy 1 to 4, and the following extension of 5:

**5 and a bit** If  $S$  is compressible, the first multicurve in the sequence can be written in the form  $\Gamma_{1,1} \cup \Gamma_{1,2}$ , where  $\Gamma_{1,1} \cup \Gamma'_{1,2}$  is also a maximal multicurve, and  $\Gamma'_{1,2}$  is the set of distinct isotopy classes in  $S$  of  $\partial(S \cap S')$  for any component  $S'$  of  $\partial V'$ , for  $V'$  a component of  $W_{i-1}$ ,  $V' \subset V$ . For any  $\gamma \in \Gamma_{1,2}$ ,  $\gamma \in \Gamma_{1,1}$  adjacent to a loop of  $\Gamma_{1,2}$  (equivalently of  $\Gamma'_{1,2}$

$$|\gamma_*| \leq L_2.$$

So suppose that  $S$  is a component of  $\partial V$ ,  $V$  a component of  $W_i$ . If  $i < n$ , suppose that the conditions are already satisfied for all components of  $V''$ , where  $V''$  is the component of  $W_{i+1}$  containing  $V$ . If  $i = n$  and  $S = S(e)$  for compressible  $e$ , then 2 is satisfied by definition. If  $V$  is incompressible, then we obtain 3 from 8.6, for any  $i$ . Then 1 is true, for  $L_0$  depending only on the topological type of  $W'$ . Satisfying 4 inductively is no problem.

So now we need to show that 5 and a bit can be satisfied. If  $i = n$ , and  $S = S(e)$  for a compressible end  $e$ , let  $\Gamma_0$  be the second multicurve in the sequence for  $S = S(e)$ . By construction,  $\Gamma_0$  is noncollapsing. If  $i + 1 < n$ , let  $S''$  be component of  $\partial V''$ ,  $V'' \subset W_{i+1}$ , which intersects  $S$ , and let  $\Gamma_0 = \partial(S \cap S'')$ . According to rule 6 of 7.10,  $S''$  is incompressible in  $N$ , and so the loops of  $\Gamma_0$  are nontrivial in both  $N$  and  $S$ . According to Rule 4 of 8.10, the second multicurve  $\Gamma_2$  in the sequence for  $S$  should include  $\Gamma_0$ . We can choose  $\Gamma_2$  so that this is true. Using 4.1, we can also ensure that  $|\gamma_*| \geq \varepsilon_0$  for  $\gamma \in \Gamma_2 \setminus \Gamma_0$ . Now let  $\Gamma'_{1,2}$  be the set of boundaries of compressing discs attached to  $S$  to form the components of  $W_{i-1}$  in  $V$ . By Rule 6 of 7.10,  $\Gamma'_{1,2}$  bounds a maximal set of disjoint compressing discs. Then every loop on  $S''$  disjoint from  $\Gamma'_{1,2}$  is nontrivial in  $V''$ , and hence in  $N$ . (It can be proved inductively that there is no compressing disc attached to the exterior of  $V''$ .) Let  $T \subset S'$  be the surface with boundary which is the convex hull of  $\Gamma'_{1,2}$  and  $\Gamma_0$ . Then all loops in  $\partial T$  (if  $\partial T \neq \emptyset$ ) are nontrivial in  $N$ , since otherwise  $\#(\Gamma'_{1,2})$  is not maximal.

Let  $f_2$  be the second pleated surface in the sequence for  $S$ . Then the conditions of 8.7 are satisfied by rule 6 of 7.10. We can therefore use 8.7 to bound  $|f_2(\Gamma'_{1,2}) \setminus A|$ , and the length of connecting geodesic segments in  $N$  where  $A_2$  is the union of badly bent annuli for  $f_2$ . We already have a bound on  $|f(\Gamma_0)| = |(\Gamma_0)_*|$  by 8.7. So let  $\tau = \Gamma_0 \cup \Gamma'_{1,2}$ . We have a bound on  $|f(\tau)|$ . Then choose  $\Gamma_{1,1}$  with  $\partial T \subset \cup \Gamma_{1,1}$  such that  $\Gamma'_{1,2} \cup \Gamma_{1,1}$  is a maximal multicurve on  $S$ .

Since  $\#(\Gamma'_{1,2})$  is maximal,  $\Gamma_{1,1}$  is noncollapsing. Then by the lemma in 3.11, we can extend  $\Gamma_{1,1}$  to a maximal noncollapsing maximal multicurve  $\Gamma_1 = \Gamma_{1,1} \cup \Gamma_{1,2}$  on  $S$ . Since  $\partial T \subset \Gamma_{1,1}$  and  $\Gamma_{1,2} \subset T$ , the loops of  $\Gamma_{1,2}$  and  $\Gamma_{1,1}$  adjacent to  $\Gamma'_{1,2}$  (equivalently  $\Gamma_{1,2}$ ) are in  $\tau$  up to homotopy. So, simply depending on the topology of  $\tau$  (that is, of the decomposition into the  $W_i$ , and, ultimately, of  $W$ ) we can bound the number of arcs of  $\tau$  making up the loops of  $\Gamma_{1,2}$ , and the arcs of  $\Gamma_{1,1}$  adjacent to  $\Gamma_{1,2}$ . So we have 5 and a bit. We can also ensure, by 4.1, that  $|\gamma_*| \geq \varepsilon_0$  for  $\gamma \in \Gamma_{1,1} \setminus \partial T$ .

## 8.12 Proof of 6 and 7 of 8.10

We have bounds on  $|f(\gamma)|$  whenever  $f : S \rightarrow N$  is a pleated surface in the family, and  $\gamma$  is an arc in  $\Sigma \cap S$ . The arcs in  $\Sigma'$  are obtained from  $\Sigma$  by successive homotopies between the boundaries of the interval bundles which make up  $W$ . The elements of  $\Sigma$  are themselves obtained from homotopy images, this time, from a certain set of closed simple loops. So it suffices, to bound the length, to find homotopies between pleated surfaces in the family with the same domain, with bounded homotopy tracks, outside badly bent annuli round short loops in the pleating locus. At the join between two or three interval bundles in  $W$ , two or three surfaces have parts of the domain in common, with a closed geodesic in common to the pleating locus of the two or three surfaces. When this happens, one surface  $S$  is a boundary component of  $\partial V$  for some component  $V$  of  $W_i$ , some  $i$ , and the other one or two surfaces are boundary components of components of  $W_{i-1}$ .

We consider first the case when  $S$  is compressible. Let  $f_1$  and  $f_2$  be the first and second pleated surfaces with domain  $S$ , as in 8.11. In this case, the domain of any  $S'$  which is attached to  $S$ , and a component of the boundary of some component of  $W_{i-1}$ , identifies with a subsurface of  $S$  bounded by loops in  $f_1$ . Now we consider the homotopy between  $f_1$  and  $f_2$ . If  $S$  is compressible, the condition of 4.3 which replaces injectivity-on- $\pi_1$  is probably not satisfied. But for  $f_2$ , the loop set  $\Gamma_0$  constructed in 8.11 decomposes  $S$  into incompressible surfaces, which, for the moment we call  $S'_j$ ,  $1 \leq j \leq r$ . Essentially, we shall apply the Short Bridge Arc Lemma to these. For  $f_1$ , we have a decomposition of  $S$  into surfaces  $S_{1,1}$  and  $S_{1,2}$  with  $\Gamma_{1,2} \cup \Gamma'_{1,2} \subset S_{1,2}$  and  $\partial S_{1,2} \subset \Gamma_{1,1}$ , and  $S_{1,1}$  is incompressible. In fact, we can further decompose  $S_{1,1}$  using  $\Gamma_{1,1}$ , using the  $S'_j$ , into surfaces  $S_{1,1,j}$  which are incompressible. The surface  $S_2$  is in  $T$  and  $\Sigma' \cap T$  can be homotoped into  $\tau$ . We apply 8.7 to  $f_1$  and deduce bounds on  $|f_1(\tau) \setminus A_1|$ , and on the geodesic arcs homotopic in  $N$  to components of  $f_1(\tau) \cap A_1$  where  $A_1$  is the union of any badly bent annuli on  $S$  for  $f$ . If there are any badly bent annuli, they are round loops of  $\Gamma_{1,1}$  adjacent to  $\Gamma_{1,2}$ , each of which actually gives two loops in the pleating locus for  $f_1$ . Note that this then automatically transfers to the next surface down. Let  $A_2$  be the set of badly bent annuli for  $f_2$ . We have chosen the pleating locus of  $f_2$  so that these only occur round loops  $\partial(S'_j \cap S)$ , where  $S'$  are the boundary components of  $V'$  which  $S$  meets, where  $V'$  is the component of  $W_{i+1}$  containing  $V$ .

First, we bound  $|f_2(\partial S_{1,j} \cap S'_k) \setminus A_2|$  and the geodesic segments homotopic

to  $f_2(\partial S_{1,j} \cap S'_k) \cap A_2$  using the Short Bridge Arc Lemma 3.5 applied to the surfaces  $S'_k$ . Similarly we bound  $|f_1(\Gamma_0 \cap S_{1,j}) \setminus A_1|$  and the geodesic segments homotopic to  $f_1(\Gamma_0 \cap S_{1,j}) \cap A_1$ . Taking the unions over all  $S'_k$ , or over  $S_{1,1}$  and  $S_{1,2}$ , we obtain bounds on  $|f_2(\partial S_{1,j}) \setminus A_2|$ ,  $|f_2(\partial S_{1,j}) \cap A_2|$ ,  $|f_1(\Gamma_0) \setminus A_1|$ ,  $|f_1(\Gamma_0) \cap A_1|$ . Then we use 3.5 to homotope  $f_2|_{\tau \setminus (f_1^{-1}(A_1) \cup f_2^{-1}(A_2))}$  to  $f_1|_{\tau \setminus (f_1^{-1}(A_1) \cup f_2^{-1}(A_2))}$ , and hence bound  $|f_1(\tau) \setminus A_1|$ , and the union of geodesic segments homotopic to  $f_1(\tau) \cap A_1$ . Similarly if  $\tau' \subset \Sigma'$  and  $|f_2(\tau') \setminus A_2|$  is known to be bounded and bounds are known on the geodesic segments homotopic to  $f_2(\tau' \cap A_2)$ , we can homotope  $f_2|_{\tau' \setminus (f_2^{-1}(A_2) \cup f_1^{-1}(A_1))}$  to  $f_1|_{\tau' \setminus (f_1^{-1}(A_1) \cup f_2^{-1}(A_2))}$ , and hence bound  $|f_1(\tau') \setminus A_1|$ , and the union of geodesic segments homotopic to  $f_1(\tau') \cap A_1$ . Assuming as we may do that  $\#(\Gamma_1 \cap \Gamma_2)$  is bounded in terms of the topological type, we can then extend the homotopy, using 3.5 as before, to produce a bounded track homotopy between  $f_1$  and  $f_2$  outside  $f_1^{-1}(A_1) \cup f_2^{-1}(A_2)$ .

We can also transfer lengths to a surface  $f_0 : S_0 \rightarrow N$ , where  $S_0$  is a component of  $\partial V'$ ,  $V'$  a component of  $W_{i-1}$ ,  $S_0$  attached to  $S$ , and incompressible. If this happens, pairs of loops in  $\Gamma_{1,1}$  on  $S$  identify in  $N$ , and also identify with a loop in the pleating locus of  $f_0$ . The second maximal multicurve on  $S_0$  is given by  $\Gamma_{1,1}$  after identification. We use all loops of  $\Gamma_{1,1}$  if  $S_0$  is connected. In general, the loops of  $\Gamma_{1,1}$ , after identifications, give the union of multicurves for the boundary components of the components of  $W_{i-1}$  attached to  $S$ . Let  $A_0$  be the union of badly bent annuli for  $f_0$ . These, again, have as cores a subset of the loops of  $\Gamma_{1,1}$  with badly bent annuli in  $A_1$ . The subsurface we are calling  $S_{1,1}$  glues up along matching loops in  $S_{1,1}$  to give  $S_0$ . Suppose that  $\tau_1 = \tau_{1,1} \cap \tau_{1,2}$ ,  $\tau_2 = \tau_{1,2} \cup \tau_{2,2} \subset S_{1,1}$  have endpoints in  $\partial S_{1,1} = \partial S_{1,2}$ , with all four graphs meeting along a loop in  $\partial S_{1,1}$ ,  $\tau_{1,1} \cup \tau_{1,2}$  identifying with a graph on  $S_0$ . Suppose also, from looking at  $f_1(\tau_1)$  and  $f_1(\tau_2)$ , we already have bounds on the length of geodesic segments making up  $\tau_1$ ,  $\tau_2$ . Suppose also that  $\tau_{1,2}$  and  $\tau_{2,2}$  lie in parts of  $S$  where the multicurve loops from  $\Gamma_{1,2}$  are identified in  $N$ , and that they are homotopic in  $N$ , under homotopy preserving endpoints. Then  $\tau_{1,1} \cup \tau_{1,2}$  identifies with a graph  $\tau_0$  on  $S_0$ . Then applying 3.5 to  $f_0$ , and using the bounds on  $\tau_1$ ,  $\tau_2$ , we can bound  $|f_0(\tau) \setminus A_0|$  and the union of geodesic segment homotopic to  $f_0(\tau) \cap A_0$  via homotopy preserving segment endpoints.

If  $S$  is incompressible, the transfers of lengths are rather similar. 3.5 can be applied directly to transfer lengths between  $f_1$  and  $f_2$ , which makes things slightly easier. Transfer of lengths around common loops is slightly different, in that  $S$  is incompressible and the one or two surfaces from boundary components of components of  $W_{i-1}$  might be compressible or incompressible. Compressible surfaces, as before, have to be decomposed into incompressible subsurfaces in order to apply 3.5.

Now we construct the map  $f : W' \rightarrow N$  which is homotopic to inclusion. We now have a connected graph of geodesic segments, each of which is either has the endpoints of  $f(\gamma)$  for some  $f : S \rightarrow N$  in the family of pleated surfaces for  $W$ , and  $\gamma \subset \Sigma' \cap S$ , or is the homotopy track between  $f_1(x)$  and  $f_2(x)$  for some endpoint  $x$  of a maximal arc in  $\Sigma' \cap S$ , where  $S$  is the domain of  $S_1$  and  $S_2$ . All these geodesic segments have length  $\leq L_1$  for some suitable  $L_1$



which, ultimately, depends only on the topological type of  $W'$ , and the constant  $c_0$ . This graph then gives a 1-complex for the cell-complex homeomorphic to  $W$ , in which the components are the components of  $W_0$ . We refine the cell decomposition to a decomposition into tetrahedra, keeping the same vertices, but adding edges. Again, the corresponding geodesic segments in  $N$  have length bounded in terms of  $L_1$  and the topological type of  $W'$ . Then we map these topological tetrahedra in the topological decomposition of  $W'$  to hyperbolic tetrahedra in  $N$ . This gives the required homotopy equivalence, except on the boundary. On  $S_d(e) \cap \partial W'$  we define  $f = f_{e,0}$ . Since  $|f_{e,0}(\tau)| \leq L_2$  for any arc  $\tau \in S(e) \cap \Sigma'$ , the arc is a bounded distance from the geodesic with the same endpoints, and we can homotope  $f_{e,0}|_{S_d(e)}$  to the map which takes the 1-complex to a union of geodesic segments, by a homotopy with bounded tracks. So putting these together, we obtain a homotopy equivalence with the desired properties.  $\square$

## 9 Combinatorially Bounded geometry case.

### 9.1 Proof of 1.3 in the combinatorial bounded geometry Kleinian surface case.

We now prove our main theorem 1.3 in the case of topological type  $S \times \mathbb{R}$  for a finite type surface  $S$ , with end invariants  $\mu(e_{\pm}) \in \mathcal{GL}_a(S)$  and with combinatorially bounded geometry, and therefore, exactly two ends. The other two main theorems follow from 1.3. We choose pleated surfaces  $f_{\pm} : S \rightarrow N$  approaching the ends, as in 3.11, where the pleating loci include maximal multicurves  $\Gamma_{\pm}$  where  $|f_+(\Gamma_+)| \leq L_0$ , and similarly for  $f_-$ ,  $\Gamma_-$ . We then fix on a choice of loop  $\zeta_{\pm} \in \Gamma_{\pm}$ . As we have seen in 7.16, we can find  $y'_{\pm} = [\varphi'_{\pm}] \in [y_-, y_+]$  such that  $|\varphi'_+(\zeta_+)| \leq L_0$ , similarly for  $y'_-$ ,  $\zeta_-$  and for a suitable  $\nu > 0$ ,

$$[y'_-, y'_+] \subset (\mathcal{T}(S))_{\geq \nu}.$$

For  $z_0 = y'_-$  and  $y_+ = y'_+$ , we then choose a sequence  $\Gamma_j$  of maximal multicurves, as in 8.2, and  $f_j$  a pleated surface with pleating locus including  $\Gamma_j$ , for  $-n_- \leq j \leq n_+$  such that  $|f_{n_+}(\zeta_+)| \leq L_0(\nu)$ , and similarly for  $f_{n_-}$ ,  $\zeta_-$ , and so that 8.1.1 to 8.1.6 are satisfied. We then know by 6.5 that every point on  $[y'_-, y'_+]$  is a bounded distance  $\leq C(\nu)$  from a point on  $[[f_{n_-}], [f_{n_+}]]$ , as noted in 8.2.

By 8.2 we can find a sequence  $f_i : S \rightarrow N$  of pleated surfaces with pleating locus of  $f_i$  including  $\Gamma_i$  for  $-n_- \leq i \leq n_+$ , such that 8.1.1-8.1.6 hold — of which 8.1.5 is redundant in this case.

It is convenient to choose the indexing of the  $z_j$  as  $-n_- \leq i \leq n_+$  — rather than  $0 \leq j \leq n_+$  as in 8.2 — so that for the corresponding sequence  $\{z_i : -n_- \leq i \leq n_+\}$  we have  $z_0$  in a bounded subset of  $\mathcal{T}(\mathcal{S})$ . We know that there is such a point  $z_0 \in [y'_-, y'_+]$  by 7.16. By definition,  $y'_- = z_{-n_-}$  and  $y'_+ = z_{n_+}$ . But at the moment we have no bound on  $d(z_i, [f_i])$  or on  $d(z_i, [f_i])$  for a general  $i$ . But by 8.1.6 we do have, for a suitable  $L_0$ , for all  $p$ ,

$$d([f_p], [f_{p+1}]) \leq L_0.$$

Now we consider  $x(., [y'_-, y'_+])$ . Now we claim that the hypotheses of 6.13 hold for  $[y'_-, y'_+]$  replacing  $[z_-, z_+]$ . So suppose that for some  $p$  and  $i$ , and  $d([f_p], z_i) \leq L$ , and  $d([f_p], z_j) \leq r' + 1$  for  $|i - j| \leq r'$ . By the definition of the sequence in 8.2,  $z_i = [\varphi_i]$  and  $\Gamma_i$  have the properties that, for a constant  $L_0$ ,  $|\varphi_i(\Gamma_i)| \leq L_0$ . So for a constant  $L'_0 = L'_0(L, L_0, r')$ , and  $|j - i| \leq r'$ ,

$$|(\Gamma_j)_*| \leq |f_p(\Gamma_j)| \leq L'_0.$$

So we also have a bound on  $|f_j(\Gamma_j)|$ . But then the bound on  $d([f_p], [f_{p+1}])$  for all  $p$  gives, for a suitable constant  $L''_0 = L''_0(L_0, L, r')$ , for  $|j - i| \leq r'$ ,

$$|f_i(\Gamma_j)| \leq L''_0.$$

If  $r'$  is large enough given  $\nu$ , by 6.2,  $\cup\{\Gamma_j : |i - j| \leq r'\}$  is cell-cutting. So then the same loops are bounded at  $z_i$  as at  $[f_i]$ . So we have a bound on  $d([f_i], z_i)$ . So the hypotheses of 6.13 are satisfied. So then by 6.13, for all  $j$ , and a suitable constant  $L_1$ , we have

$$d([f_j], z_j) \leq L_1. \quad (9.1.1)$$

In particular we have, still for a constant  $L_1$  depending only on  $\nu$ ,

$$|(\Gamma_j)_*| \leq L_1.$$

□

## 9.2 Combinatorially bounded geometry Kleinian surface case: no Margulis tubes.

Fix a basepoint  $w_0 \in N$ . Let  $\Delta_1 > 0$  be given. Then we can choose  $f_\pm$  so that the  $\Delta_1$ -neighbourhood of  $w_0$  is strictly between  $f_-(S)$  and  $f_+(S)$ . To do this, we use the Bounded Diameter Lemma of 3.3. There is  $D_0$  given by the Bounded Diameter Lemma, and an integer  $k_0$  bounded in terms of topological type, such that  $f_{n_+}(S)$  is connected to  $f_+(S)$  by a chain of at most  $k_0$  Margulis tubes and sets of diameter  $\leq D_0$ . The same is true for  $f_-(S)$ ,  $f_{-n_-}(S)$ . So we choose  $f_+$  and  $f_-$  so that  $f_\pm(S)$  cannot be connected to the  $\Delta_1$  neighbourhood of  $w_0$  by such chains. Then we claim that for some  $\varepsilon_1 > 0$  depending only on  $\nu$ , the  $\Delta_1$ -neighbourhood of a fixed basepoint  $w_0 \in N$  does not intersect any  $\varepsilon_1$ -Margulis tube. We see this as follows. For each  $i$ ,  $f_i(S)$  separates  $N$ . In particular, this is true for  $i = n_+$  and  $i = -n_-$ . The point  $w_0 \in N$  is in a component  $N_1$  of the complement in  $N$  of  $f_{n_+}(S) \cup f_{n_-}(S)$  whose boundary meets both  $f_{n_+}(S)$  and  $f_{n_-}(S)$ . The union of the homotopies between  $f_i$  and  $f_{i+1}$  is a homotopy between  $f_{n_-}$  and  $f_{n_+}$ . The basic principle we use is:

### 9.2.1

$N_1$  is contained in the image of the homotopy between  $f_{n_-}$  and  $f_{n_+}$ . Hence each point in  $N_1$  is in the image of a homotopy between  $f_i$  and  $f_{i+1}$  for some  $i$ .

For if this is not true, we have a homotopy equivalence between  $(S \times [0, 1], S \times \{0, 1\})$  and  $S \times [0, 1] \setminus \{x\}, S \times \{0, 1\})$ , for any internal point  $x$ .

But we now know, by the bound on  $d([f_i], z_i)$  that  $[f_i] \in (\mathcal{T}(S))_{\geq \varepsilon_2(\nu)}$  for some constant  $\varepsilon_2(\nu) > 0$  depending only on  $\nu$ . So since  $f_i : S(f_i) \rightarrow N$  is decreasing,  $f_i(S)$  has diameter  $\leq L_1 = L_1(\nu)$ . Also, by 8.1.5, the diameter of the homotopy between  $f_i$  and  $f_{i+1}$  is bounded by  $L_0(\nu)$ , taking  $L_0(\nu)$  sufficiently large. So for a suitable  $\varepsilon_1 = \varepsilon_1(\nu) > 0$ , the image of this homotopy does not intersect any  $\varepsilon_1$ -Margulis tube. So  $N_1$  is disjoint from all  $\varepsilon_1$ -Margulis tubes. Now if  $\Delta_1$  is sufficiently large given  $\nu$ ,  $L'_0$  and  $\Delta_2$ , for  $d(z_0, z_i) \leq \Delta_2$ ,  $|\gamma_*| \geq \varepsilon_1$  for all  $\gamma$  such that  $|\varphi_i(\gamma)| \leq L'_0$ .

### 9.3 Combinatorially bounded geometry Kleinian surface case: lower bounds between pleated surfaces.

So now we consider  $f_i : S \rightarrow N$  for  $i$  such that  $d(z_0, z_i) \leq \Delta_2$ . Then we have bounds on  $d([f_i], [f_{i+1}])$ , on the geometry of  $S(f_i)$  and also on the diameter of the homotopy in  $N$  between  $f_i$  and  $f_{i+1}$ . The following lemma will be applied more generally - always with the bounded geometry assumptions given here. It is proved in [37], essentially the same proof as given here.

**Lemma** *There is an integer  $k_0 = k_0(L_1, \varepsilon, C)$  such that the following holds. Let  $f_i : S \rightarrow N$  ( $-m_- \leq i \leq m_+$ ) be a sequence of homotopic pleated surfaces such that the pleating locus of each  $f_i$  contains a different maximal multicurve  $\Gamma_i$ , such that the following hold.*

1.  $f_i(S) \subset N_1 \subset N_{\geq \varepsilon}$  for all  $i$ .
2.  $(f_i)_* : \pi_1(S) \rightarrow \pi_1(N_1)$  is injective.
3.  $[f_i] \in (\mathcal{T}(S))_{\geq \varepsilon}$  for all  $i$ .
4.  $d([f_i], [f_{i+1}]) \leq L_0$
5.  $|\Gamma_i|_* \leq L_0$ .
6. *There is a homotopy between  $f_i$  and  $f_{i+1}$  with tracks of length  $\leq L_0$  and with image contained in  $N_1$ .*

*Then if  $f_i(S)$  and  $f_j(S)$  come within  $C$ , we have  $|i - j| \geq k_0$ .*

*Proof.* First, note that, given  $\Delta$ , there is  $n(\Delta) = n(\Delta, L_1, \varepsilon)$  such that for any  $i$  with  $m_- + n(\Delta) \leq i \leq m_+ - n(\Delta)$  the distance between  $f_i(S)$  and  $f_j(S)$  is  $> \Delta$  for at least one  $j$  with  $|i - j| < n(\Delta)$ . For if not we have a large number of distinct simple closed geodesics of length  $\leq L_1$  in a set of diameter  $\Delta$  is so large that some nontrivial loop must have length  $< \varepsilon$ , which is a contradiction. Now fix a  $\Delta$  sufficiently large given  $L_1$  and  $C$ .

Now suppose for a sequence of hyperbolic manifolds  $N_p$  all satisfying the same combinatorial bounded geometry condition and with corresponding pleated surface families  $f_i^p$ , that  $f_i^p(S)$  and  $f_j^p(S)$  come within distance  $C$  in  $N_p$  for an arbitrarily large  $|i - j|$  as  $p \rightarrow \infty$ . Then because the distance between  $f_k^p(S)$

and  $f_{k+1}^p(S)$  is bounded by  $L_0$  for all  $k$ , and  $p$ , we can find a sequence  $\Delta_p$  with  $\lim_{p \rightarrow \infty} \Delta_p = +\infty$ , and we can find  $i(p), j(p), k_q = k_q(p)$  for  $1 \leq q \leq 4$  such that

$$i(p) - k_q(p) \leq n(2\Delta), \quad q = 1, 2,$$

$$\lim_{p \rightarrow \infty} |i(p) - j(p)| = +\infty,$$

with  $f_{k_1}^p(S)$  and  $f_{k_2}^p(S)$  on opposite sides of  $f_i^p(S)$  distance  $\geq \Delta$  away from  $f_i^p(S)$ ,  $f_{k_3}^p(S)$  and  $f_{k_4}^p(S)$  on opposite sides of  $f_i^p(S)$  distance  $\geq \Delta_p$  away  $f_i^p(S)$ , but  $f_i^p(S)$  and  $f_j^p(S)$  within  $C$ .

By [21], there is  $h_q^p$  homotopic to  $f_{k_q}^p$  such that  $h_q^p : S \rightarrow N$  is an embedding and  $h_q^p(S)$  is in a small neighbourhood of  $f_{k_q}^p(S)$  and distance  $\geq \Delta$  from  $f_i^p(S)$  for  $q = 1, 2$ . Then by [63], the submanifold of  $N_p$  containing  $f_i^p(S)$  and bounded by  $h_1^p(S)$  and  $h_2^p(S)$  is homeomorphic to  $S \times [0, 1]$ , and similarly for the manifold bounded by  $h_3^p(S)$  and  $h_4^p(S)$ . (To get into the context of [63], we need to consider the horodisc deletion  $S_d$ , but this is a trivial matter.) Composing with an element of the mapping class group  $\text{Mod}(S)$ , we can assume the  $f_i^p$  all lie in a compact set. Then by the bound on  $n(\Delta)$ , the same is true of the  $f_{k_q}^p$  and the  $h_q^p$  for  $q = 1, 2$ . So we can take a geometric limit of  $(N_p, f_i^p(S))$ . The limit of  $f_i^p$  is a pleated surface  $f : S \rightarrow N'$ , where  $N'$  is the limit of  $N_p$ , taking a basepoint in  $f_i^p(S)$ . Then  $f$  is injective on  $\pi_1(S)$ , because if  $f(\gamma)$  is trivial, then for sufficiently large  $p$  given  $\gamma$ ,  $f_i^p(\gamma)$  is freely homotopic to a short loop in  $N_p$  in the region bounded by  $f_{k_3}^p(S)$  and  $f_{k_4}^p(S)$ . But there are no short loops in this region. We can also assume that  $f_{k_1}^p$  and  $f_{k_2}^p$  converge, to pleated surfaces which are homotopic to  $f$ . These limiting surfaces are homotopic to embeddings and to each other, since  $f_{k_1}^p$  and  $f_{k_2}^p$  are, and by [21], there are embeddings homotopic to them with images in arbitrarily small neighbourhoods of  $f_{k_1}^p(S)$ ,  $f_{k_2}^p(S)$ . So then, applying [21] and [63] again, the  $\Delta/2$  neighbourhood of  $f(S)$  in  $N'$  is contained in a submanifold homeomorphic to  $S \times [0, 1]$ , with  $f$  homotopic to  $z \mapsto (z, \frac{1}{2})$  under the homeomorphism. Now let  $\gamma_p$  be a loop in the pleating locus of  $f_j^p$ , which by choice of  $\Delta$  is in the submanifold of  $N'$  homeomorphic to  $S \times [0, 1]$ . Then, taking a subsequence, we can assume that  $f_j^p(\gamma_p)$  converges to a geodesic  $\gamma_*$  in  $N'$  which must be homotopic to  $f(\gamma)$  for some  $\gamma \in S$ , because  $N'$  is homeomorphic to  $S \times [0, 1]$ . Then for large  $p$ , we have two distinct closed geodesics  $\gamma_*$  and  $(\gamma_p)_*$  in  $N_p$  of length  $\leq L_1$  and with distance apart tending to 0 as  $p \rightarrow \infty$ . This is impossible.  $\square$

#### 9.4 Combinatorially bounded geometry Kleinian surface case: construction of biLipschitz map.

For  $\Delta > 0$ , we let  $p_{\pm}$  be the largest integers such that  $d(z_0, z_i) \leq \Delta$  for  $-p_- \leq i \leq p_+$ . Then we construct a map  $\Phi = \Phi_{-p_-, p_+}$  from  $M(z_{-p_-}, z_{p_+})$  to  $N$ . The map  $\Phi$  maps  $S \times \{t_j\}$  to  $f_j(S)$ , if  $d(z_0, z_j) = t_j$  for  $j > 0$  and  $-t_j$  for  $j < 0$ . Since we have a bound on  $d(z_j, [f_j])$  we can also choose  $\Phi$  so that this is boundedly Lipschitz with respect to the Riemannian metric on  $M(z_{-p_-}, z_{p_+})$ . We can then assume without loss of generality that  $\Phi(z, t_j) = f_j(z)$ . The

distance between  $f_j(z)$  and  $f_{j+1}(z)$  in  $N$  is then bounded, because a bounded loop in the homotopy class of a bounded geodesic is a bounded distance from that geodesic. So we can extend  $\Phi$  to  $S \times [t_j, t_{j+1}]$  to map  $z \times [t_j, t_{j+1}]$  to the geodesic between  $f_j(z)$  and  $f_{j+1}(z)$ . Then  $\Phi$  is coarse Lipschitz, since it is so restricted to  $S \times [t_j, t_{j+1}]$  for all  $j$ . It is also coarse biLipschitz, for the following reason. By 9.3, if  $f_i(s)$  and  $f_j(S)$  are a bounded distance apart, then we have a bound on  $|i - j|$ . So we only need to ensure that if  $\tilde{S}$  and  $\tilde{N}$  denote the universal covers of  $S$ ,  $N$ , a sufficiently long path in  $\tilde{S} \times [a, b]$  will map under  $\tilde{\Phi} : \tilde{S} \times [t_{p-}, t_{p+}] \rightarrow \tilde{N}$  to a path which is homotopic, via homotopy preserving endpoints, to a geodesic segment of length  $\geq 1$ . This is true by the same argument as in 9.3. For if this is not true, we can extend length by a bounded amount, and can find a sequence of closed geodesics such that  $|\gamma_p| \rightarrow \infty$ , in the metric on  $S(f_i^p)$ , and yet  $f_i^p(\gamma_p) \subset N_p$  is homotopic to a closed geodesic  $(\gamma_p)_*$  of length  $\leq L_2$ , where  $L_2$  is bounded in terms of  $L_1$ . Normalise as before by applying the modular group, and take a subsequence so that  $f_i^p(\gamma)$  converges for all  $\gamma$ . Since  $(\gamma_p)_*$  lies in a fixed compact set it must be  $\gamma_*$  for some fixed  $\gamma$ , for all sufficiently large  $p$ . But this again contradicts Injectivity-on- $\pi_1$ .

So  $\Phi_{-p-, p+}$  is uniformly coarse biLipschitz. The definition restricted to  $S \times [t_j, t_{j+1}]$  depends only on  $j$ , not on  $p_{\pm}$ . Taking limits as  $\Delta \rightarrow \infty$ , that is as  $p_+ \rightarrow +\infty$  and  $-p_- \rightarrow -\infty$ , we obtain a coarse biLipschitz map  $\Phi : M(\mu_-, \mu_+) \rightarrow N$ , which is onto, because for  $p_{\pm}$  sufficiently large given  $\Delta_1$ , the image contains the  $\Delta_1$ -neighbourhood of  $w_0 \in N$ .

□

## 10 Lipschitz bounds.

In this section, we do the groundwork for the construction of a biLipschitz map from model  $M$  to hyperbolic manifold  $N$  in complete generality. We follow the same general strategy as in the Kleinian surface case with combinatorial bounded geometry, which was carried out in Section 9. The main theorem from which Lipschitz bounds are deduced is 10.1 which is an analogue of the work carried out in 9.1. To simplify the statement, we give two versions, the interval bundle case and then the general case. Note, in particular, the resemblance between (9.1.1) and (10.1.2). Theorem 10.1 does, of course, take more work than 9.1. Theorem 10.2 is the start of an induction to prove it. The two theorems are proved in 10.3 to 10.6. There is also an analogy between 9.2 and 10.8, the respective bounds on Margulis tubes. In this general case, there are, of course, Margulis tubes. 10.9 gives a relationship between the geometry of model Margulis tubes in  $M$  and Margulis tubes in  $N$ . The rest of the section is the basic work which will be needed to get a map from  $M$  to  $N$  which is coarse biLipschitz, as well as Lipschitz, moreover, a map for which the biLipschitz constants will be locally uniform in the ending lamination data. 10.13 is a fairly straight analogue of 9.3.

Throughout this section, we fix a compact connected 3-dimensional submanifold-with-boundary  $W$  of  $N_c$  as in 7.6 and as in the introduction to

Section 8. As before, we define  $N_{d,W}$  to be the union of  $N_d$  and of components of  $N \setminus N_d$  disjoint from  $W$ . The conditions are such that each end  $e$  of  $N_{d,W}$  has a neighbourhood in  $N_{d,W} \setminus W$  bounded by a component  $S_d(e)$  of  $W \setminus \partial N_d$ , and  $S_d(e)$  is the horodisc deletion of a finite type surface  $S(e)$ . We denote by  $\overline{N}$  the closure of  $N$  in the projection of  $H^3 \cup \Omega$ , where  $\Omega$  is the domain of discontinuity of the covering group on  $\partial H^3$ .

**Theorem 10.1.** *For each end  $e$  of  $N_{d,W}$ , let  $f_{e,+} : S(e) \rightarrow \overline{N}$  be a generalised pleated surface as in 4.9, and homotopic to inclusion of  $S(e)$  in  $N$ . Let  $\Gamma_+(e)$  be a maximal multicurve which includes all closed loops in the pleating locus of  $f_{e,+}$ , and suppose that*

$$|f_{e,+}(\Gamma_+(e))| \leq L_0. \quad (10.1.1)$$

*Let  $z_{e,0} = [\varphi_{e,0}] \in \mathcal{T}(S)$ . Write  $y_{e,+} = [f_{e,+}]$ . If  $\Gamma_+(e)$  includes some loops which are cusps in  $N$ , let  $y_{e,+}$  be defined as an element of  $\mathcal{T}(S(e))$  so as to minimise  $d(z_{e,0}, y_{e,+})$  up to an additive constant, as in 8.4.*

*Fix ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$  and a vertically efficient ltd-bounded decomposition of  $S(e) \times [z_{e,0}, y_{e,+}]$  with respect to these. Then the following holds for a constant  $L_1$ , and for sufficiently strong ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$ , where  $L_1$  depends only on the parameter functions and  $L_0$ .*

*Interval Bundle Case.*

*Denote the ends by  $e_{\pm}$  and  $f_{\pm} = f_{e_{\pm},+}$ ,  $y_{\pm} = [f_{\pm}]$ ,  $\Gamma_+(e_{\pm}) = \Gamma_{\pm}$ . Also, in this case,  $z_{e_{+},0} = y_{-}$*

*For any  $(\alpha, \ell)$  in the decomposition of  $S(e) \times [y_{-}, y_{+}]$ , there is a pleated surface  $f : \alpha \rightarrow N$  whose pleating locus includes  $\partial\alpha$ , such that*

$$|f(\partial\alpha)| \leq L_1,$$

$$d_{\alpha}([f], y) \leq L_1 \text{ or } |\operatorname{Re}(\pi_{\alpha}([f]) - \pi_{\alpha}(y))| \leq L_1, \quad (10.1.2)$$

*depending on whether  $\alpha$  is a gap or a loop.*

*General case. Suppose that there is a constant  $c_0 > 0$  with the following property.*

*Whenever  $e$  is a compressible end,  $\gamma'$  is a simple closed loop which is nontrivial in  $S(e)$  but trivial in  $N$ , and  $\mu$  and  $\mu'$  are geodesic laminations on  $S$ , with a normalised transverse invariant measure on  $\mu$ ,*

$$i(\mu, \mu') = 0 \text{ and } i(\mu', \gamma') \leq c_0 |\gamma|,$$

*then for at least one  $\zeta \in \Gamma_+(e)$ ,*

$$i(\zeta, \mu) \geq c_0 |\zeta|. \quad (10.1.3)$$

*Compressible ends. Let  $e$  be a compressible end. Then (10.1.2) holds as above, but with  $[z_{e,0}, y_{e,+}]$  replacing  $[y_{-}, y_{+}]$ , but for a constant  $L_1$ , which, this time, depends, in addition,  $c_0$  and locally on  $z_{e,0}$ .*

*Incompressible ends. Let  $e$  be an incompressible end. Then (10.1.2) holds as in the Interval Bundle case but with  $[z_{e,0}, y_{e,+}]$  replacing  $[y_{-}, y_{+}]$ , where  $z_{e,0}$*

is chosen relative to the set of all  $y_{e',+}$  as described in 7.14, and also with  $[\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$  replacing  $[y_-, y_+]$ , whenever  $\omega = \omega(e, e')$  is the maximal (possibly empty) subsurface of  $S(e)$  which is homotopic to a subsurface  $\omega(e', e)$  of  $S(e')$ , and  $z'_{e,0}, z'_{e',0}$  are as in 7.14.

In the combinatorial bounded geometry Kleinian surface case, the interval bundle version of 10.1, is Theorem 9.1. We shall want to apply the interval bundle case of the theorem to a sequence of manifolds, all of the same interval bundle type. We shall want to apply the general case of the theorem to a sequence of manifolds of the same topological type with the same choices of  $L_0, z_{e,0}, z_{e',0}$  all along the sequence, and of  $c_0$  in the presence of compressible ends, but with  $\Gamma_+(e)$  varying. We shall then use the theorem and the follow-up results to obtain uniform biLipschitz bounds along the sequence. We shall deal later with the question of satisfying the hypotheses of this theorem.

Theorem 9.1 was proved using the first lemma of 6.13. The idea of the proof in this general case is to use the second lemma of 6.13 to start an inductive proof. The induction of 10.1 is started by proving the following theorem, which is a subset of 10.1.

**Theorem 10.2.** *Let  $f_\pm : S \rightarrow N$  be homotopic generalised pleated surfaces. Let  $\Gamma_\pm$  be maximal multicurves containing all closed loops in the pleating loci of  $\Gamma_\pm$  and let  $|f_+(\Gamma_+)| \leq L_0, |f_-(\Gamma_-)| \leq L_0$ . Let  $\omega \subset S$  be a subsurface with  $\partial\omega$  in the pleating locus of  $f_\pm$  with*

$$|(\partial\omega)_*| \leq L_0.$$

*Let  $(\alpha_i, \ell_i)$  ( $1 \leq i \leq R_0$  or  $1 \leq i \leq R_{e,0}$ ) be a totally chain of ltds for  $\omega \times [y_-, y_+]$ . Thus, the  $(\alpha_i, \ell_i)$  are as in 6.14, with  $d_\omega$ -distance  $\leq \Delta_0$  between the end of  $\ell_i$  and the start of  $\ell_{i+1}$ , for a fixed constant  $\Delta_0$  depending only on the ltd parameter functions*

*Incompressible surface case Let  $f_\pm$  be injective on  $\pi_1$ .*

*Compressible surface case Let  $S = S(e)$  for a compressible end. Let  $\Gamma_+$  satisfy (10.1.3). Suppose that for  $z_{e,0}$  as in 10.1 and for  $x(\cdot, \cdot)$  as in 6.11,*

$$d(z_{e,0}, x(z_{e,0}, [[f_-], [f_+]])) \geq D_2.$$

*Then (10.1.2) holds simply for  $(\alpha, \ell) = (\alpha_i, \ell_i)$ ,  $1 \leq i \leq R_0$ , and for a constant  $L_{1,1}$  replacing  $L_1$ .*

### 10.3 Outline proof of 10.2 in the Interval Bundle Case.

The idea is to use the second lemma of 6.13. The difficulty, as was pointed out at the time, compared with the first lemma of 6.13, is that, although we can get good bounds straight away at some points, we cannot then proceed immediately to other points, because a bound on  $d_{\alpha_j}(x(y_j), y_j)$  for gaps  $\alpha_j$  does not imply a bound on  $d(y_j, x(y_j))$ . So the idea, given  $(\alpha_{i_0}, \ell_{i_0})$  and  $y \in \ell_{i_0}$ , is to get good bounds progressively closer to  $y$ , successively using 6.13, modifying points to

get into a position to apply 6.13 again. For the moment, we concentrate on the interval bundle case. We shall consider the general case later.

So the aim is to find an integer  $m_1$ , and constants  $L_{1,1,1}$ ,  $L_{1,1,2}$  depending only on the  $(\Delta_1, r_1, s_1, K_1)$  and constant  $\kappa_1$  of 8.1.6, and finite sequences of geodesic segments  $[y_{m,-}, y_{m,+}]$ ,  $[z_{m,-}, z_{m,+}]$ , and sequences of ltd's  $(\alpha_{m,\pm}, \ell_{m,\pm})$  ( $0 \leq m \leq m_1$ ) with  $y_{0,-} = y_- = z_{0,-}$ ,  $z_{0,+} = y_{0,+} = y_+$ ,  $y_{m,\pm} = [f_{m,\pm}]$ , where  $f_{m,\pm}$  is a pleated surface, and with the following properties. We assume without loss of generality that  $f_+ = f_-$  off  $\omega$ .

1. Either  $\alpha_{m,+} = \omega$  and  $y_{m,+} = y_+$  or, for some  $n$ ,  $\alpha_{m,+} = \alpha_n$   $\ell_{m,+}$  is a segment of  $\ell_n$  including the left endpoint,  $z_{m,+}$  is the right end of  $\ell_{m,+}$ . The pleating locus of  $f_{m,+}$  includes  $\partial\omega \cup \partial\alpha_{m,+}$ . Similar properties hold for  $\alpha_{m,-}$ ,  $y_{m,-}$ ,  $f_{m,-}$ ,  $z_{m,-}$ . For all  $m$ ,  $f_{m,+} = f_{m,-} = f_+$  off  $\omega$ .

2.

$$y \in [z_{m,-}, z_{m,+}].$$

3.

$$\begin{aligned} |\varphi_{m,+}(\partial\alpha_{m,+})| &\leq L_{1,1,1} \text{ and, if } y_{m,+} \neq y_+, \\ d_{\alpha_{m,+}}(y_{m,+}, z_{m,+}) &\leq L_{1,1,1} \text{ or} \\ |\operatorname{Re}(\pi_{\alpha_{m,+}}(y_{m,+}) - \pi_{\alpha_{m,+}}(z_{m,+}))| &\leq L_{1,1,1}, \end{aligned} \quad (10.3.1)$$

depending on whether  $\alpha_{m,+}$  is a gap or a loop, and similarly with  $+$  replaced by  $-$ , and left and right interchanged, where  $L_{1,1,1}$  depends only on  $(\Delta_1, r_1, s_1, K_1)$ . Also, if  $\alpha_{m,+}$  is a gap, the pleating locus of  $f_{m,+}$  in  $\alpha_{m,+}$  is a maximal multicurve  $\Gamma_{m,+}$  with  $|\varphi_{m,+}(\Gamma_{m,+})| \leq L_{1,1}$  where  $[\varphi_{m,+}] = z_{m,+}$ , and similarly for  $f_{m,-}$ ,  $\alpha_{m,-}$ ,  $z_{m,-}$ ,  $y_{m,-}$ , if  $y_{m,-} \neq y_-$ .

4. The following two cases depend on whether or not the hypotheses of the second lemma of 6.13 are satisfied.

**Case 1.** If

$$d(y_{m,-}, y_{m,+}) \leq 2d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+})$$

then either  $\alpha_{m+1,+} = \alpha_{m,+}$  and  $f_{m+1,+} = f_{m,+}$  on  $\alpha_{m,+}$  or

$$d'_{\alpha_{m+1,+}, \alpha_{m,+}}(y_{m+1,+}, y_{m,+}) \geq \frac{1}{8}d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+}),$$

and similarly with  $+$  replaced by  $-$ , and

$$d'_{\alpha_{m+1,-}, \alpha_{m+1,+}}(y_{m+1,-}, y_{m+1,+}) \leq \frac{7}{8}d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+}), \quad (10.3.2)$$

and

$$d(y_{m+1,-}, y_{m+1,+}) \leq 2\kappa_1 d(y_{m,-}, y_{m,+}). \quad (10.3.3)$$

**Case 2.** If

$$d(y_{m,-}, y_{m,+}) \geq 2d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+})$$



then  $\alpha_{m+1,+} = \alpha_{m,+}$ ,  $f_{m+1,+} = f_{m,+}$  on  $\alpha_{m,+}$ ,  $\alpha_{m+1,-} = \alpha_{m,-}$ ,  $f_{m+1,-} = f_{m,-}$  on  $\alpha_{m,-}$  and

$$d(y_{m+1,-}, y_{m+1,+}) \leq \frac{3}{4}d(y_{m,-}, y_{m,+}). \quad (10.3.4)$$

5.

$$d(y_{m_1,-}, y_{m_1,+}) \leq L_{1,1,2}^3,$$

but for  $m < m_1$

$$d(y_{m,-}, y_{m,+}) \geq L_{1,1,2}^2.$$

If we can satisfy 1-4 then we can obviously find  $m_1$  such that 5 holds, because 4 implies that

$$d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+})$$

is nonincreasing in  $m$  and eventually decreasing. Then we obtain (10.1.2) as follows. Choose cell-cutting loop sets  $\Gamma_{m_1,+} \subset \alpha_{m_1,+}$  and  $\Gamma_{m_1,-} \subset \alpha_{m_1,-}$  such that  $|f_{m_1,+}(\Gamma_{m_1,+})| \leq L_{1,1,3}$  (for  $L_{1,1,3}$  depending on  $L_{1,1,1}$ ,  $L_{1,1,2}$ ), and similarly for  $f_{m_1,-}$ ,  $\alpha_{m_1,-}$ ,  $\Gamma_{m_1,-}$ . Because of the bound on  $d([f_{m_1,-}], [f_{m_1,+}]) = d(y_{m_1,-}, y_{m_1,+})$ , we have

$$|f_{m_1,+}(\Gamma_{m_1,-} \cup \partial\alpha_{m_1,-})| \leq L_{1,1,4}$$

for  $L_{1,1,4}$  depending on  $L_{1,1,i}$ ,  $i = 1, 2, 3$ . By 6.2 and 6.3,  $\alpha$  is in the convex hull of  $\alpha_{m_1,+}$  and  $\alpha_{m_1,-}$ . By the bound on  $d(y_{m_1,-}, y_{m_1,+})$  and (10.3.1), if we take a maximal multicurve  $\Gamma'$  on  $\alpha$  such that  $|\varphi(\Gamma')| \leq L_0$ , each loop of  $\Gamma'$  can be made out of a finite number of arcs of

$$\Gamma_{m_1,+} \cup \partial\alpha_{m_1,+} \cup \Gamma_{m_1,-} \cup \partial\alpha_{m_1,-}.$$

So then we have a bound on  $|f_{m_1,+}(\Gamma')|$ . Then we have a bound on  $|(\Gamma')_*|$ . By 4.1, we can assume that  $|\gamma_*|$  is bounded from 0, keeping the property that  $|\varphi(\Gamma')| \leq L_0$ , for  $L_0$  depending only on the topological type. Extend  $\Gamma'$  to a maximal multicurve on  $S$  with a bound on number of intersections with  $\Gamma_{m_1,+}$ , depending on  $L_{1,1,4}$ . Let  $f$  have pleating locus  $\alpha$ . We have bounds on  $d_\alpha([f_{m_1,+}], [f])$ , and hence on  $d_\alpha([f], y)$  or  $|\operatorname{Re}(\pi_\alpha([f]) - \pi_\alpha(y))|$ , (depending on whether  $\alpha$  is a gap or a loop), and on  $|f(\Gamma)|$ .

It is crucial to the construction that Case 2 can occur for many successive  $m$  with no adverse effect on  $d_{\alpha_{m,+}}(y_{m,+}, z_{m,+})$ , because  $z_{m,+}$  is constant for a string of such  $m$ , and by the observation at the end of 3.1,  $d_{\alpha_{m,+}}(y_{m,+}, z_{m,+})$  then depends only on  $f_{m,+}|\alpha_{m,+}$ : the change in  $\pi_{\alpha_{m,+}}([f_{m,+}])$  will be slightly affected by  $f_{m,+}|S \setminus \alpha_{m,+}$ , but only by a uniformly bounded amount.

## 10.4 Construction of $y_{m+1,\pm}$ from $y_{m,\pm}$ .

For the moment we are considering the Interval Bundle Case. We need to consider Cases 1 and 2 of 10.3. We choose ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$  such that all the results of Section 6 hold. Let  $\Gamma_i, f_i$  ( $0 \leq i \leq n_+$ ) be the

maximal multicurve and pleated surface sequences constructed in 8.3, and thus satisfying 8.1.1-8.1.6.

We proceed to the construction of  $y_{m+1,\pm}$  from  $y_{m,\pm}$ . First we consider Case 1.

Let  $(\alpha, y)$  be as in the statement of 10.1 and write  $y = [\varphi]$ . Write  $y_{m,0}$  for the point in  $[y_{m,-}, y_{m,+}]$  such that

$$|\varphi(\partial\alpha)| \leq C(\nu) \text{ or } |\varphi(\alpha)| \leq L_{1,1,1},$$

$$d_\alpha(y_{m,0}, y) \leq C(\nu), \text{ or } |\operatorname{Re}(\pi_\alpha(y_{m,0}) - \pi_\alpha(y))| \leq L_{1,1,1},$$

depending on whether  $\alpha$  is a gap which is long  $\nu$ -thick and dominant or a loop which is  $K_1$ -flat. Such a point  $y_{m,0}$  exists by 6.5, assuming that  $L_{1,1,2}$  is sufficiently large given the ltd parameter functions. Here,  $L_{1,1,1}$  plays the role of  $L_1$  in 6.4 and 6.5, and  $L_{1,1,2}$  plays the role of  $L_2$  in 6.5. Also by 6.5, there are segments  $\ell'_i \subset [y_{m,0}, y_{m,+}]$  a bounded  $d_{\alpha_i}$ -distance from  $\ell_i$  (or similarly if  $\alpha_i$  is a loop: precise statment in 6.5) for all  $(\alpha_i, \ell_i)$  with  $(\alpha, y_{m,0}) \leq (\alpha_i, \ell_i) \leq (\alpha_{m,+}, y_{m,+})$ , except within distance  $L_{1,1,2}$  of the endpoints, assuming  $L_{1,1,2}$  sufficiently large. Then by 6.6 and the properties of the  $(\alpha_i, \ell_i)$  we have, for  $\delta_0$  depending only on  $\Delta_0$  of 6.14 and the ltd parameter functions,

$$\delta_0 d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+}) \leq \sum_k |\ell'_k \cap [y_{m,-}, y_{m,+}]|,$$

$$\delta_0 d'_{\alpha_{m,0}, \alpha_{m,+}}(y_{m,0}, y_{m,+}) \leq \sum_k |\ell'_k \cap [y_{m,0}, y_{m,+}]|,$$

$$\delta_0 d'_{\alpha_{m,-}, \alpha_{m,0}}(y_{m,-}, y_{m,0}) \leq \sum_k |\ell'_k \cap [y_{m,-}, y_{m,0}]|.$$

Now we assume without loss of generality that

$$\sum_k |\ell'_k \cap [y_{m,0}, y_{m,+}]| \geq \frac{1}{3} \sum_k |\ell'_k \cap [y_{m,-}, y_{m,+}]|,$$

because we have either this, or the corresponding statement with  $+$  and  $-$  interchanged. In this case, we take  $\alpha_{m+1,-} = \alpha_{m,-}$ ,  $y_{m+1,-} = y_{m,-}$ . Then we apply the second lemma of 6.13 with  $x_1 = y_{m,0}$ , and  $x_2 \in \ell_j$  with  $(\alpha, x_1) < (\alpha_j, x_2) \leq (\alpha_{m,+}, y_{m,+})$ , and

$$\frac{1}{4} \sum_k |\ell'_k \cap [y_{m,-}, y_{m,+}]| \leq \sum_k |\ell'_k \cap [x_1, x_2]|.$$

In order to apply this strictly, we shall need to interpolate extra points between  $[f_k]$  and  $[f_{k+1}]$  where  $\Gamma_k$  and  $\Gamma_{k+1}$  are related by a possibly unbounded Dehn twist rather than  $\#(\Gamma_k \cap \Gamma_{k+1})$  being bounded. In this case we shall need to use some of the surfaces  $f_{k,m}$  between  $f_k$  and  $f_{k+1}$  as in 8.1.6, to keep the property of successive  $d(y_j, y_{j+1})$  being bounded, which is the hypothesis of 6.13.

Then we can find  $y_i = [f_i]$  and  $w \in \ell_j \cap [x_1, x_2]$  satisfying the conclusion of the lemma in 6.13. Then  $\alpha_{m+1,+} = \alpha_j$  and  $y_{m+1,+} = y_i$ ,  $z_{m+1,+} = w$ ,  $f_{m+1,+} = f_i$ ,  $\Gamma_{m+1,+} = \Gamma_i$  satisfy the conditions of 10.3, in particular (10.3.1). If  $\alpha_j$  is a gap, which is long,  $\nu$ -thick and dominant along  $\ell_j$ , then we get the bound on  $|(\Gamma_i)_*|$  in the same way as in Section 9. That is, we have a bound on  $d_{\alpha_j}(z_k, w)$  for some  $z_k \in \ell_j$  as in 8.3, and hence a bound on  $d_{\alpha_j}(y_i, z_k)$ . This means that we get a bound on  $|f_i(\Gamma_{k'})|$  in terms of  $\nu$  and  $p = p(\nu)$  for  $k'$  in an interval  $I$  of  $\leq p(\nu)$  integers containing  $k$ . If  $\Delta_1(\nu)$  grows sufficiently fast with  $\nu$ , as we can assume, then we can choose such an interval  $I$  with  $z_{k'} \in \ell_j$  for all  $k' \in I$ , and so that  $\cup_{k' \in I} \Gamma_{k'}$  cuts  $\alpha_j$  into cells and annuli parallel to the boundary. The bounds on  $|f_i(\Gamma_{k'})|$  for all  $k' \in I$  then give a bound, depending on  $\nu$ , on  $d_{\alpha_j}([f_k], [f_i])$ . We can then assume that  $f_i = f_k$ . We get condition 10.3.3 by the properties of the  $f_i$ , in particular 8.1.4 and 8.1.6

Now we consider case 2.

The point  $y_{m+1,\pm}$  will be obtained from a sequence  $y_{m,i,\pm}$  with  $y_{m,0,\pm} = y_{m,\pm}$ . Also,  $y_{m,i,\pm} = [f_{m,i,\pm}]$  for a pleated surface  $f_{m,i,\pm}$  with  $f_{m,0,\pm} = f_{m,\pm}$ . For each  $i$ , we shall have either  $y_{m,i,-} = y_{m,i+1,-}$  or  $y_{m,i,+} = y_{m,i+1,+}$ , but the other one will be different. There will be a finite increasing sequence of subsurfaces  $\alpha_{m,i,\pm}$  ( $i \geq 0$ ) such that the pleating locus of  $f_{m,i,+}$  includes  $\alpha_{m,i,+}$ , and either  $\alpha_{m,i+1,+}$  is strictly larger than  $\alpha_{m,i,+}$ , or  $\alpha_{m,i+1,-}$  is strictly larger than  $\alpha_{m,i,-}$ . Then for some  $r \leq -2\chi(S)$ , where  $\chi$  denotes Euler characteristic, we have  $\alpha_{m,r,\pm} = S$  and  $f_{m+1,\pm} = f_{m,r,\pm}$ . We shall always have  $f_{m,i+1,+} = f_{m,i,+}$  on  $\alpha_{m,i,+}$ , and similarly with  $+$  replaced by  $-$ . For  $i < r$   $\alpha_{m,i,+}$  will be a union of  $\beta$  such that  $(\beta, \ell')$  is an ltd for  $[y_{m,-}, y_{m,+}]$ .

For a suitable  $L_{1,1,5}$ , we shall always take

$$\begin{aligned} d(y_{m,i+1,-}, y_{m,i+1,+}) &\leq d(y_{m,i,-}, y_{m,i,+}) + L_{1,1,5} \\ &\leq d(y_{m,-}, y_{m,+}) + (i+1)L_{1,1,5}, \end{aligned}$$

and

$$\begin{aligned} &d'_{\alpha_{m,i+1,-}, \alpha_{m,i+1,+}}(y_{m,i+1,-}, y_{m,i+1,+}) \\ &\leq \text{Max} \left( d'_{\alpha_{m,i,-}, \alpha_{m,i,+}}(y_{m,i,-}, y_{m,i,+}), \frac{1}{2}d(y_{m,i,-}, y_{m,i,+}) \right) \\ &\leq \text{Max} \left( d'_{\alpha_{m,-}, \alpha_{m,+}}(y_{m,-}, y_{m,+}), \frac{2}{3}d(y_{m,-}, y_{m,+}) \right) \\ &\leq \frac{2}{3}d(y_{m,-}, y_{m,+}). \end{aligned}$$

We then take  $y_{m+1,+} = y_{m,r,+}$  and  $y_{m+1,-} = y_{m,r,-}$  for the first  $r$  such that

$$\frac{8}{9}d(y_{m,r,-}, y_{m,r,+}) \leq d'_{\alpha_{m,r,-}, \alpha_{m,r,+}}(y_{m,r,-}, y_{m,r,+}) \leq \frac{2}{3}d(y_{m,-}, y_{m,+}). \quad (10.4.1)$$

This will certainly be true if  $\alpha_{m,r,+} = S$ , which happens for some  $r \leq -2\chi(S)$ , but may happen earlier. This will ensure (10.3.4), as required, provided that  $L_{1,1,5}$  is sufficiently large given  $L_{1,1,2}$  and  $\chi(S)$ .

Now we need to consider how to define  $y_{m,i+1,\pm}$  and  $\alpha_{m,i+1,\pm}$  when (10.4.1) does not hold for  $i$  replacing  $r$ . Then we have two cases to consider: when

$$d'_{\alpha_{m,i,-}, \alpha_{m,i,+}}(y_{m,i,-}, y_{m,i,+}) \leq d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+}) - \frac{1}{18}d(y_{m,i,-}, y_{m,i,+}), \quad (10.4.2)$$

and the case when (10.4.2) does not hold, but

$$d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+}) \leq d(y_{m,i,-}, y_{m,i,+}) - \frac{1}{18}d(y_{m,i,-}, y_{m,i,+}). \quad (10.4.3)$$

If (10.4.2) holds, we take  $y_{m,i+1,-} = y_{m,i,-}$ , and need to define  $y_{m,i+1,+}$ . So now consider the case of (10.4.2) holding. The case of (10.4.3) is similar, with  $S$  replacing  $\alpha_{m,i,-}$  and  $\alpha_{m,i,-}$  replacing  $\alpha_{m,i,+}$ .

We use the geodesic segment  $[y_{m,i,-}, y_{m,i,+}]$ . We consider the set  $\mathcal{A}$  of all  $(\beta, \ell)$  such that  $\ell \subset [y_{m,i,-}, y_{m,i,+}]$  and

$$(\alpha_{m,i,-}, y_{m,i,-}) < (\beta, \ell), \quad \beta \cap \alpha_{m,i,+} = \emptyset.$$

If  $(\beta_1, \ell'_1) < (\beta_2, \ell'_2)$  and  $(\beta_1, \ell'_1) \in \mathcal{A}$  then  $(\beta_2, \ell'_2) \in \mathcal{A}$  also, by 6.3 applied to each of the triples

$$\{(\alpha_{m,i,-}, y_{m,i,-}), (\beta_1, \ell'_1), (\beta_2, \ell'_2)\}, \quad \{(\beta_1, \ell'_1), (\beta_2, \ell'_2), (\alpha_{m,i,+}, y_{m,i,+})\}.$$

Now for  $z \in [y_{m,i,-}, y_{m,i,+}]$ , let  $C(z)$  denote the convex hull (5.7) of the  $\beta$  with  $(\beta, \ell') \in \mathcal{A}$  for some  $\ell'$  with  $\ell' \cap [y_{m,i,-}, z] \neq \emptyset$ . Then  $C(z)$  increases with  $z$  and takes only finitely many values. Let  $z_{m,i,j,+}$  be the successive points on  $[y_{m,i,-}, y_{m,i,+}]$  such that  $C(z) = \omega_{m,i,j,+}$  is constant for  $z \in [z_{m,i,j-1,+}, z_{m,i,j,+}]$  with  $z_{m,i,0,+} = y_{m,i,-}$ ,  $z_{m,i,t,+} = y_{m,i,+}$ ,  $t = t(m, i, +)$ , where  $t$  is bounded by  $-2\chi(S)$ . Suppose that  $(\beta, \ell')$  is ltd for  $\ell' \subset [z_{m,i,j-1,+}, z_{m,i,j,+}]$  and  $\beta \cap \omega_{m,i,j,+} \neq \emptyset$ . Then  $\beta \cap \beta' \neq \emptyset$  for any  $(\beta', \ell'') \in \mathcal{A}$  with  $\ell'' \subset [z_{m,i,j-1,+}, z_{m,i,j,+}]$ . There is at least one such  $(\beta', \ell'')$  with  $(\beta', \ell'') \leq (\beta, \ell')$ . So  $(\beta, \ell') \in \mathcal{A}$  by the order-closure property, and hence  $\beta \subset \omega_{m,i,j,+}$ . It follows that for a constant  $L$  depending only on the ltd parameter functions,  $|\varphi(\partial\omega_{m,i,j,+})| \leq L$  for all  $[\varphi] \in [z_{m,i,j-1,+}, z_{m,i,j,+}]$ . Then, by (10.4.2),

$$d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+}) \geq \frac{1}{18}d(y_{m,i,-}, y_{m,i,+}).$$

Now

$$\begin{aligned} & d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+}) \\ &= \text{Max}(d'_{\alpha_{m,i,-}, \alpha_{m,i,+}}(y_{m,i,-}, y_{m,i,+}), d'_{\alpha_{m,i,-}, S \setminus \alpha_{m,i,+}}(y_{m,i,-}, y_{m,i,+})) + O(1) \\ &= d'_{\alpha_{m,i,-}, S \setminus \alpha_{m,i,+}}(y_{m,i,-}, y_{m,i,+}) + O(1). \end{aligned}$$

Using (10.4.2) for the last equality. So  $d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+})$  is bounded above and below, up to an additive constant depending only on the ltd parameter functions, by

$$\sum_{j=1}^t d_{\omega_{m,i,j,+}}(z_{m,i,j-1,+}, z_{m,i,j,+}).$$

So for some  $j$  and  $\lambda_1 = (.01)/(-\chi(S))$ , we have

$$d_{\omega_{m,i,j,+}}(z_{m,i,j-1,+}, z_{m,i,j,+}) \geq \frac{1}{20} \lambda_1 d(y_{m,i,-}, y_{m,i,+}),$$

$$\sum_{k=1}^{j-1} d_{\omega_{m,i,k,+}}(z_{m,i,k-1,+}, z_{m,i,k,+}) < \frac{1}{3} d(y_{m,i,-}, y_{m,i,+}).$$

Now by 6.8, the ltd's  $(\beta, \ell')$  along  $[z_{m,i,j-1,+}, z_{m,i,j,+}]$  with  $\beta \subset \omega_{m,i,j,+}$  are in natural correspondence with ltd's for  $[\pi(z_{m,i,j-1,+}), \pi(z_{m,i,j,+})]$  where  $\pi = \pi_{\omega_{m,i,j,+}}$ . So then we can apply 6.14 to  $[\pi(z_{m,i,j-1,+}), \pi(z_{m,i,j,+})]$ , to obtain a totally ordered sequence of ltd's, and then by 6.8 we have a corresponding totally ordered subset along  $[z_{m,i,j-1,+}, z_{m,i,j,+}]$ . We construct a new family of pleated surfaces for  $[y_{m,i,-}, y_{m,i,+}]$  as in 8.3, with the properties of 8.1. The hypotheses of the lemma in 6.13 holds for  $y_{\pm} = y_{m,i,\pm}$ ,  $x_1 = z_{m,i,j-1,+}$  and  $x_2 = z_{m,i,j,+}$ . Then as in case 1, we can find  $z \in [z_{m,i,j-1,+}, z_{m,i,j,+}]$  and  $\zeta$  ltd along a segment containing  $z$ ,  $\zeta \subset \omega_{m,i,j,+}$  and  $w = [f]$  a pleated surface whose pleating locus includes  $\partial\zeta$ , such that

$$d'_{\alpha_{m,i,-},\zeta}(y_{m,i,-}, z) < \frac{1}{2} d'_{\alpha_{m,i,-}}(y_{m,i,-}, y_{m,i,+}),$$

and (10.3.1) holds, with  $w$  replacing  $y_{m,+}$ ,  $z$  replacing  $z_{m,+}$  and  $\zeta$  replacing  $\alpha_{m,+}$ . Then we put  $\alpha_{m,i+1,+} = \alpha_{m,i,+} \cup \zeta$  and define  $f_{m,i+1,+} = f_{m,i,+}$  on  $\alpha_{m,i,+}$  and  $= f$  on  $S \setminus \alpha_{m,i,+}$ , which includes  $\zeta$ .  $\square$

## 10.5 Proof of 10.1 in the interval bundle case.

We shall now prove 10.1 by induction. We again restrict to the interval bundle case. We shall show that the statement of 10.1 holds for all  $(\alpha, \ell)$  in the decomposition, by induction on the topological type of  $S$ . In fact, in order to carry out the induction, we shall prove something a bit more general than 10.1. Fix a subsurface  $\omega$  of  $S$  such that all components of  $\partial\omega$  are nontrivial and  $|(\partial\omega)_*| \leq L_0$ . As usual, given a nontrivial nonperipheral loop  $\gamma \subset S$ ,  $\gamma_*$  denotes the geodesic representative in  $N$ , up to free homotopy. We shall show that 10.1 holds if  $f_{\pm} : S \rightarrow N$  are pleated surfaces whose pleating locus includes  $\partial\omega$ , and  $\alpha \subset \omega$  if  $\omega$  is a simple loop on  $S$ , then the theorem is trivially true. So now suppose inductively that 10.1 holds with  $S$  replaced by any proper essential subsurface  $\omega'$  properly contained in  $\omega$ , for any choice of  $f_{\pm}$  and  $z_0 = y_-$ . Now let  $N$  and  $f_{\pm} : S \rightarrow N$  be given, where  $\partial\omega$  is in the bending locus of both  $f_{\pm}$  and  $|(\partial\omega)_*| \leq L_0$ . By 6.14, and 6.8 with  $\alpha$  replaced by  $\omega$ , there is at least one sequence of  $(\ell_i, \alpha_i)$  as in 6.14, with  $\alpha_i \subset \omega$  for all  $i$ , to which we can apply 10.2. As already indicated, the proof of 10.1 uses 10.2 as the base for an induction. So 10.2 gives 10.1 for all  $(\alpha, \ell)$  with  $(\alpha, \ell) = (\alpha_i, \ell_i)$ ,  $1 \leq i \leq R_0$  for a totally ordered set  $\{(\alpha_i, \ell_i) : 1 \leq i \leq R_0\}$  as in 6.14, with  $L_{1,1}$  replacing  $L_{1,1,1}$ . Write  $B_{i,1} = \{(\alpha_i, \ell_i)\}$ . Inductively, for some integer  $m_1$ , we are going to define sets

$B_{i,m}$ ,  $1 \leq i \leq R_0(m)$ ,  $1 \leq m \leq m_1$  and  $B'_{i,m}$ ,  $1 \leq i \leq R'_0(m)$ ,  $2 \leq m \leq m_1$ , of pairs  $(\alpha, \ell)$  in a fixed vertically efficient decomposition of  $\omega \times [y_-, y_+]$ , with the following properties.

1.  $B_{i,m}$  and  $B'_{j,m}$  are order-closed sets of pairs  $(\ell, \beta)$  with  $\ell \subset [y_-, y_+]$  and  $\beta \subset \omega$ .
2. Let  $\alpha_{i,m}, \alpha'_{i,m}$  to be the convex hull of those  $\beta$  with  $(\ell, \beta) \in B_{i,m}$  or  $\in B'_{i,m}$ . For each  $i$  and  $m$ ,  $B_{i,m}$  is the union of a maximal number of consecutive sets  $B'_{j,m}$  for which  $\alpha_{j,m}$  are the same. and  $B'_{i,m+1}$  is the union of  $B_{i,m}$  and  $B_{i+1,m}$ , together with any  $(\ell, \beta)$  sandwiched between them. For each  $m$ , all the sets  $B_{i,m}$  are disjoint, with  $B_{i+1,m}$  the next greatest in the ordering after  $B_{i,m}$ .

Inductively, we shall show the following.

Let  $(\ell, \beta) \in B_{i,m}$  and let  $y = [\varphi] \in \ell$ . Take any collection  $\Gamma$  of disjoint simple loops in  $\beta$  with  $|\varphi(\Gamma)| \leq L_0$ . Then there is a pleated surface  $f$  with pleating locus including  $\partial\beta \cup \Gamma$  and with  $L_{1,m}$ -Lipschitz impression such that

$$|(\partial\beta \cup \Gamma)_*| \leq L_{1,m},$$

and

$$d_\beta(y, [f]) \leq L_{1,m} \text{ or } |\operatorname{Re}(\pi_\beta([g]) - \pi_\beta(y))| \leq L_{1,m}, \quad (10.5.1)$$

depending on whether  $\beta$  is a gap or a loop. Similar properties hold for  $B'_{j,m}$ .

There are pleated surfaces  $f_{i,m,\pm}$  whose pleating loci include  $\omega$  and  $\partial\alpha_{i,m}$  and pleated surfaces  $f'_{i+1,m,+}$  and  $f'_{i,m,-}$  whose pleating loci include  $\omega$  and  $\partial\alpha'_{i,m}$  such that

$$d([f_{i,m,+}], x(E_{i,m,+})) \leq L_{1,m}, \quad d([f_{i,m,-}], x(E_{i,m,-})) \leq L_{1,m}, \quad (10.5.2)$$

$$|(\partial\alpha_{i,m})_*| \leq L_{1,m},$$

$$d([f'_{i+1,m,+}], x(E'_{i,m,+})) \leq L_{1,m+1}, \quad d([f'_{i,m,-}], x(E'_{i,m,-})) \leq L_{1,m+1} \quad (10.5.3)$$

$$|(\partial\alpha'_{i,m})_*| \leq L_{1,m+1}.$$

Here,  $E_{i,m,+}$  is the order splitting defined using maximal elements of  $B_{i,m,+}$  and so on. The notation  $f'_{i+1,m,+}$  and  $f'_{i,m,-}$  is deliberate, since these are defined using respectively maximal elements of  $B_{i+1,m}$  and some maximal elements of  $B_{i,m}$ , and minimal elements of  $B_{i,m}$  and some minimal elements of  $B_{i,m,-}$ .

By 10.2, (10.5.1) and (10.5.2) are satisfied for the  $B_{i,1}$ . In general, the properties above suffice to define the  $B'_{i,m}$  from the  $B_{j,m}$ , and  $B_{i,m+1}$  from the  $B'_{j,m}$ . In order to keep the  $B_{i,m}$  disjoint, we may discard some of the sets  $B'_{j,m}$ , because the sets  $B'_{j,m}$  overlap by definition. In the case of  $B'_{1,m}$  we also include those  $(\beta, \ell)$  which are  $< (\beta_1, \ell_1)$  for some  $(\beta_1, \ell_1) \in B_{i,m} \subset B'_{1,m}$ . We make similar inclusions in  $B'_{k,m}$  for  $k = R'_0(m)$ . By 6.3,  $\alpha'_{j,m}$  is simply the convex hull of the  $\alpha_{i,m}$  with  $B_{i,m} \subset B'_{j,m}$ . Eventually we reach an  $m = m_1$  such that there is just one set  $B_{1,m_1}$ . This happens for  $m_1$  bounded in terms of the topological type of  $\omega$ , because  $\alpha_{i,t}$  is properly contained in  $\alpha_{j,t+1}$  for  $B_{i,t}$

contained in  $B_{j,t+1}$ . Then  $(\beta, \ell) \in B_{1,m_1}$  for all  $(\beta, \ell)$  in the decomposition of  $\omega \times [y_-, y_+]$  with  $\beta \subset \alpha_{1,m_1}$ . Then the proof is finished if  $\alpha'_{1,m_1} = \omega$  — and also if it is properly contained in  $\omega$ , because in the latter case we can apply the inductive hypothesis of 10.1 to  $\omega \setminus \alpha'_{1,m_1}$ .

Note that (10.5.2) for  $B_{j,m+1}$  follows from (10.5.3) for  $f'_{i,m}$  for which  $B'_{i,m} \subset B_{j,m+1}$ , since we can take  $f_{j,m,+} = f'_{i,m,+}$  for the largest  $i$  with  $B_{i,m} \subset B_{j,m}$  (so that  $B'_{i-1,m} \subset B_{j,m+1}$ ) and  $f_{j,m,-} = f'_{k,m,-}$  for the smallest  $k$  with  $B_{k,m} \subset B_{j,m+1}$ ,  $B'_{k,m} \subset B_{j,m+1}$ . We claim that (10.5.3) for  $B'_{i,m}$  follows from (10.5.1) for  $B_{i,m}$  and  $B_{i+1,m}$ . The construction of  $f'_{i+1,m,+}$  and  $f'_{i,m,-}$  are similar, so we consider  $f'_{i+1,m,+}$ . The first step is to define the pleated surfaces  $f_{i,m,+}$  and  $f_{i+1,m,-}$ . We can choose these off  $\alpha_{i+1,m}$  and  $\alpha_{i,m}$  respectively so that the pleating loci have number of intersections bounded in terms of the ltd parameter functions, and so that  $\partial\alpha'_{i,m}$  is in the pleating locus of both and so that they have no badly bent annuli. We assume now that this has been done. Then by 4.4

$$d([f_{i,m,+}], [f_{i+1,m,-}]) \leq L'_{1,m}$$

for  $L'_{1,m}$  depending on  $L_{1,m}$  and  $(\Delta_1, r_1, s_1, K_1)$ . It follows that

$$|f_{i+1,m,-}(\partial\alpha'_{i,m})| \leq L''_{1,m}$$

for  $L''_{1,m}$  depending on  $L_{1,m}$ ,  $L'_{1,m}$  and  $(\Delta_1, r_1, s_1, K_1)$ , and this gives the required bound on  $|(\partial\alpha'_{i,m})_*|$ . Then we take  $f'_{i+1,m,+} = f_{i+1,m,-}$  off  $\alpha_{i+1,m}$  and  $f'_{i+1,m,+} = f_{i+1,m,+}$  on  $\alpha_{i+1,m}$ . Then  $f'_{i+1,m,+}$  has the properties required for (10.5.3).

So it remains to prove (10.5.1) for each  $B_{i,m}$  and  $B'_{i,m}$ , by induction. The proof for  $B_{i,1}$  is given by 10.2. As already noted, (10.5.1) for  $B_{i,m}$  and  $B_{i+1,m}$  implies (10.5.2) for  $B'_{i,m}$ . So we need to obtain (10.5.1) for  $B'_{i,m}$  from (10.5.1) for  $B_{i,m}$  and  $B_{i+1,m}$ , and from (10.5.3) for  $B'_{i,m}$ . Similarly, we shall obtain (10.5.1) for  $B_{j,m+1}$  from (10.5.1) and (10.5.3) for the  $B'_{i,m}$  contained in  $B_{j,m+1}$ . First we consider (10.5.1) for  $B'_{i,m}$  when we already have (10.5.3). Now we choose two further pleated surfaces  $f_{i,m,+,+}$  and  $f_{i,m,-,-}$  as follows. An easy induction shows that each  $B_{i,m}$  contains a unique maximal element of the form  $(\ell_n, \alpha_n)$  for some  $n \leq R_0$ . Similarly,  $B_{i+1,m}$  has a unique minimal element which is then  $(\ell_{n+1}, \alpha_{n+1})$ , for the same  $n$ . Then let  $C(i, i+1, m)$  denote the convex hull of  $\alpha_n$  and  $\alpha_{n+1}$ . We can then ensure that the pleating locus of  $f'_{i,m,+}$  includes  $\partial\alpha_n$  and that the pleating locus of  $f'_{i+1,m,-}$  includes  $\partial\alpha_{n+1}$ . The distance between the right end of  $\ell_n$  and the left end of  $\ell_{n+1}$  in  $[y_-, y_+]$  is  $\leq \Delta_0$ . So we can choose  $f_{i,m,+,+}$  to have pleating locus including  $\partial\alpha_n$ ,  $\partial C(i, i+1, m)$ ,  $\partial\alpha'_{i,m}$  and  $\partial\omega$ , with (10.5.1) satisfied for  $[f] = [f_{i,m,+,+}]$ ,  $\beta = \alpha_n$  and  $y$  the right end of  $\ell_n$ . We similarly choose  $f_{i+1,m,-,-}$  to have pleating locus including  $\partial\alpha_{n+1}$ ,  $\partial C(i, i+1, m)$  (and  $\partial\alpha'_{i,m}$ ,  $\partial\omega$ ) with (10.5.1) satisfied for  $[f] = [f_{i+1,m,-,-}]$ ,  $\beta = \alpha_{n+1}$  and  $y$  the left end of  $\ell_n$ . By 4.1, 4.2, we can choose the maximal multicurves in the pleating loci of  $f_{i,m,+,+}$  and  $f_{i,m,-,-}$  to have boundedly many intersections, with bound depending on  $\Delta_0$ , and with corresponding geodesics of length bounded from 0, and hence, by 4.4,

$$d([f_{i,m,+,+}], [f_{i+1,m,-,-}]) \leq L'_0.$$

Indeed, we can choose  $\pi_{S \setminus C}([f_{i,m,+},+]) = \pi_{S \setminus C}([f_{i+1,m,-},-])$  for  $C = C(i, i+1, m)$ .

For  $i = 1$ , we can choose  $f_{1,m,-}$  to have pleating locus including  $\partial\alpha_1$  so that

$$d([f_{1,m,-},-], [f_-]) \leq L'_0.$$

We can also choose  $f_{1,m,-}$  to have pleating locus including  $\partial\alpha_1$ . We can make similar conditions on  $f_{k,m,+}$  and  $f_{k,m,+},+$  for maximal  $k$ .

So we now have pleated surfaces in order:

$$f'_{i,m,-}, f'_{i,m,+}, f_{i,m,+},+, f_{i+1,m,-},-, f'_{i+1,m,-}, f'_{i+1,m,+},$$

all of which have  $\partial\omega$  in their pleating locus and the consecutive pairs have also have as common pleating locus respectively

$$\partial\alpha_{i,m}, \partial\alpha_n \cup \partial\alpha'_{i,m}, \partial\alpha'_{i,m} \cup C(i, i+1, m),$$

$$\partial\alpha_{n+1} \cup \partial\alpha'_{i+1,m}, \partial\alpha_{i+1,m}.$$

Now take any  $(\ell, \beta) \in B'_{i,m}$ . We need to show that (10.5.1) holds for this  $(\ell, \beta)$ . We may as well assume that  $(\ell, \beta) \notin B_{i,m} \cup B_{i+1,m}$ , since otherwise there would be nothing to prove. Then we consider the five successive intervals

$$\begin{aligned} &[[f'_{i,m,-}], [f'_{i,m,+}]], [[f'_{i,m,+}], [f_{i,m,+},+]], [[f_{i,m,+},+], [f_{i+1,m,-},-]], \\ &[[f_{i+1,m,-},-], [f'_{i+1,m,-}]], [[f'_{i+1,m,-}], [f'_{i+1,m,+}]]. \end{aligned}$$

Then  $\ell$  splits into at most 5 segments, each a bounded  $d_\beta$  distance from a segment of one of these intervals, by the results of 6.4 and 6.5 with the usual modifications if  $\beta$  is a loop. The interval  $[[f_{i,m,+},+], [f_{i+1,m,-},-]]$  is bounded, so we only need to consider the other intervals, such as  $[[f'_{i,m,+}], [f_{i,m,+},+]]$  and  $[f'_{i+1,m,-}], [f'_{i+1,m,+}]]$ . On each of these we can apply the inductive hypothesis of 10.1, to  $\omega \setminus \alpha_n$  on  $[[f'_{i,m,+}], [f_{i,m,+},+]]$ , and to  $\omega \setminus \alpha_{i+1,m}$  on  $[f'_{i+1,m,-}], [f'_{i+1,m,+}]]$  — assuming as we may do that  $(\ell, \beta) \notin B_{i+1,m}$ . Then the inductive hypothesis of 10.1 gives (10.5.1) for  $(\ell, \beta)$ .

The proof of (10.5.1) for  $B_{j,m+1}$  is then immediate, because  $B_{j,m+1}$  is the union of the  $B'_{i,m}$  with both  $B_{i,m} \subset B_{j,m+1}$  and  $B_{i+1,m} \subset B_{j,m+1}$ .  $\square$

## 10.6 Modifications of 10.4, 10.5 in the case of a compressible end.

We now show how to modify the proofs of 10.2 and 10.1 for a *compressible* end  $e$ . Write  $z_{e,0} = [\varphi_{e,0}]$ . Consider the sequences  $\Gamma_i(e)$  and  $f_{e,i}$  ( $0 \leq i \leq n_+(e)$ ) of 8.9, with  $f_{e,n_+(e)} = f_{e,+}$ . We have seen in 8.9 that  $z_{e,0}$  can be chosen (moving a bounded distance if necessary) so that

$$d(z_{e,0}, x(z_{e,0})) \geq D_2,$$

where  $D_2$  is the constant of 8.8. We have also seen that, given a constant  $D_0$ , we can ensure that, moving  $z_{e,0}$  a bounded distance depending on  $D_0$  if necessary,



we assume that for all  $[\varphi] \in [z_{e,0}, y_{e,+}]$ , if  $\gamma \subset S(e)$  is nontrivial in  $S(e)$  but trivial in  $N$ , then

$$|\varphi(\gamma)| \geq D_0.$$

We assume that  $z_{e,0}$  was chosen in this way in the first place.

We now apply the proofs of 10.1 and 10.2 in 10.3 to 10.5 with  $f_{e,+}$ ,  $f_{e,0}$  replacing  $f_{\pm}$  and  $y_{e,+}$ ,  $z_{e,0}$  replacing  $y_{\pm}$ . We consider what other modifications are necessary for such ends, in both 10.2 and 10.1. First we consider 10.2. Let  $(\alpha_i, \ell_i)$ ,  $1 \leq i \leq R_0$  be exactly as in the statement of 10.2. We drop the index  $e$ . Then by 8.8, if  $D_0$  is sufficiently large given  $D_2$  and  $D_1$ ,  $|f_j(\gamma)| \geq D_1$  whenever  $T([f_j], +, [z_0, y_+])$  (in the notation of 6.10) does not contain  $(\alpha_1, \ell_1)$ . This is automatically true for  $j = n_+$ .

So then choose  $n_2$  so that containment does not happen for  $j \geq n_2$  but does happen for  $j = n_2 - 1$ . For ease of notation, renumber so that  $g_{n_2} = g_1$ . Then there are  $\ell'_j \subset [[f_0], y_+]$  for  $1 \leq j \leq R_0$  such that  $\ell'_j$  is within a bounded  $d_{\alpha_j}$ -distance of  $\ell_j$ . Then, using this sequence, we have all the properties of 8.1, including 8.1.6, for  $j \geq n_2$ . We also have

$$\sum_{j=0}^{n_+-1} d([f_j], [f_{j+1}]) \leq \kappa_1 d([f_1], y_+),$$

$$\delta_0 d([f_1], y_+) \leq \sum_{j=1}^{R_0} |\ell'_j|$$

as before. We then proceed with the construction of  $y_{m,\pm} = [f_{m,\pm}]$ ,  $z_{m,\pm}$ ,  $(\alpha_{m,\pm}, \ell_{m,\pm})$  and  $\Gamma_{m,\pm}$  as in 10.3 and 10.4, but with some minor differences. We have  $z_{0,-} = z_0$  and  $y_{0,-} = [f_1]$ , but probably not  $y_{0,-} = z_{0,-}$ . Also, so long as  $y_{k,-} = y_{0,-}$  for  $k \leq m$ , we do not try to bound  $|f_{k,-}(\Gamma_{k,-})|$  or  $d_{\alpha_{k,-}}(z_{k,-}, y_{k,-})$ . The proof of 10.2 is exactly as before, and so is the proof of 10.1. Even when we change to other sequences of pleated surfaces, as happens in case 2 in 10.5, and again in the proof of 10.1, we never change the definition of the  $f_j$  on surfaces intersecting  $\alpha_1$ . So all pleated surfaces  $[f]$  that we use have the not-containment property. So the properties of 8.1 are satisfied for all the sequences that we use, and the proofs go through as before.

This proves assumption 8.10.1 of Theorem 8.10.

## 10.7 Modifications of 10.4, 10.5 in the incompressible case.

Let  $e$  be any incompressible end. We define  $z_{e,0}$ ,  $z_{e',0}$  as in 7.14 and  $f_{e,0}$  as in 8.4. Let  $\omega(e, e')$  be as in 7.14, and as in 7.14, write

$$\beta(e) = S(e) \setminus \cup_{e'} \omega(e, e').$$

Take the sequence of multicurves and pleated surfaces  $\Gamma_{e,0}$  and  $f_{e,i}$  as in 8.4. By 8.10, for  $\Sigma'$  as defined there and in 7.11

$$|f_{e,0}(\Sigma')| \leq L_2,$$

and hence, enlarging  $L_2$  if necessary,

$$d_{\beta(e)}([f_{e,0}], z_{e,0}) = d_{\beta(e)}([f_{e,0}], z'_{e,0}) \leq L_2.$$

Now we consider the proof of 10.1. By 8.10, since the hypothesis 8.10.1 has now been proved, we have

$$|(\partial\beta(e))_*| \leq L_2,$$

and we have a pleated surface  $f_-$  with pleating locus including  $\partial\beta(e)$ , with pleating locus of length  $\leq L_2$  restricted to  $\beta(e)$ , such that

$$d_{\beta(e)}([f_-], z_{e,0}) \leq L_2.$$

By the definition of  $z'_{e,0}$  in 7.14, if  $(\alpha, \ell)$  is ltd for  $S(e) \times [z'_{e,0}, y_{e,+}]$ ,  $\alpha \cap \beta(e) \neq \emptyset$ . Take any chain of ltd's as in 6.14 for  $S(e) \times [z'_{e,0}, y_{e,+}]$ . Then 10.2 is proved as in 10.3 and 10.4, using  $f_- = f_{0,-}$ ,  $f_{e,+} = f_{0,+}$ . As in the compressible case, the fact that we do not have a bound on the pleating locus  $\Gamma_{0,-}$  of  $f_{0,-}$  off  $\beta(e)$  does not matter. We will get bounds for the first  $k$  with  $f_{k,-} \neq f_{0,-}$ . Then using this first inductive step, we can prove 10.2 for all  $(\alpha, \ell) \in \cap_{e'} E(e, e', +)$ , exactly as before.

Now, for the remaining  $(\alpha, \ell)$  in the decomposition for  $S(e) \times [z_{e,0}, y_{e,+}]$ , we also need to consider  $(\alpha, \ell)$  in a decomposition for  $\omega(e, e') \times [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$  for  $\omega = \omega(e, e')$  for varying  $e' \neq e$ . The technique of proof of 10.1 is the same as before. We start by obtaining (10.1.2) for a totally ordered set of  $(\alpha, \ell)$  with  $\alpha \cap \beta \neq \emptyset$  for all  $\alpha$  in the chain — unless  $z'_{e,0} = [f_{e,+}]$ . Having got this, we can extend to other  $(\alpha', \ell')$  which are bounded above and below by elements of the totally ordered set. When we do this, we are likely, at some point, to move into the domain of  $\omega(e, e') \times [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$  for some  $e'$ . Since we now have bounds above and below here (by  $f_{e,+}$  and  $f_{e',+}$  if not by anything lower down) we are now in a position to start the induction, applying 10.1 to a totally ordered chain for  $\omega(e, e') \times [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$ , initially using the multicurve and pleated surface sequences of 8.4. The rest of the proof is exactly as in 10.5.

## 10.8 Where short loops can occur.

Now, by the same argument as in 9.2, we can obtain information about where Margulis tubes can occur, and on the geometry of those which do occur. Precisely, we have the following. For  $\gamma \subset S \subset N$  with  $\gamma$  nontrivial in  $N$  and not represented by a parabolic element, we let  $\gamma_*$  denote the closed geodesic freely homotopic to  $S$ , as in Section 3.

It may be worth pointing out that if  $e$  is a *compressible* end, and (10.1.3) holds, and  $(\alpha, \ell)$  is ltd for  $S(e) \times [z_{e,0}, y_{e,+}]$  and  $\alpha \neq S(e)$ , then  $\alpha$  is incompressible, by taking  $\mu' = \partial\alpha/|\partial\alpha|$ . But then there cannot be another end,  $e'$ , compressible or not, with  $\alpha \subset S(e')$  up to homotopy. For if there is such an  $e'$ , any loop  $\gamma'$  on  $S(e)$  which is nontrivial on  $S$  but trivial in  $N$  must be either decomposable into two such loops, one of which is disjoint from  $\alpha$ , or is already disjoint from  $\alpha$ . Either way, this contradicts (10.1.3).

**Lemma** *We continue with the notation and hypotheses of 10.1. Fix suitable ltd parameter functions and vertically efficient ltd decompositions of  $S(e) \times [z_{e,0}, y_{e,+}]$ . Then there exists  $\varepsilon_1 > 0$  which depends only on the topological type of  $(N_{d,W}, \partial N_d)$ , and, if some end  $e$  is compressible, on the constant  $c_0$  of 10.1. Fix a Margulis constant  $\varepsilon_0$ . Let  $\gamma$  be a nontrivial nonperipheral loop in  $N$ , and suppose that  $|\gamma_*| < \varepsilon_1$  and  $T(\gamma_*, \varepsilon_0)$  is in region of  $N$  which is disjoint from the sets  $f_{e,+}(S(e))$  but bounded by all of them, and contains a tamely embedded relative Scott core. Then  $\gamma \subset \partial\alpha_j$  for some  $(\alpha_j, \ell_j)$  in the vertically efficient decomposition of  $S(e) \times [z_{e,0}, y_{e,+}]$  for some end  $e$ .*

Note that any Margulis tube  $T(\gamma_*, \varepsilon_0)$  is in such a region of  $N$  for some choice of end pleated surfaces  $f_{e,+}$ , by choosing them with images  $f_{e,+}(S(e))$  in sufficiently small neighbourhoods of the ends.

*Proof.* We use the sequence of pleated surfaces for  $[z_{e,0}, y_{e,+}]$  for each end  $e$  of 8.3, and, extending this, the family of pleated surfaces of 8.10. We assume without loss of generality that the pleated surfaces for different ends have disjoint images, by taking pleated surfaces in a sufficiently small neighbourhoods of the geometrically infinite ends. The definitions made in 8.2 mean we are using a generalised pleated surface in  $\partial N$  for each geometrically finite end. We use the same argument as in 9.2. We claim that every point in  $N$  which is in a component of the complement of the images of the boundary pleated surfaces as described in the statement of the lemma, is in the image of the homotopy between two successive pleated surfaces in an end sequence, or in the noninterval part of the core, in the image of the homotopy equivalence  $f : W' \rightarrow N$  of Theorem 8.10. The homotopies all have bounded tracks, except for the homotopy between the last two generalised pleated surfaces in a geometrically finite end — but in that case, by 4.9, the Teichmüller distance is bounded. If the claim is not true, let  $W$  be the relative Scott core in this component of the complement of the images of the pleated surfaces. We can assume an omitted point  $w_0$  of the homotopy lies in the interior of  $W$ . By [63], we can assume, after composing on the left with a homotopy which is the identity on  $W \subset N$  that the homotopy equivalence from  $W$  to  $N$  maps  $(W, \partial W)$  to  $(W, \partial W)$ . So we have a homotopy equivalence which maps  $(W, \partial W)$  into  $(W \setminus \{w_0\}, \partial W)$ . But then  $H_3(W, \partial W) \cong \mathbb{Z}$  is isomorphic to  $H_3(W \setminus \{w_0\}, \partial W) = 0$ , which is a contradiction. (This can be proved by considering the short exact sequence of chain complexes  $0 \rightarrow C_*(W \setminus \{w_0\}, \partial W) \rightarrow C_*(W, \partial W) \rightarrow C_*(W, W \setminus \{w_0\}) \rightarrow 0$ .) Now for  $\varepsilon_1$  sufficiently small, the image of the homotopy between  $f_{e,i}$  and  $f_{e,i+1}$  can only intersect a Margulis tube  $T(\gamma_*, \varepsilon_1)$  if there is no loop  $\zeta$  intersecting  $\gamma$  transversely for which  $|f_{e,i}(\zeta)| < L_1$ . Now by 10.1, in particular by (10.1.2), if  $T(\gamma_*, \varepsilon_1)$  intersects the homotopy between  $f_{e,i}$  and  $f_{e,i}$ , then  $\gamma$  has no transverse intersections with the pleating locus of  $f_{e,i}$ , nor with the pleating locus of  $f_{e,j}$  for  $|i - j| \leq t$ , if  $\varepsilon_1$  is sufficiently small given  $t$ . Now if  $t$  is sufficiently large given the ltd parameter functions, the only loops which have no transverse intersections with the pleating loci of  $2t$  successive  $g_{e,j}$  are loops which are  $\partial\alpha_k$  for some  $\alpha_k \times \ell_k$  in the decomposition.

Let  $f_{W'} : W' \rightarrow N$  be the map of 8.10. By Theorem 8.10,  $f_{W'}(W')$  has diameter  $\leq L_3$ , for some constant  $L_3$  which, ultimately, depends only on the

topological type of  $W'$  and the constant  $c_0$  of 10.1.1. So any Margulis tube  $T(\gamma_*, \varepsilon_1)$  can only intersect  $f_{W'}(W')$ , in a set of diameter  $\leq L_3$ . So then, by the first part of the proof,  $\gamma \subset \partial\alpha_j$  for some  $\alpha_j \times \ell_j$  in the decomposition for  $S(e) \times [z_{e,0}, y_{e,+}]$  for some end  $e$ . In fact, this can be true for at most two ends, and for two ends  $e, e'$  only if  $e$  and  $e'$  are both incompressible and  $\omega(e, e') \neq \emptyset$ .  $\square$

## 10.9 Bounded distance between corresponding Margulis tubes.

Let  $M$  be a model manifold of topological type  $S \times [0, 1]$ . As in 7.5, we denote by  $T(\gamma_{**})$  the Margulis tube (if any) with core loop  $\gamma_{**}$  homotopic in  $S \times [0, u]$  to  $\gamma \times \{t\}$ . In 7.4, we recalled how to parametrise Margulis tubes  $T(\gamma_*, \varepsilon_0)$  by points  $w_*(\gamma)$  in the upper half-plane  $H^2$ . In 7.5, we showed how the boundary of a model Margulis tube  $\partial T(\gamma_{**})$  also determines a point  $w_{**}(\gamma)$  of  $H^2$  up to bounded distance. The following lemma gives a bound on the distance between the corresponding points  $w_*(\gamma)$  and  $w_{**}(\gamma)$ .

We have the following, building on 10.8.

**Lemma** *We continue with the notation and hypotheses of 10.1. Fix a Margulis constant  $\varepsilon_0$ . The following holds for suitable  $\varepsilon_2 > 0$  and  $L'$  depending on  $\varepsilon_0$ , and the topological type of  $(N_{d,W}, \partial N_d)$ , and, if there are compressible ends, the constant  $c_0$  of 10.1. If one of the following holds:*

- $|\gamma_*| < \varepsilon_2$  and  $T(\gamma_*, \varepsilon_0)$  is in region of  $N$  which is disjoint from the sets  $f_{e,+}(S(e))$  but bounded by all of them, and contains a tamely embedded relative Scott core;
- $|\gamma_{**}| < \varepsilon_2$  and  $T(\gamma_{**})$  does not intersect  $\partial M$ ;

then

$$d_H(w_*(\gamma), w_{**}(\gamma)) < L',$$

where  $d_H$  denotes hyperbolic distance in  $H^2$ .

*Proof.* By 10.8, we can assume that  $\gamma = \partial\alpha_{e,j}$  for some  $\alpha_{e,j} \times \ell_{e,j}$  in a vertically efficient decomposition of  $S(e) \times [z_{e,0}, y_{e,+}]$  for some end  $e$  of  $N_{d,W}$ .

First, we assume that this is true for exactly one end  $e$ . We drop the suffix  $e$ . From the construction of the model,  $|\gamma_{**}|$  can also only be small if  $\gamma = \partial\alpha_j$  for some  $j$ . First, suppose that  $|\gamma_*| < \varepsilon_2$ , for a bound on  $\varepsilon_2$  yet to be determined, but with  $\varepsilon_2 < \varepsilon_1/2$  for  $\varepsilon_1$  as in 10.8. Let  $A$  be the set of  $k$  such that  $\alpha_k$  is a gap with  $\partial\alpha_k = \gamma$ . Let  $A_1$  be the set of  $k \in A$  such that  $\ell_k \times \alpha_k$  is ltd and  $A_2 = A \setminus A_1$ . Then by the definition of the metric in  $M$  in 7.5, the area of  $\partial T(\gamma_{**})$  is boundedly proportional to

$$\sum \{|\ell_k| : k \in A_1\} + \#(A_2). \quad (10.9.1)$$

Note that we can ignore any contribution from  $W'$ , because we have a bound on the diameter of the image of the homotopy equivalence  $f : W' \rightarrow N$  of 8.10.

Now if  $k, j \in A_2$ ,  $\alpha_k \cap \alpha_j$  contains no nontrivial subsurface in the interior, apart from  $\gamma$ , by the vertically efficient condition 5.6. Now by the last part of

the statement of 10.8, the area of  $\partial T(\gamma_*, \varepsilon_1)$  is bounded above by the number of different pleated surfaces  $f_n|_{\alpha_k}$  for which  $\partial\alpha_k$  is in the pleating locus of  $f_n$ , for  $k \in A$ , which is bounded above by a multiple of the sum in 10.9.1. But the area of  $\partial T(\gamma_*, \varepsilon_1)$  is also bounded below by a multiple of the sum in (10.9.1), with multiple depending on  $\varepsilon_1$ . This is because, by the properties of the Margulis constant, for any constant  $L'$  there is  $k(L')$  such that the following holds. For any  $x \in \partial T(\gamma_*, \varepsilon_1)$ , there are at most  $k(L')$  loops  $\zeta$  such that for some  $n$ ,  $|f_n(\zeta)| \leq L'$  and  $f_n(\zeta)$  is within distance  $L'$  of  $x$ . This works even in the case when  $S(e)$  is compressible, by Lemma 10.10 below.

So the areas of  $\partial T(\gamma_{**})$  and  $\partial T(\gamma_*, \varepsilon_1)$  are boundedly proportional as claimed, and the same is true for the areas of  $\partial T(\gamma_{**})$  and  $\partial T(\gamma_*, \varepsilon_0)$  for a constant depending on  $\varepsilon_1$ . The areas determine the imaginary coordinates of the corresponding points of  $H^2$ . So we have a bound on the ratio of  $\text{Im}(w_*(\gamma))$  and  $\text{Im}(w_{**}(\gamma))$  if one of  $|\gamma_*|$ ,  $|\gamma_{**}|$  is  $< \varepsilon_2$ .

It remains to bound the distance between  $\text{Re}(w_*(\gamma))$  and  $\text{Re}(w_{**}(\gamma))$  by a constant which is  $O(\text{Im}(w_*(\gamma))) + O(1/\varepsilon_1)$ , for  $\varepsilon_1$ . Now

$$\text{Re}(w_{**}(\gamma)) = -n_\gamma(z_2) + n_\gamma(z_1) + O(1),$$

if  $[z_1, z_2]$  is the union of all the  $\ell_k$  with  $\gamma \subset \partial\alpha_k$ . (See (7.5.1).) We recall that  $n_\gamma$  is defined relative to a fixed loop  $\zeta \subset S$  which has at least one, and at most two, essential intersections with  $\gamma$ . A different choice of  $\zeta$  does not change the quantity  $-n_\gamma(z_2) + n_\gamma(z_1)$  by more than a bounded additive constant, independent of  $\zeta$ . If the  $m$  for which  $\gamma$  is in the pleating locus of  $f_m$  include an  $m$  such that  $f_{m+1}$  and  $f_{m+2}$  are related by a Dehn twist, rather than an elementary move, then there is at most one such  $n$ , by the properties listed in 8.1, and there is a unique loop  $\zeta$  intersecting  $\gamma$  at most twice, in the pleating locus of  $f_{m+1}$ , and  $\tau_\gamma^n(\zeta)$  is in the pleating locus of  $f_{m+2}$ , where

$$n - \text{Re}(w_{**}(\gamma)) = O(1) + O(\text{Im}(w_{**}(\gamma))).$$

If there is no such  $m$ , we can still find  $m$  such that  $\gamma$  is in the pleating locus of  $f_m$  and some loop  $\zeta \subset S$  with at least one, and at most two, essential intersections with  $\gamma$ , is in the pleating locus of  $f_{m+1}$ . We choose any such  $\zeta$ , and can indeed choose such an  $m$  so that  $f_m(S)$  intersects  $T(\gamma_*, \varepsilon_1)$ , and then  $f_{m+1}(\zeta) = \zeta_*$  comes within a bounded distance of  $T(\gamma_*, \varepsilon_1)$ . By 10.8 and 10.1, if we take the minimal  $m_1$  and maximal  $m_2$  such that  $f_m(S)$  intersects  $T(\gamma_*, \varepsilon_1)$  for  $m = m_1, m_2$ , then

$$d_{\beta_j}(z_j, [f_{m_j}]) \leq L'_1$$

for a constant  $L'_1$  depending only on  $(\Delta_1, r_1, s_1, K_1)$ , where  $\beta_j = \alpha_{i(j)}$  for some  $\alpha_{i(j)}$  intersecting  $\gamma$  transversely with  $z_j \in \ell_{i(j)}$ ,  $j = 1, 2$ . Write  $n_j = n_\gamma(z_j)$ , where this is defined using  $\zeta$ . Consider the geodesics in  $S(f_{m_j})$  homotopic to

$$f_{m_j}(\tau_\gamma^{n_j}(\zeta)).$$

Take the intersection with  $T(\gamma_*, \varepsilon_0)$  of the images under  $f_{m_j}$  of these geodesics in  $N$ , and homotope, via endpoint preserving isotopy, to two segments  $\gamma_3, \gamma_4$  in

$\partial T(\gamma_*, \varepsilon_0)$  of length  $O(1/\varepsilon_1)$ . Here, we are using numbering which parallels the numbering  $\partial_k T(\alpha)$  of 7.5. Use the homotopy through the loops  $f_m(\zeta)$  to give two arcs in  $\partial T(\gamma_*, \varepsilon_0)$  of length  $O(\text{Im}(w_{**}(\gamma))) = O(\text{Im}(w_*(\gamma)))$  joining  $\gamma_1$  and  $\gamma_2$ . Then  $\gamma_1 \cup \gamma_3 \cup \gamma_2 \cup \gamma_4$  is a loop which is freely homotopic to  $\zeta_2 * \zeta_1$ , where  $\zeta_j = \tau_\gamma^{n_j}(\zeta)$ . This means that

$$\text{Re}(w_*(\gamma)) = \text{Re}(w_{**}(\gamma)) + O(\text{Im}(w_*(\gamma))) + O(1/\varepsilon_1).$$

This gives the required bound on  $d_H(w_*(\gamma), w_{**}(\gamma))$ .

Now we consider the possibility that  $\gamma \subset \partial\alpha$ ,  $\gamma \subset \partial\alpha'$  for  $(\alpha, \ell)$   $(\alpha', \ell')$  in the decompositions for two different ends  $e, e'$ . This can only happen if  $\omega(e, e') \neq \emptyset$ , and  $\alpha, \alpha' \subset \omega(e, e')$ , and up to bounded distance,  $(\alpha, \ell)$  and  $(\alpha', \ell')$  are sets in the decomposition for  $\omega \times [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$ ,  $\omega = \omega(e, e')$ . Then we can argue exactly as above, with  $\omega \times [\pi_\omega(z'_{e',0}), \pi_\omega(z'_{e,0})]$  replacing  $S(e) \times [z_{e,0}, y_{e,+}]$ .  $\square$

**Lemma 10.10.** *Let  $N_{d,W}$  be as throughout this section. We continue with the notation and hypotheses of 10.1. Fix a compressible end  $e$  of  $N_{d,W}$  bounded by  $S_d(e)$ . Let  $S = S(e)$ ,  $z_0 = z_{e,0}$ ,  $y_+ = y_{e,+}$ ,  $\Gamma_+ = \Gamma_{e,+}$  be as in 10.1, in particular satisfying (10.1.3) with respect to constants  $c_0$  and  $L_0$ . Let  $z_1 = [\varphi_1]$ ,  $z_2 = [\varphi_2] \in [z_0, y_+] \subset \mathcal{T}(S(e))$  with  $z_1 \in [z_0, z_2]$ . Let  $L$  be given. The following holds for a constant  $L'$  depending only on the topological type of  $(N_{d,W}, \partial N_{d,W})$ , on the constants  $c_0, L_0$ , on suitable ltd parameter functions. Suppose that we have fixed a vertically efficient ltd decomposition of  $S \times [z_{e,0}, y_{e,+}]$  with respect to the parameter functions. Let  $\zeta_1, \zeta_2$  be simple loops such that  $|\varphi_j(\zeta_j)| \leq L$ , and with  $\zeta_j \subset \alpha_j$ ,  $z_j \in \ell_j$  and  $z' \in \ell'$  for sets  $\alpha_j \times \ell_j$  in the ltd decomposition for  $j = 1, 2$ . Let  $d'_{\alpha_1}(z_1, z_0) \geq L'$  and let  $(\zeta_1)_* = (\zeta_2)_*$ . Then  $\zeta_1 = \zeta_2$ .*

*Proof.* Suppose that  $(\zeta_1)_* = (\zeta_2)_*$  and  $\zeta_1 \neq \zeta_2$ . We can assume without loss of generality that  $\zeta_1$  and  $\zeta_2$  have a common basepoint. So  $\zeta_1 * \overline{\zeta_2}$  is a nontrivial closed loop. Then by the Loop Theorem 4.10 of [23], we can find a loop  $\gamma \subset \zeta_1 \cup \zeta_2$  such that  $\gamma \subset S$  is a simple loop and, regarding  $S$  as a submanifold of  $N$ ,  $\gamma$  bounds an embedded disc in  $N$ . Now for any loop  $\zeta_3$ ,

$$i(\gamma, \zeta_3) \leq i(\zeta_1, \zeta_3) + i(\zeta_2, \zeta_3).$$

If  $L'$  is large enough then there is at least one ltd  $(\alpha_3, \ell_3) \leq (\alpha_1, \ell_1)$  and  $z_3 = [\varphi_3] \in \ell_3$ ,  $\zeta_3 \subset \alpha_3$  with

$$|\varphi_3(\zeta_3)| \leq L_0,$$

$$i(\zeta_1, \zeta_3) \leq C,$$

$$d'_{\alpha_3}(z_3, z_0) \geq L''.$$

Here,  $L_0$  is bounded in terms of topological type,  $C$  is bounded in terms of the ltd parameter functions, and  $L''$  can be taken arbitrarily large by choice of  $L'$ , by 6.6.

Now we claim that (10.1.3) implies that for a constant  $c_1 > 0$  depending only on  $c_0$  of (10.1.3),

$$\begin{aligned} i(\zeta, \mu) &\geq c_1 |\zeta| \text{ for some } \zeta \in \Gamma_+ \\ \text{if } i(\mu, \mu') &\leq c_1 \text{ and } i(\mu', \gamma) \leq c_1 \cdot |\gamma| \end{aligned} \quad (10.10.2)$$

For if (10.10.2) fails for a  $c_1$  depending only on  $c_0$ , we can assume that there are sequences  $\zeta_n$ ,  $\mu_n$ ,  $\mu'_n$  and  $\gamma_n$  such that  $\Gamma_{+,n}$  satisfies the condition (10.1.3), but such that  $\mu_n \rightarrow \mu_\infty$  and  $\mu'_n \rightarrow \mu_\infty$ , and

$$\lim_{n \rightarrow \infty} \text{Max}\{i(\zeta_n/|\zeta_n|, \mu_n) : \zeta_n \in \Gamma_{n,+}\} = 0$$

and

$$\lim_{n \rightarrow \infty} i(\mu_n, \mu'_n) = \lim_{n \rightarrow \infty} i(\mu'_n, \gamma_n/|\gamma_n|) = 0.$$

Then by the Lipschitz properties of  $i(\cdot, \cdot)$  ([51]), we see that (10.1.3) fails for  $\Gamma_{+,n}$  and  $\gamma_n$  for sufficiently large  $n$ , with  $\mu = \mu_\infty$  and  $\mu' = \mu'_\infty$ , giving the required contradiction.

Then by 6.6, again for  $C$  depending on the ltd parameter functions, and on  $L_0$ , for any  $\zeta \in \Gamma_+$ ,

$$|\zeta| \geq C^{-1} i(\zeta, \zeta_3) \exp d'_{\alpha_3}(z_3, z_0),$$

$$|\zeta_2| \geq C^{-1} i(\zeta_2, \zeta_3) \exp d'_{\alpha_3}(z_3, z_0),$$

$$|\zeta_3| \geq C^{-1} \exp d'_{\alpha_3}(z_3, z_0),$$

We also have, by 2.9,

$$i(\zeta_3, \zeta_2) \leq C d'_{\alpha_3, \alpha_2}(z_3, z_2),$$

again for  $C$  depending on the ltd parameter functions, because we only have a bound on  $|\varphi_2(\partial\alpha_2)|$  in terms of these. Again by 6.6, if  $z_0 = [\varphi_0]$ ,

$$|\varphi_0(\zeta_2)| \geq C^{-1} i(\zeta_3, \zeta_2) \exp d'_{\alpha_3}(z_3, z_0).$$

So we have

$$i(\zeta, \zeta_3) \leq C |\zeta| \cdot \exp -d'_{\alpha_3}(z_3, z_0),$$

$$i(\gamma, \zeta_2) \leq C + i(\zeta_3, \zeta_2) \leq C(|\zeta_3| + |\zeta_2|) \cdot \exp -d'_{\alpha_3}(z_3, z_0).$$

This contradicts (10.10.2) with  $\mu = \zeta_3/|\zeta_3|$  and  $\mu' = \zeta_2/|\zeta_2|$ , if  $L''$  is sufficiently large given  $c_1$ , that is, if  $L'$  is sufficiently large.  $\square$

## 10.11 The order on Margulis tubes is the same.

10.8 also implies the following, using the partial order of 6.3.

**Corollary** *We continue with the notation and hypotheses of 10.1. Fix an end  $e$  of  $N_{d,W}$  and write  $S(e) = S$ ,  $z_{e,0} = z_0$ ,  $y_{e,+} = y_+$ . Let  $(\gamma, \ell) < (\gamma', \ell')$  for loops  $\gamma, \gamma'$  with  $\gamma \times \ell, \gamma' \times \ell'$  in the vertically efficient decomposition of  $S \times [z_0, y_+]$ , where  $|\gamma_*| < \varepsilon_1$ ,  $|(\gamma')_*| < \varepsilon_0$ . Then the Margulis tube  $T((\gamma')_*, \varepsilon_0)$  can be homotoped to an arbitrarily small neighbourhood of  $e$ , in  $N_{d,W} \setminus T(\gamma_*, \varepsilon_0)$ . Similarly, if  $\gamma \subset \text{int}(\alpha)$ , and  $(\gamma, \ell) < (\alpha, \ell'_j)$  for  $j = 1, 2$  for  $\alpha \times \ell_j$  in the*

decomposition with  $\alpha$  a gap and  $f_k, f_m$  are pleated surfaces from the sequences of 8.4 associated to  $(\alpha, \ell'_1), (\alpha, \ell'_2)$  respectively, with pleating loci including  $\partial\alpha$ , then  $f_k(\alpha)$  can be homotoped to  $f_m(\alpha)$  via a homotopy in  $N_{d,W} \setminus T(\gamma_*, \varepsilon_0)$  and keeping  $\partial\alpha$  in  $T((\partial\alpha)_*, \varepsilon_1)$ . A similar statement holds if  $(\alpha, \ell'_j) < (\gamma, \ell)$  for  $j = 1, 2$ .

*Proof.* For the first statement, since Margulis tubes are disjoint, it suffices to find a homotopy avoiding  $T(\gamma_*, \varepsilon_1)$ , because we can then compose with a homeomorphism which expands out  $T(\gamma_*, \varepsilon_1)$  to  $T(\gamma_*, \varepsilon_0)$ . Then we simply use the homotopy defined by the sequence  $f_k$ , for  $k \geq p$  for some  $p$  such that  $\gamma'$  is in the pleating locus of  $f_p$ . By 10.8,  $f_k(S)$  does not intersect  $T(\gamma_*, \varepsilon_1)$  for  $k > p$ . The statements involving homotopies restricted to  $\alpha$  are proved similarly.  $\square$

Let  $U$  be a component of  $N \setminus f(S)$  which is a neighbourhood of the end of  $N_{d,W}$  which is the unique end of  $N_{d,W}$  in component of  $N_d \setminus W$  bounded by  $S_d$ . It is natural to say that  $T(\gamma_*, \varepsilon_0) < T(\gamma'_*, \varepsilon_0)$  in  $U$  if, as in the lemma above,  $T(\gamma'_*, \varepsilon_0)$  can be homotoped to an arbitrarily small neighbourhood of some end  $e$  in  $U$  in the complement of  $T(\gamma_*, \varepsilon_0)$ , because this order is clearly transitive. It is not yet clear that it is antireflexive.

## 10.12 First stage in the production of a geometric relative Scott core.

We already know, from 8.10 that  $N$  contains a continuous Lipschitz (but not necessarily, so far as we yet know, homeomorphic) image of the model relative Scott core in  $M$ . The following lemma gives us more information, about a relative Scott core up to homeomorphism, such that the complementary neighbourhoods of ends are products. This lemma does not say that the relative Scott core produced is biLipschitz to the model one, although we shall see later that it is. Note that in the case of a geometrically finite end, it is possible that every pleated surface in the sequence of the end intersects the  $D$ -neighbourhood of  $f_{e,0}(S(e))$

**Lemma** *We continue with the notation and hypotheses of 10.1. The following holds for a constant  $D$  depending only on the topological type of  $(N_{d,W}, \partial N_{d,W})$ , on  $c_0, L_0$ , and the loop sets  $\Gamma_0(e)$ . Fix an end  $e$  of  $N_{d,W}$  and  $S_d = S_d(e)$ . Then there is  $S' \subset N_{d,W}$  such that  $(N_{d,W}, S')$  is homeomorphic to  $(N_{d,W}, S_d)$ , and the component of  $N_{d,W} \setminus S'$  which is a neighbourhood of  $e$  is homeomorphic to  $S_d(e) \times (0, \infty)$ , each component of  $S' \cap N_{\geq \varepsilon_1}$  has diameter  $\leq D$  and for all  $f_{e,j}$  in the sequence for  $e$ , either  $f_{e,j}(S(e))$  is contained in the neighbourhood of  $e$  bounded by  $S'$ , or  $f_{e,j}(S(e))$  intersects the  $D$ -neighbourhood of  $S' \cap N_{\geq \varepsilon_1}$ .*

*Proof.* Write  $S = S(e)$ . Take the corresponding sequence  $f_{e,j} = f_j : S \rightarrow N$ ,  $0 \leq j \leq n = n_+(e)$ , of pleated surfaces, given a choice of pleated surface  $f_+ = f_n$  with image in a small neighbourhood of  $e$ , disjoint from  $W$ . For sufficiently large  $j$ , there is a bounded track homotopy between  $f_j$  and  $f_{j+1}$  which is disjoint from both  $S_d \subset \partial_d W$  and from  $f_0(S)$ . Now suppose that  $j$  is such that the homotopy between  $f_k$  and  $f_{k+1}$  is disjoint from  $f_0(S)$  for all  $k \geq j$ , but  $f_j(S)$  intersects the  $D$ -neighbourhood of  $f_0(S)$ . This is possible for  $D$  depending only on the topological type of  $S$ , by 8.1.6. Then we claim that  $f_k : S \rightarrow N \setminus f_0(S)$  is



injective on  $\pi_1$  for  $k \geq j$ . It suffices to prove this for sufficiently large  $k$ , since all the  $f_k$ ,  $k \geq j$ , are homotopic in  $N \setminus f_0(S)$ . Now for all sufficiently large  $k$ ,  $f_k$  and  $f_{k+1}$  are homotopic in  $N \setminus W$  and  $f_k : S \rightarrow N \setminus W$  is injective on  $\pi_1$ . So for sufficiently large  $k$ ,  $f_k$  is homotopic in  $N \setminus W$  to an embedding  $f'$ . It suffices to show that  $f'$  is injective on  $\pi_1$  in  $N \setminus f_0(S)$ . Suppose not. Then we can find an embedded disc  $D$  in  $N \setminus f_0(S)$ , with boundary in  $f'(S)$  but otherwise not intersecting  $f'(S)$ , which is nontrivial in  $f'(S)$ , and such that  $D$  has homotopically nontrivial intersection with  $f'(\gamma)$ , for some  $\gamma \subset S$  which is nontrivial in  $N$ . But then, since  $f'(\gamma)$  and  $f_0(\gamma)$  are homotopic in the subset of  $N$  bounded by  $f'(S)$ ,  $D$  must intersect  $f_0(\gamma)$ , which is a contradiction. Then the hypothesis of the Freedman-Hass-Scott result [21] is satisfied, and  $f_k : S \rightarrow N \setminus f_0(S)$  is homotopic to an embedding with image in an arbitrarily small neighbourhood of  $f_k(S)$ . Then we can take the image of the embedding to be  $S'$ , which is in a small neighbourhood of  $f_j(S)$  and hence intersects a  $D/2$ -neighbourhood of  $f_0(S)$  for suitable  $D$ .

Now if some  $f_k(S(e))$  is separated from  $e$  by  $S'$  and does not intersect the  $D$ -neighbourhood of  $S' \cap N_{\geq \varepsilon_1}$ , then we can repeat the construction with  $f_k$  replacing  $f_0$ , and can obtain a larger neighbourhood of  $e$ , enlarged by a set of diameter  $D/2$ . This construction can only be repeated finitely many times, because the sequence  $f_k$  is finite.  $\square$

### 10.13 Coarse biLipschitz in a ltd piece

At this point, we start to use the freedom afforded by different choices of ltd parameter functions. Let  $(\Delta_1, r_1, s_1, K_1)$  be the parameter functions and flat constant used until now, and we fix a vertically efficient partition using these. We note that, by 7.7, geometric models defined with different choice of ltd parameter functions are coarse Lipschitz equivalent. In the following lemma,  $\Delta(\nu, C)$  is likely to be much larger than  $\Delta_1(\nu)$ . There is a use of geometric limits in this lemma, but only in the context of bounded geometry.

**Lemma** *We continue with the notation and hypotheses of 10.1. Let  $\varepsilon_1$  be as in 10.8. There is a function  $\Delta(\nu, C)$ , depending on  $(\Delta_1, r_1, s_1, K_1)$ , the topological type of  $N_{d,W}$ ,  $L_0$ ,  $c_0$ , such that the following hold. Fix an end  $e$  of  $N_{d,W}$  and  $S = S(e)$ . Let  $f_{e,j} = f_j : S \rightarrow N$  be the corresponding sequence of pleated surfaces for an end, as in 8.3, between pleated surfaces  $f_- = f_0$  and  $f_+ = f_n$ , and let  $z_k = [\varphi_k] \in \mathcal{T}(S)$  be the point from which the pleating locus of  $f_k$  is constructed,  $0 \leq k \leq n$ . Let  $\alpha$  be a gap which is long  $\nu$ -thick and dominant along a segment of length  $\geq 2\Delta(\nu, C)$  centred on  $z_i$ , and let  $d(z_j, z_i) \geq \Delta(\nu, C)$ .*

*Then  $f_j(\alpha) \setminus T((\partial\alpha)_*, \varepsilon_1)$  and  $f_i(\alpha) \setminus T((\partial\alpha)_*, \varepsilon_1)$  cannot be joined by a path of length  $\leq C$  in  $N \setminus T((\partial\alpha)_*, \varepsilon_1)$ . If, in addition,  $i < j < k$ ,  $z_i, z_j, z_k$  are all in a segment of along which  $\alpha$  is long, thick and dominant, all distance  $\geq C$  from the endpoints of this segment,  $d(z_j, z_i) \geq C$ , and  $C$  is sufficiently large, then  $f_j|_\alpha$  can be homotoped to  $f_k|_\alpha$  in the complement of  $N \setminus (f_i(\alpha) \setminus T((\partial\alpha_j)_*, \varepsilon_1))$ , via a homotopy which maps  $\partial\alpha$  into  $T((\partial\alpha_j)_*, \varepsilon_1)$  and with image in  $U$ . A similar statement holds if  $k < j < i$ .*

*Proof.* Recall that in 8.3, the pleating locus of  $f_k$  was chosen to have no short

loops, and thus, with  $\partial\alpha$  not in the pleating locus, which included the maximal multicurve  $\Gamma_k$ . However,  $\Gamma_k$  was obtained by modifying a sequence of maximal multicurves  $\Gamma'_k$  which did contain  $\partial\alpha$  and such that, for each  $n$ ,  $\#(\Gamma_n \cap \Gamma'_k)$  was bounded for some  $k$ . By 4.3 and 4.4, we can return to this sequence for  $k$  such that  $z_n \in \ell$ , which we continue to call  $\Gamma_n$ , and the corresponding pleated surface  $f_n$ . But we also keep the same first pleated surface  $f_0$  in the sequence. The distance between  $f_n(\alpha)$  and  $f_{n+1}(\alpha)$  is bounded in terms of  $\nu$ , using 4.3 and the bound on  $\#(\alpha \cap \Gamma_n \cap \Gamma_{n+1})$  in terms of  $\nu$  in 8.3. If  $e$  is compressible and  $\alpha = S$ , and  $f_n(S)$  is not contained in the set  $U$  of 10.12 then by 10.12,  $f_n(S) \cap S' \neq \emptyset$ . So in this case, after discarding finitely many  $n$ ,  $f_n(S) \subset U$  for  $U$  as in 10.12. If  $\alpha \neq S$  then, as pointed out in 10.8,  $f_n|_\alpha$  is injective on  $\pi_1$ . Also,  $f_n|_\alpha$  is homotopic to an embedding via a homotopy preserving  $f_n|_{\partial\alpha}$ . We see this as follows. For sufficiently large  $p$ ,  $f_p|_S$  is homotopic to an embedding, by [21] applied to  $N \setminus W$ . Then using the homotopy through the  $f_m$  from  $F_n$  to  $f_p$ , we have a continuous map  $F : \partial\alpha \times [0, 1] \rightarrow N$  where  $F(x, 0) = f_n(x)$  and  $F(x, 1) = f_p(x)$  for all  $x \in \partial\alpha$ . Replacing  $f_p$  by  $f_q$  for an even larger  $q$  if necessary, and applying [63], we can also ensure that  $F^{-1}(F(x, j)) = F(x, j)$  for all  $x \in \partial\alpha$  and  $j = 0, 1$  and that the image of  $F$  intersects  $f_p(\alpha)$  only in  $f_p(\partial\alpha)$ . By the Generalised Loop Theorem 4.13 of [23], this map is homotopic to an embedding, keeping boundaries fixed. So then we have an embedding of  $\alpha$ , by combining  $F$  with  $f_p|_\alpha$ , and the combined map homotopic to  $f_n|_\alpha$  via a homotopy which is constant on  $\partial\alpha$ . So then the main result of [21] (as extended in Section 7 of [21]) applies and  $f_n|_\alpha$  is homotopic to an embedding, via a homotopy fixing  $\partial\alpha$ , with image in an arbitrarily small neighbourhood of  $f_p(\alpha)$ . Note that the same argument applies to  $\alpha \cup S \setminus \alpha$ . So this gives an alternative proof, in a more general setting, of Otal's proof [47] that Margulis tubes round sufficiently short loops in  $N$  are unknotted. It cannot be considered an easier proof, because we need 10.1 (or something similar) to show that  $F^{-1}F(\partial\alpha \times \{0\}) = \alpha \times \{0\}$ . We need this in order to apply the Generalised Loop Theorem.

Now we use the argument of 9.3. We fix  $\nu$  and  $C$  and subsurface  $\alpha$ , and suppose that  $\Delta(\nu, C)$  does not exist. We are in the bounded geometry case, exactly as in 9.3, except that  $\alpha$  replaces  $S$  and we have short loops corresponding to the components of  $\partial\alpha$ , rather than cusps. We have bounds on the pleated surfaces and the pleating loci of the corresponding pleated surfaces by 10.5 and 10.6, replacing the use of 9.1 for 9.3. By 10.9, we have bounds, in terms of  $\nu$ , on the Margulis tubes intersected by  $f_i(\alpha)$  with  $\partial\alpha$  in the pleating locus of  $f_i$ . If  $\Delta(\nu, C)$  does not exist, then as in 9.3, we can take geometric limits of pieces of hyperbolic manifolds  $N_p$  containing neighbourhoods of pleated surfaces  $f_i^p(\alpha)$ , bounded by pleated surfaces which are injective on  $\pi_1$  and take limits of pleated surfaces within them, exactly as in 9.3. The only difference is that we restrict the pleated surfaces to  $\alpha$  and the Margulis tubes round the loops  $(\partial\alpha)_*$ , in the different manifolds,  $\rightarrow \infty$  as  $\Delta \rightarrow \infty$ .

The statement about homotopy follows simply by using the homotopy through the pleated surfaces between  $f_j$  and  $f_k$ , since we have a bound on the diameter of the homotopy between  $f_n$  and  $f_{n+1}$ .

□

## 11 Proof of coarse biLipschitz.

In this section we prove the following theorem, which is very close to 1.3.

**Theorem 11.1.** *Let  $e_i$  ( $1 \leq i \leq n$ ) be the ends of  $N_{d,W}$ . For any choice of end pleated surfaces  $f_{e,+}$  satisfying the conditions of 10.1, and  $M = M([f_{e_1,+}], \dots [f_{e_n,+}])$ , there is a map  $\Phi : M \rightarrow N$ , with image containing  $f_{e_i,+}(S(e_i))$  for all  $i \leq n$ , which is  $\Lambda_2$ -coarse-biLipschitz, for a constant  $\Lambda_2$  which depends only on the topological type of  $N_{d,W}$  and the constant  $c_0$  of 10.1.3.*

The fact that the map is coarse Lipschitz follows from the results in Section 10. The main tools for proving coarse biLipschitz are given in 10.9 to 10.13. I believe that this may constitute an important difference from the proofs of [10] and [7]. The work done in 10.9 and 10.13 means that the map  $\Phi$ , which we are about to construct formally, already maps bits of surfaces in the ends in the correct order.

First we need to get into a position to apply 10.1 and the other results of section 10. We need the following lemma. For Theorem 1.1 we consider a sequence with  $N_n = N$  for all  $n$ . For Theorem 1.2 we take each  $N_n$  to be geometrically finite, with specified ending lamination data.

**Lemma 11.2.** *Let  $M_d$  be a topological model for the horoball deletion of a hyperbolic 3 manifold with finitely generated fundamental group with ends  $e_i$ ,  $1 \leq i \leq m$ . Let  $[\mu(e_1), \dots \mu(e_m)]$  be any permissible ending invariant. Let  $N_n$  be a sequence of hyperbolic manifolds of this topological type. Identify the ends of  $(N_n)_d$  with the ends of  $M_d$ , and label them  $e_i$ ,  $1 \leq i \leq m$ , accordingly. Let the ending invariants be  $[\mu(e_1, n), \dots \mu(e_m, n)]$  and let  $\mu(e_i, n) \rightarrow \mu(e_i)$  as  $n \rightarrow \infty$ . Regard  $S(e)$  as a subsurface of each of the  $N_n$ , bounding a neighbourhood of  $e$ . Fix an end  $e$ . Then we can find  $\Gamma_+(e, n)$ ,  $z_{e,0}$ ,  $c_0$ ,  $L_0$ ,  $f_{e,+,n}$  so that the conditions of 10.1 are satisfied for all but finitely many  $n$ , and so that any limit of  $[f_{e_i,+,n}]$  (using the topology of 3.9) is in  $\mathcal{T}(S(e_i))$  or  $\mathcal{O}_a(S(e_i, N))$  or  $\mathcal{W}(S(e_i, \Gamma))$  according to what is true for  $\mu(e_i)$ , for the same  $\Gamma$  in the last case, and all geodesic lamination components equal to the geodesic lamination components of  $\mu(e_i)$ .*

*Proof.*

We always choose  $\Gamma_+(e, n)$  to include the loops which bound the support of the geodesic lamination components of  $\mu(e)$ . If  $\mu(e, n) \in \mathcal{O}_a(S(e, N))$ , then we choose  $f_{e,n}$  with  $f(S(e, n))$  far out in the end, and with pleating locus  $|f(\Gamma_+(e, n))|$  bounded, by 3.11. This gives (10.1.1). If  $\mu(e, n) \in \mathcal{T}(S(e_i))$ , then we use 4.6. instead. In this case, we have freedom in the choice of  $\Gamma_{e,n}$ , with  $[f_{e,+,n}]$  taken to be  $[f_2]$  of 4.6, if  $\Gamma_+(e, n)$  is the  $\Gamma$  of 4.6. In the interval bundle case we are then finished.

Now suppose that  $\mu(e) \in \mathcal{W}(\Gamma, S(e))$  for some  $\Gamma \in \mathcal{O}(S, N)$ . Then 10.1.3 is satisfied with any  $\zeta \in \Gamma$  replacing  $\Gamma_+(e)$ , for some constant  $c'_0$ . If a component of

$\mu(e)$  is in  $\mathcal{T}(S(\alpha))$  for some gap  $\alpha$  of  $S(e) \setminus (\cup \Gamma(e))$ , then we choose  $\alpha \cap \Gamma_+(e, n)$  to be constant in  $n$ , and in  $\mathcal{O}(\alpha, N)$ , so that (10.1.3) is satisfied for some  $c_{0,\alpha}$ , for any  $\gamma', \mu, \mu'$  as in (10.1.3) with support in  $\alpha$ . For  $\alpha$  such that the corresponding component  $\mu(e, \alpha) \in \mathcal{O}_a(\alpha, N)$ , choose any  $\zeta_n \in \Gamma_{e,n}$ . Then we claim that (10.1.3) is satisfied for sufficiently large  $n$ , with  $\Gamma_+(e, n)$  replacing  $\Gamma_+(e)$ , for some  $c_0 > 0$ . For suppose not. Then there are geodesic laminations  $\mu'_n$  and loops  $\gamma'_n$  which are trivial in  $N$  but nontrivial in  $S(e)$  such that, after taking subsequences,

$$i(\mu'_n, \gamma'_n) = 0, \quad \lim_{n \rightarrow \infty} i\left(\mu'_n, \frac{\zeta_n}{|\zeta_n|}\right) < c_0.$$

Let  $\mu'$  and  $\mu''$  be limits of  $\mu'_n$  and  $\zeta_n/|\zeta_n|$  respectively. Then we have  $i(\mu'', \mu') = 0$ . In the notation of 4.6,  $\mu(e, n) = [f_2]$  and  $|f_2(\zeta_n)|$  is bounded. So if  $\zeta_n \subset \alpha$ , we have  $\mu'' = \mu(e, \alpha)$ . Now  $i(\zeta, \mu') \geq c'_0$  for any  $\zeta \in \Gamma$ . So  $\mu'$  has support at least  $c'_0$  in any  $\alpha$ . It is then impossible to have  $i(\mu'', \mu') = 0$  unless  $\mu'$  is a nonzero multiple of  $\mu'' = \mu(e, \alpha)$ , giving a contradiction.

□

### 11.3 Construction of Lipschitz $\Phi$ .

We construct the coarse biLipschitz map  $\Phi : M \rightarrow N$  as a limit of maps  $\Phi_n$ , where each  $\Phi_n$  is defined from a choice of pleated surfaces  $f_{e,n}$  for each end  $e$  of  $N_d$ . Let  $e_i, 1 \leq i \leq m$ , be the ends of  $N_d$ . Then we have

$$\Phi_n : M_n = M([f_{e_1,n}], \dots [f_{e_m,n}]) \rightarrow N.$$

Then by 7.17,  $M_n$  converges geometrically, with a suitable basepoint, to  $M = M(\mu(e_1), \dots \mu(e_m))$ .

Then, to prove 11.1, it suffices to show that the  $\Phi_n$  are coarse biLipschitz with respect to a constant  $\Lambda_2$  which is bounded in terms of the topological type of  $N_{d,W}$ , and the constant  $c_0$  of 11.2 (and (10.1.3)), because then we shall have a convergent subsequence to  $\Phi : M \rightarrow N$  which is coarse biLipschitz with the same bound on constant, whose image is all of the convex hull except for bounded neighbourhoods of the convex hull boundary components corresponding to geometrically finite ends. In this subsection, we complete the formal construction of the map, which, by the results of section 10 and earlier, is Lipschitz with respect to a constant  $\Lambda_1$  depending only on the topological type of  $N_{d,W}$ , and the constant  $c_0$ . From now on we drop the index  $n$  and write  $\Phi$  for  $\Phi_n$ .

We fix a Margulis constant  $\varepsilon_0$ . We fix ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$ . The geometric model is defined using this fixed choice of parameter functions and a vertically efficient decomposition of  $S(e) \times [z_0(e), y_+(e)]$ , although, as we saw in 7.7, different choices give the same model up to biLipschitz equivalence. The geometric model is a union of pieces  $M(z_0(e), y_+(e))$ , one for each end  $e$  of  $N_{d,W}$ , a model for the noninterval part  $W'$  of  $W$ , and possibly some model Margulis tubes wedged in between. If  $N$  is homeomorphic to the interior of an interval bundle, when  $W' = \emptyset$  and the two end models are combined in one.

Let  $f_j = f_{e,j}$  be the sequence of pleated surfaces constructed in 8.3, 8.4 using the ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$ . This was constructed using a sequence of points  $z_j = [\varphi_{t_j}] \in [z_0, y_+]$ ,  $0 \leq j \leq n_+$ . We have

$$d([f_j], [f_{j+1}]) \leq L_0$$

and by property 8.1.4, there is a bound by  $L_0$  on homotopy tracks of a homotopy between  $f_j$  and  $f_{j+1}$ . Let  $S_{t_j}$  be as in 7.2, that is, the hyperbolic surface determined by  $z_j$ . If  $[\varphi_t] \in \ell = \ell_k$  where  $\alpha = \alpha_k$  is a gap or loop with  $\alpha \times \ell$  in the decomposition, write  $S_{\alpha,t}$  for the subsurface  $S_{k,t}$  of  $S_t$  of 7.2. We can also assume that the homeomorphism  $\varphi_{t_k} : S \rightarrow S_{t_k}$  is part of the family  $\varphi_t$  with the properties of 7.2. The results of 10.1 give bounds on  $d_\alpha(z_j, [f_j])$ , whenever  $z_j \in \ell$  and  $\alpha \times \ell$  is a set in the vertically efficient decomposition. 10.1 also gives an upper bound on  $|(\partial\alpha)_*|$ , but there is also a better upper bound for short loops in 10.9. We recall that  $S(f_j)$  is the hyperbolic surface structure such that  $f_j : S(f_j) \rightarrow f_j(S)$  is isometric restricted to sets with totally geodesic image. (we sometimes call this map  $\text{Imp}(f_j)$ .) We write  $S(f_j, \alpha)$  for the subset of  $S(f_j)$  which is homotopic to  $f_j(\alpha)$ , and is a complementary component of  $\varepsilon(|f_j(\gamma)|)$ -neighbourhoods of the loops  $f_j(\gamma)$  for  $\gamma \subset \partial\alpha$ , where  $\varepsilon(L)$  is the function of 7.2, taken to be  $\varepsilon_0$  whenever  $L < \varepsilon_0/2$ .

Then for each  $j$  and each gap  $\alpha$  with  $\alpha \times \ell$  in the decomposition of  $S \times [z_0, y_+]$ , and for  $z_{t_j} \in \ell_j$ , we map  $\varphi_{t_j}^{-1}(S_{\alpha,t_j})$  to  $f_j(S(f_j, \alpha))$ . Because the results of 10.1, 10.6, 10.9 imply that  $S_{\alpha,t_j}$  and  $S(f_j, \alpha)$  are boundedly Lipschitz equivalent, we can make this map coarse Lipschitz. For  $t \in [t_j, t_{j+1}]$ , if  $[z_j, z_{j+1}] \subset \ell$ , we map  $\varphi_t^{-1}(S_{\alpha,t})$  to the image of the bounded track homotopy between  $f_j|_{\varphi_{t_j}^{-1}(S_{\alpha,t_j})}$  and  $f_{j+1}|_{\varphi_{t_{j+1}}^{-1}(S_{\alpha,t_{j+1}})}$ . Doing this for all  $j$  and all gaps  $\alpha$  for  $\alpha \times \ell$  in the decomposition, the map  $\Phi$  is defined on all of  $M$  except for the model Margulis tubes and is coarse Lipschitz, with respect to the hyperbolic metric on the image. For loops  $\gamma$  with model Margulis tubes for which at least one of  $|\gamma_{**}|$  or  $|\gamma_*| < \varepsilon_2$ , the map  $\Phi$  maps  $\partial T(\gamma_{**})$  to a bounded neighbourhood of  $\partial T(\gamma_*, \varepsilon_0)$  (which is nonempty). The map  $\Phi$  is then also coarse Lipschitz with respect to the induced metric on  $\partial T(\gamma_*)$ , and the metric induced from the hyperbolic metric on a bounded neighbourhood of  $\partial T(\gamma_*, \varepsilon_0)$ . Then we can extend the definition of  $\Phi$  across the model tubes  $T(\gamma_{**})$ , so that the map is Lipschitz and so that  $T(\gamma_{**})$  maps into a bounded neighbourhood of  $T(\gamma_*, \varepsilon_0)$  whenever  $|\gamma_{**}|$  or  $|\gamma_*| < \varepsilon_2$ . Defining the map in a Lipschitz way is clear when there is a bound on the diameter of  $T(\gamma_{**})$ , that is, if  $|\gamma_{**}| \geq \varepsilon_2$ . It is also clear if  $|\gamma_{**}| < \varepsilon_2$  because then, by 10.9,  $|\gamma_{**}|$  and  $|\gamma_*|$  are boundedly proportional (and more) and since the metric is chosen so that  $T(\gamma_{**})$  is isometric to a Margulis tube, the coarse Lipschitz map on  $\partial T(\gamma_{**})$  can be continuously extended to be coarse Lipschitz on the  $r$ -neighbourhood of  $\gamma_{**}$ , for each  $r \leq R$  where  $T(\gamma_{**})$  is the  $R$ -neighbourhood of  $\gamma_{**}$ .

We define  $\Phi = f_{W'}$  on  $W'$ , where  $W'$  is the non-interval-bundle part of  $W$ , and  $f_{W'}$  is the map of 8.10. This is a Lipschitz map, with Lipschitz constant  $\Lambda_1$  depending only on the topological type of  $W'$ , and on  $c_0$ , by 8.10. Also, by 8.10, the definition of  $f_{W'}$  on  $\partial W' \setminus N_d$  agrees with the definition of  $\Phi$  on

the boundaries of the model end manifolds, where the pleated surfaces  $f_{e,0}$  are used.

### 11.4 Proof of coarse biLipschitz: vertical length.

So it remains to show that  $\Phi : M \rightarrow N$  is coarse biLipschitz onto its image. Some preliminary work on this has already been done, notably in 10.11 and 10.13. We write  $\tilde{\Phi}$  for the lift  $\tilde{\Phi} : \tilde{M} \rightarrow H^3 = \tilde{N}$ . If we use  $d_{\tilde{M}}$  and  $d_H$  to denote the metric on  $\tilde{M}$  and on  $H^3$ , then all we actually need to do is to show that

$$\liminf_{d_{\tilde{M}}(x,y) \rightarrow \infty} d_H(\tilde{\Phi}(x), \tilde{\Phi}(y)) > 0, \quad (11.4.1)$$

where we take any  $x, y$  lifting points  $x', y'$  where  $\Phi(x'), \Phi(y')$  are in the region bounded by all the pleated surfaces  $f_{e,+}(S(e))$  (for all ends  $e$ , for the surfaces  $f_{e,+}$  used to define  $\Phi$ ). This is to ensure that the geodesic segment joining  $\tilde{\Phi}(x)$  and  $\tilde{\Phi}(y)$  is in the image of  $\tilde{\Phi}$ . (11.4.1) is sufficient, because if this is true, then for some  $\delta > 0$  given any points  $x, y$ , we can choose  $w_i$ ,  $0 \leq i \leq n$  with  $w_i$  successive points on the geodesic segment joining  $\tilde{\Phi}(x)$  and  $\tilde{\Phi}(y)$  with

$$\frac{\delta}{2} \leq d_H(w_i, w_{i+1}) \leq \delta$$

Then since  $w_i$  is in the image of  $\tilde{\Phi}$ , we can choose  $x_i$  with  $\tilde{\Phi}(x_i) = w_i$ . Then (11.4.1) gives an upper bound on  $d_{\tilde{M}}(x_i, x_{i+1})$  by some  $C$  and hence

$$\begin{aligned} d_H(\tilde{\Phi}(x), \tilde{\Phi}(y)) &\geq \frac{n\delta}{2} \geq \frac{\delta}{2C} \sum_{i=0}^{n-1} d_{\tilde{M}}(x_i, x_{i+1}) \\ &\geq \frac{\delta}{2C} d_{\tilde{M}}(x, y). \end{aligned}$$

We shall actually show the slightly stronger statement than (11.4.1):

$$\lim_{d_{\tilde{M}}(x,y) \rightarrow \infty} d_H(\tilde{\Phi}(x), \tilde{\Phi}(y)) = +\infty. \quad (11.4.2)$$

Already, in 10.13, we switched to a stronger set of ltd parameter functions, and we continue to do this. All our pleated surfaces, the geometric model and  $\Phi$ , are defined using the ltd parameter functions  $(\Delta_1, r_1, s_1, K_1)$ . We now use a stronger set of ltd parameter functions  $(\Delta_2, r_2, s_2, K_2)$ , such that  $\Delta_2(\nu) \geq 3\Delta(\nu, C)$  for all  $\nu$ , for a suitable  $\Delta(\nu, C)$  as in 10.13, so that the last part of 10.13 works,  $r_2(\nu) \leq \varepsilon_2$  for  $\varepsilon_2$  as in 10.9,  $s_2(\nu) \leq s_1(\nu)$  and  $K_2 \geq K_1$ . We also choose the parameter functions strong enough for all the results of Section 6 to work. Having made this second choice of ltd parameter functions, as in 7.7 we can choose a vertically efficient decomposition with respect to  $(\Delta_2, r_2, s_2, K_2)$  which is coarser than the vertically efficient decomposition used for  $(\Delta_1, r_1, s_1, K_1)$ . Naturally, the reason for choosing these new parameter functions is that we wish to use 10.9 and 10.13. The biLipschitz constant will depend on this second

choice of parameter functions, while the Lipschitz constant, which has already been found, depends on the first ltd parameter functions.

The idea of the proof of (11.4.2) is to show that, for homotopy classes of paths  $\gamma$  in  $M$  whose images under  $\Phi$  can be homotoped keeping endpoints fixed to geodesics of bounded length, first,  $\gamma$  must have bounded vertical length up to homotopy, and then, given a bound on the vertical length, we can bound what we call the *horizontal length*. In fact, the horizontal length is simply the length. The term is only used when the *vertical length* is bounded. (The terms arose in earlier, more simplistic, incomplete versions of this part of the proof. But they still have a good intuitive meaning.) In the combinatorially bounded geometry Kleinian surface case, the measure of vertical length up to homotopy is just the difference of the vertical (second) coordinates of the endpoints in  $S \times \mathbb{R}$ . Obviously this needs to be refined in the general case, when the measurement of length of the vertical coordinate varies depending on which subsurface we are in. We say that a path in  $M$  has *bounded vertical length* (up to homotopy) if it can be homotoped into a union of bounded diameter subsets of model Margulis tubes and boundedly many sets

$$M(\alpha, I) = M(\alpha, \ell) = \cup \{\varphi_t^{-1}(S_{\alpha, t}) : t \in I\}.$$

Here,  $\varphi_t$ ,  $S_{t, \alpha}$  are as in the definition of geometric model  $M(z_0, y_+)$  with respect to  $(\Delta_2, r_2, s_2, K_2)$  for some end  $e$ , and  $\ell = \{[\varphi_t] : t \in I\}$ . The set  $\alpha \times \ell$  is either in the chosen vertically efficient decomposition of  $S(e) \times [z_0, y_+]$  (for  $(\Delta_2, r_2, s_2, K_2)$ ) or is a subset of some  $\alpha, \ell'$  in the decomposition. In either case we also have that  $\ell$  has  $d_\alpha$ -length  $\leq L = L(\Delta_2, r_2, s_2, K_2)$  for suitable  $L$ .

## 11.5 Basic topological principles

We shall use the following. Let  $N$  be a hyperbolic manifold, and for  $Y \subset N$  we let  $B_\delta(Y)$  denote the  $\delta$ -neighbourhood of  $Y$ , in the hyperbolic metric. We let  $C$  be the constant of the Injectivity Radius Lemma in 3.3. As in 10.13, in the following lemma, we shall use the surface  $S'$  of 10.12, and the neighbourhood  $U$  of  $e$  bounded by  $S'$ , in the case when  $e$  is incompressible. Let  $\varepsilon_1$  be as in 10.8. By 10.8 and 10.9 we can assume, and shall do so, that  $\varepsilon_1$  is small enough that if  $|\gamma_*| < \varepsilon_1$ , then a model Margulis tube with core  $\gamma_{**}$  in the model manifold does exist, and  $|\gamma_{**}| < \varepsilon_0$ .

### Lemma

Let  $S = S(e)$ . Let  $h_i : S \rightarrow N$  ( $0 \leq i \leq 3$ ) be pleated surfaces which satisfy the conclusion of the Radius of Injectivity Lemma 3.3. Let  $\alpha, \alpha'$  be subsurfaces of  $S$ . If  $e$  is compressible and  $\alpha = S$ , let  $h_i(\alpha) \subset U$  for  $i = 0, 1, 2$ .

Let  $|(\partial\alpha)_*| < \varepsilon_1$ . and  $|(\partial\alpha')_*| < \varepsilon_1$ . Let  $\partial\alpha$  be in the pleating locus for  $h_i$ ,  $i = 0, 1, 2$ , and let  $\partial\alpha'$  be in the pleating locus of  $h_3$ .

Let  $\gamma_0, \gamma_2$  be nontrivial, nonperipheral, closed loops in  $\alpha$ , not homotopic to boundary components, such that  $h_0$  and  $h_1$  are homotopic via a homotopy which maps  $\partial\alpha$  into  $T((\partial\alpha)_*, \varepsilon_1)$ , and with image in  $U$  if  $e$  is compressible and  $\alpha = S$ , and disjoint from  $h_2(\gamma_2)$ , and similarly with  $(h_0, h_1, \gamma_2)$  replaced by  $(h_2, h_1, \gamma_0)$ .

Suppose also that any two of  $h_0, h_1, h_2$  are homotopic via a homotopy which maps  $\partial\alpha$  into  $T((\partial\alpha)_*, \varepsilon_1)$ , and with image in  $U$  if  $e$  is compressible and  $\alpha = S$ , and disjoint from  $h_3(\gamma_3)$ . Here,  $\gamma_3 \subset \overline{\alpha'}$  is a nontrivial, nonperipheral loop which is not homotopic to any component of  $\partial\alpha$ , but is allowed to be homotopic to a component of  $\partial\alpha' \setminus \partial\alpha$ .

Suppose that

$$h_3(\alpha') \cap B_\delta(h_1(\alpha)) \setminus T((\partial\alpha)_*, \varepsilon_1) \neq \emptyset \quad (11.5.1)$$

and

$$(h_i(\alpha) \cap B_\delta(h_j(\alpha)) \setminus T((\partial\alpha)_*, \varepsilon_1) = \emptyset \text{ for } (i, j) = (0, 1), (1, 2), (0, 2). \quad (11.5.2)$$

Then

$$(h_0(\alpha) \cup h_2(\alpha)) \cap h_3(\alpha') \setminus T((\partial\alpha)_*, \varepsilon_1) \neq \emptyset. \quad (11.5.3)$$

We remark that if  $\alpha = \alpha' = S(e)$  we can make a slightly simpler statement, but the above is sufficient for our purposes.

*Proof.*

Let  $C$  be the constant of the Radius of Injectivity Lemma 3.3. By our hypothesis on the Radius of Injectivity Lemma Radius of Injectivity Lemma 3.3,  $h_i$  maps the boundary components of  $(S(h_i))_{\geq \varepsilon_1}$  which are homotopic to  $\partial\alpha$  into  $T((\partial\alpha)_*, \varepsilon_1) \setminus T((\partial\alpha)_*, \varepsilon_1/C)$ . Reparametrising and taking a perturbation by a distance  $\log C$  with support inside  $T((\partial\alpha)_*, \varepsilon_1)$ , we can assume, for a subsurface  $\alpha_1$  of  $\alpha$  which is homotopic to  $\alpha$ , that  $\partial\alpha_1$  is mapped homeomorphically to  $\partial T((\partial\alpha)_*, \varepsilon_1/C)$ . As pointed out in 10.8,  $\alpha$  is incompressible unless  $\alpha = S(e)$ , even if  $e$  is compressible, and if  $e$  is compressible and  $\alpha = S(e)$ , we are assuming that  $h_i(\alpha) \subset U$ . So in all cases, as in 10.13, we can apply the results of [21]. So we can perturb to a diffeomorphism on  $\alpha_1$ , with image in an arbitrarily small neighbourhood of the original.

Then, taking another perturbation, we can assume that the image is transverse to  $\partial T((\partial\alpha)_*, \varepsilon_1)$ . Then we can reparametrise again and assume that  $\partial\alpha_1$  is contained in  $\partial T((\partial\alpha)_*, \varepsilon_1)$ . We can perturb so that  $h_i(\alpha_1) \subset N \setminus T((\partial\alpha)_*, \varepsilon_1)$ , perturbing the intersection of the image with  $N \setminus T((\partial\alpha)_*, \varepsilon_1)$  by an arbitrarily small amount. So altogether we can assume that

$$h_i : (\alpha_1, \partial\alpha_1) \rightarrow (N \setminus T((\partial\alpha)_*, \varepsilon_1), \partial T((\alpha)_*, \varepsilon_1))$$

is a homeomorphism, keeping the intersection of the image with  $N \setminus T((\partial\alpha)_*, \varepsilon_1)$  in an arbitrarily small neighbourhood of the original, for  $0 \leq i \leq 2$ , and, similarly for a subsurface  $\alpha'_1$  of  $\alpha'$  which is homotopic to  $\alpha'$ , we can assume that  $h_3$  is a homeomorphism, with

$$h_3 : (\alpha'_1, \partial\alpha'_1) \rightarrow (N \setminus T((\partial\alpha')_*, \varepsilon_1), \partial T((\alpha')_*, \varepsilon_1)).$$

We are assuming that the  $h_j(\alpha_1)$ ,  $j = 0, 1, 2$ , are disjoint. By Waldhausen's theorem [63] the closed region bounded by  $h_j(\alpha_1)$  and  $h_k(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ , for  $j, k \in \{0, 1, 2\}$  and  $j \neq k$ , is homeomorphic to  $\alpha_1 \times [0, 1]$ . Given the boundary-preserving homotopy between  $h_0(\alpha_1)$  and  $h_1(\alpha_1)$  in the complement



of  $h_2(\gamma_2)$ , it follows that  $h_2(\alpha_1)$  is not between  $h_0(\alpha_1)$  and  $h_1(\alpha_1)$ . Similarly,  $h_0(\alpha_1)$  is not between  $h_1(\alpha_1)$  and  $h_2(\alpha_1)$ . So the region bounded by  $h_0(\alpha_1)$  and  $h_1(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ , and the region bounded by  $h_1(\alpha_1)$  and  $h_2(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ , must be on opposite sides of  $h_1(\alpha_1)$ . So  $h_1(\alpha_1)$  is in the region bounded by  $h_0(\alpha_1)$  and  $h_2(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ . But because of the boundary-preserving homotopy between  $h_0(\alpha_1)$  and  $h_2(\alpha_1)$  in the complement of  $h_3(\gamma_3)$ ,  $h_3(\alpha'_1)$  is not contained in the region bounded by  $h_0(\alpha_1)$  and  $h_2(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ , and does not intersect  $T((\partial\alpha)_*, \varepsilon_1)$  except in components of  $h_3(\partial\alpha'_1)$  for components of  $\partial\alpha'_1$  which are also in  $\partial\alpha_1$ . So if, as we are assuming in (11.5.1) and (11.5.2),  $h_3(\alpha'_1)$  does intersect the interior of the region bounded by  $h_0(\alpha_1)$  and  $h_2(\alpha_1)$  and  $T((\partial\alpha)_*, \varepsilon_1)$ , then  $h_3(\alpha'_1)$  must intersect  $h_0(\alpha_1) \cup h_2(\alpha_1)$ . We are working with an arbitrarily small perturbation of the original  $h_j$ . This gives (11.5.3), as required.  $\square$

## 11.6 Cross-sections.

We extend the notion of partial order of 6.3. A *complete cross-section* (for  $S \times [z_0, y_+]$ ) is a set of  $n$ -tuples for varying  $m$ :

$$b = ((\beta_1, x_1), \dots, (\beta_m, x_m)),$$

such that the  $\beta_i$  are disjoint but the union of the  $\overline{\beta_i}$  is  $S$ , and one of the following holds.

- $(\beta_i, \ell_i)$  is an ltd in the decomposition with  $x_i \in \ell_i$ .
- $\beta_i = \cup_j \beta_{i,j}$ , where  $\beta_{i,j} \times \ell_{i,j}$  is a bounded set in the decomposition,  $x_i \in \cap_j \ell_{i,j}$ ,  $|\gamma_{**}| < \varepsilon_2$  for all  $\gamma \subset \beta_i$ , and  $|\gamma_{**}| \geq \varepsilon_2$  for all  $\gamma \subset \text{int}(\beta_i)$ .

Then we define

$$((\beta_1, x_1), \dots, (\beta_m, x_m)) < ((\beta'_1, x'_1), \dots, (\beta'_n, x'_n))$$

if the following hold.

1.  $x_i$  is strictly to the left of  $x'_j$  in  $[z_0, y_+]$  whenever  $\beta_i \cap \beta'_j \neq \emptyset$ .
2. If  $\beta_i = \beta'_j$  then  $\beta_i$  is ltd with respect to  $(\Delta_2, r_2, s_2, K_2)$  along a segment centred at  $x_i$ , and similarly for  $(\beta'_j, x'_j)$ .

A *cross-section* for  $\alpha \times \ell$  is similarly defined, whenever  $\alpha \times \ell$  is a union of sets in the vertically efficient decomposition.

A cross-section  $b = ((\beta_1, [\varphi_{t_1}]), \dots, (\beta_m, [\varphi_{t_m}]))$  for  $\alpha$  defines a surface  $S(b)$  in the model manifold which contains, in the language of 11.4,

$$\cup_{j=1}^m M(\beta_j, \{t_j\}),$$

where these are joined up by annuli in model Margulis tubes  $T(\gamma_{**})$ , for each  $\gamma$  which is a component of  $\partial\beta_j \cap \partial\beta_k$  for some  $j \neq k$ . A cross-section for  $\alpha$  also defines an element of  $x(b) \in \mathcal{T}(S(\alpha))$  up to bounded distance by

$$\pi_{\beta_j}(x(b)) = [\varphi_{t_j}]$$

if  $\beta_j$  is a gap, and if  $\beta_i$  is a loop,

$$\text{Im}(\pi_{\beta_j}(x(b))) = \text{Im}(\pi_{\beta_i}([\varphi_{t_j}]), \quad \text{Re}(\pi_{\beta_j}(x(b))) = \frac{1}{\varepsilon_0}$$

for some fixed Margulis constant  $\varepsilon_0$ .

A totally ordered chain of cross sections  $\{b_i : 1 \leq i \leq r\}$  is *maximal* if it is not possible to insert any  $b$  between any  $b_i$  and  $b_{i+1}$  with  $b_i < b < b_{i+1}$ , any  $1 \leq i < r$ .

**Lemma 11.7.** *Let  $b = ((\beta_1, x_1), \dots, (\beta_m, x_m))$ ,  $b' = ((\beta'_1, x'_1), \dots, (\beta'_n, x'_n))$  be cross-sections for  $\omega \times \lambda$  for some union of sets  $\omega \times \lambda$  in the vertically efficient decomposition. Suppose that  $b < b'$ . Then each  $\gamma \subset \partial\beta_i$ , for each  $i$ , is either equal to some  $\gamma' \subset \partial\beta'_j$ , or is contained in some  $\beta'_j$  for  $\beta'_j$  ltd along a segment in  $[x_i, x'_j]$ , or  $\gamma$  intersects  $\partial\beta'_j$  transversally. Similar properties hold for each  $\gamma' \subset \partial\beta'_j$ .*

*Proof.* Suppose not so for  $\gamma$ . Then there must be some  $j$  such that  $\gamma$  is contained in the interior of  $\beta'_j$  and  $\beta'_j \times \ell'_j$  is not ltd in the decomposition for  $x'_j \in \ell'_j$ . Consider the set of  $(\alpha, \ell)$  in the decomposition such that  $\gamma \cap \alpha \neq \emptyset$ ,  $\gamma \notin \partial\alpha$ ,  $(\gamma, x_i) < (\alpha, \ell) < (\beta'_j, x'_j)$ , using the usual ordering of 6.3. By the definition of vertically efficient, in particular, property 3 of 5.7 applied to  $\gamma$ , the set of such  $(\alpha, \ell)$  is non-empty: since  $\gamma \subset \text{int}(\beta'_j)$ ,  $\text{int}(\beta'_j) \cap \alpha \neq \emptyset$  for all such  $\alpha$ . Take a maximal such  $(\alpha, \ell)$ . If some  $\gamma' \subset \partial\beta'_j$  intersects  $\alpha$  transversally, then by 6.3 we have  $\gamma' \cap \gamma \neq \emptyset$ . So  $\alpha \subset \partial\beta'_j$ . By 6.3, there is no ltd  $(\alpha', \ell')$  with  $(\alpha, \ell) < (\alpha', \ell')$  and with  $\ell'$  to the left of  $x'_j$ , because that would contradict the maximality of  $(\alpha, \ell)$ . So by property 3 of 5.7 applied to components of  $\partial\alpha$ ,  $\beta'_j \subset \alpha$ . So  $\beta'_j = \alpha$ .  $\square$

**Corollary 11.8.** *The ordering on cross-sections is transitive.*

*Proof.* There might, perhaps, be a question about condition 2 of the ordering. So let  $b, b'$  be as in 11.7, with  $b < b'$  and  $b' < b'' = ((\beta''_1, \ell''_1), \dots, (\beta''_p, \ell''_p))$  and  $\beta_i = \beta''_k$ ,  $\beta'_j \neq \beta_i$ ,  $\beta'_j \cap \beta_i \neq \emptyset$ . if  $\beta'_j$  is ltd we have  $\beta'_j \subset \beta_i$  by 6.3. Then  $\beta'_j = \beta_i$  by 11.7, which is a contradiction. Now for any ltd  $(\alpha, \ell)$  in the decomposition with  $\ell \subset [x_i, x''_k]$  we have  $\alpha \cap \beta_i = \emptyset$  or  $\alpha \subset \beta_i = \beta''_k$  by 6.3. It follows that no component of  $\partial\beta'_j$  is transverse to  $\partial\beta_i$ , by property 3 of vertically efficient in 5.7 applied to  $\partial\beta'_j$ . So  $\beta_i \subset \beta'_j$  and  $\beta'_j \subset \beta_i$  by 11.6. Thus  $\beta_i = \beta'_j = \beta''_k$ . Then by 11.16 we have  $\beta_i = \beta'_j$  unless  $\beta_i$  is ltd on a segment centred at  $x_i$  for  $(\frac{1}{2}\Delta_2, r_2, s_2, K_2)$ .  $\square$

## 11.9 Bounded sets in the model manifold and cross-sections.

In order to bound vertical length of preimages under  $\Phi$  of bounded paths, we shall use the following. We use the language and notation of 11.4, 11.6. We continue to use the ltd parameter functions  $(\Delta_2, r_2, s_2, K_2)$ , the associated constant  $\nu_2 > 0$  and  $L = L(\Delta_2, r_2, s_2, K_2, \nu_2)$ . We also fix a model end manifold  $M(z_0, y_+)$  in  $M$ .

### Lemma

1. Given  $p_0$ , there is  $n_0$  such that the following holds. Suppose that  $(\beta, t)$  and  $(\beta', t')$  are such that  $M(\beta, \{t\})$  and  $M(\beta', \{t'\})$  are disjoint from model Margulis tubes  $T(\gamma_{**})$  with  $|\gamma_{**}| < \varepsilon_2$ .

1a) If  $\beta \cap \beta' = \emptyset$ , suppose that, for any  $\alpha$  with  $\beta \subset \alpha \subset S \setminus \beta'$ ,  $M(\beta, \{t\})$  and  $M(\beta', \{t'\})$  are not separated by totally ordered chains of  $p_0$  cross-sections for  $\alpha$ ,  $S \setminus \alpha$ , that is, a chain for  $\alpha$  above and below  $M(\beta, \{t\})$  or similarly for  $S \setminus \alpha$  and  $\beta'$ , or by a single chain of complete cross-sections.

1b) If  $\beta \cap \beta' \neq \emptyset$ , suppose that  $M(\beta, \{t\})$  is not separated from  $M(\beta', \{t'\})$  by chains of  $p_0$  cross-sections for  $\alpha$  above and below  $M(\beta, \{t\})$  for any  $\alpha \supset \beta$  with  $\alpha$  contained in the convex hull of  $\beta$  and  $\beta'$  nor by a chain of  $p_0$  complete cross-sections (that is, cross-sections for  $S$ ) if  $S$  is the convex hull of  $\beta$  and  $\beta'$ .

Then there is a connected set  $M_0 \subset M$  which is a union of  $\leq n_0$  sets which are either sets  $M(\alpha'_i, I_i)$  bounded by  $L$ , or subsets of model Margulis tubes of diameter  $\leq L$ , and  $M_0$  contains both  $M(\beta, \{t\})$  and  $M(\beta', \{t'\})$ .

2. Conversely, given  $n_0$  there exists  $p_0$ , such that, if  $M_0 \subset M$  contains a totally ordered chain of  $\geq p_0$  cross-sections for  $\alpha$ , for some  $\alpha$ , then  $M_0$  is not contained in a connected union of  $\leq n_0$  submanifolds  $M(\omega, I)$  bounded by  $L$ , and  $\leq n_0$  subsets of model Margulis tubes of diameter  $\leq L$ .

*Proof of 1.* Fix  $p_0$ , and suppose  $(\beta, t)$  and  $(\beta', t')$  are as in the statement of 1. Let  $\mathcal{P}$  be the set of all  $(\alpha, \ell)$  with  $\alpha \times \ell$  in our fixed vertically efficient decomposition of  $S \times [z_0, y_+]$  with respect to  $(\Delta_2, r_2, s_2, K_2, \nu_2)$ .

The basis of the proof is to show how to travel from  $S(b)$  to  $M(\beta, \{t\})$ , where  $b = b(E, \pm)$  is defined using the upper or lower boundary of a set  $E = E(b) \subset \mathcal{P}$ . Here,  $E$  is closed under  $\leq$  or  $\geq$ , and  $b$  is a cross-section for some  $\alpha$  which is above or below by  $(\beta, [\varphi_t])$ , and satisfies certain conditions, and similarly with  $\beta'$  replacing  $\beta$ .

To get into this situation, if  $(\beta', [\varphi_{t'}]) < (\beta, [\varphi_t])$  and  $\beta \neq \beta'$ , we take  $\alpha$  to be the convex hull of  $\beta$  and  $\beta'$ . If  $\beta'$  is not contained in  $\beta$ , then we define  $E$  to be the set of all  $(\omega, \ell) \leq (\beta, [\varphi_t])$  with  $\omega$  contained in  $\beta'$  or contained in  $\alpha \setminus \beta'$ . If  $\beta' \subset \beta$ ,  $\beta' \neq \beta$  then we interchange the role of  $\beta$  and  $\beta'$ . If  $\beta = \beta'$  and, without loss of generality,  $(\beta', [\varphi_{t'}]) \leq (\beta, [\varphi_t])$ , then we can take

$$E = \{(\omega, \ell) \in \mathcal{P} : (\omega, \ell) \leq (\beta', [\varphi_{t'}])\}$$

if there is no chain of  $p_0$  cross-sections between  $(\beta', [\varphi_{t'}])$  and  $(\beta, [\varphi_t])$ . We also define  $\alpha = \beta$  in this case. If there is such a chain, then we take  $\alpha = S$ , and

$$E = \{(\omega, \ell) \in \mathcal{P} : (\omega, \ell) \leq (\beta, [\varphi_t])\} \cup \{(\omega, \ell) : \omega \cap \beta = \emptyset\} \quad (11.9.1)$$

If  $\beta \cap \beta' = \emptyset$ , and there is no chain of  $p_0$  cross-sections for  $S \setminus \beta'$  below  $(\beta, [\varphi_t])$ , then we also define  $E$  by (11.9.1). Other cases are similarly dealt with by interchanging  $\beta$  and  $\beta'$ , or replacing  $\leq$  by  $\geq$ . In all cases, using 6.3, we see that if  $(\omega_i, \ell_i)$ ,  $i = 1, 2$ , are ltd with  $(\omega_1, \ell_1) < (\omega_2, \ell_2)$  and  $(\omega_2, \ell_2) \in E$ , then  $(\omega_1, \ell_1) \in E$  also. We then define  $b = b(E, +)$  to be the cross-section associated with the upper boundary of  $b$ .

If  $b$  is below  $(\beta, [\varphi_t])$ , we then define

$$E' = \{(\omega, \ell) \in \mathcal{P} : \omega \subset \beta, \} \cup \{(\omega, \ell) \in E : \omega \subset \alpha, \omega \cap \beta = \emptyset\}.$$

Define  $b' = b(E', +)$  to be the cross-section associated to the upper boundary of  $E'$ . Note that  $b$  has been chosen so that  $S(b)$  does not have long intersections with model Margulis tubes, except in the boundary around  $(\partial\alpha_*)$ . Then the same is true for  $S(b')$ , because if  $(\gamma, \ell) \in \mathcal{P}$  and  $(\gamma, \ell) \in E'$ , then either  $(\gamma, \ell') \in E$  for some  $\ell'$ , or  $(\gamma, \ell') \in E'$ . Also, if  $\alpha \setminus \beta \neq \emptyset$ ,  $d_{\alpha \setminus \beta}(x(E, +), x(E', +)) \leq L$ . So now, replacing  $b$  by  $b'$  if necessary, assume that any totally ordered chain of cross-sections for  $\alpha$  between  $b$  and  $(\beta, [\varphi_t])$  has  $\leq p_0$  elements and any chain between  $b$  and  $b'$  for  $\alpha_1$  properly contained in  $\alpha$  has  $\leq p_0$  elements. We continue to assume that  $b \leq b'$ , since the modifications if  $b' \leq b$  are trivial.

Now let

$$E'' = \{(\omega, \ell) \in E' : (\omega, \ell) \leq (\beta, [\varphi_t])\} \cup \{(\omega, \ell) \in E : \omega \cap \beta = \emptyset\}.$$

Let  $b'' = x(E'', +)$  be the upper boundary of  $E''$ . Then by definition of  $E'$ ,  $b \leq b'' \leq b'$ . Let  $b_0 < \dots < b_p$  be a maximal totally ordered chain of cross-sections for  $\alpha$  with  $b_0 = b$  and  $b_p = b''$ . Since the chain is maximal, there is a gap or loop  $\omega_i \subset \alpha$  such that  $(\omega_i, [x(b_i), x(b_{i+1})])$  is bounded by  $L = L(\Delta_2, r_2, s_2, K_2, \nu_2)$ . Otherwise we would be able to insert a cross-section strictly between  $b_i$  and  $b_{i+1}$ . We are now going to produce  $M_0$  by induction on  $p$  and on the topological type of  $\alpha$ . If  $p = 1$ , we simply take

$$M_0 = M(\omega, I_0) \cup M(\beta, \{t\}),$$

where

$$I_0 = \{s : \pi_\omega([\varphi_s]) \in [\pi_\omega(x(b_0)), \pi_\beta(x(b_1))]\}.$$

If  $p = 2$ , then the definitions of  $E$ ,  $E''$  ensure that we can assume that  $\beta_1 \cap \omega \neq \emptyset$  and  $\beta_1 \cap \beta \neq \emptyset$ , and we then define

$$M_0 = M(\omega, I_0) \cup M(\beta_1, I_1) \cup M(\beta, \{t\}),$$

where  $I_1$  is defined similarly to  $I_0$ , but with  $\beta_1$  replacing  $\omega$  and  $b_1, b_2$  replacing  $b_0$  and  $b_1$ . It is not immediately clear that we can proceed similarly in the case of  $p > 2$ . But let  $E_1$  be defined similarly to  $E$  in the case when  $(\beta', [\varphi_{t'}]) < (\beta, [\varphi_t])$  but using  $(\beta_1, x(b_1))$  in place of  $(\beta, [\varphi_t])$ , and then define  $E_2, E_3$  similarly to  $E', E''$ , but using  $E_1$  in place of  $E$ . These define cross-sections  $b_{1,j} = x(E_j, +)$  and  $b'_1$  with  $b_{1,1} \leq b_{1,3} < b_{1,2} \leq b'$ . Let  $\alpha_1$  be the convex hull of  $\beta_1$  and  $\beta$ . Thus,  $\alpha_1 \subset \alpha$ . A maximal chain of cross-sections between  $b_{1,1}$  and  $b_{1,3}$  for  $\alpha_1$  has

$\leq p - 1$  elements if  $\alpha_1 = \alpha$ , because otherwise the previous chain of length  $p$  is not maximal. If  $\alpha_1 = \alpha$  we can use the inductive hypothesis on  $p$ . If  $\alpha_1$  is strictly contained in  $\alpha$ , then, by the definition of the  $E_j$ ,  $d_{\alpha \setminus \alpha_1}(x(b_{1,1}), x(b_{1,2})) \leq L$  (where  $L = L(\Delta_2, r_2, s_2, K_2, \nu_2)$ ). So if  $\alpha_1 \neq \alpha$ , there is a bounded distance between the surfaces in the model manifold defined by  $b$  and  $b'$ , within the set  $M(\alpha \setminus \alpha_1, I)$  where

$$I = \{s : \pi_{\alpha \setminus \alpha_1}([\varphi_s]) \in \pi_{\alpha \setminus \alpha_1}([x(b), x(b')])\}.$$

So we are then reduced to finding a bounded path from at least one of  $S(b_{1,1})$  or  $S(b_{1,2})$  to  $M(\beta, t)$ , that is, in between two cross-sections for  $\alpha_1$ , and the proof of the inductive step is completed.

*Proof of 2.* If  $\alpha$  is a loop and we have a chain of  $p_0$  cross-sections for  $\alpha$ , then the cross-sections bound a piece of the model Margulis tube  $T(\alpha_{**})$  which can be made arbitrarily large by choice of  $p_0$ . Now suppose that  $\alpha$  is a gap. A cross-section for  $\alpha$  defines a surface in the model manifold with boundary components in the set  $(\partial\alpha)_{**}$ . The surfaces corresponding to two successive cross-sections  $b_1, b_2$  with  $b_1 < b_2$  either intersect different sets of model Margulis tubes in their interiors, or if neither intersects any Margulis tubes in the interior, both have single coordinates  $(\alpha, x_1)$  and  $(\alpha, x_2)$  such that  $\alpha$  is long,  $\nu$ -thick and dominant along  $[x_1, x_2]$ . So if  $p_0$  is sufficiently large, the set containing a chain of  $p_0$  successive cross-sections cannot be contained in  $\leq n_0$  submanifolds  $M(\omega, I)$  bounded by  $L$ .

□

## 11.10 Bounding vertical length.

We are now ready to bound vertical length of preimages under  $\Phi$ , of paths which are bounded in  $N$ . The idea is that parts of the image of  $\Phi$  are ordered in the correct way.  $\varepsilon_1$ - Margulis tubes are ordered the same way as the corresponding model Margulis tubes by 10.11. Images corresponding to ltd gaps are mapped in the correct order by 10.13. We use the

**Theorem** *Given  $\Delta_1$ , there is  $n_0$  depending only on  $\Delta_1$ , the topological type of  $N$ , and the constant  $c_0$  of 10.1.3, such that the following holds. Let  $\gamma$  be a path in the model manifold  $M$  such that  $\Phi(\gamma)$  is homotopic, fixing endpoints to a geodesic in  $N$  of length  $\leq \Delta_1$ . Then  $\gamma$  is homotopic, fixing endpoints, to a path in a union of  $\leq n_0$  sets  $M(\alpha'_i, I_i)$  which are bounded by  $L = L(\Delta_2, r_2, s_2, K_2, \nu_2)$ .*

*Proof.* It suffices to look at paths with both endpoints in a single end model manifold  $M(z_0(e), y_+(e))$ , because then  $\Phi(M(S(e), \{0\}))$  cannot separate  $\Phi(x)$  from the associated end for  $x \in M(z_0(e), y_+(e))$  sufficiently far, in the model metric, from  $M(S(e), \{0\})$ . Let one end of the path  $\gamma$  be in  $M(\beta, \{t\}) \subset M(z_0(e), y_+(e))$ , and the other end in  $M(\beta', \{t'\})$ . Suppose that  $\gamma$  is not homotopic to a path contained in  $\leq n_0$  sets. Then we apply 11.9 and use the equivalent description of bounded sets in terms of chains of cross-sections. If  $M(\beta', \{t'\})$  is in the model in a different end or in the model for  $W'$ , or  $(\beta, [\varphi_t]) < (\beta', [\varphi_{t'}])$  or  $(\beta', [\varphi_{t'}]) < (\beta, [\varphi_t])$ , then by 11.9 there is a set of  $2p_0$

cross-sections  $b_i$  ( $|i| \leq p_0$ ) either for some  $\alpha \supset \beta$  centred on a cross-section with  $(\beta, [\varphi_t])$  as a coordinate or for some  $\alpha \supset \beta'$  centred on a cross-section with  $(\beta', [\varphi_{t'}])$  as a coordinate, and in both cases with  $S(b_{\pm i})$  separating  $M(\beta, \{t\})$  from  $M(\beta', \{t'\})$ , or  $p_0$  cross-sections for  $\alpha$  below or above  $(\beta, [\varphi_t])$  and another  $p_0$  for  $S \setminus \alpha$  above or below  $(\beta', [\varphi_{t'}])$ , such that again the  $S(b_i)$  separate  $M(\beta', [\varphi_{t'}])$  from  $M(\beta, [\varphi_t])$ .

Let  $U$  be the neighbourhood of  $e$  bounded by  $S'$  of 10.12, if  $e$  is compressible.

Suppose that  $b_k$  is a totally ordered chain of cross-sections for  $\alpha$ , for  $0 \leq k \leq 3$ ,

$$b_k = ((\beta_{1,k}, x_{1,k}), \dots, (\beta_{n,k}, x_{n,k})).$$

Let  $S(b_k)$  be the surface for  $b_k$  as defined in 11.6. Then up to moving  $x_{j,k}$  a bounded  $d_{\beta_{j,k}}$  distance,  $\Phi(S(b_k))$  is a pleated surface  $h_b : \alpha \rightarrow N$  with

$$h_b|_{\omega} = f_{n(\omega)}|_{\omega}.$$

Here,  $\omega \subset \beta_{j,k}$  for some  $(j, k)$  — possibly with  $\omega = \beta_{j,k}$  — and  $\omega \times \{x_{j,k}\} \subset \omega \times \ell$ , where  $\omega \times \ell$  is one of the sets in the vertically efficient decomposition for  $(\Delta_2, r_2, s_2, K_2, \nu_2)$  and  $f_{n(\omega)}$  is one of the original pleated surfaces  $f_j$  of 8.3. So we have good bounds on the pleating locus with respect to  $(\Delta_1, r_1, s_1, K_1)$ . Then using 10.11 and 10.13, the  $h_{b_k}$  satisfy the hypotheses of the  $h_k$  in 11.5, with  $\alpha = \alpha' = S$ , apart perhaps for  $h_k(S) \subset U$ . But by 10.10, there is a bound on the number of complete cross-sections in any totally ordered chain which are not contained in  $U$ , in the cases when  $U$  is needed, that is,  $e$  is compressible.

Now suppose the first of the situations deduced from 11.9 occurs, that the  $b_i, |i| \leq p_0$  are such that  $b_0$  has  $(\beta, [\varphi_t])$  as a coordinate. The other cases are dealt with very similarly, so we shall just do this case. Fix any suitable  $\delta > 0$ . We claim that, by 11.5, there is an integer  $p_1$  which is bounded in terms of  $\varepsilon_2$  and  $(\Delta_2, r_2, s_2, K_2)$  such that, given any totally ordered chain of  $4p_1 + 1$  cross-sections for  $\alpha$ , the pleated surfaces for the first and the last cannot intersect the  $\delta$ -neighbourhood of the middle one. If  $\alpha = S$ , we can assume that the number of complete cross-sections which are not contained in  $U$ , is  $< p_1$ . So we now have at least  $p_1$  complete cross-sections on either side of the middle one. and the corresponding pleated surfaces satisfy the hypotheses of 11.10. Number cross-sections and pleated surfaces  $b_i, h_{b_i} = h_i, -p_1 \leq i \leq p_1$ . Suppose that

$$h_{-p_1}(\alpha) \cap B_\delta(h_0(\alpha)) \setminus T((\partial\alpha)_*, \varepsilon_1) \neq \emptyset \quad (11.10.1)$$

and fix  $i$  with  $-p_1 < i < 0$ . Apply 11.5 with  $h_i, h_0, h_j, h_{-p_1}$  replacing  $h_0, h_1, h_2, h_3$ , for varying  $j, 0 < j \leq p_1$ . Then either

$$B_\delta(h_i(\alpha)) \cap (h_0(\alpha) \cup h_{-p_1}(\alpha)) \setminus T((\partial\alpha)_*, \varepsilon_1) \neq \emptyset \quad (11.10.2)$$

or

$$B_\delta(h_j(\alpha)) \cap (h_0(\alpha) \cup h_{-p_1}(\alpha)) \setminus T((\partial\alpha)_*, \varepsilon_1) \neq \emptyset \text{ for } 0 < j \leq p_1. \quad (11.10.3)$$

But (11.10.3) is impossible for  $p_1$  sufficiently large, because, by the definition of a totally ordered set of cross-sections (11.6), it would imply the existence of  $\geq p_1$

different bounded geodesics and boundaries of  $\varepsilon_1$ -Margulis tubes intersecting a set of diameter  $C_2$ , for  $C_2$  depending only on the topological type and  $\varepsilon_1$ . Any two complete cross-sections differ by at least one Margulis tube or by at least one bounded loop, and recall that  $\varepsilon_1$  depends only on  $(\Delta_1, r_1, s_1, K_1)$ . Similarly, (11.10.2) cannot hold for all  $i$  if  $p_1$  is sufficiently large. So if  $p_1$  is sufficiently large, (11.10.1) cannot hold, as required. A similar argument works for  $h_{p_1}(\alpha)$ .

Then apply 11.5 again with  $h_3$  of 11.5 taken to be a pleated surface  $h'$  with domain  $\beta'$ . If  $p_1$  is sufficiently large given  $p_2$  we have, after renumberings, an ordered set of surfaces  $h_i(\alpha)$   $-2p_2 \leq i \leq 2p_2$  such that all the surfaces  $h_i(\alpha) \setminus T((\partial\alpha)_*, \varepsilon_1)$  are disjoint. Then by 11.5,  $h'(\beta')$  cannot intersect  $h_i(\alpha) \setminus T((\partial\alpha)_*, \varepsilon_1)$  for  $|i| \leq p_2$ . If  $p_2$  is sufficiently large given  $\Delta_1$  and  $\delta$ , the  $\Delta_1$  neighbourhood of  $\Phi(M(\beta, \{t\}))$  in  $N$  is contained in the region bounded by  $h_{\pm p_2}(\alpha)$  and the union of the Margulis tubes  $T(\zeta_*, \varepsilon_1)$  for  $\zeta \subset \partial\alpha$ . So then if  $\gamma_1$  is the geodesic segment homotopic to  $\Phi(\gamma)$  with endpoints fixed,  $\gamma_1$  has length  $> \Delta_1$ , giving the required contradiction.

□

### 11.11 Bounded diffeomorphism on Margulis tube boundaries.

We now claim that the map  $\Phi$  can be chosen to be a bounded diffeomorphism with bounded inverse between model and actual Margulis tubes, with respect to the model Riemannian metric and the hyperbolic metric on  $N$ . Since the model Margulis tube is an actual Margulis tube, and the right one up to bounded distortion, it suffices to make  $\Phi$  into a diffeomorphism with bounded inverse between the boundaries of the model and actual Margulis tubes. As usual, the constant  $\Lambda_1$  (enlarged from the previous one if necessary) depends, ultimately, only on the topological type of  $N_{d,W}$  and the constant  $c_0$ . So fix  $\gamma$ . It suffices to consider only  $\gamma$  with  $|\gamma_{**}| < \varepsilon_2$ . Consider all those  $(\alpha, \ell)$  in the vertically efficient partition  $\mathcal{P}$  with  $\gamma \subset \partial\alpha$ . It suffices to show that, for some  $\Delta_1$ , and given any  $M(\alpha, \ell)$  with  $\gamma \subset \partial\alpha$ , there is at least one  $(\alpha', \ell')$  between distance  $\Delta_1$  and  $2\Delta_1$  away along  $\partial T(\gamma_{**})$  which is bounded away in  $M$  itself: we can then apply 11.10 to see that  $\Phi(M(\alpha, \ell))$  and  $\Phi(M(\alpha', \ell'))$  are bounded apart, and hence the intersections with  $\Phi(T(\gamma_{**}))$  are bounded apart. But this is immediate, because if we fix  $M(\alpha, \ell)$  and some bounded neighbourhood  $M_1$  of  $M(\alpha, \ell)$  in the model manifold, only finitely many disjoint bounded sets  $M(\alpha', \ell')$  can intersect  $M_1$ . So by choosing  $\Delta_1$  large enough (assuming that  $\varepsilon_2$  is sufficiently small, as we may do) the claim is proved.

### 11.12 Bounding horizontal length in the model ends: the nature of the bounded set.

Now let  $\gamma$  be a path in  $M$  of bounded vertical length with lift  $\tilde{\gamma}$  to the universal cover. We need to show that if  $\Phi(\gamma)$  is homotopic, keeping endpoints fixed, to a geodesic  $\gamma_1$  of bounded length, then  $\gamma$  is homotopic in  $M$  to a path of bounded length. We now know that if  $\gamma_1$  is bounded, then  $\gamma$  lies in a bounded subset  $M_1$

of  $M$ , up to homotopy preserving endpoints, where the bounds on  $M_1$  depend on topological type, the ltd parameter functions (which depend only on topological type) and our constant  $c_0$  (of 10.1.3). We shall choose  $M_1$  in a particular way. First, suppose that  $M_1$  is in an end model manifold homeomorphic to  $S \times [0, 1]$  or  $S \times [0, \infty)$  for some finite type surface  $S$ . Theorem 11.10 implies that, if  $\gamma$  is sufficiently far from the model core, as we assume for the moment, it is possible to choose  $M_1$  so that the boundary of  $\partial M_1$  is the union of two sets which we call the *vertical boundary*  $\partial_v M_1$  and *horizontal boundary*  $\partial_h M_1$ , where these have the following properties. The vertical boundary  $\partial_v M_1$  is contained in the union of boundaries of model Margulis tubes. The horizontal boundary  $\partial_h M_1$  is a union of sets of the form  $\varphi_x^{-1}(S_{\beta,x}) = M(\beta, x)$ , and, by 11.9 each set  $M(\beta, x)$  is separated from  $\gamma$  by one or two totally ordered chains of cross-sections for  $\alpha$ , for some  $\alpha$ . There are three possibilities. A single totally ordered chain might encase  $M(\beta, x)$  and separate it from  $\gamma$ , or might encase  $\gamma$ , separating  $\gamma$  from  $M(\beta, x)$ , or might be a totally ordered chain for  $S$  which separates a component of  $\partial_h M_1$  from  $\gamma$ . We call components of  $\partial_h M_1$  of the first type *encased* and components of the last two types *separated*.

By 11.9, the length  $2p_1 + 1$  of the totally ordered chain or chains can be taken as large as we like be anything we like, by allowing  $M_1$  to have sufficiently large diameter in the model metric (but still bounded). We choose  $p_1 = k_0 p_0$ , where  $p_0$  is as in 11.10 and for  $k_0$  depending on the topological type of  $S$ . As in 11.10 we take  $p_0$  large enough that if a set of  $p_0$  cross-sections for  $\beta$  exists, then  $(\partial\beta)_*| < \varepsilon_1$  and there are model Margulis tubes round all components of  $\partial\beta$  — unless, of course,  $\partial\beta = \emptyset$ . Note that there is no a priori claim that the regions bounded by different chains of cross-sections are disjoint, but chains of cross-sections for  $\alpha$  and  $\alpha'$  can only interlink if one of  $\alpha$ ,  $\alpha'$  is contained in the other. So the number of interlinkings of different chains is bounded by the topological type of  $S$ . Also by 11.10, if  $p_0$  is large enough, the images under  $\Phi$  of  $S(b_{kp_0,\alpha})$ , for distinct integers  $k$ , are disjoint. We reindex these cross-sections as  $b_{k,\alpha}$ ,  $|k| \leq k_0$ . If  $k_0$  is large enough, given  $k_1$ , we can reduce some chains of cross-sections to  $b_{k,\alpha}$  for  $k \leq k_{0,\alpha}$  for some  $k_{0,\alpha} \geq k_1$  and  $\leq k_0$ , so that each chain is either encased by some other or disjoint from all others. Then we can discard any chains which does not encase  $\gamma$  and is encased by another which also does not encase  $\gamma$ , or is disjoint from some chain that does encase  $\gamma$ . Similarly if a chain for  $\alpha'$  encases a chain for  $\alpha$  which encases  $\gamma$ , then we can discard the chain for  $\alpha'$ . So now we have a set of disjoint chains  $b_{k,\alpha_i}$  ( $|k| \leq k_1$ ) for  $\alpha_i$ ,  $2 \leq i \leq r$ , some  $r$  bounded in terms of the diameter of  $M_1$ , encasing components of  $\partial_h M_1$ , and (probably) two chains  $b_{k,\alpha_1,1}$  and  $b_{k,\alpha_1,2}$  for some  $\alpha_1$  encasing  $\gamma$ . If  $\alpha_1$  does not exist, then there must be bounded paths from  $\gamma$  to  $M(S, \{z_0\})$  or  $M(S, \{y_+\})$ , possibly to both. In that case, we still take chains  $b_{k,\alpha_1,j}$  of some length  $\geq 1$ , and take  $\alpha_1 = S$ . So  $b_{0,\alpha_1,j}$  exists for  $j = 1, 2$ . Index so that  $S(b_{k,\alpha_1,j}) \subset M_1$  for  $k \geq 0$ ,  $j = 1, 2$ .

Now fix some suitable  $k_2 < k_1$ . Let  $M'_1$  be the intersection of  $M_1$  with the set bounded by  $S(b_{0,\alpha_1,1})$  and  $S(b_{0,\alpha_1,2})$  and model Margulis tubes round  $\partial\alpha_1$ . For  $2 \leq j \leq r$ , let  $M'_i$  be the set bounded by  $S(b_{\pm k_2,\alpha_i})$  and model Margulis



tubes round  $\partial\alpha_i$ . Thus

$$M'_i \subset M'_1 \text{ for } 2 \leq i \leq r.$$

We can now assume without loss of generality that

$$M_1 = M_1 \cap (M'_1 \setminus \cup_{i=2}^r M'_i), \quad (11.12.1)$$

and

$$\partial_h M_1 \subset S(b_{0,\alpha_1,1}) \cup S(b_{0,\alpha_1,2}) \cup \cup_{i=2}^r S(b_{\pm k_2,\alpha_i}). \quad (11.12.2)$$

The most obvious way to estimate length of a path in  $M_1$  is to take some cell decomposition of  $M_1$ , and count the number of cell boundaries crossed by the path. The most obvious way to make a cell decomposition is, to first decompose  $M_1$  into interval bundles by cutting along horizontal surfaces, and then to decompose each interval bundle by annuli. We can use annuli whose boundaries have bounded length. Essential intersections with the horizontal surfaces give a lower bound on length. The annuli do not have boundary in  $\partial M_1$ , but we are going to modify them to obtain surfaces with boundary in  $\partial M_1$ .

**Lemma 11.13.** *Given  $C_1$  there is  $C_2$  such that the following holds. Let  $M_2 \subset M_1$  be homeomorphic to  $\alpha \times [0, 1]$  for some  $\alpha \times [0, 1]$  and such that  $\partial\alpha \times [0, 1]$  is the intersection of  $M_2$  with  $\partial_v M_1$ . Let  $\zeta \subset M_2$  be a nontrivial simple closed loop of length  $\leq C_1$  in the model metric. Then there is a surface  $S(\zeta) \subset M_1$  of area and diameter  $\leq C_2$ , in the metric on  $S(\zeta)$  induced by the model metric, with boundary in  $\partial M_1$ , and such that  $\zeta$  is contained in  $S(\zeta)$  up to free homotopy in  $M_1$ . In fact, in the cover of  $M_1$  determined by  $M_2$ ,  $S(\zeta)$  lifts to an annulus  $A(\zeta)$  homotopic to  $\zeta$ .*

*Proof.* We start by considering an annulus  $A_1(\zeta) \subset M$  which is homotopic to  $\zeta$  and has one boundary component above  $M_1$ , that is, nearer the end, and other boundary component below  $M_1$ . For example, let  $\zeta_s$  be the geodesic on  $S_s$  which is freely homotopic to  $\varphi_s(\zeta)$ , and define

$$A_1(\zeta) = \cup_{s \in I} \varphi_s^{-1}(\zeta_s) \times \{s\}$$

for some suitable interval  $I$ . Now we consider  $T_1(\zeta) = A(\zeta) \cap M_1$ , and  $\partial T(\zeta) = A(\zeta) \cap \partial M_1$ . There is no reason why  $T(\zeta)$  should have bounded area or diameter. Nevertheless, the intersection of  $\partial T(\zeta)$  with  $\partial_v M_1$  is bounded, the number of components of intersection with  $\partial_h M_1$  is bounded, intersection with  $M_2$  is bounded and intersections with  $M_1$  of homotopy tracks between the boundary components of  $\partial A_1(\zeta)$  are bounded. The natural homotopy is given by

$$(s', z) \mapsto \varphi_{s'}^{-1} \circ \varphi_s(z) : \varphi_s^{-1}(\zeta_s) \rightarrow \varphi_{s'}^{-1}(\zeta_{s'}).$$

The definition of the model metric in Section 7 ensures that this homotopy is boundedly Lipschitz restricted to  $M_1$ . We now change  $\partial T(\zeta)$  on horizontal boundary pieces which are matched by this homotopy in  $M_1$ , and on the homotopy between them, so as to reduce length. We do not change  $T(\zeta)$  in  $M_2$ , and

do not need to, because we already have bounds there. The topological type of the surface  $T(\zeta)$  will not change, but the isotopy class of the embedded surface probably will. If  $\gamma_1$  and  $\gamma_2$  are horizontal pieces matched by the homotopy, with the homotopy entirely in  $M_1$ , we simply replace  $\gamma_j$  by a bounded path  $\gamma'_j$  with the same endpoints, such that the natural homotopy in  $M_1$  matches up  $\gamma'_1$  and  $\gamma'_2$  instead. Now there is a decomposition of the horizontal boundary of  $T(\zeta)$  into pieces which are paired by the homotopy, because every time a homotopy track enters  $M_1$ , it does so along horizontal boundary and then leaves again. The path  $\varphi_t^{-1}(\zeta)$  has bounded length so need not be involved in any changes. So performing this operation a finite number of times we have a surface  $S(\zeta)$  of bounded area and with of bounded length in the metric induced from the model metric. By construction, the components of  $\partial S(\zeta)$  in the horizontal boundary are homotopically nontrivial. The claim about  $A(\zeta)$  follows, because  $S(\zeta) \cap M_2 = A_1(\zeta) \cap M_2$  is homotopic to  $\zeta$ .  $\square$

#### 11.14 Bounding horizontal length: intersection number.

Suppose we have fixed a set of loops  $\zeta$ , such that the corresponding annuli cut the corresponding annuli into cells, and we have a bound on the length of the loops  $\zeta$  used. Then the surfaces  $S(\zeta)$ , and the horizontal surfaces used, cut  $M_1$  into cells. Let the set of surfaces be  $S_j$ ,  $1 \leq j \leq r$ . Then for a constant  $C_3 > 0$  depending only on the ltd parameter functions

$$|\gamma| \geq C_3 \text{Max}\{i(\gamma, S_j) : 1 \leq j \leq r\},$$

where  $i(.,.)$  denotes essential intersection number. The aim now is to obtain a parallel lower bound on  $|\gamma_1|$ , and hence an upper bound on  $|\gamma|$  in terms of  $|\gamma_1|$ . For this, we have to notice something about the paths  $\zeta$ , and have to be a little careful about the choices.

First, we can easily find a surface homotopic to  $\Phi(S_j)$  with boundary in  $\Phi(\partial M_1)$ , by choosing a maximal multicurve in  $S_j$  which is bounded in the model metric and using the pleated surface with this int its pleating locus, for example. We assume this surface is in fact  $\Phi(S_j)$ . If  $S_j$  is a horizontal surface then, as in 11.5, we can redefine  $\Phi$  inside Margulis tubes so that  $\Phi(S_j) \subset N_{\geq \varepsilon_1}$  and  $\Phi(\partial S_j) \subset \partial N_{\geq \varepsilon_1}$ . Then if  $T_1, T_2$  denotes the Margulis tubes intersected by  $\partial S_j$ ,  $\Phi(\partial S_j)$ ,  $(M, T_1, S_j, \gamma)$  is homotopy equivalent to  $(N, T_2, \Phi(S_j), \gamma_1)$  and so

$$i(\gamma, S_j) = i(\gamma_1, \Phi(S_j))$$

for any horizontal surface. It remains to obtain something similar for  $S(\zeta)$ . The key result is the following.

**Lemma 11.15.** *There is a constant  $L$  such that the following holds. There is a submanifold with boundary  $N_1$  of  $N$  and a continuous map  $\Phi_1 : M \rightarrow N$  homotopic to  $\Phi$  such that the following hold.  $\Phi$  is a homeomorphism in a neighbourhood of  $M$  with  $\Phi_1(\partial M_1) = \partial N_1$ ,  $\Phi_1(S(\zeta)) \subset N_1$  and  $\Phi$  is homotopic to  $\Phi_1$  under a homotopy which is trivial on  $M'_1$ .*

Note that there is no claim of a bound on the homotopy distance between  $\Phi$  and  $\Phi_1$ .

*Proof.*

We use the set-up of 11.12, in particular, (11.12.1) and (11.12.2). By [21], there is a surface  $S'(b_{k,\alpha_i})$  (or  $S'(b_{k,\alpha_1,j})$ ) in an arbitrarily small neighbourhood of  $\Phi(S(b_{k,\alpha_i}))$  (or  $\Phi(S(b_{k,\alpha_1,j}))$ ) if  $i = 1$ ) which is homeomorphic to  $\alpha_i$ . As in the proof of 11.10, we can also assume that these surfaces intersect  $\partial T(\zeta_*, \varepsilon_1)$  transversally for  $\zeta \subset \partial\alpha_i$ , and only in the homotopy class of the core loop  $\zeta$ . We take  $N_1$  to be the set homotopic to  $\Phi(M_1)$  bounded by the  $\varepsilon_1$ -model Margulis tubes which intersect  $\Phi(\partial_v M_1)$  and the surfaces  $S'(b_{\pm k_2, \alpha_i})$ ,  $i \geq 2$ , and  $S'(b_{0, \alpha_1, j})$ ,  $j = 1, 2$ . Since  $\Phi$  is  $\Lambda_1$ -Lipschitz, we can assume (by choice of  $p_0$ ) that the image under  $\Phi$  of the intersection of  $M_1$  with the region bounded by  $S(b_{k, \alpha_1, j})$  and  $S(b_{\pm k, \alpha_i})$  ( $i \geq 2$ ) is contained in the region bounded by  $S'(b_{k-1, \alpha_1, j})$  and  $S'(b_{\pm(k-1), \alpha_i})$  for all  $1 \leq k \leq k_1$ . In particular, this is true for the image under  $S(\zeta)$  of the intersection of  $S(\zeta)$  with this region. Similar facts hold for images under  $\Phi$  of interval bundles bounded by  $S(b_{k, \alpha_i})$  and  $S(b_{m, \alpha_i})$ , or by  $S(b_{k, \alpha_1, j})$  and  $S(b_{m, \alpha_1, j})$ . So we only need to change the definition of  $\Phi$  in the region bounded by  $S(b_{\pm 2, \alpha_1, j})$  and  $S(b_{\pm(k_2 \pm 2), \alpha_i})$ . We could, for example, first extend  $\Phi(S(\zeta))$  by attaching annuli and put the boundaries in  $S'(b_{-1, \alpha_1, j})$ ,  $S'(b_{\pm(k_2-1), \alpha_i})$ , and then compose with homeomorphisms to compress the interval bundles bounded by  $S'(b_{\pm 2, \alpha_1, j})$ ,  $S'(b_{\pm(k_2 \pm 2), \alpha_i})$  to the interval bundles bounded by  $S'(b_{2, \alpha_1, j})$ ,  $S(b_{0, \alpha_1, j})$ ,  $S(b_{\pm(k_2+2), \alpha_i})$ ,  $S'(b_{\pm(k_2), \alpha_i})$ . The resulting map  $\Phi_1$  is homotopic to  $\Phi$  via a homotopy which is trivial outside the region bounded by  $S(b_{\pm 2, \alpha_1, j})$  and  $S(b_{\pm(k_2 \pm 2), \alpha_i})$ .  $\square$

Now since  $\Phi_1$  is homotopic to a homeomorphism from  $M_1$  to  $N_1$ ,

$$i(\gamma_1, \Phi(S(\zeta))) = i(\gamma_1, \Phi_1(S(\zeta))) = i(\Phi_1(\gamma), \Phi_1(S(\zeta))) = i(\gamma, S(\zeta)).$$

Let  $M_2$  be any interval bundle in  $M_1$  which contains  $\zeta$  and in which  $|\zeta|$  is bounded. Let  $M'_2$  be the natural compactification of the cover of  $M_1$  corresponding to  $M_2$ . By this we mean:  $M_1$  is a sum of interval bundles, attached along subsurfaces. The universal cover of an interval bundle has natural compactification  $D \times [0, 1]$ , where  $D$  is the closed unit disc. Then (unless  $M_1 = M_2$  up to homotopy)  $M'_2$  is an infinite tree of  $M_2$  and infinitely many copies of  $\text{int}(D/\Gamma_j) \times [0, 1]$ , for subgroups  $\Gamma_j$  of  $\pi_1(M_2)$  (mostly for  $\Gamma_j$  the trivial group) attached along subsurfaces. Compactify by adding the copies of  $\partial D$  and points at each end. Then  $M'_2$  is naturally homeomorphic to  $M_2$ . Let  $\gamma_2$  be the lift of  $\gamma$  to  $M'_2$ . Then  $\gamma_2$  has infinitely many components. One component is a closed path if  $\gamma \subset M$ , but otherwise all components are paths with endpoints in  $\partial M'_2$ . The number of homotopically nontrivial paths, that is, not homotopic into the boundary, is the number of components of intersection of  $\gamma$  with  $M_2$ . As already noted, we have lift of  $S(\zeta)$  to  $A(\zeta)$  in  $M'_2$  which extends to the boundary, and  $(M'_2, A(\zeta))$  is homeomorphic to  $(M_2, A_2(\zeta))$ , where  $A_2(\zeta)$  is an annulus in  $M_2$  with boundary components in  $\partial M_2$ . Now clearly we have

$$i(\gamma, S(\zeta)) \geq i(M'_2, A(\zeta)) = i(M_2, A_2(\zeta)) = i(\gamma_2, \zeta).$$

For  $i(\gamma_2, \zeta)$ , we regard  $\gamma_2$  as an arc in  $\partial M_2$  and  $\zeta$  as being in the same boundary component. Each intersection gives a different element of the coset space  $\pi_1(M_2)/\langle \zeta \rangle$ . Each essential intersection between  $\gamma_1$  and  $S(\zeta)$  gives rise to an actual intersection, and a homotopy class of path along  $\gamma_1$  from an initial point to another point on  $\Phi_1(S) \cap \gamma_1$ , that is, from an initial point to a point on  $\Phi(S)$ . The intersections given by cosets in  $\pi_1(S)/\pi_1(S(\zeta))$  are distinct. So the bound on the length of  $\gamma_1$  and on the area of  $\Phi(S(\zeta))$  gives a bound on  $i(\gamma, \zeta)$ . We have this for all  $\zeta$ . Since we have already seen that  $i(\gamma, S_j)$  is bounded for horizontal surfaces, we have the required bound on  $|\gamma|$ .

### 11.16 Bounding horizontal length near $W'$ .

As usual, let  $W'$  denote the non-interval bundle part of the core  $W$ . Let  $\gamma$  be a path near  $W$  and  $\gamma_1$  the geodesic segment homotopic to  $\Phi(\gamma)$  with the same endpoints, as in 11.12. Choose a model piece  $M_1$  such that  $\gamma \subset M_1$  and  $\partial_h M_1$  has similar properties to those described in 11.12, but containing  $W'$  and intersecting the model manifolds of several ends. Choose the model piece large enough to include anything which was not previously included in a model piece  $M_1$  as in 11.12. So  $W'$  may be properly contained in  $M_1$ . Again, we assume that  $\gamma$  and  $\gamma_1$  are closed loops.

Now we construct surfaces in  $W'$  which decompose  $W'$  into cells, and with boundary in  $\partial W'$ . We look at intersections with the  $W_i$  of 7.10. Recall that  $W = W_n$ , and  $W_i$  decomposes into submanifolds  $W_{i-1}$ . In 7.10 we constructed a decomposition of  $W$  into submanifolds which were balls or interval bundles. Interval bundles adjacent to  $\partial W$  have been removed but there might be some interval bundles in the interior of  $W'$ . We use the annuli in the sets  $\Sigma_i$  of 7.11 to extend the decomposition into balls. So now we have an extended sequence  $W_j$ , with  $W_n = W'$  (renaming  $n$  of 7.10) such that the components of  $W_j$  decompose into the components of  $W_{j-1}$ , all components of  $W_0$  are balls, all components of  $W_r$  (for some  $r$ ) are either balls or interval bundles, and all components of  $\partial W_{j-1} \setminus \partial W_j$ , for  $j \leq r$ , are annuli. Now we want to apply the same method above with  $M_1 = W$  and suitable surfaces  $S_i$  obtained from the components of the  $\partial W_j$ . Instead of using components of  $\partial W_i \setminus \partial W_{i+1}$ , for  $i \leq n-2$ , we use surfaces with boundary in  $\partial W'$ . To do this, if  $S$  is a component of  $\partial W_i \setminus W_{i+1}$  and  $i \leq n-2$ , we form a union of two surfaces,  $T_{i+2}(S)$ , with boundary in  $\partial W_{i+2}$ , by taking a tubular neighbourhood of  $S$ , taking the boundary components of the tubular neighbourhood, adding surfaces in the boundary of tubular neighbourhoods of the adjoining components of  $\partial W_{i+1} \setminus \partial W_{i+2}$ . Inductively, if  $j < n$  we define  $T_{j+1}(S) = T_{j+1}(T_j(S))$ . We write  $T(S) = T_n(S)$ , which is a union of  $2^{n-i-1}$  surfaces, since  $S$  is two-sided and all  $W_i$  have oriented boundary.

Now we extend each surface  $T(S)$  across the remainder of  $M_1$ , using the surfaces of 11.13. We have bounds on the areas of the surfaces, as before. If these surfaces do not cut  $M_1$  into cells, then we also use some additional surfaces  $S(\zeta)$ . We may need to use some extra surfaces  $S(\zeta)$  for  $\zeta = \zeta_1 \cup \zeta_2$  of the following form:  $\zeta \subset \alpha$  for some  $M(\alpha, \ell) \subset M_1$ ,  $\zeta_1 \subset \partial T(S)$  for some surface

$T(S)$  constructed in  $W'$  and  $\zeta_2 \subset \cup_{i=1}^r \alpha_i$ , where  $\alpha_i \times \ell_i \subset S(e) \times [z_{e,0}, y_{e,+}]$ ,  $(\alpha_i, \ell_i) < (\alpha, \ell)$  and  $M_1 \cap M(\alpha_i, \ell_i) = \emptyset$ . We can choose a set of such  $\zeta$  such that the corresponding sets  $S(\zeta)$ , together with the sets  $T(S)$ , cut  $M_1$  into cells.  $\square$

## 12 Proofs of Theorems 1.1, 1.2, 1.3.

As noted in Section 1, Theorems 1.1 and 1.2 are derived from 1.3.

### 12.1 Proof of 1.3.

Item 1 of 1.3 was proved in 7.17, since we only take limits on the end parts of the geometric manifolds - and in 7.16 in the combinatorially bounded geometry Kleinian surface case. The existence of a coarse biLipschitz  $\Phi : M \rightarrow \overline{N}$  is entirely proved in sections 10 and 11, with the required bounds on the biLipschitz constant, in the case of all ends being geometrically infinite. In the case of a geometrically finite end, we need to show how to extend the map  $\Phi$  to the complement of the convex hull. For each geometrically finite end  $e_i$  of  $N_d$ , we choose an embedding  $f^i : S(e_i) \rightarrow N$  as  $f_3$  in 4.6, and the pleated surface  $f_{e_i,+}$  as  $f_4$  of 4.6. By 4.6, we therefore have bounds on  $d(\mu(e_i), [f_{e_i,+}])$ , and a bound on the hyperbolic length of homotopy tracks of a homotopy between  $f^i$  and  $f_{e_i,+}$ . So now if  $M = M(\mu(e_1), \dots, \mu(e_n))$  we have, by the results of sections 10 and 11, a coarse biLipschitz map  $\Phi : M \rightarrow N$  whose image has any surfaces  $f^i(S(e_i))$  in its boundary. The surfaces  $f^i(S(e_i))$  bound convex sets containing the convex hull of  $N$ , by construction. Perturb  $f^i$  so that  $f^i$  is smooth and the complementary components are still convex. Then perpendicular geodesics pointing outwards from  $f^i(S(e_i))$  hit the boundary of the  $t$ -neighbourhood in exactly one point. This defines a map from  $f^i(S(e_i))$  to the  $t$ -neighbourhood which expands distances by  $e^t$ . So we can adjust  $\Phi$  to a map onto  $N \cup \Omega(\Gamma)/\Gamma$  which is coarse biLipschitz with respect to the adjusted metric, and maps a set  $S(e_i) \times [u_i - 1, u_i]$  in the model manifold to the corresponding component of  $N \setminus CH(N)$ , with sets  $\{x\} \times [u_i - 1, u_i]$  mapping to geodesic rays, and mapping  $S(e_i) \times \{u_i - 1 + u'\}$  to the set distance  $t(u')$  from  $f^i(S(e_i))$  for some fixed function  $t : [0, 1] \rightarrow (0, \infty)$ . In particular,  $\Phi$  is coarse biLipschitz with respect to the hyperbolic metric on the preimage of the convex subset of  $N$  bounded by the surfaces  $f^i(S(e_i))$ .

### 12.2 Proof of 1.1.

Let  $N_1$  and  $N_2$  be two homeomorphic hyperbolic 3-manifolds with finitely generated fundamental groups and the same end invariants. Let  $M$  be the model manifold for both. Let  $\Phi_i : M \rightarrow N \cup \Omega(\Gamma_i)/\Gamma_i$  be the map of 1.3, constructed in 12.1, with lifts  $\widetilde{\Phi}_1$  and  $\widetilde{\Phi}_2$ . Then the set-valued map  $\tilde{\Phi} = \widetilde{\Phi}_2 \circ \widetilde{\Phi}_1^{-1} : H^3 \rightarrow H^3$  is coarse biLipschitz with respect to the metric which agrees with the hyperbolic metric inside the convex hulls, except in bounded (hyperbolic metric)

neighbourhoods of the convex hull boundaries, and is an adjustment of the hyperbolic metric outside. Let  $f^{j,i} : S(e_j) \rightarrow N_i$  be as in 12.1, and  $U_i$  be the lift of  $\cup_j f^{j,i}(S(e_j))$  to  $H^3$ . Then  $U_i \subset H^3 \setminus \widetilde{CH(N_i)}$  and (although we do not need this)  $U_i$  is a bounded hyperbolic distance from a component of  $\widetilde{CH(N_i)}$ . Taking a small perturbation, we can assume that  $U_1$  and  $U_2$  have smooth boundaries, and let  $U_{i,t}$  be the surface hyperbolic distance  $t$  from  $U_i$ , further from  $\widetilde{CH(N_i)}$ , as in 12.1. The component of  $H^3 \setminus U_{i,t}$  containing  $\widetilde{CH(N_i)}$  is convex. So perpendicular geodesic segments between  $U_i$  and  $U_{i,t}$  only intersect these surfaces at the endpoints. Now we have chosen  $\tilde{\Phi}_1$  and  $\tilde{\Phi}_2$  so that  $\tilde{\Phi}$  maps  $U_1$  to  $U_2$ , perpendicular segments to perpendicular segments preserving length,  $U_{1,t}$  to  $U_{2,t}$  and  $\Omega(\Gamma_1)$  to  $\Omega(\Gamma_2)$ .

Now we claim that  $\tilde{\Phi}$  is coarse biLipschitz with respect to the hyperbolic metric. It suffices to show that  $\tilde{\Phi}$  and  $\tilde{\Phi}^{-1}$  are coarse Lipschitz and we only need to consider the maps on  $H^3 \setminus \widetilde{CH(N_i)}$ ,  $i = 1, 2$ . Here, we only need to consider the norms of the derivatives with respect to the hyperbolic metric. We have a splitting of the tangent space at each point into the tangent to the geodesic segment and the tangent to the  $U_{i,t}$  foliation. The derivative of  $\tilde{\Phi}$  maps one splitting to the other and  $\tilde{\Phi}_i$  maps the tangent space to  $S(e_j) \times \{u_j - 1 + u'\}$  to the tangent space to  $U_{i,t(u')}$ , with derivative  $e^{t(u')}$  times the identity. Meanwhile, the tangent space to  $\{x\} \times [u_{j-1}, u_j]$  is mapped to the geodesic tangent space with the same derivative for both  $\tilde{\Phi}_1$  and  $\tilde{\Phi}_2$ . So the derivative of  $\tilde{\Phi}$  is actually the identity with respect to suitable framings on the domain and range tangent spaces, and  $\tilde{\Phi}$  is Lipschitz, as required.

Let  $x_t$  be a geodesic ray in  $H^3$  from a basepoint  $x_0$ , parametrised by length, and converging to a point  $x_\infty \in \partial H^3$ . First we claim that  $\lim_{t \rightarrow \infty} \tilde{\Phi}(x_t)$  exists and is a single point, remembering that  $\tilde{\Phi}(x_t)$  is set-valued in general. In fact,  $\tilde{\Phi}(x_t)$  has uniformly bounded diameter in the hyperbolic metric, because  $\tilde{\Phi}$  is coarse biLipschitz. Then the limit exists, because the coarse biLipschitz properties of  $\Phi_i$  with respect to hyperbolic metric imply that  $\tilde{\Phi}(\{x_t : t \in [n, n+1]\})$  has bounded hyperbolic diameter and geometrically small Euclidean diameter, because the distance of the set from a fixed basepoint is  $\geq n/\Lambda$ , where  $\Lambda$  is the product of the biLipschitz constants for  $\Phi_1$  and  $\Phi_2$ . Moreover, this also shows that the limit is uniform. Since  $\tilde{\Phi}$  is coarse biLipschitz with respect to the hyperbolic metric, it is biLipschitz with respect to the Euclidean metric on  $\partial H^3$ , using the unit ball model for  $H^3 \cup \partial H^3$ , and hence quasi-conformal. Then we can change the conformal structure on  $\Omega(\Gamma_2)$  so that with respect to this new conformal structure,  $\tilde{\Phi}$  is conformal, because the ending invariants of  $N_1, N_2$  are the same. So then we can find a quasiconformal map  $\psi : \Omega(\Gamma_2) \rightarrow \Omega(\Gamma_2)$  which extends continuously to the identity on the complement in  $\partial H^3$ , so that  $\psi \circ \tilde{\Phi}$  is conformal on  $\Omega(\Gamma_1)$ . It is equivariant, and hence if not conformal on the limit set, there is a nontrivial  $\Gamma_2$ -invariant line-field on the limit set, where  $\Gamma_2$  is the covering group of  $N_2$ . So then by Sullivan's theorem [57] the extension of  $\psi \circ \tilde{\Phi}$  is conformal, and  $N_1$  and  $N_2$  are isometric.

### 12.3 Proof of 1.2.

Let  $[\mu_1, \dots, \mu_r]$  be any admissible invariant for the topological type of a hyperbolic manifold  $N$  with ends  $e_i$  for  $N_d$ ,  $1 \leq i \leq r$ . Let  $[y_{1,n}, \dots, y_{r,n}] \in (\prod_{i=1}^r \mathcal{T}(S(e_i)))/G$  with  $y_{i,n} \rightarrow \mu_i$ . Then by 7.17,  $M_n = M(y_{1,n}, \dots, y_{r,n})$  converges geometrically to  $M(\mu_1, \dots, \mu_r)$  with a suitable choice of basepoint. By 1.3, if  $N_n$  denotes the hyperbolic manifold homeomorphic to  $N$  under an end-preserving homeomorphism between  $N_{n,d}$  and  $N_d$  then the map  $M_n \rightarrow N_n$  is coarse biLipschitz restricted to the preimage of  $C(N_n)$  for a biLipschitz constant uniform in  $n$ . So taking geometric limits, the  $N_n$  converge, with suitable basepoint, to a hyperbolic manifold  $N_\infty$ . The topological type remains the same, because we can find  $m_n \rightarrow \infty$  and a sequence of  $m_n$  successive pleated surfaces in each end of  $N_n$  with hyperbolic distance  $\geq 1$  between any adjacent pair. Also using these surfaces, we have closed geodesics arbitrarily far out in the end  $e_i$  converging to  $\mu_i$  in the limit. So  $(\mu_1, \dots, \mu_r)$  is the invariant of  $N_\infty$ .

## References

- [1] Abikoff, W.: The real analytic theory of Teichmüller Space. Springer Lecture Notes in Math. 820 (1980).
- [2] Agol, I.: Tameness of hyperbolic 3-manifolds, arXiv:math.GT/0405568.
- [3] Ahlfors, L. and Bers, L.: The Riemann Mapping Theorem for Variable Metrics, Ann. of Math. 72b (1960) 385-404.
- [4] Bonahon, F.: Cobordism of automorphisms of surfaces, Annales Scientifiques de l'cole Normale Suprieure Sr. 4, 16 no. 2 (1983), p. 237-270.
- [5] Bonahon, F.: Bouts des Variétés hyperboliques de dimension trois, Ann. of Math. 124 (1986) 71-158.
- [6] Bowditch, B.: Intersection numbers, and the hyperbolicity of the curve complex, to appear in J. reine angew. math..
- [7] Bowditch, B.: Geometric models for hyperbolic manifolds. Preprint Southampton 2005.
- [8] Bowditch, B.: End invariants of hyperbolic 3-manifolds. Preprint, Southampton 2005.
- [9] Brock, J. and Bromberg, K.: On the density of geometrically finite Kleinian groups. Acta Math 192 (2004) 33-93.
- [10] Brock, J., Canary, R., Minsky, Y.: The classification of Kleinian surface groups II: the Ending Lamination Conjecture, arXiv:math.GT/0412006.
- [11] Bromberg, K.: Projective structures with degenerate holonomy and the Bers density conjecture, arXiv:math.GT/0211402.

- [12] Bromberg, K. and Souto, J.: The density conjecture: A prehistoric approach, in preparation.
- [13] Buser, P.: Geometry and Spectra of Compact Riemann Surfaces. Birkhouser 1992.
- [14] Calegari, D. and Gabai, D.: Shrinkwrapping and the taming of hyperbolic 3-manifolds, J. Amer. Math. Soc. 19 (2006) 385-446.
- [15] Canary, R.D., Epstein, D.B.A. and Green, P.: Notes on notes of Thurston, Analytical and Geometric Aspects of Hyperbolic Space, C.U.P. 1987, L.M.S. Lecture Notes Series no 111, 3-92.
- [16] Canary, R.D.: Ends of hyperbolic 3-manifolds, J. Amer. Math. Soc., 6 (1993) 1-35.
- [17] Cannon, J.W. and Feustel, C.D.: Essential embedding of annuli and Möbius bands in 3-manifolds, TAMS 215 (1976), 219-239
- [18] Epstein, D.B.A. and Marden, A.: Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces, Analytical and Geometric Aspects of hyperbolic space, CUP 1987, London Math. Soc. Lecture Notes Series no. 111, 113-254.
- [19] Evans, R.: The ending lamination conjecture for hyperbolic 3-manifolds with slender end-invariants, to appear in Pac. J. Math..
- [20] Fathi, A., Laudenbach, F. and Poénaru, V. et al. Travaux de Thurston sur les surfaces, Astérisque 66-67 (1979).
- [21] Freedman, M., Hass, J. and Scott, P.: Least area incompressible surfaces in 3-manifolds, Invent. Math. 71 (1983), 609-642.
- [22] Hatcher, A. and Thurston, W.P.: A presentation of the mapping class group of an orientable surface, Topology 19 (1980) 221-237.
- [23] Hempel, J.: 3-Manifolds. Annals of Math. Studies, Princeton University Press, 1976.
- [24] Kim, I., Lecuire, C. and Ohshika, K.: Convergence of freely decomposable Kleinian groups. Preprint, 2004
- [25] Lecuire, C.: An extension of Masur domain, preprint, 2004.
- [26] , Masur, H.: Measured Foliations and handlebodies, Ergodic Theory and Dynamical Systems 6 (1986), 99-116.
- [27] Masur, H. and Minsky, Y.: Geometry of the Complex of Curves I, Hyperbolicity, Invent. Math. 138 (1999) 103-149.
- [28] Masur, H. and Minsky, Y.: Geometry of the Complex of Curves II, Hierarchical Structure, Geom. Funct. Anal. 10 (2000) 902-974.



- [29] McCullough, D.: Compact submanifolds of 3-manifolds with boundary, Quart. J. Math. Oxford Ser. 37(147) (1986) 299-307.
- [30] McCullough, D. and Miller, A.: Homeomorphisms of 3-manifolds with Compressible Boundary, Memoirs of the AMS 344 (1986).
- [31] McCullough, D., Miller, A. and Swarup, G.A.: Uniqueness of cores of non-compact 3-manifolds, J. London Math. Soc. 52 (1985) 548-556.
- [32] McMullen, C.: Iteration on Teichmüller Space. Invent. Math. 99 (1990) 425-454.
- [33] Minsky, Y.: Teichmüller geodesics and ends of hyperbolic 3-manifolds, Topology 32 (1993) 624-647.
- [34] Minsky, Y.: Harmonic maps into hyperbolic 3-manifolds, Trans Amer. Math. Soc. 332 (1992) 607-632.
- [35] Minsky, Y.: Quasi-projections in Teichmüller Space, J. Reine Angew. Math. 473 (1996), 121-136.
- [36] Minsky, Y.: The classification of punctured torus-groups, Annals of Math. 149 (1999) 559-626.
- [37] Minsky, Y.: Bounded Geometry for Kleinian Groups, Invent. Math. 146 (2001) 143-192.
- [38] Minsky, Y.: The classification of Kleinian Surface Groups, I: Models and Bounds, preprint 2002. arXiv:math.GT/0302208.
- [39] Morgan, J.W.: On Thurston's Uniformisation Theorem for Three-Dimensional Manifolds, Chapter V, The Smith Conjecture, edited by J.W. Morgan and H. Bass, Academic Press 1984.
- [40] Namazi, Hossein: Heegard splittings and hyperbolic geometry. Thesis, Stony Brook University, 2005.
- [41] Ohshika, K.: On limits of quasi-conformal deformations of Kleinian groups. Math.Z. 201 (1989) 167-176.
- [42] Ohshika, K.: Limits of geometrically tame Kleinian groups, Invent. Math. 99 (1990) 185-203.
- [43] Ohshika, K.: Ending laminations and boundaries for deformation spaces of Kleinian groups, J. London Math. Soc. 42 (1990), 111-121.
- [44] Ohshika, K.: Realising end invariants by limits of minimally parabolic, geometrically finite groups. Preprint 2005. arXiv:math.GT/0504546.
- [45] Otal, J-P.: Courant géodésiques et produits libres, Thèse d'état, Université Paris-Sud, Orsay 1988.

- [46] Otal, J-P.: Thurston's hyperbolization of Haken manifolds, *Surveys of Differential Geometry III* (1986) Int. Press.
- [47] Otal, J-P.: Les géodésiques fermées d'une variété hyperbolique en tant que noeuds, 95-104, *Kleinian Groups and Hyperbolic 3-Manifolds*, Proceedings of the Warwick workshop, September 2001, eds Y. Komori, V. Markovic, C. Series, Cambridge U.P. 2003.
- [48] Rafi, K.: *Hyperbolic 3-manifolds and Geodesics in Teichmüller Space*. Thesis, SUNY at Stony Brook 2001.
- [49] Rafi, K.: A characterization of short curves of a Teichmüller geodesic. arXiv:math.GT/0404227v2 (2004).
- [50] Rees, M.: Views of Parameter Space: Topographer and Resident. *Astérisque* 288 (2003).
- [51] Rees, M.: An alternative approach to the ergodic theory of measured foliations on surfaces, *Ergod. Th. and Dynam Sys.* 1 (1981) 461-488.
- [52] Scott, G.P.: Finitely generated 3-manifold groups are finitely presented, *J. London Math. Soc.* 6 (1973) 437-440.
- [53] Scott, G.P.: Compact submanifold of 3-manifolds, *J. London Math. Soc.* 7 (1973) 246-250.
- [54] Soma, T: Existence of polygnal wrapping in hyperbolic 3-manifolds. Preprint, Tokyo Denki (2005).
- [55] Souto, J.: A note on the tameness of compact 3-manifolds, *Topology* 44 (2005).
- [56] Sullivan, D.P.: A finiteness theorem for cusps. *Acta Math.* 147 (1981) 289-299.
- [57] Sullivan, D. P.: Quasiconformal homeomorphisms and dyanmics II. Structural Stability implies hyperbolicity for Kleinian groups. *Acta Math* 155 (1985) 243-260.
- [58] Thurston, W.T.: *The Geometry and Topology of 3-manifolds*, notes, 1979.
- [59] Thurston, W.T.: Hyperbolic structures on 3-manifolds, I: deformation of acylindrical manifolds. *Ann. of Math.* 124 (1986) 203-246.
- [60] Thurston, W.T.: Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle. arXiv:math.GT/9801045 (1998).
- [61] Thurston, W.T.: Hyperbolic structures on 3-manifolds, III: Deformations of 3-manifolds with incompressible boundary. arXiv:math.GT/9801058 (1998).

- [62] Waldhausen, F.: Eine Verallgemeinerung des Schleitfestes. *Topology* 6 (1967), 501-504.
- [63] Waldhausen, F.: On irreducible 3-manifolds which are sufficiently large, *Ann. of Math.* 87 (1968) 56-88.