

# Vector bundles near negative curves: moduli and local Euler characteristic

E. Ballico

E. Gasparim

T. Köppe

## Abstract

We study moduli spaces of vector bundles on a two-dimensional neighbourhood  $Z_k$  of an irreducible curve  $\ell \cong \mathbb{P}^1$  with  $\ell^2 = -k$  and give an explicit construction of these moduli as stratified spaces. We give sharp bounds for the local holomorphic Euler characteristic of bundles on  $Z_k$  and prove existence of families of bundles with prescribed numerical invariants.

## 1 Introduction

We study moduli spaces of rank-2 bundles on a two-dimensional neighbourhood of an irreducible curve  $\ell \cong \mathbb{P}^1$  with negative self-intersection  $\ell^2 = -k \neq 0$ . We are interested in the behaviour of bundles over a small analytic neighbourhood of  $C$  inside a smooth surface  $Z_k$ , and in coherent sheaves near the singular point of the surface  $X_k$  obtained from  $Z_k$  by contracting the curve  $\ell$ . For this “local” problem of bundles near  $\ell$  it is enough to focus on vector bundles over the total space of  $\mathcal{O}_{\mathbb{P}^1}(-k)$ . Hence we take  $Z_k := \text{Tot}(\mathcal{O}_{\mathbb{P}^1}(-k))$ , where  $\ell \subset Z_k$  is the zero section. We write  $\pi: Z_k \rightarrow X_k$  for the map that contracts  $C$  to a point. We give an explicit construction and a stratification of the moduli of rank-2 bundles on  $Z_k$ .

A bundle  $E$  over  $Z_k$  has *splitting type*  $(j_1, \dots, j_r)$  if  $E|_\ell \cong \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(j_i)$  with  $j_1 \leq \dots \leq j_r$ . For the moduli problem we concentrate on the case of rank-2 bundles  $E$  with vanishing first Chern class; in this case the splitting type of  $E$  must be  $(-j, j)$  for some  $j \geq 0$ , and for short we say that  $E$  has splitting type  $j$ . We then define

$$\mathcal{M}_j(k) = \{E \rightarrow Z_k : E|_\ell \cong \mathcal{O}(j) \oplus \mathcal{O}(-j)\} / \sim$$

for the moduli (stack) of bundles over  $Z_k$  of splitting type  $j$ . We then prove:

**Theorem 4.2.** *The generic set of  $\mathcal{M}_j(k)$  is a complex projective space  $\mathbb{P}^{2j-k-2}$  minus a closed subvariety of codimension at least 2.*

However,  $\mathcal{M}_j(k)$  is empty if  $k > 2j - 2$ , and it is non-Hausdorff for  $j > k$ . To stratify  $\mathcal{M}_j(k)$  into Hausdorff components, we need numerical invariants. For a vector bundle  $E$  over  $Z_k$ , we define the Artinian sheaf  $Q_E$  on  $X_k$  by the exact sequence

$$0 \rightarrow \pi_* E \rightarrow (\pi_* E)^{\vee\vee} \rightarrow Q_E \rightarrow 0. \quad (1)$$

$E$  has two independent numerical invariants, which we call the *height* and the *width*, by analogy with the case of instanton bundles, cf. [Ga4].

**Definition 1.1.** We define the *height* and *width* of  $E$  by

$$\mathbf{h}_k(E) := \text{length } R^1\pi_*E \quad \text{and} \quad \mathbf{w}_k(E) := \text{length } Q_E .$$

We then show:

**Theorem 4.6.** *The pair  $(\mathbf{h}_k, \mathbf{w}_k)$  stratifies  $\mathcal{M}_j(k)$  into Hausdorff components.*

We remark that this pair of invariants gives the coarsest stratification of  $\mathcal{M}_j(k)$  into Hausdorff components. One can also decompose  $\mathcal{M}_j(k)$  by local second Chern classes  $c := c_2(E) - c_2(\pi_*(E)^{\vee\vee})$ . However,  $\mathcal{M}_j(k)$  has non-Hausdorff subspaces with fixed  $c$ . See [BG1] for the case of an exceptional curve, i.e.  $k = 1$ . We explicitly calculate sharp bounds for the invariants  $(\mathbf{h}, \mathbf{w})$  as follows:

**Theorems 2.7 and 2.12.** *Let  $E$  be a holomorphic rank-2 vector bundle over  $Z_k$  with  $c_1 = 0$  and splitting type  $j > 0$ . Let  $n_1 = \lfloor \frac{j-2}{k} \rfloor$  and  $n_2 = \lfloor \frac{j}{k} \rfloor$ . Then the following bounds are sharp:*

$$\begin{aligned} j - 1 &\leq \mathbf{h}_k(E) \leq (j - 1)(n_1 + 1) - kn_1(n_1 + 1)/2 , \\ 0 &\leq \mathbf{w}_k(E) \leq (j + 1)n_2 - kn_2(n_2 + 1)/2 , \end{aligned}$$

and  $\mathbf{w}_1(E) \geq 1$ .

**Definition 1.2** ([Bl, Def. 3.9]). Let  $\sigma(\tilde{X}, \ell) \rightarrow (X, x)$  be a resolution of an isolated quotient singularity. Let  $\tilde{\mathcal{F}}$  be a reflexive sheaf on  $\tilde{X}$ , and set  $\mathcal{F} := (\sigma_*\tilde{\mathcal{F}})^{\vee\vee}$ ; notice that there is a natural injection  $\sigma_*\tilde{\mathcal{F}} \hookrightarrow \mathcal{F}$ . Then the *local holomorphic Euler characteristic* of  $E$  is

$$\chi(x, \tilde{\mathcal{F}}) := \chi((\tilde{X}, \ell), \tilde{\mathcal{F}}) := h^0(X, \mathcal{F}/\sigma_*\tilde{\mathcal{F}}) + \sum_{i=1}^{n-1} (-1)^{i-1} h^0(X, R^i\sigma_*\tilde{\mathcal{F}}) .$$

**Corollary 2.13.** *Let  $E$  be a rank-2 bundle over  $Z_k$  of splitting type  $j > 0$  and let  $j = nk + b$  such that  $0 \leq b < k$ . The following are sharp bounds for the local holomorphic Euler characteristic of  $E$ :*

$$j - 1 \leq \chi(E, \ell) \leq \begin{cases} n^2k + 2nb + b - 1 & \text{if } k \geq 2 \text{ and } 1 \leq b < k , \\ n^2k & \text{if } k \geq 2 \text{ and } b = 0 , \end{cases}$$

and

$$j \leq \chi(E, \ell) \leq j^2 \text{ for } k = 1 .$$

We then consider the question of existence of vector bundles. We recall the concept of an *admissible sequence* (Definition 3.1) and prove the following existence result:

**Theorem 3.2.** *Fix an admissible sequence  $\{a(i, l)\}_{i=1}^t$  and let  $E$  and  $F$  be rank- $r$  vector bundles on  $\hat{\ell}$  with  $\{a(i, l)\}_{i=1}^t$  as an associated admissible sequence. Then there exists a flat family  $\{E_s\}_{s \in T}$  of rank- $r$  vector bundles on  $\hat{\ell}$  parametrised by an integral variety  $T$  and  $s_0, s_1 \in T$  with  $E_{s_0} \cong E$  and  $E_{s_1} \cong F$  such that  $E_s$  has  $\{a(i, l)\}_{i=1}^t$  as admissible sequence for every  $s \in T$ .*

In Section 2 we calculate numerical invariants for bundles near negative curves. In Section 3 we describe the method of *balancing* bundles and show the existence of families of bundles with prescribed numerical invariants. In Section 4 we study moduli of rank-2 bundles on  $Z_k$ .

Our calculations were often performed on a computer using an implementation of the height and width computations as described in [GKP] written with the computer algebra software *Macaulay 2*. The program can be downloaded from <http://www.maths.ed.ac.uk/~s0571100/Instanton/>.

## 2 Bounds

Let  $Z_k$  be the total space of  $\mathcal{O}_{\mathbb{P}^1}(-k)$  and  $\ell \cong \mathbb{P}^1$  the zero section, so that  $\ell^2 = -k$ ; we write  $\ell_N$  for the  $N^{\text{th}}$  infinitesimal neighbourhood of  $\ell$ ,  $\widehat{\ell} = \varprojlim \ell_N$  for the formal neighbourhood of  $\ell$  in  $Z_k$ , and  $\mathcal{O}(j)$  for the line bundle on  $Z_k$  or on  $\widehat{\ell}$  that restricts to  $\mathcal{O}_{\mathbb{P}^1}(j)$  on  $\ell$ . A vector bundle  $E$  has *splitting type*  $(j_1, \dots, j_r)$  if  $E|_{\ell} \cong \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(j_i)$  with  $j_1 \geq \dots \geq j_r$ . A result of Griffiths [Gr] implies that  $E$  splits on  $\widehat{\ell}$  if  $j_1 - j_r \leq k + 1$ .

**Lemma 2.1.** *Let  $Z$  be a smooth surface containing a curve  $\ell \cong \mathbb{P}^1$  with  $\ell^2 = -k$ . Let  $E$  be a rank- $r$  vector bundle on  $Z$  of splitting type  $j_1 \geq j_2 \geq \dots \geq j_r$ , and assume that  $j_1 - j_r \leq k + 1$ . Then  $E$  splits on the formal neighbourhood  $\widehat{\ell}$  of  $\ell$ , that is,  $E|_{\widehat{\ell}} \cong \bigoplus_{i=1}^r \mathcal{O}_{\widehat{\ell}}(j_i)$ .*

*Proof.* This follows from [Gr, Propositions 1.1 and 1.4]: The point is that  $\text{Hom}(E, E)|_{\ell}$  is a direct sum of line bundles of degree  $\geq -(j_1 - j_r)$ , so the first-order infinitesimal extensions  $H^1(\widehat{\ell}; \text{Hom}(E, E) \otimes \mathcal{O}(-\ell))$  vanish.  $\square$

It is known that a reflexive sheaf on a surface quotient singularity  $\mathbb{C}^2/G$  is a direct sum of the tautological sheaves obtained from the irreducible representations of  $G$ . For cyclic quotient singularities  $\frac{1}{r}(1, a)$  these are just the eigensheaves  $\mathcal{O}(i)$  of the group action, such that  $\pi_*\mathcal{O} = \bigoplus_{i=0}^{r-1} \mathcal{O}(i)$ . We now study the holomorphic invariants  $\mathbf{h}_k(E) = l(R^1\pi_*E)$  and  $\mathbf{w}_k(E) = l(Q_E)$  of  $E$  as defined in Definition 1.1.

*Remark 2.2.* In principle, we need the Theorem on Formal Functions [Ha, p. 276], to calculate  $\mathbf{w}$  and  $\mathbf{h}$ . However, since holomorphic bundles on  $Z_k$  are algebraic (see [Ga3, Lemmas 2.1 and 2.3]), the limit stabilises at a finite order and it is enough to compute the cohomology on a fixed infinitesimal neighbourhood  $\ell_N$ , where  $N$  is not too small (roughly twice the degree of the extension class).

We record the obvious fact that heights and widths behave additively for bundles that split on the formal neighbourhood  $\widehat{\ell}$ :

**Proposition 2.3.** *If  $E|_{\widehat{\ell}} \cong \bigoplus_{i=1}^r \mathcal{O}(j_i)$ , then*

$$\mathbf{w}_k(E) = \sum_{i=1}^r \mathbf{w}_k(\mathcal{O}(j_i)) \quad \text{and} \quad \mathbf{h}_k(E) = \sum_{i=1}^r \mathbf{h}_k(\mathcal{O}(j_i)) . \quad \square$$

**Corollary 2.4.** *Let  $E$  be a holomorphic rank- $r$  vector bundle over  $Z_k$  such that  $E|_{\ell} \cong \bigoplus_{i=1}^r \mathcal{O}_{\ell}(j_i)$ , with  $j_1 \geq j_2 \geq \dots \geq j_r$  and  $j_r - j_1 \geq -k - 1$ . Then*

$$\mathbf{w}_k(E) = \sum_{i=1}^r \mathbf{w}_k(\mathcal{O}(j_i)) \quad \text{and} \quad \mathbf{h}_k(E) = \sum_{i=1}^r \mathbf{h}_k(\mathcal{O}(j_i)) .$$

*Proof.* By Lemma 2.1,  $E|_{\widehat{\ell}} \cong \bigoplus_{i=1}^r \mathcal{O}_{\widehat{\ell}}(j_i)$ . Hence  $E$  has the same invariants as the split bundle, and the result follows from Proposition 2.3.  $\square$

We fix once and for all coordinate charts  $U = \mathbb{C}_{z,u}^2$  and  $V = \mathbb{C}_{\zeta,v}^2$  on  $Z_k$ , glued by  $\zeta = z^{-1}$  and  $v = z^k u$ . In these charts the bundle  $\mathcal{O}(j)$  has the transition matrix  $z^{-j}$ .

Bundles with vanishing first Chern class correspond to instantons under the Kobayashi–Hitchin correspondence and are of special interest for applications to physics (see [LT]). For the case  $c_1 = 0$ , we also calculate the lower bounds for the numerical invariants  $\mathbf{h}$  and  $\mathbf{w}$ .

The second author proved ([Ga1]) that holomorphic bundles on  $Z_k$  are algebraic extensions of line bundles. By [Ga1, Theorem 3.3], a bundle  $E$  that is an extension

$$0 \rightarrow \mathcal{O}(j_1) \rightarrow E \rightarrow \mathcal{O}(j_2) \rightarrow 0 \quad (2)$$

(with  $j_1 \leq j_2$ ) has transition matrix  $T = \begin{pmatrix} z^{-j_1} & p \\ 0 & z^{-j_2} \end{pmatrix}$  in canonical coordinates, where the extension class  $[p] \in \text{Ext}^1(\mathcal{O}(j_1), \mathcal{O}(j_2))$  may be represented by a *canonical form*

$$p(z, u) = \sum_{r=1}^{\lfloor (j_2 - j_1 - 2)/k \rfloor} \sum_{s=kr+j_1+1}^{j_2-1} p_{rs} z^s u^r . \quad (3)$$

## 2.1 Heights

We begin with the computation of the height of a line bundle:

**Lemma 2.5.** *Assume  $j \geq 0$  and let  $n = \lfloor \frac{j-2}{k} \rfloor$ . Then*

$$\mathbf{h}_k(\mathcal{O}(-j)) = \begin{cases} (j-1)(n+1) - kn(n+1)/2 & \text{if } j \geq 2, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.*  $R^1\pi_*\mathcal{O}(j) = \varprojlim H^1(\mathcal{O}_{\ell_n}(j))$ , with surjective restriction maps. The result comes from the exact sequences

$$0 \rightarrow H^1(\mathcal{O}_{\ell}(-n\ell) \otimes \mathcal{O}(j)) \rightarrow H^1(\mathcal{O}_{\ell_{n+1}}(j)) \rightarrow H^1(\mathcal{O}_{\ell_n}(j)) \rightarrow 0$$

together with

$$H^1(\mathcal{O}_{\ell}(-n\ell) \otimes \mathcal{O}(j)) = H^1(\mathbb{P}^1; \mathcal{O}(j + nk)) ,$$

which gives  $\mathbf{h}_k(\mathcal{O}(j)) = \sum_{n=0}^{\infty} (j-1-nk)^+$ , where  $+$  means the sum of positive terms only.  $\square$

This result together with Proposition 2.3 allows us to compute the heights of split bundles. For non-split bundles with  $c_1 = 0$  and splitting type  $j := j_2 = -j_1$ , we use the following:

**Theorem 2.6.** *Let  $E$  be the non-split bundle represented in canonical form by  $(j, p)$ , and let  $m > 0$  be the smallest exponent of  $u$  appearing<sup>1</sup> in  $p$ . With  $\mu = \min(m, \lfloor \frac{j-2}{k} \rfloor)$ , we have*

$$l(R^1\pi_*E) = \mu \left( j - 1 - k \frac{\mu - 1}{2} \right) .$$

*Proof.* By Lemma 2.8 below, a cocycle in  $H^1(E)$  has the canonical representation on the  $U$ -chart

$$\sum_{r=0}^{\lfloor \frac{j-2}{k} \rfloor} \sum_{s=kr-j+1}^{-1} \begin{pmatrix} a_{rs} \\ 0 \end{pmatrix} z^s u^r .$$

In this representation, every monomial term  $(a_{rs}z^s u^r, 0)$  with  $r < m$  represents a non-trivial cocycle by Lemma 2.9. Lastly, all terms with  $r \geq m$  are coboundaries by Lemma 2.10.  $\square$

Putting the split and the non-split cases together, we get sharp bounds on the heights:

---

<sup>1</sup>A rank-2 bundle whose extension class is a polynomial independent of  $u$  is in fact not of splitting type  $j$ , but of a lower splitting type.

**Theorem 2.7.** *Let  $E$  be a rank-2 bundle over  $Z_k$  of splitting type  $j > 0$ . Set  $n = \lfloor \frac{j-2}{k} \rfloor$ . The following bounds are sharp:*

$$j - 1 \leq \mathbf{h}_k(E) \leq (j - 1)(n + 1) - k(n + 1)n/2 .$$

*Proof.* The upper bound is attained by the split bundle. In this case, apply Lemma 2.5 and Proposition 2.3.

For the lower bound, note that the expression in Theorem 2.6 is always less than in or equal to the split case, and that the global minimum is  $j - 1$  (for  $\mu = 1$  in Theorem 2.6, attained by  $p(z, u) = uz$ ).  $\square$

We finish this subsection by proving the details of Theorem 2.6:

**Lemma 2.8.** *Every 1-cocycle in  $H^1(E)$  has a representative of the form*

$$\sum_{r=0}^{\lfloor \frac{j-2}{k} \rfloor} \sum_{s=kr-j+1}^{-1} \begin{pmatrix} a_{rs} \\ 0 \end{pmatrix} z^s u^r ,$$

with  $a_{rs} \in \mathbb{C}$ . In particular, every 1-cochain represented by  $\begin{pmatrix} a_{rs} \\ 0 \end{pmatrix} z^s u^r$  with  $r, s \geq 0$  is a coboundary.

*Proof.* Let  $\sigma$  be a 1-cocycle and let  $\sim$  denote cohomological equivalence. A power series representative for a 1-cochain has the form

$$\sigma = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{\infty} \begin{pmatrix} a_{rs} \\ b_{rs} \end{pmatrix} z^s u^r ,$$

with  $a_{rs}, b_{rs} \in \mathbb{C}$ . The 1-cochain  $s_1 = \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \begin{pmatrix} a_{rs} \\ b_{rs} \end{pmatrix} z^s u^r$  is holomorphic in  $U$ , hence is a coboundary. Hence

$$\sigma \sim \sigma - s_1 = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \begin{pmatrix} a_{rs} \\ b_{rs} \end{pmatrix} z^s u^r .$$

Now let  $T = \begin{pmatrix} z^j & p \\ 0 & z^{-j} \end{pmatrix}$  be the transition function of  $E$ , so that after a change of coordinates,

$$T\sigma = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \begin{pmatrix} z^j a_{rs} + p b_{rs} \\ z^{-j} b_{rs} \end{pmatrix} z^s u^r .$$

However, given that  $s_2 = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \begin{pmatrix} 0 \\ z^{-j} b_{rs} \end{pmatrix} z^s u^r$  is holomorphic in  $V$ ,

$$T\sigma \sim T\sigma - s_2 = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \begin{pmatrix} z^j a_{rs} + p b_{rs} \\ 0 \end{pmatrix} z^s u^r ,$$

and going back to the  $U$ -coordinate chart,

$$\sigma = T^{-1}T\sigma \sim \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \begin{pmatrix} a_{rs} + z^{-j} p b_{rs} \\ 0 \end{pmatrix} z^s u^r .$$

But  $p$  contains only terms  $z^k$  for  $k \leq j-1$ , therefore  $z^{-j}p$  contains only negative powers of  $z$ . Renaming the coefficients we may write

$$\sigma = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \binom{a'_{rs}}{0} z^s u^r$$

for some  $a'_{rs} \in \mathbb{C}$ , and consequently  $T\sigma = \sum_{r=0}^{\infty} \sum_{s=-\infty}^{-1} \binom{z^j a'_{rs}}{0} z^s u^r$ . Here each term  $a'_{rs} z^{j+s} u^r$  satisfying  $j+l \leq kr$  is holomorphic in the  $V$ -chart. Subtracting these holomorphic terms we are left with an expression for  $a$  where the index  $s$  varies as  $kr - j + 1 \leq s \leq -1$ . This in turn forces  $r \leq \lfloor \frac{j-2}{k} \rfloor$ , giving the claimed expression for the 1-cocycle.  $\square$

**Lemma 2.9.** *Let  $E$  be the bundle over  $Z_k$  represented in canonical form by  $(j, p)$  and  $m > 0$  the smallest exponent of  $u$  appearing in  $p$ . If  $\mu = \min(m, \lfloor \frac{j-2}{k} \rfloor)$ , then*

$$l(R^1 \pi_* E) \geq \mu \left( j - 1 - k \frac{\mu - 1}{2} \right) .$$

*Proof.* Assume  $m \leq \lfloor \frac{j-2}{k} \rfloor$ . Let  $\sigma = (a, 0)$  denote a 1-cocycle where  $a = z^s u^r$  with  $0 \leq r \leq m-1$  and  $kr - j + 1 \leq s \leq -1$  (due to Lemma 2.8). We claim that  $\sigma$  represents a non-zero cohomology class. In fact, since  $\sigma$  is not holomorphic in  $U$ , for  $\sigma$  to be a coboundary there must exist a  $U$ -coboundary  $\alpha$  such that  $\sigma + \alpha$  is holomorphic in  $V$ . Hence, there must exist a polynomial  $X$ , holomorphic on  $U$ , making the expression  $z^j a + pX$  holomorphic on  $V$ . However,  $z^j a$  is not holomorphic on  $V$ . Moreover, by the choice of  $m$ , no term in  $pX$  cancels  $z^j a$ . Consequently, no choice of  $X$  solves the problem of holomorphicity on  $V$ . Hence  $l(R^1 \pi_* E)$  is at least the number of independent cocycles of the form  $\sigma = (a, 0)$ , where  $a = z^s u^r$  with  $0 \leq r \leq m-1$  and  $kr - j + 1 \leq s \leq -1$ . There are  $m((j-1) - k(m-1)/2)$  such terms.

On the other hand, if  $m > \lfloor \frac{j-2}{k} \rfloor$ , then by an analogous argument, none of the monomial terms in the canonical cocycle form of Lemma 2.8 are coboundaries, and so the same formula holds with  $m$  replaced by  $\lfloor \frac{j-2}{k} \rfloor$ .  $\square$

**Lemma 2.10.** *Under the same conditions as in Lemma 2.9,*

$$l(R^1 \pi_* E) = \mu \left( j - 1 - k \frac{\mu - 1}{2} \right) .$$

*Proof.* We need to show that all remaining terms, namely  $(a_{rs} z^s u^r, 0)$  with  $m < r \leq \lfloor \frac{j-2}{k} \rfloor$  and  $kr - j + 1 \leq s \leq -1$ , represent coboundaries. For a fixed  $\sigma = (a_{rs} z^s u^r, 0)$  we choose a polynomial  $X \in \Gamma(\mathcal{O}(U))$  as follows: We select the monomial term appearing in  $p$  whose degree in  $u$  is  $m$  and whose degree in  $z$  is minimal, say  $l_0$ , and call it  $p_{ml_0} z^{l_0} u^m$ . We write  $p = p_{ml_0} z^{l_0} u^m + p'$  and set

$$X = -\frac{a_{rs}}{p_{ml_0}} z^{j+s-l_0} u^{r-m} .$$

Notice that since  $j+s-l_0 \geq 0$  and  $r-m > 0$ ,  $X$  is holomorphic on  $U$ . Therefore  $\sigma \sim \sigma + (0, X)$ . Now, changing coordinates,

$$T(\sigma + (0, X)) \sim T \begin{pmatrix} a_{rs} z^s u^r \\ X \end{pmatrix} = \begin{pmatrix} z^j & p \\ 0 & z^{-j} \end{pmatrix} \begin{pmatrix} a_{rs} z^s u^r \\ X \end{pmatrix} = \begin{pmatrix} z^j a_{rs} z^s u^r + pX \\ z^{-j} X \end{pmatrix} = \begin{pmatrix} p'X \\ z^{-j} X \end{pmatrix} .$$

Since  $z^{-j} X = -(a_{rs}/p_{ml_0}) z^{s-l_0} u^{r-m}$  has negative degree in  $z$ , it is clearly holomorphic on  $V$ , and consequently  $T(\sigma + (0, X)) \sim (p'X, 0)$ . Moreover, any terms in  $p'X$  containing only negative powers of  $z$  are holomorphic on  $V$  and can be removed as well.

OBSERVATION:  $p'X$  is left only with terms  $z^l u^i$  for which  $l + j + s - l_0 \geq 0$ . //

Changing back to  $U$ -coordinates,

$$\sigma = T^{-1}T\sigma \sim \begin{pmatrix} z^{-j} & -p \\ 0 & z^j \end{pmatrix} \begin{pmatrix} p'X \\ 0 \end{pmatrix} = \begin{pmatrix} z^{-j}p'X \\ 0 \end{pmatrix},$$

where

$$z^{-j}p'X = -\frac{a_{rs}}{p_{ml_0}} \sum_{i=m}^{\lfloor \frac{j-2}{k} \rfloor} \sum_{l_0 \neq l=ki-j+1}^{j-1} p_{il} z^l u^i z^{s-l_0} u^{r-m}$$

is holomorphic on  $U$  if  $l + s - l_0 \geq 0$  holds, which follows exactly from the observation.  $\square$

## 2.2 Widths

Again we begin by computing the width of *line* bundles:

**Lemma 2.11.** *Assume  $j \geq 0$  and let  $n = \lfloor \frac{j}{k} \rfloor$ . Then*

$$\begin{aligned} \mathbf{w}_k(\mathcal{O}(j)) &= (j+1)n - kn(n+1)/2, \\ \mathbf{w}_k(\mathcal{O}(-j)) &= 0. \end{aligned}$$

*Proof.* The length  $l(Q)$  equals the dimension of  $Q_0^\wedge$  as a  $\mathbb{C}$ -vector space, where  $0 \in X_k$  is the singular point. Since  $Q$  is defined by the sequence (1),  $Q_0^\wedge$  is the cokernel of the map  $(\pi_*\mathcal{O}(j))_0^\wedge \rightarrow (\pi_*\mathcal{O}(j))_0^{\vee\vee\wedge}$ . By the theorem on formal functions,

$$(\pi_*\mathcal{O}(j))_0^\wedge = \varprojlim_N H^0(\ell_N; \mathcal{O}(j)|_{\ell_N}) =: M.$$

But the limit stabilises at a finite stage, and it suffices to compute  $H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})$  for large  $N$ . Therefore  $l(Q)$  equals  $\dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})^{\vee\vee} - \dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})$  as  $k_0$ -modules. Since  $\ell_N$  is closed and of cohomological dimension 1, it follows from Serre duality that

$$H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})^\vee = \text{Ext}^1(\mathcal{O}(j), \omega) = H^1(\omega \otimes \mathcal{O}(-j)) = H^1(\mathcal{O}(-j-2)).$$

Now, using Lemma 2.5 with  $j+2$  instead of  $j$ , we get

$$\dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})^\vee = \dim H^1(\mathcal{O}(-j-2)) = (j+1)(n+1) - k(n+1)n/2$$

if  $j \geq 0$ , and 0 otherwise. Consequently, for  $j \geq 0$ ,

$$\begin{aligned} l(Q) &= \dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})^{\vee\vee} - \dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N}) \\ &= \dim H^0(\ell_N; \mathcal{O}(j)|_{\ell_N})^\vee - \dim H^0(\ell; \mathcal{O}(j)|_\ell) = (j+1)(n+1) - k(n+1)n/2 - (j+1). \end{aligned}$$

$\square$

**Theorem 2.12.** *Let  $E$  be a rank-2 bundle over  $Z_k$  of splitting type  $j$ . Then the following bounds are sharp: For  $j > 0$  and with  $n = \lfloor \frac{j}{k} \rfloor$ ,*

$$0 \leq \mathbf{w}_k(E) \leq (j+1)n - kn(n+1)/2, \text{ and } \mathbf{w}_1(E) \geq 1.$$

*Furthermore, for all  $0 < j < k$ ,  $\mathbf{w}_k(E) = 0$  for all bundles  $E$  (and necessarily  $k > 1$ ).*

*Proof.* The upper bound is realised by the split bundle, so the right-hand side follows from Lemma 2.11 and Proposition 2.3.

To calculate the lower bound for  $\mathbf{w}_k(E)$ , we use the bundle given in canonical coordinates by transition matrix  $\begin{pmatrix} z^j & zu \\ 0 & z^{-j} \end{pmatrix}$ . This bundle is generic and therefore attains the lower bound. Direct computation as described in [GKP, Section 6] shows that  $\mathbf{w}_{k>1}(E) = 0$  and  $\mathbf{w}_1(E) = 1$ .  $\square$

## 2.3 Local Euler characteristic

By summing up the results for heights and widths and using Definition 1.2 we obtain bounds on the local Euler characteristic of a rank-2 bundle  $E$  on  $Z_k$  near  $\ell$ . Due to the occurrence of integer part functions, it will be useful express the splitting type  $j$  of  $E$  as  $j = nk + b$ , with  $0 \leq b < k$ .

**Corollary 2.13.** *Let  $E$  be a rank-2 bundle over  $Z_k$  of splitting type  $j$  with  $j > 0$  and let  $j = nk + b$  as above. The following are sharp bounds for the local holomorphic Euler characteristic of  $E$ :*

$$j - 1 \leq \chi(E, \ell) \leq \begin{cases} n^2k + 2nb + b - 1 & \text{if } k \geq 2 \text{ and } 1 \leq b < k, \\ n^2k & \text{if } k \geq 2 \text{ and } b = 0, \end{cases}$$

and

$$j \leq \chi(E, \ell) \leq j^2 \text{ for } k = 1.$$

*Proof.* It follows directly from the definitions that for a curve  $\ell^2 = -k$  inside a smooth surface,  $h^0(X; E/\sigma_*E) = \mathbf{w}_k(E)$  and  $h^0(X; R^1\sigma_*E) = \mathbf{h}_k(E)$ , so the result follows by adding the results of Theorems 2.7 and 2.12.

To aid the computation, note that for  $2 \leq b < k$  we have  $\lfloor \frac{j}{k} \rfloor = \lfloor \frac{j-2}{k} \rfloor = n$ , while for  $b = 0, 1$  we have  $\lfloor \frac{j}{k} \rfloor = n$  and  $\lfloor \frac{j-2}{k} \rfloor = n - 1$ . We find that for  $b = 0, 1$ ,  $\chi(E, \ell) \leq n^2k + 2nb$ , and we can absorb the case  $b = 1$  into the other case.

The upper bound is attained by the split bundle  $E = \mathcal{O}(-j) \oplus \mathcal{O}(j)$ , while the lower bound is attained by the “generic” bundle with extension  $p(z, u) = zu$  as in the proof of Lemma 2.11.  $\square$

We intend to study numerical invariants of bundles over more general exceptional loci in future papers.

## 3 Balancing

We now consider the question of constructing vector bundles with specified numerical invariants. We use the technique of *balancing bundles* to prove the existence of bundles over  $Z_k$  with certain prescribed numerical invariants. These techniques were used in [BG2] to prove the existence of bundles over  $Z_1$  with any prescribed numerically admissible invariants, and in [BG3] some properties of balancing on  $Z_2$  were given.

Given two bundles  $E$  and  $E'$  of splitting type  $(j_1, \dots, j_r)$  and  $(j'_1, \dots, j'_r)$ , we say that  $E$  is *more balanced* than  $E'$  if  $j_1 - j_r \leq j'_1 - j'_r$ . The advantage of balancing a bundle is that we control the numerical invariants at each step, and we only need to compute numerical invariants for a smaller range of bundles.

The simplest case of balancing is for rank-2 bundles and goes as follows. If  $j_1 - j_2 \leq k - 1$ , we have won and we stop. If  $j_2 \leq j_1 - k$  we make the construction of [BG3], namely an elementary transformation with respect to  $\mathcal{O}_\ell(j_2)$  and obtain a new more balanced bundle with splitting type  $(j_1, j_2 + k)$ . We may also compare the invariants of the two bundles; at the end we reduce to a case with  $j_1 - j_2 \leq k - 1$ . We now describe how to balance bundles of rank  $r \geq 2$ . Let  $E$  be a rank- $r$  vector bundle over  $Z_k$  of splitting type  $j_1 \geq j_2 \geq \dots \geq j_r$ . We say that  $E$  is *balanced* if  $j_1 \leq j_r + k - 1$ . The objective is to balance  $E$ . Balancing associates to  $E$  the following data:

1. A positive integer  $t$  (the number of steps);
2. a finite sequence of  $r$ -tuples of non-increasing integers  $\{j(i, l)\}$  with  $1 \leq i \leq t$  and  $1 \leq j \leq r$  (the splitting types) satisfying:
  - (a)  $j(1, l) = j_l$  for  $1 \leq l \leq r$  (the splitting of the bundle  $E$ ),
  - (b)  $\sum_{l=1}^r j(i, l) = \sum_{l=1}^r j(1, l) + ki - k$  for  $2 \leq i \leq t$  (change of splitting produced by an elementary transformation), and
  - (c)  $j(t, 1) \leq j(t, r) + k - 1$  (arrive at a balanced bundle);
3. a chain  $E_1, \dots, E_t$  of vector bundles, starting with  $E_1 = E$ , where  $E_i$  has splitting type  $j(i, 1) \geq j(i, 2) \geq \dots \geq j(i, r)$  for  $1 \leq i \leq t$ .

**Definition 3.1.** A sequence  $\{j(i, l)\}$  of  $r$ -tuples of integers satisfying the numerical properties 1, 2 and 3 above is called an *admissible sequence*. The sequence  $\{j(i, l)\}_{i=1}^t$  of splitting types of the bundles  $E_i$  obtained in balancing the bundle  $E$  is then called the *admissible sequence associated to  $E$* .

Balancing a bundle  $E$  proceeds as follows: Set  $E_1 := E$  and  $j(1, l) = j_l$  for  $1 \leq l \leq r$ . If  $j(1, 1) \leq j(1, r) + k - 1$  we have won, we set  $t := 1$  and stop. Otherwise,  $j(1, 1) \geq j(1, r) + k$ . Choose a surjective homomorphism  $\rho: E|_\ell \rightarrow \mathcal{O}_\ell(j_r)$  and make the corresponding elementary transformation  $\mathbf{r}: E \rightarrow \mathcal{O}_\ell(j_r)$ . Set  $E_2 := \ker \mathbf{r}$ . Since  $j_r \leq j_{r-1}$ , we have  $\ker \rho \cong \bigoplus_{i=1}^{r-1} \mathcal{O}_\ell(j_i)$ , and  $\ker \mathbf{r}|_\ell$  fits in the exact sequence

$$0 \rightarrow \mathcal{O}_\ell(j_r + k) \rightarrow \ker \mathbf{r}|_\ell \rightarrow \bigoplus_{i=1}^{r-1} \mathcal{O}_\ell(j_i) \rightarrow 0. \quad (4)$$

We call  $j(2, 1) \geq j(2, 2) \geq \dots \geq j(2, r)$  the *splitting type* of  $E_2$ . In particular,  $\sum_{l=1}^r j(2, l) = \sum_{l=1}^r j(1, l) + k$ . If  $j(2, 1) \leq j(2, r) + k - 1$ , then we have won and we stop. If  $j(2, 1) \geq j(2, r) + k$ , then we obtain in the same way a bundle  $E_3$  with splitting type  $j(3, 1) \geq \dots \geq j(3, r)$  such that  $j(3, 1) \leq j(2, 1) \leq j(1, 1)$ ,  $j(3, r) \geq j(2, r) \geq j(1, r)$ . We continue this process, thus obtaining bundles  $E_2, E_3, \dots, E_i, \dots$  with splitting type  $j(i, 1) \geq \dots \geq j(i, r)$ ,  $j(i, 1) \leq j(i-1, 1)$ ,  $j(i, r) \geq j(i-1, r)$  and  $\sum_{l=1}^r j(i, l) = \sum_{l=1}^r j(1, l) + ki - k$  for all integers  $i \geq 2$  for which the bundle  $E_i$  is defined. We must show that this procedure stops, i.e. that after finitely many steps we arrive at a bundle  $E_i$  such that  $j(i, 1) \leq j(i, r) + k - 1$ . At the same time we will show that, in a suitable sense, each bundle  $E_x$ ,  $x \geq 2$ , which is defined is more balanced than the preceding one  $E_{x-1}$ . Call  $a_1 \geq \dots \geq a_r$  the splitting type of the bundle  $T := \mathcal{O}_\ell(j_r + k) \oplus \bigoplus_{i=1}^{r-1} \mathcal{O}_\ell(j_i)$ . Since  $j_1 \geq j_r + k$  by assumption, we have  $a_1 = j_1$ . If  $j_{r-1} \geq j_r + k$ , then  $a_r = j_r + k$  and hence  $a_1 - a_r < j_1 - j_r$ . If  $j_{r-1} \leq j_r + k - 1$ , then  $a_r = j_{r-1}$  and hence  $a_1 - a_r = j_1 - j_{r-1}$ . Hence in the latter case we have  $a_1 - a_r < j_1 - j_r$ , unless  $j_{r-1} = j_r$ . By the exact sequence (4) and semi-continuity, the bundle  $E_2|_\ell$  is more balanced (in the sense of the Harder-Narasimhan filtration) than the bundle  $T$ . Hence we always have  $j(2, 1) - j(2, r) \leq j(1, 1) - j(1, r)$ , while  $j(2, 1) - j(2, r) < j(1, 1) - j(1, r)$  unless  $j(1, r-1) = j(1, r)$ . Even in this case, since  $\sum_{l=1}^r j(x, l) = k + \sum_{l=1}^r j(x-1, l)$ , we see that the procedure must stop after finitely many steps.

We now construct families of bundles having prescribed associated admissible sequence, generalising [BG3, Theorem 2.1], which constructed such families for rank-2 bundles near a  $-1$ -curve. The generalisation requires only minor modifications, but we give the details for completeness.

**Theorem 3.2.** Fix an admissible sequence  $\{j(i, l)\}_{i=1}^t$  and let  $E$  and  $F$  be rank- $r$  vector bundles on  $\widehat{\ell}$  with  $\{j(i, l)\}_{i=1}^t$  as an associated admissible sequence. Then there exists a flat family  $\{E_s\}_{s \in T}$  of rank- $r$  vector bundles on  $\widehat{\ell}$  parametrised by an integral variety  $T$  and  $s_0, s_1 \in T$  with  $E_{s_0} \cong E$  and  $E_{s_1} \cong F$  such that  $E_s$  has  $\{j(i, l)\}_{i=1}^t$  as admissible sequence for every  $s \in T$ .

*Proof.* We use induction on  $t$ . If  $t = 1$  the result is obvious because  $E$  and  $F$  are split vector bundles with the same splitting type and are hence isomorphic. Assume that  $t > 1$  and that the result is true for  $t - 1$ . Let  $E_2, F_2$  be the second bundles associated to  $E, F$  respectively. Hence  $E_2$  and  $F_2$  have  $\{j(2, l)\}_{i=2}^t$  as an associated admissible sequence. By induction, there is a flat family  $\{E'_s\}_{s \in S}$  of rank- $r$  vector bundles on  $\widehat{\ell}$  and  $m_0, m_1 \in S$  with  $E'_{m_0} \cong E_2, E'_{m_1} \cong F_2$  and such that  $E'_s$  has  $\{j(i, l)\}_{i=2}^t$  as an associated admissible sequence for every  $s \in S$ . We write  $j'_i = j(2, i)$  to simplify notation. By the balancing construction, the bundles  $E$  and  $F$  fit into exact sequences

$$\begin{aligned} 0 \rightarrow E \rightarrow E_2(\ell) \rightarrow \mathcal{O}_\ell(j_1 - k) \rightarrow 0 \\ 0 \rightarrow F \rightarrow F_2(\ell) \rightarrow \mathcal{O}_\ell(j_1 - k) \rightarrow 0 \end{aligned}$$

For every bundle  $M$  on  $\widehat{\ell}$  having  $\{j(i, l)\}_{i=2}^t$  as an associated admissible sequence, the set of surjective homomorphisms  $\mathbf{t}: M(\ell) \rightarrow \mathcal{O}_\ell(j_1 - k)$  is parametrised by an integral variety whose dimension depends only on  $j_1, j_2$  and  $j'_2 = j_1 + j_2 - j'_1 + k$ . The kernel  $\ker \mathbf{t}|_\ell$  is an extension of  $\mathcal{O}_\ell(j'_1)$  by  $\mathcal{O}_\ell(j_1)$ . This extension splits since  $j_1 \geq j'_1 + k$ , and hence the bundle  $\ker \mathbf{t}$  has  $\{j(i, l)\}_{i=1}^t$  as an admissible sequence. Varying  $M$  among bundles  $E'_s$  for  $s \in S$  we get that the set of all such surjections is parametrised by an irreducible non-empty variety  $T$ . For any fixed ample line bundle  $H$  on the  $n^{\text{th}}$  neighbourhood of  $\ell$ , it follows from the exact sequences in the balancing construction that the bundles in this family have the same Hilbert polynomial with respect to  $H$ , and therefore the family is flat.  $\square$

## 4 Moduli

**Definition 4.1.** We write

$$\mathcal{M}_j(k) = \{E \rightarrow Z_k : E|_\ell \cong \mathcal{O}(j) \oplus \mathcal{O}(-j)\} / \sim$$

for the moduli stack of bundles over  $Z_k$  of splitting type  $j$ .

Recall from (2) that a bundle  $E$  on  $Z_k$  of splitting type  $j$  is an extension of  $\mathcal{O}(j)$  by  $\mathcal{O}(-j)$  and is therefore determined by its extension class. In our choice of coordinates, this amounts to saying that  $E$  is determined by the pair  $(j, p)$ , where  $j$  is the splitting type and  $p$  is a polynomial as in (3). Let  $s = \lfloor \frac{2j-2}{k} \rfloor$ ; then  $p$  has  $N := s(2j-1) - k \binom{s+1}{2}$  coefficients. We identify  $p$  as an element in  $\mathbb{C}^N$  by writing its coefficients in lexicographical order. We then define the equivalence relation  $p \sim p'$  if  $(j, p)$  and  $(j, p')$  define isomorphic bundles over  $Z_k$ . We give  $\mathbb{C}^N$  the quotient topology. There is a bijection  $\phi: \mathcal{M}_j(k) \rightarrow \mathbb{C}^N / \sim$ . We give  $\mathcal{M}_j(k)$  the topology induced by this bijection. Here are some examples:

*Example 1.* For each  $k$ ,  $\mathcal{M}_0(k)$  contains only one point, corresponding to the trivial bundle over  $Z_k$ . In other words, if a bundle over  $Z_k$  is trivial over the zero section, it is globally trivial.

*Example 2.* For each  $k$ ,  $\mathcal{M}_1(k)$  contains only one point. In other words, a holomorphic bundle over  $Z_k$  of splitting type 1 splits. This can be verified directly from formula (3).

*Example 3.*  $\mathcal{M}_2(2)$  contains exactly 2 points ([So, Theorem 6.24]).

*Example 4.*  $\mathcal{M}_2(1) \simeq \mathbb{P}^1 \cup \{A, B\}$ , where  $A$  and  $B$  are points, with open sets  $U \subset \mathbb{P}^1$ , where  $U$  is open in the usual topology of  $\mathbb{P}^1$ ,  $\mathbb{P}^1 \cup \{A\}$ , and the whole space ([Ga2, Theorem 4.2]).

*Example 5.*  $\mathcal{M}_3(2) \simeq \mathbb{P}^2 \cup \{A, B\}$ , where  $A$  and  $B$  are points, with open sets  $U \subset \mathbb{P}^2$ , where  $U$  is open in the usual topology of  $\mathbb{P}^2$ ,  $\mathbb{P}^2 \cup \{A\}$ , and the whole space ([So, Theorem 6.35]).

*Example 6.*  $\mathcal{M}_j(k)$  is non-Hausdorff for  $j > k$ . This uses Theorem 4.3 below.

**Theorem 4.2.**  $\mathcal{M}_j(k)$  has an open, dense subspace homeomorphic to a complex projective space  $\mathbb{P}^{2j-k-2}$  minus a closed subvariety of codimension at least 2.

*Proof.* [Ga2, Theorem 3.5] showed that the generic set of  $\mathcal{M}_j(1)$  is a projective space  $\mathbb{P}^{2j-3}$  minus a closed subvariety of codimension  $\geq 2$ . The only modification needed to generalise the proof to  $k > 1$  is the calculation of dimension of the generic set. Generic points correspond to bundles that do not split on the first formal neighbourhood, and for such bundles the only equivalence relation is projectivisation, cf. [Ga3, Proposition 3.2]. The dimension count follows from formula (3), which shows that the  $u$ -coefficients are  $\sum_{s=k-j+1}^{j-1} p_{1s}$ . There are  $2j - k - 1$  coefficients, and after projectivising we obtain  $\mathbb{P}^{2j-k-2}$ . The closed subvariety to be removed contains all points having coefficients of  $u$  and of  $zu$  both zero; such points are not generic, see [BG1].  $\square$

**Theorem 4.3.** There is a topological embedding  $\Phi: \mathcal{M}_j(k) \rightarrow \mathcal{M}_{j+k}(k)$ . The image of  $\Phi$  consists of all bundles in  $\mathcal{M}_{j+k}(k)$  that split on the second formal neighbourhood of  $C$ .

*Proof.* Using the identification  $\phi: \mathcal{M}_j(k) \rightarrow \mathbb{C}^N / \sim$ , we define a map

$$\Phi: \mathcal{M}_j(k) \rightarrow \mathcal{M}_{j+k}(k) \quad \text{by} \quad (j, p) \mapsto (j+k, z^k u^2 p).$$

We want to show that  $\Phi$  defines an embedding. We first show that the map is well defined: Suppose that  $\begin{pmatrix} z^j & p \\ 0 & z^{-j} \end{pmatrix}$  and  $\begin{pmatrix} z^j & p' \\ 0 & z^{-j} \end{pmatrix}$  represent isomorphic bundles. Then there are coordinate changes  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  holomorphic in  $(z, u)$  and  $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$  holomorphic in  $(z^{-1}, z^k u)$  such that

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} z^j & p' \\ 0 & z^{-j} \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} z^{-j} & -p \\ 0 & z^j \end{pmatrix}.$$

Therefore these two bundles are isomorphic exactly when the system of equations

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} a + z^{-j} p' c & z^{2j} b + z^j (p' d - ap) - pp' c \\ z^{-2j} c & d - z^{-j} p c \end{pmatrix} \quad (*)$$

can be solved by a matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  holomorphic in  $(z, u)$  which makes  $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$  holomorphic in  $(z^{-1}, z^k u)$ .

On the other hand, the images of these two bundles are given by transition matrices  $\begin{pmatrix} z^{j+k} & z^k u^2 p \\ 0 & z^{-j-k} \end{pmatrix}$  and  $\begin{pmatrix} z^{j+k} & z^k u^2 p' \\ 0 & z^{-j-k} \end{pmatrix}$ , which represent isomorphic bundles if and only if there are coordinate changes  $\begin{pmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{pmatrix}$  holomorphic in  $(z, u)$  and  $\begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix}$  holomorphic in  $(z^{-1}, z^k u)$  satisfying the equality

$$\begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix} = \begin{pmatrix} z^{j+k} & z^k u^2 p' \\ 0 & z^{-j-k} \end{pmatrix} \begin{pmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{pmatrix} \begin{pmatrix} z^{-j-k} & -z^k u^2 p \\ 0 & z^{j+k} \end{pmatrix}.$$

That is, the images represent isomorphic bundles if the system

$$\begin{pmatrix} \bar{\alpha} & \bar{\beta} \\ \bar{\gamma} & \bar{\delta} \end{pmatrix} = \begin{pmatrix} \bar{a} + z^{-j} u^2 p' \bar{c} & z^{2k} (z^{2j} \bar{b} + z^j u^2 (p' \bar{d} - \bar{a} p) - u^4 p p' \bar{c}) \\ z^{-2j-2k} \bar{c} & \bar{d} - z^{-j} u^2 p \bar{c} \end{pmatrix} \quad (**)$$

has a solution.

Write  $x = \sum x_i u^i$  for  $x \in \{a, b, c, d, \bar{a}, \bar{b}, \bar{c}, \bar{d}\}$  and choose  $\bar{a}_i = a_{i+2k}$ ,  $\bar{b}_i = b_{i+2k} u^2$ ,  $\bar{c}_i = c_{i+2k} u^{-2}$ ,  $\bar{d}_i = d_{i+2k}$ . Then if  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  solves  $(*)$ , one verifies that  $\begin{pmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{pmatrix}$  solves  $(**)$ , which implies that the images represent isomorphic bundles and therefore  $\phi$  is well defined. To show that the map is injective, just reverse the previous argument. Continuity is obvious. Now we observe also that the image  $\phi(\mathcal{M}_j)$  is a saturated set in  $\mathcal{M}_{j+k}$ , meaning that if  $y \sim x$  and  $x \in \phi(\mathcal{M}_j)$  then  $y \in \phi(\mathcal{M}_j)$ . In fact, if  $E \in \phi(\mathcal{M}_j)$ , then  $E$  splits on the 2<sup>nd</sup> formal neighbourhood. Now if  $E' \sim E$ , then  $E'$  must also split on the 2<sup>nd</sup> formal neighbourhood, and therefore the polynomial corresponding to  $E'$  is of the form  $u^2 p'$ , and hence  $\phi(z^{-k} p')$  gives  $E'$ . Note also that  $\phi(\mathcal{M}_j)$  is a closed subset of  $\mathcal{M}_{j+k}$ , given by the equations  $p_{il} = 0$  for  $i = 1, 2$ . Now the fact that  $\phi$  is a homeomorphism over its image follows from the easy Lemma 4.5 given below.  $\square$

*Remark 4.4.* R. Moraru gave us the following coordinate-free expression of the embedding map  $\Phi: \mathcal{M}_j(k) \rightarrow \mathcal{M}_{j+k}(k)$ :

$$\Phi(E) = (- \otimes \mathcal{O}(-k)) \circ \text{Elm}_{\mathcal{O}_\ell(j+k)} \circ \text{Elm}_{\mathcal{O}_\ell(j)}(E) ,$$

where  $\text{Elm}_L$  denotes the elementary transformation with respect to the line bundle  $L$ . Using this coordinate-free expression it becomes obvious that  $\Phi$  is well defined.

**Lemma 4.5.** *Let  $X \subset Y$  be a closed subset and  $\sim$  an equivalence relation in  $Y$  such that  $X$  is  $\sim$ -saturated. Then the map  $I: X/\sim \rightarrow Y/\sim$  induced by the inclusion is a homeomorphism over the image.*

*Proof.* Denote the projections by  $\pi_X: X \rightarrow X/\sim$  and  $\pi_Y: Y \rightarrow Y/\sim$ . Let  $F$  be a closed subset of  $X/\sim$ . Then  $\pi_X^{-1}(F)$  is closed and saturated in  $X$ , and therefore  $\pi_X^{-1}(F)$  is also closed and saturated in  $Y$ . It follows that  $\pi_Y(\pi_X^{-1}(F))$  is closed in  $Y/\sim$ .  $\square$

**Theorem 4.6.** *The pair  $(\mathbf{h}_k, \mathbf{w}_k)$  stratifies  $\mathcal{M}_j(k)$  into Hausdorff components.*

*Proof.* This proof uses the same techniques as that of [BG1, Theorem 4.1]. On the first formal neighbourhood, we have two possibilities: in the first case we have bundles belonging to the open dense subset  $\mathbb{P}^{2j-k-1}$ , singled out by having the lowest possible values of the numerical invariants; in the second case, at least one of the invariants is strictly higher than the lower bound, and such bundles are separated away from the most generic stratum. On the second formal neighbourhood, the problem is solved by first separating the most generic stratum from the other ones. For the remaining part of the second neighbourhood, one divides the polynomial by  $u$  falling back to the same analysis done for the first neighbourhood. We are then left only with bundles which split on the second neighbourhood. We use induction  $j$ , assuming that the invariants stratify  $\mathcal{M}_{j-1}(k)$  into Hausdorff components together with the embedding Theorem 4.3, stating that  $\Phi(\mathcal{M}_{j-k}(k))$  is the set of bundles on  $\mathcal{M}_j(k)$  that split on the second formal neighbourhood.  $\square$

## References

- [BS] Bănică, C. and Stănăşilă, O. *Méthodes algébriques dans la théorie des espaces complexes*, Vol. I, Gauthier-Villars, Paris (1977).
- [BG1] Ballico, E. and Gasparim, E. *Numerical invariants for vector bundles on blow-ups*, Proc. Amer. Math. Soc. **130** (2002) 23–32.

- [BG2] Ballico, E. and Gasparim, E. *Vector bundles on a neighborhood of an exceptional curve and elementary transformations*, Forum Math. **15** (2003) 115–122.
- [BG3] Ballico, E. and Gasparim, E. *Vector bundles on a formal neighborhood of a curve in a surface*, Rocky Mountain J. Math. **30** (2000) 795–814.
- [BPV] Barth, W., Peters, C. and Van de Ven, A. *Compact Complex Surfaces*, Springer (1984).
- [Bl] Blache, R. *Chern classes and Hirzebruch–Riemann–Roch theorem for coherent sheaves on complex projective orbifolds with isolated singularities*, Math. Z. **222** (1996) 7–57.
- [Ga1] Gasparim, E. *Holomorphic bundles on  $\mathcal{O}(-k)$  are algebraic*, Comm. Algebra **25** (1997) 3001–3009.
- [Ga2] Gasparim, E. *Rank two bundles on the blow-up of  $\mathbb{C}^2$* , J. Algebra **199** (1998) 581–590.
- [Ga3] Gasparim, E. *Chern classes of bundles on blown-up surfaces*, Comm. Algebra **28** (2000) 4912–4926.
- [Ga4] Gasparim, E. and Ontaneda, P. *Three applications of instanton numbers*, Comm. Math. Phys. **270** (1) (2007) 1–12.
- [GKP] Gasparim, E., Köppe, T. and Majumdar, P. *Local holomorphic Euler characteristic and instanton decay*, preprint.
- [Gr] Griffiths, P. A. *The extension problem in complex analysis, II. Embeddings with positive normal bundle*, Amer. J. Math. **88** (1966) 366–446.
- [Ha] Hartshorne, R. *Algebraic Geometry*, Graduate Texts in Mathematics, Springer Verlag (1977).
- [HL] Henkin, G. and Leiterer, J. *The Oka–Grauert principle without induction over the base dimension*, Math. Ann. **311** (1998) 71–93.
- [LT] Lübke, M. and Teleman, A. *The Kobayashi–Hitchin correspondence*, World Scientific Publishing Co. Inc., River Edge, NJ (1995).
- [So] Souza, A. F. P. *Fibrados sobre  $K_{\mathbb{P}^1}$* , Master’s thesis UFPE (1999) (Portuguese).

Edoardo Ballico  
 University of Trento, Department of Mathematics  
 I-38050 Povo (Trento), Italia  
 E-mail: [ballico@science.unitn.it](mailto:ballico@science.unitn.it)

Elizabeth Gasparim and Thomas Köppe  
 School of Mathematics, The University of Edinburgh  
 James Clerk Maxwell Building, The King’s Buildings, Mayfield Road  
 Edinburgh, EH9 3JZ, United Kingdom  
 E-mail: [Elizabeth.Gasparim@ed.ac.uk](mailto:Elizabeth.Gasparim@ed.ac.uk)  
 E-mail: [t.koeppe@ed.ac.uk](mailto:t.koeppe@ed.ac.uk)